Title: Effects of Blood Sugar Levels, Kidney Disease and Medication on Anaemia
Status of Type 2 Diabetes Mellitus Patients

Malala Mulavu[1], Mildred Zulu[2], Musalula Sinkala[1], Sody Munsaka[1], Pauline Okuku[3],
Panji Nkhoma[1]

[1]Department of Biomedical Sciences, School of Health Sciences, University of Zambia,
Lusaka

[2]Department of Pathology and Microbiology, School of Medicine, University of Zambia,
Lusaka

[3]Department of Pathology and Microbiology, University Teaching Hospitals, Lusaka
12

14
15 Corresponding Author: Panji Nkhoma
16
17 E-mail:
18 panji.nkhoma@unza.zm
19
20 Postal Address:
21 The University of Zambia, School of Health Sciences, Department of Biomedical
22 Sciences, PO Box 50110, Lusaka, Zambia
23
24

ABSTRACT

Introduction: Lifestyle-related diseases like type 2 diabetes mellitus, have emerged as
a significant public health problem due to rapid urbanization and industrialization. In
2018, the prevalence of type 2 diabetes mellitus was estimated to be 500 million cases
worldwide and was comparable between high- and low-income countries. Diabetes
leads to multiple complications, including end-stage renal disease, cardiovascular
disease, infection, and death. Anaemia in diabetic individuals has severe adverse
effects on the quality of life and is associated with disease progression. A cross-
sectional study was conducted that included 101 participants. The study aimed to
determine the prevalence of anaemia and kidney disease and their interplay with
medication and/or blood glucose levels in T2D patients.

Results: seventy-one per cent of participants were females. Majority of patients 62%
were on Insulin, 30% on metformin, 7% on a combination of metformin and glimepiride
and 1% on glimepiride. Ninety-five (94%) of the participants were HIV negative. The
prevalence of anaemia among the participants was 23% out of which 56% had
moderate anaemia, and 44% had mild anaemia. Twenty-one per cent (21%) of the
participants had high creatinine levels signifying impaired kidney function or kidney
disease. Anaemia was significantly associated with kidney function, fasting blood
glucose and use of metformin; $p = 0.042$ beta = 2.5, $p = 0.025$ beta = 2.7 and $p = 0.040$
beta = -2.5 respectively.

Conclusion: The study found the prevalence of anaemia of 23%, which was of
moderate public health concern. Also, the prevalence of kidney disease was high in
patients with Type 2 Diabetes Mellitus. It also found that kidney disease and high blood
glucose levels increase the chances of developing anaemia. However, we found that
metformin had a protective role against the development of anaemia in Type 2 Diabetes
Mellitus patients.

Keywords: Type 2 diabetes mellitus, anaemia, creatinine, kidney disease, insulin,
metformin

1.0 INTRODUCTION

Lifestyle-related diseases like diabetes mellitus, have emerged as a significant public health problem due to rapid urbanization and industrialization [1]. In 2018, the prevalence of type 2 diabetes mellitus (T2D) was estimated to be 500 million cases worldwide and was comparable between high and low-income countries [2]. However, others have suggested that the condition may be up to two times more prevalent in low-income populations compared to wealthy populations [3–5]. The number of adults with diabetes mellitus in sub-Saharan Africa is predicted to rise from 19.8 million in 2013 to 41.5 million in 2035 [6–10], and in 2016, the age-standardised prevalence of diabetes in Zambia was 3.5% [11].

Diabetes is the leading causes of kidney disease [12]. About 1 out of 4 adults with diabetes have kidney disease [13]. It leads to multiple complications, including end-stage renal disease, cardiovascular disease, infection, and death [14]. In the recent past, the prevalence of diabetes and diabetic kidney disease has increased [15].

Anaemia is a reduction in the haemoglobin concentration of the blood below normal for age and sex, and the World Health Organisation defines anaemia as Hb level < 13 g/dL in men and 12 g/dL in women [16]. Anaemia in diabetic individuals has severe adverse effects on the quality of life and is associated with disease progression and the development of comorbidities [17]. Obesity and dyslipidaemias, which are strongly associated with the diabetic framework, significantly contribute to increasing the risk of cardiovascular diseases [18]. The study aimed to determine the prevalence of anaemia and kidney disease and their interplay with medication and/or blood glucose levels in

T2D patients at the University Teaching Hospital, which is the largest third-level referral

in Zambia.

2.0 MATERIALS AND METHODS

2.1 Study design

The study was cross-sectional and was carried out at the University Teaching Hospital,

a tertiary hospital in Lusaka, Zambia.

2.2 Study Population

The study population included all existing and newly diagnosed type 2 diabetes mellitus

patients visiting UTH from April to July 2019. We used convenience sampling to select

participants.

2.3 Specimen Collection

Blood samples were collected by phlebotomy. From each patient 3ml of blood, were

collected in each of two tubes; one Ethylenediaminetetraacetic acid (EDTA) for

haematological analysis, and lithium heparin tube for biochemical analysis (kidney

function tests).

2.4 Full Blood Count, Kidney Function and Fasting Blood Sugar

Full blood count was measured using Sysmex XT- 4000i (Germany) automated

analyser. Kidney function measurements were carried out using the ABX Pentra 400,

Japanese chemistry analyzer. Glucose was measured using an Accu check™

glucometer at the point of specimen collection. The patient's clinical records were

collected using the UTH laboratory information system. All quality control procedures

were followed before running the specimens.

2.5 Statistical Analysis

The data were entered in Excel & analysed using Python 3.7 for mac. Continuous variables were summarised using mean, standard deviation, percentiles, range and mode. Descriptive statistics in the form of percentages and graphs were used to show the prevalence of anaemia and kidney disease in type 2 diabetic mellitus patients. Multiple logistic regression was used to determine the effects of Kidney function, fasting blood glucose and medication on the occurrence of anaemia and the Student t-test was used to determine the mean differences in the kidney function tests between study participants and the non-diabetic control group.

2.6 Ethical Approval

Ethical approval for the study was obtained from The University of Zambia Health Science Research Ethics Committee (UNZAHSREC) (Protocol ID: 20190117001 of IORG0009227). Informed Consent was obtained from all study participants.

3.0 RESULTS

3.1 Demographic Characteristics

One hundred and one participants took part in the study out of which 71 (70%) were females. Minimum age of participants was 22 years, maximum of 83 years with a mean age of 54.43 ± 14 years. The majority of patients 63 (62%) were on Insulin, 30 (30%) on metformin, 7 (7%) on a combination of metformin and glimepiride and only 1 (1%) on glimepiride (Figure 1). Ninety-five (94%) of the participants were HIV negative.

137   *Figure 1: Distribution of Antidiabetic medication of participants*

138
139

3.2 Fasting Blood Sugar

141   The minimum fasting blood glucose recorded was 3.8mmol/l, maximum 24.9 mmol/l

142   with a mean blood glucose of 10.24 $\pm$ 4.42mmol/l. 78% had higher than normal fasting

143   blood glucose levels. 22% had normal blood glucose levels, and none had low blood

144   glucose levels (Figure 2).

Figure 2: Fasting blood glucose levels of participants

147
3.3 Anaemia

The prevalence of anaemia among the participants was 23%, of which 23 out of the 101 participants were anaemic (figure 3). The minimum haemoglobin concentration was 9.7g/dl, and the maximum was 17.2g/dl with the mean haemoglobin concentration of 13.75 $\pm$ 1.63 g/dl. Concerning the severity of anaemia, we found that 56% of the anaemic patients had moderate anaemia, 44% had mild anaemia with none having severe anaemia (Figure 4). We defined the severity and classification of anaemia based on the World Health Organization (WHO) criteria [16]

157   *Figure 3: Prevalence of anaemia*
158

159
Figure 4: Anaemia severity

Out of the 23 anaemic participants, 67% had microcytic, hypochromic anaemia, 33 %

had normochromic normocytic anaemia, and none of them had macrocytic anaemia.

3.4 Kidney Function

We found that twenty-one per cent (21%) of the participants had higher than normal

(110µmol/l) creatinine levels signifying impaired kidney function or kidney disease

(Figure 5). The minimum creatinine level that we recorded was 50.8 µmol/l, while the

maximum level was at 404.8 µmol/l with a mean of 100.77 ± 55.96 µmol/l. We used

creatine to assess kidney function because serum creatinine above specific cutoff

points reliably identifies patients with acute kidney injury or chronic kidney disease

[19,20].

174
Figure 5: Prevalence of kidney disease

Anaemia was significantly associated with kidney function (p = 0.042; beta = 2.5),

fasting blood glucose (p = 0.025; beta = 2.7) and use of metformin (p = 0.040; beta = -

2.5)

After Adjusting for the potential confounding factors; sex, alcohol consumption and HIV

status:

i. We found that the odds of having anaemia were 11.76 times higher for

participants with kidney malfunction (OR = 11.76, 95% CI =1.09 - 126.68, p =

0.042).

ii. Furthermore, we showed that the odds of having anaemia were 15.45 times

higher for participants with high fasting blood glucose (OR = 15.45, 95% CI

=1.40 - 170.37, p = 0.025).

iii. For the participants taking metformin, we found that their odds of having

anaemia were 0.09 times less, and showed a negative beta value of -2.5 (OR

= 0.09, 95% CI = 0.01 - 0.89, p = 0.040), denoting that these participants on

metformin were about 91% protected from developing anaemia.

193
We compared the plasma creatinine, urea, sodium, chloride and potassium levels

between the diabetic participants to those of the non-diabetic individuals. Here, we

found that the creatinine (p = 1.36×10^{-6}), urea (p = 0.011), sodium (p = 1.84×10^{-9}) and

chloride (p = 2.59×10^{-7}) levels were significantly higher in the diabetic study

participants than in the non-diabetic control group (Figure 6 and 7). However, potassium

ion levels between the two groups were not statistically significant, with a p-value of

0.8222.

201
202   *Figure 6: Mean Differences in creatinine and urea between diabetic participants and*
203   *non-diabetic controls*
204

205
206
207   *Figure 7: Mean Differences in sodium, potassium and chloride ions between diabetic*
208   *participants and non-diabetic controls*
209

4.0 DISCUSSION

Diabetes Mellitus is frequently accompanied by mild to moderate anaemia, which is often called anaemia of inflammation or infection or anaemia of chronic disease [21]. The reduced Hb levels identify diabetic patients at increased risk for hospitalization and premature death [22]. Our study found that nearly one in four diabetic patients have anaemia which, according to WHO is of moderate public health significance [23]. Others have also found comparable prevalences of anaemia in India (18%) and Kuwait (29.7%) [1,24].

The predominant risk factor for the development of anaemia in a diabetic population is the presence of renal disease, which manifests as either impaired renal function or albuminuria [25]. In particular, the critical factor underlying the development of anaemia is the uncoupling of the haemoglobin concentration from renal erythropoietin synthesis [24]. In our study, t[26][25]he high incidence of anaemia may also be due to other risk factors that are related to DM. Several studies have reported factors that increase the risk of anaemia, which include; damage to renal interstitium due to chronic hyperglycaemia and consequent formation of advanced glycation end products by increased reactive oxygen species, and systemic inflammation as well as reduced androgen levels induced by diabetes [27]. Here also, the increased levels of inflammatory cytokines (IL 6) may promote the apoptosis of immature erythrocytes, thus reducing the number of circulating erythrocytes and haemoglobin levels [17,28].

We revealed that the majority of participants had microcytic hypochromic anaemia. Our findings differ from those obtained in a study done in Ghana, where they found that; the

majority of participants had normochromic normocytic anaemia, followed by

hypochromic microcytic, and lastly, normochromic macrocytic anaemia [29]. Here, the

observed difference between the findings Ghana study and our study may be due to the

differences in the duration of illness. In our study, the majority of the participants had

diabetes for less than five years. Conversely, the part participants in the Ghana study

may have had diabetes for a more extended period resulting in normochromic

normocytic anaemia which is commonly associated with anaemia of chronic diseases

such as diabetes.

We found that none of our study participants had severe anaemia, but the majority had

moderate anaemia. We suggest that since our study participants were diagnosed with

diabetes within the previous five years, it is unlikely that they would develop increased

levels of inflammation with decreased erythropoietin levels both which cause severe

anaemia in diabetes. Here, others have also shown that people with diabetes have mild

to moderate anaemia [1,30].

Kidney disease is a common occurrence in patients with T2D. Approximately 30% of

patients with diabetic nephropathy eventually progress to end-stage renal failure, and

the rest usually die from cardiovascular disease before reaching end-stage [31]. The

increased blood glucose level that leads to damage of the kidney glomerular capsules is

a significant cause of kidney disease in diabetics. Here also, high blood pressure can

damage the kidney blood vessels, thereby leading to glomerular failure.

Our results revealed that 21% of participants had an elevated serum creatinine. Here,

our finding is similar to those of a study done in the USA, where they found a

prevalence of CKD of 24.1% in T2D patients [32]. Another similar study carried out in

Finland showed that 68.9% of patients presented with signs of CKD out of which 34.7%

appeared to suffer from significant CKD [33]. Another study done in Zimbabwe on

African diabetics found that 44.8% of the patients suffered from nephropathy [34]. All

the highlighted studies demonstrated a much higher prevalence of CKD compared to

our study. Here also, the majority of our study participants were diagnosed with

diabetes within five years; therefore, had little damage to the glomerulus and did not

have many complications that are associated with high blood pressure. Creatinine, urea

and electrolytes (sodium and chloride) were higher in the diabetic group compared to

the non-diabetic control group. These findings are consistent with the current literature,

which indicates that electrolyte imbalance is a common feature of diabetes mellitus

associated with hyperglycaemia [35–37]. Furthermore, the observed lower levels of

potassium in the diabetic group compared to the non-diabetic group can be explained

by the noted high levels of insulin and glucose in the diabetic group, both of which

promote the cellular influx of potassium from extracellular compartment [38–40].

We found that participants with high fasting blood glucose were significantly more likely

to develop anaemia. We suggest that this may be because chronic hyperglycaemia may

result in abnormal red blood cells, oxidative stress, and sympathetic denervation of the

kidney related to autonomic neuropathy. These factors promote a hypoxic environment

in the renal interstitium, which leads to impaired production of erythropoietin by the

peritubular fibroblasts [27]. This suggests that the incidence of anaemia is likely to

increase in poorly controlled diabetes, and therefore reducing blood glucose levels

could assist in the reduction of the risk of anaemia in diabetic populations.

We also found that the odds of having anaemia were 0.09 times less for participants

taking metformin. The odds ratio for metformin is less than one, which means that

metformin significantly reduces the chances of developing anaemia in diabetes

individuals and therefore plays a protective role. This is contrary to studies that have

suggested that metformin leads to anaemia in patients with diabetes due to the resulting

vitamin B12 deficiency that is caused by prolonged use of the drug [41]. This difference

may be due to vitamin B12 deficiency anaemia that occurs after prolonged use, while

the majority of our study participants had diabetes for less than 5yrs. Therefore, they

may not have used metformin for an extended period.

5.0 CONCLUSION

We found that there is a prevalence of 23% of anaemia and 21% of kidney disease in

patients with type 2 diabetes mellitus at the University Teaching Hospital. Furthermore,

we found that patients with kidney disease and increased blood glucose levels have an

increased chance of developing anaemia. Also, we found that metformin played a

protective role in the development of anaemia in our study participants. We suggest,

therefore, that among T2D patients, it is essential to monitor regularly for anaemia and

kidney disease the two conditions are typical ofT2D patients.

6.0 DECLARATION OF COMPETING INTEREST

none

7.0 FUNDING

8.0 AUTHOR CONTRIBUTIONS

Conceptualisation: Panji Nkhoma, Mildred Zulu, Musalula SInkala, Malala Mulavu.

Data Analysis: Panji Nkhoma, Musalula Sinkala, Sody Munsaka.

Investigation: Malala Mulavu, Pauline Okuku

Supervision: Panji Nkhoma, Mildred Zulu, Musalula Sinkala, Sody Munsaka

Writing original draft: Panji Nkhoma, Mildred Zulu, Sody Munsaka, Musalula Sinkala

9.0REFERENCES

1.   Rathod GB, Parmar P, Rathod S, Parikh A. Prevalence of anemia in patients with Type 2 diabetes mellitus at Gandhinagar, Gujarat, India. Int Arch Integ Med. 2016;3: 12–16.

2.   Kaiser AB, Zhang N, Van der Pluijm W. Global prevalence of type 2 diabetes over the next ten years (2018-2028). Am Diabetes Assoc; 2018.

3.   Bird Y, Lemstra M, Rogers M, Moraros J. The relationship between socioeconomic status/income and prevalence of diabetes and associated conditions: A cross-sectional population-based study in Saskatchewan, Canada. Int J Equity Health. 2015;14. doi:10.1186/s12939-015-0237-0

4.   Robbins JM, Vaccarino V, Zhang H, Kasl S V. Socioeconomic status and type 2 diabetes in African American and non-Hispanic White women and men: Evidence

from the Third National Health and Nutrition Examination Survey. Am J Public
Health. 2001;91: 76–83. doi:10.2105/AJPH.91.1.76

5.   Stelmach W, Kaczmarczyk-Chałas K, Bielecki W, Drygas W. How education,
income, control over life and life style contribute to risk factors for cardiovascular
disease among adults in a post-communist country. Public Health. 2005;119:
498–508. doi:10.1016/j.puhe.2004.09.006
341

342 6.   Mbanya JCN, Motala AA, Sobngwi E, Assah FK, Enoru ST. Diabetes in sub-
343   Saharan Africa. The Lancet. 2010. pp. 2254–2266. doi:10.1016/S0140-
344   6736(10)60550-8
345

346 7.   Miranda JJ, Kinra S, Casas JP, Davey Smith G, Ebrahim S. UKPMC Funders
347   Group countries⬜: context , determinants and health policy. Trop Med Int Heal.
348   2008;13: 1225–1234. doi:10.1111/j.1365-3156.2008.02116.x.Non-communicable
349

350 8.   Hall V, Thomsen R, Henriksen O, Lohse N. Diabetes in Sub Saharan Africa 1999-
351   2011: Epidemiology and public health implications. A systematic review. BMC
352   Public Health. 2011;11. doi:10.1186/1471-2458-11-564
353

354 9.   Danaei G, Finucane MM, Lu Y, Singh GM, Cowan MJ, Paciorek CJ, et al.
355   National, regional, and global trends in fasting plasma glucose and diabetes
356   prevalence since 1980: Systematic analysis of health examination surveys and
357   epidemiological studies with 370 country-years and 2·7 million participants.
358   Lancet. 2011;378: 31–40. doi:10.1016/S0140-6736(11)60679-X
359

360 10. Whiting DR, Guariguata L, Weil C, Shaw J. IDF diabetes atlas: global estimates of
361   the prevalence of diabetes for 2011 and 2030. Diabetes Res Clin Pract. 2011;94:
362   311–321.
363

364 11. Bailey S Lou, Ayles H, Beyers N, Godfrey-Faussett P, Muyoyeta M, du Toit E, et
365   al. Diabetes mellitus in Zambia and the Western Cape province of South Africa:
366   Prevalence, risk factors, diagnosis and management. Diabetes Res Clin Pract.
367   2016;118: 1–11. doi:10.1016/j.diabres.2016.05.001
368

369 12. Diabetic Kidney Disease | NIDDK. [cited 28 Dec 2019]. Available:
370   https://www.niddk.nih.gov/health-information/diabetes/overview/preventing-
371   problems/diabetic-kidney-disease
372

373 13. Afkarian M, Zelnick LR, Hall YN, Heagerty PJ, Tuttle K, Weiss NS, et al. Clinical
374   manifestations of kidney disease among US adults with diabetes, 1988-2014.
375   JAMA - J Am Med Assoc. 2016;316: 602–610. doi:10.1001/jama.2016.10924
376

377 14. Tuttle KR, Bakris GL, Bilous RW, Chiang JL, De Boer IH, Goldstein-Fuchs J, et al.
378   Diabetic kidney disease: A report from an ADA consensus conference. Diabetes
379   Care. 2014;37: 2864–2883. doi:10.2337/dc14-1296
380

15. De Boer IH, Rue TC, Hall YN, Heagerty PJ, Weiss NS, Himmelfarb J. Temporal
trends in the prevalence of diabetic kidney disease in the United States. JAMA - J
Am Med Assoc. 2011;305: 2532–2539. doi:10.1001/jama.2011.861

16. Peña-Rosas JP. WHO. Haemoglobin concentrations for the diagnosis of anaemia
and assessment of severity. Vitamin and Mineral Nutrition Information System.
Geneva, World Health Organization, 2011 (WHO/NMH/NHD/MNM/11.1).

17. Angelousi A, Larger E. Anaemia, a common but often unrecognized risk in
diabetic patients: a review. Diabetes Metab. 2015;41: 18–27.

18. Martínez-Pérez B, De La Torre-Díez I, López-Coronado M. Mobile health
applications for the most prevalent conditions by the World Health Organization:
review and analysis. J Med Internet Res. 2013;15: e120.

19. Sinkala M, Zulu M, Kaile T, Simakando M, Chileshe C, Kafita D, et al.
Performance characteristics of kidney injury molecule-1 in relation to creatinine,
urea, and microalbuminuria in the diagnosis of kidney disease. Int J Appl Basic
Med Res. 2017;7: 94. doi:10.4103/2229-516x.205811

20. Zulu M, Kaile T, Kantenga T, Chileshe C, Nkhoma P, Sinkala M. Kidney injury
molecule-1 and microalbuminuria levels in Zambian population: Biomarkers of
kidney injury. Pan Afr Med J. 2016;24. doi:10.11604/pamj.2016.24.54.8759

21. de Carvalho MC, Baracat ECE, Sgarbieri VC. Anemia ferropriva e anemia de
doença crônica: distúrbios do metabolismo de ferro. Segurança Aliment e Nutr.
2006;13: 54–63.

22. Keane WF, Lyle PA. Recent advances in management of type 2 diabetes and
nephropathy: lessons from the RENAAL study. Am J Kidney Dis. 2003;41: S22–
S25.

23. Organization WH. Haemoglobin concentrations for the diagnosis of anaemia and
assessment of severity. World Health Organization; 2011.

24. AlDallal SM, Jena N. Prevalence of Anemia in Type 2 Diabetic Patients. J
Hematol. 2018;7: 57–61.

25. Craig KJ, Williams JD, Riley SG, Smith H, Owens DR, Worthing D, et al. Anemia
and diabetes in the absence of nephropathy. Diabetes Care. 2005;28: 1118–
1123.

26. Thomas MC. Anemia in diabetes: Marker or mediator of microvascular disease?
Nature Clinical Practice Nephrology. 2007. pp. 20–30. doi:10.1038/ncpneph0378

427
428
429
430   27.   Bosman DR, Winkler AS, Marsden JT, Macdougall IC, Watkins PJ. Anemia with
431   erythropoietin deficiency occurs early in diabetic nephropathy. Diabetes Care.
432   2001;24: 495–499. doi:10.2337/diacare.24.3.495
433
434   28.   Fava S, Azzopardi J, Ellard S, Hattersley AT. ACE gene polymorphism as a
435   prognostic indicator in patients with type 2 diabetes and established renal
436   disease. Diabetes Care. 2001;24: 2115–2120.
437
438   29.   Antwi-Bafour S, Hammond S, Adjei JK, Kyeremeh R, Martin-Odoom A, Ekem I. A
439   case-control study of prevalence of anemia among patients with type 2 diabetes.
440   J Med Case Rep. 2016;10. doi:10.1186/s13256-016-0889-4
441
442   30.   Almahdi MA, Emhemmed MB, Mariam MI, Nuzha TG. Prevalence of anemia and
443   its association with renal function in diabetic patients from south of Libya. LJMR.
444   2016;10: 47–51.
445
446   31.   Atkins RC. The epidemiology of chronic kidney disease. Kidney Int. 2005;67:
447   S14–S18.
448
449   32.   Iglay K, Hannachi H, Howie PJ, Xu J, Li X, Engel SS, et al. Prevalence and co-
450   prevalence of comorbidities among patients with type 2 diabetes mellitus. Curr
451   Med Res Opin. 2016;32: 1243–1252. doi:10.1185/03007995.2016.1168291
452
453   33.   Metsärinne K, Bröijersen A, Kantola I, Niskanen L, Rissanen A, Appelroth T, et al.
454   High prevalence of chronic kidney disease in Finnish patients with type 2 diabetes
455   treated in primary care. Prim Care Diabetes. 2015;9: 31–8.
456   doi:10.1016/j.pcd.2014.06.001
457
458   34.   MacHingura PI, Chikwasha V, Okwanga PN, Gomo E. Prevalence of and factors
459   associated with nephropathy in diabetic patients attending an outpatient clinic in
460   Harare, Zimbabwe. Am J Trop Med Hyg. 2017;96: 477–482.
461   doi:10.4269/ajtmh.15-0827
462
463   35.   Ojiako OA, Chikezie PC. Blood Na+/K+ and Cl-Levels of Hyperglycemic Rats
464   Administered with Traditional Herbal Formulations. Pharmacogn Commun.
465   2015;5: 140–145.
466
467   36.   Javaid A, Hasan R, Zaib A, Mansoor S. A comparative study of the effects of
468   hypoglycemic agents on serum electrolytes in the diabetic patients. Pak J Pharm
469   Sci. 2007;20: 67–71. Available: http://www.ncbi.nlm.nih.gov/pubmed/17337432
470
471   37.   Khan RN, Saba F, Kausar SF, Siddiqui MH. Pattern of electrolyte imbalance in
472   Type 2 diabetes patients: Experience from a tertiary care hospital. Pakistan J Med

473   Sci. 2019;35: 797–801. doi:10.12669/pjms.35.3.844
474
475
476 38. Adrogué HJ, Lederer ED, Suki WN, Eknoyan G. Determinants of plasma
477   potassium levels in diabetic ketoacidosis. Medicine (Baltimore). 1986;65: 163–
478   172.
479
480 39. Li T, Vijayan A. Insulin for the treatment of hyperkalemia: a double-edged sword?
481   Clin Kidney J. 2014;7: 239–41. doi:10.1093/ckj/sfu049
482
483 40. Allon M. Hyperkalemia in end-stage renal disease: Mechanisms and
484   management. Journal of the American Society of Nephrology. 1995. pp. 1134–
485   1142.
486
487 41. Mulla K, Bradbury K. Metformin use and vitamin B12 deficiency. Society for
488   Endocrinology BES 2018. BioScientifica; 2018.
489
490