TITLE: EFFECTS OF BLOOD SUGAR LEVELS, KIDNEY DISEASE AND MEDICATION ON ANAEMIA STATUS OF TYPE 2 DIABETES MELLITUS PATIENTS

Malala Mulavu[1], Mildred Zulu[2], Musalula Sinkala[1], Sody Munsaka[1], Pauline Okuku[3], Panji Nkhoma[1]

[1]Department of Biomedical Sciences, School of Health Sciences, University of Zambia, Lusaka

[2]Department of Pathology and Microbiology, School of Medicine, University of Zambia, Lusaka

[3]Department of Pathology and Microbiology, University Teaching Hospitals, Lusaka

Corresponding Author: Panji Nkhoma

E-mail:
panji.nkhoma@unza.zm

Postal Address:
The university of Zambia, School of Health Sciences, Department of Biomedical Sciences, PO Box 50110, Lusaka, Zambia

ABSTRACT

Introduction: Lifestyle-related diseases like type 2 diabetes mellitus, have emerged as a significant public health problem due to rapid urbanization and industrialization. In 2018, the prevalence of type 2 diabetes mellitus was estimated to be 500 million cases worldwide and was comparable between high- and low-income countries. Diabetes leads to multiple complications, including end-stage renal disease, cardiovascular disease, infection, and death. Anaemia in diabetic individuals has a severe adverse effect on the quality of life and is associated with disease progression. A cross-sectional study was conducted that included 101 participants. The study aimed to determine the prevalence of anaemia and kidney disease and their interplay with medication and/or blood glucose levels in T2D patients.

Results: seventy one percent of participants were females. Majority of patients 62% were on Insulin, 30% on metformin, 7% on a combination of metformin and glimepiride and 1% on glimepiride. Ninety-five (94%) of the participants were HIV negative. The prevalence of anaemia among the participants was 23% out of which 56% had moderate anaemia, and 44% had mild anaemia. Twenty-one per cent (21%) of the participants had high creatinine levels signifying impaired kidney function or kidney disease. Anaemia was significantly associated with kidney function, fasting blood glucose and use of metformin; $p = 0.042$ beta = 2.5, $p = 0.025$ beta = 2.7 and $p = 0.040$ beta = -2.5 respectively.

Conclusion: The study found the prevalence of anaemia of 23% which was of moderate public health concern. Also, the prevalence of kidney disease was high in patients with Type 2 Diabetes Mellitus. It also found that kidney disease and high blood glucose levels increase the chances of developing anaemia. However, we found that metformin had a protective role against the development of anaemia in Type 2 Diabetes Mellitus patients.

Keywords: Type 2 diabetes mellitus, anaemia, creatinine, kidney disease, insulin, metformin

1.0 INTRODUCTION

Lifestyle-related diseases like diabetes mellitus, have emerged as a significant public health problem due to rapid urbanization and industrialization [1]. In 2018, the prevalence of type 2 diabetes mellitus (T2D) was estimated to be 500 million cases worldwide and was comparable between high and low-income countries [2]. However, others have suggested that the condition may be up to two times more prevalent in low-income populations compared to wealthy populations [3–5]. The number of adults with diabetes mellitus in sub-Saharan Africa is predicted to rise from 19.8 million in 2013 to 41.5 million in 2035 [6–10], and in 2016, the age-standardised prevalence of diabetes in Zambia was 3.5% [11].

Diabetes is the leading causes of kidney disease [12]. About 1 out of 4 adults with diabetes have kidney disease [13]. It leads to multiple complications, including end-stage renal disease, cardiovascular disease, infection, and death [14]. In the recent past, the prevalence of diabetes and diabetic kidney disease have increased [15].

Anaemia is defined as a reduction in the haemoglobin concentration of the blood below normal for age and sex, and the World Health Organisation defines anaemia as Hb level < 13 g/dL in men and 12 g/dL in women [16]. Anaemia in diabetic individuals has a severe adverse effect on the quality of life and is associated with disease progression and the development of comorbidities [17]. Obesity and dyslipidaemias, which are strongly associated with the diabetic framework, significantly contribute to increasing the risk of cardiovascular diseases [18]. The study aimed to determine the prevalence of anaemia and kidney disease and their interplay with medication and/or blood glucose

levels in T2D patients at the University Teaching Hospital, which is the largest third-level

referral in Zambia.

2.0 MATERIALS AND METHODS

2.1 Study design

The study was cross-sectional and was carried out at the University Teaching Hospital,

a tertiary hospital in Lusaka, Zambia.

2.2 Study Population

The study population included all existing and newly diagnosed type 2 diabetes mellitus

patients visiting UTH from April to July 2019. We used convenience sampling to select

participants.

2.3 Specimen Collection

Blood samples were collected by phlebotomy. From each patient 3ml of blood, were

collected in each of two tubes; one Ethylenediaminetetraacetic acid (EDTA) for

haematological analysis, and lithium heparin tube for chemistry (kidney function tests).

2.4 Full Blood Count, Kidney function and Fasting Blood Sugar

Full blood count was measured using Sysmex XT- 4000i (Germany) automated

analyser. Kidney function measurements were carried out using the ABX Pentra 400,

Japanese chemistry analyzer. Glucose was measured using an Accu check™

glucometer at the point of specimen collection. Patient clinical records were collected

using the UTH laboratory information system. All quality control procedures were

followed before running the specimens

2.5 Statistical Analysis

Data was entered in Excel & analysed using Python 3.7 for mac. Continuous variables were summarised using mean, standard deviation, percentiles, range and mode. Descriptive statistics in the form of percentage and graphs were used to show the prevalence of anaemia and kidney disease in type 2 diabetic mellitus patients. Multiple logistic regression was used to determine the effects of Kidney function, fasting blood glucose and medication on the occurrence of anaemia and a t-test was used to determine the mean differences in kidney function tests between study participants and the non-diabetic control group.

2.6 Ethical Approval

Ethical approval for the study was obtained from The University of Zambia Health Science Research Ethics Committee (UNZAHSREC) (Protocol ID: 20190117001 of IORG0009227). Informed Consent was obtained from all study participants.

3.0 RESULTS

3.1 Demographic Characteristics

One hundred and one participants took part in the study out of which 71 (70%) were females. Minimum age of participants was 22 years, maximum of 83 years with a mean age of 54.43 ± 14 years. The majority of patients 63 (62%) were on Insulin, 30 (30%) on metformin, 7 (7%) on a combination of metformin and glimepiride and only 1 (1%) on glimepiride (Figure 1). Ninety-five (94%) of the participants were HIV negative.

Figure 1: Distribution of Antidiabetic medication of participants

3.2 Fasting Blood Sugar

The minimum fasting blood glucose recorded was 3.8mmol/l, maximum 24.9 mmol/l with a mean blood glucose of 10.24 ± 4.42mmol/l. 78% had higher than normal fasting blood glucose levels. 22% had normal blood glucose levels, and none had low blood glucose levels (Figure 2).

144
145 **Figure 2: Fasting blood glucose levels of participants**

146
3.3 Anaemia

The prevalence of anaemia among the participants was 23% which means 23 out of the

101 participants had anaemia (figure 3). The minimum haemoglobin concentration was

9.7g/dl, and maximum was 17.2g/dl with the mean haemoglobin concentration of 13.75

± 1.63 g/dl. In terms of anaemia severity, 56% of the anaemic patients had moderate

anaemia, 44% had mild anaemia with none having severe anaemia (Figure 4). Anaemia

severity and classification was defined according to World Health Organization (WHO)

criteria [16]

Figure 3: Prevalence of anaemia

158
Figure 4: Anaemia severity

Out of the 23 anaemic participants, 67% had microcytic, hypochromic anaemia, 33 %

had normochromic normocytic anaemia, and none of them had macrocytic anaemia.

3.4 Kidney Function

Twenty-one per cent (21%) of the participants had higher than normal (110μmol/l)

creatinine levels signifying impaired kidney function or kidney disease (Figure 5). The

minimum creatinine level recorded was 50.8 μmol/l, maximum 404.8 μmol/l with a mean

of 100.77 ± 55.96 μmol/l.

Figure 5: Prevalence of kidney disease

Anaemia was significantly associated with kidney function (p = 0.042; beta = 2.5),

fasting blood glucose (p = 0.025; beta = 2.7) and use of metformin (p = 0.040; beta = -

2.5)

After Adjusting for the potential confounding factors; sex, alcohol consumption and HIV

status:

i.   The odds of having anaemia were 11.76 times higher for participants with

kidney malfunction (OR = 11.76, 95% CI =1.09 - 126.68, p = 0.042).

ii.   The odds of having anaemia were 15.45 times higher for participants with

high fasting blood glucose (OR = 15.45, 95% CI =1.40 - 170.37, p = 0.025).

iii.   The odds of having anaemia were 0.09 times less for participants taking

metformin which had a negative beta value of -2.5 (OR = 0.09, 95% CI = 0.01

- 0.89, p = 0.040). This means participants on metformin were about 91%

protected from developing anaemia.

We compared the plasma creatinine, urea, sodium, chloride and potassium levels

between the diabetic participants to those of the non-diabetic individuals. Here, we

found that the creatinine ($p = 1.36 \times 10^{-6}$), urea (p = 0.011), sodium ($p = 1.84 \times 10^{-9}$) and

chloride ($p = 2.59 \times 10^{-7}$) levels were significantly higher in the diabetic study

participants than in the non-diabetic control group (Figure 6 and 7). However, potassium

ion levels between the two groups were not statistically significant, with a p-value of

0.8222.

Figure 6: Mean Differences in creatinine and urea between diabetic participants and non-diabetic controls

Figure 7: Mean Differences in sodium, potassium and chloride ions between diabetic participants and non-diabetic controls

4.0 DISCUSSION

Diabetes Mellitus is frequently accompanied by mild to moderate anaemia, often called anaemia of inflammation or infection or anaemia of chronic disease [19]. The reduced Hb levels identify diabetic patients at increased risk for hospitalization and premature death [20]. Our study found that nearly one in four diabetic patients had anaemia which, according to WHO is of moderate public health significance [21]. Others have also found comparable anaemia prevalence in India (18%) and Kuwait (29.7%) [1,22].

The predominant risk factor for the development of anaemia in a diabetic population is the presence of renal disease, manifested as an impaired renal function or albuminuria [23]. In particular, the uncoupling of the haemoglobin concentration from renal erythropoietin synthesis seems to be the critical factor underlying the development of anaemia [24]. The high incidence of anaemia may also be due to other risk factors related to DM. Several studies have reported factors that increase the risk of anaemia, which include; damage to renal interstitium due to chronic hyperglycaemia and consequent formation of advanced glycation end products by increased reactive oxygen species, and systemic inflammation as well as reduced androgen levels induced by diabetes [25]. Increased levels of inflammatory cytokines (IL 6) also promotes apoptosis of immature erythrocytes causing a decrease, further, in the number of circulating erythrocytes and consequently causing a reduction of circulating haemoglobin [17,26].
We revealed that the majority of participants had microcytic hypochromic anaemia. This differs from a study carried out in Ghana, where they found that the majority of participants had normochromic normocytic anaemia, followed by hypochromic

microcytic and lastly the lowest percentage had normochromic macrocytic anaemia [27].

These differences may be due to the differences in the duration of illness. Majority of

our participants had diabetes for less than five years. The part participants in the Ghana

study may have had diabetes for a more extended period resulting in normochromic

normocytic anaemia which is commonly associated with anaemia of chronic diseases

such as diabetes.

We found that no study participants suffered from severe anaemia, with the majority

having moderate anaemia. This finding may be because most of our study participants

were diagnosed with diabetes within the last five years, therefore, are unlikely to have

developed increased levels of inflammation and decreased erythropoietin levels that

may cause severe anaemia. Here, others have also shown that people with diabetes

have mild to moderate anaemia [1,28].

Kidney disease is a common occurrence in patients with T2D. Approximately 30% of

patients with diabetic nephropathy eventually progress to end-stage renal failure, and

the rest usually die from cardiovascular disease before reaching end-stage [29]. Some

of the causes of kidney disease in people with diabetes are an increased blood

glucoselevel that results in damage to the kidney glomerular capsules. High blood

pressure is also a cause of damage to the kidney blood vessels, thereby causing

malfunctioning filtration apparatus.

Our results revealed that 21% of participants had an elevated serum creatinine. This

was similar to the study done in the USA, where they found a prevalence of CKD of

24.1% in T2D patients [30]. Another similar study carried out in Finland showed that

68.9% of patients showed some signs of CKD out of which 34.7% seemed to suffer

from significant CKD [31]. Another study done in Zimbabwe on African diabetics found

that 44.8% of patients suffered from nephropathy [32]. These studies all found a much

higher prevalence of CKD compared to our study. Here also, the majority of our study

participants were diagnosed with diabetes within five years; therefore, had little damage

to the glomerulus and other complications associated with high blood pressure.

Creatinine, urea and electrolytes (sodium and chloride) were higher in our participants

compared to the control group. Electrolyte imbalance is a common feature of diabetes

mellitus and is associated with hyperglycaemia [33–35]. The low levels of potassium in

our study participants can be explained by the high levels of insulin and glucose, both of

which promote the cellular influx of potassium from extracellular compartment [36–38].

We found that participants with high fasting blood glucose were significantly more likely

to develop anaemia. We suggest that this may be because chronic hyperglycaemia may

result in abnormal red blood cells, oxidative stress, and sympathetic denervation of the

kidney related to autonomic neuropathy. These factors promote a hypoxic environment

in the renal interstitium, which leads to impaired production of erythropoietin by the

peritubular fibroblasts [25]. This suggests that the incidence of anaemia is likely to

increase in poorly controlled diabetes, and therefore reducing blood glucose levels

could assist in the reduction of the risk of anaemia in diabetic populations.

We also found that the odds of having anaemia were 0.09 times less for participants

taking metformin. The odds ratio for metformin is less than one, which means that

metformin significantly reduces the chances of developing anaemia in diabetes

individuals and therefore plays a protective role. This is contrary to studies that have

suggested that metformin leads to anaemia in patients with diabetes due to the resulting

vitamin B12 deficiency that is caused by prolonged use of the drug [39]. This difference

may be due to vitamin B12 deficiency anaemia that occurs after prolonged use, while

the majority of our study participants had diabetes for less than 5yrs. Therefore, they

may not have used metformin for an extended period.

5.0 CONCLUSION

The study found that there is a prevalence of 23% of anaemia and 21% of kidney

disease in patients with type 2 diabetes mellitus at the University Teaching Hospital. It

also found that patients with kidney disease and increased blood glucose levels have an

increased chance of developing anaemia. The study additionally found that metformin

played a protective role against the development of anaemia in our study participants. It

is, therefore, important to regularly monitor for anaemia and kidney disease among T2D

patients as we have shown that the two conditions are common T2D patients.

291
292
293   **6.0 DECLARATION OF COMPETING INTEREST**

294   none

7.0 AUTHOR CONTRIBUTIONS

Conceptualisation: Panji Nkhoma, Mildred Zulu, Musalula SInkala, Malala Mulavu.

Data Analysis: Panji Nkhoma, Musalula Sinkala, Sody Munsaka.

Investigation: Malala Mulavu, Pauline Okuku

Supervision: Panji Nkhoma, Mildred Zulu, Musalula Sinkala, Sody Munsaka

Writing original draft: Panji Nkhoma, Mildred Zulu, Sody Munsaka, Musalula Sinkala

8.0REFERENCES

1. Rathod GB, Parmar P, Rathod S, Parikh A. Prevalence ofanemia in patients with Type 2 diabetes mellitus at Gandhinagar, Gujarat, India. Int Arch Integ Med. 2016;3: 12–16.

2. Kaiser AB, Zhang N, Van der Pluijm W. Global prevalence of type 2 diabetes over the next ten years (2018-2028). Am Diabetes Assoc; 2018.

3. Bird Y, Lemstra M, Rogers M, Moraros J. The relationship between socioeconomic status/income and prevalence of diabetes and associated conditions: A cross-sectional population-based study in Saskatchewan, Canada. Int J Equity Health. 2015;14. doi:10.1186/s12939-015-0237-0

4. Robbins JM, Vaccarino V, Zhang H, Kasl S V. Socioeconomic status and type 2 diabetes in African American and non-Hispanic White women and men: Evidence from the Third National Health and Nutrition Examination Survey. Am J Public Health. 2001;91: 76–83. doi:10.2105/AJPH.91.1.76

5. Stelmach W, Kaczmarczyk-Chałas K, Bielecki W, Drygas W. How education, income, control over life and life style contribute to risk factors for cardiovascular disease among adults in a post-communist country. Public Health. 2005;119: 498–508. doi:10.1016/j.puhe.2004.09.006

6. Mbanya JCN, Motala AA, Sobngwi E, Assah FK, Enoru ST. Diabetes in sub-Saharan Africa. The Lancet. 2010. pp. 2254–2266. doi:10.1016/S0140-6736(10)60550-8

7. Miranda JJ, Kinra S, Casas JP, Davey Smith G, Ebrahim S. UKPMC Funders
Group countries☐: context , determinants and health policy. Trop Med Int Heal.
2008;13: 1225–1234. doi:10.1111/j.1365-3156.2008.02116.x.Non-communicable
8. Hall V, Thomsen R, Henriksen O, Lohse N. Diabetes in Sub Saharan Africa 1999-
2011: Epidemiology and public health implications. A systematic review. BMC
Public Health. 2011;11. doi:10.1186/1471-2458-11-564

9. Danaei G, Finucane MM, Lu Y, Singh GM, Cowan MJ, Paciorek CJ, et al.
National, regional, and global trends in fasting plasma glucose and diabetes
prevalence since 1980: Systematic analysis of health examination surveys and
epidemiological studies with 370 country-years and 2·7 million participants.
Lancet. 2011;378: 31–40. doi:10.1016/S0140-6736(11)60679-X

10. Whiting DR, Guariguata L, Weil C, Shaw J. IDF diabetes atlas: global estimates of
the prevalence of diabetes for 2011 and 2030. Diabetes Res Clin Pract. 2011;94:
311–321.

350
351

11. Bailey S Lou, Ayles H, Beyers N, Godfrey-Faussett P, Muyoyeta M, du Toit E, et
al. Diabetes mellitus in Zambia and the Western Cape province of South Africa:
Prevalence, risk factors, diagnosis and management. Diabetes Res Clin Pract.
2016;118: 1–11. doi:10.1016/j.diabres.2016.05.001

12. Diabetic Kidney Disease | NIDDK. [cited 28 Dec 2019]. Available:
https://www.niddk.nih.gov/health-information/diabetes/overview/preventing-
problems/diabetic-kidney-disease

13. Afkarian M, Zelnick LR, Hall YN, Heagerty PJ, Tuttle K, Weiss NS, et al. Clinical
manifestations of kidney disease among US adults with diabetes, 1988-2014.
JAMA - J Am Med Assoc. 2016;316: 602–610. doi:10.1001/jama.2016.10924

14. Tuttle KR, Bakris GL, Bilous RW, Chiang JL, De Boer IH, Goldstein-Fuchs J, et al.
Diabetic kidney disease: A report from an ADA consensus conference. Diabetes
Care. 2014;37: 2864–2883. doi:10.2337/dc14-1296

15. De Boer IH, Rue TC, Hall YN, Heagerty PJ, Weiss NS, Himmelfarb J. Temporal
trends in the prevalence of diabetic kidney disease in the United States. JAMA - J
Am Med Assoc. 2011;305: 2532–2539. doi:10.1001/jama.2011.861

16. Peña-Rosas JP. WHO. Haemoglobin concentrations for the diagnosis of anaemia
and assessment of severity. Vitamin and Mineral Nutrition Information System.
Geneva, World Health Organization, 2011 (WHO/NMH/NHD/MNM/11.1).

17. Angelousi A, Larger E. Anaemia, a common but often unrecognized risk in
diabetic patients: a review. Diabetes Metab. 2015;41: 18–27.

18.   Martínez-Pérez B, De La Torre-Díez I, López-Coronado M. Mobile health
applications for the most prevalent conditions by the World Health Organization:
review and analysis. J Med Internet Res. 2013;15: e120.

19.   de Carvalho MC, Baracat ECE, Sgarbieri VC. Anemia ferropriva e anemia de
doença crônica: distúrbios do metabolismo de ferro. Segurança Aliment e Nutr.
2006;13: 54–63.

20.   Keane WF, Lyle PA. Recent advances in management of type 2 diabetes and
nephropathy: lessons from the RENAAL study. Am J Kidney Dis. 2003;41: S22–
S25.

21.   Organization WH. Haemoglobin concentrations for the diagnosis of anaemia and
assessment of severity. World Health Organization; 2011.

22.   AlDallal SM, Jena N. Prevalence of Anemia in Type 2 Diabetic Patients. J
Hematol. 2018;7: 57–61.

23.   Craig KJ, Williams JD, Riley SG, Smith H, Owens DR, Worthing D, et al. Anemia
and diabetes in the absence of nephropathy. Diabetes Care. 2005;28: 1118–
1123.

24.   Thomas MC. Anemia in diabetes: Marker or mediator of microvascular disease?
Nature Clinical Practice Nephrology. 2007. pp. 20–30. doi:10.1038/ncpneph0378

25.   Bosman DR, Winkler AS, Marsden JT, Macdougall IC, Watkins PJ. Anemia with
erythropoietin deficiency occurs early in diabetic nephropathy. Diabetes Care.
2001;24: 495–499. doi:10.2337/diacare.24.3.495

26.   Fava S, Azzopardi J, Ellard S, Hattersley AT. ACE gene polymorphism as a
prognostic indicator in patients with type 2 diabetes and established renal
disease. Diabetes Care. 2001;24: 2115–2120.

27.   Antwi-Bafour S, Hammond S, Adjei JK, Kyeremeh R, Martin-Odoom A, Ekem I. A
case-control study of prevalence of anemia among patients with type 2 diabetes.
J Med Case Rep. 2016;10. doi:10.1186/s13256-016-0889-4

28.   Almahdi MA, Emhemmed MB, Mariam MI, Nuzha TG. Prevalence of anemia and
its association with renal function in diabetic patients from south of Libya. LJMR.
2016;10: 47–51.

29.   Atkins RC. The epidemiology of chronic kidney disease. Kidney Int. 2005;67:
S14–S18.

30.   Iglay K, Hannachi H, Howie PJ, Xu J, Li X, Engel SS, et al. Prevalence and co-

prevalence of comorbidities among patients with type 2 diabetes mellitus. Curr Med Res Opin. 2016;32: 1243–1252. doi:10.1185/03007995.2016.1168291

31. Metsärinne K, Bröijersen A, Kantola I, Niskanen L, Rissanen A, Appelroth T, et al. High prevalence of chronic kidney disease in Finnish patients with type 2 diabetes treated in primary care. Prim Care Diabetes. 2015;9: 31–8. doi:10.1016/j.pcd.2014.06.001

32. MacHingura PI, Chikwasha V, Okwanga PN, Gomo E. Prevalence of and factors associated with nephropathy in diabetic patients attending an outpatient clinic in Harare, Zimbabwe. Am J Trop Med Hyg. 2017;96: 477–482. doi:10.4269/ajtmh.15-0827

33. Ojiako OA, Chikezie PC. Blood Na+/K+ and Cl-Levels of Hyperglycemic Rats Administered with Traditional Herbal Formulations. Pharmacogn Commun. 2015;5: 140–145.

34. Javaid A, Hasan R, Zaib A, Mansoor S. A comparative study of the effects of hypoglycemic agents on serum electrolytes in the diabetic patients. Pak J Pharm Sci. 2007;20: 67–71. Available: http://www.ncbi.nlm.nih.gov/pubmed/17337432

35. Khan RN, Saba F, Kausar SF, Siddiqui MH. Pattern of electrolyte imbalance in Type 2 diabetes patients: Experience from a tertiary care hospital. Pakistan J Med Sci. 2019;35: 797–801. doi:10.12669/pjms.35.3.844

36. Adrogué HJ, Lederer ED, Suki WN, Eknoyan G. Determinants of plasma potassium levels in diabetic ketoacidosis. Medicine (Baltimore). 1986;65: 163–172.

37. Li T, Vijayan A. Insulin for the treatment of hyperkalemia: a double-edged sword? Clin Kidney J. 2014;7: 239–41. doi:10.1093/ckj/sfu049

38. Allon M. Hyperkalemia in end-stage renal disease: Mechanisms and management. Journal of the American Society of Nephrology. 1995. pp. 1134–1142.

39. Mulla K, Bradbury K. Metformin use and vitamin B12 deficiency. Society for Endocrinology BES 2018. BioScientifica; 2018.