Iodine staining as a useful probe for distinguishing amyloid fibril polymorphs

Takato Hiramatsu[1], Naoki Yamamoto[2], Seongmin Ha[1], Yuki Masuda[1], Mitsuru Yasuda[3], Mika Ishigaki[4,5], Keisuke Yuzu[1], Yukihiro Ozaki[3,6], and Eri Chatani[1*]

[1]Graduate School of Science, Kobe University, 1-1 Rokkodai, Nada, Kobe, Hyogo 657-8501, Japan

[2]School of Medicine, Jichi Medical University, 3311-1 Yakushiji, Shimotsuke, 329-0498, Japan

[3]School of Science and Technology, Kwansei Gakuin University, 2-1 Gakuen, Sanda, Hyogo 669-1337, Japan

[4]Raman Project Center for Medical and Biological Applications, Shimane University, 1060 Nishikawatsu, Matsue, Shimane 690-8504, Japan

[5]Faculty of Life and Environmental Sciences, Shimane University, 1060 Nishikawatsu, Matsue, Shimane 690-8504, Japan

[6]Molecular Photoscience Research Center, Kobe University, 1-1 Rokkodai, Nada, Kobe, Hyogo 657-8501, Japan

*To whom correspondence should be addressed: Eri Chatani,

Graduate School of Science, Kobe University, 1-1 Rokkodai-cho, Nada-ku, Kobe 657-8501, Japan

chatani@crystal.kobe-u.ac.jp; Tel. +81-78-803-5673, FAX: +81-78-803-6442.

ABSTRACT: Amyloid fibrils are protein aggregates with needle-like morphology and cross-β structure that are associated with amyloidoses and neurodegenerative diseases. It is recently suggested that amyloid polymorphism, i.e., structural diversity of amyloid fibrils, has a deep relationship with pathology, attracting a great deal of attention for understanding molecular mechanism of amyloid diseases. However, its prompt recognition is almost halted due to inadequate development of analytical methods for detecting polymorphism of amyloid fibrils sensitively and quickly. Here, we propose that iodine staining, a coloring reaction that occurs by complexing iodine with host compounds to form polyiodide ions, can be used as a method for distinguishing polymorphs of amyloid fibrils. When three polymorphs of insulin fibrils, NaCl fibrils, SDS fibrils, and no-salt fibrils, were prepared and iodine-stained, they exhibited different colors. Furthermore, each of these colors was inherited to daughter fibrils by seeding reactions. The colors were fundamentally represented as a sum of three absorption bands in visible region between 400-750 nm, and these bands showed different titration curves against iodine, suggesting that there are more than one specific binding sites for the formation of polyiodide ions. The analysis of resonance Raman spectra and polarization microscope suggested the formation of several polyiodide species composed of I_3^- and/or I_5^- on the grooves or the edges of the β-sheets. It was concluded that the production of different polyiodide species and conformations are the molecular origin of the variation in colors depending on the surface structure of amyloid fibrils.

INTRODUCTION

Amyloid fibrils are protein aggregates associated with many amyloidosis and neurodegenerative diseases including Alzheimer's disease (AD) and Parkinson's disease [1]. While straight unbranched morphology and cross-β structure are known as a common basic structure of amyloid fibrils, it was revealed that amyloid fibrils show structural variations even from the same protein; this phenomenon is referred to as amyloid polymorphism [2-4]. Variations in the number and the arrangement of protofilaments are an example of polymorphism [5,6], and more microscopic variations at the level of secondary or tertiary structure of polypeptide chains have been also identified in many proteins [7-13]. In studies of prion diseases, different physicochemical properties, such as fragility and growth rates, among amyloid polymorphs have been suggested to be the molecular basis underlying the presence and propagation of distinct prion strains [14]. Amyloid polymorphs also exert different neurotoxic effects on living cells [8], and it has recently been suggested that tau protein adopted distinct amyloid conformations in the human brain in different diseases [13]. The formation of such conformationally distinct amyloid fibrils now attracts much attention for understanding amyloid-related pathologies and developing curative treatments in terms of protein structure.

Amyloid polymorphs have been successfully reproduced in vitro depending on growth conditions, e.g., temperature [14], protein concentration [15], and solvent [16]; however the most critical factors have not been fully revealed yet [4]. Subtle variations in the amino acid sequence among different species also result in different fibril structures [17], although it does not correspond to amyloid polymorphism because amino acid sequences are not the same. In another case, distinct fibril structures can be also induced in a stochastic manner even under the same conditions [18]. Recently, detailed structural investigations of amyloid fibrils are progressing by using several powerful methods, such as solid-state NMR (ssNMR) [8,10], cryo-electron microscopy (cryo-EM) [11-13], and X-ray crystallography [7]. The clarified structures have

revealed distinct molecular structures of amyloid polymorphs. The relevance of amyloid polymorphism to in vivo pathology has also been clarified by the structural investigation of $A\beta_{1-40}$ amyloid fibrils that were prepared with seeds derived from brain tissues of AD patients with distinct clinical histories [19]. However, the use of these techniques requires a lot of labor and time and therefore making them unsuitable for high throughput use. If an easy-to-use method of analyzing fibril structures can be developed, it will become a great help to cover a wide range of samples including pathological tissues as targets of investigations, thereby contributing to progress toward a unified understanding of relationships between fibril structures and pathologies.

Here, we focus on iodine as a molecular probe that recognizes structural polymorphs of amyloid fibrils. Iodine staining is a coloring reaction that occurs by complexing iodine with crystalline compounds or polymers. The molecular origin of the color is the formation and subsequent alignment and orientation of polyiodide ions. A iodine-starch complex is a good example, and from the crystal structure of iodine-doped α-cyclodextrin, it is suggested that linear chains of pentaiodide ions (I_5^-) are encapsulated in channel-like voids in the center of the helical polysaccharides [20,21]. The coloration by polyiodide ions is also observed in the iodine-doped synthetic polymers such as polyacetylene [22], nylon 6 [23] and polyvinyl alcohol [24]. Interestingly, colors formed by the iodine staining vary sensitively depending on constituting polyiodide ions and their coordination patterns inside the complex structures [25,26]. From these observations, it is postulated that the color exhibited by the iodine staining will serve as a sensitive indicator of structures of host compounds to which iodine binds. With regard to iodine staining of amyloid fibrils, Virchow firstly found that tissue deposits of amyloid fibrils were stained with iodine [27]. However, soon after this pioneering discovery in the 1850s, fluorescent dyes such as Congo red and thioflavin S or thioflavin T (ThT) with higher detection sensitivity replaced iodine for diagnosis of amyloidosis [28,29], and molecular mechanisms underlying iodine

reaction have almost not been studied; indeed, there is only one report by Dzwolak [9].

In this work, we have performed iodine staining of human insulin amyloid fibrils using three polymorphs obtained under different additive conditions (i.e., in the presence of NaCl or SDS and in the absence of additives). Insulin is a hormone protein associated with glucose metabolism and consist of two polypeptide chains linked by two intermolecular disulfide bonds, which has been used as a good model for amyloid researches in a past few decades [30]. As a result of the characterization of color properties of each type of insulin fibrils by ultraviolet-visible (UV-Vis) spectra, we have found that the color significantly varied depending on the polymorphs. Each color conserved even after the self-seeding reaction, demonstrating that the iodine staining tracked the propagation of amyloid polymorphs. Polyiodide ions formed by iodine staining were characterized by resonance Raman spectra and titration experiments. Furthermore, iodine-stained insulin spherulites were observed by polarization microscopy to gain insights into the binding manner of the polyiodide ions to amyloid fibrils. It was proposed that different polyiodide species were formed depending on the surface structures of amyloid fibrils, which resulted in different colors. From the results obtained, molecular mechanisms of iodine staining of amyloid fibrils and future perspectives for the use of iodine staining as a probe for amyloid polymorphism will be discussed.

MATERIALS AND METHODS

Materials. Recombinant human insulin was purchased from Wako Pure Chemical Industries, Ltd., Osaka, Japan. Insulin was dissolved in 25 mM HCl, and whose concentration was determined using an absorption coefficient of 1.08 for 1.0 mg/ml at 276 nm [31]. Iodine, potassium iodide, and ThT were also purchased from Wako Pure Chemical Industries, Ltd. HCl solution at a concentration of 100 mM, NaCl, and sodium dodecyl sulfate (SDS) were purchased from Nacalai Tesque (Kyoto, Japan).

Formation of amyloid fibrils of insulin. Human insulin dissolved in 25 mM HCl at a concentration of 2.0 mg/ml was used as a fundamental sample solution. Fibrillation reaction was performed by incubating this solution at 65 °C for 24 hours. For inducing polymorphism of amyloid fibrils, two types of additives, 100 mM NaCl or 100 µM SDS were used in addition to the solvent condition without additives. After the formation of amyloid fibrils, which were referred to as parent fibrils, a seed-dependent fibril formation was performed to obtain daughter fibrils. In this reaction, the fragments of fibrils were added to 2.0 mg/ml insulin dissolved in 25 mM HCl as seeds at a final concentration of 5 % (w/v), and then incubated at 37 °C for 24 hours. It should be noted that all of the seed-dependent fibrillation reactions were performed in the same solvent conditions without any additives. For the preparation of seeds, the parent fibrils were subjected to pulsed sonication using XL2000 sonicator (Misonix, San Diego, CA, USA) operating for 1 sec with power level at 2.0 W. The total number of sonication pulses was 40, and it was ensured that amyloid samples are not excessively heated during the sonication treatment. After the formation of the daughter fibrils, the same seed-dependent fibril formation was repeated using the daughter fibrils as seeds to obtain granddaughter fibrils.

ThT fluorescence assay. ThT fluorescence assay was performed according to the method of Naiki et al. [32] with a slight modification of solution; 4.5 µl aliquots of sample solution was mixed with 1.5 ml of 5 µM ThT in 50 mM Gly-NaOH buffer (pH 8.5) at room temperature. After 1 min for equilibrium, fluorescence intensity at 485 nm was measured with an excitation wavelength of 445 nm with RF-5300 PC (Shimadzu Corporation, Kyoto, Japan) to assay the formation of amyloid fibrils.

Atomic force microscopy (AFM). Samples of insulin amyloid fibrils were diluted 250

times with 25 mM HCl, and 10 µl aliquots of which were deposited onto a freshly cleaved dry mica surface. The solution was adsorbed for 1 min and then the mica surface was washed with 100 µl of water. The solution was blotted off by blotting paper and the mica plate was air-dried. AFM images were obtained by using SPA-400 and nano navi (Hitachi High-Tech Science, Tokyo, Japan). The scanning tip was a OMCL-AC160TS-C3 micro cantilever (Olympus Corporation, Tokyo, Japan; spring constant = 21-37 N/m, resonance frequency = 270-340 kHz), and the scan rate was 1.0 Hz.

Attenuated total reflection Fourier transform infrared (ATR-FTIR) absorption spectroscopy. ATR-FTIR spectra were measured with a FT/IR-6100 (JASCO, Tokyo, Japan). 2 µl of sample solution was blotted onto a germanium prism and then dried using compressed air. The spectral resolution and the number of scan were set to 4 cm^{-1} and 128, respectively. A spectrum of atmospheric air was used as a reference spectrum to subtract the contributions of water vapor and carbon dioxide from each sample spectrum.

Iodine staining of amyloid fibrils. The color formation of insulin fibrils by the addition of iodine was analyzed by using UV-Vis spectroscopy. For the preparation of a sample, iodine solution containing 24 mM KI and 3.2 mM I_2 in 25 mM HCl was initially prepared, and 0.25 mg/ml insulin fibrils containing 0.3 mM KI, and 0.04 mM I_2 in 25 mM HCl was made by mixing sample fibrils with the iodine solution. UV-Vis spectra from 250 to 750 nm were then measured at room temperature with a UV-2400PC (Shimadzu Corporation) or Jasco V-650 spectrometer (JASCO Corporation, Tokyo, Japan) with a quartz cell with 1 cm optical length. The reaction between fibrils and iodine molecules was sufficiently rapid to reach equilibrium within 1 min under the present solution conditions, and thus we performed the measurement between 1 and 10 min after preparing the sample. For tracking seed-dependent fibrillation reactions, an aliquot

of a reaction mixture was taken every one hour for sample preparation and subsequent spectral measurement, which were performed under the same conditions as those mentioned above. For titration experiments, 1 ml of 0.5 mg/ml insulin fibrils dissolved in 25 mM HCl was prepared as an analyte, to which an iodine solution containing 18 mM KI and 2.4 mM I_2 in 25 mM HCl was titrated. A UV-Vis spectrum was monitored every after the addition of 4 μl of the iodine solution to the fibrils, and this procedure was repeated until spectral saturation was observed.

Resonance Raman spectroscopy. Raman resonance spectra were measured by using two types spectrometers, HR-800-LWR (HORIBA, Kyoto, Japan) with 514.5 nm excitation and inVia Raman microscope (Renishaw, Gloucestershire, UK) with 785 nm excitation. Both of the spectrometers were equipped with notch filters for suppressing Rayleigh scattering. A sample containing 0.5 mg/ml insulin fibrils, 0.6 mM KI, and 0.08 mM I_2 in 25 mM HCl was taken in a glass capillary with its inner diameter of 1.13 mm (Drummond Scientific Company, Broomall, PA, USA) and placed on a sample stage. An excitation light was focused on the sample solution closed to the wall surface of the capillary to minimize reabsorption of Raman scattered light by the sample. Raman scattering was measured with back-scattering mode from 0 to 500 cm^{-1} at room temperature.

Polarization microscopy of spherulites. To investigate polarization properties of the polyiodide ions bound on amyloid fibrils, microscopy observation of iodine-stained insulin spherulites was performed with an ECLIPSE LV100POL (Nikon, Tokyo, Japan), in which an analyzer was placed in the optical pathway between an objective lens and an eyepiece. To prepare spherulites, insulin solution at a concentration of 5.0 mg/ml in 25 mM HCl was sealed between slide and cover glasses and incubated at 70 °C for 2 hours, according to the method reported by Rogers et al [33]. The spherulites formed were then stained with iodine by adding a

small amount of iodine solution containing 24 mM KI and 3.2 mM I_2 in in 25 mM HCl. Microscope images were recorded with a DS-L3 digital camera (Nikon) at different angles of rotations of the analyzer.

RESULTS

Formation of polymorphic insulin amyloid fibrils. To prepare polymorphic amyloid fibrils for iodine staining, fibrillation of human insulin was performed. It is already revealed that insulin amyloid fibrils are formed spontaneously at high temperature under acidic conditions [34-36]. In this work, 25 mM HCl and 65 °C were selected as standard conditions, and amyloid polymorphs were attempted to be obtained by using two types of additives, NaCl and SDS. As a result of incubating the sample solutions for 24 hours, ThT fluorescence exhibited different intensity among the amyloid fibrils formed (Figure 1A, blue bars). In light of a previous report that the affinity and stoichiometry between ThT and amyloid fibrils change depending on fibril structures [37], the difference in fluorescence intensity suggests that amyloid polymorphs were formed. Indeed, when a seed-dependent fibril formation was performed with the amyloid fibrils formed to produce daughter fibrils, the relative differences in ThT fluorescence intensity conserved (Figure 1A, red bars). The daughter fibrils all exhibited higher intensity than that of the parent fibrils, which could not be explained only by the addition of 5% seeds. This is probably because a fraction of non-fibrillar aggregates that could not self-propagate contained in the sample of the parent fibrils. Furthermore, significant differences in peak wavelength were found in addition to the differences in intensity in ThT fluorescence spectra (Figure 1B), which strongly supports the production of amyloid polymorphs. We refer to the amyloid fibrils generated in the absence and presence of NaCl and SDS as no-salt fibrils, NaCl fibrils, and SDS fibrils, respectively.

We next analyzed morphology and secondary structure of the amyloid fibrils

Figure 1. Basic properties of three types of insulin amyloid fibrils prepared in the presence of NaCl (NaCl fibrils) or SDS (SDS fibrils) or in its absence (no-salt fibrils). (A) ThT fluorescence intensity. Blue and orange bars represent fluorescence intensity of parent and daughter fibrils, respectively. All measurements were performed three times and the error bars depict the S.D. ± mean. (B) ThT fluorescence spectra of daughter fibrils. (C) AFM images. The white scale bars represent 1 μm. The scale bars on the right represent the height of the sample. (D) ATR-FTIR absorption spectra (upper panel) and their second derivatives (lower panel) in amide I region.

generated from the three different additive conditions by AFM and ATR-FTIR spectroscopy.

AFM images showed variety in fibril thickness, and the NaCl fibrils were markedly thicker (~ 10 nm) compared to SDS (~ 4 nm) and no-salt fibrils (~ 4 nm) (Figure 1C). On the other hand, when ATR-FTIR spectra were analyzed, all fibrils exhibited a main peak at 1630 cm^{-1} in amide I region, which was assigned to β-sheet structure (Figure 1D; upper panel) [38]. Second derivatives of the spectra also failed to detect difference in the peak wavelength among NaCl,

SDS, and no-salt fibrils. However, differences in the positions of minima at around 1670 cm^{-1} and 1710-1750 cm^{-1}, which are assigned to β-turn and carboxyl groups of side chains, respectively, were observed (Figure 1D; lower panel) [38]. This result suggests that, in the present case, polymorphism is derived from differences in turn and/or higher-order structures including hierarchical architecture, while the main cross-β architecture is very similar.

Color formation reaction by mixing iodine solution and insulin amyloid fibrils. To investigate iodine staining of amyloid fibrils, we tested the formation of colors by mixing iodine solution with NaCl, SDS, or no-salt fibrils. In the iodine solution, three types of iodine species, I$^-$, I$_2$, and I$_3^-$, are present in equilibrium and its absorption spectrum shows two major peaks and one broad one, which are derived from I$_3^-$ and I$_2$, respectively [39,40] (Figure S1). Immediately after the mixing, a formation of a reddish or bluish color was observed for all types of fibrils (Figure 2A). Given that native insulin mixed with iodine solution showed quite a similar UV-Vis absorption spectrum to that observed for the iodine solution-only control (Figure S1A), these observations support that iodine can specifically bind to fibril structures. Interestingly, the color tone appeared to vary among types of fibrils, and absorption spectra in UV-Vis region showed characteristic spectral shapes in a visible region (400-750 nm) depending on the types of fibrils (Figure 2B). The spectra as well as their second derivatives showed that iodine-stained NaCl fibrils had a major absorption at around 550 nm (Figure 2B, red line), whereas iodine-stained SDS and no-salt fibrils had one at around 650 nm (Figure 2B, blue and orange lines, respectively), which accounted for the color differences spectroscopically. Although a contamination of salts derived from the parent fibrils may affect the spectral shape, it was denied by confirming unchanged spectra even after the addition of salts (Figure S2).

We next performed spectral deconvolution to discuss absorption bands constructing the experimental spectrum. Each spectrum was replotted against wavenumber so that the analysis

Figure 2. Color properties of iodine-stained insulin amyloid fibrils. (A) Photographs of samples of iodine-stained NaCl fibrils, SDS fibrils, and no-salt fibrils. (B) Absorption spectra in UV-Vis region (upper panel) and their second derivatives (lower panel) of iodine-stained fibrils. For each absorption spectrum, the spectrum of unstained fibrils was subtracted to obtain net spectrum derived from iodine molecules. It should be noted that the scales of the vertical axes for the region of 400-740 nm and that of 260-400 nm are different in the lower panel. The conditions of iodine staining were 0.25 mg/ml insulin, 0.3 mM KI and 0.04 mM I_2 in 25 mM HCl.

was performed in a manner proportional to energy, and then was reproduced by a sum of

Gaussian distributions according to the following model function;

$$A(v) = \sum_{j=1}^{n} A_j \exp\left\{-\frac{(v-v_j)^2}{2\sigma_j^2}\right\} + b_0 + b_1 v \tag{1}$$

where A_j and v_j represent amplitude, center wavenumber of the jth absorption band, respectively, and σ_j is a coefficient related to the full width at half-maximum (FWHM) of the

*j*th absorption band, as is described as $2\sigma_j\sqrt{2\ln 2}$. b_0 and b_1 are intercept and slope of the baseline for Rayleigh scattering caused by fibrils, respectively. Curve fitting was performed until both measured and second derivative spectra were reproduced well at the same time, and for stable and accurate convergence, the wavenumber and FWHM of one band located at lower wavenumber were fixed to values read from the local minimum and its distance to the inflection point in the second derivative spectrum in the present study.

As a result, all of the absorption spectra were reproduced successfully by approximating with six Gaussians (i.e., $n = 6$) and a linear baseline (Figure 3). Although it was hardly noticeable in the second derivative spectrum (see Figure 2B), the installation of another band between 23×10^3 and 25×10^3 cm^{-1} (i.e., between 400 and 435 nm), which was conceived to be derived from I_2 molecules (see Figure S1) [39,40], certainly improved the reproduction of the experimental spectra. When the six absorption bands obtained were compared to the absorption bands observed in the iodine solution (Figure S1B), the three absorption bands at higher wavenumbers (dashed lines in Figure 3A-C) were observed at similar positions to those for the I_2 and I_3^- species in the iodine solution (dashed lines in Figure S1B). The remaining three absorption bands at lower wavenumbers, on the other hand, were observed only for the fibril-bound iodine (filled bands in Figure 3A-C). It was thus suggested that the color formation in iodine-amyloid fibrils complexes was represented by the appearance of these three newly emerged absorption bands. Especially in the cases of NaCl and SDS fibrils, the absorption band at $\sim 25 \times 10^3$ cm^{-1} also seemed partly responsible for the color formation, given that its intensity increased after the iodine staining. However, the color differences between amyloid polymorphs were well described with the newly emerged absorption bands. They showed significantly higher peak wavenumbers in NaCl fibrils compared to those of SDS and no-salt fibrils, and in SDS fibrils and no-salt fibrils, the relative intensities of the three bands were different while the peak positions were almost similar (filled bands in Figure 3A-C).

Figure 3. Deconvolution of UV-Vis absorption spectra of iodine-stained NaCl fibrils (A), SDS fibrils (B), and no-salt fibrils (C). In all panels, black and red solid lines represent the experimental and reproduced spectra, respectively, and the experimental spectra are the same as those shown in Figure 2B. Six Gaussian peaks represented with dashed red lines or filled areas correspond to absorption components obtained by the deconvolution using eq. 1, and dashed black lines represent baselines derived from scattering of the sample fibrils. The wavenumber of each absorption components is labeled.

Tracking structural propagation of amyloid fibrils by iodine staining. To verify whether the intrinsic iodine colors have potency to indicate the propagation of amyloid fibril structures, we tracked time course of seed-dependent elongation by iodine staining. In this experiment and thereafter, NaCl and SDS fibrils, which exhibited markedly distinct characteristics in terms of absorption band components, were subjected to the analyses (Figure 4). As seeds, the parent fibrils were used. When the color formation by iodine was examined by sampling the reaction mixture at different time points, absorption intensity in visible region gradually increased both in the NaCl and SDS fibrils as time advanced (Figure 4A,C). Spectral deconvolution was successfully achieved by assuming six absorption band components with constant wavenumbers (Figure S3). When the areas of the three absorption bands in the long wavelength (i.e., low wavenumber) region were plotted against reaction time, all of them increased synchronously with the time-dependent increase of ThT fluorescence intensity (Figure 4B,D), suggesting that the color developed in response to fibril elongation without changing its tone.

Figure 4. Tracking seed-dependent fibril growth by iodine staining. (A-D) Time dependent change in iodine color for NaCl (A,B) and SDS fibrils (C,D). In panels A and C, UV-Vis spectra of NaCl (A) and SDS fibrils (C) stained with iodine monitored at different time are shown. The conditions of iodine staining were 0.25 mg/ml insulin, 0.3 mM KI and 0.04 mM I_2 in 25 mM HCl. In panels B and D, areas of three absorption components of NaCl fibrils (B) and SDS fibrils (D) in lower wavenumber region are plotted against reaction time. For details about spectral deconvolution for the determination of these absorption bands, see Figure S3. ThT fluorescence intensity which was measured concurrently is also plotted for comparison.

Characterization of polyiodide species responsible for visible color. According to studies of iodine-doped compounds, the molecular mechanism of visible color formation by iodine is due to the formation of polyiodide ions [41,42]. In the case of iodine-doped amyloid fibrils, the formation of pentaiodide ions and several other oligoiodide ions was suggested for the first time by Dzwolak [9]. To identify polyiodide species formed on the amyloid fibrils analyzed in the present work, resonance Raman spectra were measured. Raman spectroscopy is a useful technique for detecting subunits of polyiodide chains in complexes with amylose [21,43,44] and

other organic compounds [45-48]. Here, two excitation wavelengths 514.5 nm (19436 cm^{-1}) and 785 nm (12739 cm^{-1}) were used for excitation; two out of three absorption bands (red and green bands in Figure 3A-C) are supposed to be enhanced with the 514.5 nm laser, and the remaining band (blue bands in Figure 3A-C) is supposed to be enhanced with the 785 nm laser.

As a result, resonance Raman effects were observed for both NaCl and SDS fibrils, and characteristic Raman spectra with two strong peaks at around 110 cm^{-1} and 160 cm^{-1} were observed at either excitation at 514.5 nm or 785 nm (Figure 5). Based on a previous Raman study of the crystalline compounds containing triiodide or pentaiodide ions [45], the 110 cm^{-1} and 160 cm^{-1} peaks were assigned to I_3^- and I_5^-, respectively. The present observation therefore suggests that the fundamental building blocks of polyiodide ions are dominantly I_3^- and I_5^-. Unstained fibrils showed no significant peaks in the same way as those of iodine solution without fibrils, indicating that the peaks are derived from iodine species bound to the fibrils. When two spectra excited at different wavelengths were compared for each type of fibrils (i.e., Figure 5A,C or 5B,D), the positions the I_3^- and I_5^- peaks were different, and their intensity ratio was reversed. This observation suggests that the conformations and the composition ratio of the I_3^- and I_5^- subunits constituting the polyiodide ions vary among the absorption bands. Especially when excited at 514.5 nm, two weak bands were additionally observed at ~ 220 cm^{-1} and ~ 320 cm^{-1}, which were assumed to be overtone bands of the I_3^- and I_5^- bands, respectively. A very weak combination band of the I_3^- and I_5^- bands was also found at around 270 cm^{-1} for SDS fibrils, whereas it was undetectable for NaCl fibrils, probably due to low intensity of the 160 cm^{-1} band. When excited at 785 nm, an overtone band of the I_5^- band as well as a combination band were observed both for NaCl and SDS fibrils. When the spectra of NaCl and SDS fibrils were compared at the same excitation wavelength (i.e., Figure 5A,B or 5C,D), the intensity ratio of the I_3^- and I_5^- peaks was significantly different at both excitation wavelengths, and the positions of the peaks were also slightly different between NaCl and SDS fibrils. This

Figure 5. Resonance Raman spectra of iodine-stained insulin amyloid fibrils with excitation of 514.5 nm (A,B) and 785 nm(C,D). The spectra of NaCl fibrils (A,C) and SDS fibrils (B,D) are shown. Samples containing 0.5 mg/ml amyloid fibrils, 0.6 mM KI, and 0.08 mM I_2 in 25 mM HCl were subjected to the measurements. Solid and dotted lines represent spectra of iodine-stained fibrils and unstained fibrils measured as a control, respectively, and the positions of peaks are indicated by black dashed lines.

observation implies that different polyiodide ions are formed depending on amyloid polymorphs.

Titration of iodine to insulin amyloid fibrils. We further performed titration experiments to identify iodine concentration dependence on the absorbance of each spectral band. With stepwise addition of the iodine solution to amyloid fibrils, gradual intensification of the absorption spectrum was observed both for NaCl and SDS fibrils. The absorption in the low wavenumber region finally saturated after the addition of excess amount of iodine (Figure 6A,B). Interestingly, it seemed that absorption at lower wavenumbers grew slightly later compared to that at higher wavenumbers, and indeed, distinct shapes of the titration plots were

Figure 6. Titration of insulin fibrils with iodine solution. 0.5 mg/ml insulin fibrils in 25 mM HCl and iodine solution containing 18 mM KI and 2.4 mM I_2 in 25 mM HCl were used as an analyte and a titrant, respectively. (A,B) Examples of absorption spectra of NaCl (A) and SDS fibrils (B) measured at different concentrations of iodine solution. (C,D) Iodine concentration dependence of area of the three absorption bands of iodine-stained NaCl or SDS appearing in the lower wavenumber region. In these plots, the concentration of I_2 in the titrant iodine solution is represented as the scale of horizontal axes. For details about spectral deconvolution for the determination of these absorption bands, see Figure S4. Black lines represent fitted curves using eq. 2 and parameters obtained from the fitting are represented inside the panels.

observed when spectral deconvolution was performed by using eq. 1 and the areas ($A(c)$) of the three absorption bands in the low wavenumber region (the bands with filled area in Figure S4) was plotted against iodine concentration c. The titration plots showed a cooperative properties and as a result of curve fittings with Hill equation [49],

$$A(c) = \frac{A_{max}[c]^n}{K_A + [c]^n} \qquad (2)$$

where A_{max}, K_A, and n represent band area attained after saturation, iodine concentration at which half of the binding sites are occupied, and Hill coefficient, respectively, resultant curves exhibited different parameters in their binding affinity and cooperativity among the three absorption bands both in NaCl (Figure 6C) and SDS fibrils (Figure 6D). This result suggests that there are at least three different iodine-binding sites. Furthermore, all plots exhibited $n > 1$ (Figure 6C,D), from which some cooperative mechanisms were suggested in the iodine binding to amyloid fibrils. It is deduced that small iodide species, such as I_3^- and I^- with negligible absorption in visible region initially bind and then facilitates subsequent iodine binding to form polyiodide ions showing visible color.

Investigation of the orientation of polyiodides on the surface of amyloid fibrils.
It is often revealed that polyiodide chains formed by complexation of iodine/iodide with various compounds lie in an oriented manner [20,48,50]. To gain insights into whether the polyiodides bound to amyloid fibrils have some specific orientation, a polarization property of the iodine-stained fibrils was investigated. Linear polyiodides ions have transition moment in a direction parallel to their long axes [45], and thus the aligned polyiodides ions are estimated to absorb polarized light, as seen in unidirectionally stretched films of iodine-doped polyvinyl alcohol [51]. In this analysis, amyloid spherulites, a spherical hierarchical structure composed of radially oriented amyloid fibrils [33], were selected as a good model for evaluating the orientation of fibril-bound iodine molecules. Amyloid spherulites of insulin were formed, treated with iodine

solution, and then observed by a microscope equipped with an analyzer to assess whether the transmitted light of the iodine-stained amyloid spherulites show a polarized property.

As a result, the transmission microscope images of the iodine-stained amyloid spherulites showed two cone-shaped dark sections along the analyzer axis, and their direction changed with response to the rotation of the analyzer (Figure 7). In contrast, no dark section was observed for unstained spherulites (Figure S5), verifying that the polarized nature of the iodine-stained spherulites was derived from polyiodide. Given the radial orientation of fibril axes, the direction of the dark sections and that of the analyzer axis coincided well, which suggests that polyiodide chains are oriented along the fibril axis.

Figure 7. Anisotropic property of polyiodides on the spherulite. In each image, the direction of the analyzer slit is represented with a double-headed arrow in the upper right. Scale bars represent 10 μm. The images of an unstained spherulite recorded by the same procedure are shown in Figure S5 as a control.

DISCUSSION

Amyloid structure-dependent coloration by iodine staining. We have investigated color properties of iodine-stained amyloid fibrils of human insulin mainly by using UV-Vis absorbance spectroscopy. When iodine molecules bound on the surface of the insulin fibrils, they showed clear coloration like iodine-starch reactions. An interesting feature of the iodine staining of amyloid fibrils was that different colors were observed among the three types of insulin amyloid fibrils (i.e., NaCl, SDS, and no-salt fibrils) upon the addition of iodine solution. The differences in color tone were spectroscopically demonstrated in visible region, and

furthermore, the colors formed by iodine staining could track the propagation of amyloid fibril structures by seed-dependent growth reactions. The present observation suggests that iodine staining can serve as a new probe indicating structural polymorphs of amyloid fibrils.

There are several chemical dyes that have been widely used for detection and tracking of amyloid fibril formation. The most widely used one is ThT, which exhibits enhanced fluorescence upon binding to amyloid fibrils. ThT has been used for many assays as well as extended applications such as microscopic observation of amyloid fibril growth and diagnosis of fibrils in tissue sections [32,52]. Several ThT derivatives with high affinity toward amyloid fibrils have also been designed, and recently, radioactively labeled Pittsburgh compound-B is widely used for non-invasive in vivo positron emission tomography imaging for diagnosis of AD [53,54]. Congo red, which shows characteristic apple-green birefringence under polarized light, is the most histological amyloid dye used for the diagnosis of amyloidosis for several decades [55,56]. 8-Anilino-1-naphthalenesulfonic acid has also been used for detecting fibrils and prefibrillar intermediates, although specificity is not high [57]. These amyloid dyes are very good at detecting the presence and quantifying the amount of amyloid fibrils. However, they are not so sensitive to differences in structure of amyloid fibrils and fail to probe amyloid polymorphism in many cases. Although NaCl, SDS, and no-salt fibrils showed different peak wavelengths in ThT fluorescence spectrum (Figure 1), ThT fluorescence is typically considered to show little change in peak wavelength and only fluorescence intensity can be used for detecting amyloid polymorphs [58]. Compared to these conventionally used dyes, it is expected that the color formed by iodine staining is highly sensitive to fibril structures, and in the case of the insulin fibril samples used in this study, clear differences in color, which were obvious even with the naked eye, were observed. The present results have indicated that iodine staining can be used for identifying polymorphs of amyloid fibrils.

Fundamental scheme of iodine staining of amyloid fibrils. The coloration of amyloid fibrils by iodine staining is thought to be derived from polymerization of iodine upon binding to amyloid fibrils [9]. In water solution without any host compounds, polyiodide ions larger than I_3^- are absent in an iodine solution [59]. However, a wide range of polyiodides structures have been observed in a supramolecular crystalline compounds and polymers [22-26]. As for amyloid fibrils, it is predicted that, because the cross-β structure of amyloid fibrils has high periodicity along the fibril axis, they can act as a scaffold of iodine complexation in a similar manner to other compounds. Indeed, resonance Raman spectroscopy identified the presence of I_3^- and I_5^- species as fibril-bound forms of iodine (Figure 5). Considering that I_3^- itself does not absorb visible light so strongly [39,40], it is predicted that I_5^- and other polyiodide ions composed of I_3^- and I_5^- as fundamental building blocks, $(I_3^-)_n$ and $(I_5^-)_n$, and possibly some other types of polyiodides composed of both of I_3^- and I_5^-, $(I_3^-)_m(I_5^-)_n$, serve as the molecular basis of color formation.

Interestingly, the resonance Raman spectra excited at 785 nm were similar to the previously reported iodine-amylose pattern in terms of positions and relative intensities of fundamental bands as well as their overtone and combination bands [21,44]. According to the crystal structure of iodine-doped α-cyclodextrin, a structural model that resembles the iodine-amylose complex, it is proposed that a chain-like I_5^- ion is encapsulated inside a channel-like cavity of the helical structure of amylose [26]. It is therefore suggested that linearly aligned $(I_5^-)_n$ polyiodides contribute to the formation of the absorption band at the lowest wavenumber. On the other hand, the band at ~110 cm^{-1} was rather stronger than that at ~160 cm^{-1} when excited at 514.5 nm (Figure 5A,B), implying that other types of polyiodides containing I_3^-, possibly $(I_3^-)_n$, might also be involved in the formation of reddish color. Alternatively, the increased Raman band at ~110 cm^{-1} in the spectra may result from impairment of linear array of $(I_5^-)_n$ chains, in light of a previous report on the analysis of $(I_5^-)_n$ chains with different levels of alignment [45].

Figure 8 is a proposed model of the binding of polyiodides on amyloid fibrils. The crystal structures of several iodine-doped compounds, such as α-cyclodextrin [20] and an inorganic-organic compound $\{(C_4H_{12}N_2)_2[Cu^I I_4](I_2)\}_n$ [60], have already been revealed at atomic level. According to these structures, iodine molecules are encapsulated or buried in gap structures of the host molecules. In contrast, Dzwolak revealed that iodine species bound to bovine insulin fibrils are easily reduced by ascorbic acids [9], suggesting that iodine molecules are located on the solvent-accessible surface of amyloid fibrils. Given that anisotropic coordination of polyiodides was suggested by the observation of iodine-stained spherulites by polarization microscopy, it is estimated that line shaped polyiodides are formed and stabilized on the grooves of the pleated cross-β structure running in parallel to the long axis of fibrils (model I in Figure 8). The space between two side chains of the β-strand is approximately 7 Å, which is suitable size for iodine atoms of 4 Å in diameter to get stuck, and it is thus conceived to act as a scaffold to form line shaped polyiodide species. In addition to the surface of cross-β structure, the typical distance between two laminated β-sheet layers is approximately 10 Å. The edge of β-sheets could be another candidate for the binding site of polyiodides in a similar manner to the case of ThT binding [61] (model II in Figure 8). Twist grooves formed by intertwining protofilaments may also play a role as iodine binding sites suitable for the fixation of polyiodides. Further investigation is needed for more detailed specification of the binding sites.

Molecular origin of different colors shown by amyloid polymorphs. In the system of insulin amyloid fibrils, spectral deconvolution of the UV-Vis spectra has clarified that the color formed by iodine staining of amyloid fibrils is represented as the sum of three major absorption bands in visible region. The titration of iodine solution to amyloid fibrils revealed that each of these absorption bands had its own specific binding constant for iodine binding

Figure 8. Schematic illustration for the binding of the polyiodides on the surface of amyloid fibrils. In this figure, I_5^- and $(I_3^-)_2$ are shown as tentative examples of polyiodides, and two possible candidates for the binding site of polyiodides, i.e., the grooves of the pleated cross-β structure (model I) and the edge of β-sheets (model II), are represented.

(Figure 6). This result indicates that insulin amyloid fibrils basically have three independent iodine-binding sites, to each of which a specific and different type of polyiodide ion binds; while the polyiodide ions are commonly composed of I_3^- and I_5^- building blocks, the conformations of the building blocks as well as their composition are different, as deduced from the result of resonance Raman spectra obtained at two different excitation wavelengths (Figure 5). Additionally, degree of polymerization may also be different depending on the binding sites, in view of previous studies on amylose with different degree of polymerization in which absorption band shifted effectively to longer wavelength as the chain length of polyiodides increased [25].

As shown in Figure 3, differences in the peak positions and intensity ratio of the three absorption bands in visible region were the spectroscopic entities underlying the color

difference in iodine staining among polymorphs. The characteristics of the three absorption bands seem to change sensitively depending on structural properties of fibril surface. It is anticipated that characteristic protein topology of amyloid polymorphs on cross-β structure and/or at its edge parts determines the structure and degree of polymerization of polyiodides on the binding sites, and eventually, leads to different coloration. Furthermore, when the absorption spectra of SDS and no-salt fibrils were compared, the wavenumbers of the three constituent absorption bands were similar to those of SDS fibrils; however, the relative ratio of their absorbance intensities were markedly different (Figure 3). This observation may suggest that differences in factors related to absorption intensity, i.e., the number of binding site and their binding constant of polyiodides, are also available for detecting polymorphic structures of amyloid fibrils, even though iodine binding sites are very similar.

Conclusion and future perspectives of iodine staining as a tool for distinguishing amyloid polymorphism. Through this work, it was clarified that coloration by iodine staining has a promising characteristics in identifying polymorphism of amyloid fibrils. Considering that the colors successfully reported the differences in structure of insulin amyloid fibrils despite that their secondary structures were very similar (Figure 1-3), it is expected that iodine staining works as a sensitive probe for amyloid polymorphs by discriminating their surface structures. The iodine color also successfully tracked the propagation process of amyloid structures (Figure 4). Although iodine staining alone cannot function as a technical strategy for clarifying amyloid structures, its colors can act as a fingerprint of amyloid polymorphs. If color information on a particular type of amyloid fibrils of interest is investigated beforehand, its quick identification will be achieved by matching the color features. Moreover, given the recent situation that a growing number of the structures of amyloid fibrils are determined by cryo-EM and ssNMR spectroscopy, valuable insights into the

relationships between the surface topology of amyloid fibrils and the corresponding color property by iodine staining are obtained practically by analyzing the color properties of amyloid fibrils with known structures. This will provide perspective on clarifying the molecular mechanism of iodine staining, and consequently, will boost the elucidation of molecular details for amyloid polymorphism underlying amyloid diseases.

Recently, luminescent conjugated polythiophenes and oligothiophenes have been developed as a molecular probe of amyloid fibrils [62] and very recently, succeeded in differentiate pathological amyloid subtypes of Aβ in AD brains [63]. This observation has revealed the presence of structurally distinct conformations in the affected area of amyloid diseases, arguing the importance of characterizing amyloid polymorphs in biological samples. Although iodine staining is based on absorption and is inferior in terms of signal intensity compared to fluorescent probes, it has potency of being applied to all of the amyloidogenic proteins, and covering a wide range of samples that have not been analyzed so far because of problems with amount that can be supplied for the analysis as well as inhomogeneity. Furthermore, in light of the fact that the iodine staining of amyloid fibrils was firstly discovered in the field of anatomy and is still used in part for diagnosis of amyloidoses, the present work also suggests that it will be worth revisiting the usefulness of the iodine staining of pathological tissues for evaluating the conditions and progress of diseases on the basis of amyloid polymorphism. Further data accumulation connecting the iodine colors and fibril structures will open up new perspectives of the usage of iodine staining in various field of amyloid researches.

SUPPORTING INFORMATION

Figure S1. UV-vis spectra showing negligible interaction between iodine molecules and native insulin.

Figure S2. UV-vis spectra showing negligible effects of the presence of salts on iodine staining.

Figure S3. Representative results of curve fitting of the UV-vis absorption spectra in the seeding reaction.

Figure S4. Representative results of curve fitting of the UV-vis absorption spectra in the iodine titration experiment.

Figure S5. Images of an unstained spherulite observed by polarization microscopy.

ACKNOWLEDGMENTS

We thank Tomomi Kozu (Renishaw K.K.) for help of resonance Raman measurements and Prof. Masahide Yamamoto (Kyoto University) for valuable comments on iodine-polymer complexes. This work was supported by JSPS Core-to-Core Program, A. Advanced Research Networks. This work was funded by JSPS KAKENHI Grant Numbers JP16H04778, JP16H00772, and JP17H06352.

REFERENCES

(1)   Knowles, T. P.; Vendruscolo, M.; Dobson, C. M. The amyloid state and its association with protein misfolding diseases. *Nat. Rev. Mol. Cell Biol.* **2014**, *15*, 384-396.

(2)   Pedersen, J. S.; Andersen, C. B.; Otzen, D. E. Amyloid structure-one but not the same: The many levels of fibrillar polymorphism. *FEBS J.* **2010**, *277*, 4591-4601.

(3)   Tycko, R. Physical and structural basis for polymorphism in amyloid fibrils. *Protein Sci.* **2014**, *23*, 1528-1539.

(4)   Gallardo, R.; Ranson, N. A.; Radford, S. E. Amyloid structures: Much more than just a cross-β fold. *Curr. Opin. Struct. Biol.* **2019**, *60*, 7-16.

(5)   Seuring, C.; Verasdonck, J.; Ringler, P.; Cadalbert, R.; Stahlberg, H.; Bockmann, A.; Meier, B. H.; Riek, R. Amyloid fibril polymorphism: Almost identical on the atomic level, mesoscopically very different. *J. Phys. Chem. B* **2017**, *121*, 1783-1792.

(6)   Close, W.; Neumann, M.; Schmidt, A.; Hora, M.; Annamalai, K.; Schmidt, M.; Reif, B.; Schmidt, V.; Grigorieff, N.; Fandrich, M. Physical basis of amyloid fibril polymorphism. *Nat. Commun.* **2018**, *9*, 699.

(7)   Nelson, R.; Sawaya, M. R.; Balbirnie, M.; Madsen, A. O.; Riekel, C.; Grothe, R.; Eisenberg, D. Structure of the cross-β spine of amyloid-like fibrils. *Nature* **2005**, *435*, 773-778.

(8)   Petkova, A. T.; Leapman, R. D.; Guo, Z.; Yau, W. M.; Mattson, M. P.; Tycko, R. Self-propagating, molecular-level polymorphism in Alzheimer's β-amyloid fibrils. *Science* **2005**, *307*, 262-265.

(9)   Dzwolak, W. Insulin amyloid fibrils form an inclusion complex with molecular iodine: A misfolded protein as a nanoscale scaffold. *Biochemistry* **2007**, *46*, 1568-1572.

(10)   Xiao, Y.; Ma, B.; McElheny, D.; Parthasarathy, S.; Long, F.; Hoshi, M.; Nussinov, R.; Ishii, Y. Aβ(1-42) fibril structure illuminates self-recognition and replication of amyloid in Alzheimer's disease. *Nat. Struct. Mol. Biol.* **2015**, *22*, 499-505.

(11)   Gremer, L.; Scholzel, D.; Schenk, C.; Reinartz, E.; Labahn, J.; Ravelli, R. B. G.; Tusche, M.; Lopez-Iglesias, C.; Hoyer, W.; Heise, H.; Willbold, D.; Schroder, G. F. Fibril structure of amyloid-β(1-42) by cryo-electron microscopy. *Science* **2017**, *358*, 116-119.

(12)   Fitzpatrick, A. W. P.; Falcon, B.; He, S.; Murzin, A. G.; Murshudov, G.; Garringer, H. J.; Crowther, R. A.; Ghetti, B.; Goedert, M.; Scheres, S. H. W. Cryo-EM structures of tau filaments from Alzheimer's disease. *Nature* **2017**, *547*, 185-190.

(13)   Falcon, B.; Zhang, W.; Murzin, A. G.; Murshudov, G.; Garringer, H. J.; Vidal, R.; Crowther, R. A.; Ghetti, B.; Scheres, S. H. W.; Goedert, M. Structures of filaments from Pick's disease reveal a novel tau protein fold. *Nature* **2018**, *561*, 137-140.

(14)   Tanaka, M.; Chien, P.; Naber, N.; Cooke, R.; Weissman, J. S. Conformational variations in an infectious protein determine prion strain differences. *Nature* **2004**, *428*, 323-328.

(15)   Pedersen, J. S.; Dikov, D.; Flink, J. L.; Hjuler, H. A.; Christiansen, G.; Otzen, D. E. The changing face of glucagon fibrillation: Structural polymorphism and conformational imprinting. *J. Mol. Biol.* **2006**, *355*, 501-523.

(16)   Chatani, E.; Yagi, H.; Naiki, H.; Goto, Y. Polymorphism of β2-microglobulin amyloid fibrils manifested by ultrasonication-enhanced fibril formation in trifluoroethanol. *J. Biol. Chem.* **2012**, *287*, 22827-22837.

(17)   Jones, E. M.; Surewicz, W. K. Fibril conformation as the basis of species- and strain-dependent seeding specificity of mammalian prion amyloids. *Cell* **2005**, *121*, 63-72.

(18)   Dzwolak, W.; Loksztejn, A.; Galinska-Rakoczy, A.; Adachi, R.; Goto, Y.; Rupnicki, L. Conformational indeterminism in protein misfolding: Chiral amplification on amyloidogenic pathway of insulin. *J. Am. Chem. Soc.* **2007**, *129*, 7517-7522.

(19)   Lu, J. X.; Qiang, W.; Yau, W. M.; Schwieters, C. D.; Meredith, S. C.; Tycko, R. Molecular structure of beta-amyloid fibrils in Alzheimer's disease brain tissue. *Cell* **2013**, *154*,

1257-1268.

(20)   Noltemeyer, M.; Saenger, W. X-Ray Studies of linear polyiodide chains in α-cyclodextrin channels and a model for starch-iodine complex. *Nature* **1976**, *259*, 629-632.

(21)   Teitelbaum, R. C.; Ruby, S. L.; Marks, T. J. A resonance Raman/iodine Mossbauer investigation of the starch-iodine structure. Aqueous-solution and iodine vapor preparations. *J. Am. Chem. Soc.* **1980**, *102*, 3322-3328.

(22)   Matsuyama, T.; Sakai, H.; Yamaoka, H.; Maeda, Y.; Shirakawa, H. Mossbauer Spectroscopic study of iodine-doped polyacetylene. *J. Phys Soc. Jpn.* **1983**, *52*, 2238-2245.

(23)   Murthy, N. S.; Khanna, Y. P. Transformations between polyiodide structures in a nylon-6 matrix. *Chem. Mater.* **1993**, *5*, 672-677.

(24)   Yokoyama, T.; Kaneyuki, K.; Sato, H.; Hamamatsu, H.; Ohta, T. Structure, composition, and vibrational property of iodine-doped polyvinyl-alcohol studied by temperature-dependent I K-edge extended X-ray-absorption fine-structure. *Bull. Chem. Soc. Japan* **1995**, *68*, 469-475.

(25)   Ono, S.; Tsuchihashi, S.; Kuge, T. On the starch-iodine complex. *J. Am. Chem. Soc.* **1953**, *75*, 3601-3602.

(26)   Noltemeyer, M.; Saenger, W. Structural chemistry of linear α-cyclodextrin-polyiodide complexes. X-ray crystal-structures of (α-cyclodextrin)$_2$·LiI$_3$·I$_2$·8H$_2$O and (α-cyclodextrin)$_2$·Cd$_{0.5}$·I$_5$·27H$_2$O. Models for the blue amylose-iodine complex. *J. Am. Chem. Soc.* **1980**, *102*, 2710-2722.

(27)   Virchow, R. Nouvelles observation sur la substance animale analogou a la cellulose vegetale. *C.R. Acad. Sci. (Paris)* **1853**, *37*, 860-861.

(28)   Sipe, J. D.; Cohen, A. S. Review: History of the amyloid fibril. *J Struct. Biol.* **2000**, *130*, 88-98.

(29)   Buxbaum, J. N.; Linke, R. P. A molecular history of the amyloidoses. *J. Mol. Biol.* **2012**, *421*, 142-159.

(30)   Groenning, M.; Frokjaer, S.; Vestergaard, B. Formation mechanism of insulin fibrils and structural aspects of the insulin fibrillation process. *Curr. Protein Pept. Sci.* **2009**, *10*, 509-528.

(31)   Nielsen, L.; Frokjaer, S.; Carpenter, J. F.; Brange, J. Studies of the structure of insulin fibrils by Fourier transform infrared (FTIR) spectroscopy and electron microscopy. *J Pharm. Sci.* **2001**, *90*, 29-37.

(32)   Naiki, H.; Higuchi, K.; Hosokawa, M.; Takeda, T. Fluorometric determination of amyloid fibrils in vitro using the fluorescent dye, thioflavin T. *Anal. Biochem.* **1989**, *177*, 244-249.

(33)   Rogers, S. S.; Krebs, M. R.; Bromley, E. H.; van der Linden, E.; Donald, A. M. Optical microscopy of growing insulin amyloid spherulites on surfaces in vitro. *Biophys. J.* **2006**, *90*, 1043-1054.

(34)   Waugh, D. F. A Fibrous modification of insulin. I. The heat precipitate of insulin. *J. Am. Chem. Soc.* **1946**, *68*, 247-250.

(35)   Brange, J.; Andersen, L.; Laursen, E. D.; Meyn, G.; Rasmussen, E. Toward understanding insulin fibrillation. *J. Pharm. Sci.* **1997**, *86*, 517-525.

(36)   Whittingham, J. L.; Scott, D. J.; Chance, K.; Wilson, A.; Finch, J.; Brange, J.; Dodson, G. G. Insulin at pH 2: Structural analysis of the conditions promoting insulin fibre formation. *J. Mol. Biol.* **2002**, *318*, 479-490.

(37)   Groenning, M. Binding mode of Thioflavin T and other molecular probes in the context of amyloid fibrils-current status. *J. Chem. Biol.* **2010**, *3*, 1-18.

(38)   Bandekar, J. Amide modes and protein conformation. *Biochim. Biophys. Acta* **1992**, *1120*, 123-143.

(39)   Awtrey, A. D.; Connick, R. E. The Absorption Spectra of I$_2^-$, I$_3^-$, I$^-$, IO$_3^-$, S$_4$O$_6^=$ and

$S_2O_3^=$. Heat of the reaction $I_3^- = I_2^- + I^-$. *J. Am. Chem. Soc.* **1951**, *73*, 1842-1843.

(40)   Pursell, J. L.; Pursell, C. J. Host-guest inclusion complexation of α-cyclodextrin and triiodide examined using UV-Vis spectrophotometry. *J. Phys. Chem. A* **2016**, *120*, 2144-2149.

(41)   Svensson, P. H.; Kloo, L. Synthesis, structure, and bonding in polyiodide and metal iodide-iodine systems. *Chem. Rev.* **2003**, *103*, 1649-1684.

(42)   Kupper, F. C.; Feiters, M. C.; Olofsson, B.; Kaiho, T.; Yanagida, S.; Zimmermann, M. B.; Carpenter, L. J.; Luther, G. W., 3rd; Lu, Z.; Jonsson, M.; Kloo, L. Commemorating two centuries of iodine research: An interdisciplinary overview of current research. *Angew. Chem. Int. Ed. Engl.* **2011**, *50*, 11598-11620.

(43)   Yu, X. C.; Houtman, C.; Atalla, R. H. The complex of amylose and iodine. *Carbohydr. Res.* **1996**, *292*, 129-141.

(44)   Teitelbaum, R. C.; Ruby, S. L.; Marks, T. J. On the structure of starch-iodine. *J. Am. Chem. Soc.* **1978**, *100*, 3215-3217.

(45)   Mizuno, M.; Tanaka, J.; Harada, I. Electronic-spectra and structures of polyiodide chain complexes. *J. Phys. Chem.* **1981**, *85*, 1789-1794.

(46)   Burzynski, R.; Prasad, P. N.; Murthy, N. S. Structure of the iodine columns in iodinated nylon-6. *J. Polym. Sci. Polym. Phys.* **1986**, *24*, 133-141.

(47)   Moulay, S. Molecular iodine/polymer complex. *J. Polym. Eng.* **2013**, *33*, 389-443.

(48)   Madhu, S.; Evans, H. A.; Doan-Nguyen, V. V.; Labram, J. G.; Wu, G.; Chabinyc, M. L.; Seshadri, R.; Wudl, F. Infinite polyiodide chains in the pyrroloperylene-iodine complex: Insights into the starch-iodine and perylene-iodine complexes. *Angew. Chem. Int. Ed. Engl.* **2016**, *55*, 8032-8035.

(49)   O'Riordan, J. F.; Goldstick, T. K.; Ditzel, J.; Ernest, J. T. Characterization of oxygen-hemoglobin equilibrium curves using nonlinear regression of the Hill equation: Parameter values for normal human adults. *Adv. Exp. Med. Biol.* **1983**, *159*, 435-444.

(50)   Reddy, J. M.; Knox, K.; Robbin, M. B. Crystal structure of $HI_3 \cdot 2C_6H_5CONH_2$: A model of the starch-iodine complex. *J. Chem. Phys.* **1964**, *40*, 1082.

(51)   Seto, M.; Maeda, Y.; Matsuyama, T.; Yamaoka, H.; Sakai, H. Light polarization in iodine doped polyvinyl-alcohol films. *Hyperfine Interact.* **1991**, *68*, 221-224.

(52)   Vassar, P. S.; Culling, C. F. Fluorescent stains, with special reference to amyloid and connective tissues. *Arch. Pathol.* **1959**, *68*, 487-498.

(53)   Klunk, W. E.; Engler, H.; Nordberg, A.; Wang, Y.; Blomqvist, G.; Holt, D. P.; Bergstrom, M.; Savitcheva, I.; Huang, G. F.; Estrada, S.; Ausen, B.; Debnath, M. L.; Barletta, J.; Price, J. C.; Sandell, J.; Lopresti, B. J.; Wall, A.; Koivisto, P.; Antoni, G.; Mathis, C. A.; Langstrom, B. Imaging brain amyloid in Alzheimer's disease with Pittsburgh Compound-B. *Ann. Neurol.* **2004**, *55*, 306-319.

(54)   McNamee, R. L.; Yee, S. H.; Price, J. C.; Klunk, W. E.; Rosario, B.; Weissfeld, L.; Ziolko, S.; Berginc, M.; Lopresti, B.; Dekosky, S.; Mathis, C. A. Consideration of optimal time window for Pittsburgh compound B PET summed uptake measurements. *J. Nucl. Med.* **2009**, *50*, 348-355.

(55)   Bennhold, H. Specific staining of amyloid by Congo Red. *Munch. Med. Wochenschr.* **1922**, *69*, 1537-1538.

(56)   Yakupova, E. I.; Bobyleva, L. G.; Vikhlyantsev, I. M.; Bobylev, A. G. Congo Red and amyloids: History and relationship. *Biosci. Rep.* **2019**, *39* BSR20181415.

(57)   Lindgren, M.; Sorgjerd, K.; Hammarstrom, P. Detection and characterization of aggregates, prefibrillar amyloidogenic oligomers, and protofibrils using fluorescence spectroscopy. *Biophys. J.* **2005**, *88*, 4200-4212.

(58)   Singh, P. K.; Kumbhakar, M.; Pal, H.; Nath, S. Ultrafast bond twisting dynamics in amyloid fibril sensor. *J. Phys. Chem. B* **2010**, *114*, 2541-2546.

(59)   Ramette, R. W.; Sandford, R. W. Thermodynamics of iodine solubility and triiodide

ion formation in water and in deuterium oxide. *J. Am. Chem. Soc.* **1965**, *87*, 5001-5005.

(60)   Redel, E.; Rohr, C.; Janiak, C. An inorganic starch-iodine model: The inorganic-organic hybrid compound $\{(C_4H_{12}N_2)_2[Cu^II_4](I_2^-)\}_n$. *Chem. Commun.* **2009**, 2103-2105.

(61)   Morii, H.; Saiki, M.; Konakahara, T.; Ishimura, M. Peripheral region for core cross-beta plays important role in amyloidogenicity. *Biochem. Biophys. Res. Commun.* **2006**, *342*, 808-816.

(62)   Fandrich, M.; Nystrom, S.; Nilsson, K. P. R.; Bockmann, A.; LeVine, H., 3rd; Hammarstrom, P. Amyloid fibril polymorphism: A challenge for molecular imaging and therapy. *J. Intern. Med.* **2018**, *283*, 218-237.

(63)   Rasmussen, J.; Mahler, J.; Beschorner, N.; Kaeser, S. A.; Hasler, L. M.; Baumann, F.; Nystrom, S.; Portelius, E.; Blennow, K.; Lashley, T.; Fox, N. C.; Sepulveda-Falla, D.; Glatzel, M.; Oblak, A. L.; Ghetti, B.; Nilsson, K. P. R.; Hammarstrom, P.; Staufenbiel, M.; Walker, L. C.; Jucker, M. Amyloid polymorphisms constitute distinct clouds of conformational variants in different etiological subtypes of Alzheimer's disease. *Proc. Natl. Acad. Sci. U.S.A.* **2017**, *114*, 13018-13023.