Neural processing genes are linked to divergence in visual mate preference

in sympatric *Heliconius* butterflies

Matteo Rossi[a,b], Timothy J. Thurman[b,c], Alexander E. Hausmann[a], Stephen H.

Montgomery[d], Riccardo Papa[b,e,f], Chris D. Jiggins[g] , W. Owen McMillan[b] & Richard M.

Merrill[a,b]

[a] Division of Evolutionary Biology, Ludwig Maximilian University of Munich, Großhaderner Str. 2, 82152

Planegg-Martinsried, Germany

[b] Smithsonian Tropical Research Institute, Gamboa, 0843-03092, Panama

[c] Division of Biological Sciences, University of Montana, 32 Campus Dr., Missoula, MT 59812, US

[d] School of Biological Sciences, University of Bristol, 24 Tyndall Ave, Bristol BS8 1TQ, UK

[e] Department of Biology, University of Puerto Rico, Rio Piedras, San Juan 00925, Puerto Rico

[f] Molecular Sciences and Research Center, University of Puerto Rico, San Juan 00926, Puerto Rico

[g] Department of Zoology, University of Cambridge, Downing Street, Cambridge CB2 3EJ, UK

Corresponding author: merrill@bio.lmu.de

Short title: Genetics of visual mate preference in *Heliconius*

Abstract

Divergence in mating behaviors plays a major role during speciation, but we know little about the genetic mechanisms underlying the evolution of such traits. Reproductive isolation between the Neotropical butterflies *Heliconius melpomene* and *H. cydno* depends on shifts in mimetic warning patterns, which also act as mating cues. Preferences for conspecific warning patterns have a strong genetic component, but unlike the warning pattern cues, the exact genes responsible remain unknown. Here, we reduce the number of candidate genes associated with a large effect QTL for changes in visual mate preference from 200 to just 6. First, we confirm that genomic regions previously associated with male courtship modulate preference behaviors, rather than other traits that may influence courtship time. We then investigate gene expression in the brains of *H. melpomene*, *H. cydno*, and their hybrids, across development. We identify two genes, *regucalcin2* and an ionotropic glutamate receptor, that are differentially expressed both in species and hybrid comparisons, in a manner consistent with known patterns of dominance. We also find substitutions in the protein-coding regions of 4 genes predicted to affect protein function. These six candidates are located in regions resistant to interspecific gene flow, as expected for genes that contribute to reproductive isolation. Our candidates have been implicated in key components of neural signaling, suggesting shifts in visual mating preference do not rely on primary photoreceptor genes, but instead involve changes in visual integration or processing. Such genetic changes would allow mate preference evolution without altering perception of the wider environment.

Keywords: speciation | behavioral isolation | mate preferences | behavioral genetics

Significance statement

Many species remain separate not because they fail to produce viable hybrids but because they "choose" not to mate. Although these changes in behavior can be hardwired in the genome, we know little about the specific genes involved. *Heliconius* butterflies display a striking diversity of warning color patterns, which they use as cues to recognize conspecifics. We analyze a genomic region associated with changes in these visual behaviors in two *Heliconius* species. We couple quantitative trait locus (QTL) studies with gene expression and population genomic analyses to identify candidate genes associated with these visual preferences. This is an important step towards understanding how behavioral differences, crucial to speciation, are generated both during development and across evolutionary time.

Introduction

The evolution and maintenance of new animal species often relies on the emergence of divergent mating preferences[1,2]. Changes in sensory perception or other neural systems must underlie differences in innate behaviors between species, and will ultimately have a genetic basis. However, although the significance of behavioral barriers for speciation has been recognized since the Modern Synthesis[3], we still know little of the genetic mechanisms underlying changes in mating preferences, or variation in behaviors across natural populations more broadly[4,5]. Identifying the genes underlying species-specific mating preferences provides an important route towards understanding how behavioral differences are generated, both during development and across evolutionary time.

Previous genetic studies of isolating preference behaviors have largely been limited to the identification of causal genomic regions, which almost invariably contain many genes[6,7,8,9]. Only a handful of studies have identified likely candidate genes that contribute to behavioral preferences causing reproductive isolation between species. These are largely limited to studies of chemosensory-guided mating preferences[10,11,12,13], which have identified changes at chemoreceptor genes as the underlying evolutionary mechanisms. To our knowledge, only two studies – in incipient fish species – have identified candidates for visual mate preference evolution, albeit indirectly, both suggesting a role for sensory perception mediated by changes in the peripheral visual system[14,15]. However, whether or not visual mate preferences generally involve shifts at the sensory periphery, or shifts in downstream processing, is not known.

The Neotropical *Heliconius* butterflies are well known for their diversity of warning color patterns. These patterns are often associated with Müllerian mimicry, where distantly related species converge on the same warning signals[16,17]. However, within a community, closely related species often look very different. Not only are warning patterns under disruptive selection due to increased predation of intermediate hybrid phenotypes[18], but these divergent patterns are also used as mating cues. In particular, males almost invariably prefer to court and attempt to mate live females, and artificial models, that share their own warning pattern[19]. This leads to varying degrees of premating isolation between taxa with divergent warning patterns[20,21].

The closely related species *Heliconius melpomene* and *H. cydno* differ in warning patterns, and although male hybrids are fertile, the two species remain separate largely due to strong assortative mating[22]. In central Panama, *Heliconius melpomene rosina* shares the black, red and yellow pattern of its local *Heliconius erato* co-mimic. In contrast, *Heliconius*

cydno chioneus mimics the black and white patterns of *H. sapho*. Differences in warning

pattern between *melpomene* and *cydno* are largely due to expression differences in just three

genes, specifically *optix*[23], *WntA*[24] and *cortex*[25]. Visual preferences for these divergent

patterns are particularly apparent in males, which strongly prefer to court conspecific

females[26,27,28].

Quantitative trait locus (QTL) mapping of *H. melpomene* and *H. cydno* has revealed

three genomic regions of major effect that influence the relative time males spend courting

red *melpomene* or white *cydno* females[28]. Notably, the best supported QTL was in the same

genomic region (within 4.8cM) as *optix*, the gene responsible for presence or absence of the

red color pattern elements in *Heliconius*[23]. Genetic linkage between ecological and mating

traits will facilitate speciation by acting against breakdown of these genetic associations by

recombination[29]. Nevertheless, this QTL, and its associated candidate region, contain

hundreds of genes, and the exact genes responsible for differences in preference behavior

remain unknown.

To identify specific genes that may underlie behavioral differences, we first confirm

that the QTLs identified previously are associated with variation in male courtship initiation

behaviors. We next identified genes within the major behavioral QTL, which were i)

differentially expressed in the neural tissue (central brain, optic lobes and ommatidia) of *H.*

melpomene and *H. cydno*, or ii) have protein coding changes predicted to alter protein

function. Out of 200 genes within the QTL region, we identify just 6 candidates likely to

underlie assortative mating behaviors.

Results

Chromosome 18 is associated with differences in courtship initiation.

Our previous results reveal that QTLs on chromosomes 1, 17 and 18 influence the relative

time hybrid males spend courting red *melpomene* or white *cydno* females[28]. However, the

time males spend courting a particular female might depend not only on male attraction, but

also on the female's response (and in turn his response to her behavior). To confirm that these

previously reported QTLs influence male approach *behaviors* (as opposed to other traits that

may influence courtship, for example male morphology[30]), we reanalyzed our previous data,

this time explicitly considering whether males initiated courtship towards *melpomene*, *cydno*

or both types of female during choice trials. Consistent with our previous analyses[28], we

found that F1 and backcross-to-*melpomene* males initiated courtship more frequently with

melpomene females, whereas courtship initiation behaviors segregate in the backcrosses to

Figure 1. Genotype at the preference QTL on chromosome 18 influences courtship initiation. Ternary plots showing the number of 15-minute choice trials in which courtship was initiated towards *melpomene, cydno* or both females for different male types. Left ternary axis shows proportion of trials where courtship was initiated towards *H. cydno* female only, bottom axis towards *H. melpomene* female only, and right axis towards both female species. Orange points represent individuals that have inherited at least one *melpomene* derived allele at the preference QTL on chromosome 18 (i.e. either *melp/melp* or *cyd/melp)*; and blue points represent individuals that are homozygous for *cydno* alleles at the preference QTL on chromosome 18 (i.e. *cyd/cyd*). Point size is scaled to the number of trials in which the male showed a response and a 'jitter' function has been applied. 95% credibility intervals (CrIs) for all three proportions are shown as hexagons around predictors with lines projecting to corresponding values on the three axes.

cydno (**Figure 1**). Notably, backcrosses-to-*cydno* males heterozygous at the QTL on chromosome 18 (*i.e.* with a *melpomene* allele derived from the F1 father) initiated courtship towards *melpomene* females more frequently than males homozygous for the *cydno* allele (**Figure 1**, bottom left; *n* = 139, ΔELPD: -10.9 (S.E.±5.1), *i.e.* a change of 2.14 SE units). Together with previous evidence that male hybrids bearing *melpomene* alleles at *optix* prefer to court the artificial models of *melpomene* females over those of *cydno*[27], these results suggest that the QTL on chromosome 18 harbors genes for visual attraction behaviors towards females with the red pattern. Consequently, we focused our subsequent analyses on this QTL

on chromosome 18 (and also because tight linkage of *optix* allowed us to track the alleles at

preference-color locus in hybrid crosses). The QTL on chromosome 1 was also retained in our

model of initiation behaviors (**Supporting figure 1;** n = 139, ΔELPD = -13.6 (SE±5.7), *i.e.* a

change of 2.34 SE units), in contrast to the QTL on chromosome 17 which was not retained (n

= 139, ΔELPD = -2.1 (S.E.±3.0)). Results for the QTL on chromosome 1 are reported in the

supporting materials (**Supporting table 1**).

27 *genes* within the major QTL are differentially expressed in the brains and eyes of *H.*

***cydno* and *H. melpomene*.**

We hypothesized that changes in gene regulation that determine differences in visual mate

preference behaviors might occur during pupal development (for instance, during visual

circuit assembly) or in the imago, and must involve changes in the peripheral and/or central

nervous system[31]. Therefore, we generated RNA-seq libraries for combined eye and brain

tissue, across two pupal stages (around the time of ommochrome pigment deposit and half-

way through pupal development) and one adult stage, for *H. melpomene* and *H. cydno* and

compared their gene expression levels. We found considerable differential expression at the

QTL on chromosome 18 (the QTL spans 2.75 Mb, and contains 200 genes). We identified 27

genes within the QTL region that show differential expression between *melpomene* and

cydno, in at least one of the three developmental stages, and with a minimum fold change

threshold of 2. The same genes were frequently differentially expressed across development

(**Supporting table 1**), with 11 genes being differentially expressed in more than one stage.

Differentially expressed genes were mostly located within the QTL peak (*i.e.* the genomic

region with strongest statistical association with male preference) or in close proximity to

optix (**Figure 2**).

The genomic region between the start of chromosome 18 and *optix* (comprising the

QTL peak) is highly divergent between *melpomene* and *cydno*[32], and divergent coding

sequences within this region could also introduce mapping biases of RNA-seq reads. To

account for this, we repeated the analysis having mapped to both the *H. melpomene* reference

genome[33] and to a *H. cydno* genome[34]. Generally, we found similar patterns of differential

expression when mapping to the *H. cydno* genome (**Supporting figure 2, Supporting table**

2). Nevertheless, in subsequent analyses we excluded two genes, HMEL034187g1 and

HMEL034229g1, which showed reversal of the fold change or did not show differential

expression when mapping to the *H. cydno* genome respectively.

Figure 2. Differential expression at the preference QTL region on chromosome 18. Left: Summary of the comparative transcriptomic analyses with stage, number of samples and chromosome 18 composition. Right: the corresponding results, zooming in on the QTL region on chromosome 18. The *x*-axis represents physical position. The QTL peak, and the rest of the QTL 1.5 LOD candidate region (from[28]) are shown in green and purple, respectively. Points correspond to individual genes, with the *y*-axis indicating the *log₂*(fold-change) for each comparison. The two horizontal dashed lines (at *y*-values of 1 and -1) indicate a 2-fold change in expression. Genes showing a significant 2-fold+ change in expression level between groups are highlighted in orange and blue, where orange indicates higher levels in *melpomene* or in the hybrids *cyd/melp* (blue if in *cydno* – hybrids *cyd/cyd*). Vertical dashed lines highlight those genes that are differentially expressed between *melpomene* and *cydno* AND between *cyd/melp* vs *cyd/cyd* individuals, at the same stage. Two genes highlighted by dashed fuchsia vertical lines were excluded because they did not show differential expression, or showed reversal of the fold change when mapping RNA-seq reads to the *H. cydno* genome. Note that *Heliconius* brain (reconstruction) images, added for reference, do not include the eyes (ommatidia and retinal membrane).

A *regucalcin* and an *ionotropic glutamate receptor* are upregulated in both *H. melpomene* and F1 hybrid males.

Our previous behavioral experiments suggest that the alleles for the *melpomene* behavior are dominant over the *cydno* alleles[27,28] (**Figure 1**). Given this pattern of dominance, we predicted that genes underlying variation in male preference to be up- or down- regulated in the brains of both *melpomene* and first generation (F1) hybrid males, with respect to *cydno*. Of the putative genes differentially expressed between *cydno* and *melpomene* reported above, only four, within the QTL candidate region, were differentially expressed between the F1 hybrids and *cydno* (**Figure 2**). These included two *regucalcins* (also called *senescence marker proteins-30*: HMEL013552g1, HMEL034199g1), an *ionotropic glutamate receptor* (HMEL009992g4), which is a putative ortholog of *Grik2*, and one gene with no annotated

function (HMEL009992g1). We obtained the same results regardless of whether we

considered both males and females together, or males alone.

Further inspection of spliced mRNA-reads indicated that the two annotated

regucalcins were in fact a single gene (from now on referred to as *regucalcin2*). This was also

the case for the *ionotropic glutamate receptor* and the gene with no annotated function (from

now on referred to as *Grik2*). To ensure that apparent fragmentation of a few gene models in

the *H. melpomene* (Hmel2.5) annotation[35], in which exons belonging to the same isoform are

incorrectly considered distinct genes, did not introduce inaccuracies in estimates of

differential gene expression, we produced a new transcript-based annotation of the

melpomene genome using Cufflinks's RABT[36]. Repeating all comparative transcriptomic

analyses using this new annotation (where exons were correctly considered as part of single

genes), we confirmed that both *regucalcin2* and *Grik2* were differentially expressed in both

species and hybrids comparisons.

Figure 3. *Grik2* and *Regucalcin2* show evidence of allele specific expression. Points indicate the value, and bars
the standard error, of the (base 2) logarithmic fold change in expression between parental species (horizontal) and
the alleles in F1 hybrids (vertical), for candidate genes (as defined in the transcript-guided annotation). Dashed
lines indicate the threshold for a 2-fold change in expression for the genes in the species (horizontal), and for the
alleles in the hybrids (vertical).

Differential expression of *Grik2* in adults is likely due to cis-regulatory effects.

To determine whether differences in gene expression levels between parental species were

due to *cis*- or *trans*-regulatory changes, we conducted allele specific expression (ASE)

analyses in adult F1 hybrids. In F1 hybrids, both parental alleles are exposed to the same

trans-environment, and consequently *trans*-acting factors will act on alleles derived from each

species equally (unless there is a change in the cis-regulatory regions of the respective

alleles). Therefore, differences in allele specific expression should indicate changes in *cis*-

regulatory regions[37]. Both candidate genes (*Grik2* and *regucalcin2*) had a very low number of

SNPs that could differentiate the *melpomene* and *cydno* allele (using both gene models of the

Hmel2.5 annotation and RABT annotation) and few reads mapped to these SNPs.

Nevertheless, for *Grik2,* the *melpomene* allele was significantly more highly expressed

relative to the *cydno* allele (p=0.017, Wald test), suggesting cis-regulatory effects (**Figure 3**).

For *regucalcin2,* although there was a tendency towards up-regulation of the *melpomene*

allele, consistent with *cis*-regulation, we did not have sufficient power to rule out *trans*-only

regulatory effects (**Figure 3;** p=0.108, Wald test).

***Grik2* is differentially expressed in hybrids that essentially differ only for allelic**

composition at the behavioral QTL region.

In order to study the specific effects that *melpomene* derived alleles at the QTL on

chromosome 18 had on gene-expression, we introgressed this region into a *cydno* background

through multiple backcrosses (crossing design in **Supporting figure 3**). We wanted to

investigate whether differences at this QTL regulated expression of any specific genetic

pathway during development, and more generally what changes in genome-wide transcription

were observed in hybrids differing (mostly) just at this QTL region.

Notably, in these third-generation backcross hybrid comparisons, across the entire

genome only 23 and 29 genes were differentially expressed (at 156h after pupal formation

(APF) and at 60hAPF, respectively). Of these, 20 and 19 genes (at 156hAPF and at 60hAPF,

respectively) were located on chromosome 18, indicating that gene expression differences in

these comparisons were mostly restricted to the preference-color region on chromosome 18,

segregating for *cyd/melp* or *cyd/cyd* alleles. No genetic pathway was enriched for gene

expression differences between these hybrids at either pupal stage (PANTHER enrichment

test[38]), suggesting that overall this QTL harbors a few, modular changes in gene regulation in

the developing brain/eyes of *cydno* and *melpomene***.** *Grik2* was the only gene detected as

differentially expressed between species and hybrids at these pupal stages (**Supporting figure**

4).

To verify that differential expression of candidate genes at the QTL region is driven

by *melpomene* alleles on chromosome 18 and not by other *melpomene* alleles at *trans*-acting

genes on other chromosomes, we compared gene expression levels between hybrids carrying

cyd/melp vs. *cyd/cyd* regions on chromosomes chr1, chr4 and chr15, chr20 (**Supporting**

figure 5A). In these comparisons, there was no signal of differential expression on

chromosome 18**.** This supports the *cis*-regulatory activity of the *melpomene* allele of

candidate genes on chromosome 18. To test this further, we conducted another allele specific

expression study in the BC3 hybrids, which suggested *trans*-regulatory effects for *Grik2* at

these pupal stages, but were less conclusive with regard to *regucalcin2* (**Supporting figure**

6). Since causal gene/s might exert an effect on behavior due to their action during

development or in adult form, and this action might in turn be differently (cis- vs trans-)

regulated, we still considered both genes as strong candidates.

291
Figure 4. **Gene flow at the QTL region for behavior.** Admixture proportion (f_d) values estimated in overlapping
100kb (top) and 20kb (bottom) windows for chromosome 18 (top) and the QTL region (bottom) between *H.*
*melpomene rosin*a and *H. cydno chioneus*, with candidate genes positions highlighted by a vertical dashed line,
and *optix* location displayed for reference. The *x*-axis represents physical position, the *y*-axis indicates the f_d value.
f_d values close to zero indicate that the proportion of shared derived alleles, and consequently gene flow, between
melpomene and *cydno* is small (or zero), implying localized selection against foreign alleles that introgress between
the two species.

300
301

4 genes with protein-coding substitutions within the QTL candidate region have

predicted effects on protein function.

Because shifts in behavioral phenotypes could be due to changes in protein-coding regions,

we additionally considered protein-coding substitutions between *melpomene* and *cydno*.

Overall, we found 152 protein-coding substitutions, spanning 54 of the 200 genes across the

entire QTL candidate region. We then studied whether these variants were predicted to have

non-neutral effects on protein function with PROVEAN[39]. The PROVEAN algorithm predicts

the functional effect of protein sequence variations based on how they affect alignments to

different homologous protein sequences**.** We found 4 genes with such predicted effects

(PROVEAN score < -2.5): Specifically, a *WD40*-repeat domain containing protein

(HMEL013551g3), a *cysteine protease* (HMEL009684g2), a *MORN* motif containing protein

(HMEL006660g1), and another *regucalcin* (HMEL013551g4) adjacent to, but distinct from,

that found to be differentially expressed above (from now on referred to as *regucalcin1*).

Candidate genes occur in regions with reduced gene flow.

Of our six candidate genes for preference behaviors that contribute to reproductive isolation

between *H. cydno* and *H. melpomene* (*regucalcin2*, *Grik2* and the four genes with protein

coding modifications), five are found within the QTL peak (**Figure 4**). Genetic changes

causing reproductive isolation between populations are expected to reduce localized gene

flow in their genomes. Therefore, we compared the position of our candidate genes to

estimated levels of admixture proportions (f_d)[40] between *H. melpomene* and *H. cydno* across

the QTL candidate region[41]. We found that all candidate genes were located in genomic

regions with low f_d values (**Figure 4**), suggesting localized resistance to gene flow between

melpomene and *cydno* at these genes and their putative *cis*-regulatory regions.

Discussion

Behavioral isolation is frequently implicated in the formation of new species, and involves the

correlated evolution of both mating cues and mating preference. Here we have analyzed a

genomic region in a pair of closely related sympatric butterflies, *H. cydno* and *H. melpomene*,

that contains genes for divergence in both an ecologically relevant mating cue and the

corresponding preference. Physical linkage between ecological and mating traits will facilitate

speciation by allowing different barriers to act in concert to restrict gene flow[42,43]. Although

the genes underlying changes in the warning pattern cue in *Heliconius* are well characterized

(e.g. *optix*), those underlying the corresponding shift in behavior have not been identified. We

have pinpointed a small number of genes that fall within the QTL peak, which show either

expression (*regucalcin2* and *Grik2*) or protein coding differences (HMEL013551g3,

HMEL009684g2, HMEL006660g1, and *regucalcin1*) and fall within a region of reduced

admixture, that are strong candidates for modulating mating behavior.

Two broad neural mechanisms could underlie the evolution of divergent visual

preferences, involving changes in either i) detection at the sensory periphery or ii) the

processing and/or integration of visual information. Although *H. melpomene* and *H. cydno*

have the same retinal mosaics/class of photoreceptors[44], spectral sensitivity in *Heliconius* eyes

could be altered by filtering pigments[45], or other physiological processes taking place at the

photoreceptors/sensory periphery, eventually shifting sensitivity towards different

wavelengths (and possibly color patterns). It has previously been hypothesized that the gene

regulatory networks for ommochrome deposition in the *Heliconius* eyes might have been co-

opted in the wings[46], where *optix* plays a central role, and therefore that *optix* might play a

role in eye pigmentation in *Heliconius*. However, the protein product of *optix* has not been

detected in pupal or adult retinas of various *Heliconius* species tested[47], and therefore has no

obvious link to ommochrome deposition in the eyes. More generally, the underlying

evolutionary mechanism is unlikely to involve detection at photoreceptors, as this would

probably have a broad effect on downstream processing[2] and alter the visual perception of the

animal's wider environment.

The second mechanism, involving changes in the processing, and/or integration, of

visual information, could act through an alteration of neuronal activity or connectivity. For

instance, different levels of gene expression in conserved neural circuits between *melpomene*

and *cydno* may affect overall synaptic weighting and determine whether a signal (e.g. color

and motion) elicits a motor pattern (response towards a female) or not. Consistent with this

scenario, the composition of ionotropic receptors at post-synapses is a key modulator of

synaptic transmission[48], implicating *Grik2*. Interestingly, differential expression of ionotropic

glutamate receptors is also associated with variation in social and aggressive behaviors in

vertebrates[49,50]. *Regucalcins* are involved in calcium signaling[51], which regulates synaptic

excitability and plasticity[52], and has an important role in axon guidance[53] (albeit alongside

additional roles across a broad range of biological processes), making the two *regucalcins* we

identify strong candidates for behavior.

Changes in the regulation of genes with pleiotropic effects are likely to be less

detrimental compared to changes in their protein-coding sequences[54] (although emerging

evidence has begun to suggest that enhancer/repressor elements may be more pleiotropic than

previously thought[55,56]). Furthermore, there is considerable evolutionary potential in the co-

option of transcription factors/networks[54] that regulate neural patterning or neuron-type

activity, possibly resulting in novel adaptive expression patterns. In line with this,

Regucalcin2 and *Grik2,* which are differentially expressed in the eyes and brain in both our

species and hybrid comparisons, are likely to be involved in multi-functional processes, such

as calcium signaling and ion transport, and likely have pleiotropic alleles. We also found

evidence of cis-regulatory effects for both genes (albeit not significant for *regucalcin2*),

which would be required of the causal genetic change within the QTL, if it were to be in gene

regulation.

Despite expectations that non-coding, regulatory loci may provide a flexible route to

divergent mating preferences, we also found substitutions in coding regions at the QTL,

which are predicted to have an effect on protein functioning and therefore remain strong

candidates. These genes include *regucalcin1*, which is distinct from, but located next to,

regucalcin2 (which is differentially expressed). Notably, the eye transcript of *regucalcin1* was

recently characterized as fast-evolving across *Heliconius* species[57]. Other candidates include a

cysteine protease, which functions in protein degradation, and might be linked to behavior for

example through degradation of neurotransmitters, a *MORN* motif containing protein

(function unknown), and a *WD40* containing protein. *WD-repeat* containing proteins have

been implicated in a wide array of functions ranging from signal transduction to apoptosis

(https://www.ebi.ac.uk/interpro).

Although preference for red coloration and the *optix* gene are tightly linked, we find no

evidence that *optix* is differentially expressed in the eyes or brains of our two species. It is

also not located within the QTL peak (and it contains no non-synonymous changes in protein

coding regions[23]). It seems unlikely therefore that changes in cue and preference are

pleiotropic effects of the same allele. More generally, although we have pinpointed the

strongest candidates yet identified for assortative mating behaviors in *Heliconius*, it is

possible that actual causal changes in gene regulation are restricted to developmental stages

other those sampled, or restricted to a few neuronal populations not detected with

transcriptomic data from eyes and whole brain tissue. Nonetheless, by sampling at two pupal

stages (around the time of *optix* expression/ommochrome pigment deposit in the wing/eye and

halfway through pupal development) and at the adult stage, we should have captured

important transitions for the behavioral programming of the two species.

Functional studies of candidate genes would of course be desirable in order to clarify

the link between individual genes and behavioral divergence. For example, can a distinct

genetic change at a single locus trigger a shift in behavior, or are multiple interacting loci

necessary to affect it? Although CRISPR-Cas9 technology has been successfully used to

produce knock-out individuals[54,58,59] for color patterning genes in *Heliconius*, "behavioral

genes" will present a greater challenge. Gene knock-outs could broadly impair an individual's

behavioral phenotype, permitting only tentative inference about the link between the action of

a gene and a specific behavior/perception. While posing a significant challenge, in the future,

this or other emerging technologies that allow switching of specific alleles will allow direct

functional experiments on behavioral phenotypes for the candidates we have identified.

Work in the past decade has shown that complex innate behavioral differences between

species can be encoded in relatively few genetic modules[60,61], but very few studies[62,63,64] have

identified specific genes underlying behavioral evolution. Although traditional laboratory

organisms continue to provide important insights into the evolution and genetics of behavior,

comparative approaches are required to determine if developmental principles can be broadly

applied, and also to incorporate a wider range of phenotypic variation. The challenge now is

to increase the resolution of studies in non-traditional systems, in order to link individual

genetic elements to behaviors, *and* the sensory and/or neurological structures through which

they are mediated. In this light, we have identified a small handful of strong candidate genes

associated with the evolution of visual mate preference behaviors in *Heliconius*. These genes

are in tight physical linkage with the locus for the corresponding shifts in an ecologically

relevant mating cue, providing an important opportunity to investigate the build-up of genetic

barriers crucial to speciation. The candidate genes identified seem more likely to alter visual

processing or integration, rather than detection at photoreceptors, consistent with permitting

changes in mate preference without altering perception of the animal's wider environment.

Materials and Methods

Courtship initiation analyses. Butterfly rearing, crossing design and genotyping are

described in detail elsewhere[28]. In brief, we assayed male preference behaviors for *H.*

melpomene, *H. cydno,* their first generation (F1) hybrids and backcross to hybrids to both

parental species in standardized choice trials. Males were introduced into outdoor

experimental cages (1x1x2m) with a virgin female of each species and courtship behaviors

recorded. Whenever possible, trials were repeated for each male (median = 5 trials). To

determine whether previously identified QTLs for courtship time contribute to variation in

courtship *initiation* behaviors, we performed a *post-hoc* analysis using categorical models in a

Bayesian framework with a multinomial error structure, using the R package *brms*[65]. In

contrast to our previous analysis[28], in which we considered the number of minutes (*i.e.* time)

for which courtship was directed towards *H. cydno* or *H. melpomene* females, here the

response variable was number of trials in which male courtship was *initiated* towards *H.*

cydno females only, *H. melpomene females* only*, or both female types (hereafter referred to as

"*initiation*")*. Across males the median number of trials with a response was 3. Using

backcross-to-*cydno* males only, we fitted *initiation* as a response variable to genotype

(*cyd*/*cyd* or *cyd*/*melp*) at each QTL, which were included as separate fixed effects. Individual

ID was fitted as random factor. To test the effect of each QTL on male *initiation*, we

compared the saturated model incorporating all three QTL with reduced models excluding

each QTL in turn, using approximate leave-one-out (LOO) cross-validation as implemented in

brms[66], and based on expected log pointwise predictive density (ELPD). Normal distribution

of ELPD can be a straightforward approximation given our large samples sizes (n=139)[66].

Therefore, we considered an absolute value of ELPD greater than 1.96 units of its standard

error as indicative of the reduced model being less-informative than the saturated model (95%

confidence). Males that did not initiate courtship to any female across trials were excluded

from analyses, resulting in a dataset of 139 males, from a total of 146 backcross males for

which we had genotype data. Finally, we extracted predictors and credibility intervals for

backcross males with differing genotypes from the minimum adequate model. Credibility

intervals for *H. melpomene, H. cydno,* F1 hybrid and backcross to *melpomene* males

displayed in Figure 1 were generated following the same procedures. Raw data and analysis

code are available in the following github repository:

https://github.com/SpeciationBehaviour/neural_genes_heliconius.git

Butterfly collection, rearing and crossing design for expression analyses. Wild *H.*

melpomene rosina and *H. cydno chioneus* individuals were caught along Pipeline Road near

Gamboa, Panama, in the Soberania National Park, and used to establish stocks at the

Smithsonian Tropical Research Institute insectaries in Gamboa. Butterflies were reared in

common garden conditions, in 2x2x2m cages, and provided with fresh *Psiguria* flowers and

10% sugar solution. Larvae were reared on fresh *Passiflora* shoots/leaves until pupation. *H.*

cydno, *H. melpomene* and hybrid individuals used for RNA-seq (see below) were reared

concurrently and under the same conditions. F1 hybrids were obtained by crossing a wild-

caught *H. m. rosina* male to an insectary-bred virgin *H. c. chioneus* female.

The introgression line was generated by outcrossing a hybrid male with a red forewing

band (crossing design shown in **Supporting figure 3**) to virgin *H. cydno* females, over three

generations. The peak of the behavioral QTL reported previously[28] on chromosome 18 (at

0cM) is in very tight linkage with the *optix* color pattern locus (at 1.2cM), which controls for

the presence and absence of the red forewing band seen in *H. melpomene rosina*. Presence of

the red forewing band is dominant over its absence so that segregation of the red band can be

used to infer genotype at the *optix* locus. Specifically, hybrid individuals with a red forewing

band are heterozygotes for *H. melpomene/H. cydno* alleles at the *optix* locus, whereas

individuals lacking the red band are homozygous for the *cydno* allele. Due to the tight linkage

we expected little recombination between *optix* and QTL peak even after three generations of

introgression, allowing us to infer genotype at the preference-*optix* locus (which we

confirmed with genetic data, see below).

Tissue dissection, RNA extraction and mRNA sequencing. Eye (ommatidia and retinal

membrane) and brain tissue (central brain and optic lobes) were dissected out of the head

capsule in cold (4 ºC) 0.01M PBS solution, at two pupal stages: 60 hours after pupal

formation (60h APF) and 156h APF; and in adults aged 9 - 13 days. We sampled adults at

around 10 days of age because by this stage males are mature and frequently court females[17].

Adult males and females sampled were sexually naive. We decided to sample at 60h APF

because this is the developmental stage at which *optix* is expressed in the wing, so we

hypothesized that it might had also been when *optix* is expressed in the brain. We sampled at

156h APF as a putative stage halfway through pupal development, and at this stage most of

the major neural connections have just been established in the *Heliconius* brain (Stephen

Montgomery, unpublished data).

Tissues were stored in RNAlater at 4 ºC for 24 hours, and subsequently at -20 ºC, until RNA

extraction. Total RNA was extracted using TRIzol Reagent (Thermo Fisher, Waltham, MA,

USA) and a RNeasy Mini kit (Qiagen, Valencia, CA, USA). Samples were treated with

DNase I (Ambion, Darmstadt, Germany). Integrity of total RNA was checked either on an

agarose gel or using an Agilent Bioanalyzer 2100 (Agilent Technologies, Santa Clara, CA,

USA). RNA concentration was measured on a Nanodrop spectrophotomer. Illumina TruSeq

RNA-seq libraries were prepared and sequenced at Edinburgh Genomics (Edinburgh, UK)

with 100 bp paired-end reads. To avoid lane effects the distribution of the species samples

was randomized on the sequencing platform. More detailed information about individuals and

sequencing yields can be found in the Supporting file 1.

RNA-seq read mapping and differential gene expression analyses. After a quality control

of RNA-seq reads with FastQC, we trimmed adaptor and low-quality bases using TrimGalore

v.0.4.4 (https://www.bioinformatics.babraham.ac.uk/projects/). RNA-seq reads were mapped

to the *H. melpomene* 2.5 genome[33]/annotation[35] using STAR v.2.4.2a[67] in 2-pass mode. We

only kept reads that mapped in 'proper pairs' using Samtools[68]. The number of reads mapping

to each gene were estimated with HTseq v. 0.9.1[69] with model "union", thus excluding

ambiguously mapped reads. Differential gene expression analyses between species/hybrids

were conducted in DESeq2[70]. We considered only those genes showing a 2-fold change in

expression level, and at adjusted (false discovery rate 5%) p-values < 0.05, to be differentially

expressed, to exclude expression differences caused by known differences in brain

morphology[71] (Montgomery et al., in prep).

Sexing pupae. In all DESeq2 analyses, sex was included as a random factor. To sex pupae,

we first marked duplicate RNA mapped reads with Picard

(https://broadinstitute.github.io/picard/), and used GATK 3.8[72] to split uniquely mapped reads

into exon segments and trim sequences overhanging the intronic regions. We then used

Haplotype Caller on each individual, using calling and filtering parameters according to the

GATK Best Practices for variant calling on RNA-seq data. The sex of pupal samples was

inferred from the proportion of heterozygous (biallelic) SNPs using the R package SNPstats.

Males (ZZ) were expected to have $\gg$ 0% heterozygous sites, whereas females (ZW) to have

0%. Z-linked heterozygosity of the pupal samples (**Supporting table 3**) were in line with

expectations (either ~ 0 for females or an order of magnitude higher for males), and matched

heterozygosity of either adult males or females, for which the sex was determined from

external morphology.

Inference of gene function and transcript-based annotation. Biological functions of

annotated genes were inferred with InterProScan v5[73], using the corresponding Hmel2.5

predicted protein sequences. InterProScan uses different databases like InterPro, Pfam,

PANTHER, and others, to infer functional protein domains and motifs (based on homology).

To study whether specific biological functions were enriched among genes showing

differential expression among hybrid types, we conducted the PANTHER enrichment test[38]

(with Bonferroni correction for multiple testing) using *Drosophila melanogaster* as the

reference gene function database.

536   Upon detailed inspection of the mapping coverage of spliced RNA-seq reads to the

Hmel2.5 gene annotation, we noticed that some gene models were fragmented, namely, a few

exons that appeared to be spliced together were incorrectly considered distinct genes. To

check that this did not introduced inaccuracies in our differential gene expression analyses,

we re-annotated the *melpomene* genome using the Cufflinks reference annotation-based

transcript (RABT) assembly tool[36]. We used the transcriptomic data from both *melpomene*

and *cydno* to reannotate the *melpomene* genome, separately for every developmental stage,

and reconducted the differential gene expression analyses in DESeq2 as described above.

Inference of BC3 hybrids genome composition. In order to perform comparative transcriptomic analyses between third-generation backcross hybrids (BC3) segregating at the QTL on chromosome 18 (crossing design in **Supporting figure 3**), we first determined which genomic regions in these hybrids were heterozygous (*cyd/melp*) or homozygous (*cyd/cyd*). For this, we inferred variants from RNA-seq reads for each BC3 hybrid (individually as above), and from the combined *melpomene* and *cydno* samples. For the species, we used HaplotypeCaller[72] on RNA-seq samples from all developmental stages of either species, to produce individual genomic records (gVCF), and then jointly genotyped *melpomene* and *cydno* gVCFs (separately for the two species) using genotypeGVCFs with default parameters. Genotype calls were filtered for quality by depth (QD) > 2, strand bias (FS) < 30 and allele depth (DP) > 4. For further analyses we kept *biallelic* genotypes only. We then used the *intersect* function of *bcftools*[68] to infer variants exclusive to the *cydno* and to the *melpomene* samples.

We calculated the fraction of variants that each BC3 hybrid individual shared with the *melpomene* and with the *cydno* samples, in non-overlapping 100kb windows. We compared these to the fraction of variants that a F1 hybrid and a *H. cydno* individual (not included in the combined genotyping of the *cydno* samples), shared with the same species samples, and found that they matched either one of them, indicating heterozygous (*cyd/melp*) or homozygous (*cyd/cyd*) regions (**Supporting figure 5B**). In this analysis, we considered only those 100kb windows where BC3 hybrids/F1 hybrid/*H. cydno* individuals shared more than 30 variants with the *melpomene*/*cydno* samples.

To corroborate our findings, we repeated the same type of analysis, this time calling species-specific variants for *melpomene* and *cydno* using 10 *H. melpomene rosina* and 10 *H. cydno chioneus* genome resequencing samples[41]. For this, genomic reads were mapped to the *melpomene* genome with BWA v0.7.2[74] using default parameters, genotypes were called with GATK's HaplotypeCaller and GenotypeGVCFs with default parameters, and we considered only *biallelic* genotype calls that had 10 < DP < 100 and genotype quality (GQ) > 30. With this analysis we found the same heterozygous and homozygous regions in BC3 hybrids**.**

The size and number of the introgressed regions were in line with expectations about 3rd generation backcross hybrids following our crossing design: segregating at the level of chromosome 18 and at four other chromosomes. For the BC3 hybrids sampled at 156 hours after pupal formation (APF) we had 6 *cyd/melp* and 10 *cyd/cyd* at the QTL region on chromosome 18 (**Supporting figure 5A**), for those at 60h APF, 8 *cyd/melp* and 9 *cyd/cyd* hybrids at the same region.

Allele-specific expression (ASE) in hybrids. In order to conduct ASE analyses we first identified species specific variants, fixed in either *melpomene* and *cydno*. For this, we took the quality filtered variants inferred from the species genome resequencing data, and assigned those genotype calls in *cydno* and *melpomene* for which allele frequency (AF) was > 0.9 as homozygous (we did not consider indels in this analysis). We then used *bcftools intersect*[68] to get only those variants for which *cydno* and *melpomene* had opposite alleles.

At the same time, we called variants from RNA-seq reads of F1 hybrid individuals, again according to the GATK Best Practices (with the exception of parameters -window 35 -cluster 3, to increase SNPs density), and selected only heterozygous SNPs in F1s that matched the species-specific variants. Finally, we used GATK's ASEReadCounter, with default parameters, to count RNA reads in the F1 hybrids (and later on in BC3 hybrids) that mapped to either the *cydno* or the *melpomene* allele. We summed all reads mapping to either the *cydno* or *melpomene* allele/variant within the same gene (both for gene models of the Hmel2.5 gene annotation and for the Cufflinks annotation we assembled previously). To test for allele specific expression (diffASE) we fitted the model "~0 + individual + allele" in DESeq2, setting library size factors to 1 (thus not normalizing between samples, as the test for diffASE is conducted within individuals). We only considered those alleles showing at least a 2-fold change in expression and $p < 0.05$, as differentially expressed.

In order to check that there were no biases in alleles assignment to one of the two species, we analyzed the ratios of the species alleles, for every gene, and checked that they were not systematically biased to either one of the two species. The $\log_2$ fold-changes of the species alleles were centered around 0, suggesting no obvious bias in alleles assignment[75] (**Supporting figure 7**).

Protein-coding substitutions and predicted effects on protein-function. We inferred fixed variants in protein-coding regions from the combined *melpomene* and *cydno* RNA samples in order to include variants from genes for which we detected expression in the brain/eyes across the 3 stages. We took the quality filtered variants called from the joint genotyping of RNA-seq data of *cydno* and *melpomene* (from all stages), and selected those genotype calls for which allele frequency (AF) > 0.8, and where the allelic variant was present in at least 7 individuals of the ~30 samples (for each species). We retained those substitutions/indels validated with the genome resequencing data. For this, of the genotype calls found in RNA reads from brain/eyes of different stages, we kept only those that were also called in at least 8

of the 10 genome resequencing samples of each species. We considered this overlapping set of variants as being fixed in *H. melpomene rosina* or *H. cydno chioneus*. Following a similar approach to Bendesky *et al.*[64], we then restricted this set of substitutions between *cydno* and *melpomene* to protein-coding regions, and selected those non-synonymous substitutions that were considered to have moderate or high effect on protein function, with SNPeff[76]. Finally, we used the PROVEAN algorithm[39], to further study the functional effects of these substitutions on protein function. The PROVEAN algorithm predicts the functional effect of protein sequence variations based on how they affect alignments to homologous protein sequences (for this we used the PROVEAN protein database online). We selected those amino acid changes with the suggested PROVEAN score < -2.5, indicating non-neutral effect on protein function.

Admixture analyses. We retrieved estimated admixture proportions between *H. melpomene rosina* and *H. cydno chioneus*, for 100kb and 20kb windows, from Martin et al.[41]

Data accessibility. RNA-seq data will be deposited on a public database (https://www.ebi.ac.uk/ena) on acceptance. Analysis scripts and behavioral data are available at: https://github.com/SpeciationBehaviour/neural_genes_heliconius.git

Acknowledgments

We thank Liz Evans and Adriana Tapia for assistance in the insectaries. We are very grateful to Ana Pinharanda for sharing the *H. cydno* genome assembly and annotation and to Simon Martin for sharing admixture proportions and comments on the manuscript. We thank Ricardo Pereira, Nicolas Gompel and Jochen Wolf for input on the project. Jochen Wolf also kindly provided computational resources. We thank the Smithsonian Tropical Research Institute for providing research infrastructure, the Ministerio del Ambiente for permission to collect butterflies in Panama, and Edinburgh Genomics for sequencing support. MR, AEH and RMM are supported by an Emmy Noether fellowship and research grant awarded to RMM by the Deutsche Forschungsgemeischaft (DFG) (Grant Number: GZ: ME 4845/1-1). RMM was also supported by a Junior Research Fellowship from King's College Cambridge, an Ernst Mayr fellowship from the Smithsonian Tropical Research Institute, a Varley-Gradwell fellowship from the Oxford University Museum of Natural History, and the Balfour-Browne Fund, University of Cambridge. SHM was supported by a NERC IRF (NE/N014936/1). This project was also supported by an Institutional Development Award (IdeA) INBRE from the National

Institute of General Medical Sciences (NIGMS) awarded to RP (Grant number:

P20GM103475), and a European Research Council (ERC) awarded to CDJ (Grant number:

339873).

References

1. J. A. Coyne, H. A. Orr, *Speciation* (Sunderland, MA: Sinauer 2004).

2. G. Rosenthal, *Mate choice* (Princeton University Press, 2017).

3. E. Mayr, *Animal species and evolution* (Harvard University Press, 1963).

4. J. Cande, B. Prud'homme, N. Gompel, Smells like evolution: The role of chemoreceptor
evolution in behavioral change. *Curr. Opin. Neurobiol.* **23**, 152–158 (2013).

5. J. R. Arguello, R. Benton, Open questions: Tackling Darwin's "instincts": The genetic basis
of behavioral evolution. *BMC Biol.* **15**, 8–10 (2017).

6. A. J. Moehring, *et al.*, Quantitative trait loci for sexual isolation between Drosophila
simulans and D. mauritiana. *Genetics 167*, 1265-1274 (2004).

7. K. L. Shaw, S. C. Lesnick, Genomic linkage of male song and female acoustic preference
QTL underlying a rapid species radiation. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 9737-9742
(2009).

8. R. A. Bay, *et al.*, Genetic Coupling of Female Mate Choice with Polygenic Ecological
Divergence Facilitates Stickleback Speciation. *Curr. Biol.* **27**, 3344-3349 (2017).

9. M. P. Shahandeh, A. Pischedda, J. M. Rodriguez, T. L. Turner, The genetics of male
pheromone preference difference between Drosophila melanogaster and Drosophila
simulans. *G3 Genes, Genomes, Genet.* **10**, 401–415 (2020).

10. F. Gould, *et al.*, Sexual isolation of male moths explained by a single pheromone response
QTL containing four receptor genes. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 8660–8665 (2010)

11. G. P. Leary, *et al.*, Single mutation to a sex pheromone receptor provides adaptive
specificity between closely related moth species. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 14081–
14086 (2012).

12. P. Fan, *et al.*, Genetic and neural mechanisms that inhibit drosophila from mating with other
species. *Cell* **154**, 89–102 (2013).

13. P. Brand, *et al.*, The evolution of sexual signaling is linked to odorant receptor tuning in
perfume-collecting orchid bees. *Nat. Commun.* **11** (2020).

14. O. Seehausen, *et al.*, Speciation through sensory drive in cichlid fish. *Nature* **455**, 620–626
(2008).

15. K. Hench, M. Vargas, M. P. Höppner, W. O. McMillan, O. Puebla, Inter-chromosomal
coupling between vision and pigmentation genes during genomic divergence. *Nat. Ecol.*
Evol. **3**, 657–667 (2019).

16. R. M. Merrill, *et al.*, The diversification of Heliconius butterflies: What have we learned in
150 years? *J. Evol. Biol.* **28**, 1417–1438 (2015).

17. C. D. Jiggins, *The ecology and evolution of Heliconius butterflies* (Oxford University Press,
2016).

18. R. M. Merrill, *et al.*, Disruptive ecological selection on a mating cue. *Proc. R. Soc. B Biol.*
Sci. **279**, 4907–4913 (2012).

19. C. D. Jiggins, R. E. Naisbit, R. L. Coe, J. Mallet, Reproductive isolation caused by colour
pattern mimicry. *Nature* **411**, 302–305 (2001).

20. R. M. Merrill, *et al.*, Mate preference across the speciation continuum in a clade of mimetic
butterflies. *Evolution*, *65*, 1489-1500 (2011).

21. C. Mérot, C. Salazar, R. M. Merrill, C. D. Jiggins, M. Joron, What shapes the continuum of
reproductive isolation? Lessons from Heliconius butterflies. *Proc. R. Soc. B Biol. Sci.* **284**
(2017).

22. C. D. Jiggins, Ecological Speciation in Mimetic Butterflies. *Bioscience* **58**, 541–548 (2008).

23. R. D. Reed, *et al.,* Optix drives the repeated convergent evolution of butterfly wing pattern
mimicry. *Science 333*, 1137-1141 (2011).

24. A. Martin, *et al.*, Diversification of complex butterfly wing patterns by repeated regulatory
evolution of a Wnt ligand. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 12632–12637 (2012).

25. N. J. Nadeau, *et al.*, The gene cortex controls mimicry and crypsis in butterflies and moths.
Nature **534**, 106–110 (2016).

26. C. D. Jiggins, C. Estrada, A. Rodrigues, Mimicry and the evolution of premating isolation
in Heliconius melpomene Linnaeus. *J. Evol. Biol.* **17**, 680–691 (2004).

27. R. M. Merrill, B. Van Schooten, J. A. Scott, C. D. Jiggins, Pervasive genetic associations
between traits causing reproductive isolation in Heliconius butterflies. *Proc. R. Soc. B Biol.*
Sci. **278**, 511–518 (2011).

28. R. M. Merrill, *et al.*, Genetic dissection of assortative mating behavior. *PLoS Biol 17*,
e2005902 (2018).

29. J. Felsenstein, Skepticism Towards Santa Rosalia, or Why Are There So Few Kinds of
Animals? *Evolution* **35**, 124–138 (1981).

30. J. H. Massey, D. Chung, I. Siwanowicz, D. L. Stern, P. J. Wittkopp, The yellow gene
influences drosophila male mating success through sex comb melanization. *Elife* **8**, 1–20
(2019).

31. L. F. Seeholzer, M. Seppo, D. L. Stern, V. Ruta, Evolution of a central neural circuit
underlies Drosophila mate preferences. *Nature* **559**, 564–569 (2018).

32. S. H. Martin, *et al.*, Genome-wide evidence for speciation with gene flow in Heliconius
butterflies. *Genome Res.* **23**, 1817–1828 (2013).

33. J. Davey, *et al.*, Major improvements to the Heliconius melpomene genome assembly used
to confirm 10 chromosome fusion events in 6 million years of butterfly evolution. *G3* **6**,
695–708 (2015).

34. K. Darragh, *et al.*, A novel terpene synthase produces an anti-aphrodisiac pheromone in the
butterfly Heliconius melpomene. *bioRxiv*, 779678 (2019).

35. A. Pinharanda, *et al.*, Sexually dimorphic gene expression and transcriptome evolution
provide mixed evidence for a fast-Z effect in Heliconius. *J. Evol. Biol.* **32**, 194–204 (2019).

36. A. Roberts, H. Pimentel, C. Trapnell, L. Pachter, Identification of novel transcripts in
annotated genomes using RNA-seq. *Bioinformatics* **27**, 2325–2329 (2011).

37. P. J. Wittkopp, B. K. Haerum, A. G. Clark, Evolutionary changes in cis and trans gene
regulation. *Nature* **430**, 85–88 (2004).

38. P. D. Thomas, *et al.*, Applications for protein sequence-function evolution data:
mRNA/protein expression analysis and coding SNP scoring tools. *Nucleic Acids Res.* **34**,
645–650 (2006).

39. Y. Choi, G. E. Sims, S. Murphy, J. R. Miller, A. P. Chan, Predicting the Functional Effect
of Amino Acid Substitutions and Indels. *PLoS One* **7** (2012).

40. S. H. Martin, J. W. Davey, C. D. Jiggins, Evaluating the use of ABBA-BABA statistics to
locate introgressed loci. *Mol. Biol. Evol.* **32**, 244–257 (2015).

41. S. H. Martin, J. W. Davey, C. Salazar, C. D. Jiggins, Recombination rate variation shapes
barriers to introgression across butterfly genomes. *PLoS Biol.* **17**, 1–28 (2019).

42. M. Kopp, *et al.*, Mechanisms of assortative mating in speciation with gene flow: Connecting
theory and empirical research. *Am. Nat.* **191**, 1–20 (2018).

43. R. K. Butlin, C. M. Smadja, Coupling, reinforcement, and speciation. *Am. Nat.* **191**, 155–
172 (2018).

44. K. J. McCulloch, *et al.*, Sexual Dimorphism and Retinal Mosaic Diversification following
the Evolution of a Violet Receptor in Butterflies. *Mol. Biol. Evol.* **34**, 2271–2284 (2017).

45. G. Zaccardi, A. Kelber, M. P. Sison-Mangus, A. D. Briscoe, Color discrimination in the red
range with only one long-wavelength sensitive opsin. *J. Exp. Biol.* **209**, 1944–1955 (2006).

46. A. Monteiro, Gene regulatory networks reused to build novel traits. *BioEssays* **34**, 181–186
(2012).

47. A. Martin, *et al.*, Multiple recent co-options of Optix associated with novel traits in adaptive
butterfly wing radiations. *Evodevo* **5** (2014).

48. E. R Kandel, J. H. Schwartz, T. M. Jessell, S. A. Siegelbaum, A. J. Hudspeth. *Principles of*
Neural Science, 2012th Ed. (New York: McGraw Hill, 2000).

49. X. Wang, *et al.*, Genomic responses to selection for tame/aggressive behaviors in the silver
fox (Vulpes vulpes). *Proc. Natl. Acad. Sci. U. S. A.* **115**, 10398–10403 (2018).

50. N. I. Bloch, *et al.*, Early neurogenomic response associated with variation in guppy female
mate preference. *Nat. Ecol. Evol.* **2**, 1772–1781 (2018).

51. M. Yamaguchi, Role of regucalcin in brain calcium signaling. *Integr. Biol.* **4**, 825–837
(2012).

52. M. J. Berridge, Neuronal calcium signaling. *Neuron*, **21**, 13-26 (1998).

53. G. J. Bashaw, R. Klein, Signaling from axon guidance receptors. *Cold Spring Harb.*
Perspect. Biol. **2**, 1–17 (2010).

54. B. Prud'homme, N. Gompel, S. B. Carroll, Emerging principles of regulatory
evolution. *Proc. Natl. Acad. Sci. U. S. A.* ***104***, 8605-8612 (2007).

55. E. Preger-Ben Noon, *et al.*, Comprehensive Analysis of a cis-Regulatory Region Reveals
Pleiotropy in Enhancer Function. *Cell Rep.* **22**, 3021–3031 (2018).

56. J. Lewis, *et al.*, Parallel evolution of ancient, pleiotropic enhancers underlies butterfly wing
pattern mimicry. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 24174–24183 (2019).

57. W. Zhang, *et al.*, Comparative Transcriptomics Provides Insights into Reticulate and
Adaptive Evolution of a Butterfly Radiation. *Genome Biol. Evol.* **11**, 2963–2975 (2019).

58. A. Mazo-Vargas, *et al.*, Macroevolutionary shifts of WntA function potentiate butterfly
wing-pattern diversity. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 10701–10706 (2017).

59. C. Concha, *et al.*, Interplay between Developmental Flexibility and Determinism in the
Evolution of Mimetic Heliconius Wing Patterns. *Curr. Biol.* **29**, 3996-4009 (2019).

60. J. N. Weber, B. K. Peterson, H. E. Hoekstra, Discrete genetic modules are responsible for
complex burrow evolution in Peromyscus mice. *Nature* **493**, 402–405 (2013).\

61. J. Cande, P. Andolfatto, B. Prud'homme, D. L. Stern, N. Gompel, Evolution of Multiple
Additive Loci Caused Divergence between Drosophila yakuba and D. santomea in Wing
Rowing during Male Courtship. *PLoS One* **7**, 1–10 (2012).

62. C. S. McBride, *et al.*, Evolution of mosquito preference for humans linked to an odorant
receptor. *Nature* **515**, 222–227 (2014).

63. Y. Ding, A. Berrocal, T. Morita, K. D. Longden, D. L. Stern, Natural courtship song
variation caused by an intronic retroelement in an ion channel gene. *Nature* **536**, 329–332
(2016).

64. A. Bendesky, *et al.*, The genetic basis of parental care evolution in monogamous mice.
Nature **544**, 434–439 (2017).

65. P. C. Bürkner, Advanced Bayesian Multilevel Modeling with the R Package brms. *The R*
Journal, **10**, 395-411 (2018).

66. A. Vehtari, A. Gelman, J. Gabry, Practical Bayesian model evaluation using leave-one-out
cross-validation and WAIC. *Stat. Comput.* **27**, 1413–1432 (2017).

67. A. Dobin, *et al.*, STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21
(2013).

68. H. Li, *et al.*, The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
2078–2079 (2009).

69. S. Anders, P. T. Pyl, W. Huber, HTSeq-A Python framework to work with high-throughput
sequencing data. *Bioinformatics* **31**, 166–169 (2015).

70. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol.* **15**, 1–21 (2014).

71. S. H. Montgomery, J. E. Mank, Inferring regulatory change from gene expression: the
confounding effects of tissue scaling. *Mol. Ecol.* **25**, 5114–5128 (2016).
72. A. McKenna, *et al.*, The genome analysis toolkit: A MapReduce framework for analyzing
next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).
73. R. D. Finn, *et al.*, InterPro in 2017-beyond protein family and domain annotations. *Nucleic*
Acids Res. **45**, D190–D199 (2017).
74. H. Li, R. Durbin, Fast and accurate long-read alignment with Burrows-Wheeler transform.
Bioinformatics **26**, 589–595 (2010).
75. R. A. York, *et al.*, Behavior-dependent cis regulation reveals genes and pathways associated
with bower building in cichlid fishes. *Proc. Natl. Acad. Sci. U. S. A.* 115, 1081–1090 (2018).
76. P. Cingolani, *et al.*, A program for annotating and predicting the effects of single nucleotide
polymorphisms, SnpEff *Fly.* 6, 80–92 (2012).

Supporting information: Neural processing genes are linked to divergence in
visual mate preference in sympatric *Heliconius* butterflies.

Matteo Rossi[a,b], Timothy J. Thurman[b,c], Alexander E. Hausmann[a], Stephen H.
Montgomery[d], Riccardo Papa[b,e,f], Chris D. Jiggins[g] , W. Owen McMillan[b] & Richard M.
Merrill[a,b]

[a] Division of Evolutionary Biology, Ludwig Maximilian University of Munich, Grosshaderner Str. 2, 82152
Planegg-Martinsried, Germany
[b] Smithsonian Tropical Research Institute, Gamboa, 0843-03092, Panama
[c] Division of Biological Sciences, University of Montana, 32 Campus Dr., Missoula, MT 59812, US
[d] School of Biological Sciences, University of Bristol, 24 Tyndall Ave, Bristol BS8 1TQ, UK
[e] Department of Biology, University of Puerto Rico, Rio Piedras, San Juan 00925, Puerto Rico
[f] Molecular Sciences and Research Center, University of Puerto Rico, San Juan 00926, Puerto Rico
[g] Department of Zoology, University of Cambridge, Downing Street, Cambridge CB2 3EJ, UK

                                    Supporting methods and results

Mapping RNA-seq reads to the *Heliconius cydno* genome. To determine whether the *H.*

melpomene reference genome introduced mapping biases of RNA-seq reads, possibly

affecting differential expression estimates, we also mapped to a *H. cydno*

assembly/annotation. Generally, we found similar patterns of differential expression when

mapping to the two genomes. Since **i)** we observed an equal decrease (~ 40 %) of genes

showing 2-fold changes in *melpomene* and *cydno* when mapping to *H. cydno,* at every stage

(p-value=0.317 at adult stage, p-value=0.800 at 156h APF, p-value=0.897 at 60h APFP,

Fisher's Exact test, **Table S2**), and **ii)** this decrease was widespread throughout the genome,

we concluded that the *melpomene* reference genome did not bias differential gene expression

analyses.

Allele-specific expression in the introgression line. BC3 hybrids had different combinations

of chromosomes segregating for the *melpomene* alleles in a *cydno* background. Therefore, in

principle, we could not infer *cis-* or *trans-* gene regulatory effects genome-wide from the

profiles of allele specific expression (ASE) in these hybrids as for F1 hybrids, due to the

diverse trans-acting environments. However, previous analyses (comparing gene expression

levels between hybrids carrying *cyd/melp* vs. *cyd/cyd* regions on chromosomes other than 18)

imply that differential expression of the candidate genes seems to be driven by the *melpomene*

copy difference within the introgressed region on chromosome 18. Therefore, ASE analyses

of candidate genes in BC3 hybrids carrying *cyd/melp* alleles on chromosome 18 should

indicate whether the differences are due to cis- or trans-regulatory effects from within the

introgressed region (**Figure S6**).

In BC3 hybrids sampled at 156h APF and 60hAPF, the *melpomene* and *cydno* alleles of the

ionotropic glutamate receptor (*Grik2*) are expressed at very similar levels (at 156hAPF p

value=0.841, at 60hAPF p value=0.579, Wald test), suggesting trans-only regulatory effects at

these stages for *Grik2*. For *regucalcin1* we again had very few allele-informative read counts

in hybrids at 156h APF. Although there was a tendency towards up regulation of the

melpomene allele, there was no statistical significance to support this (p=0.174, Wald test).

We detected diffASE expression of *regucalcin2* only at 60h APF (p <0.001, Wald test), but at

this stage *regucalcin2* was not detected as differentially expressed between pure species.

Thus, although there is tentative evidence for *cis*-regulatory effects for species differences in

regucalcin expression during development, it is not conclusive.

Supporting figures

Figure S1. Ternary plots showing the number of 15-minute choice trials in which courtship
was initiated towards *melpomene, cydno* or both females for backcross-to-*cydno* males, with
different genotypes at the two QTLs retained in our model (on chromosome 1 and chromosome
18). Left ternary axis shows proportion of trials where courtship was initiated towards *H. cydno*
female only, bottom axis towards *H. melpomene* female only, and right axis towards both
female species. Each point corresponds to a male (and a 'jitter' function applied). Point size is
scaled to the number of trials in which the male showed a response. 95% credibility intervals
(CrIs) for all three proportions are shown as hexagons around predictors extrapolated from the
statistical model with the best fit, with lines projecting to corresponding values on the three
axes.

Figure S2. Results of comparative transcriptomic analyses between *melpomene* and *cydno* (in the imago) when mapping RNA-seq reads to the *H. melpomene* assembly/annotation (top), and to the *H. cydno* assembly/annotation (bottom), zooming in on the QTL region on chromosome 18. The *x*-axis represents physical position. Points correspond to individual genes, with the *y*-axis indicating the log_2(fold-change) for each comparison. The two horizontal dashed lines (at *y*-values of 1 and -1) indicate a 2-fold change in expression. Genes showing a significant 2-fold+ change in expression level between groups are highlighted in orange and blue, where orange indicates higher levels in *melpomene*, blue if in *cydno*. Genes detected as differentially expressed mapping to both *melpomene* and *cydno* genomes are labelled with gene names. dehydr.=2-oxoisovalerate dehydrogenase, OBP=odorant-binding protein.

Figure S3. Crossing design for producing backcross hybrids segregating at the QTL on chromosome 18. This introgression line was created by outcrossing a male hybrid to *H. cydno* females over three generations, selecting a hybrid male that showed a red band on the wing at each generation. This meant that these males carried one copy of the *H. melpomene* allele at the *optix* locus. We expected that, following recombination (which occurs in males), by the fourth generation we would remain with two types of individuals: either *cyd/melp* or *cyd/cyd* at the level of the *optix* region (which approximately corresponds to the region associated with male preference behavior).

Figure S4. Differential gene expression at the QTL region at pupal stages. Left: summary of the comparative transcriptomic analyses with stage, number of samples and chromosome 18 composition. Right: the corresponding results, zooming in on the QTL region on chromosome 18. The *x*-axis represents physical position. The QTL peak, and the rest of the QTL 1.5 LOD candidate region (from[28]) are shown in green and purple, respectively. Points correspond to individual genes, with the *y*-axis indicating the *log₂*(fold-change) for each comparison. The two horizontal dashed lines (at *y*-values of 1 and -1) indicate a 2-fold change in expression. Genes showing a significant 2-fold+ change in expression level between groups are highlighted in orange and blue, where orange indicates higher levels in *melpomene* or in the hybrids *cyd/melp* (blue if in *cydno* − hybrids *cyd/cyd*). Vertical dashed lines highlight those genes that are differentially expressed between *melpomene* and *cydno* AND between *cyd/melp* vs *cyd/cyd* individuals, at the same stage. One gene highlighted by a dashed fuchsia vertical line was excluded because it showed reversal of the fold change when mapping RNA-seq reads to the *H. cydno* genome.

Figure S5. A) Schematic representation of hybrid pupae (sampled at 156h APF) genome composition. Columns represent chromosomes, rows represent individuals, orange indicates *cyd/melp* regions, blue indicates *cyd/cyd* regions. **B)** Genome composition of (a subset of) BC3 hybrids. We calculated the fraction of SNPs and indels that each BC3 hybrid, one *cydno* and one F1 hybrid samples shared with *melpomene* and *cydno* samples, in non-overlapping 100kb windows. x-axes represent physical position (for each chromosome), y-axes fractions of shared variants with *melpomene* (in gold) and with *cydno* (in light blue). Matching variant fractions between BC3 hybrids and the F1 hybrid, indicating heterozygous regions, are highlighted with a gold bar underneath. Note that the general trend of higher number of variants shared with *H. cydno* in heterozygous regions is due to the fact that we inferred variants by mapping to the *H. melpomene* genome (and used variant sites only for this analysis).

A)

B)

Figure S6. Allele specific expression profiles of candidate genes at pupal stages. Points indicate the value, and bars the standard error, of the (base 2) logarithmic fold change in expression between parental species (horizontal) and the alleles in F1 hybrids (vertical), for candidate genes (as defined in the transcript-guided annotation). Dashed lines indicate the threshold for a 2-fold change in expression for the genes in the species (horizontal), and for the alleles in the hybrids (vertical).

Figure S7. Distribution of the (base 2) logarithmic fold change in allele expression. Colored bars indicate the number of genes showing a bias in expression for the *cydno* allele (in blue) and for the *melpomene* allele (in yellow). Values departing from 0 on the x-axis, indicate an increase in the fold change for the cydno allele (negative values) or for the *melpomene* allele (positive values), respectively.

1051
1052

Table S1. List of differentially expressed genes in species and hybrids comparisons.

1053
1054

A) QTL chromosome 1. Orange indicates genes up-regulated in *H. melpomene*, and blue those up-regulated in *H. cydno.*

1055
1056
1057
1058

#	Gene name (Hmel2.5)	Annotated function	Species comparison		
			60h APF	156h APF	Imago
1	HMEL002973g1	No match	✓	✓	✓
2	HMEL003796g1	Regulation of enolase protein 1	✓	✗	✓
3	HMEL011272g1	no match	✗	✓	✓
4	HMEL030024g1	Ribonuclease H superfamily	✓	✓	✓
5	HMEL030042g1	SWR1-complex protein 5	✗	✗	✓
6	HMEL030052g1	reverse transcriptase	✓	✓	✓
7	HMEL005260g1	unknown	✗	✓	✗
8	HMEL030040g1	No match	✗	✓	✗
9	HMEL030037g1	No match	✓	✓	✗
10	HMEL010076g1	Amino acid transporter	✓	✗	✗

B) QTL chromosome 18. Those genes found to be differentially expressed when also mapping to the *H. cydno* genome are highlighted in bold. Genes annotated as distinct but sharing the same number in the table (#) were later found to be single genes (see second paragraph of the Results section).

#	Gene name (Hmel2.5)	Annotated function	Species comparison			Hybrids comparison		
			60h APF	156h APF	Imago	60h APF	156h APF	Imago
1	**HMEL009992g1**	**No match**	✓	✗	✓	✗	✗	✓
1	**HMEL009992g4**	**Ionotropic glutamate receptor**	✓	✗	✓	✓	✓	✓
2	**HMEL009996g1**	**Gag-related protein**	✓	✓	✓	✗	✗	✗
3	HMEL034168g1	unknown	✗	✗	✓	✗	✗	✗
4	HMEL034173g1	SWR1-complex protein 5	✗	✗	✗	✗	✗	✗
5	HMEL034176g1	Aspartic peptidase	✓	✓	✓	✗	✗	✗
6	HMEL034184g1	No match	✗	✗	✓	✗	✗	✗
7	**HMEL034185g1**	**No match**	✓	✓	✓	✗	✗	✗
8	HMEL034187g1	Major facilitator superfamily (MFS) transporter	✓	✓	✓	✗	✓	✓
9	**HMEL003176**	**Odorant binding protein**	✗	✓	✓	✓	✗	✗
10	HMEL013551g1	2-oxoisovalerate dehydrogenase	✗	✗	✗	✓	✓	✗
10	**HMEL013551g2**	**2-oxoisovalerate dehydrogenase**	✗	✗	✓	✓	✗	✗
11	HMEL013551g4	SMP-30/regucalcin	✗	✗	✓	✗	✗	✗
12	**HMEL013552g1**	**SMP-30/regucalcin**	✗	✗	✗	✗	✓	✓
12	**HMEL034199g1**	**SMP-30/regucalcin**	✗	✗	✓	✗	✓	✓
13	HMEL014202g1	Catalase	✗	✗	✓	✗	✗	✗
14	**HMEL014202g3**	**Catalase**	✗	✓	✗	✗	✗	✗
15	**HMEL034201g1**	**No match**	✗	✗	✓	✗	✗	✗
16	**HMEL034205g1**	**No match**	✓	✗	✓	✗	✗	✗
17	HMEL034227g1	Ribonuclease H superfamily	✓	✓	✓	✗	✗	✗
18	HMEL034229g1	Endonuclease/exonuclease/phosphatase superfamily	✓	✓	✓	✗	✗	✓
19	HMEL034230g1	No match	✓	✓	✓	✗	✗	✗
20	**HMEL003863g1**	**Vacuolar protein sorting-associated (VPS) protein**	✓	✓	✗	✗	✗	✗
21	HMEL003863g3	No match	✓	✗	✗	✗	✗	✗
22	**HMEL006662g1**	**Serpin family protein**	✗	✓	✗	✗	✗	✗
23	**HMEL006663**	**Odorant binding protein**	✗	✓	✗	✗	✗	✗
24	**HMEL022553**	**Odorant binding protein**	✗	✓	✗	✗	✗	✗
25	HMEL001038g1	Monocarboxylate transporter	✓	✗	✗	✗	✗	✗
26	HMEL014190g1	unknown function	✓	✗	✗	✗	✗	✗
27	HMEL034236g1	No match	✓	✗	✗	✗	✗	✗
28	HMEL034189g1	PiggyBac transposable element-derived protein	✗	✗	✗	✗	✗	✓
29	HMEL034246g1	No match	✗	✗	✗	✗	✗	✓
30	HMEL034195g1	Gag-related protein	✗	✗	✗	✓	✓	✗

Table S2. Number of genes showing significant >2-fold change in expression, at different stages, mapping to the *melpomene* and to the *cydno* genomes. Note that the considerable reduction in the number of genes detected as differentially expressed when mapping to *H. cydno* is most likely a result of the lower quality/completeness of the *H. cydno* genome assembly.

Stage	Mapping to:	Up-regulated in *H. melpomene*	Up-regulated in *H. cydno*
Adult	*H. melpomene*	694	733
	H. cydno	390	451
156h APF	*H. melpomene*	837	667
	H. cydno	518	403
60h APF	*H. melpomene*	846	642
	H. cydno	490	376

1125 **Table S3.** Heterozygosity on the *Z*-chromosome. Heterozygosity is calculated as proportion of
1126 variants (SNPs and indels) which are heterozygous, in each sample, rounded at the second
1127 decimal place (note that variant sites were inferred having mapped to the *H. melpomene*
1128 genome).

1129

H. melpomene				*H. cydno*			
Males		Females		Males		Females	
ID	Het.	**ID**	Het.	**ID**	Het.	**ID**	Het.
Adults							
45	0.49	**53**	0.04	**57**	0.23	**50**	0.02
47	0.46	**78**	0.05	**82**	0.23	**51**	0.02
70	0.47	**80**	0.04	**98**	0.25	**58**	0.02
71	0.48	**128**	0.04	**99**	0.24	**67**	0.02
83	0.46	**218**	0.05			**68**	0.01
100	0.49					**81**	0.02
104	0.46					**84**	0.02
156h APF							
5	0.47	**6**	0.05	**4**	0.26	**13**	0.02
14	0.49	**18**	0.05	**8**	0.26	**21**	0.04
17	0.49	**24**	0.05	**142**	0.30	**30**	0.03
184	0.50	**150**	0.05	**151**	0.29	**137**	0.04
		220	0.06	**156**	0.29	**168**	0.04
60h APF							
92	0.48	**87**	0.06	**85**	0.26	**90**	0.02
97	0.49	**95**	0.05	**86**	0.27	**118**	0.02
115	0.47	**117**	0.05	**119**	0.27	**125**	0.02
		124	0.04	**146**	0.26	**144**	0.02
		149	0.06			**162**	0.02
		164	0.05			**200**	0.02
		208	0.06				

1130

Hybrids							
Males		Females		Males		Females	
ID	Het.	**ID**	Het.	**ID**	Het.	**ID**	Het.
F1 hybrids (adults)							
42	0.64	**56**	0.05				
49	0.64	**69**	0.04				
Introgression line -156h APF				**Introgression line – 60h APF**			
105	0.26	**108**	0.02	**152**	0.27	**161**	0.02
116	0.27	**123**	0.02	**193**	0.27	**165**	0.02
126	0.27	**136**	0.02	**198**	0.25	**179**	0.02
131	0.27	**139**	0.02	**199**	0.27	**187**	0.02
133	0.26	**140**	0.02	**201**	0.26	**188**	0.02
154	0.26	**166**	0.02	**212**	0.28	**192**	0.02
155	0.26			**215**	0.27	**197**	0.02
183	0.27					**209**	0.03
185	0.27					**214**	0.02
189	0.27					**224**	0.02

1131

1132

1133