Visual mate preference evolution during butterfly speciation is linked to neural processing genes

Matteo Rossi[1,2], Alexander E. Hausmann[1], Timothy J. Thurman[2,3], Stephen H. Montgomery[4], Riccardo Papa[2,5,6], Chris D. Jiggins[7], W. Owen McMillan[2] & Richard M. Merrill[1,2*]

1. Division of Evolutionary Biology, LMU, Munich, Germany; 2. Smithsonian Tropical Research Institute, Panama City, Panama; 3. Division of Biological Sciences, University of Montana, USA; 4. School of Biological Sciences, University of Bristol, Bristol, UK; 5. Department of Biology, University of Puerto Rico, San Juan, Puerto Rico; 6. Molecular Sciences and Research Center, University of Puerto Rico, San Juan, Puerto Rico; 7. Department of Zoology, University of Cambridge, Cambridge UK. *corresponding author**: merrill@bio.lmu.de

Many animal species remain separate not because they fail to produce viable hybrids, but because they "choose" not to mate. However, we still know very little of the genetic mechanisms underlying changes in these mate preference behaviours. *Heliconius* butterflies display bright warning patterns, which they also use to recognize conspecifics. Here, we couple QTL for divergence in visual preference behaviours with population genomic and gene expression analyses of neural tissue (central brain, optic lobes and ommatidia) across development in two sympatric *Heliconius* species. Within a region containing 200 genes, we identify five genes that are strongly associated with divergent visual preferences. Three of these have previously been implicated in key components of neural signalling (specifically an *ionotropic glutamate receptor* and two *regucalcins*), and overall our candidates suggest shifts in behaviour involve changes in visual integration or processing. This would allow preference evolution without altering perception of the wider environment.

The evolution and maintenance of new animal species often relies on the emergence of divergent mating preferences[1,2]. Changes in sensory perception or other neural systems must underlie differences in innate behaviours between species, and will ultimately have a genetic

basis. However, although the significance of behavioural barriers for speciation has been

recognized since the Modern Synthesis[3], we know almost nothing of the genes underlying

changes in mating preferences, or variation in behaviours across natural populations more

broadly[4,5]. Identifying these genes will provide an important route towards understanding how

behavioural differences are generated, both during development and across evolutionary time.

Previous studies of isolating preference behaviours have largely been limited to the

identification of causal genomic regions, which almost invariably contain many genes[6,7,8,9].

Only a handful of studies have identified likely candidate genes that contribute to species

behavioural preferences. These are largely limited to chemosensory-guided mating

preferences[10,11,12,13], and have identified changes at chemoreceptor genes. To our knowledge,

only two studies – in incipient fish species – have identified candidates for visual preference

evolution, albeit indirectly, both suggesting a role for sensory perception mediated by changes

in the peripheral visual system[14,15]. Whether or not visual preference evolution generally

involves shifts at the sensory periphery, or in downstream processing, remains unknown.

The closely related species *Heliconius melpomene* and *H. cydno* differ in warning

patterns, which are both under disruptive selection for mimicry[16] and are important mating

cues [17]. In central Panama, *H. melpomene* shares the black, red and yellow pattern of its local

Heliconius erato co-mimic. In contrast, *H. cydno* mimics the black and white patterns of *H.*

sapho. The two species remain separate largely due to strong assortative mating[18]. Visual

preferences for divergent patterns are particularly apparent in males, which strongly prefer to

court conspecific females[17,19,20]. Differences in warning pattern between *melpomene* and

cydno are largely due to expression differences in just three genes, specifically *optix*[21],

WntA[22] and *cortex*[23].

Quantitative trait locus (QTL) mapping of *H. melpomene* and *H. cydno* has revealed

three genomic regions of major effect that influence the relative time males spend courting

red *melpomene* or white *cydno* females[20]. Notably, the best supported QTL was in the same

genomic region as *optix*, the gene responsible for presence of the red colour pattern elements

in *H. melpomene*[21]. Genetic linkage will facilitate speciation by impeding the breakdown of

genetic associations between ecological and mating traits[24]. Nevertheless, this QTL, and its

associated candidate region, contain hundreds of genes, and the exact genes responsible for

differences in preference behaviour are not known.

Here, we first confirm that the behavioural QTLs identified previously are associated

with variation in male courtship initiation. We then identify genes within the major QTL,

which were differentially expressed in the neural tissue (central brain, optic lobes and

ommatidia) of *H. melpomene* and *H. cydno*, or have protein coding changes predicted to alter

protein function. Out of 200 genes within the QTL region, we identify just five candidates

likely to underlie assortative mating behaviours.

Results

Chromosome 18 is associated with differences in courtship initiation.

Our previous results reveal that QTLs on chromosomes 1, 17 and 18 influence the relative

time hybrid males spend courting red *melpomene* or white *cydno* females[20]. However, the

time males spend courting a particular female might depend not only on male attraction, but

on the female's response (and in turn his response to her behaviour). To confirm that these

previously reported QTLs influence male approach behaviours (as opposed to other traits that

may influence courtship, for example male morphology[25]), we reanalysed our previous data,

this time explicitly considering whether males initiated courtship towards *melpomene*, *cydno*

or both types of female during choice trials. Consistent with our previous analyses[20], we

found that F1 and backcross-to-*melpomene* prefer to court *melpomene* females, whereas

courtship initiation behaviours segregate in the backcrosses to *cydno* (Figure 1). Notably,

backcrosses-to-*cydno* males heterozygous at the QTL on chromosome 18 (*i.e.* with a

melpomene allele derived from the F1 father) initiated courtship towards *melpomene* females

more frequently than males homozygous for the *cydno* allele (Figure 1, bottom left; *n* = 139,

ΔELPD: -10.9 (S.E.±5.1), *i.e.* a change of 2.14 SE units). Together with previous evidence

that male hybrids bearing *melpomene* alleles at *optix* prefer to court the artificial models of

melpomene females over those of *cydno*[26], these results suggest that the QTL on chromosome

18 harbours genes for visual attraction behaviours towards females with the red pattern.

Consequently, we focused our subsequent analyses on this QTL on chromosome 18 (and also

because tight linkage of *optix* allowed us to track the alleles at preference-colour locus in

hybrid crosses). The QTL on chromosome 1 was also retained in our model of initiation

behaviours (Supplementary figure 1; *n* = 139, ΔELPD = -13.6 (SE±5.7), *i.e.* a change of 2.34

SE units), in contrast to the QTL on chromosome 17 which was not retained (*n* = 139,

ΔELPD = -2.1 (S.E.±3.0)). Results for the QTL on chromosome 1 are reported in the

supplementary materials (Supplementary table 1).

27 *genes* within the major QTL are differentially expressed in the brains and eyes of *H.*

***cydno* and *H. melpomene*.**

We hypothesized that changes in gene regulation that determine differences in visual mate

preference behaviours might occur during pupal development (for instance, during visual

circuit assembly) or in the imago, and must involve changes in the peripheral and/or central

nervous system[27]. Therefore, we generated RNA-seq libraries for combined eye and brain

tissue, across two pupal stages (around the time of ommochrome pigment deposit and half-

way through pupal development) and one adult stage, for *H. melpomene* and *H. cydno* and

compared their gene expression levels. We found considerable differential expression at the

QTL on chromosome 18 (the QTL spans 2.75 Mb, and contains 200 genes). We identified 27

genes within the QTL region that show differential expression between *melpomene* and

cydno, in at least one of the three developmental stages. These were mostly located within the

QTL peak (*i.e.* the genomic region with strongest statistical association with male preference)

or in close proximity to *optix* (Figure 2). The same genes were frequently differentially

expressed across development (Supplementary table 1), with 11 genes being differentially

expressed in more than one stage.

The genomic region between the start of chromosome 18 and *optix* (comprising the

QTL peak) is highly divergent between *melpomene* and *cydno*[28], and divergent coding

sequences within this region could also introduce mapping biases of RNA-seq reads. To

account for this, we repeated the analysis having mapped to both the *H. melpomene* reference

genome[29] and to a *H. cydno* genome[30]. Generally, we found similar patterns of differential

expression when mapping to the *H. cydno* genome (Supplementary figure 2, Supplementary

table 2). Nevertheless, in subsequent analyses we excluded two genes, HMEL034187g1 and

HMEL034229g1, which showed reversal of the fold change or did not show differential

expression when mapping to the *H. cydno* genome respectively.

A *regucalcin* and an *ionotropic glutamate receptor* are upregulated in both *H. melpomene*

and F1 hybrid males.

Our previous behavioural experiments suggest that the alleles for the *melpomene* behaviour

are dominant over the *cydno* alleles[20,26] (Figure 1). Given this pattern of dominance, we

predicted that genes underlying variation in male preference to be up- or down- regulated in

the brains of both *melpomene* and first generation (F1) hybrid males, with respect to *cydno*.

Of the putative genes differentially expressed between *cydno* and *melpomene* reported above,

only four, within the QTL candidate region, were differentially expressed between the F1

hybrids and *cydno* (Figure 2). These included two *regucalcins* (also called *senescence marker*

proteins-30: HMEL013552g1, HMEL034199g1), an *ionotropic glutamate receptor*

(HMEL009992g4), which is a putative ortholog of *Grik2*, and one gene with no annotated

function (HMEL009992g1). We obtained the same results regardless of whether we

considered both males and females together, or males alone. Further inspection of spliced

mRNA-reads indicated that the two annotated *regucalcins* were in fact a single gene (from

now on referred to as *regucalcin2*). This was also the case for the *ionotropic glutamate*

receptor and the gene with no annotated function (from now on referred to as *Grik2*).

Differential expression of *Grik2* in adults is likely due to cis-regulatory effects.

To determine whether differences in gene expression levels between parental species were

due to *cis*- or *trans*-regulatory changes, we conducted allele specific expression (ASE)

analyses in adult F1 hybrids. In F1 hybrids, both parental alleles are exposed to the same

trans-environment, and consequently *trans*-acting factors will act on alleles derived from each

species equally (unless there is a change in the cis-regulatory regions of the respective

alleles). Therefore, differences in allele specific expression indicate changes in *cis*-regulatory

regions[31]. Both candidate genes (*Grik2* and *regucalcin2*) had a very low number of SNPs that

could differentiate the *melpomene* and *cydno* allele (using both gene models of the Hmel2.5

annotation and RABT annotation) and few reads mapped to these SNPs. Nevertheless, for

Grik2, the *melpomene* allele was significantly more highly expressed relative to the *cydno*

allele (p=0.017, Wald test), suggesting cis-regulatory effects (Figure 3). For *regucalcin2,*

although there was a tendency towards up-regulation of the *melpomene* allele, consistent with

cis-regulation, we did not have sufficient power-to rule out *trans*-only regulatory effects

(Figure 3; p=0.108, Wald test).

***Grik2* is differentially expressed in hybrids that essentially differ only for allelic**

composition at the behavioural QTL region.

In order to study the specific effects that *melpomene* derived alleles at the QTL on

chromosome 18 had on gene-expression, we introgressed this region into a *cydno* background

through multiple backcrosses (crossing design in Supplementary figure 3). We wanted to

investigate whether differences at this QTL regulated expression of any specific genetic

pathway during development, and more generally what changes in genome-wide transcription

were observed in hybrids differing (mostly) just at this QTL region.

Notably, in these third-generation backcross hybrid comparisons, across the entire

genome only 23 and 29 genes were differentially expressed (at 156h after pupal formation

(APF) and at 60hAPF, respectively). Of these, 20 and 19 genes (at 156hAPF and at 60hAPF,

respectively) were located on chromosome 18, indicating that gene expression differences in

these comparisons were mostly restricted to the preference-colour region on chromosome 18,

segregating for *cyd/melp* or *cyd/cyd* alleles. No genetic pathway was enriched for gene

expression differences between these hybrids at either pupal stage (PANTHER enrichment

test[32]), suggesting that overall this QTL harbours a few, modular changes in gene regulation

in the developing brain/eyes of *cydno* and *melpomene*. *Grik2* was the only gene detected as

differentially expressed between species and hybrids at these pupal stages (Supplementary

figure 4).

To verify that differential expression of candidate genes at the QTL region is driven

by *melpomene* alleles on chromosome 18 and not by other *melpomene* alleles at *trans*-acting

genes on other chromosomes, we compared gene expression levels between hybrids carrying

cyd/melp vs. *cyd/cyd* regions on chromosomes chr1, chr4 and chr15, chr20 (Supplementary

figure 5A). In these comparisons, there was no signal of differential expression on

chromosome 18. This supports the *cis*-regulatory activity of the *melpomene* allele of

candidate genes on chromosome 18. To test this further, we conducted another allele specific

expression study in the BC3 hybrids, which suggested *trans*-regulatory effects for *Grik2* at

these pupal stages, but were less conclusive with regard to *regucalcin2* (Supplementary figure

6). Since causal gene/s might exert an effect on behaviour due to their action during

development or in adult form, and this action might in turn be differently (cis- vs trans-)

regulated, we still considered both genes as strong candidates.

4 genes with protein-coding substitutions within the QTL candidate region have

predicted effects on protein function.

Because shifts in behavioural phenotypes could be due to changes in protein-coding regions,

we additionally considered protein-coding substitutions between *melpomene* and *cydno*.

Overall, we found 152 protein-coding substitutions, spanning 54 of the 200 genes across the

entire QTL candidate region. We then studied whether these variants were predicted to have

non-neutral effects on protein function with PROVEAN[33]. The PROVEAN algorithm predicts

the functional effect of protein sequence variations based on how they affect alignments to

different homologous protein sequences. We found 4 genes with such predicted effects

(PROVEAN score < -2.5): Specifically, a *WD40*-repeat domain containing protein

(HMEL013551g3), a *cysteine protease* (HMEL009684g2), a *MORN* motif containing protein

(HMEL006660g1), and another *regucalcin* (HMEL013551g4) adjacent to, but distinct from,

that found to be differentially expressed above (from now on referred to as *regucalcin1*).

Candidate genes occur in regions with reduced gene flow.

Of our six candidate genes for preference behaviours that contribute to reproductive isolation

between *H. cydno* and *H. melpomene* (*regucalcin2*, *Grik2* and the four genes with protein

coding modifications), five are found within the QTL peak (Figure 4). Genetic changes

causing reproductive isolation between populations are expected to reduce localized gene

flow in their genomes. Therefore, we compared the position of our candidate genes to

estimated levels of admixture proportions (f_d)[34] between *H. melpomene* and *H. cydno* across

the QTL candidate region[35]. We found that all candidate genes were located in genomic

regions with low f_d values (Figure 4), suggesting localized resistance to gene flow between

melpomene and *cydno* at these genes and their putative *cis*-regulatory regions.

Discussion

Behavioural isolation is frequently implicated in the formation of new species, and involves the correlated evolution of both mating cues and mating preference. Here we have analysed a genomic region in a pair of closely related sympatric butterflies, *H. cydno* and *H. melpomene*, that contains genes for divergence in both an ecologically relevant mating cue and the corresponding preference. Physical linkage between ecological and mating traits will facilitate speciation by allowing different barriers to act in concert to restrict gene flow[36,37]. Although the genes underlying changes in the warning pattern cue in *Heliconius* are well characterized[21,22,23,38] (e.g. *optix*), those underlying the corresponding shift in behaviour have not previously been identified[20,39,40]. We have pinpointed a small number of genes that fall within the QTL peak, which show either expression (*regucalcin2* and *Grik2*) or protein coding differences (HMEL013551g3, HMEL009684g2, HMEL006660g1, and *regucalcin1*) and fall within a region of reduced admixture, that are strong candidates for modulating mating behaviour.

Two broad neural mechanisms could underlie the evolution of divergent visual preferences, involving changes in either i) detection at the sensory periphery or ii) the processing and/or integration of visual information. Although *H. melpomene* and *H. cydno* have the same retinal mosaics/class of photoreceptors[41], spectral sensitivity in the *Heliconius* eyes could be altered by filtering pigments[42], or other physiological processes taking place at the photoreceptors/sensory periphery, eventually shifting sensitivity towards different wavelengths (and possibly colour patterns). It has previously been hypothesized that the gene regulatory networks for ommochrome deposition in the *Heliconius* eyes might have been co-opted in the wings[43], where *optix* plays a central role, and therefore that *optix* might play a role in eye pigmentation in *Heliconius*. However, the protein product of *optix* has not been detected in pupal or adult retinas of various *Heliconius* species tested[44], and therefore has no obvious link to ommochrome deposition in the eyes. More generally, the underlying

evolutionary mechanism is unlikely to involve detection at photoreceptors, as this would

probably have a broad effect on downstream processing[2] and alter the visual perception of the

animal's wider environment.

The second mechanism, involving changes in the processing, and/or integration, of

visual information, could act through an alteration of neuronal activity or connectivity. For

instance, different levels of gene expression in conserved neural circuits between *melpomene*

and *cydno* may affect overall synaptic weighting and determine whether a signal (e.g. colour

and motion) elicits a motor pattern (response towards a female) or not. Consistent with this

scenario, the composition of ionotropic receptors at post-synapses is a key modulator of

synaptic transmission[45], implicating *Grik2*. Interestingly, differential expression of ionotropic

glutamate receptors is also associated with variation in social and aggressive behaviours in

vertebrates[46,47]. *Regucalcins* are involved in calcium signalling[48], which regulates synaptic

excitability and plasticity[49], and has an important role in axon guidance[50] (albeit alongside

additional roles across a broad range of biological processes), making the two *regucalcins* we

identify strong candidates for behaviour.

Changes in the regulation of genes with pleiotropic effects are likely to be less

detrimental compared to changes in their protein-coding sequences[51] (although emerging

evidence has begun to suggest that enhancer/repressor elements may be more pleiotropic than

previously thought[52,53]). Furthermore, there is considerable evolutionary potential in the co-

option of transcription factors/networks[51] that regulate neural patterning or neuron-type

activity, possibly resulting in novel adaptive expression patterns. In line with this,

Regucalcin2 and *Grik2,* which are differentially expressed in the eyes and brain in both our

species and hybrid comparisons, are likely to be involved in multi-functional processes, such

as calcium signalling and ion transport, and likely have pleiotropic alleles. We also found

evidence of cis-regulatory effects for both genes (albeit not significant for *regucalcin2*),

which would be required of the causal genetic change within the QTL, if it were to be in gene

regulation.

Despite expectations that non-coding, regulatory loci may provide a flexible route to

divergent mating preferences, we also found substitutions in coding regions at the QTL,

which are predicted to have an effect on protein functioning and therefore remain strong

candidates. These genes include *regucalcin1*, which is distinct from, but located next to,

regucalcin2 (which is differentially expressed). Notably, the eye transcript of *regucalcin1* was

recently characterized as fast-evolving across *Heliconius* species[54]. Other candidates include a

cysteine protease, which functions in protein degradation, and might be linked to behaviour

for example through degradation of neurotransmitters, a *MORN* motif containing protein

(function unknown), and a *WD40* containing protein. *WD-repeat* containing proteins have

been implicated in a wide array of functions ranging from signal transduction to apoptosis

(https://www.ebi.ac.uk/interpro).

Although preference for red colouration and the *optix* gene are tightly linked, we find

no evidence that *optix* is differentially expressed in the eyes or brains of our two species. It is

also not located within the QTL peak (and it contains no non-synonymous changes in protein

coding regions[21]). It seems unlikely therefore that changes in cue and preference are

pleiotropic effects of the same allele. More generally, although we have pinpointed the

strongest candidates yet identified for assortative mating behaviours in *Heliconius*, it is

possible that actual causal changes in gene regulation are restricted to developmental stages

other those sampled, or restricted to a few neuronal populations not detected with

transcriptomic data from eyes and whole brain tissue. Nonetheless, by sampling at two pupal

stages (around the time of *optix* expression/ommochrome pigment deposit in the wing/eye and

halfway through pupal development) and at the adult stage, we should have captured

important transitions for the behavioural programming of the two species.

Work in the past decade has shown that complex innate behavioural differences

between species can be encoded in relatively few genetic modules[55,56], but very few

studies[57,58,59] have identified specific genes underlying behavioural evolution. In particular,

traditional laboratory organisms continue to provide important insights into the evolution and

genetics of behaviour[27,58,60], however, comparative approaches are required to determine if

developmental principles can be broadly applied, and also to incorporate a wider range of

phenotypic variation and sensory modalities. The challenge now is to increase the resolution

of studies in non-traditional systems, in order to link individual genetic elements to

behaviours, *and* the sensory and/or neurological structures through which they are mediated.

In this light, we have identified a small handful of strong candidate genes associated with the

evolution of visual mate preference behaviours in *Heliconius*. These genes are in tight

physical linkage with the locus for the corresponding shifts in an ecologically relevant mating

cue, providing an important opportunity to investigate the build-up of genetic barriers crucial

to speciation. The candidate genes identified seem more likely to alter visual processing or

integration, rather than detection at photoreceptors, consistent with permitting changes in

mate preference without altering perception of the animal's wider environment.

Materials and Methods

Courtship initiation analyses. Butterfly rearing, crossing design and genotyping are

described in detail elsewhere[20]. In brief, we assayed male preference behaviours for *H.*

melpomene, *H. cydno,* their first generation (F1) hybrids and backcross to hybrids to both

parental species in standardized choice trials. Males were introduced into outdoor

experimental cages (1x1x2m) with a virgin female of each species and courtship behaviours

recorded. Whenever possible, trials were repeated for each male (median = 5 trials). To

determine whether previously identified QTLs for courtship time contribute to variation in

courtship *initiation* behaviours, we performed a *post-hoc* analysis using categorical models in

a Bayesian framework with a multinomial error structure, using the R package *brms*. All

models were run under default priors (non- or very weakly informative). In contrast to our

previous analysis[20], in which we considered the number of minutes (*i.e.* time) for which

courtship was directed towards *H. cydno* or *H. melpomene* females, here the response variable

was number of trials in which male courtship was *initiated* towards *H. cydno* females only, *H.*

melpomene females only, or both female types (hereafter referred to as "*initiation*"). Across

males the median number of trials with a response was 3. Using backcross-to-*cydno* males

only, we fitted *initiation* as a response variable to genotype (*cyd*/*cyd* or *cyd*/*melp*) at each

QTL, which were included as separate fixed effects. Individual ID was fitted as random

factor. To test the effect of each QTL on male *initiation*, we compared the saturated model

incorporating all three QTL with reduced models excluding each QTL in turn, using

approximate leave-one-out (LOO) cross-validation[61] as implemented in *brms*, and based on

expected log pointwise predictive density (ELPD). Normal distribution of ELPD can be a

straightforward approximation given our large samples sizes (n=139)[61]. Therefore, we

considered an absolute value of ELPD greater than 1.96 units of its standard error as

indicative of the reduced model being less-informative than the saturated model (95%

confidence). Males that did not initiate courtship to any female across trials were excluded

from analyses, resulting in a dataset of 139 males, from a total of 146 backcross males for

which we had genotype data. Finally, we extracted predictors and credibility intervals for

backcross males with differing genotypes from the minimum adequate model. Credibility

intervals for *H. melpomene*, *H. cydno,* F1 hybrid and backcross to *melpomene* males

displayed in Figure 1 were generated following the same procedures. Raw data and analysis

code are available in the following github repository:

https://github.com/SpeciationBehaviour/neural_genes_heliconius.git

Butterfly collection, rearing and crossing design for expression analyses. Wild *H. melpomene rosina* and *H. cydno chioneus* individuals were caught along Pipeline Road near Gamboa, Panama, in the Soberania National Park, and used to establish stocks at the Smithsonian Tropical Research Institute insectaries in Gamboa. Butterflies were reared in common garden conditions, in 2x2x2m cages, and provided with fresh *Psiguria* flowers and 10% sugar solution. Larvae were reared on fresh *Passiflora* shoots/leaves until pupation. *H.* cydno, *H. melpomene* and hybrid individuals used for RNA-seq (see below) were reared concurrently and under the same conditions. F1 hybrids were obtained by crossing a wild-caught *H. m. rosina* male to an insectary-bred virgin *H. c. chioneus* female.

The introgression line was generated by outcrossing a hybrid male with a red forewing band (crossing design shown in Supplementary figure 3) to virgin *H. cydno* females, over three generations. The peak of the behavioural QTL reported previously[20] on chromosome 18 (at 0cM) is in very tight linkage with the *optix* colour pattern locus (at 1.2cM), which controls for the presence and absence of the red forewing band seen in *H. melpomene rosina*. Presence of the red forewing band is dominant over its absence so that segregation of the red band can be used to infer genotype at the *optix* locus. Specifically, hybrid individuals with a red forewing band are heterozygotes for *H. melpomene/H. cydno* alleles at the *optix* locus, whereas individuals lacking the red band are homozygous for the *cydno* allele. Due to the tight linkage we expected little recombination between *optix* and QTL peak even after three generations of introgression, allowing us to infer genotype at the preference-*optix* locus (which we confirmed with genetic data, see below).

Tissue dissection, RNA extraction and mRNA sequencing. Eye (ommatidia and retinal membrane) and brain tissue (central brain and optic lobes) were dissected out of the head capsule in cold (4 ºC) 0.01M PBS solution, at two pupal stages: 60 hours after pupal formation (60h APF) and 156h APF; and in adults aged 9 - 13 days. We sampled adults at

around 10 days of age because by this stage males are mature and frequently court females[62].

Adult males and females sampled were sexually naive. We decided to sample at 60h APF

because this is the developmental stage at which *optix* is expressed in the wing, so we

hypothesized that it might had also been when *optix* is expressed in the brain. We sampled at

156h APF as a putative stage halfway through pupal development, and at this stage most of

the major neural connections have just been established in the *Heliconius* brain (Stephen

Montgomery, unpublished data).

Tissues were stored in RNAlater at 4 ºC for 24 hours, and subsequently at -20 ºC, until

RNA extraction. Total RNA was extracted using TRIzol Reagent (Thermo Fisher, Waltham,

MA, USA) and a RNeasy Mini kit (Qiagen, Valencia, CA, USA). Samples were treated with

DNase I (Ambion, Darmstadt, Germany). Integrity of total RNA was checked either on an

agarose gel or using an Agilent Bioanalyzer 2100 (Agilent Technologies, Santa Clara, CA,

USA). RNA concentration was measured on a Nanodrop spectrophotomer. Illumina TruSeq

RNA-seq libraries were prepared and sequenced at Edinburgh Genomics (Edinburgh, UK)

with 100 bp paired-end reads. To avoid lane effects the distribution of the species samples

was randomized on the sequencing platform. More detailed information about individuals and

sequencing yields can be found in the Supplementary dataset.

RNA-seq read mapping and differential gene expression analyses. After a quality control

of RNA-seq reads with FastQC, we trimmed adaptor and low-quality bases using TrimGalore

v.0.4.4 (https://www.bioinformatics.babraham.ac.uk/projects/). RNA-seq reads were mapped

to the *H. melpomene* 2.5 genome[29]/annotation[63] using STAR v.2.4.2a[64] in 2-pass mode. We

only kept reads that mapped in 'proper pairs' using Samtools[65]. The number of reads mapping

to each gene were estimated with HTseq v. 0.9.1[66] with model "union", thus excluding

ambiguously mapped reads. Differential gene expression analyses between species/hybrids

were conducted in DESeq2[67]. We considered only those genes showing a 2-fold change in

expression level, and at adjusted (false discovery rate 5%) p-values < 0.05, to be differentially

expressed, to exclude expression differences caused by known differences in brain

morphology[68] (Montgomery et al., in prep).

Sexing pupae. In all DESeq2 analyses, sex was included as a random factor. To sex pupae,

we first marked duplicate RNA mapped reads with Picard

(https://broadinstitute.github.io/picard/), and used GATK 3.8[69] to split uniquely mapped reads

into exon segments and trim sequences overhanging the intronic regions. We then used

Haplotype Caller on each individual, using calling and filtering parameters according to the

GATK Best Practices for variant calling on RNA-seq data. The sex of pupal samples was

inferred from the proportion of heterozygous (biallelic) SNPs using the R package SNPstats.

Males (ZZ) were expected to have >> 0% heterozygous sites, whereas females (ZW) to have

0%. Z-linked heterozygosity of the pupal samples (Supplementary table 3) were in line with

expectations (either ~ 0 for females or an order of magnitude higher for males), and matched

heterozygosity of either adult males or females, for which the sex was determined from

external morphology.

Inference of gene function and transcript-based annotation. Biological functions of

annotated genes were inferred with InterProScan v5[70], using the corresponding Hmel2.5

predicted protein sequences. InterProScan uses different databases like InterPro, Pfam,

PANTHER, and others, to infer functional protein domains and motifs (based on homology).

To study whether specific biological functions were enriched among genes showing

differential expression among hybrid types, we conducted the PANTHER enrichment test[38]

(with Bonferroni correction for multiple testing) using *Drosophila melanogaster* as the

reference gene function database.

Upon detailed inspection of the mapping coverage of spliced RNA-seq reads to the

Hmel2.5 gene annotation, we noticed that some gene models were fragmented, namely, a few

exons that appeared to be spliced together were incorrectly considered distinct genes. To

check that this did not introduced inaccuracies in our differential gene expression analyses,

we re-annotated the *melpomene* genome using the Cufflinks reference annotation-based

transcript (RABT) assembly tool[71] We used the transcriptomic data from both *melpomene* and

cydno to reannotate the *melpomene* genome, separately for every developmental stage, and

reconducted the differential gene expression analyses in DESeq2 as described above.

Repeating all comparative transcriptomic analyses using these new annotations (where exons

were correctly considered as part of single genes), we confirmed that both *regucalcin2* and

Grik2 were differentially expressed in both species and hybrids comparisons.

Inference of BC3 hybrids genome composition. In order to perform comparative

transcriptomic analyses between third-generation backcross hybrids (BC3) segregating at the

QTL on chromosome 18 (crossing design in Supplementary figure 3), we first determined

which genomic regions in these hybrids were heterozygous (*cyd/melp*) or homozygous

(*cyd/cyd*). For this, we inferred variants from RNA-seq reads for each BC3 hybrid

(individually as above), and from the combined *melpomene* and *cydno* samples. For the

species, we used HaplotypeCaller[69] on RNA-seq samples from all developmental stages of

either species, to produce individual genomic records (gVCF), and then jointly genotyped

melpomene and *cydno* gVCFs (separately for the two species) using genotypeGVCFs with

default parameters. Genotype calls were filtered for quality by depth (QD) > 2, strand bias

(FS) < 30 and allele depth (DP) > 4. For further analyses we kept *biallelic* genotypes only.

We then used the *intersect* function of *bcftools*[65] to infer variants exclusive to the *cydno* and

to the *melpomene* samples.

We calculated the fraction of variants that each BC3 hybrid individual shared with the

melpomene and with the *cydno* samples, in non-overlapping 100kb windows. We compared

these to the fraction of variants that a F1 hybrid and a *H. cydno* individual (not included in the

combined genotyping of the *cydno* samples), shared with the same species samples, and found

that they matched either one of them, indicating heterozygous (*cyd/melp*) or homozygous

(*cyd*/*cyd*) regions (Supplementary figure 5B). In this analysis, we considered only those

100kb windows where BC3 hybrids/F1 hybrid/*H. cydno* individuals shared more than 30

variants with the *melpomene*/*cydno* samples.

To corroborate our findings, we repeated the same type of analysis, this time inferring

species-specific variants for *melpomene* and *cydno* using 10 *H. melpomene rosina* and 10 *H.*

cydno chioneus genome resequencing samples. Variant calling files (vcf) were retrieved from

Martin et al[35]. We considered only *biallelic* genotype calls that had $10 < DP < 100$ and

genotype quality (GQ) > 30. With this analysis we found the same heterozygous and

homozygous regions in BC3 hybrids**.**

The size and number of the introgressed regions were in line with expectations about

3rd generation backcross hybrids following our crossing design: segregating at the level of

chromosome 18 and at four other chromosomes. For the BC3 hybrids sampled at 156 hours

after pupal formation (APF) we had 6 *cyd/melp* and 10 *cyd*/*cyd* at the QTL region on

chromosome 18 (Supplementary figure 5A), for those at 60h APF, 8 *cyd/melp* and 9 *cyd*/*cyd*

hybrids at the same region.

Allele-specific expression (ASE) in hybrids. In order to conduct ASE analyses we first

identified species specific variants, fixed in either *melpomene* and *cydno*. For this, we took

the quality filtered variants inferred from the species genome resequencing data, and assigned

those genotype calls in *cydno* and *melpomene* for which allele frequency (AF) was > 0.9 as

homozygous (we did not consider indels in this analysis). We then used *bcftools intersect*[65] to

get only those variants for which *cydno* and *melpomene* had opposite alleles.

At the same time, we called variants from RNA-seq reads of F1 hybrid individuals,

again according to the GATK Best Practices (with the exception of parameters -window 35 -

cluster 3, to increase SNPs density), and selected only heterozygous SNPs in F1s that matched

the species-specific variants. Finally, we used GATK's ASEReadCounter[69], with default

parameters, to count RNA reads in the F1 hybrids (and later on in BC3 hybrids) that mapped

to either the *cydno* or the *melpomene* allele. We summed all reads mapping to either the *cydno*

or *melpomene* allele/variant within the same gene (both for gene models of the Hmel2.5 gene

annotation and for the Cufflinks annotation we assembled previously). To test for allele

specific expression (diffASE) we fitted the model "~0 + individual + allele" in DESeq2[67],

setting library size factors to 1 (thus not normalizing between samples, as the test for diffASE

is conducted within individuals). We only considered those alleles showing at least a 2-fold

change in expression and $p < 0.05$, as differentially expressed.

In order to check that there were no biases in alleles assignment to one of the two

species, we analyzed the ratios of the species alleles, for every gene, and checked that they

were not systematically biased to either one of the two species. The $\log_2$ fold-changes of the

species alleles were centered around 0, suggesting no obvious bias in alleles assignment[72]

(Supplementary figure 7).

Protein-coding substitutions and predicted effects on protein-function. We inferred fixed

variants in protein-coding regions from the combined *melpomene* and *cydno* RNA samples in

order to include variants from genes for which we detected expression in the brain/eyes across

the 3 stages. We took the quality filtered variants called from the joint genotyping of RNA-

seq data of *cydno* and *melpomene* (from all stages), and selected those genotype calls for

which allele frequency (AF) > 0.8, and where the allelic variant was present in at least 7

individuals of the ~30 samples (for each species). We retained those substitutions/indels

validated with the genome resequencing data. For this, of the genotype calls found in RNA

reads from brain/eyes of different stages, we kept only those that were also called in at least 8

of the 10 genome resequencing samples of each species. We considered this overlapping set

of variants as being fixed in *H. melpomene rosina* or *H. cydno chioneus*. Following a similar

approach to Bendesky *et al.*[59], we then restricted this set of substitutions between *cydno* and

melpomene to protein-coding regions, and selected those non-synonymous substitutions that

were considered to have moderate or high effect on protein function, with SNPeff[73]. Finally,

we used the PROVEAN algorithm[33], to further study the functional effects of these

substitutions on protein function. The PROVEAN algorithm predicts the functional effect of

protein sequence variations based on how they affect alignments to homologous protein

sequences (for this we used the PROVEAN protein database online). We selected those amino

acid changes with the suggested PROVEAN score < -2.5, indicating non-neutral effect on

protein function.

Admixture analyses. We retrieved estimated admixture proportions between *H. melpomene*

rosina and *H. cydno chioneus*, for 100kb and 20kb windows, from Martin et al.[35]

Data accessibility. RNA-seq data will be deposited on a public database

(https://www.ebi.ac.uk/ena) on acceptance. Analysis scripts and behavioural data are

available at: https://github.com/SpeciationBehaviour/neural_genes_heliconius.git

Author contributions: R.M.M. and M.R. conceived the study and designed the experiments,

with input from W.O.M. and C.D.J; M.R. analysed expression and sequence data; A.E.H.

analysed the behavioural data; T.J.T., S.H.M. and R.M.M. reared butterflies, dissected neural

tissue and extracted RNA; R.M.M., C.D.J. and W.O.M. secured funding, contributed

resources and provided supervision; R.P. additionally secured funding and contributed

resources; M.R. and R.M.M. wrote the manuscript with contributions from all authors.

Acknowledgments

We thank Liz Evans and Adriana Tapia for assistance in the insectaries. We are grateful to

Ana Pinharanda for sharing the *H. cydno* genome assembly and annotation and to Simon

Martin for sharing admixture proportions and comments on the manuscript. We thank the

Smithsonian Tropical Research Institute for providing research infrastructure, the Ministerio

del Ambiente for permission to collect butterflies in Panama, Edinburgh Genomics for

sequencing support and Jochen Wolf who kindly provided computational resources. MR,

AEH and RMM are supported by an Emmy Noether fellowship and research grant awarded to

RMM by the Deutsche Forschungsgemeischaft (DFG) (Grant Number: GZ: ME 4845/1-1).

RMM was also supported by a Junior Research Fellowship from King's College Cambridge,

an Ernst Mayr fellowship from the Smithsonian Tropical Research Institute, a Varley-

Gradwell fellowship from the Oxford University Museum of Natural History, and the

Balfour-Browne Fund, University of Cambridge. SHM was supported by a NERC IRF

(NE/N014936/1). This project was also supported by an Institutional Development Award

(IdeA) INBRE from the National Institute of General Medical Sciences (NIGMS) awarded to

RP (Grant number: P20GM103475), and a European Research Council (ERC) awarded to

CDJ (Grant number: 339873).

References
1. Coyne, J. A., Orr, H. A. *Speciation* (Sunderland, MA: Sinauer 2004).
2. Rosenthal, G. G. *Mate choice* (Princeton University Press, 2017).
3. Mayr, E. *Animal species and evolution* (Harvard University Press, 1963).
4. Cande, J., Prud'homme, B. & Gompel, N. Smells like evolution: The role of chemoreceptor
evolution in behavioural change. *Curr. Opin. Neurobiol.* **23**, 152–158 (2013).
5. Arguello, J. R. & Benton, R. Open questions: Tackling Darwin's "instincts": The genetic
basis of behavioural evolution. *BMC Biol.* **15**, 8–10 (2017).

6. Moehring, A. J. *et al.* Quantitative trait loci for sexual isolation between Drosophila
simulans and D. mauritiana. *Genetics 167*, 1265-1274 (2004).
7. Shaw, K. L. & Lesnick S. C. Genomic linkage of male song and female acoustic preference
QTL underlying a rapid species radiation. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 9737-9742
(2009).
8. Bay, R. A. *et al.* Genetic Coupling of Female Mate Choice with Polygenic Ecological
Divergence Facilitates Stickleback Speciation. *Curr. Biol.* **27**, 3344-3349 (2017).
9. Shahandeh, M. P., Pischedda, A., Rodriguez, J. M. & Turner, T. L. The genetics of male
pheromone preference difference between Drosophila melanogaster and Drosophila
simulans. *G3 Genes, Genomes, Genet.* **10**, 401–415 (2020).
10. Gould, F. *et al.* Sexual isolation of male moths explained by a single pheromone response
QTL containing four receptor genes. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 8660–8665 (2010)
11. Leary, G. P. *et al.* Single mutation to a sex pheromone receptor provides adaptive specificity
between closely related moth species. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 14081–14086
(2012).
12. Fan, P. *et al.* Genetic and neural mechanisms that inhibit drosophila from mating with other
species. *Cell* **154**, 89–102 (2013).
13. Brand, P. *et al.* The evolution of sexual signaling is linked to odorant receptor tuning in
perfume-collecting orchid bees. *Nat. Commun.* **11** (2020).
14. Seehausen, O. *et al.* Speciation through sensory drive in cichlid fish. *Nature* **455**, 620–626
(2008).
15. Hench, K., Vargas, M., Höppner, M. P., McMillan, W. O. & Puebla, O. Inter-chromosomal
coupling between vision and pigmentation genes during genomic divergence. *Nat. Ecol.*
Evol. **3**, 657–667 (2019).
16. Merrill, R. M. *et al.* Disruptive ecological selection on a mating cue. *Proc. R. Soc. B Biol.*
Sci. **279**, 4907–4913 (2012).
17. Jiggins, C. D., Naisbit, R. E., Coe, R. L. & Mallet, J. Reproductive isolation caused by
colour pattern mimicry. *Nature* **411**, 302–305 (2001).
18. Jiggins, C. D. Ecological Speciation in Mimetic Butterflies. *Bioscience* **58**, 541–548 (2008).
19. Jiggins, C. D., Estrada, C. & Rodrigues, A. Mimicry and the evolution of premating
isolation in Heliconius melpomene Linnaeus. *J. Evol. Biol.* **17**, 680–691 (2004).
20. Merrill, R. M. *et al.* Genetic dissection of assortative mating behaviour. *PLoS Biol* **17**,
e2005902 (2018).
21. Reed, R. D. *et al.* Optix drives the repeated convergent evolution of butterfly wing pattern
mimicry. *Science 333*, 1137-1141 (2011).
22. Martin, A. *et al.* Diversification of complex butterfly wing patterns by repeated regulatory
evolution of a Wnt ligand. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 12632–12637 (2012).
23. Nadeau, N. J. *et al.* The gene cortex controls mimicry and crypsis in butterflies and moths.
Nature **534**, 106–110 (2016).
24. Felsenstein, J. Skepticism Towards Santa Rosalia, or Why Are There So Few Kinds of
Animals? *Evolution* **35**, 124–138 (1981).
25. Massey, J. H., Chung, D., Siwanowicz, I., Stern, D. L. & Wittkopp, P. J. The yellow gene
influences drosophila male mating success through sex comb melanization. *Elife* **8**, 1–20
(2019).
26. Merrill, R. M., Van Schooten, B., Scott, J. A. & Jiggins, C. D. Pervasive genetic
associations between traits causing reproductive isolation in Heliconius butterflies. *Proc.*
R. Soc. B Biol. Sci. **278**, 511–518 (2011).
27. Seeholzer, L. F., Seppo, M., Stern, D. L. & Ruta, V. Evolution of a central neural circuit
underlies Drosophila mate preferences. *Nature* **559**, 564–569 (2018).
28. Martin, S. H. *et al.* Genome-wide evidence for speciation with gene flow in Heliconius
butterflies. *Genome Res.* **23**, 1817–1828 (2013).

29. Davey, J. *et al.* Major improvements to the Heliconius melpomene genome assembly used to confirm 10 chromosome fusion events in 6 million years of butterfly evolution. *G3* **6**, 695–708 (2015).

30. Darragh, K. *et al.* A novel terpene synthase produces an anti-aphrodisiac pheromone in the butterfly Heliconius melpomene. *bioRxiv*, 779678 (2019).

31. Wittkopp, P. J., Haerum, B. K. & Clark, A. G. Evolutionary changes in cis and trans gene regulation. *Nature* **430**, 85–88 (2004).

32. Thomas, P. D. *et al.* Applications for protein sequence-function evolution data: mRNA/protein expression analysis and coding SNP scoring tools. *Nucleic Acids Res.* **34**, 645–650 (2006).

33. Choi, Y., Sims, G. E., Murphy, S, Miller, J. R. & Chan, A. P. Predicting the Functional Effect of Amino Acid Substitutions and Indels. *PLoS One* **7** (2012).

34. Martin, S. H., Davey, J. W. & Jiggins, C. D. Evaluating the use of ABBA-BABA statistics to locate introgressed loci. *Mol. Biol. Evol.* **32**, 244–257 (2015).

35. Martin, S. H., Davey, J. W., Salazar, C. & Jiggins, C. D. Recombination rate variation shapes barriers to introgression across butterfly genomes. *PLoS Biol.* **17**, 1–28 (2019).

36. Kopp, M. *et al.* Mechanisms of assortative mating in speciation with gene flow: Connecting theory and empirical research. *Am. Nat.* **191**, 1–20 (2018).

37. Butlin, R. K. & Smadja, C. M. Coupling, reinforcement, and speciation. *Am. Nat.* **191**, 155–172 (2018).

38. Westerman, E. L. *et al.* Aristaless controls butterfly wing color variation used in mimicry and mate choice. *Current Biology*, *28*, 3469-3474 (2018).

39. Kronfrost, M. R *et al.* Linkage of butterfly mate preference and wing color preference cue at the genomic location of wingless. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 6575-6580 (2006).

40. Chamberlain, N. L., Hill, R. I., Kapan, D. D., Gilbert, L. E., & Kronforst, M. R. Polymorphic butterfly reveals the missing link in ecological speciation. *Science*, *326*, 847-850 (2009).

41. McCulloch, K. J. *et al.* Sexual Dimorphism and Retinal Mosaic Diversification following the Evolution of a Violet Receptor in Butterflies. *Mol. Biol. Evol.* **34**, 2271–2284 (2017).

42. Zaccardi, G., Kelber, A., Sison-Mangus, M. P. & Briscoe, A. D. Colour discrimination in the red range with only one long-wavelength sensitive opsin. *J. Exp. Biol.* **209**, 1944–1955 (2006).

43. Monteiro, A. Gene regulatory networks reused to build novel traits. *BioEssays* **34**, 181–186 (2012).

44. Martin, A. *et al.* Multiple recent co-options of Optix associated with novel traits in adaptive butterfly wing radiations. *Evodevo* **5** (2014).

45. Kandel, E. R, Schwartz, J. H., Jessell, T. M., Siegelbaum, S. A. & Hudspeth. A. J. *Principles of Neural Science*, 2012th Ed. (New York: McGraw Hill, 2000).

46. Wang, X. *et al.* Genomic responses to selection for tame/aggressive behaviours in the silver fox (Vulpes vulpes). *Proc. Natl. Acad. Sci. U. S. A.* **115**, 10398–10403 (2018).

47. Bloch, N. I. *et al.* Early neurogenomic response associated with variation in guppy female mate preference. *Nat. Ecol. Evol.* **2**, 1772–1781 (2018).

48. Yamaguchi, M. Role of regucalcin in brain calcium signaling. *Integr. Biol.* **4**, 825–837 (2012).

49. Berridge, M. J. Neuronal calcium signaling. *Neuron*, *21*, 13-26 (1998).

50. Bashaw, G. J. & Klein, R. Signaling from axon guidance receptors. *Cold Spring Harb. Perspect. Biol.* **2**, 1–17 (2010).

51. Prud'homme, B., Gompel, N. & Carroll, S. B. Emerging principles of regulatory evolution. *Proc. Natl. Acad. Sci. U. S. A.* *104*, 8605-8612 (2007).

52. Preger-Ben Noon, E. *et al.* Comprehensive Analysis of a cis-Regulatory Region Reveals Pleiotropy in Enhancer Function. *Cell Rep.* **22**, 3021–3031 (2018).

53. Lewis, J. *et al.* Parallel evolution of ancient, pleiotropic enhancers underlies butterfly wing
pattern mimicry. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 24174–24183 (2019).
54. Zhang, W. *et al.* Comparative Transcriptomics Provides Insights into Reticulate and
Adaptive Evolution of a Butterfly Radiation. *Genome Biol. Evol.* **11**, 2963–2975 (2019).
55. Weber, J. N., Peterson, B. K. & Hoekstra, H. E. Discrete genetic modules are responsible
for complex burrow evolution in Peromyscus mice. *Nature* **493**, 402–405 (2013).
56. Cande, J., Andolfatto, P., Prud'homme, B., Stern, D. L. & Gompel, N. Evolution of Multiple
Additive Loci Caused Divergence between Drosophila yakuba and D. santomea in Wing
Rowing during Male Courtship. *PLoS One* **7**, 1–10 (2012).
57. McBride, C. S. *et al.* Evolution of mosquito preference for humans linked to an odorant
receptor. *Nature* **515**, 222–227 (2014).
58. Ding, Y., Berrocal, A., Morita, T., Longden, K. D. & Stern, D. L. Natural courtship song
variation caused by an intronic retroelement in an ion channel gene. *Nature* **536**, 329–332
(2016).
59. Bendesky, A. *et al.* The genetic basis of parental care evolution in monogamous mice.
Nature **544**, 434–439 (2017).
60. Auer, T. O. *et al.* Olfactory receptor and circuit evolution promote host
specialization. *Nature*, 1-7 (2020).
61. Vehtari, A., Gelman, A. & Gabry, J. Practical Bayesian model evaluation using leave-one-
out cross-validation and WAIC. *Stat. Comput.* **27**, 1413–1432 (2017).
62. Jiggins, C. D. *The ecology and evolution of Heliconius butterflies* (Oxford University Press,
2016).
63. Pinharanda, A. *et al.* Sexually dimorphic gene expression and transcriptome evolution
provide mixed evidence for a fast-Z effect in Heliconius. *J. Evol. Biol.* **32**, 194–204 (2019).
64. Dobin, A. *et al.* STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21
(2013).
65. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–
2079 (2009).
66. Anders, S., Pyl, P. T. & Huber, W. HTSeq-A Python framework to work with high-
throughput sequencing data. *Bioinformatics* **31**, 166–169 (2015).
67. Love, M. I. Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol.* **15**, 1–21 (2014).
68. Montgomery, S. H. & Mank, J. E. Inferring regulatory change from gene expression: the
confounding effects of tissue scaling. *Mol. Ecol.* **25**, 5114–5128 (2016).
69. McKenna, A. *et al.* The genome analysis toolkit: A MapReduce framework for analyzing
next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).
70. Finn, R. D. *et al.* InterPro in 2017-beyond protein family and domain annotations. *Nucleic*
Acids Res. **45**, D190–D199 (2017).
71. Roberts, A., Pimentel, H., Trapnell, C. & Pachter, L. Identification of novel transcripts in
annotated genomes using RNA-seq. *Bioinformatics* **27**, 2325–2329 (2011).
72. York, R. A. *et al.* Behaviour-dependent cis regulation reveals genes and pathways
associated with bower building in cichlid fishes. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 1081–
1090 (2018).
73. Cingolani, P. *et al.* A program for annotating and predicting the effects of single nucleotide
polymorphisms, SnpEff *Fly.* **6**, 80–92 (2012).

Figures

Figure 1. Genotype at the preference QTL on chromosome 18 influences courtship initiation. Ternary plots showing the number of 15-minute choice trials in which courtship was initiated towards *melpomene, cydno* or both females for different male types. Left ternary axis shows proportion of trials where courtship was initiated towards *H. cydno* female only, bottom axis towards *H. melpomene* female only, and right axis towards both female species. Orange points represent individuals that have inherited at least one *melpomene* derived allele at the preference QTL on chromosome 18 (i.e. either *melp/melp* or *cyd/melp)*; and blue points represent individuals that are homozygous for *cydno* alleles at the preference QTL on chromosome 18 (i.e. *cyd/cyd*). Point size is scaled to the number of trials in which the male showed a response and a 'jitter' function has been applied. 95% credibility intervals (CrIs) for all three proportions are shown as hexagons around predictors with lines projecting to corresponding values on the three axes.

Figure 2. Differential expression at the preference QTL region on chromosome 18. Left: Summary of the comparative transcriptomic analyses with stage, number of samples and chromosome 18 composition. Right: the corresponding results, zooming in on the QTL region on chromosome 18. The *x*-axis represents physical position. The QTL peak, and the rest of the QTL 1.5 LOD candidate region (from[28]) are shown in green and purple, respectively. Points correspond to individual genes, with the *y*-axis indicating the *log₂*(fold-change) for each comparison. The two horizontal dashed lines (at *y*-values of 1 and -1) indicate a 2-fold change in expression. Genes showing a significant 2-fold+ change in expression level between groups are highlighted in orange and blue, where orange indicates higher levels in *melpomene* or in the hybrids *cyd/melp* (blue if in *cydno* – hybrids *cyd/cyd*). Vertical dashed lines highlight those genes that are differentially expressed between *melpomene* and *cydno* AND between *cyd/melp* vs *cyd/cyd* individuals, at the same stage. Two genes highlighted by dashed fuchsia vertical lines were excluded because they did not show differential expression, or showed reversal of the fold change when mapping RNA-seq reads to the *H. cydno* genome. Note that *Heliconius* brain (reconstruction) images, added for reference, do not include the eyes (ommatidia and retinal membrane).

Figure 3. *Grik2* and *Regucalcin2* **show evidence of allele specific expression.** Points indicate
the value, and bars the standard error, of the (base 2) logarithmic fold change in expression
between parental species (horizontal) and the alleles in F1 hybrids (vertical), for candidate
genes (as defined in the transcript-guided annotation). Dashed lines indicate the threshold for a
2-fold change in expression for the genes in the species (horizontal), and for the alleles in the
hybrids (vertical).
750

Figure 4. **Gene flow at the QTL region for behaviour.** Admixture proportion (f_d) values
estimated in overlapping 100kb (top) and 20kb (bottom) windows for chromosome 18 (top) and
the QTL region (bottom) between *H. melpomene rosin*a and *H. cydno chioneus*, with candidate
genes positions highlighted by a vertical dashed line, and *optix* location displayed for reference.
The *x*-axis represents physical position, the *y*-axis indicates the f_d value. f_d values close to zero
indicate that the proportion of shared derived alleles, and consequently gene flow, between
melpomene and *cydno* is small (or zero), implying localized selection against foreign alleles
that introgress between the two species.
762

763
764

1 **Supplementary information:** Visual mate preference evolution

2 during butterfly speciation is linked to neural processing genes

4 Matteo Rossi[1,2], Alexander E. Hausmann[1], Timothy J. Thurman[2,3], Stephen H. Montgomery[4],

5 Riccardo Papa[2,5,6], Chris D. Jiggins[7], W. Owen McMillan[2] & Richard M. Merrill[1,2]*

*1. Division of Evolutionary Biology, LMU, Munich, Germany; 2. Smithsonian Tropical Research Institute,
Panama City, Panama; 3. Division of Biological Sciences, University of Montana, USA; 4. School of Biological
Sciences, University of Bristol, Bristol, UK; 5. Department of Biology, University of Puerto Rico, San Juan,
Puerto Rico; 6. Molecular Sciences and Research Center, University of Puerto Rico, San Juan, Puerto Rico;
7. Department of Zoology, University of Cambridge, Cambridge UK.* *corresponding author**:**
merrill@bio.lmu.de

Supplementary methods and results

Mapping RNA-seq reads to the *Heliconius cydno* genome. To determine whether the *H. melpomene* reference genome introduced mapping biases of RNA-seq reads, possibly affecting differential expression estimates, we also mapped to a *H. cydno* assembly/annotation. Generally, we found similar patterns of differential expression when mapping to the two genomes. Since **i)** we observed an equal decrease (~ 40 %) of genes showing 2-fold changes in *melpomene* and *cydno* when mapping to *H. cydno,* at every stage (p-value=0.317 at adult stage, p-value=0.800 at 156h APF, p-value=0.897 at 60h APFP, Fisher's Exact test, **Table S2**), and **ii)** this decrease was widespread throughout the genome, we concluded that the *melpomene* reference genome did not bias differential gene expression analyses.

Allele-specific expression in the introgression line. BC3 hybrids had different combinations of chromosomes segregating for the *melpomene* alleles in a *cydno* background. Therefore, in principle, we could not infer *cis-* or *trans-* gene regulatory effects genome-wide from the profiles of allele specific expression (ASE) in these hybrids as for F1 hybrids, due to the diverse trans-acting environments. However, previous analyses (comparing gene expression levels between hybrids carrying *cyd/melp* vs. *cyd/cyd* regions on chromosomes other than 18) imply that differential expression of the candidate genes seems to be driven by the *melpomene* copy difference within the introgressed region on chromosome 18. Therefore, ASE analyses of candidate genes in BC3 hybrids carrying *cyd/melp* alleles on chromosome 18 should indicate whether the differences are due to cis- or trans-regulatory effects from within the introgressed region (**Figure S6**).

In BC3 hybrids sampled at 156h APF and 60hAPF, the *melpomene* and *cydno* alleles of the

ionotropic glutamate receptor (*Grik2*) are expressed at very similar levels (at 156hAPF p

value=0.841, at 60hAPF p value=0.579, Wald test), suggesting trans-only regulatory effects at

these stages for *Grik2*. For *regucalcin1* we again had very few allele-informative read counts

in hybrids at 156h APF. Although there was a tendency towards up regulation of the

melpomene allele, there was no statistical significance to support this (p=0.174, Wald test).

We detected diffASE expression of *regucalcin2* only at 60h APF (p <0.001, Wald test), but at

this stage *regucalcin2* was not detected as differentially expressed between pure species.

Thus, although there is tentative evidence for *cis*-regulatory effects for species differences in

regucalcin expression during development, it is not conclusive.

Supplementary figures

Figure S1. Ternary plots showing the number of 15-minute choice trials in which courtship was initiated towards *melpomene, cydno* or both females for backcross-to-*cydno* males, with different genotypes at the two QTLs retained in our model (on chromosome 1 and chromosome 18). Left ternary axis shows proportion of trials where courtship was initiated towards *H. cydno* female only, bottom axis towards *H. melpomene* female only, and right axis towards both female species. Each point corresponds to a male (and a 'jitter' function applied). Point size is scaled to the number of trials in which the male showed a response. 95% credibility intervals (CrIs) for all three proportions are shown as hexagons around predictors extrapolated from the statistical model with the best fit, with lines projecting to corresponding values on the three axes.

Figure S2. Results of comparative transcriptomic analyses between *melpomene* and *cydno* (in
the imago) when mapping RNA-seq reads to the *H. melpomene* assembly/annotation (top), and
to the *H. cydno* assembly/annotation (bottom), zooming in on the QTL region on chromosome
18. The *x*-axis represents physical position. Points correspond to individual genes, with the *y*-
axis indicating the *log₂*(fold-change) for each comparison. The two horizontal dashed lines (at
y-values of 1 and -1) indicate a 2-fold change in expression. Genes showing a significant 2-
fold+ change in expression level between groups are highlighted in orange and blue, where
orange indicates higher levels in *melpomene*, blue if in *cydno*. Genes detected as differentially
expressed mapping to both *melpomene* and *cydno* genomes are labelled with gene names.
dehydr.=2-oxoisovalerate dehydrogenase, OBP=odorant-binding protein.
118

Figure S3. Crossing design for producing backcross hybrids segregating at the QTL on chromosome 18. This introgression line was created by outcrossing a male hybrid to *H. cydno* females over three generations, selecting a hybrid male that showed a red band on the wing at each generation. This meant that these males carried one copy of the *H. melpomene* allele at the *optix* locus. We expected that, following recombination (which occurs in males), by the fourth generation we would remain with two types of individuals: either *cyd/melp* or *cyd/cyd* at the level of the *optix* region (which approximately corresponds to the region associated with male preference behaviour).

Figure S4. Differential gene expression at the QTL region at pupal stages. Left: summary of the comparative transcriptomic analyses with stage, number of samples and chromosome 18 composition. Right: the corresponding results, zooming in on the QTL region on chromosome 18. The *x*-axis represents physical position. The QTL peak, and the rest of the QTL 1.5 LOD candidate region are shown in green and purple, respectively. Points correspond to individual genes, with the *y*-axis indicating the *log2*(fold-change) for each comparison. The two horizontal dashed lines (at *y*-values of 1 and -1) indicate a 2-fold change in expression. Genes showing a significant 2-fold+ change in expression level between groups are highlighted in orange and blue, where orange indicates higher levels in *melpomene* or in the hybrids *cyd/melp* (blue if in *cydno* – hybrids *cyd/cyd*). Vertical dashed lines highlight those genes that are differentially expressed between *melpomene* and *cydno* AND between *cyd/melp* vs *cyd/cyd* individuals, at the same stage. One gene highlighted by a dashed fuchsia vertical line was excluded because it showed reversal of the fold change when mapping RNA-seq reads to the *H. cydno* genome.

Figure S5. A) Schematic representation of hybrid pupae (sampled at 156h APF) genome composition. Columns represent chromosomes, rows represent individuals, orange indicates *cyd/melp* regions, blue indicates *cyd/cyd* regions. **B)** Genome composition of (a subset of) BC3 hybrids. We calculated the fraction of SNPs and indels that each BC3 hybrid, one *cydno* and one F1 hybrid samples shared with *melpomene* and *cydno* samples, in non-overlapping 100kb windows. x-axes represent physical position (for each chromosome), y-axes fractions of shared variants with *melpomene* (in gold) and with *cydno* (in light blue). Matching variant fractions between BC3 hybrids and the F1 hybrid, indicating heterozygous regions, are highlighted with a gold bar underneath. Note that the general trend of higher number of variants shared with *H. cydno* in heterozygous regions is due to the fact that we inferred variants by mapping to the *H. melpomene* genome (and used variant sites only for this analysis).

A)

B)

Figure S6. Allele specific expression profiles of candidate genes at pupal stages. Points indicate
the value, and bars the standard error, of the (base 2) logarithmic fold change in expression
between parental species (horizontal) and the alleles in F1 hybrids (vertical), for candidate
genes (as defined in the transcript-guided annotation). Dashed lines indicate the threshold for a
2-fold change in expression for the genes in the species (horizontal), and for the alleles in the
hybrids (vertical).
223
224

225
226

227 **Figure S7.** Distribution of the (base 2) logarithmic fold change in allele expression. Coloured
228 bars indicate the number of genes showing a bias in expression for the *cydno* allele (in blue)
229 and for the *melpomene* allele (in yellow). Values departing from 0 on the x-axis, indicate an
230 increase in the fold change for the cydno allele (negative values) or for the *melpomene* allele
231 (positive values), respectively.
232

240 **Table S1**. List of differentially expressed genes in species and hybrids comparisons.

242 **A)** QTL chromosome 1. Orange indicates genes up-regulated in *H. melpomene,* and blue
243 those up-regulated in *H. cydno.*

#	Gene name (Hmel2.5)	Annotated function	Species comparison		
			60h APF	156h APF	Imago
1	HMEL002973g1	No match	✓	✓	✓
2	HMEL003796g1	Regulation of enolase protein 1	✓	✗	✓
3	HMEL011272g1	no match	✗	✓	✓
4	HMEL030024g1	Ribonuclease H superfamily	✓	✓	✓
5	HMEL030042g1	SWR1-complex protein 5	✗	✗	✓
6	HMEL030052g1	reverse transcriptase	✓	✓	✓
7	HMEL005260g1	unknown	✗	✓	✗
8	HMEL030040g1	No match	✗	✓	✗
9	HMEL030037g1	No match	✓	✓	✗
10	HMEL010076g1	Amino acid transporter	✓	✗	✗

B) QTL chromosome 18. Those genes found to be differentially expressed when also mapping to the *H. cydno* genome are highlighted in bold. Genes annotated as distinct but sharing the same number in the table (#) were later found to be single genes (see second paragraph of the Results section).

#	Gene name (Hmel2.5)	Annotated function	Species comparison			Hybrids comparison		
			60h APF	156h APF	Imago	60h APF	156h APF	Imago
1	**HMEL009992g1**	**No match**	✓	✗	✓	✗	✗	✓
1	***HMEL009992g4***	***Ionotropic glutamate receptor***	✓	✗	✓	✓	✓	✓
2	***HMEL009996g1***	***Gag-related protein***	✓	✓	✓	✗	✗	✗
3	HMEL034168g1	unknown	✗	✗	✓	✗	✗	✗
4	HMEL034173g1	*SWR1-complex protein 5*	✗	✗	✗	✗	✗	✗
5	HMEL034176g1	*Aspartic peptidase*	✓	✓	✓	✗	✗	✗
6	HMEL034184g1	No match	✗	✗	✓	✗	✗	✗
7	**HMEL034185g1**	**No match**	✓	✓	✓	✗	✗	✗
8	HMEL034187g1	*Major facilitator superfamily (MFS) transporter*	✓	✓	✓	✗	✓	✓
9	**HMEL003176**	***Odorant binding protein***	✗	✓	✓	✓	✗	✗
10	HMEL013551g1	*2-oxoisovalerate dehydrogenase*	✗	✗	✗	✓	✓	✗
10	***HMEL013551g2***	***2-oxoisovalerate dehydrogenase***	✗	✗	✓	✗	✗	✗
11	HMEL013551g4	*SMP-30/regucalcin*	✗	✗	✓	✗	✗	✗
12	***HMEL013552g1***	***SMP-30/regucalcin***	✗	✗	✓	✗	✓	✓
12	***HMEL034199g1***	***SMP-30/regucalcin***	✗	✗	✓	✗	✓	✓
13	HMEL014202g1	*Catalase*	✗	✗	✓	✗	✗	✗
14	***HMEL014202g3***	***Catalase***	✗	✓	✗	✗	✗	✗
15	***HMEL034201g1***	***No match***	✗	✗	✓	✗	✗	✗
16	***HMEL034205g1***	***No match***	✓	✗	✓	✗	✗	✗
17	HMEL034227g1	*Ribonuclease H superfamily*	✓	✓	✓	✗	✗	✗
18	HMEL034229g1	*Endonuclease/exonuclease/phosphatase superfamily*	✓	✓	✓	✗	✗	✓
19	HMEL034230g1	*No match*	✓	✓	✓	✗	✗	✗
20	***HMEL003863g1***	***Vacuolar protein sorting-associated (VPS) protein***	✓	✓	✗	✗	✗	✗
21	HMEL003863g3	*No match*	✓	✗	✗	✗	✗	o
22	***HMEL006662g1***	***Serpin family protein***	✗	✓	✗	✗	✗	✗
23	**HMEL006663**	***Odorant binding protein***	✗	✓	✗	✗	✗	✗
24	**HMEL022553**	***Odorant binding protein***	✗	✓	✗	✗	✗	✗
25	HMEL001038g1	*Monocarboxylate transporter*	✓	✗	✗	✗	✗	✗
26	HMEL014190g1	*unknown function*	✓	✗	✗	✗	✗	✗
27	HMEL034236g1	*No match*	✓	✗	✗	✗	✗	✗
28	HMEL034189g1	*PiggyBac transposable element-derived protein*	✗	✗	✗	✗	✗	✓
29	HMEL034246g1	*No match*	✗	✗	✗	✗	✗	✓
30	HMEL034195g1	*Gag-related protein*	✗	✗	✗	✓	✓	✗

Table S2. Number of genes showing significant >2-fold change in expression, at different stages, mapping to the *melpomene* and to the *cydno* genomes. Note that the considerable reduction in the number of genes detected as differentially expressed when mapping to *H. cydno* is most likely a result of the lower quality/completeness of the *H. cydno* genome assembly.

Stage	Mapping to:	Up-regulated in *H. melpomene*	Up-regulated in *H. cydno*
Adult	*H. melpomene*	694	733
	H. cydno	390	451
156h APF	*H. melpomene*	837	667
	H. cydno	518	403
60h APF	*H. melpomene*	846	642
	H. cydno	490	376

312 **Table S3.** Heterozygosity on the *Z*-chromosome. Heterozygosity is calculated as proportion of
313 variants (SNPs and indels) which are heterozygous, in each sample, rounded at the second
314 decimal place (note that variant sites were inferred having mapped to the *H. melpomene*
315 genome).
316

H. melpomene				*H. cydno*			
Males		Females		Males		Females	
ID	Het.	ID	Het.	ID	Het.	ID	Het.
Adults							
45	0.49	53	0.04	57	0.23	50	0.02
47	0.46	78	0.05	82	0.23	51	0.02
70	0.47	80	0.04	98	0.25	58	0.02
71	0.48	128	0.04	99	0.24	67	0.02
83	0.46	218	0.05			68	0.01
100	0.49					81	0.02
104	0.46					84	0.02
156h APF							
5	0.47	6	0.05	4	0.26	13	0.02
14	0.49	18	0.05	8	0.26	21	0.04
17	0.49	24	0.05	142	0.30	30	0.03
184	0.50	150	0.05	151	0.29	137	0.04
		220	0.06	156	0.29	168	0.04
60h APF							
92	0.48	87	0.06	85	0.26	90	0.02
97	0.49	95	0.05	86	0.27	118	0.02
115	0.47	117	0.05	119	0.27	125	0.02
		124	0.04	146	0.26	144	0.02
		149	0.06			162	0.02
		164	0.05			200	0.02
		208	0.06				

Hybrids							
Males		Females		Males		Females	
ID	Het.	ID	Het.	ID	Het.	ID	Het.
F1 hybrids (adults)							
42	0.64	56	0.05				
49	0.64	69	0.04				
Introgression line -156h APF				Introgression line – 60h APF			
105	0.26	108	0.02	152	0.27	161	0.02
116	0.27	123	0.02	193	0.27	165	0.02
126	0.27	136	0.02	198	0.25	179	0.02
131	0.27	139	0.02	199	0.27	187	0.02
133	0.26	140	0.02	201	0.26	188	0.02
154	0.26	166	0.02	212	0.28	192	0.02
155	0.26			215	0.27	197	0.02
183	0.27					209	0.03
185	0.27					214	0.02
189	0.27					224	0.02