Diversified regulation of circadian clock gene expression following whole genome

duplication

Alexander C. West[1], Marianne Iversen[1], Even H. Jørgensen[1], Simen R. Sandve[2], David G.

Hazlerigg[1], Shona H. Wood[1]*

1 Arctic chronobiology and physiology research group, Department of Arctic and Marine

Biology, UiT – The Arctic University of Norway, Tromsø NO-9037, Norway

2 Centre for Integrative Genetics (CIGENE), Department of Animal and Aquacultural Sciences

(IHA), Faculty of Life Sciences (BIOVIT), Norwegian University of Life Sciences (NMBU), 5003

Ås, Norway

* Corresponding author shona.wood@uit.no

Keywords: Whole genome duplication, Circadian, Photoperiod, Rediploidization, Anadromy,

Smoltification, Cortisol, Atlantic salmon, *Salmo salar*.

Short title: Circadian clock gene regulation and whole genome duplication

Abstract

Across taxa, circadian control of physiology and behavior arises from cell-autonomous oscillations in gene expression, governed by a networks of so-called 'clock genes', collectively forming transcription-translation feedback loops. In modern vertebrates, these networks contain multiple copies of clock gene family members, which arose through whole genome duplication (WGD) events during evolutionary history. It remains unclear to what extent multiple copies of clock gene family members are functionally redundant or have allowed for functional diversification. We addressed this problem through an analysis of clock gene expression in the Atlantic salmon, a representative of the salmonids, a group which has undergone at least 4 rounds of WGD since the base of the vertebrate lineage, giving an unusually large complement of clock genes. By comparing expression patterns across multiple tissues, and during development, we present evidence for gene- and tissue-specific divergence in expression patterns, consistent with functional diversification of clock gene duplicates. In contrast to mammals, we found no evidence for coupling between cortisol and circadian gene expression, but cortisol mediated non-circadian regulated expression of a subset of clock genes in the salmon gill was evident. This regulation is linked to changes in gill function necessary for the transition from fresh- to sea-water in anadromous fish. Overall, this analysis emphasises the potential for a richly diversified clock gene network to serve a mixture of circadian and non-circadian functions in vertebrate groups with complex genomes.

Author Summary

The generation of daily (circadian) rhythms in behaviour and physiology depends on the activities of networks of so-called clock genes. In vertebrates, these have become highly complex due to a process known as whole genome duplication, which has occurred repeatedly during evolutionary history, giving rise to additional copies of key elements of the clock gene network. It remains unclear whether this results in functional redundancy, or whether it has permitted new roles for clock genes to emerge. Here, based on studies in the Atlantic salmon, a species with an unusually large complement of clock genes, we present evidence in favour of the latter scenario. We observe marked tissue-specific, and developmentally-dependent differences in the expression patterns of duplicated copies of key clock genes, and we identify a subset of clock genes whose expression is associated with the physiological preparation to migrate to sea, but is independent of circadian regulation. Associated with this, cortisol secretion is uncoupled from circadian organisation, contrasting with the situation in mammals. Our results indicate that whole genome duplication has permitted clock genes to diversify into non-circadian functions, and raise interesting questions about the ubiquity of mammal-like coupling between circadian and endocrine function.

Introduction

Circadian control of metabolic physiology and behaviour is a ubiquitous characteristic across taxa [1–3]. In eukaryotes, circadian control derives from a cell-autonomous molecular oscillator, assembled from a regulatory network of transcription factors, co-factors, (co-) regulators, chromatin modifiers and an array of post-translational regulators of protein function, often described collectively as 'clock genes' [1]. Clock gene oscillations coordinate the transcription of multiple genes to exert effects on global cell metabolism [1]. While the molecular clock is conserved between insects and mammals [2], the mammalian network contains many duplicated components as a consequence of both local and whole genome duplication (WGD).

Two rounds of WGD preceded the establishment of the tetrapod lineage 500 million years ago (MYA) (Figure 1A), and gave rise to the complement of clock genes seen in mammals, including multiple paralogues of *Period* and *Cryptochrome* genes. Paralogues arising from WGD are known as 'ohnologues', after Susumu Ohno, who wrote a seminal monograph hypothesising that the genetic redundancy proceeding WGD facilitates evolutionary innovation [4,5]. Nevertheless, the evolutionary importance and extent to which clock gene ohnologues are functionally divergent largely remains unclear [6–10]. Indeed the retention of multiple redundant ohnologues of core clock genes is puzzling given that the essential role of the circadian clock has not changed during the course of evolutionary history [1,2,11,12]. Conceivably, functional differences between ohnologues, achieved either by coding sequence differences or by promoter-based differences in expression level, could enable tissue-specific optimization of function, but evidence for this is sparse [11,12]. It has been suggested that preferential interactions of ancient duplicated

mammalian PERIOD proteins with specific duplicated mammalian CRYPTOCHROME proteins

may affect photic entrainment [13], but experimental evidence is lacking [14]. Tissue-specific

functions of mammalian CKIδ/ε ohnologues in regulation of PERIOD protein stability have

been suggested [15], and alterations in period (tau), amplitude and clock resetting behavior

have been observed but clear distinctions of function between the ohnologues are lacking

[8,16–18].

Following the two basal vertebrate WGD events (Figure 1A), subsequent rounds of

WGD have occurred in several linages, resulting in highly complex genomes containing

thousands of ohnologue pairs. This is exemplified by the situation found in the salmonids,

which underwent two additional rounds of WGD compared to basal vertebrates; a third

WGD (Ts3R) shared by all teleost fish, and a more recent, the salmonid-specific fourth round

of duplication (Ss4R) taking place some 100 MYA (Figure 1A) [19]. The Ss4R event is a

defining characteristic of the salmonid group and is theorized to have led to the evolution of

anadromy; an adaptation of freshwater salmonids to spend part of their life-cycle at sea

[20]. Genome-scale analysis in salmonids has begun to provide new insights into the

evolutionary significance of ohnologue divergence [21,22]. Impressively, even though gene

loss often occurs following duplication events (reviewed in:[23]), there remains a rich

complexity of clock genes in teleosts compared to mammals. Of the 18 clock genes (as

defined in Figure 1B) identified in laboratory mice the zebrafish genome contains 30, and the

Atlantic salmon genome contains 61 clock genes (S1 Table, S1 appendix).

To understand why so many additional copies of core clock genes are retained, we

have undertaken a comprehensive analysis of clock gene expression in the Atlantic salmon,

exploring temporal regulation in different tissues and responsiveness to different

environmental stimuli. Here, we show diversified regulation of clock ohnologues as a result

of WGD, reflecting the fundamental differences in temporal organization of metabolism

between tissues.

Results

Tissue-specific expression of clock gene ohnologues indicates regulatory divergence

To identify all conserved clock genes in the Atlantic salmon we extracted amino acid sequences from the well-characterized mouse clock gene network (Figure 1B) then searched for homologous sequences in Atlantic salmon [22] and 12 other vertebrates (S1 appendix, Materials and Methods, *Evolutionary analysis*). Homology relationships between protein sequences were traced back to the root of the vertebrate tree, revealing 61 canonical clock genes in the Atlantic salmon. Comparing the repertoire of clock genes in spotted gar (2 WGDs), zebrafish (3 WGDs), and Atlantic salmon (4 WGDs) (Figure 1A) we find no difference in gene retention for loop 1 versus loop 2 genes (Figure 1B). Forty-two of the 61 salmon circadian genes are duplicates arising from the salmonid specific genome duplication and can be assigned to 21 Ss4R specific ohnologue pairings (referred to as Ss4R pairs from here on), while for the remaining 19 genes no extant Ss4R duplicate can be identified suggesting gene loss after WGD (S1 Table, S1 appendix). The chromosomal locations of the Ss4R pairs are shown on Figure 1C, and for ease of comparison the genes will be referred to by their gene name and chromosome. The S1 Table lists the specific gene identifiers and orthologues to zebrafish, medaka, spotted gar and mouse.

Following an autopolyploidization, such as the Ss4R, the tetraploid genome will accumulate mutations which block recombination and thereby accelerates duplicate divergence (referred to as rediploidization) [24]. This process has occurred at different rates in different genomic regions in salmonids. Using published data [22] on sequence similarity in 1Mbp windows across syntenic Ss4R regions we could classify the rediploidization times for our 21 Ss4R pairs from early (approx. 87% sequence similarity) to late (>95% sequence

similarity)(Figure 1C) and assess whether the history of rediploidization was associated with

regulatory divergence.

RNA profiling from 13 different tissues [22] demonstrated tissue-specificity of clock

gene expression, and particularly highlighted the wide variety and high abundance of clock

genes in the brain (S1 Figure). To assess the divergence between Ss4R pairs we calculated an

expression divergence index (EDI), based on the relative expression of each member of a

pair across all tissues expressed as a ratio (Figure 1D). This revealed evidence for divergent

tissue-specific expression within multiple Ss4R pairs but no clear relationship to approximate

time of rediploidization (Figure 1D). The Cry3-Ch12/Ch22 pair had the highest EDI, largely

attributable to divergent expression in the brain and gill (S1 Figure). The three Ss4R pairs of

Nr1d1 (*Rev-erbα*) and *Nr1d2* (*Rev-erbβ*), which encode transcriptional repressors linking the

circadian clock to energy metabolism [25], were also highly divergently expressed genes,

again due to differences in the brain and gill (Figure 1D, S1 Figure). Hence tissue-specific

expression divergence is a feature of particular aspects of the circadian clockwork.

Circadian and light-regulated Ss4R pair expression differs between tissues but is highly

similar within tissues

To test for circadian regulation of gene expression we collected samples from fish kept in a

light dark cycle (diel), constant light (LL) and constant dark (DD) (Figure 2A, see Circadian

experiment I, Materials and Methods). To avoid unintended rhythmical stimuli (zeitgebers),

fish were fasted from 48 hours before the first sampling point and temperature was held

constant. We focussed on three tissues with distinctive roles in salmonid physiology: the

optic tectum (OT) of the brain, because it is linked to visual processing and is coupled

directly and indirectly to light input [26–29]; the saccus vasculous (SV) because it has been

proposed as a mediator of photoperiodic responses [30]; and the gill because it is essential

for respiratory gas exchange, ion- and water balance [31]. We hypothesized that expression

profiles of clock genes in these three tissues would differ reflecting tissue-specific

differences in temporal metabolic demand. We analysed RNA transcript profiles using a

bespoke NanoString CodeSet which could specifically identify 46 clock gene targets including

17 Ss4R pairs (S2 Table, S2A Figure & S2 appendix).

In diel conditions, we identified 28 oscillating transcripts (JTK-cycle adj.p<0.05, S1

Table) [32] (Figure 2B). Of the three tissues studied, the OT showed by far the strongest

oscillations in gene expression, both under diel and constant conditions (Figure 2B & C). For

half of the genes identified, oscillation was only observed in the OT, and even for genes

showing significant oscillation across tissues (e.g. Per1-Ch4) the amplitude of oscillation was

clearly highest in the OT (Figure 2B & D).

In contrast, rhythmicity in both the SV and gill was much less robust. In the SV, only

Nr1d1-Ch6 maintained rhythmicity and phase under DD, while in the gill only Arntl1-Ch10

maintained rhythmicity and phase under DD (S2B-C Figure). Hence robust circadian

rhythmicity is a feature of the salmon brain, but gene expression rhythms are severely

dampened in the peripheral tissues we studied.

Although differences in absolute expression levels were widely seen within Ss4R pairs

- both across and within tissues, when comparing temporal dynamics of expression, within a

given tissue they were typically similar (Figure 2D, S1 Table, S2 appendix). This is exemplified

by the almost superimposable expression patterns seen for the Per1-Ch4/11 pair (Figure 2D),

and for the Tef-Ch3/6 pair (Figure 2D). Indeed, only two significant within-pair differences in

expression profile were observed (non-linear regression p-value <0.01, S1 Table, S2D-E

Figure): the Arntl1-Ch10/16 pair, with Arntl1-Ch10 showing more robust and higher

amplitude rhythmicity than Arntl1-Ch16 in the OT (Figure 2D & S2D-E Figure), and the

arrhythmic Cry3-Ch12/22 pair, with Cry3-Ch22 showing a light-induced increase in

expression following transfer to LL in the SV (Figure 2D & S2E Figure). Interestingly, we do

not observe light responses through Tef as reported in zebrafish [33].

Regulatory divergence of clock gene ohnologues within a tissue during a developmental

transition

The lack of circadian regulatory divergence among Ss4R pairs led us to consider whether

retention of duplicates might be related to developmental changes in tissue function. One

striking example of this in salmon is the transformation of gills during smoltification from a

salt retaining, water excreting organ in freshwater to a salt excreting water retaining organ

in seawater [34]. This developmental transition during the anadromous lifecycle relies on

hormonally-driven changes in physiology, dependent on seasonally changing day-length

(photoperiod) [34]. We therefore performed a photoperiod manipulation experiment, over

110 days, to assess the impact of photoperiod-dependent developmental changes in juvenile

salmon (parr) (Figure 3A, see smoltification experiment, Materials and Methods). This

protocol produces a seawater-tolerant (smolt) phenotype within 4-6 weeks of return to LL

(S3 Figure) (reviewed in:[34]).

We identified 30 clock genes showing significant changes in expression over the 110

days of the experiment (FDR<0.01, S1 Table, Figure 3B); 3 clock genes were undetectable by

RNAseq, while a further 28 were present but did not change significantly over time. Amongst

the differentially regulated genes we found 9 Ss4R pairs (18 genes) where both copies within

a pair showed smoltification-associated regulation, while for 3 pairs only one copy of the

pair showed significant changes in expression during smoltification.

Gene expression correlation cluster analysis of the differentially expressed genes

revealed 5 distinct patterns of expression over the experiment indicating that different

regulatory pathways are directing the expression of particular clock genes during

smoltification (Figure 3B). Based upon this analysis we identified 6 Ss4R pairs within which

evidence of developmental regulatory divergence could be observed (Figure 3C). For 3 of

these Ss4R pairs only one member showed a developmental change in expression (Figure 3C,

Cry3-Ch12/22, Rora-Ch2/5, Rora-Ch10/16) , while for the 3 remaining Ss4R pairs the

dynamics of expression, as defined by cluster analysis, differed within the pairing (Figure 3C,

Nr1d2-Ch2/5, Per1a-Ch4/11, CKIδ-Ch1/28). Therefore we see strong developmental

regulation of clock genes in the gill over 110 days, contrasting with the lack of circadian

regulation.

Glucocorticoid signaling induces clock ohnologue expression and accounts for regulatory

divergence observed in the Ss4R pair Tef-Ch3/6

While glucocorticoids play a major role in the circadian organization of mammals (reviewed

in: [35]), the evidence for an analogous role in fish is unclear (References summarized in: S3

Table). Nevertheless, cortisol is a major hormonal regulator of smoltification in Atlantic

salmon, steadily rising during this photoperiod-driven seasonal process [36]. We collected

blood samples from fish kept in a light dark cycle (LD - 6:18) and in constant conditions (LL or

DD) and found no evidence of diel or circadian rhythmicity in cortisol secretion (Figure 4A,

see Circadian experiment II, Materials and Methods, S4 Table) along with weak or absent

peripheral tissue clock gene circadian oscillation (S2B-C Figure). We hypothesized that the

changes in gene expression observed in Figure 3 may be due to seasonally increasing cortisol

during smoltification, therefore if we induced cortisol through a simple stress test we may

induce the same clock genes seen during smoltification.

To test this we conducted a 24 hour seawater (SW) challenge test in freshwater-

adapted fish (Figure 4B, see Smoltification experiment, Materials and Methods) eliciting an

osmotic stress-mediated increase in cortisol secretion (Figure 4C). Gills were collected from

SW and fresh water (FW) groups. We identified 15 clock genes showing significant changes

in expression in response to SW by RNAseq (FDR<0.01, S1 Table, Figure 4D). Importantly,

87% of acutely SW-responsive clock genes (13/15) also change over the chronic

developmental time-scales of smoltification (Figure 3, S1 Table). Amongst the SW responsive

genes we found 3 Ss4R pairs (6 genes) where both copies responded to SW, and 6 pairs

where only one of the pair changed expression in SW. To assess regulatory divergence within

these 9 pairs we plotted the fold change in response to SW for and determined that 5 of the

pairs showed significant regulatory divergence (two-way ANOVA <0.01, S1 Table, Figure 4E).

To further examine if glucocorticoid signaling, via cortisol, was responsible for the

induction of clock genes in the gill we used transcription factor binding site analysis [37] on

clock genes induced by SW (15 genes) compared to 43 clock genes that were SW-insensitive.

SW-induced circadian genes promoters were highly enriched for HSF1 (heat shock factor 1),

FOXO1 (forkhead box O1), MAX1 (myc-assocated factor X1) and glucocorticoid receptor

response elements (GR) (Figure 4F, S5 Table). Smoltification and responses to SW-exposure

are coordinated by multiple endocrine factors including cortisol, growth hormone (GH) and

IGF-1 [38,39]. HSF-1 and FOXO-1 elements are regulated by IGF1 signaling, during stress,

cellular metabolism and development [40–44]. Furthermore, the enrichment of GR

implicates non-circadian glucocorticoid signaling in the induction of clock genes in the gill

during smoltification and stress.

Next we wanted to experimentally test the *in-silico* association with glucocorticoid

signaling and ask whether differential sensitivity to glucocorticoid signaling might account

for the differential regulation observed within Ss4R pairs. We treated isolated gill arches

with dexamethasone (DEX; a GR agonist) for 24h and then measured the gene expression of

the 5 Ss4R pairs using qPCR (Figure 5A, S6 Table). We validated the experiment by

assessment of a positive control gene: *Tsc22d3* (*Gilz*) [45](S4 Figure). We found that whilst

some of the genes are glucocorticoid sensitive (3/10), supporting the *in-silico* association,

the majority are not induced by glucocorticoids (Figure 5B-D). However, we do demonstrate

divergence in the regulation of the Tef-Ch3/6 pair in response to glucocorticoids (Figure 5D).

Discussion

Divergent regulation of gene expression is a major contributor to the evolution of novel adaptations and species diversity [46,47]. Our analysis of the circadian clock network in the Atlantic salmon reveals clear evidence of diversified expression regulation among the many retained copies of canonical clock genes that have arisen through successive WGD events. Diversified regulation is seen in tissue-specific expression patterns (Figure 1), and in response to developmental and environmental stimuli (Figures 3, 4, 5). Ultimately, these differences likely reflect the uneven influence of diel factors on metabolism and cellular processes in central and peripheral tissues in a cold blooded fish. Altered sensitivity to GR-signalling, likely mediated through the cortisol axis, emerges as one proximate cause of differences in tissue-specific expression dynamics (Figures 4, 5). Overall this analysis emphasises the potential for a richly diversified clock gene network to serve non-circadian functions in vertebrate groups with complex genomes.

A striking contrast emerges between clock gene expression regulation in the OT and the gill. The OT is a brain site which is coupled to light input both through retinal afferents, photoreceptor expression, and indirect melatonin-mediated input via OT melatonin receptors [26–29]. In contrast, the gill is directly exposed to the water environment and continuously handles the osmotic and energetic challenges that this presents [48–50]. Thus while both tissues are highly metabolically active and heavily reliant on ATPase activity to maintain function, the principal environmental influence for the OT is rhythmic light dark input, while for the gill it is continuous osmotic challenge.

Given these differences in tissue function, it is not surprising to observe different complements of clock genes in the two tissues and dramatic differences in temporal

dynamics (Figure 2). In the diel-sensitive OT a dominant diel / circadian influence on

expression is seen, while in the gill this is a much weaker influence. Nonetheless, clock gene

expression in the gill is dynamically regulated, both over the chronic developmental time-

scales of smoltification, and acutely in response to osmotic stress. The finding that the

complement of genes showing highly sensitive changes in expression in the gill is largely

non-overlapping with that under light / circadian control in the OT is clear evidence for

diverged expression regulation within the circadian network.

Previous work has identified several Atlantic salmon clock genes that cycle in diel

conditions [51–53]. While confirming these earlier findings, our analysis goes far beyond

them by showing how the entire clock network is regulated under diel and constant

conditions in serval tissues. We show that only the OT shows persistent rhythmicity under

constant conditions, not the SV or gill (Figure 2). These results sit in contrast to the zebrafish

circadian system in which tissue autonomous clock gene oscillation are seen in all tissues

investigated [54]. Specific elements of the zebrafish clock appear to be fundamentally

different to the Atlantic salmon clock. For example zebrafish Cry1a is acutely light-induced

and appears to pause the circadian mechanism under LL [9], whereas the closest salmon

orthologues to zebrafish Cry1a, Cry1-Ch10 and Cry1-Ch16, are not light-induced and show

persistent rhythmicity in the OT under LL (S1 Table). Instead the salmon orthologues to

zebrafish Cry3b, Cry3-Ch22 and Cry3-Ch12, are acutely induced by light and salt,

respectively. Light induction of Zebrafish Tefa [33] is not seen in the closest salmon

orthologues Tef-Ch1 and Tef-Ch28, however, the zebrafish Tefb orthologues, Tef-Ch3 and

Tef-Ch6 are seawater induced. In zebrafish, Tef transcription is induced in response to light

and UV exposure producing reactive oxygen species (ROS) which act on the D-box in the Tef

promoter, initiating a DNA damage repair cascade [55]. Importantly this light-mediated ROS

inducibility of Tef appears to be absent in cave fish and mammals [55], suggesting a

zebrafish-specific adaptation to UV light exposure, potentially related to spawning

environment [56]. In our study we found no enrichment for genes associated with oxidative

stress in either the seawater challenge or photoperiodic treatment groups but the

photoperiod-driven smoltification induction of all four Tef genes is noteworthy, as is the

seawater specific induction of the Ss4r pair Tef-Ch3 and Ch6, and the apparent regulatory

divergence of these duplicates in glucocorticoid responsiveness. These species differences

indicate plasticity of the circadian clock in response to different environmental stressors

depending on life history.

Within the Atlantic salmon the distinct tissue-specific expression dynamics we

observe and the smoltification and stress-related induction of clock genes focuses attention

on cortisol [34,35,57]. In mammals cortisol is a major player in circadian organization, acting

as an internal zeitgeber through which the hypothalamic–pituitary–adrenal (HPA) axis can

coordinate daily changes in tissue activity throughout the organism (reviewed in: [35]).

Because the teleost hypothalamic–pituitary–interrenal (HPI) axis is functionally analogous to

the HPA axis [57], there has been a widespread assumption that cortisol plays a similar

circadian coordination function in teleosts [58], although definitive evidence for this is

lacking (S3 Table)[59]. Indeed our study implies that it is unlikely that cortisol plays a

circadian role in salmonids – we found no evidence for circadian or even diel changes in

cortisol secretion in Atlantic salmon (Figure 4A), and GREs are less enriched in circadian

oscillating clock genes than in non-oscillating, development- and SW-sensitive clock genes

(Figure 4F). Hence while our data strongly implicate cortisol in the dynamic expression of a

subset of Atlantic salmon clock genes, this seems to have nothing to do with circadian

function *per se*, and more to do with a role for these genes in non-circadian influences of the

HPI axis. It is interesting to speculate that this shifting relationship between glucocorticoids

and clock genes could be a contributory factor for the evolution of anadromy and the

regulation of its seasonal timing. Furthermore, this finding raises interesting questions about

the ubiquity and evolutionary origins of mammal-like coupling between the HPA axis and

circadian function.

While evidence of functional divergence in ancient duplicated cryptochromes, and,

sequence divergence of Ts3R cryptochromes in zebrafish have been observed [60] we

expected that Ss4R generated significant genetic complexity in the salmonid circadian clock

and therefore asked how expression patterns diverged within Ss4R pairs. Intriguingly, the

answer to this question was highly dependent upon context. We saw many examples of

pronounced within-Ss4R pair differences in terms of tissue-specific expression, and some

50% of identifiable Ss4R pairs showed within-pair divergence in expression during

smoltification. But within-pair divergence in daily expression patterns was hardly observed.

Why might this be so? We suggest this may reflect a difference in the way that selection

pressures have operated on promoter regions to, on the one hand modulate tissue-specific

expression and, on the other daily temporal regulation. According to this view duplication

would confer freedom to diverge, thereby meeting differing tissue-specific requirements.

Conversely, the daily temporal patterning may be so fundamental to cell function that any

mutations leading to deviation from the ancestral dynamics were strongly selected against.

This conjecture will require detailed analysis of regions of promoter conservation /

divergence among Ss4R pairs.

Materials and Methods:

Evolutionary analysis

To identify gene orthologs and ohnologs we generated protein sequence homology based

orthogroups using the Orthofinder pipeline [61]. For each orthogroup we used the resulting

protein tree topology to manually annotate pairs of salmon ohnologs based on the following

criteria: (i) salmon ohnologs should form a monophyletic clade only containing genes from

other salmonid species, (ii) this monophyletic salmonid clade must have Northern pike as the

sister group, and (iii) putative ohnolog pairs had to be conserved in minimum one other

salmonid species. Finally, we only retained putative ohnolog if their genomic positions were

defined as syntenic regions originating from the Ss4R as defined in Lien *et al.* [22]. Maximum

likelihood gene trees with bootstrap values for all genes included in the study are presented

in S1 appendix. These trees include the following species: human, mouse, spotted gar,

zebrafish, stickleback, medaka, northern pike, grayling, danube salmon, arctic charr, coho

salmon, rainbow trout, Atlantic salmon. The relevant scripts, input files and resulting trees

are also available here. For quick reference S1 table presents the salmon gene loci and their

corresponding orthologues in Zebrafish, spotted gar, medaka and mouse.

Multi-tissue analysis

Publically available data was used to assess the multi-tissue expression in the Atlantic

salmon (one adult male, kept in freshwater), these data can be found in the NCBI Sequence

Read Archive (SRA): PRJNA72713 and PRJNA260929.

Ethics statement

Fish handling and euthanasia was performed by competent persons and in accordance with

the European Union Regulations concerning the protection and welfare of experimental

animals (European directive 91/492/CCE). The experiment was approved by the Norwegian

Committee on Ethics in Animal Experimentation (ID 3630).

Animal husbandry

Juvenile Atlantic salmon (*Salmo salar*, Linnaeus, 1758) of the Aquagene commercial strand

(Trondheim, Norway) were used in all experiments. Fish were held under constant light (LL;

>200 lux), at 10° from hatching onwards, and kept in 500 L tanks from first feeding. The fish

were approximately 7 months old when the experiments were initiated. Up until that time

the fish had been feed continuously with pelleted salmon feed (Skretting, Stavanger,

Norway), from automatic feeders.

Circadian experiment I

Fish were maintained in 500L freshwater were transferred from LL to a short photoperiod

(SP; 6L:18D) light schedule for 8 weeks before the start of the experiment. Short

photoperiods provide a strong zeitgeber (timing signal) entraining the fish to a light dark

cycle. In order to test for a circadian rhythm it is necessary to remove any zeitgebers, light,

temperature and food are all potential zeitgebers which we controlled for. Two weeks

before sampling, fish were distributed to two separate 150L tanks and water temperature

was maintained at 14°C (±0.5). Fish were fasted for 48 hours prior to the experiment and

throughout the sampling. To identify genes which were driven by an endogenous circadian

rhythm we sampled under diel (SP; 6L:18D), constant light (LL) and constant dark conditions

(DD). Sampling was at 4 hour resolution starting at ZT5 (zeitgeber time; time since lights on)

in diel conditions (n= 3 per time point, 7 time-points), transitioning onto either LL or DD. In

LL sampling was from CT9 (circadian time) at a 4 hour resolution (n = 3 per time-point, 6

time-points) because ZT 1 and 5 in this design can also be defined as CT1 and 5. In DD

samples were collected from CT1 at a 4 hour resolution (n = 3 per timepoint, 8 time-points)

(Figure 2A). For the statistical analysis ZT5 to ZT5 was used for diel conditions, and CT9 to

CT29 for constant conditions. Collections during the dark phase were conducted under dim

red light. During sampling fish were netted out and euthanized by an overdose of

benzocaine (150ppm). Weight and length were recorded and no significant variation noted.

Optic tectum, gill and saccus vasculous were dissected and snap frozen on dry ice. RNA was

extracted for subsequent nanostring profiling.

Circadian experiment II

Fish were maintained in 500L freshwater were transferred from LL to a short photoperiod

(SP; 6L:18D) light schedule for 20 weeks before the start of the experiment. Two weeks

before sampling, fish were distributed to two separate 150L tanks. Fish were fasted for 48

hours prior to the experiment and throughout the sampling. Temperature was maintained at

an average of 8.5^{o}C ±1. Sampling was at a 4 hour resolution starting one hour after lights on

(ZT1), maintaining the fish under diel conditions (SP; 6L:18D) for 24 hours and then switching

to either LL or DD, and sampling at 4h resolution for a further 29 hours (n = 4 per time-point)

(Figure 4A). Collections during the dark phase were conducted under dim red light. During

sampling fish were netted out and euthanized by an overdose of benzocaine (150ppm).

Weight and length were recorded and no significant variation noted. Blood was collected at

each time-point from the caudal vein in heparinized vacutainers. Blood samples were

centrifuged at 500 x g for 15 min to collect plasma for subsequent hormone analysis.

Smoltification experiment and seawater tests

Using a standard aquaculture and research method we used photoperiod to induce the

developmental transition of juvenile salmon (parr) to seawater prepared fish (smolts). This

method requires switching parr on constant light to short photoperiod (SP; 6L:18D) for at

least 8 weeks before returning the fish to constant light where over 4 weeks they gain osmo-

regulatory capacity in seawater (Figure 3A). Fish were maintained in 150L freshwater tanks

at an average 8.5^{o}C ±1 and were transferred from LL to SP (8L:16:D) for 8 weeks before

return to LL for 7 weeks. Fish were sampled at six time-points in fresh-water shown in Figure

3A (T1-T6, n = 6 per time-point), these time-points are designed to capture the

photoperiodically induced transition from parr to smolt. Fish were fasted for 48 hours prior

to sampling. Gills were collected into RNAlater (Sigma-Aldrich, St. Louis, Missouri, USA),

storing at 4°C for 24 hours, before being transferred to -80°C. RNA was extracted and used

for RNA-seq (Data shown in Figure 3).

To validate the photoperiod protocol seawater challenge tests were conducted on a

randomly selected subgroup that were transferred to a 100L tank supplied with seawater

(34% salinity) for 24 hours. Fish were netted out after 24 hours and euthanized by an

overdose of benzocaine (150ppm). Blood samples were collected from the caudal vein in

heparinized vacutainers and plasma was collected and stored at -20⬚. To test the osmolality

of blood thawed plasma samples were analysed for osmolyte content using a Fiske One-Ten

Osmometer (Fiske Associates, Massachusetts, USA, ± 4 mOsm kg-1). These tests were

conducted from T2 to T6 (n = 12 per time-point), and confirm that from T5 (29 days after the

return to LL) the fish had osmo-regulatory capacity in seawater (S3 Figure). Gills were

collected at each time-point after seawater challenge (n= 6 per time-point), RNA was

extracted and RNA-seq preformed as above. All data are available from the European

nucleotide archive under project number: PRJEB34224. This study presents data from the

freshwater groups at all time-points (Figure 3) and seawater challenged fish at T4 only

(Figure 4B). T4 is 8 days after return to LL and therefore these fish do not have osmo-

regualtory capacity in seawater and are used as a stress test comparison to fish from the

same time-point kept in freshwater (Figure 4B – F).

In-vitro Gill Culture

Juvenile Atlantic Salmon were prepared as in the smoltification experiment, sampling at the

equivalent of T4 (8 days after return to LL) (Figure 5A). Following euthanasia whole gill

arches were rapidly dissected (biological replicates, n=4 per treatment), excess mucus was

removed by careful blotting onto tissue paper before the arches were transferred

individually into 50 ml of pre-prepared control or treatment media. The prepared media

consisted of Leibovitz L-15 (Lonza) supplemented with non-essential amino acids (1%, 100x

Lonza), sodium-pyruvate (1%, 100x Lonza), 0.05 mg/ml gentamycin (Sigma) and 20% fetal

bovine serum (FBS, sigma). The experimental group was supplemented with 0.1M

dexamethasone diluted in DMSO (dimethyl-sulphoxide, Sigma) to a final concentration of

0.1μM. The control group contained an equivalent concentration of DMSO (0.1%). The

excised gill arches were incubated for 24h at 4°C, gill filaments were removed with a scalpel

and snap frozen on dry ice before being stored at -80°C. RNA was extracted for qPCR

analysis.

RNA extraction

RNA extraction for RNAseq was performed using a TRIzol-based method (Invitrogen, Thermo

Fisher, Waltham, Massachusetts, USA), and in accordance with the manufacturers

recommendation. Resulting RNA concentrations and quality were checked using a NanoDrop

spectrophotometer (NanoDrop Technologies, Wilmington, DE, USA). RNA was stored at -

80°C.

For nanostring and qPCR, SVs were extracted using QIAgen RNeasy micro kit, OT and gill

tissues were extracted using QIAgen RNeasy mini kit according to the manufacturers

instructions. RNA concentration was quantified and quality confirmed using the Experion

Automated Electrophoresis System (BioRad).

Nanostring

Custom nanostring codesets were designed by Nanostring Technologies Inc. using the

Atlantic Salmon reference genome (Cigene), accession numbers and target sequences are

shown in S2 Table. Codesets were processed by the Univerisity of Manchester Genomic

Technologies Core Facility. This technology is based on the use of fluorescent barcoded

probes which bind specifically to the target molecule. Importantly these barcodes should

only bind one at a time to each target molecule, therefore the number of fluorescent

barcodes reflects the number of RNA molecules of your target gene. A spike control with a

known number of RNA molecules was also used to normalise across samples and runs.

Therefore the units are counts normalised to spike-in positive controls. Data was processed

using nSolver 4.0 software (Nanostring). Data can be accessed on GEO under the project

identifier GSE146530.

Transcriptome sequencing and assembly

Libraries were prepared using TruSeq Stranded mRNA HS kit (Illumina, San Diego, California,

USA). Mean library length was determined using the 2100 Bioanalyzer with the DNA 1000 kit

(Agilent Technologies, Santa Clara, California, USA). Library concentrations was determined

using the Qubit BR kit (Thermo Scientific, Waltham, Massachusetts, USA). Samples were

barcoded with Illumina unique indexes. The Illumina HiSeq 2500 was used to perform single-

end 100-bp sequencing of samples at the Norwegian Sequencing Centre (University of Oslo,

Oslo, Norway).

Cutadapt (ver. 1.8.1) was used for removal of sequencing adapters and trimming of low

quality bases (parameters –q 20, -O 8 -minimum-length 40). Quality control was performed

with FastQC software. Reads were mapped onto the references genome using STAR

software (ver. 2.4.2a). Read counts for annotated genes were generated using the HTSEQ-

count software (ver. 0.6.1p1).

All RNAseq data for the smoltification experiment is available in the European nucleotide

archive under project number: PRJEB34224.

Analysis of differentially expressed genes

Analysis of differential gene expression was performed with package edgeR (ver. 3.14.0)

using R (ver. 3.4.2) and RStudio (ver. 1.0.153). Prior to analysis of differential expression, the

raw counts were filtered, setting an expression level threshold of a minimum of one count

per million reads (cpm) in five or more libraries, resulting in a list of 33 951 expressed genes.

The counts were scaled by applying trimmed means of M-values (TMM) scaling. Exact tests

were then performed to find genes that were differentially expressed between FW-kept and

24-h SW challenged fish. An ANOVA-like test was performed to find genes that were

differentially expressed over T1-T6 FW time-points. The test results were filtered for a false

discovery rate (FDR) to be less than 0.01 to identify significantly differentially expressed

genes. Clustering analysis was performed using Pearson correlation.

Heatmaps were generated in R using custom scripts for pheatmap. Transcription factor

binding site analysis was conducted using SalmotifDB [37].

qPCR

cDNA was synthesised from sample total RNA using High capacity RNA to cDNA kit (Applied

Biosystems). qPCR was performed using GoTaq Master Mix (Promega) and a 96 well thermal

cycler (Applied Biosystems). Relative gene expression was quantified by the ΔΔCT method

using *Ef1a* as reference gene. Primer sequences are listed in S2 Table and source data is in S6

Table.

Hormone Assays

Cortisol ELISA assays were performed by Stockgrand (UK). Source data in S4 Table.

Statistical analyses

RNAseq analysis is detailed above. Mean difference comparisons were carried out using

Student's t-test (two-sided, unpaired), two-way ANOVA with post hoc tests as appropriate

(Graphpad Prism 8.1.2). The expression divergence index (EDI) index was calculated as

follows: EDI= abs(log2[Gene1/Gene2)].

The R package JTK cycle was used to assess rhythmicity of transcripts under LD and constant

light or dark conditions [32]. For statistical comparison of gene expression between

ohnologue pairs in the circadian experiment, expression was normalized to group mean then

best fit sixth-order centered polynomial curves were generated by non-linear regression

analysis and shared characteristics tested with extra sum of squares F test (Graphpad Prism

8.0).

References

1. Shi M, Zheng X. Interactions between the circadian clock and metabolism: there are

good times and bad times. Acta Biochim Biophys Sin (Shanghai). 2013;45: 61–69.

doi:10.1093/abbs/gms110

2. Rosbash M. The Implications of Multiple Circadian Clock Origins. Michael Young,

editor. PLoS Biol. 2009;7: e1000062. doi:10.1371/journal.pbio.1000062

3. Young MW, Kay SA. Time zones: a comparative genetics of circadian clocks. Nat Rev

Genet. 2001;2: 702–715. doi:10.1038/35088576

4. Taylor JS, Raes J. Duplication and divergence: the evolution of new genes and old

ideas. Annu Rev Genet. 2004;38: 615–43.

doi:10.1146/annurev.genet.38.072902.092831

5. Ohno S. Evolution by Gene Duplication. Berlin, Heidelberg: Springer Berlin Heidelberg;

1970. doi:10.1007/978-3-642-86659-3

6. Horst GTJ van der, Muijtjens M, Kobayashi K, Takano R, Kanno S, Takao M, et al.

Mammalian Cry1 and Cry2 are essential for maintenance of circadian rhythms.

Nature. 1999;398: 627–630. doi:10.1038/19323

7. Erzberger A, Hampp G, Granada AE, Albrecht U, Herzel H. Genetic redundancy

strengthens the circadian clock leading to a narrow entrainment range. J R Soc

Interface. 2013;10: 20130221. doi:10.1098/rsif.2013.0221

8. Pilorz V, Cunningham PS, Jackson A, West AC, Wager TT, Loudon ASI, et al. A Novel

Mechanism Controlling Resetting Speed of the Circadian Clock to Environmental

Stimuli. Curr Biol. 2014;24: 766–773. doi:10.1016/j.cub.2014.02.027

9. Tamai TK, Young LC, Whitmore D. Light signaling to the zebrafish circadian clock by

Cryptochrome 1a. Proc Natl Acad Sci. 2007;104: 14712–14717.

doi:10.1073/pnas.0704588104

10. Shi S, Hida A, McGuinness OP, Wasserman DH, Yamazaki S, Johnson CH. Circadian

Clock Gene Bmal1 Is Not Essential; Functional Replacement with its Paralog, Bmal2.

Curr Biol. 2010;20: 316–321. doi:10.1016/j.cub.2009.12.034

11. Tauber E, Last KS, Olive PJW, Kyriacou CP. Clock Gene Evolution and Functional

Divergence. J Biol Rhythms. 2004;19: 445–458. doi:10.1177/0748730404268775

12. Looby P, Loudon ASI. Gene duplication and complex circadian clocks in mammals.

Trends Genet. 2005;21: 46–53. doi:10.1016/j.tig.2004.11.012

13. Daan S, Albrecht U, Van der Horst GTJ, Illnerová H, Roenneberg T, Wehr TA, et al.

Assembling a Clock for All Seasons: Are There M and E Oscillators in the Genes? J Biol

Rhythms. 2001;16: 105–116. doi:10.1177/074873001129001809

14. Hastings M. Modeling the Molecular Calendar. J Biol Rhythms. 2001;16: 117–123.

doi:10.1177/074873001129001818

15. Meng Q-J, Logunova L, Maywood ES, Gallego M, Lebiecki J, Brown TM, et al. Setting

Clock Speed in Mammals: The CK1☐ tau Mutation in Mice Accelerates Circadian

Pacemakers by Selectively Destabilizing PERIOD Proteins. Neuron. 2008;58: 78–88.

doi:10.1016/j.neuron.2008.01.019

16. Meng Q-J, Maywood ES, Bechtold DA, Lu W-Q, Li J, Gibbs JE, et al. Entrainment of

disrupted circadian behavior through inhibition of casein kinase 1 (CK1) enzymes. Proc

Natl Acad Sci. 2010;107: 15240–15245. doi:10.1073/pnas.1005101107

17.   Lee H, Chen R, Lee Y, Yoo S, Lee C. Essential roles of CKIδ and CKIε in the mammalian

circadian clock. Proc Natl Acad Sci. 2009;106: 21359–21364.

doi:10.1073/pnas.0906651106

18.   Etchegaray J-P, Machida KK, Noton E, Constance CM, Dallmann R, Di Napoli MN, et al.

Casein Kinase 1 Delta Regulates the Pace of the Mammalian Circadian Clock. Mol Cell

Biol. 2009;29: 3853–3866. doi:10.1128/MCB.00338-09

19.   Macqueen DJ, Johnston IA. A well-constrained estimate for the timing of the salmonid

whole genome duplication reveals major decoupling from species diversification. Proc

R Soc B Biol Sci. 2014;281: 20132881. doi:10.1098/rspb.2013.2881

20.   Alexandrou MA, Swartz BA, Matzke NJ, Oakley TH. Molecular Phylogenetics and

Evolution Genome duplication and multiple evolutionary origins of complex migratory

behavior in Salmonidae. Mol Phylogenet Evol. 2013;69: 514–523.

doi:10.1016/j.ympev.2013.07.026

21.   Robertson FM, Gundappa MK, Grammes F, Hvidsten TR, Redmond AK, Lien S, et al.

Lineage-specific rediploidization is a mechanism to explain time-lags between genome

duplication and evolutionary diversification. Genome Biol. 2017;18: 111.

doi:10.1186/s13059-017-1241-z

22.   Lien S, Koop BF, Sandve SR, Miller JR, Matthew P, Leong JS, et al. The Atlantic salmon

genome provides insights into rediploidization. Nature. 2016;533: 200–205.

doi:10.1038/nature17164

23.   Moriyama Y, Koshiba-takeuchi K. Significance of whole-genome duplications on the

emergence of evolutionary novelties. 2018;17: 329–338. doi:10.1093/bfgp/ely007

24.   Li WH. Rate of gene silencing at duplicate loci: a theoretical study and interpretation

588   of data from tetraploid fishes. Genetics. 1980;95: 237–58. doi:7429144

589   25.   Cho H, Zhao X, Hatori M, Yu RT, Barish GD, Lam MT, et al. Regulation of circadian

590   behaviour and metabolism by REV-ERB-α and REV-ERB-β. Nature. 2012;485: 123–127.

591   doi:10.1038/nature11048

592   26.   Holmqvist BI, Östholm T, Ekström P. Neuroanatomical analysis of the visual and

593   hypophysiotropic systems in Atlantic salmon (Salmo salar) with emphasis on possible

594   mediators of photoperiodic cues during parr-smolt transformation. Aquaculture.

595   1994;121: 1–12. doi:10.1016/0044-8486(94)90003-5

596   27.   Meyer RL. Eye-in-water electrophysiological mapping of goldfish with and without

597   tectal lesions. Exp Neurol. 1977;56: 23–41. doi:10.1016/0014-4886(77)90136-4

598   28.   Sandbakken M, Ebbesson L, Stefansson S, Helvik JV. Isolation and characterization of

599   melanopsin photoreceptors of atlantic salmon (Salmo salar). J Comp Neurol.

600   2012;520: 3727–3744. doi:10.1002/cne.23125

601   29.   Mazurais D, Brierley I, Anglade I, Drew J, Randall C, Bromage N, et al. Central

602   melatonin receptors in the rainbow trout: Comparative distribution of ligand binding

603   and gene expression. J Comp Neurol. 1999;409: 313–324. doi:10.1002/(SICI)1096-

604   9861(19990628)409:2<313::AID-CNE11>3.0.CO;2-1

605   30.   Nakane Y, Ikegami K, Iigo M, Ono H, Takeda K, Takahashi D, et al. The saccus

606   vasculosus of fish is a sensor of seasonal changes in day length. Nat Commun. 2013;4:

607   2108. doi:10.1038/ncomms3108

31.   Evans DH, Piermarini PM, Choe KP. The Multifunctional Fish Gill: Dominant Site of Gas

Exchange, Osmoregulation, Acid-Base Regulation, and Excretion of Nitrogenous

Waste. Physiol Rev. 2005;85: 97–177. doi:10.1152/physrev.00050.2003

32.   Hughes ME, Hogenesch JB, Kornacker K. JTK_CYCLE: An Efficient Nonparametric

Algorithm for Detecting Rhythmic Components in Genome-Scale Data Sets. J Biol

Rhythms. 2010;25: 372–380. doi:10.1177/0748730410379711

33.   Vatine G, Vallone D, Appelbaum L, Mracek P, Ben-Moshe Z, Lahiri K, et al. Light Directs

Zebrafish period2 Expression via Conserved D and E Boxes. Kramer A, editor. PLoS

Biol. 2009;7: e1000223. doi:10.1371/journal.pbio.1000223

34.   Björnsson BT, Stefansson SO, McCormick SD. Environmental endocrinology of salmon

smoltification. Gen Comp Endocrinol. 2011;170: 290–298.

doi:10.1016/j.ygcen.2010.07.003

35.   Dumbell R, Matveeva O, Oster H. Circadian Clocks, Stress, and Immunity. Front

Endocrinol (Lausanne). 2016;7. doi:10.3389/fendo.2016.00037

36.   Langhorne P, Simpson TH. The interrelationship of cortisol, Gill (Na + K) ATPase, and

homeostasis during the Parr-Smolt transformation of atlantic salmon (Salmo salar L.).

Gen Comp Endocrinol. 1986;61: 203–213. doi:10.1016/0016-6480(86)90198-X

37.   Mulugeta TD, Nome T, To T-H, Gundappa MK, Macqueen DJ, Våge DI, et al.

SalMotifDB: a tool for analyzing putative transcription factor binding sites in salmonid

genomes. BMC Genomics. 2019;20: 694. doi:10.1186/s12864-019-6051-0

38.   McCormick SD. Effects of Growth Hormone and Insulin-like Growth Factor I on Salinity

Tolerance and Gill Na+, K+-ATPase in Atlantic Salmon (Salmo salar): Interaction with

Cortisol. Gen Comp Endocrinol. 1996;101: 3–11. doi:10.1006/gcen.1996.0002

39.   McCormick SD. Endocrine Control of Osmoregulation in Teleost Fish. Am Zool.

2001;41: 781–794. doi:10.1093/icb/41.4.781

40.   Accili D, Arden KC. FoxOs at the Crossroads of Cellular Metabolism, Differentiation,

and Transformation. Cell. 2004;117: 421–426. doi:10.1016/S0092-8674(04)00452-0

41.   Barthel A, Schmoll D, Unterman TG. FoxO proteins in insulin action and metabolism.

Trends Endocrinol Metab. 2005;16: 183–189. doi:10.1016/j.tem.2005.03.010

42.   Dong XC, Copps KD, Guo S, Li Y, Kollipara R, DePinho RA, et al. Inactivation of Hepatic

Foxo1 by Insulin Signaling Is Required for Adaptive Nutrient Homeostasis and

Endocrine Growth Regulation. Cell Metab. 2008;8: 65–76.

doi:10.1016/j.cmet.2008.06.006

43.   Li J, Labbadia J, Morimoto RI. Rethinking HSF1 in Stress, Development, and Organismal

Health. Trends Cell Biol. 2017;27: 895–905. doi:10.1016/j.tcb.2017.08.002

44.   Barna J, Princz A, Kosztelnik M, Hargitai B, Takács-Vellai K, Vellai T. Heat shock factor-1

intertwines insulin/IGF-1, TGF-β and cGMP signaling to control development and

aging. BMC Dev Biol. 2012;12: 32. doi:10.1186/1471-213X-12-32

45.   Ronchetti S, Migliorati G, Riccardi C. GILZ as a Mediator of the Anti-Inflammatory

Effects of Glucocorticoids. Front Endocrinol (Lausanne). 2015;6.

doi:10.3389/fendo.2015.00170

46.   Carroll SB. Evo-Devo and an Expanding Evolutionary Synthesis: A Genetic Theory of

Morphological Evolution. Cell. 2008;134: 25–36. doi:10.1016/j.cell.2008.06.030

47.   Carroll SB. Endless Forms. Cell. 2000;101: 577–580. doi:10.1016/S0092-

8674(00)80868-5

48.   McCormick SD, Saunders R. Preparatory Physiological Adaptations for Marine Life of

654   Salmonids: Osmoregulation, Growth, and Metabolism. Am Fish Soc Symp. 1987;1:

655   211–229.

656   49.   McCarthy ID. Temporal repeatability of relative standard metabolic rate in juvenile

657   Atlantic salmon and its relation to life history variation. J Fish Biol. 2000;57: 224–238.

658   doi:10.1111/j.1095-8649.2000.tb00788.x

659   50.   McCormick SD, Moyes CD, Ballantyne JS. Influence of salinity on the energetics of gill

660   and kidney of Atlantic salmon (Salmo salar). Fish Physiol Biochem. 1989;6: 243–254.

661   doi:10.1007/BF01875027

662   51.   Davie A, Minghetti M, Migaud H. Seasonal Variations in Clock-Gene Expression in

663   Atlantic Salmon (Salmo salar). Chronobiol Int. 2009;26: 379–395.

664   doi:10.1080/07420520902820947

665   52.   Huang T, Ruoff P, Fjelldal PG. Diurnal expression of clock genes in pineal gland and

666   brain and plasma levels of melatonin and cortisol in Atalantic salmon parr and smolts.

667   Chronobiol Int. 2010;27: 1697–1714. doi:10.3109/07420528.2010.514630

668   53.   Huang T, Ruoff P, Fjelldal PG. Effect of continuous light on daily levels of plasma

669   melatonin and cortisol and expression of clock genes in the pineal gland, brain, and

670   liver in Atalantic salmon postsmolts. Chronobiol Int. 2010;27: 1715–1734.

671   doi:10.3109/07420528.2010.521272

672   54.   Davies WIL, Tamai TK, Zheng L, Fu JK, Rihel J, Foster RG, et al. An extended family of

673   novel vertebrate photopigments is widely expressed and displays a diversity of

674   function. Genome Res. 2015;25: 1666–1679. doi:10.1101/gr.189886.115

675   55.   Pagano C, Siauciunaite R, Idda ML, Ruggiero G, Ceinos RM, Pagano M, et al. Evolution

676   shapes the responsiveness of the D-box enhancer element to light and reactive

677   oxygen species in vertebrates. Sci Rep. 2018;8: 13180. doi:10.1038/s41598-018-

678   31570-8

679   56.   Gavriouchkina D, Fischer S, Ivacevic T, Stolte J, Benes V, Dekens MPS. Thyrotroph

680   Embryonic Factor Regulates Light-Induced Transcription of Repair Genes in Zebrafish

681   Embryonic Cells. Riley BB, editor. PLoS One. 2010;5: e12542.

682   doi:10.1371/journal.pone.0012542

683   57.   Wendelaar Bonga SE. The stress response in fish. Physiol Rev. 1997;77: 591–625.

684   doi:10.1152/physrev.1997.77.3.591

685   58.   Isorna E, Pedro N De, Valenciano AI, Alonso-gómez ÁL. Interplay between the

686   endocrine and circadian systems in fishes. 2017. doi:10.1530/JOE-16-0330

687   59.   Mommsen TP, Vijayan MM, Moon TW. Cortisol in teleosts: dynamics, mechanisms of

688   action, and metabolic regulation. Rev Fish Biol Fish. 1999;9: 211–268.

689   60.   Liu C, Hu J, Qu C, Wang L, Huang G, Niu P, et al. Molecular evolution and functional

690   divergence of zebrafish (Danio rerio) cryptochrome genes. Sci Rep. 2015;5: 8113.

691   doi:10.1038/srep08113

692   61.   Emms DM, Kelly S. OrthoFinder: solving fundamental biases in whole genome

693   comparisons dramatically improves orthogroup inference accuracy. Genome Biol.

694   2015; 1–14. doi:10.1186/s13059-015-0721-2

Acknowledgements

The authors thank all of the animal staff at Kårvik havbruksstasjonen for their expert care of their research animals. The work was supported by grants from the Tromsø forskningsstiftelse (TFS) starter grant TFS2016SW, awarded to SHW. ACW is supported by the Tromsø forskningsstiftelse (TFS) grant awarded to DGH (TFS2016DH). Research council of Norway grant number 241016 awarded to DGH. The publication charges for this article have been funded by a grant from the publication fund of UiT-The Arctic University of Norway.

Author contributions

ACW – conceived and designed the circadian and *in-vitro* experiments, collected samples, performed RNA preparation, gill culture, qPCR, bioinformatics analysis, analysed/interpreted data, prepared the manuscript and figures. **MI–** Bioinformatic analysis, osmolality measurements, data analysis and reviewed manuscript. **EHJ –** collected samples, designed and conducted the smoltification experiment, and reviewed the manuscript. **SS –** Conducted the evolutionary analysis of clock genes and revised the manuscript. **DGH** - collected samples, analysed/interpreted data, designed and conducted the smoltification experiment, and revised the manuscript. **SHW–** conceived and designed the circadian and *in-vitro* experiments, collected samples, performed RNA preparation, gill culture, analysed/interpreted data, and prepared the manuscript.

 Figure legends

 Figure 1. Clock gene Ss4R duplicate pairs are divergently expressed in different tissues.

A. Phylogenetic tree highlighting WGD events and evolution of anadromy. All species

represented (excluding the lamprey, ground sharks and coelacanth) were included in the

phylogenetic identification of Atlantic salmon clock genes. B. The mammalian molecular

clock network. C. Circos plot representing the Atalantic salmon chromosomes and the large

collinear blocks of the genome that are duplicated (grey blocks/lines). Overlayed are the

chromosomal positions of all Ss4R clock ohnologues, duplicated clock genes are connected

by a coloured line. The colour of this line represents the different orthogroups. Around the

outside of the circos plot sequence similarity of the loci in a 1mb window is shown as a

coloured bar; high similarity >95% (red), medium 95-90% (blue), low ~87% (green). Sequence

similarity on a genome wide level relates to approximate rediploidization time of Atlantic

salmon chromosomes (see grey box). D. Differential regulation of Ss4R pairs in a panel of 11

different tissues. For each Ss4R pair, in each tissue, an expression divergence index (EDI)

index was calculated (EDI= abs(log2[Gene1/Gene2]). The graph shows a violin plot of the

distribution of EDI values across all tissues. The vertical black bar represents the median

value. Approximate rediploidisation time of the loci each pair is found on is represented by a

colour: red – late, blue – mid, green – early.

Figure 2. The molecular clock network regulation in the Atlantic salmon.

A. Diel and circadian regulation experiment design, arrows indicate tissue collection points.

Fish were maintained in a light dark cycle ("Diel" - short photoperiod 6 hours light: 18 dark)

transitioning onto either constant light (LL) or constant dark (DD). Sampling in diel conditions

was at a 4h resolution starting from 5 hours after lights on (Zeitgeber time - ZT5) for 24

hours. The green line represents the time-points used in the statistical analyses of diel

rhythmicity. Constant conditions are defined by circadian time (CT) which is relative to lights

on in the preceding cycle. The assessment of DD or LL rhythmicity was made using samples

from CT9 to CT29, indicated by the red and blue lines. N = 3 for all time-points.

B. Heatmap displays all diel rhythmic genes in OT, SV and gill. Overlap between tissues is

shown in the venn diagram. C. Peak-phase aligned LL and DD rhythmic genes in the OT.

Shaded area shows 95% confidence limit. D. Example duplicate comparisons from OT, SV and

gill. JKTcycle (adjP<0.05*, adjP<0.01**, adjP<0.001***, adjP<0.0001****).

Figure 3. The molecular clock network shows ohnologue specific differences during a

photoperiodically driven developmental transistion.

A. Photoperiodic gene expression experiment design. LL – constant light, SP – short

photoperiod. Time-points for sampling (n=6) indicated by arrow, sampling was conducted

once per time-point in mid-light phase, time scale shown is experimental days. B. Expression

heatmap of significantly photoperiodic (FDR<0.01) clock genes in the gill. Significant genes

cluster into five distinct expression patterns. Individual profiles are represented. Shaded area

shows 95% confidence limit where applicable. C. RNAseq counts per million profiles for

divergently regulated ohnologue pairs.

Figure 4. Stress response implicates glucocorticoid receptor signaling in clock ohnologue

regulation. A. Diel and circadian profile of plasma cortisol (n=4). Time axis is given in

continuous hours since the start of the experiment, therefore 1 to 21 represent zeitgeber

time (ZT) and 25 onwards is equivalent to circadian time (CT)1 to CT29. Due to the sampling

protocol time-points 1 and 5 are replotted from time-points 25 and 29. B. Sea-water stress

experiment design. LL – constant light, SP – short photoperiod, SW- sea-water challenge. C.

Plasma cortisol concentration in blood plasma in sea-water stress experiment (n=11-12). D.

Sea-water stress regulation of clock genes (n=6). Significantly regulated transcripts

(FDR<0.01) are shown in red. FC – fold change. E. Differential sea-water stress regulation of

ohnologue pairs. Significantly different pairs (Analysis of genes where one or both genes are

significantly regulated by seawater (FDR<0.01), then submitted to a two-way ANOVA, with

sea-water regulation and interaction, p<0.05) are shown in red. F. Predicted transcription

factor promoter binding analysis. Both sea-water induced and not-induced gene cohorts

were analysed. 50 motifs were specific to the sea-water induced cohort. The top four motifs

in each group are displayed.

Figure 5: In-vitro validation of glucocorticoid stimulated clock ohnologue expression

A. Dexamethasone-dependent transcript regulation experimental design. After photoperiod

manipulation whole gill arches (n=4) were removed and treated for 24h with glucocorticoid

receptor agonist dexamethasone (DEX), or dimethlysulphoxide (DMSO; control). B.

Comparative regulation of Cry3-Ch12/22 taken from *in vivo* sea-water stress experiment and

in vitro dexamethasone treatment. C. As in B for Nr1d2-Ch2/5 & Nr1d2-Ch14/27. D. As in B

for Tef-Ch1/28 & Tef-Ch3/6. Braches indicate phylogenetic relationship between gene sets.

S1 Figure: Tissue specific expression of clock ohnologues

A. PCA plot showing the relative tissue differences when considering clock ohnologue

expression. B. Heatmap showing the tissue specific expression of clock ohnologues.

S2 Figure: Nanostring clock gene expression and circadian phase aligned plots

A. Heatmap showing the mesor expression for each clock ohnologue in three tissues. Grey

indicates the gene is not expressed. B. Phase aligned plots for the gill. C. Phase aligned plot

for the SV. D. Arntl1-Ch10/16 comparison: plot of non-linear regression using a sixth-order

centered polynomial to fit the data and compare individual curves. P-value is the result of

extra sum-of-squares F test. E. As above for Cry3-Ch12/22.

S3 Figure: Osmoregulatory capacity during the smoltification experiment

A. Osmolality (mOsm kg-1) is displayed for fish in freshwater (FW – blue) and seawater (SW –

green) (n=6). This plot show osmoregulatory capacity develops by the two latest timepoints

(T5 and T6).

S4 Figure: Gene expression of Tsc22d3-Ch3, a positive control gene for DEX treatment

A. Gene expression of Tsc22dd3-Ch3 *in vivo* sea-water stress experiment (RNAseq counts per

million (cpm)) and B. *in vitro* dexamethasone treatment (qPCR).

S1 Table: Clock genes identified in Atlantic Salmon, orthogroups, duplicates, significances

for the circadian, smoltification and seawater challenge experiments

S2 Table: Nanostring codeset design and qPCR primers

S3 Table: Summary of previous studies measuring cortisol in fish

S4 Table: Cortisol source data for Figure 4A

S5 Table: SalmotifDB results – transcription factor binding site analysis

S6 Table: qPCR source data for Figure5

S1 appendix: Evolutionary gene trees for circadian clock genes

S2 appendix: Nanostring circadian profiles for all genes

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Supplementary Fig. 1

Supplementary Fig. 2

Supplementary Fig. 3

A.

Tsc22d3_Ch3

B.

Tsc22d3_Ch3

Supplementary Fig. 4