Cooperative genetic networks drive a mammalian cell state transition

Andreas Lackner[#1], Robert Sehlke[#2], Marius Garmhausen[#2], Giuliano Giuseppe Stirparo[#3], Michelle Huth[$1], Fabian Titz-Teixeira[$2], Petra van der Lelij[1], Julia Ramesmayer[1], Henry Fabian Thomas[1], Meryem Ralser[3], Laura Santini[1], Mihail Sarov[4], A. Francis Stewart[4,5], Austin Smith[3], Andreas Beyer*[2], Martin Leeb*[1]

[1] Max Perutz Laboratories Vienna, University of Vienna, Vienna Biocenter, Dr Bohr Gasse 9, 1030 Vienna

[2] CECAD, Faculty of Mathematics and Natural Sciences, Medical Faculty Cologne, University of Cologne, Joseph-Stelzmann-Str. 26 50931 Köln, Germany

[3] Wellcome - MRC Cambridge Stem Cell Institute, University of Cambridge, Cambridge CB2 0AW, UK

[4] Max Planck Institute of Molecular Cell Biology and Genetics, Pfotenhauerstraße 108

01307 Dresden, Germany

[5] Biotechnology Center, Center for Molecular and Cellular Bioengineering, Technische Universität Dresden, Tatzberg 47, 01307 Dresden, Germany

[6] Center for Molecular Medicine (CMMC), University of Cologne, Cologne, Germany.

# Equal contribution

$ Equal contribution

* Corresponding authors

Abstract

In the mammalian embryo, epiblast cells must exit their naïve state and acquire formative pluripotency. This cell state transition is recapitulated by mouse embryonic stem cells (ESCs), which undergo pluripotency progression in defined conditions *in vitro*. However, our understanding of the molecular cascades and gene-networks involved in the exit from naïve pluripotency remains fragmented. Here we employed a combination of genetic screens in haploid ESCs, CRISPR/Cas9 gene disruption, large-scale transcriptomics and computational systems-biology to delineate the regulatory circuits governing naïve state exit. Transcriptome profiles for 73 knockout ESC lines predominantly manifest delays on the trajectory from naive to formative epiblast. We find that gene networks operative in ESCs are active during transition from pre- to post-implantation epiblast *in utero*. We identified 374 naïve-associated genes tightly connected to epiblast state and largely conserved in human ESCs and primate embryos. Integrated analysis of mutant transcriptomes revealed funneling of multiple gene activities into discrete regulatory modules. Finally, we delineate how intersections with signaling pathways direct this pivotal mammalian cell state transition.

1 Introduction

Mouse embryonic stem cells (ESCs) can self-renew in defined conditions in a ground state of

naïve pluripotency [1]. The ESC exit from naïve pluripotency provides an amenable

experimental system for dissection of a cell fate decision paradigm [2, 3]. Naïve pluripotency is

under control of a gene regulatory network (GRN) containing the core pluripotency

transcription factors (TFs) *Pou5f1*, *Sox2* and naïve-specific TFs like *Nanog*, *Esrrb*, *Klf4* and

others [4-6]. In defined cell culture conditions that include inhibitors against Mek1/2

(PD0325901) and Gsk3 (CHIR990201, CH; collectively termed '2i'), ESCs can be homogenously

maintained in the naïve state [7]. Within 24 to 36 hours after withdrawal of 2i, ESCs transit into

formative pluripotency, entirely losing naïve identity [2]. During this transition, the naïve GRN

is extinguished and expression of formative factors like *Otx2*, *Pou3f1*, *Dnmt3a/b* and *Fgf5* is

initiated. A similar transition is evident during peri-implantation development, where the TF-

network maintaining naïve pluripotency dissolves between embryonic day (E) 4.5 and E5.5 [8-

10].

The speed of the naïve to formative GRN transition is notable because: i) the cell cycle is

around 12 hours long; ii) all factors that are required to establish and maintain naïve

pluripotency are expressed robustly in naïve cells; and (iii) the naïve pluripotency network is

recursively self-reinforcing. The rapid dissolution of naïve pluripotency implies the existence

of circuit-breaking mechanisms. In recent years, we and others have identified various factors

promoting ESC differentiation using screens in haploid and diploid ES cells [11-15].

Robust assays employing ESCs expressing a Rex1-promoter driven destabilised GFP reporter

ESC line (Rex1::GFPd2) enable the dissection of the exit from naïve pluripotency in high

resolution [2, 16]. Rex1-GFP downregulation is initiated within 24h after 2i withdrawal (N24) and

completed after 48h (N48). Nevertheless, the exact nature, mechanistic underpinnings and

sequence of events during exit from naïve pluripotency remain only partially understood. In

particular, we lack insight into how the different molecular components of the system co-

operate to elicit proper cell fate transition.

Here, we have driven a Rex1-GFP reporter screen to saturation, thus providing an extensive

list of genes and pathways involved in the exit from naïve pluripotency. We utilized this

information in a systems biology approach to explore regulatory principles of the exit from

naïve pluripotency. To evaluate dependencies and causal relationships within the

pluripotency and differentiation circuitries, we probed the response of the differentiation

program to a comprehensive series of exit factor gene knockouts. Through computational

integration of molecular profiling data with regulatory networks and *in vivo* GRN-trajectories,

we expose the regulatory foundations of a cell fate choice paradigm at a pivotal junction in

early mammalian development.

Results

Haploid ES cell saturation screen

Haploid ES cells are an efficient platform for insertional mutagenesis-based screens [11, 17, 18]. We previously reported a medium-scale screen comprising approximately 5×10^4 mutagenic events to identify factors regulating the exit from naïve pluripotency [12]. We have now driven this approach to saturation by assaying approximately 1.2 million mutations in receptive genomic regions that cause delays in Rex1 downregulation in two independent Rex1-reporter cell lines (Figure 1ab), utilizing three different mutagenic vectors in 35 independent screens (Figure 1cd).

Stringent filtering resulted in a candidate list comprising 489 genes (Supplementary Table 1). Reassuringly, the known exit from naïve pluripotency regulators *Tcf7l1, Fgfr2, Jarid2* and *Mapk1* (*Erk2*) [19] were among the highest ranked (Supplementary Figure 1a). The candidate hit genes were enriched for processes involved in transcription regulation, epigenetics and signalling-related functions (Supplementary Figures 1bc; Supplementary Table 1), as well as RNA binding functions in line with emerging evidence of RNA regulatory mechanisms in cell fate control [20].

These screens generated a candidate inventory of the machinery that mediates exit from naïve pluripotency. Many identified genes are not specific to pluripotency and have functions in common pathways and processes, implying that the exit from naïve pluripotency utilizes

widely expressed cellular machinery. Therefore, mechanisms mediating ESC transition might

also be utilised in other differentiation processes.

Establishment of a mutant ESC library for systematic transcriptional profiling

To characterise deficiencies in naïve exit, we generated KO ESC lines deficient for 73 selected

genes, comprising top ranked genes from the mutagenic screen together with components

from pathways and protein complexes for which multiple members were recovered, even if

just below the cut-off threshold (e.g. the Paf complex member *Leo1*, the mTORC1 regulator

Tsc2 and the NMD component *Smg6*), and *Mbd3*, *Zfp281* and *L3mbtl3* as known players in

the exit from naïve pluripotency [14]. Three control genes were included that are either not

expressed in ES cells (Nestin), expected to be neutral (Hprt), or whose ablation was expected

to accelerate differentiation (cMyc). Paired gRNAs were used to disrupt target genes in a

diploid <u>R</u>ex1::GFPd2 reporter ES cell line carrying a <u>Cas9</u>-transgene (henceforth termed RC9

cells) (Figure 1a). Following an efficient parallelized approach, we established passage

matched and isogenic cell lines, thus maximising comparability (Supplementary Figure 1de).

Full protein deficiency was validated for 14 KOs (*Eed, Suz12, Jarid2, Kdm6a, Smg5, Smg6,*

Smg7, Tsc2, Pten, Raf1, Tcf7l1, Leo1, Nmt1, and *Csnk1a1*), while reduced protein expression

in concordance with a heterozygote phenotype was confirmed for *Mapk1* (Supplementary

Figure 1f). For five further genes (*Alg13, Dido1, Msi2, Etv5, Jmjd1c*) we confirmed absence of

the corresponding transcripts or specific out of frame deletion of an exon by RT-qPCR or

Sanger sequencing of RT-PCR products. Successful rescue experiments using 3xflag-tagged

transgenes for six genes (*Rbpj, Etv5, Fgfr1, Jarid2, Mbd3,* and *Tcf7l1*) established causality

between the observed genotype and phenotype (Supplementary Figures 1gh). Thus, all tested

knockouts showed the expected impact on RNA or protein expression. However, we cannot

exclude the possibility of hypomorphic phenotypes in some cases.

ESC differentiation behaviour is highly dependent on cell density and timing of medium

changes. To enable robust comparison of the differentiation of multiple KO ESC lines in

parallel, we performed differentiations batch wise in duplicates, always including WT ESCs

and negative controls across seven experiments. At N24 we assayed the differentiation status

by FACS analysis (Figure 1e) and extracted RNA for transcriptome analysis.

The exit machinery is already poised in 2i

Batch corrected RNA-seq data comprising 14 replicates of WT ES cells indicated 3058

differentially expressed genes (DEG) between 2i and N24 (H0: |FC| < 1.5, FDR ≤ 0.05; Figure

1f, Supplementary Figure 1i,j and Supplementary Table 1). Interestingly, most of the 489

genes identified in the haploid screens including the 73 genes selected for KO did not change

significantly in transcript expression between 2i and N24 and were not present in the list of

DEG (Figures 1f and S1K), with only *Jarid2*, *L3mbtl3*, *Pten* and *Zfp423* showing upregulation at

N24 (FC ≥ 2; adj.p. ≤ 0.05). This implies that the exit machinery is already embedded in the

ground state and ESCs are poised for rapid decommissioning of naïve identity and entry into

differentiation [19].

To facilitate interrogation of the KO gene expression datasets, we developed an interactive

online   tool   (GENEES - Genetic Network Explorer for Embryonic Stem Cells –

http://shiny.cecad.uni-koeln.de:3838/stemcell_network/;   User: stemcells;   Pw:

S3m7KmIfHr9).

Using t-Distributed Stochastic Neighbour Embedding (t-SNE), we visualized similarities

between KOs based on expression of the DEG in 2i and at N24. We observed clustering of

members of the same complex or pathway: *Eed-Suz12* (PRC2), *Ptpn11-Raf1-Fgfr1-Etv5*

(Fgf/ERK), *Smg5-Smg6-Smg7* (NMD; nonsense mediated decay), *Mta3-Mbd3* (NuRD), *Pten-*

Tsc2 (mTorc1 signaling) (Figure 1gh). The transcription profiles of the KO ESCs clustered by

culture condition (2i and N24) and to a lesser extent by genotype (Supplementary Figure 1i).

This is consistent with the phenotypic observation that despite manifesting differentiation

delays at N24, all of the KO ESCs ultimately departed from the naïve state during longer

differentiation timecourses, as measured by loss of Rex1GFP. Furthermore, even KOs that

showed extensive Rex1GFP downregulation delays at N24 displayed transcription profiles

that were globally adjusted towards differentiation. Therefore, the knockout of a single gene

is not sufficient to permanently block exit from naïve pluripotency in culture, in accord with

the finding that ternary depletion of *Tcf7l1*, *Etv5* and *Rbpj* is required for sustained self-

renewal in the absence of 2i or LIF [21].

The exceptional role of *Csnk1a1* and the involvement of compensatory mechanisms

The *Csnk1a1* KO is an exception to this generalization, clustering with 2i samples at N24

(Supplementary Figure 1i). Csnk1a1 is a serine threonine kinase and a component of the beta-

catenin destruction complex. Although KO of another destruction complex member *Apc*, or

of the downstream repressor *Tcf7l1* resulted in the upregulation of similar gene-sets

(Supplementary Figure 1l), we observed stronger differentiation defects and larger amplitude

of gene deregulation in two independently derived *Csnk1a1* mutants. However, these

mutants also exhibited markedly reduced population doubling rates, which was not shown by

Apc or *Tcf7l1* mutants. Upon continuous culture in 2i medium (~5 passages), proliferation was

restored in *Csnk1a1* KO cells and differentiation potential was regained, suggesting up-

regulation of compensatory mechanisms and a likely effect of proliferation rate on

differentiation kinetics. Csnk1a1 can be chemically inactivated using Epiblastin A, [22]. We found

that Epiblastin A partially blocked exit from naïve pluripotency, without affecting proliferation

within the duration of the assay (Supplementary Figure 1m). A second case of phenotype

adaptation was observed with Pum1. *Pum1* KOs showed pronounced differentiation delays

during early passages (Supplementary Figure 1n), as also seen for acute Pum1 depletion by

siRNA and in previously generated CRISPR KO ESCs [12]. However, the phenotype was lost in

later passages and *Pum1* KO cells showed WT like expression-levels of naïve marker genes

(Figures 2a and S1a).

Robust feedback wiring in the naïve TF network

Interestingly, the transcriptome data revealed that exit factors do not, in general, reduce

naïve transcription factor (TF) expression in the ground state (Supplementary Figure 2a). *Rbpj*

and *Trim71* KO resulted in moderate increases in *Klf4*, while the aforementioned *Csnk1a1* KO

ESCs showed limited up-regulation of both *Klf4* and *Tbx3* in 2i. Other KOs had only minor

effects on the naïve network in 2i. These data are consistent with robust feedback wiring in

the naïve TF network [4, 5] and neutralisation of most 'differentiation drivers' in 2i in culture [23].

In contrast, we observed a more extensive impact on formative markers. Several KO cell lines

showed lower baseline expression in 2i of *Otx2, Fgf5, Dnmt3a/b* and *Pou3f1* (Oct6)

(Supplementary Figure 2b). In line with recent results, we noted that depletion of several

Fgf/ERK components resulted in reduced *Dnmt3a/b, Pou3f1* and *Fgf5* expression in 2i [21].

Although Fgf/ERK signalling is effectively inhibited in 2i [7], our data suggest that either residual

pathway activity or potential moonlighting functions of pathway components mediate poised

expression of the formative pluripotency programme in 2i.

Clustering based on the expression of ten naïve pluripotency marker genes showed that the

downregulation of the naïve pluripotency TFs during formative differentiation is defective

across multiple KOs (Figure 2a). Although expression of the naïve TFs was highly correlated,

Klf2 appeared to be an exception. *Klf2* downregulation was notably impaired in *Tcf7l1* KO

ESCs, whereas it was unaffected by several KOs, including *Jarid2,* despite a comparable extent

of deregulation of most other naïve marker genes*.* This indicates that *Klf2* expression can be

uncoupled from the core naïve network. Forced *Klf2* expression stabilizes self-renewal [24, 25].

Klf2 depletion destabilizes mouse ESC identity [26] and increases the speed of Rex1-GFP

downregulation upon 2i withdrawal (Supplementary Figure 2c). The differentiation delay in

Tcf7l1 KO ES cells is partially dependent on *Klf2* (Figure 2b), consistent with direct regulation

[27]. Conversely, *Klf2* depletion in *Jarid2* KOs did not restore differentiation timing, indicating

that separable gene-networks contribute to dismantling naïve pluripotency.

Effects of exit gene depletion on global gene regulation

We used two measures to gauge differentiation delays in KOs compared to WT cells: first, the

average expression levels of a set of naïve marker genes (defined throughout this manuscript

as the average deregulation of the naïve marker genes *Esrrb, Nanog, Tfcp2l1, Tbx3, Prdm14*

and *Klf4* at N24); second, the extent of global transcriptome adjustments between 2i and N24.

Generally, the delayed extinction of the naïve TF network in KOs at N24 was accompanied by

reduced gene expression change between 2i and N24 (R = -0.91; Figure 2c). However, some

KOs deviated from this pattern. *Alg13, Jmjd1c* and *Jarid2* mutants showed a less pronounced

change in naïve marker gene down-regulation than expected based on the reduced level of

global gene expression adjustments towards an N24 profile. *Trim71* KO cells in contrast

displayed modest global transcription changes but a more evident effect on naïve TF

expression levels. Thus, the RNA binding protein Trim71 appears to be focused on the

regulation of naïve pluripotency TF genes, although the ensuing exit delay is modest.

A general correlation was observed between the numbers of genes deregulated in 2i and at

N24 (Figure 2d). Mutants with the strongest impact on the transcriptome (e.g. *Csnk1a1, Eed,*

Suz12, Zfp281 and Smg5) showed deregulation of several thousand genes both in 2i and at

N24. *Fgfr1* and *Tcf7l1* KOs were two exceptions to this correlation with relatively few genes

deregulated in 2i, but several hundred at N24. We surmise that the effect of these two genes

in 2i is largely masked because they are in the same pathways as the inhibitor targets Mek1/2

(PD0325901) and Gsk3 (Chiron 990201) [7, 23]. Defective differentiation could simply equate

with the extent of overall gene deregulation. We therefore mapped the average deregulation

of naïve marker genes onto Figure 2d. Most KOs showing the strongest deregulation of naïve

marker genes correlated with large-scale gene deregulation (both in 2i and at N24). However,

there were several exceptions: for example, KO of *Eed*, *Suz12*, *Trim71*, and *Ctbp2* affected a

substantial number of genes, but caused naïve marker deregulation that was weaker than

expected assuming a direct correlation between the number of deregulated genes and the

differentiation phenotype. Vice versa, KO of *Mbd3, Cdk8* or *Pten* showed marked naïve

marker downregulation delays, despite a relatively mild impact on overall transcription

(Figures 2d and Supplementary Figure 2d).

Transcriptome analysis reveals a genetic interaction between *Jmjd1c* and *Tcf7l1*

The putative histone 3 lysine 9 (H3K9) demethylase *Jmjd1c (Kdm3c)* KO is a case of particular note. Few genes were significantly deregulated at N24 (Figure 2d), with a mild global neutralisation of differentiation-induced transcriptome changes (Figure 2c). Neither the role of *Jmjd1c* in pluripotency regulation, nor its mode of action are known. We used multiple regression analysis to determine the similarity of the *Jmjd1c* KO to all other KO RNA-seq profiles. At N24, the *Jmjd1c* KO transcriptome was most similar to the *Tcf7l1* KO profiles (Supplementary Figure 2e), suggesting a potential functional connection between *Jmjd1c* and *Tcf7l1*. Indeed, Jmjd1c has recently been reported as a high-confidence protein-interactor of Tcf7l1 [28]. The 2i component Chiron 99021 is a specific inhibitor of Gsk3 and phenocopies deletion of *Tcf7l1*, which is the downstream repressor in ESCs [29, 30]. Accordingly, Chiron 99021 delays the differentiation of WT ESCs, but had little additional effect on *Tcf7l1* KOs (Figure 2e). *Jmjd1c* deficient ESCs did not show a discernible phenotype at N72, in line with minimal deregulation of the naïve TF network. Addition of Chiron 99021 had a stronger than expected effect on Jmjd1c KOs and resulted in a synthetic enhanced delay phenotype (Figure 2e), suggesting a cooperative activity of *Jmjd1c* and the Wnt/Tcf7l1 axis in the exit from naïve pluripotency.

Relative quantification of differentiation delays *in vitro*

To quantify differentiation delays, we obtained RNA-seq data from a 2h-resolved WT ESC differentiation time course (Figures 3a and Supplementary Figure 3a). We then compared the expression patterns of specific gene-sets in a given KO at N24 to the expression of the same gene-set along the WT 2i to N32 trajectory. This enabled us to 'position' each KO along the trajectory and thus quantify the differentiation delay with a precision of about 4h (Figure 3b,

Supplementary Table 2). In a complementary approach, we used all 3058 DEG between 2i and

N24 as an alternative reference set. This yielded similar, but on average slightly less

pronounced differentiation delays (Figures 3c and Supplementary Figure 3b), supporting the

hypothesis that the naïve TF network is regulated in concordance with but partially

independently from the rest of the transcriptome during exit from naïve pluripotency.

The regulatory program for pre- to post-implantation epiblast transition is preserved in

most mutants

To explore the extent to which our *in vitro* data captures *in vivo* regulation, we first compared

global transcriptomes of WT and KO ESCs to single cell transcriptome data from the *in vivo*

pre- (E4.5) and post-implantation (E5.5 and E6.5) mouse epiblast [10] (Figure 4a,B). As expected

[8], WT and KO cells in 2i showed greater similarity (fraction of identity - FOI) to the E4.5 than

to the E5.5 epiblast. Cells at N24, in contrast, showed higher FOI to the E5.5 epiblast. Cells in

2i and at N24 showed low FOI to E6.5 epiblast, underscoring that 24h 2i withdrawal models

the E4.5 to E5.5 epiblast transition [2]. Interestingly, some KO ESCs showed expression features

in 2i with increased similarity to the *in vivo* epiblast (Figures 4c,d and Supplementary Table

3). Among those were KOs showing strong *in vitro* differentiation defects, such as *Zfp281,*

Tsc2 and *Etv5,* but also *Trim71* and *Smg7*, which showed only a modest differentiation delay

in culture.

At N24 several KOs, such as *Zfp281*, *Tsc2*, *Smg5* and *Smg6*, retained strong similarity to the

E4.5 epiblast (Figures 4c,e and Supplementary Figure 4a). Strikingly, *Zfp281* KO profiles at N24

showed similarity to the E4.5 epiblast on par with WT cells cultured in 2i (Figures 4c and

Supplementary Figure 4a), consistent with an overt differentiation delay phenotype [31].

Overall, there was a good correlation between the *in vitro* differentiation delay and the

similarity with pre-implantation epiblast cells: the N24 KO transcriptomes that were more

similar to the E4.5 epiblast exhibited stronger exit-delay phenotypes (Figure 4e,

Supplementary Figure 4b). Together, this indicates that similar transcriptional networks are

regulated by similar mechanisms during *in vivo* and *in vitro* transitions to formative

pluripotency. However, at N24 three KOs (*Jarid2*, *Tcf7l1*, and *Trp53*) showed a similarity to

the E4.5 epiblast which was smaller than expected based on their *strong in vitro*

differentiation defects (Figures 2a and 4e).

We then performed principal component analysis (PCA) for genes variably expressed in

embryo development, using averaged values for inner cell mass (ICM), primitive endoderm

(PrE) and E4.5, 5.5 and 6.5 epiblast [10]. The PCA separated the 2i and N24 transcriptomes into

two clusters with proximity to PrE/EPI and E5.5 respectively. Notably*, in vitro* differentiation

defects were reflected by closer proximity of mutant N24 profiles to the E4.5 epiblast (Figure

4f). Overlaying the PCA with a colour gradient indicating similarity to PrE shows that multiple

KOs in 2i gained similarity to PrE (Supplementary Figure 4d). Concordantly, the expression of

PrE-defining transcription factors *Gata4*, *Gata6*, *Sox7* and *Sox17,* was readily detectable in

Eed, Suz12, Zfp423 and *Csnk1a1* KOs in 2i (Supplementary Figure 4e). A shift toward PrE-

identity was most pronounced for the PRC2 KOs *Eed* and *Suz12*. Notably, this group of KOs

(Csnk1a1, PRC2, Zfp423, which exhibited substantial *in vitro* differentiation defects while

showing reduced similarity to the E4.5 epiblast, also showed decreased similarity between

their N24 profiles and the E5.5 epiblast (Supplementary Figure 4d). We therefore suggest that

deregulation of the PrE-programme above a certain threshold has disengaged these KOs from

the normal embryonic developmental trajectory. In contrast KOs showing relatively increased

similarity to the E4.5 in 2i and decreased similarity to the E5.5 epiblast at N24 are deficient

for genes regulating the formative transition during development (e.g. *Zfp281*, *Etv5*, *Rbpj* and,

Trim71, Supplementary Figure 4d).

Correspondence to primate embryogenesis

Despite morphological and timing differences between rodent and primate peri-implantation

development, embryos of both orders appear to transit through similar pluripotency states [1,

32]. To examine this issue, we compared transcriptional profiles of our KO series and cells of

the macaque *in vivo* naive- and post-implantation epiblast [33]. In general, ESCs in 2i were more

similar than cells at N24 to the pre-implantation macaque epiblast (Figure 4g).

Correspondingly, N24 cells were closer to the macaque post implantation epiblast.

Interestingly, KOs showing higher identity with the mouse *in vivo* epiblast also displayed an

increased similarity with the macaque pre-implantation epiblast (Supplementary Figures

4f,h). We also compared our datasets to data obtained from *in vitro* capacitation during

human naïve to primed ESC differentiation [34]. KOs displaying strong differentiation defects in

mouse cell culture had higher correspondence to naïve human ESCs and, at N24, lower

correspondence to primed human ESCs (Figures 4h and Supplementary Figure 4g,I). These

observations suggest that the overall GRN-redeployment during the naïve to formative

transition is conserved between rodents and primates.

Identification of an extended naïve pluripotency network

ESC differentiation requires fine-tuned coordination between extinction of the naïve and

initiation of the formative transcription networks. To date, only incomplete inventories of the

genes that functionally define these two states have been made. These genes cannot be

defined based simply on differential expression between 2i and N24, because not all DEG will

be functionally linked to the naïve GRN. To identify those genes that show specific linkage to

the core naïve pluripotency network, we trained regression models to predict expression

changes across all KOs (in 2i and at N24) as a function of naïve marker log-fold-changes. The

374 genes whose expression was tightly correlated ($R^2 > 0.7$) with one or several of the seven

core pluripotency markers (*Nanog, Esrrb, Tbx3, Tfcp2l1, Klf4, Prdm14 and Zfp42)* were defined

as 'Naive-Associated Genes' (NAGs; Supplementary Table 3). The naïve pluripotency TFs not

represented in the defining TF-set were correctly identified as NAGs, including *Klf5, Nr0b1*

and *Nr5a2*. However, *Klf2* is not one of the NAGs being only weakly associated ($R^2=0.53$) with

the naïve network. This concords with our earlier observation that *Klf2* expression can be

uncoupled from the naïve TF network. Of further note, *Klf2* expression is barely detectable in

marmoset or human pre-implantation epiblast cells (Supplementary Figure 5b).

The NAG showing the highest correlation with the naïve core network is *Pdgfa*, which is

relatively highly expressed in ES cells (FPKM ~70). *Pdgfa* has no known role in ESC self-

renewal, but functions in segregation of the primitive endoderm [35]. To examine whether the

link between *Pdgfa* and the naïve transcription factor network is maintained *in vivo*, we

utilized GRAPPA, a tool to visualize single cell expression data from preimplantation

development [36]. Indeed, *Pdgfa* is uniformly expressed in the E4.5 epiblast (Supplementary

Figure 5c). Interestingly, its cognate receptor *Pdgfra* is neither expressed in ES cells nor the

naïve epiblast (FPKM in WT ESCs <0.5), but specifically marks the neighbouring primitive

endoderm at E4.5 [37].

We surveyed the expression of NAGs at the transition from naïve to post-implantation

pluripotency in the single cell RNA-seq datasets from E4.5, E5.5 and E6.5 epiblast cells [10]. We

detected a clear enrichment of NAGs within genes that separate the pre- from the post-

implantation epiblast in the differentiation-state resolving dimension of a principal

component analysis (PCA) (Figures 5a and Supplementary Figure 5d), highlighting that NAGs

are a set of indicators for the naïve epiblast state *in vitro* and *in vivo*. Strikingly, NAGs showed

a near identical fold change behaviour *in vitro* and *in vivo* during the epiblast transition from

E4.5 to E5.5 (Figure 5b,e,f). Notably, this effect was not observed for the top 250 differentially

expressed genes in ESC differentiation *in vitro*. Furthermore, NAG orthologues showed

significantly co-regulated expression dynamics during macaque pre- to post-implantation

epiblast differentiation *in vivo* (Figure 5c,e,f) and during capacitation of human ESCs *in vitro*

(Figures 5d,e,f), which further underscores their relevance. We thus propose that NAGs

constitute an integral component of the naïve transcriptional network with potential

functional relevance *in vivo* across mammalian species.

Deregulation of signaling cascades is a hallmark of differentiation delay

We next analysed the extent of deregulation of five key signalling pathways known to be

active in pluripotent cells: LIF/Stat3, mTOR, Wnt/β-catenin, Fgf/ERK and Notch [7, 38-42]. Changes

in the activity of a signalling pathway are not necessarily reflected in expression changes of

pathway member transcripts. Thus, in order to quantify pathway activities, we employed

'expression footprints'. We identified pathway specific marker gene-sets, each containing 50

genes, reporting pathway activity changes. These marker sets were determined using N24

derived KO transcription-profiles of ESC lines deficient for key signalling pathway

components. Thereby we defined key mTOR (affected by *Tsc2 KO*), Wnt/β-catenin signaling

(*Tcf7l1 KO*), Fgf/ERK (*Fgfr1* and *Ptpn11 KOs*), and Notch (*Rbpj KO*) pathway targets

(Supplementary Figure 6a, Supplementary Table 4). Overlaps with available Tcf7l1 and Rbpj

ChIP and activity profiles [27, 43] supported the reliability of this approach (Supplementary Figure

7E). For LIF-signaling, we compared RC9 ESCs grown in 2i in the presence and absence of LIF

for 24 hours. This resulted in a list of LIF-sensitive genes including known targets like *Socs3,*

Gbx2, Junb, Tfcp2l1, Klf4 and *Klf5* [27, 44] (Supplementary Table 4). In summary, the expression

footprints present non-overlapping sets of marker genes whose expression state is indicative

of the activity of the respective signaling pathway.

Using the expression footprints, we then asked whether we could detect a preferential

deregulation of one or more of these signalling footprints in specific KOs either in 2i (for LIF)

or at the N24 time-point (for all other pathway-profiles) (Figure 6a, Supplementary Table 4).

Surprisingly, we detected mis-regulated LIF target genes in several KO ESC lines cultured in 2i

in the absence of LIF (Figure 6a). Activation of such a 'LIF-like'-profile in 2i was closely

correlated with the extent of differentiation delay observed at N24 (Supplementary Figure

6b). Notably, presence of LIF together with 2i before induction of differentiation slows down

naïve state exit (Supplementary Figure 6c) [4, 16]. *Ptpn11, Zfp281, Tsc2* and *Trim71* KO ESCs

showed the greatest similarity to the LIF profile in 2i; in contrast, *Tcf7l1* KO ESCs, despite

showing a pronounced differentiation defect, lack the 'LIF footprint' (Supplementary Figure

6b). Addition of a JAK inhibitor to several KOs showing a LIF-like profile resulted in no or only

minor amelioration of the differentiation defect. This suggests that these KOs do not directly

activate Jak/Stat signalling, but that a LIF-like expression profile reflects a consolidated naïve

network that is resilient to dismantling.

As expected, *Raf1*, *Fgfr2*, *Mapk1* and *Etv5* KO ESCs showed an Fgf-signalling footprint at N24.

Mbd3, *Mta3*, *Nsd1* and *Arid5b* KOs showed Fgf/ERK target deregulation similar to reference

KOs, indicating an involvement of chromatin regulators in ERK target-gene control (Figure 6a).

Tsc2 deficiency lead to constitutive activity of the mTORC1 pathway, which has previously

been associated with an exit from naïve pluripotency phenotype [45]. Many KOs (including *Pten*)

exhibited deregulated Tsc2 responsive genes, suggesting that regulation of mTORC1 (or its

downstream effectors) is a common node to gate the exit from naïve pluripotency. *Tsc2*

depletion resulted in partial activation of a LIF footprint, suggesting that a component of the

LIF response is mediated through Akt [39, 46]. *Rbpj* and *Tcf7l1* KOs did not show a correlation

with each other or with Tsc*2* KO, consistent with independent mechanisms [21]. The beta-

catenin destruction complex member *Apc* and the previously discussed *Jmjd1c* showed the

expected similarities with the *Tcf7l1* profile. Interestingly, several KOs with strong phenotypes

showed footprints similar to *Tcf7l1* KOs at N24. This is in line with evidence that ⍰-catenin

constitutes or regulates a major differentiation switch during the exit from naïve pluripotency

and is influenced by multiple exit KO genes [29, 47]. Genes downstream of the Notch pathway

were most strongly affected in a group of mutants deficient for mRNA homeostasis (*Smg5*

and *Smg6*) and chromatin regulators, including *Jarid2, L3mbtl3 and Mbd3*. The latter suggests

an interaction between the Notch pathway (*Rbpj*) with the Polycomb and Nurd complexes to

modulate network rewiring during the exit from naïve pluripotency. Cooperativity between

Rbpj and the Polycomb-associated protein L3mbtl3 has been reported in *Drosophila* and *C.*

elegans [48].

We asked to what extent the alteration of any of the five signalling pathways was predictive

for the strength of the differentiation defect. We found that whereas aberrant Fgf/ERK or

mTORC1 activity were unreliable predictors, transcriptional similarity to *Rbpj* or *Tcf7l1* KOs at

N24 or deregulation of LIF target genes in 2i correlated with differentiation delay

(Supplementary Figure 6d). These data indicate that many KOs with a delay phenotype have

altered activity of at least one of the five key signalling pathways. Strikingly, however, there

was no single signalling pathway perturbed downstream of all KOs. We observed that KOs

showing multiple pathway footprints are rare, but most KOs showing a delayed naïve marker

gene downregulation at N24 deregulate at least one specific pathway. *Trp53*, *Ncor1* and

Usp9x constitute exceptions, which show appreciable differentiation defects without

deregulating of any of the tested signalling cascades. We conclude that signalling pathways

are harnessed in parallel and that most KO genes contribute directly or indirectly to at least

one pathway activity that promotes the exit from naïve pluripotency.

Identification of gene-networks downstream of multiple KO genes

To explore common regulatory targets of multiple KO-factors, we classified deregulated genes

into two groups: (i)**,** deregulated by a KO in both 2i and at N24, termed 'constitutive KO

response'; (ii) not, or only weakly, deregulated by a KO in 2i, but significantly deregulated at

N24, termed 'N24 induced KO response'. Genes in both groups are affected by knockouts, but

genes in the second group deviate from wild-type expression only under conditions of

differentiation. We identified all genes that showed either type of behaviour in at least one

KO (Supplementary Figure 6e; Supplementary Table 5). Genes deregulated exclusively in PRC2

core-components (*Eed* or *Suz12* KOs) or *Csnk1a1* KOs were excluded from further analysis,

because the deregulation of thousands of genes in these KOs had a disproportionate impact

on the resulting gene-lists. In total, the constitutive and the N24 induced KO response groups

contained 1,837 and 720 genes, respectively. Genes belonging to the N24 induced KO

response group were strongly enriched for factors that are dynamically regulated during

normal differentiation, whereas genes of the constitutive response group were not

(Supplementary Figure 6f). Thus, naïve marker genes were absent from the constitutive

group. Together with NAGs, they were strongly enriched in the N24 induced group, because

deregulation of the naïve network only initiates upon 2i withdrawal.

To characterize the constitutive KO response genes, we grouped them based on their log-fold-

changes between KO and WT at N24. This resulted in twelve clusters, ranging in size from 20

to 597 genes (Supplementary Table 5) (Figure 6b). By Gene Ontology (GO) analysis, eight of

the twelve clusters showed a significant enrichment of at least one GO term containing more

than five genes (Figure 6c, Supplementary Table 5). Many of these GO terms relate to cell fate

specification and development. Notably, the constitutive response clusters often shared

identical or similar GO terms, suggesting functional convergence. NAGs were

underrepresented in constitutive clusters and only contained in clusters 1 (const c1) and 2,

for which expression profiles across all KOs correlated with naïve marker expression (Figure

6b). Therefore, deregulation of genes within const c1 and c2 in 2i is proposed to directly

translate to differentiation defects observed at N24. Cluster 1 genes were enriched for

general differentiation-related factors and response to Bmp signalling. Constitutive cluster 2

contained a significant number of Fgf/ERK pathway defining genes (Supplementary Figure 7e)

and was consistently enriched for Fgf/ERK signalling related terms (Supplementary Table 5).

Intriguingly, constitutive cluster 10 contained multiple *Zscan4* isoforms [49] and 14 out of 45

genes in this cluster are bound by the co-repressor Trim28, which is involved in silencing of

endogenous retroviruses [50-52]. Strong upregulation of this cluster was observed in *Csnk1a1*

and *Zfp281* KOs, but strong downregulation in *Rbpj* and *Jarid2* KOs. Overall, cluster 10

regulation showed no strict correlation with a differentiation delay phenotype.

Analysis of the N24 induced response genes yielded 12 clusters ranging from 8 to 103 genes

(Figure 6d and Supplementary Table 5). All naïve marker genes were detected in N24 induced

cluster 1 (N24i c1), with the notable exception of *Klf2*. *Klf2* was part of N24 induced cluster

10, within which more than half the genes are Tcf7l1 ChIP targets [27] (Supplementary Figure

7E). This provides further support for the notion that Wnt/Tcf7l1 controls *Klf2* expression

independently of the rest of the naïve network. Interestingly, twelve percent of genes in N24

induced cluster 10 were screen hit genes, suggesting a major impact of this cluster on the exit

from naïve pluripotency (Supplementary Figure 7e). NAGs were distributed in eight out of the

twelve N24 induced clusters. In marked contrast to the constitutive KO response, individual

N24-induced clusters displayed specific functional enrichments (Figure 6e). Therefore, the

N24 induced response clusters represent distinct molecular pathways and cellular functions

that were altered in multiple KOs. N24 induced cluster 7 was significantly regulated in *Etv5*,

Trim71 and *Zfp281* KOs at N24 (Figure 6d). Cluster 7 contains pivotal cell fate switch genes

like *Fgf4*, *Tfe3* [14] and *Tcf7l1* itself, but NAGS were underrepresented in this cluster. Notably,

Etv5, Trim71 and *Zfp281* are the three KO lines that show the strongest exit delay while

retaining E4.5 identity at N24 (Supplementary Figure 4d), suggesting that N24 induced cluster

7 contains crucial cell fate switch genes that are modulated during *in vitro* and *in vivo* exit

from naïve pluripotency. In summary, constitutive and N24 induced clusters define crucial

genetic modules that are co-regulated by multiple exit factors.

We asked whether the expression status of these genetic modules correlated with the activity

of differentiation regulating pathways. We utilized pathway activity levels and cluster specific

expression levels across all KOs to build a model predicting the impact of pathway activity on

cluster-gene expression (Figures 7A and S7A to F). Adding further evidence to the pathway to

cluster relationships, we found that in most cases activity changes of the regulatory pathways

preceded the expression changes of their putative target clusters over the 2h differentiation

time course. The Notch and mTor pathways showed high connectivity to N24-induced clusters

(8 and 6 out of 12 clusters, respectively), whereas Wnt/⬚catenin signalling was specifically

correlated with clusters containing high levels of NAGs (N24i c1 and c10) and the only

constitutive cluster containing an appreciable number of NAGs (const c1) (Figure 7A).

Together, this shows that at least 24 distinct transcription modules can be defined, whose

expression is deregulated upon exit gene depletion and which are largely under the control

of five key signalling pathways gating differentiation.

Discussion

Ordered progression through pluripotency requires shutdown of the naïve TF-network and concomitant large-scale rewiring to establish the formative GRN [2, 3, 19]. Following Harald Weintraub's pioneering work with muscle differentiation and Thomas Graf's work on haematopoiesis, the ability of certain transcription factors to change cell identity without a developmental context gained central prominence with somatic cell reprogramming to pluripotency [53-55]. However, most of these studies involved forced expression of selected transcription factors to achieve synthetic cell state transitions *in vitro*. Here we utilized the remarkable properties of ESCs to recapitulate the pivotal embryonic transition from naïve to formative pluripotency. Notably, we show that this developmental cell state transition is not based on the action of one or two master transcription factors. Although crucial for defining lineage trajectories, transcription factors do not initiate cell fate decisions. Instead, the robust stability of naïve pluripotent stem cells is sustained by multiple crosstalking components and is dismantled by at least four diverse signalling inputs.

Our dissection of an authentic cell state transition indicates that the exit from naïve pluripotency is largely guided along a trajectory constrained by signalling pathways and funnelled through a handful of genetic modules. These adaptations of cellular networks establish the opportunity for TFs to initiate and consolidate cell identity. In the specific case of the exit from naïve pluripotency, expression of Oct4 and Sox2, which are key TFs for both the naïve and formative states, provide cornerstones for both the naïve and formative regulatory networks to ensure lineage fidelity during differentiation (Figure 7B). Our data support a cell fate transition model in which extinction of cell identity induced by the interplay

of signalling and the chromatin modification machinery precedes the initiation of a novel cell

fate.

Several signalling-cascades have been implicated in ESC self-renewal and naïve pluripotency

exit, including pathways controlled by LIF, Akt/mTOR, Wnt, Fgf/ERK and Notch. We derived

transcription footprints of these pathways and scrutinized the extent to which they were

deregulated by the different KOs to find that virtually all KOs showing a strong differentiation

defect also showed deregulation of at least one of these five pathways. Notably, however, we

observed segregation in terms of pathway deregulation. Thus, KOs that induce e.g. a *Tcf7l1*-

like profile were less likely to show a signature of another pathway. This suggests that ESC

differentiation requires coordinated activity changes of independent pathways under control

of separable genetic networks. The observation that single gene depletion is not sufficient to

prevent exit from the naïve state is consistent with this interpretation [21]. Our analyses

exposed discrete pathway and GRN features that mediate timely and robust mammalian cell

state transitions. We found that 12 genetic sub-networks (clusters) with non-overlapping

cellular functions emerge during the exit from naïve pluripotency. Interestingly, signalling

pathway activity was closely co-regulated with these clusters suggesting a functional linkage.

This implies that formative differentiation is driven by a limited number of parallel acting

genetic modules under control of, or controlling, key signalling pathway activities.

By examining co-regulation with the core naïve network across 146 perturbations (73 KOs in

2i and at N24) we identified the NAGs. NAGs obey strikingly similar expression dynamics *in*

vitro and *in vivo*. Moreover, in the macaque pre- to post-implantation transition and in human

naïve cell capacitation, orthologous NAGs show similar behaviour as during mESC exit from

naïve pluripotency. Therefore, this cohort of genes constitutes a layer of the pluripotency

network that is tightly linked to, and likely acts in conjunction with, the core naïve TF-network

in mammals [4]. We suggest that collective modulation of NAG expression will propel naïve cells

into formative differentiation and conversely, that collective NAG deregulation will delay

proper differentiation. The control of NAG expression appears to be interwoven with the

signalling pathways involved in controlling the naïve to formative transition. The N24 induced

clusters enriched for NAGs showed high connectivity to all tested pathways. The functional

significance of NAGs likely extends beyond self-renewal and naïve identity. Indeed, most

NAGs are not transcription factors. The growth factor Pdgfa is a case in point. *Pdgfa* is the

most strongly associated NAG, but has no known activity on ESCs, which do not express the

cognate receptor, Pdfgra. Within the ICM, Pdgfa expression is restricted to the naïve epiblast,

whereas Pdgfra is present exclusively in the primitive endoderm. This reciprocal expression

pattern is consistent with the known paracrine action of Pdgfa to promote primitive

endoderm segregation [35]. Thus, conserved linkage of NAGs to the naïve TF network may

mediate paracrine communication to other lineages within the blastocyst as well as

consolidate naïve epiblast fate.

Whereas most naïve TFs were tightly co-regulated across the 73 KOs, *Klf2* expression was

uncoupled from other members of the core network. Interestingly, while *Klf2* is highly

expressed in mouse ESCs and naïve epiblast, it is very lowly expressed in primate blastocysts,

and also in the porcine pluripotent compartment [56]. *Klf2* may therefore be a rodent specific

addition to the naïve TF-network. A further significant species difference is the very low

expression of *TCF7L1* in primate naïve cells, which underlies the differential responsiveness

of mouse and human naïve cells to GSK3 or Wnt pathway inhibition [34, 57]. *Klf2* is a genomic

target of Tcf7l1 [27]. Interestingly, *Klf2* and *Tcf7l1* are found in the same N24 induced gene-

expression cluster (cluster 10), suggesting coregulation of both components of this mouse

specific naïve pluripotency feedback loop. For most KOs the degree of failure in naïve TF

downregulation and the neutralisation of global differentiation-related transcription changes

are comparable. However, the relatively weak *Trim71* phenotype despite specific

upregulation of some naïve TF-genes, suggests that up-regulation solely of core naïve factors

must reach a certain threshold to stably maintain naïve pluripotency, consistent with negative

feedback constraints within the network [4, 5, 58].

The remarkable advantages of ESCs as an experimental venue include the utilisation of well-

defined culture conditions to recapitulate key features of *in utero* developmental progression

[1, 2, 59, 60]. Our high-resolution transcriptome analysis confirms that differentiation over 24

hours *in vitro* mimics the peri-implantation transition from naïve to formative epiblast *in*

utero. These data provide rich resources for further studies. In particular, many strong

differentiation delay KO transcriptome profiles retained transcriptome resemblance to the

pre-implantation epiblast. This relationship held true in 2i and at N24 and indicates that the

KO of these genes perturb the operative *in vivo* cell state transition machinery. In general, we

surmise that stalling the activity of the exit machinery is critical to *in vitro* capture and self-

renewal of naïve stem cells, consistent with empirical requirements for signal inhibition in

both mouse and human [7, 57]. Interestingly, deficiency for some differentiation drivers

increases similarity of 2i profiles to the *in vivo* pre-implantation epiblast. Therefore, we

propose that some of the mechanisms driving exit from naïve pluripotency contribute to the

lack of complete identity between *in vivo* E4.5 epiblast and ES cells in 2i. This is in line with

ESCs representing an engineered interruption in developmental progression and suggests

that state-of-the-art self-renewal culture conditions may not yet capture the naïve state

perfectly [1].

Our analyses provide a comprehensive inventory of genetic factors and regulatory networks

governing the major cell fate transition from pre- to post-implantation epiblast (Figure 7B).

We present evidence that mammalian development, employs similar regulatory networks

operating with similar mechanisms [61, 62]. These programmed cell state transitions in

development do not rely upon instructions delivered by one or two master transcription

factors. Rather, a cloud of activity, involving multiple co-ordinated inputs, serves to destabilise

an existing, multiply stabilised GRN. Thereby a consolidated cell state can be rapidly

dismantled by a combination of signals, triggering transition to the next developmental

waystation.

Acknowledgements

We thank Johanna Stranner, Thomas Sauer and Andy Riddell for help with flow cytometry.

Meng Li and Kosuke Yusa for sharing RC9 cells, Andreas Dahl for NGS-sequencing support and

Christa Bücker for critical comments on the manuscript. ML is funded by a WWTF-VRG grant

(VRG14-006). This study was supported by an FWF/DFG DACH grant to AB and ML (FWF grant

number: I 3786; DGF grant number 398882498). AB and MG were supported by the BMBF

(Sybacol). RS received support by the Cologne Graduate School of Ageing Research. Wellcome

and the Medical Research Council provide core support for the Wellcome-MRC Cambridge

Stem Cell Institute. AS is an MRC Professor. This project was initiated within the EU FP7

integrated project SyBoSS (Grant agreement: 242129) co-ordinated by AFS and AB including

partners ML and AS.

Author contributions:

AB, ML, AS and AFS conceptualized the study. AB, ML designed the study. AL, ML, AB, RS, MG

and FTT designed experiments. AL, MH, JR, PvdL, HFT, LS and MS carried out wet-lab

experiments. RS, MG, GGS, FTT and AL performed bioinformatic analyses. ML, AB, AS, AFS

wrote the paper with input from AL, RS, MG, MH, FTT and GG. Funding was acquired by ML,

AB, AFS and AS.

Figure 1

Supplementary Figure 1

Supplementary Figure 1

Figure 2

Supplementary Figure 2

Figure 3

a

b

Figure 4

Supplementary Figure 4

Figure 5

Supplementary Figure 5

Figure 6

Supplementary Figure 6

a

NAG enrichment
low ▭ high

const c2

NAGs, Fgf/ERK — const c1

const c9

const c6

const c7

Zscan4, Trim28 targets — const c10

LIF

Wnt *(Tcf7l1)*

Fgf/ERK *(Fgfr1 & Ptpn11)*

mTOR *(Tsc2)*

Notch *(Rbpj)*

N24i c6

N24i c10 — Klf2, >50% Tcf7l1 ChIP targets, exit genes

N24i c1 — most NAGs & naive genes

N24i c11

N24i c9

N24i c8

N24i c3

N24i c2

unconnected clusters

const c3	const c4	many exit genes
const c12	N24i c7	Etv5, Trim71 & Zfp281 KOs
const c5	N24i c5	
const c8	N24i c4	
const c11	N24i c12	

b

naïve GRN on

formative GRN on

LIF

Klf2, Klf4, Nanog, Prdm14, Oct4, Sox2, Tfcp2l1, Esrrb, Tbx3
NAGs

Wnt *(Tcf7l1)* — Jmjd1c

Apc

Notch *(Rbpj)* — Jarid2 Mbd3

Pten — mTor *(Tsc2)* — Zfp281

Raf1 — Fgf/Erk *(Fgfr1 & Ptpn11)*

Etv5

exit network modules (clusters)

Dnmt3a, Dnmt3b, Sox3, Oct4, Sox2, Fgf5, Oct6, Otx2

naïve → formative

in vitro

or

in vivo

very strong exit gene KO (e.g. Zfp281, Smg5)

strong exit gene KO (e.g. Ptpn11, Smg6)

intermediate exit gene KO (e.g. Nmt1, Cdk8)

weak exit gene KO (e.g. Jmjd1c, Leo1)

Figure 7

Supplementary Figure 7

 # Figure Legends

Figure 1

a, Illustration of the Rex1::GFPd2 reporter cell line and its' exit from naïve pluripotency. Rex1-

GFPd2 (in short Rex1-GFP) expression is tightly linked to naïve pluripotency. Shutdown of GFP

expression indicates commitment to differentiation.

b, FACS analysis of Rex1-GFPd2 reporter levels throughout a 72h differentiation time-course

after 2i withdrawal.

c, Scheme of the screening strategy to identify candidate genes involved in exit from naïve

pluripotency. After random insertional mutagenesis using piggyBac transposon gene-traps,

haploid Rex1::GFPd2 ESCs were released into differentiation. Cells that maintain GFP

expression after exposure to differentiation conditions were isolated and the gene-trap

insertion sites mapped.

d, Plot showing cumulative number of hits and the cumulative number of novel hits over 35

independent insertional mutagenesis screens in haploid ESCs.

e, Representative Rex1-GFP FACS plots showing the differentiation delays 30h after 2i

withdrawal of *Tcf7l1*, *Trim71* and *Mbd3* KOs. A Myc KO served as a negative control. Blue

indicates the KO FACS profiles and dashed lines indicates WT Rex-GFP.

f, Differential expression of genes at N24 vs. 2i in WT RC9 cells. Black dots show significance

(FDR ≤ 0.05, H0: |FC| > 1.5). Pluripotency genes are red dots, haploid screen hits are blue

dots, the 73 KO-genes are green dots.

g, t-SNE projection of the 73 KOs in 2i, based on expression of 3058 differentiation-associated

genes (RC9^{N24} vs. RC9^{2i}, FDR ≤ 0.05, H0: |FC| > 1.5). Differentiation defects observed at N24

in the respective KOs are indicated by a colour gradient, measured as average naive marker

log2fold change (based on expression levels of *Esrrb, Nanog, Tfcp2l1, Tbx3, Prdm14* and *Klf4*)

in the respective KO at N24. Red: delayed differentiation; blue: accelerated differentiation -

see key in **h,**.

h, Similar to **g,** for KOs at N24

Supplementary Figure 1

a, The top 15 candidate genes from the haploid screen ordered by the number of times the

hit was found in the 35 screens, including the number of independent integration sites and

the calculated FDR.

b, Significantly enriched pathways among screen hits ranked according to fold enrichment;

colours represent FDR in screen analysis. Numbers next to bars indicate number of hit-genes

within category,

c, Significantly enriched GO terms among screen hits as in **b,**

d, Workflow to generate Cas9 knockouts in RC9 ESCs.

e, Idealized strategy illustrating gRNAs and genotyping primers.

f, Western analysis using indicated KOs and indicated antibodies. Tubulin or Gapdh were used

as loading controls, as indicated.

g, Anti-flag specific Westerns in indicated KO^rescue ESCs upon stable forced expression of 3xflag

rescue cDNAs driven from a CAG promoter.

h, Rex1GFP levels measured by FACS showing restoration of differentiation behaviour in the

indicated rescue cell lines at N30.

i, t-SNE visualization of RNA-seq profiles in 2i and at N24 before and after batch correction.

j, GO enrichment analysis of the top 10% of genes (by absolute logFC) that were differentially

expressed between 2i and N24 in WT ESCs (FDR ≤ 0.05, H0: |FC| < 1.5).

k, Exit from pluripotency factors showing significant expression changes between 2i and N24

(lfc≥0.8; adj. p-value≤0.05). Log2FC, adj. p value and quantification of the molecular

phenotype using the average upregulation of a set of naïve marker genes (*Esrrb, Nanog,*

Tfcp2l1, Tbx3, Prdm14, Klf4) are shown.

l, Dot-plot showing regression analysis of KO-induced changes at N24, comparing *Tcf7l1* KO

to Apc and Csnk1a1 KOs. Total least squares (TLS) analysis results are indicated.

m, Rex1-GFP levels measured by FACS at N24 showing the effect of Epiblastin A addition

during differentiation.

n, Rex1GFP analysis at N24 of *Pum1* KO cells, showing a relatively strong differentiation defect

at early passages.

Figure 2

a, Naïve (top) and formative (bottom) marker gene expression changes at N24 compared to

WT in all 73 KO ESCs. Clustering based on naïve marker gene expression shows a wide

distribution of differentiation delay phenotypes.

b, Rex1GFP FACS analysis of *Jarid2* and *Tcf7l1* KO ESCs after transfection with negative control

or *Klf2* specific siRNAs.

c, For each of the 73 KOs, the change of mean naïve marker gene expression at N24 (KO vs

WT) was plotted against the change of the global transcriptome at N24 state (correlation of

KO-induced change at N24 with differentiation-induced changes in WT differentiation, N24 vs

2i).

d, Comparison of number of genes deregulated in 2i and at N24 in all KOs (FDR ≤ 0.05, H0:

|FC| < 1.5). Differentiation phenotypes are colour coded according to mean naïve marker

gene expression change in KOs at N24.

e, Rex1GFP FACS analysis of WT, *Jmjd1c* and *Tcf7l1* KO cells at N48 cultured with and without

the Gsk3 inhibitor, CH.

Supplementary Figure 2

a, RNA-seq based fold changes (FC, relative to WT) of indicated naïve marker genes in

indicated KOs in 2i (only FCs with adj. p ≤ 0.05 are shown).

b, RNA-seq derived data showing FCs of indicated formative marker genes in indicated KOs in

2i (only FCs with adj. p ≤ 0.05 are shown).

c, Rex1-GFP analysis of WT cells at N24, transfected with negative control or *Klf2* specific

siRNAs. *Klf2* depletion increases the speed of Rex1GFP downregulation.

d, Comparison of the number of differentially expressed genes at N24 (FDR ≤ 0.05, H0: |FC|

< 1.5) to the mean naïve marker gene expression at N24 (phenotype strength). Stronger

differentiation phenotypes correlate with more differentially expressed genes.

e, Expression log2 fold changes (log2FC) at N24 between knockout and RC9 control,

comparing *Tcf7l1* KOs to *Jmjd1c* KOs. Each dot corresponds to one gene. Only genes showing

significance (FDR ≤ 0.05, H0: |FC| < 1.5) in either one of the KOs are plotted. Red line: total

least square regression; regression coefficients are shown. Convergence on similar target

genes is shown by largely identical directionality of regulation of significantly regulated genes

in both KOs.

Figure 3

a, Scheme illustrating comparisons of the 73 KO gene expression profiles at N24 with a 2h-

resolution *in vitro* differentiation time course of WT RC9 ESCs.

b, Heatmap showing differentiation delays of the 73 KO lines quantified by average naïve

marker gene expression. Red bars indicate closest correlation to that specific time-point.

Positive values indicate delayed differentiation. Negative values more rapid differentiation.

Each line corresponds to one KO. Selected KOs are indicated. See Supplementary Table 2 for

the full hierarchy of the 73 KOs.

c, Plot comparing the differentiation delays calculated using naïve markers as in **b,** or using all

3058 genes changed in WT differentiation from 2i to N24.

Supplementary Figure 3

a, Principle component analysis (PCA) of the 2h resolved WT differentiation time course

sampled by RNA-seq. 0h corresponds to cells in 2i.

b, Heatmap showing differentiation delays of the 73 KO lines quantified using expression of

all 3058 genes differentially expressed in WT between 2i and N24. Red bars indicate closest

correlation to that specific time-point. Positive values indicate delayed differentiation.

Negative values more rapid differentiation compared to WT. Each line corresponds to one KO.

Selected KOs are indicated.

Figure 4

a, Scheme illustrating the comparison of gene expression profiles of 73 KOs in 2i and at N24

with *in vivo* derived averaged single cell expression data comprising peri-implantation epiblast

tissues. Specifically, transcription profiles of each KO in 2i and at N24 were compared with in

expression datasets obtained from the E4.5, E5.5 and E6.5 epiblast.

b, Fraction of identities (FOI) between 2i and N24 expression data and mouse E4.5, E5.5 and

E6.5 epiblast, computed using all expressed genes (log2FPKM > 0) for all KOs. Each dot

represents one KO. Average naïve marker expression in KOs at N24 is indicated by a colour

gradient.

c, Plot showing a comparison of FOIs of WT (RC9) and KO cells in 2i (green) and at N24 (grey)

to E4.5 and E5.5 mouse epiblast, computed using all expressed genes (log2FPKM > 0).

(d,e) Data shown in Figure 4c separated by culture condition. Comparison of FOIs of WT and

KO cells in 2i **d,** and at N24 **e,** to E4.5 and E5.5 mouse epiblast; average naïve marker

expression in KOs at N24 is indicated by a colour gradient.

f, PCA showing all KOs in 2i (triangles) and at N24 (diamonds) and embryo derived datasets

(ICM, PrE, E4.5Epi, E5.5Epi, E6.5Epi). As in **d,,** blue to red colour gradient indicates naïve

marker expression at N24 of the respective KO.

g, Similar to Figure 4b, showing FOIs between 2i and N24 datasets and macaque pre- and

post-implantation embryos.

h, Similar to Figure 4b, showing FOIs between 2i and N24 datasets and hESC capacitation

datasets comprising hESCs in naïve conditions (day0) and hESCs 20 days after release into

primed conditions (day20).

Supplementary Figure 4

a, Zoom in of the plot in Figure 4c, focusing on the overlap of several KOs in 2i (green) and in

N24 (grey). N24 samples of *Etv5, Smg7, Tsc2, Smg5* and *Zfp281* KOs show FOIs to the E4.5 EPI

similar to KOs cultured in 2i.

b, Comparison of FOIs to the E4.5 epiblast of all KOs in 2i compared to FOI of all KOs at N24;

average naïve marker expression in KOs at N24 is indicated as colour gradient. With some

exceptions, stronger *in vitro* phenotypes translate to increased similarity to the E4.5 epiblast

in 2i and at N24.

c, FOIs of KOs in 2i to ICM, E4.5 trophectoderm (TE), E4.5 primitive endoderm (PrE), E4.5

epiblast (Epi), E5.5 Epi and E 6.5 Epi. Analysis extended but similar to Figure 4b.

d, Plot comparing FOIs of 2i samples to E4.5 epiblast (x-axis) to FOIs of N24 samples to E5.5

epiblast(y-axis). Dotted grey lines indicate 95% confidence intervals. Colour gradient shows

similarity of 2i expression profiles to PrE.

e, Correlation plot of FOIs between 2i profiles and mouse naïve (E4.5) epi and FOIs between

2i profiles and macaque naive epi (correlation significance is indicated).

f, Correlation plot of FOIs between 2i profiles and mouse naive epi and FOIs between 2i

profiles and human naïve ESCs (day0) (correlation significance is indicated).

g, Correlation plot of FOIs between N24 profiles and mouse E5.5 epi and FOIs between N24

profiles and macaque post-implantation epi (correlation significance is indicated).

h, Correlation plot of FOIs between N24 profiles and mouse E5.5 epi and FOIs between N24

profiles and human ESCs at day 20 after release into primed culture conditions (correlation

significance is indicated).

Figure 5

a, x-axis displays all 3rd dimension (Dim3) genes, which separates mouse naïve from post-

implantation epiblast (the inset shows the full PCA plot, see also Supplementary Figure 5d) in

order of extent of contribution to Dim3. The contribution to Dim3 of NAGs (orange) and the

top 250 differentially regulated genes (green) are plotted in a cumulative manner.

b, Plots comparing relative log2FC in mouse ESC differentiation (2i vs N24) to relative log2FC

in mouse *in vivo* transition from pre- to post-implantation epiblast (E4.5 vs E5.5). Selected

gene groups are plotted: i) NAGs, ii) all 3058 differentially expressed genes in mESC

differentiation and iii) the top 250 up and downregulated genes in mESC differentiation. Rho

values indicate the level of correlation between *in vivo* and *in vitro* differentiation of given

gene groups. Alpha indicates the angle between the x-axes and the orthogonal regression.

Genes with log2FC > 0.5 and < -0.5 in both the x- and y-axes are highlighted in orange and

blue respectively.

c, As for **b,** except plots comparing log2FCs in mouse ESC differentiation (2i vs N24) to

macaque *in vivo* transition from pre- to post-implantation epiblast cells.

d, As for **b,** except plots compare log2FCs in mouse ESC differentiation (2i vs N24) to human

naïve ESC capacitation (d0 to d20).

e, Graphs showing the frequency of co-regulation of NAGs or the top 250 differentially

expressed genes (2i vs N24) with differentially expressed genes in mouse and macaque *in vivo*

pre- to post-implantation epiblast differentiation, and human naïve ESC capacitation. Two-

tailed chi-square test (CI 95%) was used to compute significance levels between the expected

and the observed number of modulated genes.

f, One-way hierarchical clustering showing expression changes of NAGs (left) and top

differentially expressed genes (right) in the indicated differentiation models.

Supplementary Figure 5

a, Expression change of NAGs during naïve to formative differentiation in WT ESCs (y-axis)

versus the strength of link to the naïve network (naïve association, x-axis). The orange line

indicates the linear regression. The R^2 value is indicated in the plot.

b, *Klf2* expression at the indicated stages. Compacted morula (cMOR), expanded ICM (eICM),

E4.5 epiblast (EPI) derived from single cell RNA-seq datasets from human, marmoset and

mouse preimplantation development.

c, Plots derived from the GRAPPA visualisation-app showing expression in yellow of *Pdgfa* and

its cognate receptor *Pdgfra* during mouse preimplantation embryo development.

d, Principal component analysis of published single cell RNA-seq datasets showing that

Dimension 3 (Dim 3) separates pre- from post-implantation epiblast *in vivo* (as shown in Figure

5a).

Figure 6

a, Relationship between the KO expression profiles at N24 (or in 2i for the LIF pathway) to the

expression footprints of the five signalling pathways as indicated (see also Supplementary

Figure 6a). Relative pathway activity is -1 for Fgf/ERK (defined by Fgfr1 & Ptpn11 KOs), Wnt

(Tcf7l1 KO) and Notch (Rbpj KO) and +1 for mTOR (Tsc2) and LIF (defined by presence of LIF

in 2i medium) for effects equally strong as in the representative KO. Pathway activity is

indicated by colour code (green for activity, red for inactivity). All tiles with an absolute

pathway activity smaller 0.3 are grey. The size of the tiles corresponds to the spearman

correlation of pathway footprints between each KO and the corresponding representative

KOs. The colour gradient on the sidebar at the left indicates mean change of naïve marker

gene expression (as a measure of phenotypic strength) for the corresponding KO at N24.

b, Heatmap showing average log2FCs (KO^{N24} vs. RC9^{N24}) of constitutive response gene

clusters. Asterisks indicate that the absolute row-wise z-value of the respective mean

expression value was above 1.96 ($p \leq 0.05$), indicating extreme values within the context of

that row. For reference, average naive marker log2FCs at N24 vs WT are shown in a separate

row below (similar to Figure 2a).

c, GO enrichment analysis per 'constitutive KO response' cluster. Dot-size scales with log2

enrichment. Adj. p vales are clour coded in shades of red.

d, As for **b,** except heatmap showing average log2FCs (KO^{N24} vs. $RC9^{N24}$) of N24 induced KO

response gene-clusters.

e, As for **c,** except GO enrichment analysis per 'N24 induced KO response' cluster.

Supplementary Figure 6

a, Schematised computational strategy to derive non-overlapping gene-expression footprints

for the five signalling pathways.

b, Average change in naive marker expression induced by KOs at N24 (y-axis) and their 'LIF-

likeness' in 2i (x-axis; correlation of RC9 LIF response with knockout induced response in 2i).

Overall correlation across these values is r=0.69. The KOs showing the strongest correlations

(*Tsc2*, *Ptpn11* and *Trim71* KOs) are indicated in red.

c, Flow analysis showing delayed Rex1-GFP downregulation in cells starting from cultures in

2i/LIF compared to cells starting from 2i.

d, Multiple regression coefficients of pathway activity profiles across all KOs when predicting

the differentiation delay phenotype (defined by mean naïve marker gene expression change

at N24). Error bars indicate the standard error of the coefficient.

e, Euler diagram showing the overlap of defined gene groups. Panels on the right show

exemplary gene expression behaviours. The constitutive and the N24 induced KO response

group of genes were used for further analysis.

f, Percentages of genes in the N24 induced KO response (N24 induced) or constitutive KO

response (constitutive) clusters that significantly change during WT differentiation (FDR ≤

0.05, H0: |FC| < 1.5).

Figure 7

a, Visualization of connections between pathways (central ellipses) and gene clusters

(rectangles) based on multiple regression models. The colour gradient of the clusters indicates

enrichment for NAGs. Dashed connections link pathways to clusters that fit time-course

validated regression models (see Supplementary Figure 7A and C). Continuous connections

are further supported by ChIP or pathway defining gene enrichments (see Supplementary

Figure 7E,F). Connections are coloured by source pathway. Unconnected clusters are listed at

the right.

b, Summary model of the genetic and signaling pathway control of naïve to formative

transition. Between the naïve and formative gene regulatory networks, the five signaling

pathways largely operate through partially overlapping separable gene clusters.

Supplementary Figure 7

a, Heatmap illustrating multiple regression models of pathway activities predicting cluster

expression. Only significant (adj. p. ≤ 0.01) interactions are shown. Colour intensity represents

the strength of the interaction. Red tiles indicate that higher activity of a certain pathway

results in higher expression of the cluster, while blue tiles indicate the opposite.

b, Pathway activities (average expression of the 50 specific pathway-defining genes) across a

WT differentiation time-course for all 5 pathways. y-axis shows average log2FC of pathway

defining genes relative to the mean expression throughout the time-course.

c, Observed cluster expression change (red) and cluster expression change predicted by

regression models (blue) throughout the 32h WT differentiation time course for all induced

clusters. Predicted cluster expression is calculated using pathway activities over the WT time

course and significant (adj. p ≤ 0.01) interactions from the multiple regression models. Graphs

show log2FC of cluster expression relative to the mean cluster expression across the time

course. Red cluster names indicate that cluster-pathway connections were not predicted in

Supplementary Figure 7A or validation through 2h time-course expression failed (predicted

expression did not precede observed expression).

d, Similar to Supplementary Figure 7C for constitutive clusters.

e, Enrichment from Fisher test of ChIP-targets, NAGs and pathway defining gene sets in

induced clusters. Enrichment is indicated by red circles, depletion by blue squares. Size

indicates strength of over- or underrepresentation and colour gradient encodes adj. p values.

f, Similar to Supplementary Figure 7E for constitutive clusters.

Methods

Cloning

Guide RNAs (gRNAs) were designed using the computational pipeline described below, or by

http://crisprscan.org. To clone the gRNAs for ESC KO of the 154 selected genes, we used RecET

recombineering to insert the guide RNA protospacer encoding sequence between the U6

promoter and the universal sgRNA scaffold in pBR322-U6-cm-ccdB-sgRNA-amp, as described

previously (Baker et al, 2016). We generated 331 sgRNA expressing plasmids for 149 genes (2

sgRNAs for 131 genes, 3 sgRNAs for 5 genes, 4 sgRNAs for 12 gens and 6 sgRNAs for 1 gene).

In brief, the plasmid was linearized at the point of insertion by BstZ17I digestion, purified

using the Invitrogen Charge Switch PCR Purification Kit and dissolved in water; 80-mer

oligonucleotides containing the protospacer sequence, flanked by 30bp of homology to the

target plasmid were dissolved in water. *E. coli* GB05 transformed with the pSC101-Prha-ETgA-

tet plasmid [63], was cultured to OD600 ~0.5 in deep-well 96 well plates and then induced for 1

hour for expression of the ETgA operon with L-rhamnose. Electroporation with 200 ng (0.1

pmol) of the linearised vector and 50 pmol of oligonucleotide, using BTX 96 well

electroporation plates (MOS96, 2mm gap,) and the BTX ECM630 electroporator with HT-200

adapter, was described previously [64]. After one hour of recovery time, 100 microliters (1/10th)

of the culture was transferred to a new plate with 900 microliters of LB media plus 100

micrograms/ml ampicillin. After overnight selection the saturated cultures were plated for

single colonies on L-agar + 100 micrograms/ml ampicillin. A single colony for each construct

was picked and the clones were grown under selection in 96 well plates, plasmid DNA was

isolated, and Sanger sequenced from both directions using the M13F and M13R primers

(flanking the U6-protospacer-sgRNA scaffold cassette, Supplementary Table 6). Sequence

analysis confirmed the correct engineering of 316 out 331 plasmids. After sequencing of a

second colony or repeat cloning for the remaining 15 clones, all the desired constructs were

obtained. Alternatively, annealed oligonucleotides (Supplementary Table 6) were cloned into

a BsaI site of a gRNA expression vector (Addgene Plasmid number 41824) [65] and correct

insertion was determined by Sanger sequencing with the SP6 primer (Supplementary Table

6). For generating rescue cell lines, indicated coding sequences were cloned into a pCAG-

3xFLAG-empty-pgk-hph vector [14] after PCR amplification and correct insertion was detected

by restriction digest and Sanger sequencing with the 3xFlag_seq primer (Supplementary Table

6).

Cell culture

Diploid (biparental) ES cells were routinely cultured in gelatin coated 25cm^2 flasks in DMEM

supplemented with 15% FCS (batch tested, Biowest), 1x Penicillin-Streptomycin (Sigma),

0.1mM NEEA, 1mM sodium pyruvate, 1mM L-glutamine, 0.05mM 2-mercatpoethanol

(Gibco), 10ng/mL LIF (batch tested, in-house), 1.5µM CHIRON and 0.5µM PD0325901 (termed

ESDMEM-2i) on gelatin coated plates. The basal medium during all differentiation assays and

for haploid ES cell culture (N2B27) consisted of a 1:1 mixture of DMEM/F12 and Neurobasal

medium supplemented with 1x B27 (Gibco), 0.5x N2 (homemade), 0.1mM NEAA, 1mM L-

glutamine, 1x Penicillin-Streptomycin and 0.05mM 2-mercaptoethanol. Haploid ES cells were

routinely cultured in N2B27 supplemented with 3µM CHIR99201, 1µM PD0325901 and

10ng/ml LIF and regularly sorted based on FSC/SSC parameters by FACS [66]. Haploid

Rex1GFPd2-IRES-BSD (hRex1GFPd2, [67] and haploid Rex1::mKO2-IRES-BSD/pOct4-GFP-IRES-

Puro ESCs (generated in this study from a double reporter mouse) were derived from

activated oocytes in 2i/LIF medium and used for screens. Biparental diploid ES cells with a GFP

driven by the endogenous Rex1 promoter and EF1a driven Cas9 targeted to the Rosa26 locus

(RC9 cells, [13] served as parental cell line for the KO lines generated in this study. A list of KO

cell lines can be found in Supplementary Table 6.

Transposon-based saturation screen for key players of the exit from pluripotency

The genetrap vectors 5'-PTK-3', TNN, TNP (Horie et al. 2011, Allan Bradley Lab) together with

haploid Rex1::GFPd2 ES cells (hRex1GFPd2, Leeb et al. 2014) and haploid Rex1::mKO2-IRES-

BSD/pOct4-GFP-IRES-Puro ES cells (hOR, this study) were used in 35 independent experiments

to drive the screen to saturation. Mutant pools were generated by electroporation of 10^7

haploid ESCs using a GenePulser Xcell (270 V, 500 µF, ∞ Ω, Biorad) with 0.5 µg genetrap

plasmid and 10 µg hyperactive transposase (hyPBase) [68]. 24h after electroporation, selection

was started for 4 days using 1µg/ml Puromycin. Thereafter cells were plated at a density of

10^4 cells/cm^2 in N2B27 medium in the absence of LIF or inhibitors to allow differentiation.

After 7–10 days in differentiation conditions, GFP-positive cells were sorted and replated at a

density of 10^4 cells/cm^2. After culture in N2B27 medium for a further 7–10 days, GFP-positive

ESCs were resorted, expanded for 48h in 2i/LIF medium and DNA was isolated using the

PureGene kit (Qiagen) according to the manufacturer instructions. Sequencing libraries were

prepared using an optimized Splinkerette PCR protocol (Leeb et al. 2014) with a modified and

improved set of adapters and primers (see Supplementary Table 6). Adapters were annealed

at 50µM each in T4 DNA ligase buffer by incubating at 97.5°C for 150 sec followed by a

temperature decrease of 0.1°C/5sec for 775 cycles. Ready-to-use adapters were stored at -

20°C. Genomic DNA was isolated using the PureGene kit (Qiagen) according to the

manufacturers recommendations and quantified on the Qubit 2 fluorometer (Life

Technologies) using the Qubit dsDNA Broad Range Assay (Life Technologies, Q32853). 2µg of

genomic DNA was diluted to a total volume of 120µl with Low TE buffer (10mM Tris-HCl, pH

8.0, 0.1mM EDTA) and sheared to 250bp on a Covaris S-series Sample Preparation System

with the following settings: Duty Cycle – 20%, Intensity – 5, Cycles per Burst – 200, Time – 60

seconds, Temperature – 4° to 7°C. After shearing DNA was isolated with the QIAquick PCR

purification kit (Qiagen) and quality assessed using the Agilent Bioanalyzer with the DNA high

sensitivity kit (Agilent). End-repair was performed using the NEBNext End Repair Module

(E6050S) by mixing 100µl sheared DNAl, 11.6µl buffer and 6µl enzyme followed by incubation

at 20°C for 60min. After cleanup with the PureLink PCR Purification Kit (K3100-01), 42µl of the

end repaired DNA was A-tailed with the NEBNext dA-Tailing Module (E6053S) with 5µl buffer

and 3µl enzyme at 37°C for 60 minutes. Following PureLink purification of to the A-tailed DNA,

the annealed adapters were ligated using theNEBNext Quick Ligation (E6056S) kit at 20°C for

30min. Products were cleaned up using Ampure XP beads at a 1:1 ratio. Libraries were PCR

amplified using Kapa HIFI (Roche) with 10µM PB5_pr1 and 10µM SplAP1 during 18 cycles of

20sec 98°C, 20sec 63°C and 40sec 72°C with an initial denaturation at 95°C for 2minutes and

a final extension at 72°C for 5min. Samples were cleaned up using Ampure XP beads at a ratio

of 0.8. A second round of PCR was conducted using xxul from PCR1 using the Kapa HIFI

polymerase with 10µM SplAP2 and 10µM index primer for 13 cycles at an annealing

temperature of 60°C with all other settings as in PCR1. Final libraries were quantified using

the KAPA SYBR Fast qPCR Mix with primers SYB7 and SYB5 at 10µM each and a final quality

control was conducted with the Agilent Bioanalyzer DNA High Sensitivity Assay.I ntegration

sites were mapped as in [12] and described below.

Generation of KO ESCs

Direct comparison between KOs necessitates a similar culture and passaging history. We

therefore established an experimental pipeline to parallelise the generation of multiple KO

ESC lines per week. 2x10^5 RC9 cells were transfected in a 6-well format with a set of two gRNA

containing vectors (Supplementary Table 6; (1μg each), together with 0.5μg of pCAG-dsRed

by Lipofectamine2000 (3μl per reaction). 12-16h after transfection, medium was replaced. To

enrich for transfectants, cultures were sorted for GFP/DsRed double positive cells on a BD

FACS Aria III 48h after transfection and plated at clonal density in 6cm dishes (100 to 800 cells

per dish) in ESDMEM2i. Approximately one week after sorting, 48 colonies per gene KO were

picked into 96-well plates, trypsinized and split into one expansion-plate and one plate for

PCR genotyping   after boiling lysis and proteinase K treatment (see PCR genotyping).

Identified KO clones were expanded to 12-well or 6-well plates after 3 days of growth and the

remaining plate was frozen in 50%FCS/10% DMSO as a backup. Clones were frozen from 12-

or 6-well plates in freezing medium containing 50% FCS and 10% DMSO. Of the attempted

154 gene disruptions, we achieved 115 homozygous knockout cell lines. In 16 cases, only

heterozygous clones were obtained (indicating essentiality of these genes), and in 23 cases

we were unable to recover gene disruptions, despite multiple attempts (Supplementary Table

6).

PCR genotyping

Crude DNA lysates were generated by pelleting half of a picked colony in a PCR plate at 500g

at 4°C, followed by two washes with PBS. 25μl of water were added and pellets were boiled

for 5min at 95°C. After cooling, Proteinase K was added at a final concentration of 3μg/μl and

incubated at 65°C for 1h followed by inactivation at 95°C for 10min. Successful KO generation

was confirmed by PCR, employing a three-primer strategy. Specific reverse primers for the

deletion event and a possible wildtype allele were used in combination with a common

forward primer (Supplementary Table 6). Cell lines with indels caused by a NHEJ event around

the exonic gRNA1 target site, were detected by Sanger sequencing of the "wildtype"-band.

Verification of the KO-events was performed by the same PCR strategy on DNA, isolated using

the PureGene DNA isolation kit (Qiagen) according to the manufacturer recommendations.

For genotyping PCR, we used OneTaq (NEB) or JumpStart RedTaq (Sigma) PCR Master Mix

according to the manufacturers recommendations for cycling and a standard annealing

temperature of 55°C and 35-40 cycles.

Immunoblotting

Whole cell fractions were isolated using RIPA buffer and protein concentrations were

determined using a Bradford Assay (Bio-Rad). 20μg whole cell lysate were separated on 8-

12% SDS-PAGE gels (depending on the molecular weight of the target proteins), and

subsequently blotted on 0.2μm nitrocellulose membranes (Amersham). Membranes were

blocked in TBS or PBS containing 5% milk and 0.1% Tween-20 or 3% BSA and 0.1% Tween 20

for 1h. Primary antibodies were incubated overnight at 4°C, at a dilution of 1/1000 with the

following exceptions: Anti-Csnk1a1 1/200, Anti-Flag M2 1/1000 Anti-Gapdh 1/25000, Anti-

Tubulin 1/5000, Anti-Smg5 1/200, Smg7 1/2000. Secondary goat anti-mouse IgG HRP

(1:10000) or goat anti-rabbit IgG HRP (1:15000) were incubated at RT for 1h in blocking

solution. Antibody binding was detected using the ECL select detection kit (Amersham). A list

of antibodies can be found in Supplementary Table 6.

1003

Parallel Differentiation

ES Cells were differentiated in seven batches, always including WT controls, control KO ESCs

(no phenotype expected) and a range of ESCs with expected weak to strong phenotypes.

Where applicable two independent KO clones were used, other KO ESCs were cultured in

parallel replicates before intiation of differentiation. 24h before starting the differentiation

assay, medium was changed from ESDMEM2i to N2B27 based 2i/LIF medium. Cells were then

trypsinized, counted and plated in 2i (without LIF) at a density of 10^4 cells/cm^2 in two replicate

wells of a 6-well plate for differentiation, one 6-well as undifferentiated control and 2 wells

of a 12-well plate (undifferentiated control and differentiated sample) to monitor

differentiation by FACS. 12h after plating, cells were washed twice with PBS and medium was

changed to unsupplemented N2B27. For the undifferentiated controls medium was replaced

with fresh 2i. 24h later, at N24, cultures were harvested in RLT buffer and stored at -80°C

before isolation of RNA using the RNeasy Mini Kit (Qiagen). mRNA was isolated from 1 ug total

RNA by poly-dT enrichment using the NEBNext Poly**a,** mRNA Magnetic Isolation Module

according to the manufacturer's instructions. Samples were eluted in 15ul 2x first strand

cDNA synthesis buffer (NEBnext, NEB). After chemical fragmentation by incubating for 15 min

at 94°C the sample was directly subjected to the workflow for strand specific RNA-Seq library

preparation (Ultra Directional RNA Library Prep, NEB). For ligation custom adapters were

(Supplementary Table 6). After ligation, adapters were depleted by an XP bead purification

(Beckman Coulter) adding beads in a ratio of 1:1, followed by an index PCR (15 cycles) using

Illumina compatible index primer. After double XP beads purifications (with beads added in a

ratio 1:1) libraries were quantified on a Fragment Analyzer run with a NGS Assay Kit (Agilent)

and loaded on a HiSeq 3000 flowcell with 50 cycles single end sequencing by pooling the

samples based on molarity aiming for 30 mio. reads per sample.

1028

2h resolved differentiation timecourse

Medium was changed to N2B27-2iLIF 24h before the assay. Cells were trypsinized, counted

and plated at a density of 10^4 cells/cm^2 in 6-well plates in 2i (without LIF). 12h later medium

was changed to unsupplemented N2B27 (or 2i for the undifferentiated controls) and cells

were harvested in buffer RLT every 2 hours for the next 32h (N32). RNA was isolated using

the RNeasy Mini Kit (Qiagen). All libraries were generated from 500ng RNA using the TruSeq

Stranded mRNA Library Prep Kit (Illumina) and analysed on a HiSeq4000, Paired End 75bp.

1036

LIF signature

$1x10^4$cells/cm^2 were plated in N2B27-2i and N2B27-2iLIF and grown for 24h after which RNA

was isolated using the RNeasy Mini Kit (Qiagen). Libraries were made from 1µg high quality

RNA using the Quantseq 3' mRNA-Seq Library Prep Kit FWD for Illumina (Lexogen) according

to the manufacturer's instructions. Libraries were quantified using qPCR according to the

Quantseq Library Prep Kit and with the included primers, pooled and quality checked using

an Agilent Bioanalyzer DNA High Sensitivity Assay (Agilent). Multiplexes were sequenced on

a HiSeq4000, Single End 50bp.

1045

RNAi experiments

For RNAi FlexiTube siRNAs against Klf2 were used with AllStars Negative Control siRNAs

(Qiagen). 20ng siRNAs/4x10^4 cells were transfected in 2i using DharmaFect (Dharmacon). 12h

later medium was changed to N2B27 after two PBS washes. At N24 and N32 Rex1GFPd2

expression was determined by flowcytometry (see Flowcytometry Analysis) and cells were

harvested in RLT buffer and isolated with the RNeasy mini kit (Qiagen) according to the

manufacturer's recommendations. cDNA was transcribed using the SensiFAST cDNA Synthesis

Kit (Bioline). Naïve and primed marker expression was determined by qPCR using the Sensifast

SYBR No Rox-Kit (Bioline). Primers are listed in Supplementary Table 6.

1055

Flow cytometry

Cells were dissociated in 0.25% trypsin/EDTA and trypsin was neutralized with DMEM

supplemented with 15% FCS. After passing through a 40μm mesh, Rex1-reporter activity was

measured using a BD Fortessa machine. High-throughput-measurements were performed in

96-well plates using the HTS unit of the BD Fortessa. Data were analyzed using FlowJo

software.

Data Analysis

Transposon Mutagenesis Screen in Haploid Murine Embryonic Stem Cells

Integration mapping was performed as previously described [12]. Most gene-trap insertions

were found in intronic sequences, consistent with the relative length distributions of genomic

features. However, after normalising to available TTAA sites per region, TTAA sites in 5'UTRs

and promoters (<-500bp) were overrepresented, whereas TTAA sites in introns were

relatively depleted. 7760 independent integrations passing cut-off criteria were mapped to

3469 genes. Of those, 232 Genes (2474 independent integration sites) were hit in five or more

independent screens, indicating that those are integrations causative for the detected

differentiation defect. To complement the candidate list and correct for gene specific biases

we further developed a strategy to assign statistical strength to candidate genes by assessing

the number of integrations per gene in relation to available TTAA transposon integration sites

We utilized the number of different TTAA-sites within each gene to calculate the probability

of having integrations in k or more different locations by chance. All TTAA sites in mm10 were

assessed using bowtie and the analysis was restricted to genes classified as protein coding,

long non-coding RNAs and micro RNAs. We computed the probability that a gene is hit k times

given that it contains n TTAA transposon integration sites assuming a binomial distribution

(while k is the observed number of integrations).

1080
$$P[X \geq i] = \sum_{i=k}^{n} P[X = i] = \sum_{i=k}^{n} B(i|p_0, n) = \sum_{i=k}^{n} \binom{n}{i} p_0^i (1 - p_0)^{n-i}$$

p_0 was then calculated by $\frac{All\ hit\ TTAA-sites\ in\ genes}{All\ TTAA-sites\ in\ genes}$

We applied this test to genes with at least 26 TTAA sites, as genes with fewer TTAA sites would

be significant (p-value < 0.05) with a single integration. Candidate gene lists between both

analyses showed overlapping results and top hits were largely identical. This resulted in a list

of 421 significant genes (adjusted p-value < 0.01 and hit in at least two independent screens).

Guide RNA design criteria

To automate the design of sgRNAs as far as possible we used the R package CRISPRseek (Zhu

et al., 2014) to score the efficiency and likely off-target binding of candidate sgRNA sequences.

For a selected gene the algorithm proceeds as follows:

1090

1. Retrieve the sequence of the first protein coding exon

2. Sequence has to be at least 50 bases long

3. Find sgRNAs within the sequence and score them.

4. If no appropriate sgRNA could be found, move to next exon

5. Still in the first third of the coding sequence?

6. Find a second sgRNA in an intron at least 5kb away, but no more than 30kb

7. Score the sgRNAs

A minimal efficiency of 0.2 and an off-target score < 100 was required.

PCR Primer Design Strategy for Simplified Knock-out Validation

The surrounding regions of the sgRNAs (min distance 20bp) were used as input for PRIMER3

primer predictions. Primers were designed to have a length of 18-27 bases with an optimum

of 20 and to have melting temperatures between 57°C and 63°C with an optimum of 60°C. All

primers were checked for off-target binding sites by using Blastn. Blastn was called with the

parameters:

-task megablast -use_index false -word_size 7 -perc_identity 65 –evalue 30000 -

max_target_seqs 50 -num_threads 4 –outfmt 7.

RNA quantification, RNAseq

Quality control was performed using fastQC (version 0.11.5). Transcripts were mapped to the

mm10 mouse reference genome and counts determined using STAR (version 2.5.3). RPKM

values were calculated using DeSEQ2 (version 1.20.0).

RNA quantification, QuantSeq

Transcripts from RC9 cells in 2i and 2i/LIF medium were measured using QuantSeq (Lexogen)

and quantified as described. Reads were trimmed with bbduk (version 35.92). Quality control

was performed using fastQC (version 0.11.5). Transcripts were mapped to the mm10 mouse

reference genome using STAR (version 2.5.3). After indexing with samtools (version 1.3) reads

in genes were counted using HTSeq-count (version 0.6.0). Counts were subjected to

differential expression analysis.

Differential expression analysis

Differential expression analysis was carried out using limma (version 3.30.13), after

transforming transcript counts using the included voom function. We then fitted a linear

model to sample measurements from each combination of knockouts (including RC9 control)

and conditions (2i and N24). RNAseq batches were included in the model as a confounding

factor. Contrasts were fitted to determine the expression differences between a) each

knowckout sample in 2i and the RC9 control, b) each knockout sample in N24 and the RC9

control, and c) the difference between the log-fold-changes calculated in a) and b)

(interaction effect). We also fitted a contrast to determine changes during normal

differentiation, i.e. the differential expression between RC9N24 *versus* RC92i. All fitted

contrasts were tested for differential expression using moderated t-statistics through the

limma functions eBayes (H0 of no change) or treat (H0 interval [-log2(1.5), log2(1.5)]). p-

Values were corrected for multiple testing using the Benjamini Hochberg method (FDR). For

subsequent analyses the more stringent p-values and FDR values as calculated by treat were

used (unless noted otherwise). Differential expression analysis of 2i/LIF vs. 2i RC9

measurements was carried out separately, but in an identical fashion.

GO enrichment analysis

Mouse gene GO annotations were extracted from the R package org.Mm.eg.db (version

3.4.0). GO terms that included between 5 and 500 genes were selected for further analysis.

For each gene list of interest, significance of functional enrichment of GO terms compared to

the background list was determined using Fisher's exact tests. The background was restricted

to genes whose transcripts were detected at a median count above 5. To reduce redundancy

of functional categories, we clustered GO terms that differed in 5 or fewer genes of interest

(using the R base function hclust on the L1-distance of the binary membership matrix). The

smallest GO term by total annotations was selected independently from Fisher test results as

the primary, i.e. most specific term to represent each cluster. We then carried out correction

for multiple hypothesis testing using the Benjamini-Hochberg method on all primary GO

terms to obtain adjusted p-values.

Naive marker-dependency of N24 KO expression patterns (NAGs)

To quantify the similarity of each gene's differential expression (DE) pattern across KOs at N24 with the DE pattern of seven core pluripotency markers (Nanog, Esrrb, Tbx3, Tfcp2l1, Klf4, Prdm14 and Zfp42), multiple regression analysis was performed. I.e., each gene's DE pattern was modelled as a linear combination of marker DE patterns, quantifying the strength of association as their shared variance (multiple-R^2). NAGs were defined as having a naïve marker R^2 of ≥ 0.7.

Cluster analysis (constitutive & N24 induced)

Based on the responses to the knockout conditions (excluding *Suz12*, *Eed* & *Csnk1a1* KOs) in both N24 and 2i and their interactions with the RC9 control, we defined two main categories of responsive genes: (1) the constitutive knockout response: genes that were significantly changed (adj. p-≤ 0.05) in the same direction in both N24 and 2i, in at least one knockout condition. Genes from the N24 induced KO response category were excluded. (2) N24 induced knockout response: genes that were significantly changed in both N24 and the knockout:RC9 interaction term, in at least one knockout condition. To determine whether genes in the N24 induced knockout response subset were overall more correlated with naive pluripotency markers, we checked the distribution of naive marker multiple-R2 values in the 'N24 induced knockout response' set compared to both the background distribution and genes classified as part of the constitutive KO response.

To determine if distinct functional clusters of genes that may be activated in all, single, or sub-groups of knockouts, we clustered genes in either of the categories (constitutive and N24 induced response) based on their KO^{N2} vs. $RC9^{N24}$ log-fold-changes, using the R base package

hclust with the "Ward.D2" method. We determined the total within-cluster variance at

different numbers of clusters. The number of clusters was chosen such that the model

improved strongly by increasing the number of clusters up to that value, but only marginal

gains result from increasing it further. Thereby 12 gene clusters in the constitutive response

and 12 clusters in the induced response were identified. GO enrichment analysis was carried

out as described above for member genes of each cluster

Differentiation pathway correlation analysis

We defined downstream responses of key differentiation pathways using the most specific

responses to the knockouts of their upstream regulators: *Tsc2* (mTOR), *Ptpn11* & *Fgfr1*

(Fgf/Erk signalling), *Tcf7l1* (Wnt signalling), and *Rbpj* (Notch signalling). In addition, LIF-specific

changes were defined as the response of genes in RC9 cells in the 2i/LIF compared to the 2i

condition. To ensure specificity, we required that the log-fold-change of each response gene

for a given regulator (in N2B27 medium) exceeded that observed in the remaining conditions

by a factor of two. In the case of Ptpn11 & Fgfr1, this criterion had to be met by the lower of

the two log-fold-changes for each gene. Additionally, the genes were required to satisfy a

FPKM (CPM in case of LIF) cut-off of at least 1 in either the KO or the WT. From all remaining,

significantly expressed genes (FDR ≤ 0.05) we then selected, by absolute log-fold-change in

descending order, the top 50 most specific genes for each pathway (Supplementary Figure

6a).

1188

Pathway activity was determined across all other knockouts at N24. For each pathway, the

log-fold-changes of the 50 pathway-specific response genes from the associated regulator

knockout (KO^{N24} vs. $RC9^{N24}$) were selected. We then correlated these log-fold-changes to the

log-fold-changes of the same genes in all other knockouts. A positive correlation indicates

that the pathway is disrupted in a manner similar to knocking out its upstream regulator,

whereas a negative correlation indicates the opposite. In the case of LIF-specific genes, a

positive correlation indicates an effect similar to that of adding LIF. The analysis was repeated

for all 2i experiments (using the same lists of specific genes as in N24). P-values of correlation

values were calculated based on Fisher's Z transform. P-values were then corrected for

multiplicity using the Bonferroni method.

1199

To determine how predictive the calculated correlations were of the differentiation

phenotype (mean naive marker logFC vs. RC9 in N2B27), we carried out multiple linear

regression and extracted the coefficients of the model to establish the relative contribution

of each pathway signature to naïve marker deregulation.

Quantification of differentiation delay in knockouts

Raw RNA sequencing data was aligned to the mm10 genome and read counts were obtained

using the STAR aligner (version2.5.2b). Gene expression for the following biotypes was

quantified: protein_coding, misc_RNA, miRNA, scaRNA, scRNA, snoRNA, snRNA, sRNA. Read

counts were normalized using DESeq2 (version 1.22.2) and subsequently the log2fc between

all samples and the mean expression of the 2i samples was calculated. This resulted in

expression profiles for each gene over 32 hours of differentiation.

1211

Gaussian process regression was used to smoothen the profiles of each gene (R package tgp

version 2.4-14). The expression profiles of 73 knockouts at N24 were mapped on the

differentiation axis from the time course to quantify differentiation delays per KO, i.e. we

aimed to quantify how many hours the KO expression pattern is 'behind' the expected

differentiation in WT cells. This 'mapping' to the time axis was done first computing the logFCs

between N24 versus 2i for each KO. The resulting logFC profiles were compared to the logFCs

at each time point during the WT differentiation. We assumed that the time point at which

the difference between the WT profile and the KO profile is minimized best reflects the

molecular differentiation state of the respective KO.

We used the Euclidean distance for this purpose. To make distances better interpretable

between the knockouts the distances were set to:

1223

1224 $$\frac{Euclidean\ distance - min\ Euclidean\ distance}{max\ Euclidean\ distance - min\ Euclidean\ distance}$$

1225

1226   where the maximum and minimum distances refer to the respective maximum and minimum

1227   Euclidean distances of the respective KO across all time points (i.e. the worst and the best

1228   matching time point). Thus, the best matching time point will get a distance score of 0.

1229

In order to further increase the time resolution, additional time points were imputed via linear

interpolation. Here we interpolated expression profiles every 15 minutes between the two

neighbouring time points of the time point with the smallest Euclidean distance. Timing of

knockouts was repeated on the interpolated time points and the new minimal distance was

used to quantify the differentiation delay.

Prediction of cluster expression by pathway activity

Pathway activities for each pathway in all KOs were calculated using following formulas:

1237

$$directed\ FC(KO) = FC\left(\frac{KO}{WT}\right) * \begin{cases} LIF \begin{cases} +1\ same\ direction\ as\ in\ \frac{LIF}{2i}\ comparison \\ -1\ opposite\ direction\ as\ in\ \frac{LIF}{2i}\ comparison \end{cases} \\ Tsc2,P11F1,Tcf7l1,Rbpj \begin{cases} +1\ same\ direction\ as\ in\ \frac{rep.KO}{WT}\ comparison \\ -1\ opposite\ direction\ as\ in\ \frac{rep.KO}{WT}\ comparison \end{cases} \end{cases}$$

1238

1239

$$directed\ avg.logFC_{KO} = \frac{\sum_n^{PW\ specific\ marker\ genes} directed\ FC(KO)_n}{\sum_n^{PW\ specific\ marker\ genes} directed\ FC(rep.KO)_n}$$

1240

1241

$$pathway\ activity_{KO} = directed\ avg.logFC_{KO} * \begin{cases} 1\ for\ LIF,Tsc2 \\ -1\ for\ P11F1,Tcf7l1,Rbpj \end{cases}$$

1242

The combinations of pathway activities and average cluster expression changes over all KOs

were used to build a multiple regression model to predict the expression changes in each

cluster by the five pathway activities. Connections between pathway activities and cluster

expression changes not making a significance cut-off (padj <= 0.01) of the multiple regression

models were excluded from each model. The remaining model was tested in the time course

data. Connections from clusters where prediction of expression changes based on pathway

activities did follow or contradict the observed expression changes in the time course were

excluded.

1251

Embryo comparison

Sequencing data were obtained from the European Nucleotide Archive [69] from single-cell

mouse embryo profiling studies [10, 70]. Human capacitation dataset was downloaded from

Gene Expression Omnibus website (GSE123055). The macaque FKPM expression dataset was

provided by [33]. Orthologues genes mapping in 1-to-1 fashion were used during comparative

analysis. Alignments to gene loci were quantified with htseq-count [71] based on annotation

from Ensembl 87 [69]. Principal component and cluster analysis were based on log2 FPKM values

computed with the Bioconductor package DESeq [72], custom scripts and FactoRmineR

packages [73]. High variable genes were calculated by fitting a non-linear regression curve

between average log2 FPKM and the square of coefficient of variation. Specific thresholds

were applied along the x-axis (log2FPKM) and y-axis (coefficient of variation) to identify the

most variable genes. Fractional identity between the bulk RNAseq (2i, N24) and the mouse,

macaque embryo or human capacitation dataset were computed using R package

DeconRNASeq [74].

GENEES (Genetic Network Explorer for Embryonic Stem Cell Differentiation)

To both visualize and interrogate gene expression analysis of all KO ES cells we have set up

the online tool GENEES (Genetic Network Exploration of Embryonic Stem Cell Differentiation).

It is accessible at [http://shiny.cecad.uni-koeln.de:3838/stemcell_network/; User: stemcells;

Pw: S3m7KmIfHr9]. The app integrates multiple visualisations and summary statistics to make

all information, from the gene level to network modules / pathways / GO terms, directly

available for exploratory analysis. The app has 4 different tabs for inspection of single genes,

predefined genesets, pre-computed clusters and custom genesets. Single genes can be

inspected in the "Genes tab", while predefined genesets are available for selection under the

"Genesets" tab. The previously described constitutive clusters and InteractionN2 clusters can

be further inspected in the "Pre-computed clusters" tab. In the "Select custom geneset" tab,

a set of genes of interest subjected to analysis.

References

1.   Smith, A. Formative pluripotency: the executive phase in a developmental continuum.
Development **144**, 365-373 (2017).

2.   Kalkan, T. *et al.* Tracking the embryonic stem cell transition from ground state pluripotency.
Development **144**, 1221-1234 (2017).

3.   Buecker, C. *et al.* Reorganization of Enhancer Patterns in Transition from Naive to Primed
Pluripotency. *Cell Stem Cell* **14**, 838-853 (2014).

4.   Dunn, S.J., Martello, G., Yordanov, B., Emmott, S. & Smith, A.G. Defining an essential
transcription factor program for naive pluripotency. *Science* **344**, 1156-1160 (2014).

5.   Niwa, H. The principles that govern transcription factor network functions in stem cells.
Development **145**, dev157420 (2018).

6.   Chen, X. *et al.* Integration of external signaling pathways with the core transcriptional
network in embryonic stem cells. *Cell* **133**, 1106 - 1117 (2008).

7.   Ying, Q.L. *et al.* The ground state of embryonic stem cell self-renewal. *Nature* **453**, 519-523
(2008).

8.   Boroviak, T., Loos, R., Bertone, P., Smith, A. & Nichols, J. The ability of inner-cell-mass cells to
self-renew as embryonic stem cells is acquired following epiblast specification. *Nature cell*
biology **16**, 516-528 (2014).

9.   Acampora, D. *et al.* Loss of the Otx2-Binding Site in the Nanog Promoter Affects the Integrity
of Embryonic Stem Cell Subtypes and Specification of Inner Cell Mass-Derived Epiblast. *Cell*
reports **15**, 2651-2664 (2016).

10.   Mohammed, H. *et al.* Single-Cell Landscape of Transcriptional Heterogeneity and Cell Fate
Decisions during Mouse Early Gastrulation. *Cell reports* **20**, 1215-1228 (2017).

11.   Leeb, M. & Wutz, A. Derivation of haploid embryonic stem cells from mouse embryos.
Nature **479**, 131-134 (2011).

12.   Leeb, M., Dietmann, S., Paramor, M., Niwa, H. & Smith, A. Genetic exploration of the exit
from self-renewal using haploid embryonic stem cells. *Cell Stem Cell* **14**, 385-393 (2014).

13.   Li, M. *et al.* Genome-wide CRISPR-KO Screen Uncovers mTORC1-Mediated Gsk3 Regulation
in Naive Pluripotency Maintenance and Dissolution. *Cell reports* **24**, 489-502 (2018).

14.   Betschinger, J. *et al.* Exit from Pluripotency Is Gated by Intracellular Redistribution of the
bHLH Transcription Factor Tfe3. *Cell* **153**, 335-347 (2013).

15.   Guo, G., Huang, Y., Humphreys, P., Wang, X. & Smith, A. A PiggyBac-based recessive
screening method to identify pluripotency regulators. *PloS one* **6**, e18189 (2011).

16.   Mulas, C., Kalkan, T. & Smith, A. NODAL Secures Pluripotency upon Embryonic Stem Cell
Progression from the Ground State. *Stem cell reports* **9**, 77-91 (2017).

17.   Kokubu, C. & Takeda, J. When half is better than the whole: advances in haploid embryonic
1313   stem cell technology. *Cell Stem Cell* **14** (2014).

1314   18.   Elling, U. *et al.* Forward and reverse genetics through derivation of haploid mouse embryonic
1315   stem cells. *Cell Stem Cell* **9**, 563-574 (2011).

1316   19.   Kalkan, T. & Smith, A. Mapping the route from naive pluripotency to lineage specification.
1317   *Philosophical Transactions of the Royal Society of London B: Biological Sciences* **369** (2014).

1318   20.   Ye, J. & Blelloch, R. Regulation of pluripotency by RNA binding proteins. *Cell Stem Cell* **15**,
1319   271-280 (2014).

1320   21.   Kalkan, T. *et al.* Complementary Activity of ETV5, RBPJ, and TCF3 Drives Formative Transition
1321   from Naive Pluripotency. *Cell Stem Cell* **24**, 785-801.e787 (2019).

1322   22.   Ursu, A. *et al.* Epiblastin A Induces Reprogramming of Epiblast Stem Cells Into Embryonic
1323   Stem Cells by Inhibition of Casein Kinase 1. *Cell Chem Biol* **23**, 494-507 (2016).

1324   23.   Martello, G. & Smith, A. The nature of embryonic stem cells. *Annual review of cell and
1325   developmental biology* **30**, 647-675 (2014).

1326   24.   Qiu, D. *et al.* Klf2 and Tfcp2l1, Two Wnt/β-Catenin Targets, Act Synergistically to Induce and
1327   Maintain Naive Pluripotency. *Stem cell reports* **5**, 314-322 (2015).

1328   25.   Hall, J. *et al.* Oct4 and LIF/Stat3 additively induce Kruppel factors to sustain embryonic stem
1329   cell self-renewal. *Cell Stem Cell* **5**, 597-609 (2009).

1330   26.   Yeo, J.C. *et al.* Klf2 is an essential factor that sustains ground state pluripotency. *Cell Stem
1331   Cell* **14**, 864-872 (2014).

1332   27.   Martello, G. *et al.* Esrrb is a pivotal target of the Gsk3/Tcf3 axis regulating embryonic stem
1333   cell self-renewal. *Cell Stem Cell* **11**, 491-504 (2012).

1334   28.   Moreira, S. *et al.* Endogenous BioID elucidates TCF7L1 interactome modulation upon GSK-3
1335   inhibition in mouse ESCs. *bioRxiv*, 431023 (2018).

1336   29.   Wray, J. *et al.* Inhibition of glycogen synthase kinase-3 alleviates Tcf3 repression of the
1337   pluripotency network and increases embryonic stem cell resistance to differentiation.
1338   *Nature cell biology* **13**, 838-845 (2011).

1339   30.   Shy, B.R. *et al.* Regulation of Tcf7l1 DNA binding and protein stability as principal
1340   mechanisms of Wnt/β-catenin signaling. *Cell reports* **4**, 1-9 (2013).

1341   31.   Mayer, D. *et al.* Zfp281 orchestrates interconversion of pluripotent states by engaging Ehmt1
1342   and Zic2. *The EMBO journal* **39**, e102591 (2020).

1343   32.   Rossant, J. & Tam, P.P.L. Exploring early human embryo development. *Science* **360**, 1075-
1344   1076 (2018).

1345   33.   Nakamura, T. *et al.* A developmental coordinate of pluripotency among mice, monkeys and
1346   humans. *Nature* **537**, 57-62 (2016).

34.   Rostovskaya, M., Stirparo, G.G. & Smith, A. Capacitation of human naïve pluripotent stem
cells for multi-lineage differentiation. *Development* **146**, dev172916 (2019).

35.   Artus, J., Kang, M., Cohen-Tannoudji, M. & Hadjantonakis, A.-K. PDGF signaling is required
for primitive endoderm cell survival in the inner cell mass of the mouse blastocyst. *Stem cells*
(Dayton, Ohio) **31**, 1932-1941 (2013).

36.   Boroviak, T. *et al.* Single cell transcriptome analysis of human, marmoset and mouse
embryos reveals common and divergent features of preimplantation development.
Development **145** (2018).

37.   Plusa, B., Piliszek, A., Frankenberg, S., Artus, J. & Hadjantonakis, A.-K. Distinct sequential cell
behaviours direct primitive endoderm formation in the mouse blastocyst. *Development* **135**,
3081-3091 (2008).

38.   Smith, A.G. *et al.* Inhibition of pluripotential embryonic stem cell differentiation by purified
polypeptides. *Nature* **336**, 688-690 (1988).

39.   Watanabe, S. *et al.* Activation of Akt signaling is sufficient to maintain pluripotency in mouse
and primate embryonic stem cells. *Oncogene* **25**, 2697-2707 (2006).

40.   Kunath, T. *et al.* FGF stimulation of the Erk1/2 signalling cascade triggers transition of
pluripotent embryonic stem cells from self-renewal to lineage commitment. *Development*
134, 2895-2902 (2007).

41.   Lowell, S., Benchoua, A., Heavey, B. & Smith, A.G. Notch promotes neural lineage entry by
pluripotent embryonic stem cells. *PLoS biology* **4**, e121 (2006).

42.   Molotkov, A., Mazot, P., Brewer, J.R., Cinalli, R.M. & Soriano, P. Distinct Requirements for
FGFR1 and FGFR2 in Primitive Endoderm Development and Exit from Pluripotency.
Developmental cell **41**, 511-526 e514 (2017).

43.   Lake, R.J., Tsai, P.-F., Choi, I., Won, K.-J. & Fan, H.-Y. RBPJ, the Major Transcriptional Effector
of Notch Signaling, Remains Associated with Chromatin throughout Mitosis, Suggesting a
Role in Mitotic Bookmarking. *PLoS genetics* **10**, e1004204 (2014).

44.   Ohtsuka, S., Nakai-Futatsugi, Y. & Niwa, H. LIF signal in mouse embryonic stem cells. *JAK-*
STAT **4**, e1086520-e1086520 (2015).

45.   Villegas, F. *et al.* Lysosomal Signaling Licenses Embryonic Stem Cell Differentiation via
Inactivation of Tfe3. *Cell Stem Cell* **24**, 257-270 e258 (2019).

46.   Niwa, H., Ogawa, K., Shimosato, D. & Adachi, K. A parallel circuit of LIF signalling pathways
maintains pluripotency of mouse ES cells. *Nature* **460**, 118-122 (2009).

47.   Hoffman, J.A., Wu, C.I. & Merrill, B.J. Tcf7l1 prepares epiblast cells in the gastrulating mouse
embryo for lineage specification. *Development* **140**, 1665-1675 (2013).

48.   Xu, T. *et al.* RBPJ/CBF1 interacts with L3MBTL3/MBT1 to promote repression of Notch
signaling via histone demethylase KDM1A/LSD1. *The EMBO journal* **36**, 3232-3249 (2017).

49.   Amano, T. *et al.* Zscan4 restores the developmental potency of embryonic stem cells. *Nature*
Communications **4**, 1966 (2013).

50. Macfarlan, T.S. *et al.* Embryonic stem cell potency fluctuates with endogenous retrovirus
activity. *Nature* **487**, 57-63 (2012).

51. Chen, E.Y. *et al.* Enrichr: interactive and collaborative HTML5 gene list enrichment analysis
tool. *BMC Bioinformatics* **14**, 128 (2013).

52. Quenneville, S. *et al.* In Embryonic Stem Cells, ZFP57/KAP1 Recognize a Methylated
Hexanucleotide to Affect Chromatin and DNA Methylation of Imprinting Control Regions.
Molecular Cell **44**, 361-372 (2011).

53. Takahashi, K. & Yamanaka, S. Induction of pluripotent stem cells from mouse embryonic and
adult fibroblast cultures by defined factors. *Cell* **126**, 663-676 (2006).

54. Xie, H., Ye, M., Feng, R. & Graf, T. Stepwise Reprogramming of B Cells into Macrophages. *Cell*
117, 663-676 (2004).

55. Lassar, A.B., Paterson, B.M. & Weintraub, H. Transfection of a DNA locus that mediates the
conversion of 10T1 2 fibroblasts to myoblasts. *Cell* **47**, 649-656 (1986).

56. Ramos-Ibeas, P. *et al.* Pluripotency and X chromosome dynamics revealed in pig pre-
gastrulating embryos by single cell analysis. *Nature Communications* **10**, 500 (2019).

57. Bredenkamp, N. *et al.* Efficient RNA-mediated reprogramming of human somatic cells to
naïve pluripotency facilitated by tankyrase inhibition. *bioRxiv*, 636670 (2019).

58. Chambers, I. & Tomlinson, S.R. The transcriptional foundation of pluripotency. *Development*
136, 2311-2322 (2009).

59. Nichols, J. & Smith, A. Pluripotency in the embryo and in culture. *Cold Spring Harb Perspect*
Biol **4**, a008128 (2012).

60. Boroviak, T. & Nichols, J. The birth of embryonic pluripotency. *Philosophical Transactions of*
the Royal Society B: Biological Sciences **369** (2014).

61. Boroviak, T. & Nichols, J. Primate embryogenesis predicts the hallmarks of human naïve
pluripotency. *Development (Cambridge, England)* **144**, 175-186 (2017).

62. Rossant, J. & Tam, P.P.L. New Insights into Early Human Development: Lessons for Stem Cell
Derivation and Differentiation. *Cell Stem Cell* **20**, 18-28 (2017).

63. Baker, O. *et al.* RAC-tagging: Recombineering And Cas9-assisted targeting for protein tagging
and conditional analyses. *Sci Rep* **6**, 25529 (2016).

64. Sarov, M. *et al.* A genome-scale resource for in vivo tag-based protein function exploration
in C. elegans. *Cell* **150**, 855-866 (2012).

65. Mali, P. *et al.* RNA-Guided Human Genome Engineering via Cas9. *Science* **339**, 823-826
(2013).

66. Leeb, M., Perry, A.C.F. & Wutz, A. Establishment and Use of Mouse Haploid ES Cells, in
Current Protocols in Mouse Biology (John Wiley & Sons, Inc., 2015).

67. Leeb, M. *et al.* Germline potential of parthenogenetic haploid mouse embryonic stem cells.
Development **139**, 3301-3305 (2012).

68. Yusa, K., Zhou, L., Li, M.A., Bradley, A. & Craig, N.L. A hyperactive piggyBac transposase for
mammalian applications. *Proceedings of the National Academy of Sciences of the United*
States of America **108**, 1531-1536 (2011).

69. Toribio, A.L. *et al.* European Nucleotide Archive in 2016. *Nucleic Acids Res* **45**, D32-D36
(2017).

70. Deng, Q., Ramsköld, D., Reinius, B. & Sandberg, R. Single-Cell RNA-Seq Reveals Dynamic,
Random Monoallelic Gene Expression in Mammalian Cells. *Science* **343**, 193-196 (2014).

71. Anders, S., Pyl, P.T. & Huber, W. HTSeq—a Python framework to work with high-throughput
sequencing data. *Bioinformatics* **31**, 166-169 (2014).

72. Anders, S. & Huber, W. Differential expression analysis for sequence count data. *Genome*
biology **11**, R106 (2010).

73. Lê, S., Josse, J. & Husson, F. FactoMineR: An R Package for Multivariate Analysis. *2008* **25**, 18
(2008).

74. Gong, T. & Szustakowski, J.D. DeconRNASeq: a statistical framework for deconvolution of
heterogeneous tissue samples based on mRNA-Seq data. *Bioinformatics* **29**, 1083-1085
(2013).