Sex determination through X-Y heterogamety in *Salix nigra*

Brian J. Sanderson[1,8,9], Guanqiao Feng[1], Nan Hu[1], Julianne Grady[2], Craig H. Carlson[3], Lawrence

B. Smart[3], Ken Keefover-Ring[4], Tongming Yin[5], Tao Ma[6], Jianquan Liu[6,7], Stephen P. DiFazio[2],

Matthew S. Olson[1]

[1]Department of Biological Sciences, Texas Tech University, Lubbock, TX 79409-3131 USA

[2]Department of Biology, West Virginia University, Morgantown, WV, 26506 USA

[3]Horticulture Section, School of Integrative Plant Science, Cornell University, New York State

Agricultural Experiment Station, Geneva, New York 14456 USA

[4]Departments of Botany and Geography, University of Wisconsin-Madison, 430 Lincoln Dr.,

Madison, WI 53706, USA

[5]Key Laboratory of Tree Genetics and Biotechnology of Jiangsu Province and Education

Department of China, Nanjing Forestry University, Nanjing, China

[6]Key Laboratory of Bio-Resource and Eco-Environment of Ministry of Education & College of

Life Sciences, Sichuan University, Chengdu 610065, China

[7]State Key Laboratory of Grassland Agro-Ecosystem, Institute of Innovation Ecology & College

of Life Sciences, Lanzhou University, Lanzhou 730000, China

[8]Author for correspondence: brian@biologicallyrelevant.com

[9]Current address: Department of Biology, West Virginia University, Morgantown, WV, 26506-

6057 USA

Abstract

The development of non-recombining sex chromosomes has radical effects on the evolution of

discrete sexes and sexual dimorphism. Although dioecy is rare in plants, sex chromosomes have

evolved repeatedly throughout the diversification of angiosperms, and many of these sex

chromosomes are relatively young compared to those found in vertebrates. The Salicaceae, the

family of plants that includes poplars and willows, contains predominantly dioecious species and

dioecy likely evolved prior to the split of willows and poplars over 70 million years ago. Yet,

despite this shared origin of dioecy both the chromosomal location of the sex determination

region and the pattern of sex chromosome heterogamety differ between and within these genera.

In this study we identified a novel sex-linked region (SLR) in *Salix nigra*, a basal species in the

willow clade, and demonstrated that this species has XY heterogamety. We did not detect any

genetic overlap with the previously characterized ZW SLRs in willows, which map to a different

chromosome. The *S. nigra* SLR is characterized by strong recombination suppression across a

2MB region. The non-recombining portion of the X chromosome also exhibits a pattern of low

Tajima's D, consistent with a recent selective sweep of the X chromosome in this region.

Key words: Sex determination; sex chromosome; *Salix*; Salicaceae; targeted sequence capture

Introduction

Some groups of animals exhibit surprisingly frequent movement in the genomic locations of sex
determination loci (Gammerdinger & Kocher, 2018; Miura, 2007), but it is not yet clear whether
similar patterns are present in plants, where the evolution of separate sexes (called dioecy in
plants) is more frequent than in most animal groups (Beukeboom & Perrin, 2014). In mammals
(Cortez et al., 2014; Lahn, 1999) and birds (Ellegren, 2010; Shetty et al., 1999; but see Sigeman
et al., 2019) most species have maintained similar sex chromosomes for over 80-130 My. In other
animal groups, however, sex chromosome transitions appear more common. For instance, among
several genera of cichlid fishes the sex determination regions reside on 12 different sex
chromosomes, with both male and female heterogametic systems (XY and ZW, respectively;
Gammerdinger & Kocher, 2018). Similarly, Dipteran flies harbor more than six different sex
chromosomes, and multiple sex chromosome transitions have occurred within the genus
Drosophila (Vicoso & Bachtrog, 2015). In plants, however, few taxonomic groups with the same
origin of dioecy have been studied in detail. One notable exception is in octoploid *Fragaria*
(strawberry), where a cassette of sex determination genes has moved among homeologs and
picked up linked genomic regions as it has moved, likely all within the last 1My (Tennessen et
al., 2018). In this regard, the willows (genus *Salix*) are of particular interest because they are
exclusively dioecious and diverged from a likely dioecious ancestor in the Cretaceous, over 70
million years ago (Zhang et al., 2018).

Chromosome 15 was originally identified as the location of the SLR in *Salix* using
anonymous sex-linked markers (Temmel et al., 2007), which was later confirmed in three species
of willow from subgenus Vetrix using genome resequencing and mapping (*Salix viminalis*:

Pucholt et al., 2015; *S. suchowensis*: Hou et al., 2015; and *S. purpurea*: Zhou et al., 2018). The

most intensively studied of these species is *S. purpurea*, in which the sex-linked region (SLR) is a

6.7 MB nonrecombining region (Zhou et al., 2018). The location and pattern of heterogamety of

the SLR in these species of *Salix* differ from those found in the sister genus *Populus*, where the

SLR is commonly found on chromosome 19 with XY heterogamety (Geraldes et al., 2015; Pakull

et al., 2011, 2009). These differences in the location of the SLR and pattern of heterogamety have

led to the hypothesis that dioecy evolved independently in these two genera (Hou et al., 2015).

An alternative hypothesis of a single origin of dioecy and the movement of a common set of sex

determination genes that are shared across both genera, however, is supported by a set of three

genes found in the SLRs of both *S. purpurea* and *P. trichocarpa* (Zhou et al., 2018, 2020).

The goal of the present study was to identify an SLR in a novel subgenus of *Salix* to

address whether its location and heterogamety is common among widely diverged willow

species. *Salix nigra* is a tree-form willow in the subgenus Protitea that is basal in the *Salix* lineage

(Argus, 1997; Barkalov & Kozyrenko, 2014). Using targeted sequence capture to selectively

genotype 24 males and 24 females with genome-wide markers, we show that a region on

chromosome 7 exhibits highly divergent genotypes between males and females, indicating that

this interval harbors the SLR. A comparison of heterozygosity between males and females, as

well as slow decay of linkage disequilibrium in this region among males, is consistent with an

XY sex determination system, which we confirmed with amplicon sequencing. Assuming that

dioecy arose only once within the genus, these results suggest that sex chromosome transitions

have occurred in the genus *Salix* and that differences in the chromosomal location of the SLR do

not reflect independent origins of dioecy in the family, but rather a movement of the SLR

between chromosomes.

Methods

Probe Design

To consistently genotype common sites in a reduced representation of the genome, we used a

targeted sequence capture array (RNA bait hybridization) that we custom designed to efficiently

capture exonic regions throughout the genome across species in the *Populus + Salix* clade of the

Salicaceae. Because the efficiency of RNA bait binding, and thus capture efficiency, is reduced

as target regions diverge due to sequence polymorphism (Lemmon & Lemmon, 2013), we

quantified sequence polymorphism among whole-genome resequencing data from an outgroup,

Idesia polycarpa, and a diverse array of *Populus* and *Salix* species, including *P. tremuloides*, *P.*

trichocarpa, *P. mexicana*, *P. euphratica*, *P. deltoides*, and *S. purpurea*. The whole-genome short

reads of the *Populus* and *Salix* species were aligned to the *Populus trichocarpa* genome assembly

version 3 (Tuskan et al., 2006) using bwa mem (Li, 2013), single nucleotide polymorphisms

(SNPs) and insertion-deletion mutations (indels) were identified using samtools mpileup (Li,

2011), and read depth for the variant calls was quantified using vcftools v. 0.1.15 (Danecek et al.,

2011). Custom Python and R scripts were used to identify variant and indel frequencies for all

exons in the *P. trichocarpa* genome annotation.

Orthologs for our candidate loci in the in the *Salix purpurea* genome assembly version 1

(Zhou et al., 2018) were identified using a list of orthologs between the *P. trichocarpa* and *S.*

purpurea genomes prepared by Phytozome (Goodstein et al., 2012) using a tree-based approach.

We further screened candidate regions to exclude high-similarity duplicated regions by accepting

only loci with single BLAST hits against the highly contiguous assembly of *S. purpurea* version

5 (Zhou et al., 2020), which is less fragmented than the *S. purpurea* version 1. For the remaining

loci, we selected regions across which at least 360 base pairs (bp) of exon sequence contained 2-

12% difference in single nucleotides and fewer than 2 indels across all aligned species. For a

majority of these genes, we identified multiple regions within each exon. These candidate regions

were sent to Arbor Biosciences (Ann Arbor, MI, USA) for probe synthesis. Probes were designed

with 50% overlap across the targeted regions, so that each targeted nucleotide position would

potentially be captured by two probes. The final capture array consisted of 60,000 probes that

target exonic sequences for 16,580 genes.

Library Preparation and Sequence Capture

The sex of 24 males and 24 females of *Salix nigra* was identified from flowering catkins in a wild

population near Dickens Springs, TX, USA. Leaf tissues were collected and dried on silica beads.

DNA was extracted from leaf tissue using the Qiagen DNeasy Plant Minikit (Qiagen, Hilden,

Germany) and fragmented using sonication with the Covaris E220 Focused Ultrasonicator

(Covaris, Inc., Woburn, MA, USA). Libraries were prepared using the NEBNext Ultra II DNA

Prep Kit (New England Biolabs, Ipswitch, MA, USA), and quantified using an Agilent

Bioanalyzer 2100 DNA 1000 kit (Agilent Technologies, Santa Clara, CA, USA). Libraries were

pooled at equimolar concentrations into sixteen pools of six prior to probe hybridization to our

targeted capture baits following the Arbor Biosciences myBaits protocol v 3.0.1. The hybridized

samples were subsequently pooled at equimolar ratios and sequenced at the Oklahoma Medical

Research Foundation (Oklahoma, OK, USA) using a HiSeq 3000 (Illumina, Inc., San Diego, CA,

USA).

Variant calls and hard filtering

Read data were aligned to the version 5 assembly of *Salix purpurea*, using bwa mem (Li, 2013).

We estimated the depth of read coverage across all targeted genes using bedtools intersect v.

2.25.0 (Quinlan & Hall, 2010). The alignments were screened for optical duplicates following the

Broad Institute's best practices recommendations (DePristo et al., 2011). Variants were identified

using HaplotypeCaller in GATK v4.0.8.1 (McKenna et al., 2010) and initially called separately

for each individual as GVCFs. The GVCFs were merged and variant sites and indels were called

for all individuals using GenotypeGVCFs in GATK. Based on the distribution of quality scores,

variant sites were screened with the hard filters: "$MQ < 40.00 \; || \; SOR > 3.000 \; || \; QD < 2.000 \; || \; FS$

$> 60.000 \; || \; MQRankSum < -12.500 \; || \; ReadPosRankSum < -8.000 \; || \; ReadPosRankSum > 8.000$".

Individual genotypes were filtered if they were supported by < 6 reads, or > 125 reads. Finally,

genotypes that did not pass filtering criteria were set to no call genotypes (./.).

Location of the sex determination region

To determine the genomic regions exhibiting the greatest differences between males and females,

we performed a genome-wide association study (GWAS) in which we regressed sex (male or

female) on the filtered variants using the program emmax (Kang et al., 2010) using a Bonferroni-

adjusted significance threshold of $P < 0.05$ to assess the significance of sex linkage. We

visualized the results with a Manhattan plot using the R package ggplot2 (Wickham, 2016). We

quantified heterozygosity of genotypes at the loci identified with significant sex linkage directly

from the filtered VCF file using the program vcftools v. 0.1.15 (Danecek et al., 2011).

To confirm the sex association we analyzed an independent population of 16 male and 16

female *S. nigra* trees collected from the vicinity of Morgantown, WV. Primers were designed in

conserved regions that flanked two loci that showed significant sex-linkage in the Texas

population. These were sequenced using Sanger chemistry on an ABI3130XL sequencer. Base

calls were made using PHRED (Ewing & Green, 1998) and sequences were assembled using

PHRAP and Consed (Gordon et al., 1998). Polymorphisms were identified using PolyPhred

(Nickerson, 1997) and confirmed by visualization in Consed. The frequency of heterozygosity for

males and females was compared for all loci with minor allele frequency > 0.35, which

effectively screened out mutations that have arisen since establishment of the SLR, as well as

instances of genotyping or sequencing error.

Population genomics statistics

Kinship was estimated using the relatedness2 algorithm in vcftools, which applies the methods

used in KING (Manichaikul et al., 2010). Linkage disequilibrium (LD) for all pairs of loci was

calculated as r^2 using vcftools v. 0.1.15 (Danecek et al., 2011). LD decay was calculated using

formulas in Hill & Weir (1988). To compare genetic diversity within the *S. nigra* SLR to

diversity other regions of the genome, we calculated per-site nucleotide diversity as π (Nei & Li,

1979) using vcftools v. 0.1.15 (Danecek et al., 2011) after filtering for only biallelic SNPs. We

calculated π separately for males, which carried both X and Y chromosomes, and females, which

carried two X chromosomes. The average π for 5kb sliding windows were calculated using R (R

Core Team, 2018). To assess whether non-neutral processes may have influenced the SLR, we

calculated values of Tajima's D (Tajima, 1989) using the sliding window method implemented in

vcftools v. 0.1.15 (Carlson, 2005; Danecek et al., 2011), with a window size of 25Kb.

The scripts described above as well as the full details of these analyses including the

alignment and variant calling pipeline are available in Jupyter notebooks at

https://github.com/BrianSanderson/salix-nigra-slr.

Results

The sequence capture reads aligned to 184 MB of the *S. purpurea* reference genome before

filtering, of which 21,229,252 sites met our quality and depth filtering criteria. This corresponded

to an average read depth of 6.30 ± 2.00X (mean $\pm$ sd) at sites genome-wide, and an average read

depth of 44.68 ± 2.68X (mean $\pm$ sd) at on-target sites across the 16,580 targeted genes (Table

S1). There were a total of 6,826,811 SNPs among the 48 individuals, of which 2,011,511 passed

the filtering criteria, for an average SNP density in the sample of 95 SNP/kb. The average kinship

among individuals in the population was 0.176, indicating an average degree of kinship between

half-sibs and first cousins in this population (Fig. S1).

A total of 38 SNPs on chromosome 7 between 4.88 Mb and 6.81 Mb exhibited significant

associations with sex ($P_{FDR} < 0.05$; Fig. 1;). We refer to this region as the sex-linked region

(SLR). The average heterozygosity for males among these SNPs was 0.875 ± 0.218, and the

average heterozygosity for females was 0.072 ± 0.148 (mean $\pm$ sd), suggesting that the SLR in

this species exhibits XY heterogamety (Fig. 2; Table S2). Amplicons from two loci designed to

target areas of this region confirmed these patterns in 32 independent individuals, with an average

female heterozygosity of 0.021 ± 0.088, and average male heterozygosity of 0.959 ± 0.067

(mean $\pm$ sd) for 18 SNPs with a MAF > 0.2 (Table S3).

Recombination is suppressed between the X and Y chromosomes in the SLR as shown by

the slow decay in linkage disequilibrium exhibited in males (XY) compared to females (XX) and

the recombining end of chromosome 7 from 7-12 MB (Fig. 3). When mapped onto the annotation

from *S. purpurea*, the SLR includes 111 genes with *Arabidopsis* orthologs (Table S4), including

homologs to cytokinin oxidase (AT5G56970.1) and a SAUR-like auxin responsive protein family

gene (AT3G12830.1) which are both possible candidates in the metabolic pathway controlling

sex determination (Feng et al., 2020).

In males the SLR exhibited elevated heterozygosity and pairwise nucleotide diversity

compared to the average in autosomes (Fig. 2A), a pattern consistent with XY heterogamety. In

females, the SLR exhibited high homozygosity and low pairwise nucleotide diversity compared

to the average in autosomes (Fig. 2B), a pattern consistent with the female carrying two X

chromosomes. Because the female carries two X chromosomes, we can use the population of

females to assess deviations from neutral evolution for the SLR of the X chromosome. The SLR

on the X chromosome exhibited a large negative deviation in Tajima's D compared to the rest of

the genome (Figs. 4B & S2), indicating there is an excess of low- frequency polymorphisms in

this region for females. When the population of males was analyzed, the SLR exhibited a large

positive deviation in Tajima's D (Figs. 4B & S2), consistent with the lack of recombination

between the X and Y chromosomes.

Discussion

We identified a new sex-linked region (SLR) in willows that exhibits XY heterogamety on

chromosome 7 in *Salix nigra*. This result establishes that sex chromosomes in *Salix* have

undergone changes in the genomic location as well as patterns of heterogamety. The previously

identified sex chromosomes in *Salix* have all been in the subgenus Vetrix, and the location and

heterogamety of all of these SLRs was on chromosome 15 ZW (Hou et al., 2015; Pucholt et al.,

2015; Temmel et al., 2007; Zhou et al., 2018). The sex chromosomes of poplars (*Populus* spp.),

the sister genus to *Salix*, have been mapped to two different SLRs on chromosome 19, with *P.*

trichocarpa and *P. tremuloides* exhibiting XY sex determination (Geraldes et al., 2015; Pakull et

al., 2011, 2009) and *P. alba* exhibiting ZW sex determination (Paolucci et al., 2010). We estimate

that the size of the non-recombining SLR in *Salix nigra* is approximately 2MB and contains over

100 genes. However, these estimates are based on a comparison with the *S. purpurea* reference

genome, and thus require confirmation by comparison with an assembled *Salix nigra* genome.

Nonetheless, this SLR appears to be substantially smaller than the 6.7 MB SLR in *S. purpurea*

(Zhou et al., 2020) and 20 times larger than the 100kb SLR in *P. trichocarpa* (Geraldes et al.,

2015).

Dioecy (separate male and female individuals) is found in only 5-6% of angiosperms

(Renner & Ricklefs, 1995) and is unevenly distributed across the angiosperm orders (Renner,

2014). Because many dioecious species have congeners and sister species that are

hermaphroditic, it has been posited the dioecy is an evolutionary dead end (reviewed in Käfer et

al., 2017). The Salicaceae is among a small group of angiosperm families in which all of the

members are dioecious, which provides an opportunity to test whether the location of sex

determination regions is evolutionarily stable, as in mammals (Cortez et al., 2014; Lahn, 1999)

and birds (Ellegren, 2010; Shetty et al., 1999), or it is variable, as in cichlid fishes

(Gammerdinger & Kocher, 2018), frogs (Miura, 2007), and strawberries (Tennessen et al., 2018).

Previous studies have hypothesized that dioecy and the sex determination regions in *Salix* and

Populus evolved independently (Hou et al., 2015), which was previously supported by the

observation that all *Salix* sex chromosomes in the subgenus Vetrix were 15 ZW and most

Populus sex chromosomes were 19 XY. Thus, our discovery of the 7 XY SLR in *Salix nigra*

indicates a previously unknown dynamism in the movement of sex chromosomes in *Salix* and

supports the hypothesis of a single origin of dioecy and sex chromosomes in the Salicaceae. The

mechanism of sex determination in *Populus* is likely controlled by a Type A cytokinin response

regulator (a homology of the *Arabidopsis* gene ARR17) which shows sex association in both

Populus and *Salix* (Bräutigam et al., 2017; Xue et al., 2020; Zhou et al., 2020). Although there is

no homolog of this gene within the corresponding genome interval in *S. purpurea* to which the *S.*

nigra reads mapped, it is intriguing that the region does contain genes involved with cytokinin

and auxin signaling, raising the possibility that a different but related mechanism of sex

determination may be present in this species.

243   The high level of heterozygosity in the 4.8-6.8 MB SLR region in *S. nigra* males is

consistent with XY sex determination, and likely results from the evolutionary divergence

between the non-recombining regions of the X and Y chromosomes. The suppression of

recombination between the X and Y chromosomes in the SLR allows independent evolution of

these two regions. Sequence differences that underlie the high levels of heterozygosity likely

arose from both fixation of polymorphism that was present prior to recombination suppression

and from the occurrence of new mutations since suppression. The pattern of elevated Tajima's D

in the SLR of males directly reflects the high heterozygosity in this region and the independent

evolution of the X and Y chromosomes. The females, however, carry two X chromosomes which

recombine freely at the same rate as the non-SLR region of chromosome 7. We found that

Tajima's D among females was depressed relative to regions flanking the SLR on chromosome 7

and compared to the autosomes (Fig. 4; Fig. S2B), indicating that alleles at segregating sites of

the SLR were rare compared to neutral expectations. This pattern is inconsistent with a recent

reduction in the effective population size on the X, which would result in an increase in Tajima's

D (Wakeley, 2009). The lower Tajima's D on the X also could not have been caused purely by

demographic effects because these would have influenced the entire genome, including the

autosomes. The elevated frequency of rare alleles underlying the reduced levels of Tajima's D is

consistent with a recent selective sweep in the SLR on the X chromosome (Simonsen et al.,

1995). Just after a selective sweep, Tajima's D is expected to decrease dramatically, and as new

mutations are introduced into the region, Tajima's D rapidly recovers to neutral expectations

(Simonsen et al., 1995).

Conclusion

This report of a previously unidentified SLR in *Salix nigra* on chromosome 7 with XY

heterogamety illuminates the dynamic history of the shifting position and dominance

relationships of sex chromosomes in the Salicaceae family, which includes both poplars (*Populus*

spp.) and willows (*Salix* spp.). Patterns of Tajima's D suggest that the X chromosome was

recently impacted by a selective sweep, underscoring the evolutionary dynamism in this region

and suggesting that selective factors have influenced the movement of the SLRs in this plant

family. This discovery indicates that the different locations of sex chromosomes in *Populus* and

Salix more likely result from the movement of the sex chromosome than multiple independent

origin of dioecy in the Salicaceae. This study places the Salicaceae among a small number of

taxonomic groups with high dynamism in the turnover of sex chromosomes. Future research

should focus on understanding whether additional shifts in the location and dominance

relationships of the sex chromosomes have occurred within the family and what mechanisms are

driving these changes.

Acknowledgements

We thank Rao Kottapalli, Pratibha Kottapalli, and Sandra Simon for assistance with library

preparation, and Haley Hale for assistance with DNA extraction. We thank the Dickens County

Judge for permission to sample this *S. nigra* population. This research was supported by grants

from the US National Science Foundation (1542509, 1542599, 1542479, 1542486) and the

National Natural Science Foundation of China (31561123001).

Author Contributions

B.J.S., S.P.D, and M.S.O designed the experiment; S.P.D., L.B.S., K.K-R., T.Y., J.L., and M.S.O.

secured funding for this project. L.B.S., T.M., and J.L. contributed whole-genome data for

sequence capture design; B.J.S and S.P.D developed the targeted sequence capture array; M.S.O.

collected plant materials; B.J.S., G.F., N.H., and J.G. carried out the experiment; B.J.S, S.P.D.,

G.F., C.H.C., and M.S.O. analyzed the data and prepared figures; B.J.S. and M.S.O. drafted the

manuscript, and all authors contributed to revisions of the manuscript.

Data Accessibility Statement

Upon acceptance of this manuscript the short read data from the targeted sequence capture will be

uploaded to the Genbank Sequence Read Archive. All custom scripts used, as well as full Jupyter

analysis notebooks will be made available at https://github.com/BrianSanderson/salix-nigra-slr/,

and upon acceptance of this manuscript a DOI will be generated for this repository to create a

DOI that will reference the state of the repository at the time of acceptance.

Literature Cited

Argus, G. W. (1997). Infrageneric Classification of Salix (Salicaceae) in the New World. *Systematic Botany Monographs*, *52*, 1. https://doi.org/10.2307/25096638

Barkalov, V. Y., & Kozyrenko, M. M. (2014). Phylogenetic relationships of Salix L. subg. Salix species (Salicaceae) according to sequencing data of intergenic spacers of the chloroplast genome and ITS rDNA. *Russian Journal of Genetics*, *50*(8), 828–837. https://doi.org/10.1134/S1022795414070035

Beukeboom, L., & Perrin, N. (2014). *The Evolution of Sex Determination*. Oxford University Press.

Bräutigam, K., Soolanayakanahally, R., Champigny, M., Mansfield, S., Douglas, C., Campbell, M. M., & Cronk, Q. (2017). Sexual epigenetics: gender-specific methylation of a gene in the sex determining region of Populus balsamifera. *Scientific Reports*, *7*(1), 45388. https://doi.org/10.1038/srep45388

Carlson, C. S. (2005). Genomic regions exhibiting positive selection identified from dense genotype data. *Genome Research*, *15*(11), 1553–1565. https://doi.org/10.1101/gr.4326505

Cortez, D., Marin, R., Toledo-Flores, D., Froidevaux, L., Liechti, A., Waters, P. D., Grützner, F., & Kaessmann, H. (2014). Origins and functional evolution of Y chromosomes across mammals. *Nature*, *508*(7497), 488–493. https://doi.org/10.1038/nature13151

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Handsaker, R. E., Lunter, G., Marth, G. T., Sherry, S. T., McVean, G., & Durbin, R. (2011). The variant call format and VCFtools. *Bioinformatics*, *27*(15), 2156–2158. https://doi.org/10.1093/bioinformatics/btr330

DePristo, M. A., Banks, E., Poplin, R., Garimella, K. V., Maguire, J. R., Hartl, C., Philippakis, A. A., Angel, G. del, Rivas, M. A., Hanna, M., McKenna, A., Fennell, T. J., Kernytsky, A. M., Sivachenko, A. Y., Cibulskis, K., Gabriel, S. B., Altshuler, D., & Daly, M. (2011). A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nature Genetics*, *43*(5), 491–498. https://doi.org/10.1038/ng.806

Ellegren, H. (2010). Evolutionary stasis: the stable chromosomes of birds. *Trends in Ecology & Evolution*, *25*(5), 283–291. https://doi.org/10.1016/j.tree.2009.12.004

Ewing, B., & Green, P. (1998). Base-Calling of Automated Sequencer Traces Using Phred. II. Error Probabilities. *Genome Research*, *8*(3), 186–194. https://doi.org/10.1101/gr.8.3.186

Feng, G., Sanderson, B. J., Keefover-Ring, K., Liu, J., Ma, T., Yin, T., Smart, L. B., DiFazio, S. P., & Olson, M. S. (2020). Pathways to sex determination in plants: how many roads lead to Rome? *Current Opinion in Plant Biology*, *54*, 61–68. https://doi.org/10.1016/j.pbi.2020.01.004

Gammerdinger, W. J., & Kocher, T. D. (2018). Unusual Diversity of Sex Chromosomes in
African Cichlid Fishes. *Genes*, *9*(10), 480. https://doi.org/10.3390/genes9100480

Geraldes, A., Hefer, C. A., Capron, A., Kolosova, N., Martinez-Nuñez, F., Soolanayakanahally,
R. Y., Stanton, B., Guy, R. D., Mansfield, S. D., Douglas, C. J., & Cronk, Q. (2015). Recent Y
chromosome divergence despite ancient origin of dioecy in poplars (Populus). *Molecular*
Ecology, *24*(13), 3243–3256. https://doi.org/10.1111/mec.13126

Goodstein, D. M., Shu, S., Howson, R., Neupane, R., Hayes, R. D., Fazo, J., Mitros, T., Dirks,
W., Hellsten, U., Putnam, N., & Rokhsar, D. S. (2012). Phytozome: a comparative platform for
green plant genomics. *Nucleic Acids Research*, *40*(D1), D1178–D1186.
https://doi.org/10.1093/nar/gkr944

Gordon, D., Abajian, C., & Green, P. (1998). Consed: A Graphical Tool for Sequence Finishing.
Genome Research, *8*(3), 195–202. https://doi.org/10.1101/gr.8.3.195

Hill, W., & Weir, B. (1988). Variances and covariances of squared linkage disequilibria in finite
populations. *Theoretical Population Biology*, *33*(1), 54–78. https://doi.org/10.1016/0040-
5809(88)90004-4

Hou, J., Ye, N., Zhang, D., Chen, Y., Fang, L., Dai, X., & Yin, T. (2015). Different autosomes
evolved into sex chromosomes in the sister genera of Salix and Populus. *Scientific Reports*, *5*(1),
9076. https://doi.org/10.1038/srep09076

Kang, H. M., Sul, J. H., Service, S. K., Zaitlen, N. A., Kong, S.-y., Freimer, N. B., Sabatti, C., &
Eskin, E. (2010). Variance component model to account for sample structure in genome-wide
association studies. *Nature Genetics*, *42*(4), 348–354. https://doi.org/10.1038/ng.548

Käfer, J., Marais, G. A. B., & Pannell, J. R. (2017). On the rarity of dioecy in flowering plants.
Molecular Ecology, *26*(5), 1225–1241. https://doi.org/10.1111/mec.14020

Lahn, B. T. (1999). Four Evolutionary Strata on the Human X Chromosome. *Science*, *286*(5441),
964–967. https://doi.org/10.1126/science.286.5441.964

Lemmon, E. M., & Lemmon, A. R. (2013). High-Throughput Genomic Data in Systematics and
Phylogenetics. *Annual Review of Ecology, Evolution, and Systematics*, *44*(1), 99–121.
https://doi.org/10.1146/annurev-ecolsys-110512-135822

Li, H. (2013). *Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM*.
00(00), 1–3. https://doi.org/arXiv:1303.3997

Li, H. (2011). A statistical framework for SNP calling, mutation discovery, association mapping
and population genetical parameter estimation from sequencing data. *Bioinformatics*, *27*(21),
2987–2993. https://doi.org/10.1093/bioinformatics/btr509

Manichaikul, A., Mychaleckyj, J. C., Rich, S. S., Daly, K., Sale, M., & Chen, W.-M. (2010).
Robust relationship inference in genome-wide association studies. *Bioinformatics*, *26*(22), 2867–
2873. https://doi.org/10.1093/bioinformatics/btq559

McKenna, A., Hanna, M., Banks, E., Sivachenko, A. Y., Cibulskis, K., Kernytsky, A. M.,
Garimella, K. V., Altshuler, D., Gabriel, S. B., Daly, M., & DePristo, M. A. (2010). The Genome
Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data.
Genome Research, *20*(9), 1297–1303. https://doi.org/10.1101/gr.107524.110

Miura, I. (2007). An Evolutionary Witness: the Frog *Rana rugosa* Underwent Change of
Heterogametic Sex from XY Male to ZW Female. *Sexual Development*, *1*(6), 323–331.
https://doi.org/10.1159/000111764

Nei, M., & Li, W. H. (1979). Mathematical model for studying genetic variation in terms of
restriction endonucleases. *Proceedings of the National Academy of Sciences*, *76*(10), 5269–5273.
https://doi.org/10.1073/pnas.76.10.5269

Nickerson, D. (1997). PolyPhred: automating the detection and genotyping of single nucleotide
substitutions using fluorescence-based resequencing. *Nucleic Acids Research*, *25*(14), 2745–
2751. https://doi.org/10.1093/nar/25.14.2745

Pakull, B., Groppe, K., Mecucci, F., Gaudet, M., Sabatti, M., & Fladung, M. (2011). Genetic
mapping of linkage group XIX and identification of sex-linked SSR markers in a Populus tremula
× Populus tremuloides cross. *Canadian Journal of Forest Research*, *41*(2), 245–253.
https://doi.org/10.1139/X10-206

Pakull, B., Groppe, K., Meyer, M., Markussen, T., & Fladung, M. (2009). Genetic linkage
mapping in aspen (Populus tremula L. and Populus tremuloides Michx.). *Tree Genetics &*
Genomes, *5*(3), 505–515. https://doi.org/10.1007/s11295-009-0204-2

Paolucci, I., Gaudet, M., Jorge, V., Beritognolo, I., Terzoli, S., Kuzminsky, E., Muleo, R.,
Scarascia Mugnozza, G., & Sabatti, M. (2010). Genetic linkage maps of Populus alba L. and
comparative mapping analysis of sex determination across Populus species. *Tree Genetics &*
Genomes, *6*(6), 863–875. https://doi.org/10.1007/s11295-010-0297-7

Pucholt, P., Rönnberg-Wästljung, A.-C., & Berlin, S. (2015). Single locus sex determination and
female heterogamety in the basket willow (Salix viminalis L.). *Heredity*, *114*(6), 575–583.
https://doi.org/10.1038/hdy.2014.125

Quinlan, A. R., & Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing
genomic features. *Bioinformatics*, *26*(6), 841–842. https://doi.org/10.1093/bioinformatics/btq033

R Core Team. (2018). *R: A Language and Environment for Statistical Computing*. https://www.r-
project.org/

Renner, S. S. (2014). The relative and absolute frequencies of angiosperm sexual systems:
Dioecy, monoecy, gynodioecy, and an updated online database. *American Journal of Botany*,
101(10), 1588–1596. https://doi.org/10.3732/ajb.1400196

Renner, S. S., & Ricklefs, R. E. (1995). Dioecy and Its Correlates in the Flowering Plants.
American Journal of Botany, *82*(5), 596. https://doi.org/10.2307/2445418

Shetty, S., Griffin, D. K., & Graves, J. A. (1999). Comparative painting reveals strong
chromosome homology over 80 million years of bird evolution. *Chromosome Research*, *7*(4),
289–295. https://doi.org/10.1023/1009278914829

Sigeman, H., Ponnikas, S., Chauhan, P., Dierickx, E., Brooke, M. de L., & Hansson, B. (2019).
Repeated sex chromosome evolution in vertebrates supported by expanded avian sex
chromosomes. *Proceedings of the Royal Society B: Biological Sciences*, *286*(1916), 20192051.
https://doi.org/10.1098/rspb.2019.2051

Simonsen, K. L., Churchill, G. A., & Aquadro, C. F. (1995). Properties of statistical tests of
neutrality for DNA polymorphism data. *Genetics*, *141*(1), 413–429.
http://www.ncbi.nlm.nih.gov/pubmed/8536987
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC1206737

Tajima, F. (1989). Statistical method for testing the neutral mutation hypothesis by DNA
polymorphism. *Genetics*, *123*(3), 585–595. http://www.ncbi.nlm.nih.gov/pubmed/2513255
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC1203831

Temmel, N. A., Rai, H. S., & Cronk, Q. (2007). Sequence characterization of the putatively sex-
linked Ssu72 -like locus in willow and its homologue in poplarThis note is one of a selection of
papers published in the Special Issue on Poplar Research in Canada. *Canadian Journal of
Botany*, *85*(11), 1092–1097. https://doi.org/10.1139/B07-058

Tennessen, J. A., Wei, N., Straub, S. C. K., Govindarajulu, R., Liston, A., & Ashman, T.-L.
(2018). Repeated translocation of a gene cassette drives sex-chromosome turnover in
strawberries. *PLOS Biology*, *16*(8), e2006062. https://doi.org/10.1371/journal.pbio.2006062

Tuskan, G. A., DiFazio, S. P., Jansson, S., Bohlmann, J., Grigoriev, I., Hellsten, U., Putnam, N.,
Ralph, S., Rombauts, S., Salamov, A., Schein, J., Sterck, L., Aerts, A., Bhalerao, R. R., Bhalerao,
R. P., Blaudez, D., Boerjan, W., Brun, A., Brunner, A., … Rokhsar, D. S. (2006). The Genome of
Black Cottonwood, Populus trichocarpa (Torr. & Gray). *Science*, *313*(5793), 1596–1604.
https://doi.org/10.1126/science.1128691

Vicoso, B., & Bachtrog, D. (2015). Numerous Transitions of Sex Chromosomes in Diptera.
PLOS Biology, *13*(4), e1002078. https://doi.org/10.1371/journal.pbio.1002078

Wakeley, J. (2009). *Coalescent Theory: An introduction*. Roberts; Company.

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag.

Xue, L., Wu, H., Chen, Y., Li, X., Hou, J., Lu, J., & Charlesworth, D. (2020). Two antagonistic
effect genes mediate separation of sexes in a fully dioecious plant. *bioRxiv*.
https://doi.org/https://doi.org/10.1101/2020.03.15.993022

Zhang, L., Xi, Z., Wang, M., Guo, X., & Ma, T. (2018). Plastome phylogeny and lineage
diversification of Salicaceae with focus on poplars and willows. *Ecology and Evolution*, *8*(16),
7817–7823. https://doi.org/10.1002/ece3.4261

Zhou, R., Macaya-Sanz, D., Carlson, C. H., Schmutz, J., Jenkins, J. W., Kudrna, D., Sharma, A.,
Sandor, L., Shu, S., Barry, K., Tuskan, G. A., Ma, T., Liu, J., Olson, M. S., Smart, L. B., &
DiFazio, S. P. (2020). A willow sex chromosome reveals convergent evolution of complex
palindromic repeats. *Genome Biology*, *21*(1), 38. https://doi.org/10.1186/s13059-020-1952-4

Zhou, R., Macaya-Sanz, D., Rodgers-Melnick, E., Carlson, C. H., Gouker, F. E., Evans, L. M.,
Schmutz, J., Jenkins, J. W., Yan, J., Tuskan, G. A., Smart, L. B., & DiFazio, S. P. (2018).
Characterization of a large sex determination region in Salix purpurea L. (Salicaceae). *Molecular*
Genetics and Genomics, *293*(6), 1437–1452. https://doi.org/10.1007/s00438-018-1473-y

Figure 1. Genome wide associations with sex for 24 male and 24 female *Salix nigra* individuals, expressed in terms of $-\log_{10}$ P-values. The relative positions of each SNP on each of the 19 chromosomes and scaffolds as mapped against the *S. purpurea* genome are shown on the x-axis. Alternating black and grey dots represent different chromosomes. Red dashed line indicates a Bonferroni-adjusted $P_{FDR} = 0.05$ ($-\log_{10} P_{FDR} = 7.91$).

Figure 2. Patterns of nucleotide diversity on *Salix nigra* chromosome 7. A) Dots represent Nei's π for all SNPs in males across chromosome 7. Darkness of the dots represent the relative density of overlapping SNPs plotted in the same are on the figure. The black line represents the rolling mean of 5000 base pair windows. Red horizontal lines indicate the 99% and 1% quantiles of non-overlapping 5000 base pair windows across the genome. B) π for all SNPs in females across chromosome 7.

Figure 3. Patterns of linkage disequilibrium decay on *Salix nigra* chromosome 7. The blue line is the estimated LD-decay calculated in the SLR from males (non-recombining regions of the X & Y chromosomes). The black line is estimated LD-decay from 7-12MB in the recombining region of chromosome 7. The red dashed line is estimated LD-decay calculated in the SLR from females (only X chromosomes).

Figure 4. Patterns of Tajima's D on *Salix nigra* chromosome 7. A) Tajima's D for 25 kb windows for males (both X and Y chromosomes) across chromosome 7. Dots represent the mean Tajima's D for non-overlapping 25kb windows across the chromosome. The black line is the rolling mean of 22 25kb windows, the approximate size of the non-recombining sex linked region(SLR). Red horizontal lines indicate the 99% and 1% quantiles of the Tajima's D calculated from non-overlapping 25kb windows across the genome. B) Tajima's D for 25 kb windows for females (only X chromosomes) across chromosome 7.

Figure S1. Pairwise kinship estimates between males and females of *Salix nigra* in this study. Kinship was estimated using the relatedness2 algorithm in vcftools, which applies the methods used in KING (Manuchaikul et al. 2010) .

A. Males only (XY)

B. Females only (XX)

Figure S2. Patterns of Tajima's D (TajD) across the *Salix nigra* genome. A) Tajima's D for males. Dots represent values of Tajima's D within 25kb across the genome. The blue line is the rolling mean of 22 25kb windows, the approximate size of the non-recombining sex linked region(SLR). Red horizontal lines indicate the 99% and 1% quantiles of the Tajima's D calculated from non-overlapping 25kb windows across the genome. B) Tajima's D for females only.