Modality-independent effect of gravity in shaping the internal representation of 3D space

Morfoisse Theo[1,#], Herrera Altamira Gabriela[1,#], Angelini Leonardo[2,3], Clément Gilles[4], Beraneck Mathieu[1], McIntyre Joseph[1,5,6], Tagliabue Michele[1*]

# These authors contributed equally

Université Paris Cité, CNRS UMR 8002, INCC - Integrative Neuroscience and Cognition Center, F-75006, Paris, France.
HumanTech Institute, University of Applied Sciences Western Switzerland//HES-SO, 1700 Fribourg, Switzerland
School of Management Fribourg, University of Applied Sciences Western Switzerland//HES-SO, 1700 Fribourg, Switzerland
Université de Caen Normandie, Inserm, COMETE U1075, CYCERON, CHU de Caen, Normandie Univ, 14000, Caen, France.
Ikerbasque Science Foundation, 48013 Bilbao, Spain
Health Division, Tecnalia Research & Development, E-20009 San Sebastian, Spain

* **Corresponding author:** Michele Tagliabue; 45 rue des St-Pères; Université Paris Cité, Integrative Neuroscience and Cognition Center, CNRS UMR 8002, F-75270 Paris, France. E-mail: michele.tagliabue@parisdescartes.fr

Acknowledgment: The authors thank M. Patrice Jegouzo for his technical help in designing the experiments. This work was supported by the Centre National des Etudes Spatiales. This study contributes to the IdEx Université de Paris ANR-18-IDEX-0001.

Abstract

Human 3D perception of visual objects is flawed by distortions, which are influenced by non-visual factors, such as gravitational vestibular signals. Whether gravity acts specifically on the visual system or at a higher, modality-independent, level of information processing remains unknown. To test these modality-specific vs modality-independent hypotheses, we performed experiments comparing visual versus haptic 3D shape perception in normo-gravity and microgravity. The results obtained for upright and supine posture in 1g show that visual and haptic perceptual anisotropies are systematically in opposing ego-centered, but not gravity-centered, directions suggesting they share a common origin. On the other hand, microgravity significantly modulates both visual and haptic perceptual distortion in the same direction. Overall, our results show a clear link between the visual and haptic

perceptual distortions and demonstrate a role of gravity-related signals on a modality-independent internal representation of 3D space used to interpret incoming sensory inputs.

1 Introduction

Perception of tridimensional (3D) objects includes the ability to determine an item's location in space, as well as its geometrical properties, such as the relative size along each of three dimensions and the relative orientation of its edges. Given its importance for interacting with the physical world, 3D object perception has been deeply investigated. Visual perception has received the most attention, showing how various features of the stimuli such as disparities, size, occlusions, perspective, motion, shadows, shading, texture and blur all influence 3D visual perception (Welchman 2016) and how *internal models* shape the interpretation of the sensory signals (Curry 1972, Kersten and Yuille 2003, Kersten et al. 2004, Lee 2015).

Despite its critical importance to perception and action, visual perception suffers from measurable distortions: height underestimation with respect to width, also known as the horizontal-vertical, or "L", illusion (Avery and Day 1969) and a systematic depth underestimation (Loomis and Philbeck 1999, Todd and Norman 2003). Non-visual factors, such as gravity, also appear to affect visual perception. For example, tilting the body with respect to gravity affects objects recognition (Leone 1998, Barnett-Cowan et al. 2015), orientation and distance perception (Marendaz et al. 1993, Harris and Mander 2014), and other phenomena such as the tilted frame illusion (Goodenough et al. 1981, Howard 1982), the oblique effect (Lipshits and McIntyre 1999, Luyat and Gentaz 2002, McIntyre and Lipshits 2008) and some geometric illusions (Prinzmetal and Beck 2001, Clément and Eckardt 2005). Furthermore, weightlessness significantly alters the perception of stimulus size and shape, especially in tasks involving depth, during both short-term (Villard et al. 2005, Clément and Bukley 2008, Clément et al. 2008, Harris et al. 2010, Clément and Demel 2012, Clément et al. 2016, Bourrelly et al. 2016) and long-term (Clément et al. 2012-2013, De Saedeleer et al. 2013, Bourrelly et al. 2015) exposure.

One hypothesis to explain gravity-related changes in visual perception is that gravity affects both the eye movements underlying visual exploration (Clément et al. 1986, Reschke et al. 2017-2018) and eye positioning that contribute to the estimation of the visual eye-height, a key reference within the visual scene (Goltz et al. 1997, Bourrelly et al. 2016). Gravity's

influence on oculomotor control should specifically affect visual perception, although

weightlessness might also induce distinct distortions in other sensory modalities. An

alternative hypothesis is that gravity does not affect visual signals *per se,* but rather affects

an internal representation of space (Clément et al. 2009, 2012), based on prior knowledge,

that serves to interpret those signals, independent of the sensory system from which they

come (Wolbers et al. 2011, Loomis et al. 2013). An example, among many, of the use of an

internal model of space for perception is the famous 'Ames room' illusion, where the

persons' size is misperceived due to the use of the inappropriate prior that the room is

rectangular (O'Reilly et al. 2012). A direct implication of this second hypothesis is that mi-

crogravity should distort all spatial perceptions in the same way, regardless of the sensory

modality. Because previous studies in microgravity were focused on visual tasks only,

however, these proposed hypotheses have never been tested.

To investigate these two assumptions, we first compared distortions of visual versus haptic

perception of 3D shape in a normal, upright posture on Earth. Next, we studied the effect

of changing the subject's orientation with respect to gravity to assess whether any visual

or haptic distortions are egocentric or gravity-centric. Third, we tested the consequences

of removing the effects of gravity by performing both haptic and visual experiments in

weightlessness during parabolic flight.

Materials and Methods

In an analogy with previous experiments on visual perception (Clément et al. 2008, 2013),

our paradigm was conceptually designed to detect distortions in the perception of a

purported 3D cube. The sequential nature of haptic perception induced us, however, to

focus each trial on the comparison of the relative size between two out of three possible

dimensions. In both the visual and the haptic cases, the task consisted of adjusting one

side of the rectangle to match the other, to form a square. The adjustments were performed

using a trackball hold in the left hand. Subjects pressed a button on a trackball when they

perceived the object to be perfectly square.

For the haptic tasks, subjects were asked to close their eyes and to feel, through haptic

sense only, a rectangular cutout in a rigid, virtual plank generated by a Force Dimension

Omega.3 haptic robot (Figure 1A). This manipulandum was able to simulate the presence

of a 3D object by applying the appropriate contact forces on the right hand of the subject

when he/she performed exploration movements aimed at perceiving its shape and size.

During each trial the robot constrained the subject's hand movement to lie within the plane

of the virtual plank and to remain inside the rectangle prescribed by the virtual cutout. To

allow direct comparisons between the experimental results from haptic and visual tests, an

analogous bi-dimensional task was also used for visual perception. Subjects were shown

planar rectangles with different orientations in 3D space, without being able to manually

explore it. For trials involving visual perception, an Oculus Rift virtual reality headset was

used to provide a stereoscopic view of the virtual object. The visual environment was dark

and the shapes were represented by light-gray frames. For both sensory conditions, the

virtual object was located approximatively 40 cm in front of the subject's right shoulder.

Although there were no instructions to work quickly, subjects were asked to attempt to

perform each trial in a fixed time window (20 s for all experiments except those performed

on board the parabolic flight plane, for which a 10 s time window was used). An audible

cue indicated to the subject when the end of the allotted time was approaching. The

apparatus recorded the subject's final responses (dimensions of each rectangle judged to

be square), which is the main output of the tests. For the haptic tasks, the movements of

the subject's hand and the contact forces applied against the virtual constraints were also

recorded via the haptic device.

The use of bi-dimensional tasks allowed the estimation of the perceptive distortion in one

plane at a time. Subjects in our experiments judged the squareness of rectangles lying in

each of three anatomical planes: frontal, sagittal, or transversal (see bottom part of Figure

1C). The combination of the three possible planes and the two rectangle dimensions

resulted in six different geometric configurations that the subject had to deal with. They are

represented in the upper part of Figure 2. At the beginning of each trial, an audio command

told the subject in which anatomical plane the rectangle was lying and which of the two

dimensions of the rectangle had to be adjusted. In our paradigm, the reference dimension

was always 40 mm, but subjects were not aware of this fact. The initial length of the

adjustable side was randomly selected between 15, 25, 35, 45, 55, and 65 mm. Subjects

performed five series of trials in all; each series being composed of a random permutation

of the six geometric configurations (total number of trials per condition: 30).

Figure 1: A) Haptic device and virtual reality headset used for the haptic and visual experiments, respectively. In panels B and C are reported the name of the orthogonal directions defined in an external, gravity-centered (Up-Down, UD; East-West, EW; North-South, NS) and egocentric, body-centered (Longitudinal, Lateral, and Anterior-Posterior) reference frames respectively. The bottom part of the figure represents the planes in which the task is performed expressed in the gravity-centered (Horizontal, Meridian and Latitudinal) and body-centered (Transversal, Sagittal and Frontal) reference frames.

Experiment 1: Effect of Sensory Modality

To study the differences and similarities between haptic and visual perception of 3D shapes in normo-gravity, 18 seated subjects (8 males, 10 females, aged 29±9) performed the task for all six geometrical configurations in each of the two sensory conditions: Haptic and Visual. The order of the two sensory conditions was randomized across subjects.

Experiment 2: Effect of Body Orientation

To study the perceptive distortions of both haptic and visual senses and whether the information is encoded in an ego-centric (body-centered) or exo-centric (gravity-centered) reference frame, a group of 18 subjects (9 males and 9 females, aged 25.5±5 years) performed the haptic task while seated (Upright) and while lying on the back (Supine), while a second group of 18 subjects (11 male and 7 female, aged 24±4 years) performed the visual task in the same two postures (Upright and Supine). For the Supine posture, subjects laid on a medical bed. The two postures are represented in Figure 2 together with the respective correspondence between exo-centric and ego-centric references. The virtual object was placed always at the same distance from the subject's shoulder, independent

of the posture. In order to compensate for possible learning effects, the order of the postural conditions was randomized in both sensory conditions.

		Plane	Latitudinal		Horizontal		Meridian	
		Adj. Dim.	East-West	Up-Down	East-West	North-South	Up-Down	North-South
		Ref. Dim.	Up-Down	East-West	North-South	East-West	North-South	Up-Down
Upright		Plane	Frontal		Transversal		Sagittal	
		Adj. Dim.	Lateral	Longitudinal	Lateral	Ant-Post	Longitudinal	Ant-Post
		Ref. Dim.	Longitudinal	Lateral	Ant-Post	Lateral	Ant-Post	Longitudinal
Supine		Plane	Transversal		Frontal		Sagittal	
		Adj. Dim.	Lateral	Ant-Post	Lateral	Longitudinal	Ant-Post	Longitudinal
		Ref. Dim.	Ant-Post	Lateral	Longitudinal	Lateral	Longitudinal	Ant-Post

Figure 2: Geometrical Configurations. The first row represents the six geometric configurations, which correspond to the combination of the three planes in which the rectangle could lie and the two different dimensions of the rectangle that the subject had to adjust. The black bold text corresponds to the definitions, in the gravity-centered reference frame, of the task planes, as well as of the adjustable and reference dimensions of each rectangle. Those definitions are independent of the postural condition of Experiment 2 (Upright and Supine) represented in the lower part of the table. For each combination of geometric and postural conditions, the table reports with gray bold text the anatomical plane in which the task is performed as well as the anatomical direction of the adjustable and reference dimensions of the rectangles.

Experiment 3: Effect of Weightlessness

To study the role of gravitational cues in the encoding of haptic and visual signals we performed the haptic (18 subjects: 10 males, 8 females, aged 38±11 years) and visual (18 subjects: 9 males, 9 females, aged 41±11 years) paradigm in normal gravity (1G) and during the weightlessness phases of parabolic flight (0G). For the haptic experiment, a third condition was added: the subjects were also tested in normal gravity, but with the arm supported by a strap (Supp.), to differentiate the biomechanical effect of gravity on the arm from the gravitational stimulation of graviceptors, such as the otoliths.

Parabolic flight provides short intervals (~20s) of weightlessness within a stable visual

environment inside the airplane, bracketed by periods of hyper-gravity (1.6 - 1.8 G) just

before and just after each period of weightlessness. Given the short duration of 0G phases

during parabolic flight, the subjects were trained to perform the task in about 10 seconds

(two tasks per parabola). Since each subject performed the experiment during 15

consecutive parabolas, he or she could perform all 30 trials per condition.

All experimental conditions were performed inflight onboard the Novespace Zero-G

airplane in order to minimize possible undesired changes in uncontrolled factors. The 1G

and Support conditions were tested during the level-flight phase just preceding the first

parabola or just following the last parabola of its session, depending on the subject. The

subjects were very firmly restrained with belts so that their relative position with respect to

the apparatus and the virtual rectangles did not vary between gravitational conditions.

Data analysis

For each trial, t, the error, ε, between the length of the adjustable and reference sides of

the rectangle was computed. If the exo-centered definition of the three dimensions (*EW*,

NS, and *UD*) of Figure 1B is used, the errors of the six geometric configurations are defined

as *EW-UD*, *UD-EW*, *EW-NS*, *NS-EW*, *UD-NS,* and *NS-UD*, where the minuend and the

subtrahend are the adjustable and reference dimensions respectively.

Table 1: Definition of the squaring errors for all six
geometrical configurations of the task.

Plane	Adjustable dimension	Reference dimension	Task error
Latitudinal	EW	UD	$\varepsilon_{EW-UD} = \varepsilon_{EW} - \varepsilon_{UD}$
	UD	EW	$\varepsilon_{UD-EW} = \varepsilon_{UD} - \varepsilon_{EW}$
Horizontal	EW	NS	$\varepsilon_{EW-NS} = \varepsilon_{EW} - \varepsilon_{NS}$
	NS	EW	$\varepsilon_{NS-EW} = \varepsilon_{NS} - \varepsilon_{EW}$
Meridian	UD	NS	$\varepsilon_{UD-NS} = \varepsilon_{UD} - \varepsilon_{NS}$
	NS	UD	$\varepsilon_{NS-UD} = \varepsilon_{NS} - \varepsilon_{UD}$

Table *1* shows how the perceptive distortion associated with each of the three dimensions

contributes to the error made on the six geometric configurations. Positive errors

correspond to underestimations of the adjustable dimension and/or to overestimations of

the reference dimension. Thus, the present experimental paradigm, similar to the one
previously used by Clément et al. (2008, 2013), allows the quantification of the perceptive
distortions of one dimension relative to another, but cannot lead to a measure of the
absolute perceptive distortions for each dimension separately. Nevertheless, we have
devised a methodology that allows one to visualize the 3D patterns of distorsion as an
elongated box compared to an ideal cube (see Appendix I).

Estimation of 3D perceptual distortion

Table *1* shows that the error in estimating one dimension has opposite effects for the two
tasks performed within a given plane. For instance, an overestimation of the NS dimension
should result in negative and positive errors in the NS-EW and EW-NS tasks, respectively.
These relationships appear to be confirmed by the experimental results (Figure 4a),
because this hypothesis accounts for 96% of the data variance. It follows that the theo-
retical relationships below are valid:

$$\varepsilon_{EW\text{-}NS} = - \varepsilon_{NS\text{-}EW}$$

$$\varepsilon_{EW\text{-}UD} = - \varepsilon_{UD\text{-}EW} \tag{1}$$

$$\varepsilon_{UD\text{-}NS} = - \varepsilon_{NS\text{-}UD}$$

Exploiting this property, it was possible to combine the five errors obtained for one
geometric condition, with the additive inverse of the five errors obtained for the other
geometric condition performed in the same plane. This allowed computing the combined
mean and the variance of the errors for each of the three planes (Horizontal, Latitudinal,
and Meridian), instead of individually for each of the 6 geometrical configurations of the
task. This technique has the considerable advantage of being more robust, because it is
based on 10 samples instead of only 5.

$$\overline{\varepsilon}_{Hor} = \frac{\sum_{t=1}^{5}\left(\varepsilon_{EW-NS,t} - \varepsilon_{NS-EW,t}\right)}{10}$$

$$\sigma_{Hor}^2 = \frac{\sum_{t=1}^{5}\left(\left(\varepsilon_{EW-NS,t} - \overline{\varepsilon}_{Hor}\right)^2 + \left(-\varepsilon_{NS-EW,t} - \overline{\varepsilon}_{Hor}\right)^2\right)}{10} \tag{2}$$

$$\overline{\varepsilon}_{Lat} = \frac{\sum_{t=1}^{5}\left(\varepsilon_{EW-UD,t} - \varepsilon_{UD-EW,t}\right)}{10}$$

$$\sigma_{Lat}^2 = \frac{\sum_{t=1}^{5}\left(\left(\varepsilon_{EW-UD,t} - \overline{\varepsilon}_{Lat}\right)^2 + \left(-\varepsilon_{UD-EW,t} - \overline{\varepsilon}_{Lat}\right)^2\right)}{10}$$

$$\overline{\varepsilon}_{Mer} = \frac{\sum_{t=1}^{5}\left(\varepsilon_{NS-UD,t} - \varepsilon_{UD-NS,t}\right)}{10}$$

$$\sigma_{Mer}^2 = \frac{\sum_{t=1}^{5}\left(\left(\varepsilon_{NS-UD,t} - \overline{\varepsilon}_{Mer}\right)^2 + \left(-\varepsilon_{UD-NS,t} - \overline{\varepsilon}_{Mer}\right)^2\right)}{10}$$

With the above formulas, one can characterize perceptual distortions in each of the three

different planes as illustrated in Figure 3. By our convention, a rectangle lying in one of the

two vertical planes (Meridian or Latitudinal) is associated with a positive error (stubby

rectangle) if the vertical dimension is smaller than the other dimension. In the horizontal

plane, a positive error (stubby rectangle) corresponds to the depth (NS dimension) being

smaller than the width (EW dimension). It is worth noting that if the subject produced a

"stubby" rectangle (positive errors) this means that he/she perceived a square to be

"slender", and vice versa. The global variance was computed as the average of the three

planar variances.

Figure 3: Sign conventions for the errors in the Horizontal, Latitudinal and Meridian planes. The gray squares represent the correct answer (i.e. a square). The black lines represent the distorted answers. Positive planar error values correspond to "stubby" rectangles. Negative values correspond to "slender" rectangles. The same conventions are used for the error expressed in the body-centered planes. In this case, Anterior-Posterior, Lateral and Longitudinal directions replace NS, EW and UD, respectively. Transversal, Frontal and Sagittal replace Horizontal, Latitudinal and Meridian planes, respectively.

170 The estimation of the three planar errors is then improved by considering that if the
171 (distorted) metrics used to compare distances in 3D space are locally smooth and
172 consistent for the different dimensions in space, the three planar errors ε are not
173 independent and that, given the sign conventions of Figure 3, they should fulfill the following
174 relationship

$$\bar{\varepsilon}_{Mer} + \bar{\varepsilon}_{Hor} = \bar{\varepsilon}_{Lat} \tag{3}$$

Note that equation 3 is a particular case of the formula describing a plane, $ax + by + cz =$
d, where $a = b = 1$, $c = -1$ and $d = 0$. Thus, if the metrics in each plane are consistent with
each other, the vectors of measured planar errors $\bar{\varepsilon} = [\bar{\varepsilon}_{Mer}\ \bar{\varepsilon}_{Hor}\ \bar{\varepsilon}_{Lat}]$ should fall on that
plane and points outside the plane can be considered to be noise. By projecting the
individual vectors $\bar{\varepsilon}$ onto the plane corresponding to equation 3, this noise is effectively
filtered out. Using the resulting 2D representation of the distortion is a conservative choice,
especially when comparing their orientation in different conditions, because the 3D
representation may lead to consider distortion directions and components of data
variability that have no functional meaning. On average, the data projected on the plane of
equation 3 account for 98% of the variance of the original data, suggesting that the
recorded responses tend to well fulfill this constraint.

We used the same equations (1-3) to compute the analogous parameter in the ego-centric
reference frame after having replaced the externally defined planes and directions with the
body-centered planes (*Transversal*, *Frontal,* and *Sagittal*) and directions (*Lateral*, *Anterior-*
Posterior, and *Longitudinal*) as shown in Figure 2. Table *2* shows the relationships between
the planar distortions defined in the body-centered and gravity-centered reference frame
for the Upright and Supine posture.

Table 2: Relationship between ego- and exo-
centrically defined distortions for the Upright and
Supine condition.

Upright	$\bar{\varepsilon}_{Mer} = \bar{\varepsilon}_{Sag}$	$\bar{\varepsilon}_{Lat} = \bar{\varepsilon}_{Fro}$	$\bar{\varepsilon}_{Hor} = \bar{\varepsilon}_{Tra}$
Supine	$\bar{\varepsilon}_{Mer} = -\bar{\varepsilon}_{Sag}$	$\bar{\varepsilon}_{Lat} = \bar{\varepsilon}_{Tra}$	$\bar{\varepsilon}_{Hor} = \bar{\varepsilon}_{Fro}$

Distortion direction

The 2D vector resulting from the projection of $\bar{\varepsilon}$ to the plane of equation 3 was computed for each subject. To estimate the mean direction of the distortion in each experimental condition, the vectorial average of the 2D vectors was computed.

Reaction forces during haptic task

To estimate changes of the contact forces between gravitational conditions in the haptic tasks, we computed the average of the reaction forces generated by the haptic device when the subject's hand was in contact with the edges of the virtual cutout or when the hand tried to move out of the task plane.

Microgravity effect and modeling

To quantify the effect of microgravity on the perceptive distortions, for each subject the mean planar error in 1G was subtracted from the corresponding error in 0G:

$$\Delta\bar{\varepsilon} = \bar{\varepsilon}_{0G} - \bar{\varepsilon}_{1G}$$

computed separately for the visual and the haptic modalities. As for the mean planar error, a 2D vector representation was computed projecting each individual $\Delta\bar{\varepsilon}$ onto the plane of equation 3. To predict the perceptive distortion expected in microgravity under the hypothesis that the 0G effect was identical for the haptic and visual modalities, we averaged all 2D vectors representing the measured individual microgravity effects from both the haptic and visual experiments, combined. The obtained vector was then added to the individual visual and haptic distortions measured in normo-gravity conditions. We then compared these predictions to the distortions measured in 0G for the visual and haptic modalities.

Statistical analysis

For each experiment, we first tested the significance of the distortions by testing for each plane whether the constant errors were on average different from zero (two-sided Student's t-test). Then, we performed repeated-measures ANOVA on the constant and variable errors. The sign conventions (Figure 3) being arbitrary, they allow a rigorous comparison of the perceptive distortion within a given plane, but they do not allow the comparison

between different planes. For this reason, in the statistical analyses, the results on each
plane were tested with independent ANOVAs.

<u>Experiment 1:</u> For each of the 3 task planes we tested for an effect of Sensory Modality on
the perceptive distortion as a single within-subject independent factor with two levels
(Haptic, Visual).

<u>Experiment 2:</u> We tested for an effect of Body Posture as a within-subject independent
factor with two levels (*Upright*, *Supine*) in separate ANOVAs for each group/sensory
modality (Visual and Haptic). Note that this separation is justified by the hypotheses being
tested, for which cross effects between posture and modality would have little meaning. To
test whether distortions are tied to an ego-centric or gravity-centric reference frame, we
defined the task planes both in terms of anatomical axes and world axes. Invariance of
distortion (lack of a statistical difference) for the anatomically defined planes, but not the
world-defined frames, would indicate that the distortions are primarily egocentrically rather
than exocentrically aligned and vice-versa.

<u>Experiment 3:</u> For each of the 3 task planes we tested for an effect of Gravity on shape
perception as a single within-subject independent factor with three (1G, 0G, Supported)
and two (1G, 0G) levels for the haptic and visual experiment respectively.

Before performing each ANOVA, we tested for normality and homogeneity of the
distributions using the Kolmogorov-Smirnov and Levenes tests, respectively. To achieve
the normal distribution for the response variability, the standard deviations were
transformed by the $\log(\sigma+1)$ function (Tagliabue and McIntyre 2011). For the errors
expressed in both exocentric and ego-centric reference frames the data were distributed
normally (all $p>0.20$) and the data variability was similar among all conditions (all $p>0.50$).

To test whether the direction of the distortion (2D vectorial average) differs between two
conditions, a bootstrap technique was used: 10000 re-samplings were used to estimate
the statistical distribution of the angle θ between the vectorial average of two conditions,
and to compute the probability of error in rejecting the null hypothesis, H_0, that $\theta=0$. The
same technique was used to compare the global amplitude of the distortions.

Following a Bayesian approach, taking into account a prior uniform distribution of all possible angles (θ range $\pm180°$), we evaluated the ratio, $R_{0/1}$, between the probability to obtain the observed data under the null hypothesis, H_0, and the probability under the alternative hypothesis, H_1, that $\theta\neq0$ (Wagenmakers et al. 2018).

Results

Experiment 1: Haptic and Visual Perception

Figure 4A shows that for the six geometric configurations of the squaring task (see methods) the subjects made systematic errors in both visual and haptic conditions. The comparison of the errors made using haptic information alone versus visual information alone shows consistent, opposing results for the two sensory modalities. Hence, in each task, when subjects made on average significant positive errors in the haptic condition, they made negative errors in the visual condition, and vice-versa. Figure 4B represents the more robust evaluation of the planar distortions obtained by considering the constraints existing between the errors performed in the six squaring tasks (see methods equation 1-3). The amplitude of the distortion was significantly different from zero for both visual and haptic perception in the Sagittal (visual: $F_{(17)}=5.86$, $p<10^{-4}$, haptic: $F_{(17)}=-8.10$, $p<10^{-6}$) and Transversal plane (visual: $F_{(17)}=-7.22$, $p<10^{-5}$, haptic: $F_{(17)}=9.22$, $p<10^{-6}$), but in the Frontal plane none of the modality was significantly different from zero (visual: $F_{(17)}=-1.26$, $p=0.22$, haptic $F_{(17)}=-0.57$, $p=0.58$). Sensory modality had a significant effect in the Sagittal ($F_{(1,17)}=60.8$, $p<10^{-5}$) and Transversal ($F_{(1,17)}=94.96$, $p<10^{-6}$) planes, but not in the Frontal plane ($F_{(1,17)}=0.14$, $p=0.71$). Remarkably, the significant perceptive distortions in the Sagittal and Transversal planes were in opposite directions between the two sensory conditions. When using haptic sense, subjects produced rectangles with the Anterior-Posterior dimension smaller than the Longitudinal and Lateral dimension, while, when using vision, they made rectangles with the Anterior-Posterior dimension larger than the Longitudinal and Lateral dimension.

The vectorial representation of the individual errors in Figure 4C shows that the perceptive distortions for the two sensory modalities appear along the same axis, but in opposite directions: average angle θ between the visual and haptic distortion is $174\pm5°$ and not significantly different from $180°$ (bootstrap $p=0.13$). Taking into account all possible

orientations for the two groups of vectors, the observed results are 14 times more likely

under the hypothesis that distortions of the two senses are in opposite directions (H_0: θ

=180˚), than under the alternative hypothesis (H_1), i.e. $\theta \neq 180°$. The amplitude of the

distortion is larger for visual than for haptic sense (bootstrap p=0.02). The illustration of the

perceptive distortion corresponding to the two sensory modalities is reported in Figure 4D

and suggests that the haptic and visual perceptive distortions would mainly consist of a

depth overestimation and underestimation for the haptic and visual sense, respectively.

The method used to compute the parallelepiped dimensions is described, with its

limitations, in Appendix 1.

If the amplitude of the perceptive biases (constant components of the errors reported in

Figure 4 A-C) appear smaller for the haptic than for the visual modality, the latter is

characterized by a clearly smaller intra-personal variability of the responses (σ_{hapt}=6.1±2.6

mm, σ_{vis}=4.2±2.2 mm, sensory modality effect: $F_{(1,17)}$=12.02, $p<10^{-2}$), corresponding to a

higher precision for the visual than for the haptic task.

Figure 4: **A**) Errors for the task performed in each of the six geometrical conditions using haptic information only (light blue bars) or visual information only (red bars). Each geometrical condition is characterized by the plane in which the rectangle lies (sagittal, transversal, frontal), and by which direction within the plane was adjustable or held constant: Longitudinal (Lo), Anterior-Posterior (AP), and Lateral (La). Positive errors correspond to the final size of the adjustable dimension being greater than the reference dimension. Vertical wiskers represent 95% confidence intervals. A significant difference between the two tasks performed in the same plane is indicative of an important perceptive distortion in that specific plane. **B**) Distortions in the three task planes for haptic and visual conditions. *** : $p < 10^{-3}$ in the ANOVA testing the modality effect. ‡ : $p < 10^{-3}$ for the t-test ascertaining differences from zero. **C**) Vectorial representation of individual responses in the Sagitt+Transv=Frontal errors space. Thicker vectors represent the average vectorial response. Dashed line indicates the Sagitt+Transv=0 axis. **D**) Illustration of how a cube (gray shape) would be perceived by the subjects when using haptic or visual information alone, respectively. For illustration purposes, the distortions of this panel are scaled up by a factor of 5.

295 In summary, Experiment 1 shows clear differences in the patterns of visual and haptic

296 distortions. Distortion appeared primarily in the sagittal and transversal planes and were in

297 opposite direction for the two sensory modalities (contraction and expansion of perceived

298 depth for visual and haptic, respectively).

Experiment 2: Effect of Body Orientation

The responses of the subjects upright were characterized by constant errors similar to

those observed in Experiment 1 (Experiment effect: Wilks' Lambda=0.85, $F_{(4,32)}$=1.35,

p=0.27). The left columns of Table 3 and left panels of Figure 5 show that for both haptic

and visual experiments the planar distortions appear consistent between postures if

expressed ego-centrically: we observed no statistically significant effects of posture on the

errors for any of the three planes when expressed in body-centered reference frame, nor

on the distortion magnitude (boostrap for haptics p=0.76 and vision p=0.75) and direction

(boostrap for haptics p=0.52 and vision p=0.36; $R_{0/1}$=10.1 for haptics and $R_{0/1}$=21.8 for

vision). On the other hand, if the errors are represented in terms of exo-centrically defined

planes, i.e. fixed with respect to gravity (last three columns of Table 3 and right panels of

Figure 5), a clear effect of posture can be observed in all planes for both sensory modalities

and on the distortion orientation (bootstrap $p<10^{-4}$ for haptics and vision; $R_{0/1}<10^{-4}$ for

haptics and vision).

Figure 5: Errors within each plane when the subjects are seated normally (Upright) or lying Supine. The upper (**A**) and lower (**B**) panels represent the results for the Haptic and Visual modalities, respectively. The left panels represent the errors per anatomical, ego-centered plane. The right panels represent the data per exo-centered (fixed with respect to gravity) plane. ** : $p<10^{-2}$ and *** : $p<10^{-3}$ in the ANOVA. † and ‡ : $p<10^{-2}$ and $p<10^{-3}$ for the t-test ascertaining differences from zero. Vertical wiskers represent 95% confidence intervals. In each barplot the inset illustrates the corresponding 3D perceptive distortion (amplified x5) of a cube. The circular vectorial plots represent individual responses in the Sagitt+Transv=Frontal errors space. Thicker vectors represent the average vectorial response. Dashed diameters indicate the Sagitt+Transv=0 (or the Merid+Horiz=0) axis. Dotted diameters indicate the Saggitt=Frontal (or Merid=Latitud) axis, corresponding to a 90° rotation of the distortions pattern.

313
314

Table 3: Results of ANOVA for the posture effect on the planar perceptive distortion.

	Sagittal	Transversal	Frontal	Meridian	Horizontal	Latitudinal
Haptic	$F_{(1,17)}=0.40$ $p=0.53$	$F_{(1,17)}=0.58$ $p=0.46$	$F_{(1,17)}=0.001$ $p=0.97$	$F_{(1,17)}=52.28$ $p<10\text{-}5$	$F_{(1,17)}=13.01$ $p=0.002$	$F_{(1,17)}=12.18$ $p=0.003$
Visual	$F_{(1,17)}=2.00$ $p=0.18$	$F_{(1,17)}=1.32$ $p=0.27$	$F_{(1,17)}=0.15$ $p=0.70$	$F_{(1,17)}=25.46$ $p<10\text{-}3$	$F_{(1,17)}=19.92$ $p<10\text{-}3$	$F_{(1,17)}=22.87$ $p<10\text{-}3$

The intra-personal variability of the responses was not affected by the posture for the haptic modality ($\sigma_{upright}=6.2\pm6.1$ mm, $\sigma_{supine}=6.6\pm6.0$ mm, posture effect: $F_{(1,17)}=0.12$, $p=0.73$), but significantly increased in the supine position for the visual experiment ($\sigma_{upright}=3.5\pm3.2$ mm, $\sigma_{supine}=4.8\pm4.7$ mm, posture effect: $F_{(1,17)}=6.81$, $p=0.02$).

In conclusion, in this experiment we found that patterns of distortion of both visual and haptic perception were invariant when expressed in an egocentric reference frame, but not when expressed in an exocentric one.

Experiment 3: Gravity's Effect on Visual and Haptic Perception

While the visual inputs are not different on ground and in weightlessness, the forces exerted against the virtual constraints during haptic exploration might be different in 0g due to biomechanical and neurophysiological effects. We therefore first analyze the changes in the contact forces between the subject's hand and the virtual object and then the pattern of perceptive distortions (Figure 6A-C). The left plot of Figure 6A shows that vertical forces applied by the subjects on the upper and lower edge of the sensed object were modulated ($F_{(2,34)}=3.9$, $p=0.02$) by the experimental conditions (1G, 0G, Supported). As expected, upward and downward forces increased and decreased respectively in microgravity (post-hoc 1G Vs 0G, $p=0.02$), coherent with a reduction of the weight of the upper limb. When the weight of the arm was supported (see methods), the vertical forces also tended to differ from 1G condition (post-hoc Supp Vs 1G $p=0.09$) and were modulated in the same direction as in 0G (post-hoc Supp Vs 0G, $p=0.29$). Horizontal forces were also significantly affected by the experimental condition ($F_{(2,34)}=6.32$, $p<0.01$; Figure 6A, right plot), with a significant increase of the contact forces in microgravity with respect to the 1G and Support conditions.

This increase of the contact force in 0G, similar to what was previously observed in haptic tests during parabolic flights (Mierau et al. 2008), could be the result of a specific strategy

aimed at keeping muscular tension, and hence muscle spindle sensitivity, similar to normal

gravity conditions. This strategy would avoid the decrease in proprioception precision

previously observed in weightlessness for 'open-chain' motor tasks, for which the same

strategy could not be adopted, resulting in a decrease in muscle tension (Clément and

Reschke 2010). This hypothesis well matches the fact that the precision of haptic

responses was not significantly affected by the experimental condition (response variability:

1G 6.8±2.6, 0G 7.1±3.1, Sup 6.4±2.9; $F_{(2,34)}$=1.75, p=0.19), suggesting that neither

microgravity nor the arm support significantly interfered with the subjects' ability to perform

the task. This lack of microgravity effect on haptic precision appears in line with the results

of previous orbital experiments (McIntyre and Lipshits 2008).

Importantly, the results about the vertical contact forces and responses' variability suggest

that the 'arm support' condition successfully mimicked the expected lightening of the arm

observed in microgravity. Therefore, if haptic perceptive distortions (constant errors) are

affected by microgravity, but not by the arm support, they would not be directly ascribable

to the biomechanical action of microgravity on the arm.

Figure 6: Results of the microgravity experiments for the haptic (A-C and F panels) and visual (D-E and G panels) tasks. A) Contact forces in the three experimental conditions: normogravity (1G), microgravity (0G) and with a mechanical support of the arm (Supp). Left: Vertical forces generated against the upper and lower edges of the rectangle. Right: Horizontal forces generated against all other edges of the rectangle. B) and D) Errors in the three task planes. C) and E) are circular vectorial plots representing individual responses in the Sagitt+Transv=Frontal errors space. Thicker vectors represent the average vectorial response. Dashed diameters indicate the Sagitt+Transv=0 axis. Within-subject variability of the responses when considering all three planes together. F-G) Illustration of the perceptive-distortion (experimental results scaled up by 5) of reference cube (gray) in normal gravity and in microgravity. * : p<0.05 and ** : p<10^{-2} in the ANOVA. ⱡ, † and ‡ : p<0.05, p<10^{-2} and p<10^{-3} for the t-test ascertaining difference from zero.

The comparison of the constant errors in the three experimental conditions, reported in

Figure 6B, clearly shows that the perceptive distortion characterizing haptic perception in

the Sagittal plane was significantly amplified (became more negative) by microgravity, but

was not affected by the arm support (condition effect $F_{(2,34)}$=12.49, p<10^{-4}), suggesting a

perceptive rather than biomechanical effect. Similarly, the haptic distortion in the

Transversal plane was amplified (became more positive) in 0G and was not affected by the

support, either (condition effect $F_{(2,34)}$=11.13, p=<10^{-3}). Finally, the lack of distortion in the

Frontal plane persisted independent of the gravitational and support condition ($F_{(2,34)}$=0.33,

p=0.71). Figure 6C shows a clear increase of the global haptic distortion amplitude in 0G

(bootstrap $p<10^{-4}$) and very similar direction between the haptic distortions observed in the

two gravitational conditions (bootstrap on the angle θ between the two vectors: p=0.64;

$R_{0/1}$=7.2).

For the visual tasks, Figure 6D shows that, as for the haptic sense, microgravity significantly

modulated the perceptive distortions. More precisely, the large errors characterizing both

sagittal and transversal planes in 1G were significantly reduced in weightlessness

($F_{(1,17)}$=15.41, p=0.0011 and $F_{(1,17)}$=7.87, p=0.012 respectively). In the frontal plane, a small

but significant height underestimation appeared in 0G ($F_{(1,17)}$=9.531, p=0.007). Figure 6E

shows that the global visual distortion amplitude tends to decrease in microgravity, but not

significantly (bootstrap p=0.15) and thatthe angle between the distortions in the two gravity

conditions (6.7±5°) is not significantly larger than zero (bootstrap p=0.08): the probability of

observing these data is eleven times larger under the hypothesis that the two vectors'

orientation representing the direction of the distortion are identical compared to the

alternative hypothesis that they are different ($R_{0/1}$=11.2). The analysis of the variable

component of the errors shows that microgravity did not significantly affect subjects' visual

precision ($F_{(1,17)}$=4.3, p=0.054), although the response variability tended to increase from

4.4±2.5 to 5.2±2.4 mm.

In neither haptic nor visual 0G tasks the amplitude of the distortions appears to change

over the parabolas (trial number effect on haptic errors: Sagittal $F_{(4,60)}$=0.79, p=0.54;

Transversal $F_{(4,60)}$=0.23, p=0.92; Frontal $F_{(4,60)}$=0.49, p=0.74; and on visual errors Sagittal

$F_{(4,68)}$=1.23, p=0.30; Transversal $F_{(4,68)}$=0.60, p=0.67; Frontal $F_{(4,68)}$=0.63, p=0.64) suggesting

a lack of significant adaptation to microgravity during the experiment duration.

The qualitative comparison of Figure 6F and Figure 6G shows that the effect of gravity on

both sensory modalities mainly consists of a stretch of depth perception with respect to

normo-gravity conditions (an increase in slenderness for haptic; a decrease in stubbiness

for visual). The direct quantitative comparison of the effect of microgravity between the two

groups of subjects of the visual and haptic experiments (Figure 7A) shows similar

modulations of the perceptual distortion for both senses (Wilks' Lambda=0.91, $F_{(3,32)}$=0.96,

p=0.42). Although the amplitude of the microgravity effect tends to be larger for haptic than

for visual perception (bootstrap, p=0.06), the average directions of the microgravity effect

on visual and haptic sense appear very similar (Figure 7B): the angle θ between the two

vectors representing the average effect of gravity on the two modalities is only 15.6±15.6°

and not significantly larger than zero (bootstrap, p=0.14). When considering the range of all

possible θ (±180°), Bayesian statistics suggest that the observed data are 5.2 times more

likely under the hypothesis that θ=0° (H$_0$) than under the hypothesis θ≠0° (H$_1$). As shown in

Figure 7C, assuming that the gravity effect is identical for the haptic and visual modality

(both in terms of direction and amplitude) predicts indeed distortions in microgravity

indistinguishable from the observed results (Wilks' Lambda=0.82, $F_{(4,14)}$=0.77, p=0.56).

Figure 7: Comparison of the effect of microgravity on the Haptic and Visual senses. A) Difference between the constant errors made by the subjects in the 0G and 1G conditions for the tasks in the three anatomical planes. B) Vectorial representation of the gravity effect in the *Sag+Trans=Front* errors space. Thicker vectors represent the average response. Dashed line indicates the *Sag=-Trans* and *Sag=Trans* axes. C) Errors observed in 0G and predicted from 1G data assuming an identical effect of microgravity on both haptic and visual perception. The values correspond to the error projection on the *Sag=-Tr* and *Sag=Tr* axes of panel B. Vertical wiskers represent 95% confidence interval.

To summarize, the parabolic flight experiments show that, although very different

perceptive distortions characterize vision and haptic sense in normal gravity conditions, the

effects of microgravity on each of those distortions are in the same direction for the two

sensory modalities.

Results Summary

Experiment 1 revealed strong and opposite distortions between haptic and visual

perception of 3D geometry. Subjects visually underestimated an object's depth with

respect to both height and width, whilst overestimating depth when exploring the object haptically. In Experiment 2 the comparison of seated versus supine body orientation clearly showed that both visual and haptic distortions align with the subject's body rather than with gravity. Experiment 3, conducted during parabolic flight, showed a clear effect of microgravity on both haptic and visual distortion while preserving their respective 'terrestrial' direction. Importantly, the incremental effect of microgravity was along the same axis and in the same direction for visual and haptic tasks, despite the fact that the perceptive errors in normo-gravity were in diametrically opposite directions: weightlessness *increases* the haptic *over*-estimation of depth with respect to width and height and *decreases* the visual *under*-estimation of depth with respect to width and height.

Discussion

The experiments presented here aimed to understand how gravity affects the perception of 3D shapes. We extend previous studies restricted to vision to include haptic sensation, by using the same experimental paradigm for the two modalities. Hereafter, we discuss our empirical results in normal gravity and in microgravity with respect to the literature and how they support the idea of a modality-independent role of gravity in interpreting incoming sensory signals.

Haptic and Visual perception in normo-gravity conditions

Individually, the visual and haptic distortions observed here are consistent with previous findings obtained without using head-mounted displays or haptic devices, supporting the validity of the present experimental paradigms. Our haptic results concur with overestimation in the radial dimension observed for haptic tasks performed in the horizontal plane (Lipshits et al. 1994, Armstrong and Marks 1999, Fasse et al. 2000, Henriques and Soetching 2003). Similarly, the reported visual underestimation of depth corresponds to what has been previously observed for this modality in the horizontal plane (Wagner 1985, Loomis and Philbeck 1999). Surprisingly, we observed no significant distortion in the frontal plane, as one might have expected given the well-known horizontal-vertical, or "L", illusion (Avery and Day 1969). The object being situated in front of the right shoulder, rather than straight ahead, might explain this discrepancy: in the latter position, vertical lines would more likely be associated with depth (Girgus and Coren 1975), than in the former, where horizontal lines may also contribute to depth perception.

Collectively, these results represent, to our knowledge, the first direct quantitative

comparison of visual and haptic perception of 3D shapes. Although perceptual biases were

known to differ between visually and haptically guided pointing (vanBeers et al 1999, Liu et

al. 2018), no previous experiment has shown, as we do here, that the perceptual distortions

are exactly in the opposite direction for the two senses.

Moreover, our second set of experiments on body orientation shows that visual and haptic

distortions remain aligned to the subject anterior-posterior axis and opposite to each other

when supine. Although in apparent contradiction with the effect of body tilt shown for a

variety of visual tasks (Marendaz et al. 1993, Leone 1998, Barnett-Cowan et al. 2015), and

with the effect of the direction of external forces in haptic tasks (Wydoodt et al. 2006), the

present invariance of the ego-centric perceptive distortions is consistent with previously

reported body-centered haptic perception (Gurfinkel et al. 1993, Dupin et al. 2018) and eye-

centered encoding of shape/position of visual object (Averly and Day 1969, Howard et al.

1990, McIntyre et al. 1997, Henriques et al. 1998, Vetter et al. 1999). Furthermore, the

present lack of effects of the body orientation on the egocentric visual and haptic 3D

shapes perception is consistent with the previously reported absence of effects of tilting

the body on unimodal visual and proprioceptive tasks, as opposed to cross-modal tasks

(Bernard-Espina et al. 2022).

Our empirical observations of invariant, diametrically opposite visual and haptic egocentric

distortions represent a novel and intriguing finding. It suggests that, in addition to causes

associated with modality-specific mechanisms, such as eye vergence signals integration

for vision (Murdison et al. 2019) and kinematics of the exploratory movements for haptic

(Armstrong and Marks 1999, McFarland and Soechting 2007), the distortions characterizing

the two modalities might have a common origin, probably linked to the continuous

interaction between the information originating from the two sensory systems during

everyday object manipulation: haptic distortions might compensate the perceptive visual

bias, and viceversa, but this specific hypothesis remains to be tested.

The role of gravity in 3D object perception

Although the ego-centric patterns observed for visual and haptic distortions would suggest

that an external cue, such as gravity, should not influence shape perception, the strong

microgravity effects observed in our parabolic-flights experiments clearly show that the

contrary is, in fact, true. How can these apparently contradictory results be reconciled? We

have shown that the observed effects of microgravity on both haptic and visual perceptive

distortions are not directly ascribable to a decrease in their precision, or, specifically for the

haptic task, to the mechanical action of gravity on the arm (arm support condition and

supine haptic experiment). Moreover, the remarkable similarity between microgravity's

effects on visual and haptic distortions makes it unlikely that they might be caused by

independent effects of gravity on the two sensory systems, as has been proposed for the

results of previous microgravity experiments: modification of proprioceptive-tactile

receptors' output for haptic tasks (Lipshits et al. 1994) and deviation of the vertical position

of the eye for visual tests (Clément et al. 2008, 2013, Bourrelly et al. 2015, 2016). A more

parsimonious and likely explanation is that microgravity's effect on the two senses has a

common origin. This modality-independent phenomenon could be due to an effect of

gravity on an internal representation of the egocentric 3D space used to interpret the

object-related inputs, independent of the sensory system from which they originate. Clear

evidence that internal models of space affect the interpretation of incoming sensory

information in a Bayesian fashion has been extensively reported. Good examples include

the 'Ames room' and the Müller-Lyer visual illusions based on prior knowledge about the

geometry of constructed environments (O'Reilly et al. 2012) or the cutaneous Rabbit illusion

(Goldreich et al. 2007). The idea that the same internal representation of space is used for

both visual and haptic perception is also consistent with Müller-Lyer, inverted-T and L-

shape illusions observed for both modalities (Taylor 2001, Gentaz and Hatwell 2004).

In conclusion, the present microgravity experiments testing visual and haptic shape

perception with identical tasks provide the first direct experimental evidence of an effect of

the gravitational signals on the modality-independent processing of spatial information,

which could be only hypothesized in previous unimodal studies (Clement et al. 2009, 2012,

2013).

Neural mechanisms

To understand how gravity affects this internal representation of space, it is important to

consider the neural structures involved in 3D shapes perception. Brain imaging and

electrophysiological studies have shown complex neural networks underlying object

perception (Figure 8A): although 3D object images are known to activate the visual Lateral

Occipital Complex, the same area is active during haptic shape recognition. Similarly, the

areas commonly associated with haptic object perception, (primary and secondary somatosensory cortex, ventral premotor region, and Brodmann area 5) are activated also by images of manipulable objects. These cross-modal activations are mediated by the intraparietal sulcus (IPS) activity which is even enhanced during cross-modal, visuo-haptic object recognition (Grefkes et al. 2002). Electrophysiological studies have shown IPS activities consistent with recurrent neural networks able to perform cross-modal sensory transformations (Pouget et al. 2002, Avillac et al 2005), supporting its role in reconstructing a visual representation of a haptically sensed object and vice-versa (Figure 8B). The coexistence of a visual and a haptic object representation is consistent with the concurrent representations of reaching/grasping tasks aimed at statistically maximizing precision (McGuire and Sabes 2009, 2011, Tagliabue and McIntyre 2011-2014).

To transform a visually-acquired object into a stable haptic representation (and vice-versa), despite potential independent movements of the two sensory systems, IPS must constantly integrate information about the eye-hand kinematic chain and the body position in space. In other words, the IPS contributes to build a stable internal representation of the egocentric space despite eye, head, and body movements (Andersen et al. 1997, Cohen and Andersen 2002, Land 2014).

The similar effect of gravity on both visual and haptic object perception observed here could then be explained by a role of gravitational/otolithic inputs in building the IPS internal representation of space. This idea is in line with the results of several electrophysiological studies in monkeys and humans reporting otolithic projections to IPS area (Figure 8) and with the observed head-tilt effect on the ability to perform visuo-proprioceptive transformations (Tagliabue and McIntyre 2011, 2013, Burns et al. 2011, Bernard-Espina et al. 2022).

A

1. Sakata et al. 1973
2. Koch & Fuster 1989
3. Moore & Engel 2001
4. James et al. 2002
5. Grefkes et al. 2002
6. Amedi et al. 2002
7. Grill-Spector 2003
8. Deshpande et al. 2008

9. Stilla & Sathian 2008
10. Vingerhoets 2008
11. Lacey et al. 2009
12. Meyer et al. 2011
13. Snow et al. 2014
14. Sun et al. 2016
15. Yau et al. 2016

a. Blanke et al. 2000
b. Miyamoto et al. 2007
c. Schlindwein et al.2008

d. Chen et al. 2011
e. Chen et al. 2013

B

Figure 8: A) Brain cortical areas associated to haptic (blue), visual (red) and cross-modal (violet) objects' perception. The regions primarily involved in haptic objects representation are the primary and secondary somatosensory areas (S1 and S2), the Brodmann area 5 (BA5), and the ventral premotor (vPM) area. The 3D object visual representation is known to reside in the lateral occipital complex (LOC). Numbers' location represents the areas in which the corresponding study observed a neural activation during object perception tasks. Green letters represent studies reporting otolithic projection in the intraparietal sulcus (IPS) area. B) Simplified scheme of a putative neural network involved in objects perception. The IPS is connected to both haptic and visual object representation areas (S1/S2/BA5/vPM and LOC respectively). Its interaction with each of the two internal object representations would contribute to spatially interpret the haptic or visual sensory inputs during objects acquisition. IPS would also allow building a visual representation of the object from haptic signals and vice-versa. Otolithic signal affect IPS activity, thus indirectly both visual and haptic object representations.

Conclusions

Our study provides a better understanding of human perception of 3D geometry. For the first time, we have shown a tight link between the axes of distortions characterizing visual and haptic object perception, even if the distortions are in opposite directions. This novel finding, together with the state-of-the-art knowledge about the neural circuitry underlying object perception with reciprocal activations of the visual and haptic systems, suggests a common origin for the observed perceptual biases. This idea is further strengthened by the observation that microgravity has the same incremental effect on visual and haptic object perception, consistent with the existence of a modality-independent perceptive

mechanism: independently from the originating sensory system, the incoming object

information would be treated by neural networks of the parietal cortex and interpreted

through an internal representation of the egocentric 3D space, shaped by gravity (otolithic)

signals. These microgravity experiments, therefore, provide fundamental cues to better

understand the neurophysiology of perception "on Earth". They suggest that unimodal 3D

object perception might not actually exist, suggesting that restraining future investigations

to the brain areas associated with a single sensory modality would be a clear limiting factor

in understanding the neural mechanisms underlying 3D perception.

Bibliography

Armstrong, L. & Marks, L. E. (1999), 'Haptic perception of linear extent.', Perception & psychophysics 61, 1211–1226.

Arnoux, L.; Fromentin, S.; Farotto, D.; Beraneck, M.; McIntyre, J. & Tagliabue, M. (2017), 'The visual encoding of purely proprioceptive intermanual tasks is due to the need of transforming joint signals, not to their interhemispheric transfer.', Journal of neurophysiology 118, 1598--1608.

Avery, G. C. & Day, R. H. (1969), 'Basis of the horizontal-vertical illusion.', Journal of experimental psychology 81, 376–380.

Barnett-Cowan, M.; Snow, J. C. & Culham, J. C. (2015), 'Contribution of Bodily and Gravitational Orientation Cues to Face and Letter Recognition.', Multisensory research 28, 427–442.

Bernard-Espina, J.; Dal Canto, D.; Beraneck, M.; McIntyre, J. & Tagliabue, M. (2022) How Tilting the Head Interferes With Eye-Hand Coordination: The Role of Gravity in Visuo-Proprioceptive, Cross-Modal Sensory Transformations. *Frontiers in Integrative Neuroscience, 16: 788905.*

Blanke, O.; Perrig, S.; Thut, G.; Landis, T. & Seeck, M. (2000), 'Simple and complex vestibular responses induced by electrical cortical stimulation of the parietal cortex in humans.', *Journal of neurology, neurosurgery, and psychiatry* 69: 553--556.

Bourrelly, A.; McIntyre, J. & Luyat, M. (2015), 'Perception of affordances during long-term exposure to weightlessness in the International Space station.', Cognitive processing 16 Suppl 1, 171–174.

Bourrelly, A.; McIntyre, J.; Morio, C.; Despretz, P. & Luyat, M. (2016), 'Perception of Affordance during Short-Term Exposure to Weightlessness in Parabolic Flight.', PloS one 11, e0153598.

Burns, J. K. & Blohm, G. (2010), 'Multi-sensory weights depend on contextual noise in reference frame transformations.', Front Hum Neurosci 4, 221.

Chen, A.; DeAngelis, G. C. & Angelaki, D. E. (2011), 'Representation of vestibular and visual cues to self-motion in ventral intraparietal cortex.', *The Journal of neuroscience : the official journal of the Society for Neuroscience* 31, 12036--12052.

Clément, G.; Vieville, T.; Lestienne, F. & Berthoz, A. (1986), 'Modifications of gain asymmetry and beating field of vertical optokinetic nystagmus in microgravity.', Neuroscience letters 63, 271–274.

Clément, G. & Eckardt, J. (2005), 'Influence of the gravitational vertical on geometric visual illusions.', Acta Astronaut 56(9-12), 911–917.

Clément, G.; Lathan, C. & Lockerd, A. (2008), 'Perception of depth in microgravity during parabolic flight', Acta Astronautica 63(7), 828 – 832.

Clément, G. & Bukley, A. (2008), 'Mach's square-or-diamond phenomenon in microgravity during parabolic flight.', Neurosci Lett 447(2-3), 179–182.

Clément, G.; Fraysse, M.J. & Deguine, O. (2009), 'Mental representation of space in vestibular patients with otolithic or rotatory vertigo.', Neuroreport 20, 457–461.

Clément, G. & Reschke, M. F. (2010), Neuroscience in space, Springer Science & Business Media.

Clément, G.; Skinner, A.; Richard, G. & Lathan, C. (2012), 'Geometric illusions in astronauts during long-duration spaceflight.', Neuroreport 23(15), 894–899.

Clément, G. & Demel, M. (2012), 'Perceptual reversal of bi-stable figures in microgravity and hypergravity during parabolic flight.', Neuroscience letters 507, 143–146.

Clément, G.; Skinner, A. & Lathan, C. (2013), 'Distance and Size Perception in Astronauts during Long-Duration Spaceflight', Life 3(4), 524–537.

Clément, G.; Loureiro, N.; Sousa, D. & Zandvliet, A. (2016), 'Perception of Egocentric Distance during Gravitational Changes in Parabolic Flight.', PloS one 11, e0159422.

Curry, R. E. (1972), A Bayesian Model for Visual Space Perception, in 'Seventh Annual Conference on Manual Control', pp. 187.

De Saedeleer, C.; Vidal, M.; Lipshits, M.; Bengoetxea, A.; Cebolla, A. M.; Berthoz, A.; Cheron, G. & McIntyre, J. (2013), 'Weightlessness alters up/down asymmetries in the perception of self-motion.', Experimental brain research 226, 95–106.

Deshpande G, Hu X, Stilla R, Sathian K (2008) Effective connectivity during haptic perception: a study using Granger causality analysis of functional magnetic resonance imaging data, *NeuroImage* 40: 1807-1814.

Dupin, L.; Hayward, V. & Wexler, M. (2018), 'Radial trunk-centred reference frame in haptic perception.', Scientific reports 8, 13550.

Ernst MO, Banks MS (2002) Humans integrate visual and haptic information in a statistically optimal fashion. Nature 415:429 – 433.

Fasse, E. D.; Hogan, N.; Kay, B. A. & Mussa-Ivaldi, F. A. (2000), 'Haptic interaction with virtual objects. Spatial perception and motor control.', Biological cybernetics 82, 69–83.

Girgus, J. S. & Coren, S. (1975), 'Depth cues and constancy scaling in the horizontal-vertical illusion: the bisection error.', Canadian journal of psychology 29, 59–65.

Goldreich, D. A (2007) Bayesian perceptual model replicates the cutaneous rabbit and other tactile spatiotemporal illusions. *PLoS ONE, 2*, e333

Goltz, H. C.; Irving, E. L.; Steinbach, M. J. & Eizenman, M. (1997), 'Vertical eye position control in darkness: orbital position and body orientation interact to modulate drift velocity.', Vision research 37, 789–798.

Goodenough, D. R.; Oltman, P. K.; Sigman, E. & Cox, P. W. (1981), 'The rod-and-frame illusion in erect and supine observers.', Perception & psychophysics 29, 365–370.

Grefkes, C.; Weiss, P. H.; Zilles, K. & Fink, G. R. (2002), 'Crossmodal processing of object features in human anterior intraparietal cortex: an fMRI study implies equivalencies between humans and monkeys.', *Neuron* 35, 173--184.

Grill-Spector, K. (2003), 'The neural basis of object perception.', *Current opinion in neurobiology* **13**, 159--166.

Gurfinkel, V. S.; Lestienne, F.; Levik, Y. & Popov, K. E. (1993), 'Egocentric references and human spatial orientation in microgravity. I. Perception of complex tactile stimuli.', Exp Brain Res 95(2), 339-42.

Harris, L. R.; Jenkin, M.; Jenkin, H.; Dyde, R.; Zacher, J. & Allison, R. S. (2010), 'The unassisted visual system on earth and in space.', J Vestib Res 20(1), 25–30.

Harris, L. R. & Mander, C. (2014), 'Perceived distance depends on the orientation of both the body and the
visual environment.', Journal of vision 14.

Henriques, D. Y.; Klier, E. M.; Smith, M. A.; Lowy, D. & Crawford, J. D. (1998), 'Gaze-centered remapping of
remembered visual space in an open-loop pointing task.', J Neurosci 18(4), 1583–1594.

Henriques, D. Y. P. & Soechting, J. F. (2003), 'Bias and sensitivity in the haptic perception of geometry.',
Experimental brain research 150, 95–108.

Howard, I. (1982), Human visual orientation, New York: Wiley.

Howard, I. P.; Bergström, S. S. & Ohmi, M. (1990), 'Shape from shading in different frames of reference.',
Perception 19, 523–530.

Kersten, D. & Yuille, A. (2003), 'Bayesian models of object perception.', Current opinion in neurobiology 13,
150–158.

Kersten, D.; Mamassian, P. & Yuille, A. (2004), 'Object perception as Bayesian inference.', Annu Rev Psychol
55, 271–304.

Kim, J. J. J., McManus, M. E. & Harris, L. R. (2022), 'Body Orientation Affects the Perceived Size of Objects.',
Perception 51, 25--36.

Koch, KW. & Fuster, JM. (1989), 'Unit activity in monkey parietal cortex related to haptic perception and
temporary memory.', Experimental brain research 76: 292--306.

Lacey, S.; Tal, N.; Amedi, A. & Sathian, K. (2009), 'A putative model of multisensory object representation.',
Brain topography 21, 269--274.

Lee, T. S. (2015), 'The visual system's internal model of the world.', Proceedings of the IEEE. Institute of
Electrical and Electronics Engineers 103, 1359–1378.

Leone, G. (1998), 'The effect of gravity on human recognition of disoriented objects.', Brain research. Brain
research reviews 28, 203–214.

Lipshits, M. I.; Gurfinkel, E. V.; McIntyre, J.; Droulez, J.; Gurfinkel, V. S. & Berthoz, A. (1994), Influence of
weightlessness on haptic perception, in 'Life Sciences Research in Space', pp. 367.

Lipshits, M. & McIntyre, J. (1999), 'Gravity affects the preferred vertical and horizontal in visual perception of
orientation.', Neuroreport 10(5), 1085–1089.

Liu, Y.; Sexton, B. M. & Block, H. J. (2018), 'Spatial bias in estimating the position of visual and proprioceptive
targets.', Journal of neurophysiology 119, 1879–1888.

Loomis, J. M. & Philbeck, J. W. (1999), 'Is the anisotropy of perceived 3-D shape invariant across scale?',
Perception & psychophysics 61, 397–402.

Loomis, J. M.; Klatzky, R. L. & Giudice, N. A. (2013), Representing 3D space in working memory: Spatial
images from vision, hearing, touch, and language 'Multisensory imagery', Springer, pp. 131–155.

Luyat, M. & Gentaz, E. (2002), 'Body tilt effect on the reproduction of orientations: studies on the visual oblique
effect and subjective orientations.', J Exp Psychol Hum Percept Perform 28(4), 1002-1011.

Marendaz, C.; Stivalet, P.; Barraclough, L. & Walkowiac, P. (1993), 'Effect of gravitational cues on visual
search for orientation.', Journal of experimental psychology. Human perception and performance 19,
1266–1277.

McFarland, J. & Soechting, J. F. (2007), 'Factors influencing the radial-tangential illusion in haptic perception.',
Experimental brain research 178, 216–227.

McGuire, L. M. M. & Sabes, P. N. (2009), 'Sensory transformations and the use of multiple reference frames
for reach planning.', Nat Neurosci 12(8), 1056–1061.

McGuire, L. M. M. & Sabes, P. N. (2011), 'Heterogeneous representations in the superior parietal lobule are
common across reaches to visual and proprioceptive targets.', *J Neurosci* 31(18), 6661--6673.

McIntyre, J.; Stratta, F. & Lacquaniti, F. (1997), 'Viewer-centered frame of reference for pointing to memorized
targets in three-dimensional space.', J Neurophysiol 78(3), 1601–1618.

McIntyre, J. & Lipshits, M. (2008), 'Central processes amplify and transform anisotropies of the visual system
in a test of visual-haptic coordination.', J Neurosci 28(5), 1246–1261.

Mierau, A.; Girgenrath, M. & Bock, O. (2008), 'Isometric force production during changed-Gz episodes of
parabolic flight'. *Eur J Appl Physiol 102*, 313-318.

Miyamoto, T.; Fukushima, K.; Takada, T.; de Waele, C. & Vidal, P.-P. (2007), 'Saccular stimulation of the
human cortex: a functional magnetic resonance imaging study.', *Neuroscience letters* 423: 68--72.

Moore, C. & Engel, S. A. (2001), 'Neural response to perception of volume in the lateral occipital complex.',
Neuron 29, 277--286.

Murdison, T.S.; Leclercq, G.; Lefèvre, P. & Blohm, G. (2019) Misperception of motion in depth originates from
an incomplete transformation of retinal signals., *Journal of Vision*, 19(12): 21, 1—15

O'Reilly, J. X.; Jbabdi, S. & Behrens, T. E. J. (2012) How can a Bayesian approach inform neuroscience?
Eur. J. Neurosci., Wiley, 35, 1169-1179

Paillard, J.J. Paillard, ed., (1991), Brain and Space, Oxford University Press, chapter Knowing where and
knowing how to get there, pp. 461-481.

Prinzmetal, W. & Beck, D. M. (2001), 'The tilt-constancy theory of visual illusions.', J Exp Psychol Hum Percept
Perform 27(1), 206–217.

Reschke, M. F.; Kolev, O. I. & Clément, G. (2017), 'Eye-Head Coordination in 31 Space Shuttle Astronauts
during Visual Target Acquisition.', Scientific reports 7, 14283.

Reschke, M. F.; Wood, S. J. & Clément, G. (2018), 'Ocular Counter Rolling in Astronauts After Short- and
Long-Duration Spaceflight.', Scientific reports 8, 7747.

Sakata, H.; Takaoka, Y.; Kawarasaki, A. & Shibutani, H. (1973), Somatosensory properties of neurons in the
superior parietal cortex (area 5) of the rhesus monkey., *Brain research* 64: 85-102.

Schlindwein, P.; Mueller, M.; Bauermann, T.; Brandt, T.; Stoeter, P. & Dieterich, M. (2008), 'Cortical
representation of saccular vestibular stimulation: VEMPs in fMRI.', *NeuroImage* 39, 19--31.

Smeets JBJ, van den Dobbelsteen JJ, de Grave DDJ, van Beers RJ, Brenner E (2006), Sensory integration
does not lead to sensory calibration., Proc Natl Acad Sci USA 103(49): 18781–18786.

Sun, H.-C.; Welchman, A. E.; Chang, D. H. F. & Di Luca, M. (2016), 'Look but don't touch: Visual cues to
surface structure drive somatosensory cortex.', *NeuroImage* 128, 353--361.

Tagliabue, M. & McIntyre, J. (2011), 'Necessity is the Mother of Invention: Reconstructing Missing Sensory
Information in Multiple, Concurrent Reference Frames for Eye-Hand Coordination.', J Neurosci 31(4),
1397–1409.

Tagliabue, M. & McIntyre, J. (2012), 'Eye-hand coordination when the body moves: Dynamic egocentric and
exocentric sensory encoding.', Neurosci Lett 513, 78-83.

Tagliabue, M.; Arnoux, L. & McIntyre, J. (2013), 'Keep your head on straight: Facilitating sensori-motor
transformations for eye-hand coordination.', Neuroscience 248, 88-94.

Tagliabue, M. & McIntyre, J. (2013), 'When Kinesthesia Becomes Visual: A Theoretical Justification for
Executing Motor Tasks in Visual Space', PLoS ONE 8(7), e68438.

Tagliabue, M. & McIntyre, J. (2014), 'A modular theory of multisensory integration for motor control.', Front
Comput Neurosci 8, 1.

Taylor, C. M. (2001) Visual and haptic perception of the horizontal-vertical illusion. Perceptual and motor skills,
92, 167-170.

Todd, J. T. & Norman, J. F. (2003), 'The visual perception of 3-D shape from multiple cues: are observers
capable of perceiving metric structure?', Perception & psychophysics 65, 31–47.

van Beers, R. J.; Sittig, A. C. & Gon, J. J. (1999), 'Integration of proprioceptive and visual position-information:
An experimentally supported model.', J Neurophysiol 81(3), 1355–1364.

Vetter, P.; Goodbody, S. J. & Wolpert, D. M. (1999), 'Evidence for an eye-centered spherical representation of
the visuomotor map.', Journal of neurophysiology 81, 935–939.

Villard, E.; Garcia-Moreno, F. T.; Peter, N. & Clément, G. (2005), 'Geometric visual illusions in microgravity
during parabolic flight.', Neuroreport 16(12), 1395–1398.

Vingerhoets, G. (2008), 'Knowing about tools: neural correlates of tool familiarity and experience.', *NeuroImage*
40, 1380--1391.

Wagner, M. (1985), 'The metric of visual space', Perception & psychophysics 38(6), 483–495.

Welchman, A. E. (2016), 'The Human Brain in Depth: How We See in 3D.', Annual review of vision science 2,
345–376.

Wolbers, T.; Klatzky, R. L.; Loomis, J. M.; Wutte, M. G. & Giudice, N. A. (2011), 'Modality-independent coding
of spatial layout in the human brain.', Current biology : CB 21, 984–989.

Wydoodt, P.; Gentaz, E. & Streri, A. (2006), 'Role of force cues in the haptic estimations of a virtual length.',
Experimental brain research 171, 481–489.

Appendix I: Graphically Representing Distortion

As we stated in the methods section, it is not possible to univocally quantify the absolute distortion for each of the three dimensions, but only with respect to the other two dimensions. However, in order to provide in the results section a graphical representation of the perceptive distortions, the following method was used to compute the dimensional errors. We first solved the system of equations of Table *1* reported below in the matrix form.

$$
\begin{bmatrix}
1 & 0 & -1 \\
-1 & 0 & 1 \\
1 & -1 & 0 \\
-1 & 1 & 0 \\
0 & -1 & 1 \\
0 & 1 & -1
\end{bmatrix}
\cdot
\begin{bmatrix}
\varepsilon_{EW} \\
\varepsilon_{NS} \\
\varepsilon_{UD}
\end{bmatrix}
=
\begin{bmatrix}
\varepsilon_{EW-UD} \\
\varepsilon_{UD-EW} \\
\varepsilon_{EW-NS} \\
\varepsilon_{NS-EW} \\
\varepsilon_{UD-NS} \\
\varepsilon_{NS-UD}
\end{bmatrix}
$$

If we call A the matrix of linear coefficient, then the solutions of this underdetermined problem are

$$
\begin{bmatrix}
\varepsilon_{EW} \\
\varepsilon_{NS} \\
\varepsilon_{UD}
\end{bmatrix}
= A^{\dagger} \cdot
\begin{bmatrix}
\varepsilon_{EW-UD} \\
\varepsilon_{UD-EW} \\
\varepsilon_{EW-NS} \\
\varepsilon_{NS-EW} \\
\varepsilon_{UD-NS} \\
\varepsilon_{NS-UD}
\end{bmatrix}
+ (I - A^{\dagger}A) *
\begin{bmatrix}
\varepsilon_{EW} \\
\varepsilon_{NS} \\
\varepsilon_{UD}
\end{bmatrix}
= A^{\dagger} \cdot
\begin{bmatrix}
\varepsilon_{EW-UD} \\
\varepsilon_{UD-EW} \\
\varepsilon_{EW-NS} \\
\varepsilon_{NS-EW} \\
\varepsilon_{UD-NS} \\
\varepsilon_{NS-UD}
\end{bmatrix}
+ (I - A^{\dagger}A)w
$$

$$
= A^{\dagger} \cdot
\begin{bmatrix}
\varepsilon_{EW-UD} \\
\varepsilon_{UD-EW} \\
\varepsilon_{EW-NS} \\
\varepsilon_{NS-EW} \\
\varepsilon_{UD-NS} \\
\varepsilon_{NS-UD}
\end{bmatrix}
+
\begin{bmatrix}
w \\
w \\
w
\end{bmatrix}
$$

Where the $A^{\dagger}$ is the pseudo inverse of A and w is a free scalar parameter that reflects the fact that the observed results can be explained by an infinity of triplets of dimensional distortions differing by isotropic component, w, only (underdetermination of the problem).

To define a set of dimensional distortions, $(\varepsilon_{EW}, \varepsilon_{NS}, \varepsilon_{UD})$ to be used for a graphical representation, we arbitrary decided to select the solution that minimizes the Euclidean norm of the distortion vectors.

Although the *w* parameter cannot be univocally defined, the difference between the errors along the three dimensions are correctly quantified and then used to test the anisotropy of the perceptive errors. The dimensional errors, however, cannot be rigorously compared between postures or gravitational conditions, because the differences between

experimental conditions could be due to differences in defining the w parameter for each condition.