T606-phosphorylation deprives the function of Kaiso as a transcription repressor

Wei Tian[1,2], Sisi Qin[1],Wensu Liu, Baozhen Zhang, Liankun Gu, Jing Zhou, Dajun Deng[2]

Key Laboratory of Carcinogenesis and Translational Research (Ministry of Education/Beijing),

Division of Cancer Etiology, Peking University Cancer Hospital and Institute, China

Running title: Phosphorylation of Kaiso at T606 by AKT1

[1] These authors contributed equally to this work.

[2] Correspondence: Key Laboratory of Carcinogenesis and Translational Research (Ministry of Education/Beijing), Division of Cancer Etiology, Peking University Cancer Hospital and Institute, Fu-Cheng-Lu #52, Beijing, 100142, China, Dajun Deng (E-mail: dengdajun@bjmu.edu.cn, Tel:86-10-88196752, Fax: 86-10-88122437) or Wei Tian (tianwei@bjmu.edu.cn, Tel:86-10-88196721, Fax: 86-10-88122437)

Abbreviations

KBS: Kaiso binding sequence

pT606: T606-phosphorylated

Phos-tag: Phos-tag molecular binds specifically to phosphate group in proteins via metal ions

CIAP: calf intestinal alkaline phosphatase

pAKT-Sub: phosphorylated AKT substrate

wt: wildtype

T606A: replacement of the 606th amino acid residue threonine with alanine

Y442F: replacement of the 442th tyrosine with phenylalanine

14-3-3 BM mAb: 14-3-3 binding motif antibody

IP: immunoprecipitation

ELISA: enzyme-linked immunosorbent assay

BSA: bovine serum albumin

PBS: Phosphate-buffered saline

DAPI: 4', 6-diamidino-2-phenylindole

FITC: fluorescein isothiocyanate

CY5: Cyanine5

HC: Heavy chain of IgG

Abstract

It is well known that Kaiso protein encoded by *ZBTB33* gene is a transcription repressor and that

Kaiso-P120ctn interaction increases the shift of Kaiso from the nucleus into the cytoplasm. However,

the regulatory mechanisms of Kaiso compartmentalization are far from clear. Here, we reported that

AKT1 could phosphorylate 606-threonine residue (T606) within the RSSTIP motif of Kaiso in the

cytoplasm. The pT606-Kaiso could directly bind to 14-3-3 family proteins and the loss of T606

phosphorylation byT606A mutation abolished most of the Kaiso-14-3-3 binding. In addition, the

Kaiso-P120ctn interaction was essential for the pT606-Kaiso accumulation in the cytoplasm. Notably,

enforced *14-3-3σ* overexpression could increase the pT606-Kaiso accumulation in the cytoplasm and

de-repress the transcription of Kaiso target genes, including *CDH1*. Decreased amounts of both

T606-phosphorylated Kaiso (pT606-Kaiso) and CDH1 were frequently observed in human gastric and

colon cancer tissues relative to paired normal controls. The mRNA levels of *14-3-3σ/SFN* and Kaiso

target gene *CDH1* were positively and significantly correlated with each other in bioinformatics

analyses using publicly available RNA-seq datasets for human normal tissues ($n=11688$, $r=0.60$,

$p<0.001$) in the GTEx project and for cancer cell lines ($n=1156$, $r=0.41$, $p<0.001$) in the CCLE project.

In conclusion, Kaiso could be phosphorylated by AKT1 at the T606 and the pT606-Kaiso accumulates

in the cytoplasm through binding to 14-3-3/P120ctn that de-represses the expression of Kaiso target

genes and may contribute to the development of gastrointestinal cancer.

Keywords: Kaiso, phosphorylation, AKT1, 14-3-3, P120ctn, transcription factor, cancer

Introduction

Kaiso protein encoded by the *ZBTB33* gene is a classic transcription repressor containing a zinc-finger domain and a BTB/POZ domain (1). The zinc-finger domain of Kaiso can bind to both methylated CGCG-containing and non-methylated Kaiso binding-specific sequences of Kaiso target genes, and the BTB/POZ domain can further recruit the complex of NCoR1 corepressor and histone deacetylases to target genes and repress their transcription in the nucleus (2-4). Recent studies show that Kaiso may also act as a transcription activator in the promoter context-dependent manner (5, 6).

As a transcription repressor, Kaiso controls the cell cycle through repressing *CCND1* and *CCNE1* expression, affects Notch signaling pathway in intestinal cells through targeting *DLL1* and *JAG1* promoter, and inhibits the proliferation and invasion of tumor cells through downregulating *MMP7*, *MTA2* and other genes (7-10). Kaiso also represses *CDH1* and *CDKN2A* expression (11,12). Interestingly, it has been reported that the amount of nuclear Kaiso, but not total Kaiso, is correlated with the invasion or prognosis of cancers (13, 14). Kaiso-deficient mice show resistance to intestinal cancer (15). Apparently, the expression and subcellular location states of Kaiso determine its normal functions and roles in cancer development.

Kaiso is also a cytoplasm protein, which regulates WNT-related pathway through interacting with P120ctn protein (16). Differences of subcellular locations of Kaiso are observed between cultured cells and tissues (12, 17). The P120ctn-Kaiso complexes could shift from the nucleus to the cytoplasm (18). Kaiso nuclear-cytoplasmic trafficking could be affected by environmental factors, such as cigarette smoke, through MUC1 and P120ctn binding (19). However, detailed regulation machinery for the compartmentalization of Kaiso remains far from clear.

The 14-3-3 proteins are originally identified in the brain (20). There are seven human 14-3-3 isoforms (α/β, ε, η, δ/γ, τ, ζ, σ). These 14-3-3 isoforms are homologous proteins with approximately 50% amino acid identity, capable of forming either homo- or hetero-dimers (21-24). Recent findings have implicated 14-3-3 as a key regulator of signal transduction events (25). Among the family, 14-3-3γ and 14-3-3σ have been confirmed to play important roles in cancer development (26-29).

In the present study, we found, for the first time, that Kaiso could be phosphorylated at Thr-606 (T606) within the RSS<u>T</u>IP motif by the protein serine-threonine kinase AKT1 and

T606-phosphorylated Kaiso (pT606-Kaiso) could efficiently interact with 14-3-3 family member and P120ctn proteins, and accumulate in the cytoplasm that in turn blocks transcription repressor function of Kaiso and de-repressed the expression of its target gene *CDH1*. Relative to the paired normal control tissues, the level of pT606-Kaiso was markedly decreased in human gastric and colon cancer tissues that could lead to transcriptional repression of the *CDH1* gene.

Materials and Methods

Cell lines and culture

The gastric cancer cell lines MGC803, BGC823 and SGC7901 were kindly provided by Dr. Yang Ke at Peking University Cancer Hospital; MKN45 cell line was purchased from the National Infrastructure of Cell Line Resource (Beijing, China). The human embryonic kidney HEK293T cell line was kindly provided by Professor Yasuhito Yuasa at Tokyo Medical and Dental University. The human colon cancer cell line HCT116 was kindly provided by Dr. Yuanjia Chen, Peking Union Hospital; RKO cell line, by Dr. Guoren Deng, University of California. HEK293T cells was cultured in DMEM medium containing 10% FBS, and all the others were cultured in RPMI 1640 medium containing 10% FBS and 100 U/mL penicillin/streptomycin (Life Technologies, Carlsbad, CA, USA) at 37 °C in a humidified incubator with 5% CO_2. These cell lines were tested and authenticated by Beijing JianLian Gene Technology Co., before use. Short tandem repeat (STR) patterns were analyzed using GoldeneyeTM20A STR Identifiler PCR Amplification Kit.

Gastric and colon carcinoma tissues and ethical issue

Gastric and colon cancer tissues and the paired normal surgical margin tissues were respectively collected from 4 patients with gastric carcinoma and 4 patients with colon carcinoma at Peking University Cancer Hospital from 2000 to 2001 and stored at −70 °C. The Institutional Review Board of the Peking University Cancer Hospital approved the study. All of the patients provided written informed consent.

Plasmids and reagents

The full-length Kaiso-coding sequence of the *ZBTB33* gene was amplified from human cDNA of

MGC803 cells with a primer set (forward 5'-attaaactcgaggcatggagagtagaaaactga-3' and reverse

5′-cgcttcgaattcgtttagtaagactctggtattat-3'), then inserted into *Xho*I and *Eco*RI sites of pEGFP-C1 vector

to generate pEGFP-C1-Kaiso expression vector. pEGFP-C1-Kaiso-T606 mutants were obtained by

mutation PCRs using a primer set (forward 5'-gatagatcaagcgctattcctgcaatg -3' and reverse

5'-cattgcaggaatagcgcttgatctatc-3') for 606Thr → Ala (T606A) mutation. Plasmid

pCMV-3Tag-2C-Kaiso was generated by inserting the full-length Kaiso-coding sequence into *BamH*I

and *Eco*RI sites of pCMV-3Tag-2C vector. Plasmid pEBG-GST-Kaiso was generated by inserting the

full-length Kaiso coding sequence into *BamH*I and *Not*I sites of pEBG vector. Kaiso fragment

(aa346-638) coding sequence without BTB/POZ domain was amplified with a primer set (forward

5'-tatcggaattccctgactcggccgtcagt-3' and reverse 5'-attgcctcgagcattggcttgttctgagt-3'), then inserted into

*Eco*RI and *Xho*I sites of vector pGEX-4T-1.

pcDNA3.1-HA-AKT1 vector was purchased from Addgene (#9008, MA, USA);

pEZ-M56-14-3-3γ-mCherry, pEZ-M56-14-3-3σ-mCherry vectors, from FulenGen (EX-T4084-M56,

EX-T4084-M98-5, EX-C0507-M98, Guangzhou, China); pENTER-Flag-14-3-3 isoforms (α /β, ε, η,

δ/γ, τ, ζ, σ) were purchased from Vigene Bioscience (CH867785, CH897212, CH845486, CH898602,

CH878525, CH824520, CH890307, Shandong, China); Insulin (P3376, Beyotime, Shanghai, China),

IL-6 (Cat. 200-06, Proteintech, NJ, USA), EGF (PHG6045, Thermo Fisher Scientific, MA, USA),

MK2206 (HY-10358, MedChemExpress, NJ, USA) were also used in the study. *P120ctn*-specific

siRNAs (#1: sense 5'-gaaugugaugguuuaguuuu-3' and antisense 5'-aacuaaaccaucacauucuu-3'; #2: sense

5'-uagcugaccuccugacuaauu-3' and antisense 5'-uuagucaggaggucagcuauu-3'; #3: sense

5'-ggaccuuacugaaguuauuuu-3' and antisense 5'-aauaacuucaguaagguccuu-3') were synthesized by

Genepharma (Shanghai, China). The quantitative RT-PCR primer sequences for detection the level of

CDH1 mRNA were: forward 5'-gaacgcattgccacatacac-3' and reverse 5'-gaattcgggcttgttgtcat-3'

(Tm=58 °C). The *Alu* RNA was used as the reference (forward 5'-gaggctgaggcaggagaatcg-3' and

reverse 5'-gtcgcccaggctggagtg-3', Tm=60°C), as previously described (30).

Cell transfection and xenografts in nude mice

X-tremeGENE siRNA Transfection Reagent or X-tremeGENE HP DNA Transfection Reagent (Cat.

04476093001, Cat. 06366236001, Roche, Mannheim, Germany) were used in cell transfection with

siRNAs for *p120ctn* (final concentration, 100 nM) or Kaiso or its mutant expression vectors (2 μg/well

in 6 wells plate) following manufacturer's instructions. The efficiency of gene overexpression or

knockdown was determined 48 or 72 hrs post transfection by Western blotting. For stable transfection,

G418 was added into the medium to select consistent GFP-Kaiso expressing MGC803 cells (final

concentration, 750 μg/mL). These cells resuspended in 0.2 mL PBS (1×10^7 cells/mL) were inoculated

subcutaneously into the right/left pads of 6 week-old male BALB/c nude mice. Mice were sacrificed on

days 21 after inoculation, and xenografts were separated and cut into about 1-mm^3 pieces for protein

fractionation.

Subcellular fractionation and de-phosphorylation treatment (31, 32)

To prepare cytoplasmic and nuclear extracts, small pieces of tissue or cultured cells at 80% confluence

were homogenized in ice-cold buffer A (REF04693159001, Roche, Mannheim, Germany) on ice for 20

min. Samples were then supplemented with NP-40 to 0.1% (v/v), vortexed, and centrifuged at 14,000 *g*

for 15 min at 4 °C. The supernatants were recovered to obtain the cytosolic extracts. The pellets were

washed sequentially with ice-cold buffer A and then incubated in buffer B on ice for 20 min. After

centrifuging at 14,000 g for 15 min, the extract was collected as nuclear proteins. The purities of

cytoplasmic and nuclear extracts were respectively verified by probing with anti-β-Tubulin and

anti-Lamin B antibodies.

For the calf intestinal alkaline phosphatase (CIAP)-catalyzed de-phosphorylation, the cytoplasmic

and nuclear extracts were aliquoted into two centrifuge microtubes. CIAP (p4978, Merck, Darmstadt,

Germany) (1 U/μg) of protein was added into one aliquot in the CIAP buffer and incubated for 30 min

at 37 °C.

Immunoprecipitation (IP) and Western blotting

Antibodies for Kaiso (sc-365428, Santa Cruz, USA), P120ctn (66208-1, Proteintech, IL, USA),

Pan-14-3-3 (sc-629, Santa Cruz), 14-3-3 family (α/β, ε, η, δ/γ, τ, ζ, σ) kits (#9769, CST, USA),

14-3-3σ (sc-100638, Santa Cruz), Ser/Thr/Tyr phosphor-protein (ab15556, Abcam, UK),

phosphorylated AKT substrate [(R/K)X(R/K)XX(pT/pS)] (pAKT-sub; #9611, Cell Signaling

Technology), CDH1 (#3195, CST), Lamin B1 (66095-1, Proteintech), β-Tublin (66240-1, Proteintech), HA (M20003, Abmart, Shanghai, China), FLAG (66008-2, Proteintech), GFP (NB100-1614, Novus, CO, USA), GST (66001-1, Proteintech), GAPDH (660004-1, Proteintech) were used in the IP and Western blotting analyses.

After being pre-cleared with protein A/G-coupled Sepharose beads (Cat. 11134515001 and 11243233001, Roche, Mannheim, Germany) for 2 hrs, the nuclear or cytoplasmic lysate was immunoprecipitated with mouse anti-Kaiso antibody or anti-Phosphoserine/threonine/tyrosine antibody plus protein A/G Sepharose for 8 hrs at 4 °C. Mouse IgG was used as a negative control. The precipitates were washed six times with lysis buffer, and boiled in $1 \times$ loading buffer. Protein samples were resolved by SDS–PAGE, and electroblotted onto nitrocellulose membranes, which were blocked in 5% skim milk in PBST and probed with antibodies according to the Instruction Manual.

Phos-tag SDS-PAGE assay

This is a modified SDS-PAGE method based on the novel Phos-tag (33), which can bind to phosphorylated proteins and decrease their migration speed. Thus, this assay is often used to distinguish dephosphorylated proteins from phosphorylated proteins on the case of that a phosphorylation-specific antibody is not available. 50 μM Phos-tag (final concentration, Phos-tag Acrylamide AAL-107, WAKO, Japan) and 100 μM $MnCl_2$ were mixed to prepare SDS-PAGE gel.

GST-Pull down (34)

The day after pEBG-GST-Kaiso transfection, MGC803 cells (in 10-cm dishes, eight dishes/group) were further transfected with pENTER-Flag-14-3-3 members (α/β, ε, η, δ/γ, τ, ζ, σ) or negative control vector, respectively. 48 hrs post transfection, MGC803 cells were harvested and used to prepare lysate with cell lysis buffer with $1\times$ Protease Inhibitor Cocktail (REF04693159001, Roche, Mannheim, Germany). The lysate was incubated with Glutathione Sepharose beads (20 μL for one group, 17-0756-01, GE healthcare, Sweden) at 4 °C overnight. Beads were washed six times with 500 μL cell lysis buffer. After the last centrifuging, the supernatant was removed as clean as possible, and the pellet was suspended in 20 μL 1×SDS sample buffer. Antibodies for GST-Kaiso and Flag-14-3-3 family proteins were used to detect the pulled down precipitant.

Co-immunoprecipitation (Co-IP)

After pre-cleared with protein A/G-coupled Sepharose beads for 2 hrs, the soluble proteins from whole cell lysate were immunoprecipitated with anti-Kaiso (sc-365428, Santa Cruz) or anti-GFP (ab290, Abcam) or other antibodies plus protein A/G Sepharose overnight at 4 °C. Mouse IgG or rabbit IgG was used as a negative control. The precipitates were washed six times with lysis buffer, and boiled after 1 × SDS loading buffer was added. Protein samples were resolved by SDS–PAGE, and electroblotted onto nitrocellulose membranes, which were blocked with 5% skim milk in PBST and probed with the interacted protein antibodies.

***In vitro* kinase assay** (35)

1 µg of the substrate (GST or GST-Kaiso-wt/T606A) protein per reaction was extracted from pGEX-4T1 plasmid transfected *BL21 E. coli* induced with IPTG 0.5 mM at 37 °C and purified with Glutathione Sepharose beads. Active kinase AKT1 protein (PKBα, 15 µg) was purchased from Merck (#14-276, Darmstadt, Germany). 10× Kinase buffer (1 × concentration: 25 mM Tris (pH 7.5), 5 mM β-glycerophosphate, 2 mM DTT, 0.1 mM Na_3VO_4, 10 mM $MgCl_2$) was prepared when used. Glutathione Sepharose beads were washed with substrate twice with 500 µL of 1 × Kinase Buffer on ice. The suspended pellet in 30 µL of 1 × Kinase buffer supplemented with 1 µL of 10 mM ATP and 0.5 µg AKT1 kinase were incubated at 30°C for 30 min. Reaction was terminated with 30 µL 2 × SDS Sample buffer. Samples were vortexed, then microcentrifuged for 30 sec at 12,000 *g*. The reaction products were then resolved in SDS-PAGE gel and Western blotted to detect the phosphorylated Kaiso using AKT1 substrate antibody.

Preparation of pT606-Kaiso-specific antibody

Anti-pT606-Kaiso polyclonal antibodies were raised in rabbits challenged with the synthesized phosphorylated peptide LSD**RSSpTIP**AM, a sequence corresponding to amino acids 600-610 of wildtype Kaiso (Kaiso-wt), absorbed with non-phosphorylated peptide LSD**RSSTIP**AM, and enriched by the phosphorylated peptide LSD**RSSpTIP**AM. Peptide synthesis and immunization of the animals were done by YouKe (Shanghai, China). The non-phos-peptide was also used to prepare the control polyclonal antibodies against total Kaiso at the same time. The specificity of pT606-Kaiso and control

antibodies against the corresponding peptide (LSDRSSpTIPAM or LSDRSSTIPAM) was detected with ELISA assay (Fig. S1).

ELISA analyses for polyclonal antibodies against pT606-Kaiso and total Kaiso

Polystyrene plates were coated with 1 μg/mL synthetic phosphorylated peptide LSDRSSpTIPAM or non-phosphorylated peptide LSDRSSTIPAM link-coupled by bovine serum albumin (BSA) in $1 \times$ CBS buffer overnight at 4 °C, respectively, and were washed three times with PBS containing 0.05% Tween 20. Unbinding sites were blocked with 5% milk at room temperature for 2 hrs. The purified antibodies, diluted as described in Figure S1, were added (100 μL/well) and incubated at 37 °C for 1 hr. After being washed with 0.05% Tween/PBS, plates were added HRP-labeled goat anti-Rabbit IgG (100 μL/well) and incubated at room temperature for 30 min. Peroxidase activity was measured with 0.15 mg/mL TMB substrate solution (100 μL/well). After 15 min at room temperature, the reaction was stopped by 2M H_2SO_4 (50 μL/well). Optical density absorbance (OD) at 450 nm was determined using a model 550 microplate reader.

Immunohistochemical staining (IHC)

Rabbit pT606-Kaiso polyclonal antibody (1:100) and mouse monoclonal antibody for total Kaiso (1:50, sc-23871, Santa Cruz) were used in the IHC analysis. The REAL™ EnVision™ Detection system including anti-rabbit/mouse IgG secondary antibodies and peroxidase/diaminobenzidine (Dako, Agilent Technologies) was used to visualize the primary antibody-binding cells according to the manufacturer's protocol. Briefly, paraffin sections (4 μm) were dewaxed and rehydrated in xylene and ethanol. For antigen retrieval these sections were autoclaved for 3 min in 10 mM sodium citrate buffer containing 0.05% Tween-20 (pH 6.0) for pT606-Kaiso and in 1 mM EDTA buffer (pH 8.0) for total Kaiso. Then, these sections were immersed in 3% H_2O_2 for 10 min to block endogenous peroxidase. Following submerging in 5% bovine serum albumin (BSA) (cat. no. A1933; Sigma Life Science; Merck KGaA) for 60 min, the sections were incubated with the primary antibody overnight at 4 ˚C. The PBS washed sections were then treated with the REAL™ EnVision™ Detection system and counterstained with hematoxylin (0.125%; ZhongShan Jinqiao Biotechnology) at room temperature for 1 min. Normal rabbit IgG and normal mouse IgG (cat. no. ZDR 5003 and cat. no. ZDR 5006; ZhongShan Jinqiao

Biotechnology) were used as negative controls. The former was diluted and incubated as for the pT606-Kaiso antibody, and the latter was diluted and incubated as for total Kaiso antibody).

Confocal analysis

For mCherry-14-3-3, direct fluorescence was detected with laser confocal microscope assay. MGC803 cells with mCherry-14-3-3 overexpression were rinsed for three times with PBS, fixed with 1% paraformaldehyde in PBS 30 min at 37 °C, punched with 0.5% triton X-100 for 10 min at 37 °C, incubated with respective primary and secondary antibodies, washed for three times in PBS, counterstained with DAPI (1 µg/mL) for 5 min, and then examined with Leica SP5 Laser Scanning Confocal Microscopy. The antibodies for pT606-Kaiso (1 µg/µL, 1:100), for total Kaiso (commercial antibody (0.2 µg/µL, 1:20, sc-365428, Santa Cruz), and for p120ctn (1:100, 66208-1, Proteintech) were used as the primary antibodies; the FITC-labeled antibody against rabbit IgG (1:100, ab6717, Abcam) and CY5-labeled antibody against mouse IgG (1:100, Cat. No. 072-02-18-06, KPL Gaithersburg, MD, USA) were used as secondary antibodies for observation under Leica SP5 Laser Scanning Confocal Microscope and were analyzed with ImageXpress Micro high content screening System.

For the detection of Kaiso in tissues, fresh cryostat sections (4 μm) from gastric carcinoma and the paired normal tissues were fixed with 4% paraformaldehyde for 10 min at 37°C and treated with 0.5% triton X-100 for 10 min at 37 °C, and then incubated with respective primary, secondary antibodies, DAPI, and then examined as described above.

Publicly available RNA-Seq datasets

The RNA sequencing datasets in CCLE and GTEx projects were downloaded from official websites (portals.broadinstitute.org/ccle and www.gtexportal.org) (36, 37). All raw data were transferred to Transcripts per kilobase of exon model per Million mapped reads (TPM) using uniform names for each protein-coding gene in the human genome. The Pearson correlation coefficient (r) of genes expression was calculated by R statistical software (version 3.6.1) with a set of procedures (38). Briefly, raw data was read through "read.table()" function, and an array was established to store data by "array()" function. Then the Pearson correlation coefficient was calculated by "cor()" function, and finally saving the data to files in csv format by the function of "write.table()".

Results

Discovery of phosphorylated Kaiso in the cytoplasm

Immunoaffinity profiling of phosphorylated proteins in cancer cells suggests there may be putative

phosphorylation sites within Kaiso in proteomic mass spectrometry analysis (Fig. S2) (39, 40). We

wondered if Kaiso compartmentalization was related to the statuses of the putative phosphorylation. In

Phos-tag SDS-PAGE analyses, a Phos-tag could bind specifically to a phosphate group in proteins via

metal ions, such as Zn^{2+} or Mn^{2+}, which could be used to separate phosphorylated proteins from

non-phosphorylated proteins (33). Thus, the Phos-tag assay was preliminarily utilized to analyze the

phosphorylation status of Kaiso in human cells. We found that while the cytoplasmic and nuclear Kaiso

from MGC803 and HEK293T cells migrated at the same speed in regular SDS-PAGE gel,

approximately all the cytoplasmic Kaiso migrated much slower than the nuclear Kaiso in the Phos-tag

gel (Fig. 1A). While the migration of nuclear Kaiso was not affected by the CIAP de-phosphorylation

treatment, the delayed cytoplasmic Kaiso migration was partially concealed by the CIAP treatment,

suggesting that the endogenous cytoplasmic Kaiso in these cells was mostly phosphorylated.

To validate the different phosphorylation statuses of Kaiso in the cytoplasm and nucleus, an

anti-phosphoserine/threonine/tyrosine universal antibody was then used to precipitate global

phosphorylated proteins and a Kaiso-specific antibody was used to visualize the possible

phosphorylated Kaiso, and vice versa (Fig. 1B). Again, phosphorylated Kaiso was observed only in the

cytoplasmic precipitates, but not in the nuclear counterpart. Furthermore, the phosphorylated

GFP-Kaiso was consistently detected in the cytoplasm of MGC803 cells stably transfected with

GFP-Kaiso, whether these cells were cultured *in vitro* or transplanted into nude mice as a xenograft in

the Phos-tag SDS-PAGE analyses, and no phosphorylated Kaiso was detected with the CIAP

de-phosphorylation (Fig. 1C). All the above results suggest there may be phosphorylation of Kaiso in

the cytoplasm. Thus, Kaiso phosphorylation kinase and target site were characterized in details as

described below.

Kaiso is phosphorylated at T606 by AKT1 kinase

Protein kinase B AKT1 is a typical kinase for the (RX)RXXpS/pT motif in multiple proteins such as mTOR, GSK-3β, AMPKA, Catenin-β1 (41-44). Human Kaiso contains a conservative RSS<u>T</u>IP motif with the threonine-606 (T606) residue (Fig.2A). According to the results of liquid chromatography-mass spectrum (LC/MS) analyses for protein modifications, the T606 was one of phosphorylation sites in Kaiso with a low frequency (Fig. S2) (40). To verify if Kaiso T606 could be a true phosphorylation site for the Kinase AKT1, we stimulated the activity of AKT signaling of MGC803 cells with insulin, IL-6, and FBS after overnight starvation, as Hong et al. reported (45). As expected, the phosphorylation level of endogenous Kaiso was markedly increased in MGC803 cells stimulated with insulin, IL-6, and FBS, but not with EGF (Fig. 2B).

Using the immunoprecipitation assay employing a phospho-(Ser/Thr) AKT substrate (pAKT-sub)-specific antibody, endogenous Kaiso signal could also be detected in the immunoprecipitated pAKT-sub protein complexes, and vice versa (Fig. S3A). The amounts of both global pAKT-sub and Kaiso-antibody-immunoprecipitated pAKT-sub were also significantly increased by insulin stimulation and reversed by AKT inhibitor MK2206 treatment (Fig. 2C). A similar effect was also observed in MGC803 cells with *AKT1* overexpression (Fig. 2D).

We further found that active AKT1 recombinant kinase could directly phosphorylate GST-Kaiso recombinant protein (aa346-638 in Kaiso-wt) in an *in vitro* kinase assay. When the phosphorylation site T606 was mutated as A606, which is not a phosphorylation site for any kinase, most Kasio-T606A protein could not be phosphorylated by the AKT1 recombinant kinase (Fig. 2E). Together, these results suggest that AKT1 could directly phosphorylate Kaiso at T606.

Then, a pT606-Kaiso-specific polyclonal antibody was prepared from rabbit using the phos-peptide LSD**RSSpTIP**AM as antigen (Fig. S1) and used to directly detect phosphorylated Kaiso. As expected, the amount of pT606-Kaiso was significantly increased in MGC803 and BGC823 cells with the insulin stimulation (or *AKT1* overexpression) and reversed by the AKT inhibitor MK2206 treatment (Fig. 2F-G).

In consistent with the results described above, most pT606-Kaiso was detected in the cytoplasm of human cancer cell lines (MGC803, BGC823, HCT116, and RKO) and the CIAP de-phosphorylation

treatment markedly decreased the amount of pT606-Kaiso in the cytoplasm in Western blotting (Fig.

3A). The confocal microscopy results further confirmed the cytoplasmic compartmentalization of

pT606-Kaiso in MGC803 cells (Fig. 3B). While total Kaiso was mainly detected in the nucleus of

human gastric mucosal tissues, pT606-Kaiso was only detected in the cytoplasm in the IHC and

confocal microscopy analyses (Fig. 3C-D). Collectively, the above results demonstrate that AKT1 can

phosphorylate Kaiso at T606 and pT606-Kaiso mainly located in the cytoplasm of human cells.

pT606-Kaiso interacted with 14-3-3 family members

It is well known that RSXpSXP is a 14-3-3 phosphoserine binding consensus motif (49). To study

whether the Kaiso RSSpTIP is a 14-3-3 binding motif, we performed GST-pull down and Co-IP

experiments. The results of GST-pull down assay showed that GST-Kaiso could strongly pull down

14-3-3σ, moderately pull down 14-3-3ε, 14-3-3γ, 14-3-3ζ, and weakly pull down 14-3-3η in MGC803

cells lysate (Fig. 4A). The Co-IP results confirmed that endogenous Kaiso could bind to endogenous

pan-14-3-3 or 14-3-3σ (SFN) protein in MGC803 cells (Fig. 4B). As expected, the T606A mutation

abolished most Kaiso-14-3-3 interaction (Fig. 4C), suggesting that the phosphorylation site T606

within the RSSTIP motif may play a main role in determining the Kaiso-14-3-3 interaction. In addition,

AKT1 overexpression could slightly increase the Kaiso-14-3-3 interaction (Fig. S3C). These data

indicate that Kaiso could interact with 14-3-3 family proteins in a T606 phosphorylation-dependent

manner.

Increased cytoplasmic accumulation of pT606-Kaiso by the 14-3-3σ interaction

Kaiso usually distributes in both the cytoplasm and the nucleus. Intracellular Kaiso

compartmentalization is affected by growth conditions (13, 17). Our indirect immunofluorescence

confocal microscope analyses showed that 81% endogenous Kaiso was observed in the nucleus of

MGC803 cells (Fig.5A-B). Notably, enforced mCherry-14-3-3γ or -14-3-3σ overexpression

significantly increased the proportion of endogenous Kaiso in the cytoplasm, compared to the mCherry

control vector (from 19% to 37% or 36%) and Kaiso was mainly co-localized with mCherry-14-3-3γ or

-14-3-3σ in the cytoplasm of these cells in the confocal analysis. Western blotting confirmed that

mCherry-14-3-3σ overexpression increased the cytoplasmic accumulation of endogenous Kaiso (Fig.

5C). In addition, 14-3-3σ only promoted the cytoplasmic accumulation of wildtype Kaiso (Kaiso-wt),

but not the T606A mutant (Fig. 5D, red arrowed). 14-3-3σ or 14-3-3γ overexpression also induced

more cytoplasmic pT606-Kaiso in MGC803 and BGC823 cell lines in Western blotting (Fig. S3D-E).

These results solidly confirm that the pT606-Kaiso can interact with 14-3-3 and in turn accumulate in

the cytoplasm of human cancer cells.

The P120ctn interaction is essential for pT606-Kaiso-14-3-3 accumulation in the cytoplasm

As a given Kaiso interacting protein, the P120ctn binding is essential for the trafficking of Kaiso from

the nucleus to the cytoplasm (18, 19). Thus, we further studied whether 14-3-3 bound to the

pT606-Kaiso-P120ctn complex or individual pT606-Kaiso molecule in the cytoplasm. Interestingly, we

found that 14-3-3σ overexpression markedly increased the amount of P120ctn in the anti-Kaiso

antibody precipitated complexes, while the levels of total endogenous Kaiso and P120ctn proteins were

not changed in MGC803 cells (Fig. 6A). Further analysis showed that more Kaiso-P120ctn binding was

detected in the cytoplasm, but not in the nucleus of MGC803 cells with 14-3-3σ overexpression (Fig.

6B). Immunofluorescence confocal microscopy showed that P120ctn, Kaiso, and 14-3-3σ were

co-localized in the cytoplasm (Fig. 6C). Moreover, when P120ctn was knocked down by siRNA, the

cytoplasmic Kaiso accumulation promoted by 14-3-3σ disappeared in the MGC 803 cells (Fig. 6D)

while the level of total Kaiso was not changed (Fig. 6E). These results suggest that the P120ctn

interaction is essential for the pT606-Kaiso-14-3-3σ accumulation in the cytoplasm.

De-repression of Kaiso target genes by T606 phosphorylation

It was reported that Kaiso could bind to both methylated and non-methylated Kaiso binding sequences

in target gene promoters and suppress their transcription (11). We wondered whether T606

phosphorylation of Kaiso and its bindings with 14-3-3/P120ctn proteins could affect expression of

Kaiso target genes through the cytoplasmic accumulation. Through re-analyzing the RNA-seq datasets

for 14-3-3 family members and Kaiso target genes (including *CDH1*, *CCND1*, *CCNE1, MTA2, DLL1,*

and *DAG1*) from Genotype-Tissue Expression (GTEx) and Cancer Cell Line Encyclopedia (CCLE)

(38-40), we found that the level of 14-3-3σ mRNA was positively and most significantly correlated

with that of *CDH1* in both normal human tissues (n=11688, r=0.60, P<0.001 in Pearson correlation

analysis) in GTEx datasets and cancer cell lines in CCLE datasets (n=1156, r=0.41, P<0.001) (Fig. 7A and Fig. S4A), suggesting an exact effect of 14-3-3σ expression on regulation of *CDH1* transcription, probably through the pT606-Kaiso-14-3-3σ interaction. Interestingly, association between the levels of *CDH1* and *Kaiso* mRNA was only observed in normal tissues in the GTEx datasets, but not in cancer cell lines in the CCLE datasets (Fig. S4B), suggesting a dysregulation of *CDH1* transcription by Kaiso in cancer cells.

Then, two gastric cancer cell lines SGC7901 and MKN45 with active *CDH1* expression were used to study effects of Kaiso and 14-3-3σ expression changes on the function of Kaiso as a transcription repressor. As expectedly, Kaiso overexpression alone indeed decreased the amount of CDH1 whereas 14-3-3σ co-overexpression could abolish Kaiso-induced *CDH1* repression in SGC7901 and MKN45 cells in Western blotting (Fig. 7B). These results were confirmed in qRT-PCR analyses (Fig. 7C). A similar relationship was observed in gastric and colon carcinoma tissues. While the amount of pT606-Kaiso was much lower in these cancer tissues than in the paired normal tissues, the amount of CDH1 was also much lower in these cancer tissues than in the paired normal tissues (Fig. 7D and Fig. S5). Taken together, the above results indicate that pT606-Kaiso-14-3-3σ binding in the cytoplasm can deprive the function of Kaiso as a transcription repressor and lead to de-repression of the transcription of Kaiso target gene *CDH1*.

Discussions

The subcellular locations of Kaiso determine its roles in normal cell differentiation and cancer development. However, detailed regulation machinery for the compartmentalization of Kaiso is far from clear. In this study we demonstrated, for the first time, that Kaiso could be phosphorylated at T606 by AKT1, the pT606-Kaiso could interact with 14-3-3 and P120ctn in the cytoplasm and promote the shift of Kaiso from the nucleus to the cytoplasm. The phosphorylation of Kaiso finally leads to deprivation of Kaiso as a transcription repressor and de-repression of Kaiso target gene *CDH1*.

It's well known that 14-3-3 family proteins bind to common phosphoserine/phosphothreonine -containing peptide motifs corresponding to Mode-1 (RSXpSXP) or Mode-2 (RXY/FXpSXP) sequences (46). We found that Kaiso contains a very conservative RS<u>S</u>TIP motif that could be phosphorylated by AKT1 in both *in vivo* and cell-free system at RSS<u>T</u>IP-T606 site. This is consistent

with the report that T606 is one of phosphorylation sites of Kaiso in mass spectrometry analysis (40). That pT606-Kaiso could directly bind 14-3-3 family proteins and the T606A mutation could abolish most Kaiso-14-3-3 binding indicate the T606 phosphorylation is essential for Kaiso-14-3-3 binding.

It has been reported that a region consisting of 1-200 amino acid residues of Kaiso directly interacts with P120ctn (1) and Kaiso-P120cnt interaction promotes cytoplasmic-nuclear trafficking of Kaiso (18, 19). In addition, P120ctn could bind to CDH1 and modulate its function and stability (47) and WNT-stimulated P120ctn phosphorylation could promote P120ctn releasing from the E-Cadherin-P120ctn complexes, enhances Kaiso-P120ctn interaction (48). It is well known that most 14-3-3σ proteins localize in the cytoplasm (49) while Kaiso mainly localizes in the nucleus (50). Here, we observed that pT606-Kaiso, 14-3-3, and P120ctn proteins could collocate in the cytoplasm and siRNA-knockdown of *P120ctn* abolished the 14-3-3-induced cytoplasmic Kaiso-14-3-3 binding. These results demonstrate that Kaiso-14-3-3σ interaction may be dependent on the Kaiso-P120ctn binding and that Kaiso, 14-3-3σ, and P120ctn may form a triplex in the cytoplasm. How Kaiso, 14-3-3σ, and P120ctn interacting with each other is worth studying.

It is controversy on Kaiso's roles in cancer development. In the absence of the tumor suppressor *APC*, Kaiso-deficient mice were resistant to intestinal cancer, suggesting that Kaiso might be an oncogene (15). On the contrary, Kaiso has also been suggested to be a potential tumor suppressor, which repressed the transcription of *MMP7*, *CCND1*, and *WNT11* genes involved in oncogenesis and metastasis (7, 9, 51). Functions of Kaiso are tightly related and significantly influenced by microenviromental factors (17). We found here that increase of the level of pT606-Kaiso or pan-14-3-3 (especially 14-3-3σ) expression could lead to Kaiso accumulation in the cytoplasm, deprive transcriptional function of Kaiso, and de-repression of Kaiso target gene *CDH1*. This indicates that Kaiso phosphorylation is involved in cancer development through de-repression of tumor related genes.

In conclusion, Kaiso can be phosphorylated by AKT1 at T606 within the very conservative motif RSS<u>T</u>IP in the cytoplasm. pT606-Kaiso can directly interact with 14-3-3 and P120ctn proteins and accumulated in the cytoplasm that consequently lead loss of Kaiso's function as a transcription repressor. The decreased phosphorylation of Kaiso may contribute to cancer development through downregulation of *CDH1* expression in the stomach and colon.

Acknowledgements

We gratefully acknowledge the critical reading of the manuscript by Dr. Jian-Ping Guo in Sun Yat-Sen University. This work is supported by National Natural Science Foundation of China (81402343).

Competing Interest

The authors declare that they have no conflict of interest.

Author contribution

D. Deng, W. Tian and S. Qin. designed research; W. Tian, S. Qin and B. Zhang performed research; D. Deng , W. Tian, L. Gu and J. Zhou analyzed data; D. Deng, W. Tian and S. Qin wrote the paper.

References

1. Daniel JM, Reynolds AB. (1999) The catenin p120(ctn) interacts with Kaiso, a novel BTB/POZ domain zinc finger transcription factor. *Mol Cell Biol* 19: 3614-3623.

2. Prokhortchouk A, Hendrich B, Jørgensen H, Ruzov A, Wilm M, Georgiev G, *et al.* (2001) The p120 catenin partner Kaiso is a DNA methylation-dependent transcriptional repressor. *Genes Dev* 15: 1613-1618.

3. Daniel JM, Spring CM, Crawford HC, Reynolds AB, Baig A. (2002) The p120(ctn)-binding partner Kaiso is a bi-modal DNA-binding protein that recognizes both a sequence-specific consensus and methylated CpG dinucleotides. *Nucleic Acids Res* 30: 2911-2919.

4. Yoon HG, Chan DW, Reynolds AB, Qin J, Wong J. (2003) N-CoR mediates DNA methylation-dependent repression through a methyl CpG binding protein Kaiso. *Mol Cell* 12: 723-734.

5. Rodova M, Kelly KF, VanSaun M, Daniel JM, Werle MJ. (2004) Regulation of the rapsyn promoter by kaiso and delta-catenin. *Mol Cell Biol* 24: 7188-7196.

6. Blattler A, Yao L, Wang Y, Ye Z, Jin VX, Farnham PJ. (2013) ZBTB33 binds unmethylated regions of the genome associated with actively expressed genes. *Epigenetics Chromatin* 6: 13.

7. Pozner A, Terooatea TW, Buck-Koehntop BA. (2016) Cell-specific Kaiso (ZBTB33) Regulation of Cell Cycle through Cyclin D1 and Cyclin E1. *J Biol Chem* 291: 24538-24550.

8. Robinson SC, Klobucar K, Pierre CC, Ansari A, Zhenilo S, Prokhortchouk E, et al. (2017) Kaiso differentially regulates components of the Notch signaling pathway in intestinal cells. *Cell Commun Signal* 15: 24.

9. Ogden SR, Wroblewski LE, Weydig C, Romero-Gallo J, O'Brien DP, Israel DA, *et al.* (2008) p120 and Kaiso regulate Helicobacter pylori-induced expression of matrix metalloproteinase-7. *Mol Biol Cell* 19: 4110-4121.

10. Dai SD, Wang Y, Zhang JY, Zhang D, Zhang PX, Jiang GY, *et al.* (2011) Upregulation of δ-catenin is associated with poor prognosis and enhances transcriptional activity through Kaiso in non-small-cell lung cancer. *Cancer Sci* 102: 95-103.

11. Buck-Koehntop BA, Stanfield RL, Ekiert DC, Martinez-Yamout MA, Dyson HJ, Wilson IA, Wright PE. (2012) Molecular basis for recognition of methylated and specific DNA sequences by the zinc finger protein Kaiso. Proc Natl Acad Sci U S A. Sep 18;109: 15229-15234

12. Lopes EC, Valls E, Figueroa ME, Mazur A, Meng FG, Chiosis G, Laird PW, Schreiber-Agus N, Greally JM, Prokhortchouk E, Melnick A. (2008) Kaiso contributes to DNA methylation-dependent silencing of tumor suppressor genes in colon cancer cell lines. Cancer Res. Sep 15;68 (18):7258-63

13. Jones J, Wang H, Zhou J, Hardy S, Turner T, Austin D, *et al.* (2012) Nuclear Kaiso indicates aggressive prostate cancers and promotes migration and invasiveness of prostate cancer cells. *Am J Pathol* 181: 1836-1846.

14. Jones J, Wang H, Karanam B, Theodore S, Dean-Colomb W, Welch DR, *et al.* (2014) Nuclear localization of Kaiso promotes the poorly differentiated phenotype and EMT in infiltrating ductal carcinomas. *Clin Exp Metastasis* 31: 497-510.

15. Prokhortchouk A, Sansom O, Selfridge J, Caballero IM, Salozhin S, Aithozhina D, *et al.* (2006) Kaiso-deficient mice show resistance to intestinal cancer. *Mol Cell Biol* 26: 199-208.

16. Iioka H, Doerner SK, Tamai K. (2009) Kaiso is a bimodal modulator for Wnt/beta-catenin signaling. *FEBS Lett* 583: 627-632.

17. Soubry A, van Hengel J, Parthoens E, Colpaert C, Van Marck E, Waltregny D, *et al.* (2005) Expression and nuclear location of the transcriptional repressor Kaiso is regulated by the tumor microenvironment. *Cancer Res* 65: 2224-2233.

18. Zhang PX, Wang Y, Liu Y, Jiang GY, Li QC, Wang EH. (2011) p120-catenin isoform 3 regulates subcellular localization of Kaiso and promotes invasion in lung cancer cells via a phosphorylation-dependent mechanism. *Int J Oncol* 38: 1625-1635.

19. Zhang L, Gallup M, Zlock L, Feeling Chen YT, Finkbeiner WE, McNamara NA. (2016) Cigarette Smoke Mediates Nuclear to Cytoplasmic Trafficking of Transcriptional Inhibitor Kaiso through MUC1 and P120-Catenin. *Am J Pathol*; 186: 3146-3159.

20. Moore, B. W. and Perez, V. J. (1967) Specific Acid Proteins in the Nervous System. *Physiological and Biochemical Aspects of Nervous Integration.* p343–359. New Jersey: Prentice-Hall. Englewood Cliffs

21. Dougherty MK, Morrison DK. (2004) Unlocking the code of 14-3-3. *J Cell Sci* 117: 1875-1884.

22. Coblitz B, Wu M, Shikano S, Li M. (2006) C-terminal binding: an expanded repertoire and function of 14-3-3 proteins. *FEBS Lett* 580: 1531-1535

23. Mori M, Vignaroli G, Botta M. (2013) Small molecules modulation of 14-3-3 protein-protein interactions. *Drug Discov Today Technol* 10: e541-547.

24. Wilker EW, Grant RA, Artim SC, Yaffe MB. (2005) A structural basis for 14-3-3 sigma functional specificity. *J Biol Chem* 280: 18891-18898.

25. Mackintosh C. (2004) Dynamic interactions between 14-3-3 proteins and phosphoproteins regulate diverse cellular processes. *Biochem J* 381: 329-342.

26. Mühlmann G, Ofner D, Zitt M, Müller HM, Maier H, Moser P, *et al.* (2010) 14-3-3 sigma and p53 expression in gastric cancer and its clinical applications. *Dis Markers* 29: 21-29.

27. Ko S, Kim JY, Jeong J, Lee JE, Yang WI, Jung WH. (2014) The role and regulatory mechanism of 14-3-3 sigma in human breast cancer. *J Breast Cancer* 17: 207-218.

28. Radhakrishnan VM, Putnam CW, Martinez JD. (2012) Activation of phosphatidylinositol 3-kinase (PI3K) and mitogen-activated protein kinase (MAPK) signaling and the consequent induction of

transformation by overexpressed 14-3-3γ protein require specific amino acids within 14-3-3γ N-terminal variable region II. *J Biol Chem* 287: 43300-43311.

29. Radhakrishnan VM, Martinez JD. (2010) 14-3-3gamma induces oncogenic transformation by stimulating MAP kinase and PI3K signaling. *PLoS One* 5: e11433.

30．Marullo M, Zuccato C, Mariotti C, Lahiri N, Tabrizi SJ, Donato SD(2010), Cattaneo E. Expressed Alu repeats as a novel, reliable tool for normalization of real-timequantitative RT-PCR data. GenomeBiology2010,11:R9

31. Samaddar S, Dutta A, Sinharoy S, Paul A, Bhattacharya A, Saha S, *et al.* (2013) Autophosphorylation of gatekeeper tyrosine by symbiosis receptor kinase. *FEBS Lett* 587: 2972-2979.

32. Lynes M, Narisawa S, Millán JL, Widmaier EP. (2011) Interactions between CD36 and global intestinal alkaline phosphatase in mouse small intestine and effects of high-fat diet. *Am J Physiol Regul Integr Comp Physiol* 301: R1738-1747.

33. Kinoshita E, Kinoshita-Kikuta E, Koike T. (2009) Separation and detection of large phosphoproteins using Phos-tag SDS-PAGE. *Nat Protoc* 4: 1513-1521

34. Margret B E, Elena N P, Jason R O (2007) GST Pull-down. CSH Protoc. Aug 1;2007:pdb.prot4757

35. Hisamoto K, Ohmichi M, Kurachi H, Hayakawa J, Kanda Y, Nishio Y, *et al.* (2001) Estrogen induces the Akt-dependent activation of endothelial nitric-oxide synthase in vascular endothelial cells. *J Biol Chem* 276: 3459-3467.

36. Ghandi M, Huang FW, Jane-Valbuena J, Kryukov GV, Lo CC, McDonald ER, *et al*. (2019) Next-generation characterization of the Cancer Cell Line Encyclopedia. *Nature* 569: 503-508.

37. Melé M, Ferreira PG, Reverter F, DeLuca DS, Monlong J, Sammeth M, *et al*. (2015) Human genomics. The human transcriptome across tissues and individuals. *Science* 348: 660-665.

38. Liu WS, Tian W, Zhou J, Gu LK, Deng DJ. (2020) The RNA transcriptome database was used to mine the candidate transcription factors of EZH2 gene in tumor cells. *Chinese J Biochem Mol Biol* 36: 665-673.

39. Rush J, Moritz A, Lee KA, Guo A, Goss VL, Spek EJ, *et al.* (2005) Immunoaffinity profiling of tyrosine phosphorylation in cancer cells. *Nat Biotechnol* 23: 94-101.

40. Hornbeck PV, Zhang B, Murray B, Kornhauser JM, Latham V, Skrzypek E. (2015) PhosphoSitePlus, 2014: mutations, PTMs and recalibrations. *Nucleic Acids Res*. 43: D512-520.

41. Navé BT, Ouwens M, Withers DJ, Alessi DR, Shepherd PR. (1999) Mammalian target of rapamycin is a direct target for protein kinase B: identification of a convergence point for opposing effects of insulin and amino-acid deficiency on protein translation. *Biochem J* 344(Pt 2): 427-431.

42. Salas TR, Reddy SA, Clifford JL, Davis RJ, Kikuchi A, Lippman SM, *et al.* (2003) Alleviating the suppression of glycogen synthase kinase-3beta by Akt leads to the phosphorylation of cAMP-response element-binding protein and its transactivation in intact cell nuclei. *J Biol Chem* 278: 41338-41346.

43. Fang D, Hawke D, Zheng Y, Xia Y, Meisenhelder J, Nika H, *et al.* (2007) Phosphorylation of beta-catenin by AKT promotes beta-catenin transcriptional activity. *J Biol Chem* 282: 11221-11229.

44. Horman S, Vertommen D, Heath R, Neumann D, Mouton V, Woods A, *et al.* (2006) Insulin antagonizes ischemia-induced Thr172 phosphorylation of AMP-activated protein kinase alpha-subunits in heart via hierarchical phosphorylation of Ser485/491. *J Biol Chem* 281: 5335-5340.

45. Hong F, Nguyen VA, Shen X, Kunos G, Gao B. (2000) Rapid activation of protein kinase B/Akt has a key role in antiapoptotic signaling during liver regeneration. *Biochem Biophys Res Commun* 279: 974-979.

46. Yaffe MB, Rittinger K, Volinia S, Caron PR, Aitken A, Leffers H, *et al.* (1997) The structural basis for 14-3-3:phosphopeptide binding specificity. *Cell* 91: 961-971.

47. Venhuizen JH, Jacobs FJC, Span PN, Zegers MM. (2019) P120 and E-cadherin: Double-edged swords in tumor metastasis. *Semin Cancer Biol* 60: 107-120.

48. del Valle-Pérez B, Casagolda D, Lugilde E, Valls G, Codina M, Dave N, *et al.* (2016) Wnt controls the transcriptional activity of Kaiso through CK1ε-dependent phosphorylation of p120-catenin. *J Cell Sci* 129: 873.

49. Yang H, Zhang Y, Zhao R, Wen YY, Fournier K, Wu HB, *et al.* (2006) Negative cell cycle regulator 14-3-3 sigma stabilizes p27 Kip1 by inhibiting the activity of PKB/Akt. *Oncogene* 25: 4585-4594.

50. Qin S, Zhang B, Tian W, Gu L, Lu Z, Deng D. (2015) Kaiso mainly locates in the nucleus in vivo and binds to methylated, but not hydroxymethylated DNA. *Chinese Journal of Cancer Research* 27: 148-155.

51. Kim SW, Park JI, Spring CM, Sater AK, Ji H, Otchere AA, *et al.* (2004) Non-canonical Wnt signals are modulated by the Kaiso transcriptional repressor and p120-catenin. *Nat Cell Biol* 6: 1212-1220.

Figures and Legends

Figure 1. Different phosphorylation states of Kaiso protein in the cytoplasm and nucleus of cells *in vitro* and *in vivo*.

(**A**) Different phosphorylation statuses of endogenous Kaiso in the cytoplasm and nucleus of MGC803 and HEK293T

cells in the Phos-tag and regular SDS-PAGE analyses. The purities of cytoplasmic and nuclear extracts were

respectively verified using anti-β-Tubulin and anti-Lamin B antibodies. Kaiso in both protein extracts with and

without the de-phosphorylation treatment by CIAP were showed. (**B**) The phosphorylation statuses of endogenous

cytoplasmic and nuclear Kaiso in MGC803 cells were validated using anti-phosphoSer/Thr/Tyr universal antibody. (**C**)

The phosphorylation statuses of GFP-Kaiso stably transfected MGC803 cells and the corresponding xenograft tissues

in the Phos-tag SDS-PAGE analysis.

Figure 2. AKT1 increases the phosphorylation of Kaiso at T606. (**A**) A conservative RSX$\underline{T}$XP motif within Kaiso of human and other species (**B**) After starvation overnight, treatments of Insulin (100 ng/mL), IL-6 (10 ng/mL) and fetal bovine serum (FBS, 1:10000 v/v) for 15 min increased the phosphorylation level of endogenous Kaiso in MGC803 cells. (**C**) Effects of AKT inhibitor MK2206 treatment (10 μmol/mL) for 30 min blocked the promotion of Insulin induced Kaiso phosphorylation as pAKT substrate in MGC803 cells. (**D**) *AKT1* overexpression at different doses increased the amount of phosphorylated AKT substrate (pAKT-sub) in Kaiso complexes immunoprecipitated by Kaiso antibody in MGC803 cells. (**E**) GST-Kaiso-wt recombinant treated with *AKT1* was identified by AKT substrate-specific antibody in an *in vitro* kinase assay and T606A mutation abolished AKT1's impact on Kaiso. (**F**) The T606-phosphorylation status of endogenous Kaiso in MGC803 and BGC823 with treatment of Insulin and its combination with AKT inhibitor MK2206 after starvation overnight. (**G**) The T606-phosphorylation status of endogenous Kaiso in MGC803 and BGC823 with *AKT1* overexpression after starvation overnight.

Figure 3. Subcellular location of endogenous pT606-Kaiso and total Kaiso. (A) Amounts of pT606-Kaiso and total

Kaiso in cytoplasmic and nuclear proteins (with and without de-phosphorylation treatment by CIAP for 30 min)

extracted from 4 cancer cell lines in Western blotting. (B) Locations of pT606-Kaiso and total Kaiso in the cytoplasm

and nucleus in an indirect immunofluorescence confocal analysis. (C) Locations of pT606-Kaiso and total Kaiso in a

representative gastric carcinoma and the paired normal tissues in IHC analysis. (D) Locations of pT606-Kaiso and

total Kaiso in human gastric tissues in the indirect immunofluorescence confocal analysis.

Figure 4. Kaiso interacted with 14-3-3 family members depending on Kaiso pT606. (A) After the plasmids of 14-3-3 family members and GST-Kaiso were co-transfected into MGC803 cells, GST-Kaiso pulled down various isoforms of 14-3-3 family, especially 14-3-3 σ. (B) Endogenous Kaiso immunoprecipitated pan-14-3-3 or 14-3-3σ protein immunoprecipitated endogenous Kaiso in MGC803 cell in Co-IP analysis. (C) T606A mutation of Kaiso concealed its interaction with pan-14-3-3 in MGC803 cells in Co-IP analysis.

Figure 5. 14-3-3 proteins promote the cytoplasmic accumulation of Kaiso in Kaiso T606 phosphorylation-dependent

manner. (**A**) The subcellular location of endogenous Kaiso in MGC803 cells with or without 14-3-3γ or 14-3-3σ

overexpression by indirect immunofluorescence staining assay. (**B**) Proportion of Kaiso in the nucleus and in

cytoplasm of MGC803 cells with and without 14-3-3γ or 14-3-3σ overexpression. (**C**) Western blot for detecting the

amounts of endogenous Kaiso in cytoplasm and nucleus protein in MGC803 cells with 14-3-3σ overexpression. (**D**)

Comparison of the levels of GFP-Kaiso (wild type) or T606A mutant in the cytoplasm and nucleus in MGC803 cell

with and without 14-3-3σ overexpression. Lamin B and β-Tublin were used as nuclear and cytoplasmic protein

controls.

Figure 6. 14-3-3σ promotes the interaction of Kaiso and p120ctn in the cytoplasm. (**A**) More Kaiso-p120 complex was immunoprecipitated by Kaiso antibody in MGC803 cells with 14-3-3σ overexpression in Co-IP assay. Total protein levels of p120ctn and 14-3-3σ in the lysate of MGC803 cells with *14-3-3σ* overexpression were displayed in the right side. (**B**) Alterations of interactions between Kaiso and p120ctn proteins in the cytoplasm and nucleus of these cells with *14-3-3σ* overexpression in the Co-IP assay using Kaiso antibody. (HC: Heavy chain of IgG). (**C**) The subcellular locations of Kaiso, p120ctn, and mCherry-14-3-3σ in MGC803 cells. (**D**) Western blotting images for detecting effect of *p120ctn* knockdown and *14-3-3σ* overexpression on distribution of Kaiso in the cytoplasm and nucleus of MGC803 cells. (**E**) The levels of total p120ctn and Kaiso proteins in the lysate of MGC803 cells with *14-3-3σ* overexpression and siRNAs knockdown of *p120ctn* expression.

Figure 7. Effects of 14-3-3σ on inhibition of Kaiso's target gene *CDH1* expression. (**A**) The Pearson correlation

coefficient between mRNA levels of 14-3-3 family members and Kaisos' target genes in public RNA-seq datasets

from Cancer Cell Line Encyclopedia (CCLE, 1156 cancer cell lines) and Genotype-Tissue Expression (GTEx, 11688

human normal tissues). (**B**) The level of *CDH1* in SGC7901 and MKN45 cells with or without wild type Kaiso and

14-3-3σ overexpression. (**C**) The level of the *CDH1* mRNA in SGC7901 and MKN45 cells with or without wild type

Kaiso and 14-3-3σ overexpression (**D**) The amounts of pT606-Kaiso, total Kaiso, 14-3-3 σ, and CDH1 proteins in

gastric carcinoma (T) and the paired normal tissues (N) from 4 patients by Western blotting

Figure S1. Characterizing the specificity of pT606-Kaiso polyclonal antibody. ELISA results for the specificity of

pT606-Kaiso and control antibodies against the pT606-Kaiso peptide (LSDRSSpTIPAM, the top chart) and for the

specificity of pT606-Kaiso and control antibodies for the nonphosphorylated control peptide (LSDRSSTIPAM, the

bottom chart).

Figure S2. The status of Kaiso protein modifications detected by LC/MS. Phosphorylation at the Thr606 residue in

the LSDRSS<u>T</u>IPAM motif was illustrated in the bottom chart in details. The image was adapted with graphs for Kaiso

modifications from the web site (www.phosphosite.org; Ref.:40).

Figure S3. AKT1 and 14-3-3 regulate the T606-phosphorylation and subcellular localization of endogenous Kaiso. **(A)**

Endogenous Kaiso in MGC803 cells immunoprecipitated by Kaiso antibody was identified by the antibody specific

for AKT substrate motif, and the immunoprecipitation by AKT substrate antibody was identified by antibody against

Kaiso. **(B)** After starvation overnight, overexpressed *AKT1* at different dose in HCT116 cells promoted the

phosphorylation of Kaiso in phos-tag SDS PAGE gel. **(C)** *AKT1* overexpression increased endogenous

Kaiso-14-3-3 interaction in MGC803 cells in Co-IP assay. **(D** and **E)** The T606-phosphorylation states of endogenous

Kaiso in the cytoplasm and nucleus of MGC803 and BGC823 cells with 14-3-3σ and 14-3-3γ overexpression,

respectively.

Figure S4. Correlation between the levels of *CDH1* and *14-3-3σ* (A) or *Kaiso* (B) mRNA in RNA-seq and cDNA

array datasets for human normal tissues in the Genotype-Tissue Expression (GTEx) projects and cancer cell lines in

Cancer Cell Line Encyclopedia (CCLE) project.

2 **Figure S5**. The amounts of pT606-Kaiso, total Kaiso, 14-3-3 σ, and CDH1 proteins in colon carcinoma (T) and the

3 paired normal tissues (N) from 4 patients by Western blotting