Paternal exposure to DDT programs breast cancer development in daughters via sperm

miRNA in a mouse model

Raquel Santana da Cruz[1], Hong Cao[1], Camile C. Fontelles[1], Apsra Nasir[1], Alexandra

Krebs[1], M. Idalia Cruz[1], Lu Jin[1] and Sonia de Assis[1]*

[1]Department of Oncology, Lombardi Comprehensive Cancer Center, Georgetown University,

Washington, DC, USA.

Corresponding author: Sonia de Assis, Ph.D., 3970 Reservoir Road, NW, The Research Building,

Room E410, Washington, DC 20057, USA; Phone: (202) 687-9518, Fax: (202) 687-7505, Email:

deassiss@georgetown.edu, ORCID: 0000-0001-5053-0614

16 **The authors declare they have no actual or potential competing financial interests**
17

Abstract

Background: DNA sequence accounts for the majority of disease heritability, including

cancer. However, it is becoming clear that environmentally-induced epigenetic inheritance

can also occur. Epidemiological studies have shown that maternal exposure to the pesticide

DDT in pregnancy is associated with increased breast cancer risk in women. Yet, the effects of

paternal exposure to this pesticide on the progeny's breast cancer development has not been

investigated.

Methods: Male mice (*c57bl/6*) were exposed to DDT or to a control-vehicle (CO) solution and

used for sperm collection or mating with unexposed females to produce the DDT or CO

daughters. In another experiment, normal mouse embryos (zygote stage) were injected with

miRNA-10b and implanted into surrogate mothers to produce miR-10b offspring. DDT

daughters or miRNA-10b females were used to study breast cancer development and metabolic

parameters. Paternal sperm was used for RNA-seq analysis and miRNA expression levels.

Results: Pre-conception paternal DDT exposure altered the sperm small non-coding RNA load,

with a notable increase in miRNA-10b levels. DDT offspring weighed less at birth and at

weaning, but became overweight and showed metabolic dysfunction in adulthood compared to

CO. DDT daughters also showed increased mammary tumorigenesis, developing more

aggressive tumors that grew faster than in CO. This tumor phenotype was linked to suppression

of the AMPK energy sensing pathway and mTOR activation in mammary tissues. Remarkably,

embryonic injection of miRNA-10b recapitulated the mammary gland and tumor phenotypes

observed in DDT daughters.

Conclusions: To our knowledge, this is the first report of an association between paternal DDT

exposure and breast cancer in offspring. Paternal DDT-induced programming of breast cancer

development in daughters is mechanistically linked to sperm miRNA-10b. The impact of DDT

and other endocrine disrupting chemicals on sperm and programming of breast and other

cancers in offspring needs to be evaluated in humans.

Introduction

Dichlorodiphenyltrichloroethane (DDT), one of the first synthetic organochlorine pesticides, was widely used in the United States (U.S.) from the 1940s to the 1970s and up to the 1980s in Europe (Knower KC 2014; SE 2013). While the use of DDT has been restricted in the U.S. and Europe for decades, developing countries with endemic malaria and other vector-borne diseases continue to use it (Cohn BA 2015; Kabasenche and Skinner 2014; Knower KC 2014). This pesticide has a prolonged half-life and is lipophilic, bio-accumulating in the food chain, and can still be detected in fatty tissues and circulation of animals and humans (Knower KC 2014). In the U.S., the highest concentrations of DDT are detected in minority populations and recent immigrants (Daniels et al. 2018; Ruiz et al. 2018). Like other organochlorines, DDT is an endocrine disruptor (Knower KC 2014; Kuiper et al. 1998) and it has been linked to metabolic syndrome (La Merrill M 2014), obesity (Skinner et al. 2013) and cancer (Cohn et al. 2007; Cohn BA 2015).

DNA sequence is responsible for the majority of disease heritability, including cancer. It is becoming clear, however, that epigenetic inheritance can also occur. At conception, parents not only contribute their genome but also transmit a molecular memory of past environmental experiences to their progeny (YW Chen Q, Duan W 2016; OJ 2012). This can result in offspring's predisposition to chronic diseases later in life (MK 2007). This environmentally-induced risk of diseases, including breast cancer, has been shown to be transmitted to the offspring via epigenetic mechanisms through both the female and male germ-lines (de Assis S 2012; Gapp K 2014; MK 2007). Although most of the evidence for this mode of disease inheritance comes from maternal exposures in pregnancy, we (da Cruz et al. 2018; da Cruz et al. 2019; Fontelles et al. 2016a; Fontelles et al. 2016b) and others (Caronne BR 2010; Guido et al.

2016; Ng SF 2010) have shown that paternal exposures in the pre-conception window are also

important in determining disease outcomes in the offspring.

Several recent published reports demonstrated that the small RNA load in paternal sperm can

transmit environmentally-induced phenotypes to the offspring (YM Chen Q, Cao Z, Li X, Zhang

Y, Shi J, Feng GH, Peng H, Zhang X, Zhang Y, Qian J, Duan E, Zhai Q, Zhou Q 2016; YW

Chen Q, Duan W 2016; Gapp K 2014; Sharma U 2016). Some of those studies implicate specific

classes of small RNA and both miRNAs and tRFs obtained from sperm can recapitulate the

effect of specific paternal exposures in offspring when injected into normal embryos following in

vitro fertilization (YM Chen Q, Cao Z, Li X, Zhang Y, Shi J, Feng GH, Peng H, Zhang X, Zhang

Y, Qian J, Duan E, Zhai Q, Zhou Q 2016; Gapp K 2014; Rodgers AB 2015; Sharma U 2016).

Because the mammary gland bud first emerges during fetal development, it has been proposed

that some breast cancers may originate *in utero* (Trichopoulos 1990) and multiple reports on

maternal exposure in pregnancy have offered support to this hypothesis (Cohn BA 2015; de

Assis S 2006; de Assis S 2012; de Oliveira Andrade et al. 2014). In line with that, a recent

epidemiological analysis using the Child Health and Development Studies cohort (CHDS)

showed that daughters of women exposed to high levels of DDT in pregnancy have increased

rates of breast cancer (Cohn BA 2015) and are more likely to be diagnosed with advanced stage

and HER-2 positive tumors. Other studies in the same cohort found that women exposed to this

pesticide before age 14 have increased breast cancer risk, with the highest risk if the exposure

occurred in early infancy (Cohn et al. 2007; Cohn et al. 2019).

While there is a clear link between maternal exposure to DDT and breast cancer in the next

generation, no studies to date have investigated the contribution of pre-conception paternal

exposure to this or other pesticides to offspring's breast cancer development. Here, we tested the

hypothesis that paternal exposure to DDT can alter the sperm non-coding RNAs and program

breast cancer development in daughters, using a mouse model of this disease. We also examined

whether paternal DDT-induced programming of breast cancer development in daughters is

mechanistically linked to sperm miRNAs.

Materials and Methods

DDT exposures and breeding: Animals were purchased from Taconic Biosciences. The *c57bl/6*

strain of mice was used in all experiments. Adult male mice (8 weeks of age) were either treated

with a DDT solution (1.7 mg/kg) or vehicle-control solution (vegetable oil) via oral gavage for

two weeks. The DDT solution used mimics the formulation of DDT before its ban in the U.S.:

77.2% p,p'-DDT and 22.8% o,p'-DDT as described before (La Merrill M 2014). Animals body

weight was monitored weekly. At the end of the exposure period, DDT-exposed and control

male mice were mated to unexposed females to generate the female offspring (**Fig. S1**). Female

mice were kept on a standard chow diet during the breeding period, for the extent of pregnancy

(21 days) and after giving birth. The birth weight of pups and number of pups per litter was

determined. To avoid litter-effect, pups were cross-fostered one day after birth. Pups from 2-3

dams were pooled and housed in a litter of 8-10 pups per nursing dam. All pups were weaned on

postnatal day (PND) 21. Offspring's body weight was recorded weekly. The female offspring of

control or DDT-exposed fathers were used to study body weight, metabolic function, and

mammary tumorigenesis and for mammary gland collection as described in the following

sections. The number of contributing fathers and female offspring used in each experiment is

listed in **Table S1**. All animal procedures were approved by the Georgetown University Animal

Care and Use Committee, and the experiments were performed following the National Institutes

of Health guidelines for the proper and humane use of animals in biomedical research.

DDT and Phenobarbital (PB) treatment: PB treatment is known to promote hepatic enzymes

activation, accelerating DDT metabolism and excretion in humans (Davies et al. 1969; Kwalick

1971) and animals (Alary et al. 1971; Lambert and Brodeur 1975, 1976). A sub-set of control-

vehicle or DDT exposed males were concomitantly treated with phenobarbital (20 mg/kg of

body weight, i.p.) or saline and used for sperm harvesting.

Mature spermatozoa collection and purification: Control and DDT-exposed males as well as

those treated with PB were euthanized and caudal epididymis dissected for sperm collection. The

epididymis was collected, punctured, and transferred to tissue culture dish containing M2 media

(M2 Medium-with HEPES, without penicillin and streptomycin, liquid, sterile-filtered, suitable

for mouse embryo, SIGMA, product #M7167) where it was incubated for 1 hour at 37°C. Sperm

samples were isolated and purified from somatic cells. Briefly, the samples were washed with

PBS, and then incubated with SCLB (somatic cell lysis buffer, 0.1% SDS, 0.5% TX-100 in

Diethylpyrocarbonate water) for 1 hour. SCLB was rinsed off with 2 washes of PBS and the

somatic cell-free purified spermatozoa sample pelleted and used for RNA extraction.

Measurement of DDT metabolites: Liver tissues from control, DDT and DDT-PB treated males

(n=3 pooled samples/group), collected within 24 h after last dose, were used to determine the

levels of DDT main metabolites. Measurements were performed commercially by Pacific Rim

Labs, a fully accredited laboratory providing analysis of DDT and other persistent organic

pollutants. Briefly, samples were fortified with each of the six 13C-DDT isotopes and then

extracted with hexane. The extract was columned on Florisil and analyzed by GC-MS/MS. The

instrument was calibrated with a five-point calibration covering the range of 0.2-400 ng/g or mL.

Body weight monitoring: DDT female offspring's body weight was measured at birth (n=8-17

litters/group) and at weaning and weekly after that (n=25/group). Birth weights were analyzed by

t-test and longitudinal body weight analyzed using two-way ANOVA (group, time), followed by

post-hoc analyses.

Small RNA-Seq Analysis: Total RNA was isolated from paternal sperm (n=4/group) using

Qiagen's miRNeasy extraction kit, according to the manufacturer's instructions. Small non-

coding RNA transcript libraries were constructed according to Illumina TrueSeq Small RNA

Pre-Kit. Indexed, paired-end sequencing libraries were prepared from quality total RNA (RIN $\geq$

8). For each library, 10 million reads (raw data) were generated by Illumina Hi-Seq 4000. The

raw reads were subject to adapter trimming and low quality filtering using Trimmomatic

program. The high quality clean reads were aligned on the mouse genome. Non-coding RNA

tags were mapped to the mouse genome (GRCm38/mm10 reference genome) in order to analyze

their expression and distribution on the genome. Small RNA tags were annotated with miRNA,

t-RNA and pi-RNA to miRBase, Ref-seq, GenBank and Rfam database using blastn with

standard parameters. To analyze the differential expression of miRNA between control and DDT

groups, miRNAs were normalized to TPM (Transcripts Per Kilobase Million). miRNAs with a P

value less than 0.05 were considered significant, with an appropriate correction for multiple

testing (Clarke et al. 2008).

Quantitative real time PCR validation (qRT-PCR): cDNA was synthesized from total RNA

samples using the TaqMan™ Advanced miRNA cDNA Synthesis Kit (Applied Biosystems).

PCR products were amplified from cDNA using TaqMan® Fast Advanced Master Mix and

sequence-specific primers from TaqMan Assays (Applied Biosystems). Fold change was

calculated from Ct values and the expression levels of specific miRNAs or genes were

determined by normalizing these values with the fold change values for appropriate endogenous

controls (miR-26a or *GAPDH*). Fold differences between the groups were analyzed using t-test

or two-way ANOVA (group, treatment), followed by post-hoc analyses.

Carcinogen-induced mammary tumorigenesis: Mammary tumors were induced in control and

DDT female offspring (n=25/group) by administration of MPA (medroxyprogesterone acetate,

15 mg, s.c.) at 6 weeks of age, followed by three weekly doses of 1mg of 7,12-

dimethylbenz[a]anthracene (DMBA) (Sigma, St. Louis, MO) dissolved in peanut oil by oral

gavage. This established model of breast cancer has been used by us and others (Apostoli et al.

2014; Fontelles et al. 2016a) . Mice were examined for mammary tumors by palpation once per

week, starting on week two after the last dose of DMBA and continue for a total of 20 weeks.

Tumor growth was measured using a caliper and the width and height of each tumor were

recorded. Other end-points for data analysis were latency to tumor appearance, the number of

animals with tumors (tumor incidence), and the number of tumors per animal (tumor

multiplicity). During follow-up, those animals in which tumor burden approximated 10% of total

body weight were euthanized, as required by our institution. Histopathology of tumors was

determined by our pathologist. All tumors included in our analyses were classified as mammary

carcinomas. Differences in tumor latency and multiplicity were analyzed by t-test (or

corresponding non-parametric test). Kaplan-Meier survival curves were used to compare

differences in tumor incidence, followed by the log-rank test. Tumor growth was analyzed using

two-way ANOVA (group, time), followed by post-hoc analyses.

Embryonic miRNA microinjections. To induce super-ovulation, female mice (*c57bl/6*) received

intra-peritoneal injections of 5 IU of pregnant mare's serum gonadotropin (PMSG). About 48

hours after the PMSG injection, mice were injected with 5 IU of human chorionic gonadotropin

(hCG). Super-ovulated females were mated with *c57bl/6* male mice. About 16 hours later,

fertilized eggs were harvested. Zygotes were injected with either synthetic miRNA-10b

(*mir*Vana®, ThermoFisher) diluted in TE solution (10mM Tris-HCl, pH7.4, 0.25 EDTA) or TE

solution only (to control for microjection). Each zygote received ~10 pL of TE or miRNA

preparation at 2ng/µL. Microinjections were confirmed by observing the swelling of the male

pro-nucleus. Injected zygotes were transferred into recipient female mice to produce offspring as

described before (Rodgers AB 2015). Briefly, recipient females were mated with vasectomized

males overnight and identified by copulation plug before the zygote transfer. Each recipient

female was implanted with 25-30 zygotes. Dams were housed in groups of two with free access

to food and water. Female offspring were used to study tumor growth, metabolic function and for

mammary tissue collection. Mature sperm cells contain approximately 20 femtograms of total

RNA, of which about 20% are small non-coding RNAs and 0.4% are miRNAs (Kawano M

2012). Thus, the amount of miRNA injected into zygotes did not exceed physiological levels.

Embryonic injections were performed at the Johns Hopkins Transgenic Core Facility. Resulting

offspring were subsequently transferred to our institution for follow-up studies.

Syngeneic orthotopic tumor implantation: Mouse EO771 breast cancer cells were cultured in

complete medium [Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10%

(v/v) FBS and 1% L-glutamine, at 37°C in 5% CO_2 (v/v)] to 80% confluence, trypsinized,

counted, checked for viability and re-suspended in PBS for injection into both 4[th] inguinal

mammary glands. A total of 1×10^6 cells were injected into each of the 4[th] inguinal mammary

gland fat pads of miRNA-10b females (n=6-8/group). Tumor growth was monitored for at least

six weeks.

Immunohistochemistry   ***analysis.***   Cell   proliferation   (Ki-67)   was   evaluated   by

immunohistochemistry in mammary tumors (n=5-7/group). Briefly, tissues were fixed in 10%

buffered formalin, embedded in paraffin, and sectioned (5 µm). Sections were deparaffinized

with xylene and rehydrated through a graded alcohol series. Antigen retrieval was performed by

immersing the tissue sections at 98°C for 40 minutes in 1X Diva Decloaker (Biocare). Tissue

sections were treated with 3% hydrogen peroxide and 10% normal goat serum for 10 minutes,

and were incubated with the primary antibody (**Table S2**), overnight at 4°C. After several

washes, sections were treated to the appropriate HRP labeled polymer for 30 min and DAB

chromagen (Dako) for 5 minutes. Slides were counterstained with Hematoxylin (Fisher, Harris

Modified Hematoxylin), blued in 1% ammonium hydroxide, dehydrated, and mounted with

Acrymount. The sections were photographed using an Olympus IX-71 Inverted Epifluorescence

microscope at 40x magnification. Proliferation index (Ki-67 staining) was determined by

immunoRatio, a plugin Image J software (National Institute of Health, Bethesda, MD, USA) to

quantify hematoxylin and DAB-stained cells. Results were analyzed by t-test (or corresponding

non-parametric test).

Analysis of cell apoptosis: Cell apoptosis analysis was conducted in mammary tumors (n=14-

17/group) by morphological detection. Tissues were fixed in neutral buffered 10% formalin and

stained with hematoxylin and eosin (H&E). Cells presenting loss of adhesion between adjacent

cells, cytoplasmic condensation and formation of apoptotic bodies were considered apoptotic as

described before (Elmore et al. 2016). The sections were photographed using a Olympus IX-71

Epifluorescence microscope at 40x magnification. Twenty areas were photographed randomly

and the number of apoptotic bodies counted. The apoptotic index was determined as the

percentage of apoptotic bodies per total number of cells. Images were evaluated with Image J

software (NIH, USA). Results were analyzed by t-test (or corresponding non-parametric test).

Mammary gland development: Inguinal mammary glands (4[th] pair, n=5-16/group) collected on

post-natal day 50 (PND 50) were stretched onto a slide, placed in a fixative solution and stained

with a carmine aluminum solution (Sigma Chemical Co.) as previously described (de Assis et al.

2010). Whole mounts were examined under the microscope (AmScope) and ductal elongation

and number of TEBs (undifferentiated structure considered to be the targets of malignant

transformation), as previously described (de Assis et al. 2010). In addition, whole-mount slides

were digitized (Olympus SZX12 Stereomicroscope) and analyzed using the ImageJ software

(National Institute of Health, Bethesda, MD, USA): The area and integrated density of a

manually drawn perimeter around the mammary epithelial tree of each whole-mount were

measured. The average area as well integrated density measurements are reported for each group.

Differences were statistically tested using t test (or corresponding non-parametric test).

Analysis of protein expression: Protein levels were assessed by western blot on PND50

mammary tissues (n=6-7 group) and tumors (n=6/group). Total protein was extracted

from mammary tissues using RIPA buffer with Halt™ Protease Inhibitor Cocktail (Thermo

Fisher). Ten micrograms of protein extracts were resolved on a 4-12% denaturing

polyacrylamide gel (SDS-PAGE). Proteins were transferred using the iBlot® 7-Minute Blotting

System (Invitrogen, USA) and blocked with 5% non-fat dry milk for 1h at room

temperature. Membranes were incubated with the specific primary antibodies (for antibody

specifications and dilutions see **Table S2**) at 4°C overnight. After several washes, the

membranes were incubated with horseradish peroxidase (HRP)-conjugated secondary antibody

at room temperature for 1h. Membranes were developed using the Chemiluminescent HRP

antibody detection reagent (Denville Scientific Inc., USA), and exposed to blot imaging systems

(Amersham[TM] Imager 600, GE Healthcare Life Sciences). Optical density of the bands was

quantified using Quantity-one software (BIO-RAD, USA). To control for equal protein loading,

expression of the proteins of interest was normalized to the β-actin or β-tubulin signal.

Metabolic Parameters: Glucose tolerance test (GTT) was performed after overnight fasting in

mice. Glucose (0.75 U/kg body weight) was administered intraperitoneally and tail blood

samples were collected at 0, 15, 30 and 60 minutes after injection (n=8/group). Circulating

insulin levels (n=6-7/group) were measured using an ELISA kit (Ultra Mouse insulin EIA Kit,

Crystal Chem). Differences in GTT were analyzed using two-way ANOVA (group, time),

followed by post-hoc analyses. Insulin levels were statistically tested using t test (or

corresponding non-parametric test).

Results

DDT offspring have altered body weight gain and metabolic function

Male mice were either exposed to a vehicle-control or an environmentally relevant dose of DDT

(as described in the Material and Methods section) and mated with unexposed females to

generate the offspring (**Fig. S1**). The phenotypes in offspring from control and DDT-exposed

male mice were then examined. No differences in gender distribution or litter size were

observed between offspring born from DDT and control males (**Table S3**). However, paternal

DDT exposure was associated with a significant decrease in offspring's birthweight and weaning

weights (P<0.05, P<0.01 **Fig. 1a-b**). After weaning, DDT daughters underwent a catch-up

growth period and by 6 weeks of age no differences in body weight were observed between the

groups. In fact, DDT females became significantly heavier than controls by 28-weeks of age

(P<0.05, P<0.01, **Fig. 1c**).   They also showed metabolic dysfunction with higher circulating

levels of insulin and impaired glucose tolerance compared to controls in adulthood (**Fig**. **2a-b**,

P<0.05).

DDT daughters have accelerated breast cancer growth

Maternal DDT exposure has been linked to a near 4-fold increase in breast cancer in daughters

(Cohn BA 2015). Low birthweight, a phenotype observed in offspring DDT-exposed fathers,

has also been linked with higher risk for breast cancer in humans (Mellemkjaer L 2003;

Sanderson et al. 1996) and animal models (da Cruz et al. 2018; Fernandez-Twinn DS 2010).

Thus, we studied mammary cancer development in control and DDT daughters using the well-

established (Apostoli et al. 2014; Fontelles et al. 2016a) DMBA-induced mouse model of breast

cancer. Tumor growth was significantly increased in DDT daughters compared to controls

(*P*<0.001, **Fig. 3a**), with tumors weighing significantly more at the end of the monitoring period

(*P*<0.01, **Fig. 3b**). The incidence of mammary tumors in the DDT offspring was higher and

tumor latency slightly shorter than in controls but results did not reach statistical significance

(**Fig. 3c-d**).

Given the increased mammary tumor growth in the DDT daughters, we assessed the cell

proliferation and apoptosis rates in their tumors. We found a slight increase in cell proliferation

rates (*P*<0.1, **Fig. S2**) as well as a significant decrease in apoptosis rates in tumors of DDT

daughters compared to controls (*P*<0.05, **Fig. S2**).

DDT daughters' mammary gland morphological development is mostly normal

Because DDT daughters showed accelerated breast cancer growth, we next analyzed the effects

paternal DDT on offspring's mammary morphological development in post-natal day (PND)50

inguinal mammary glands. We assessed number of terminal end buds (TEB), ductal elongation

and epithelial area/density. No major morphological changes were detected, except for a slight

increase in mammary ductal elongation in DDT daughters compared to controls (**Fig. S3**).

DDT daughters' mammary tissues have rewired nutrient sensing mechanisms

We previously showed that low birth weight is linked to accelerated tumor growth, a phenotype

associated with suppression of the nutrient/energy sensing AMPK pathway (**Fig. 4b**) in

mammary tissues and tumors (da Cruz et al. 2018). In support of that, we found that both AMPK

expression and phosphorylation levels were decreased in DDT daughters' mammary tissues. We

also detected an increase in expression and activity of mTOR and its down-stream target, 4E-

BP1 (**Fig. 4b, Fig. S4a-f**). Similar patterns of mTOR and 4E-BP1 expression and activity were

also found in tumors of DDT daughters, although results were not statistically significant (**Fig.**

4c, Fig. S4g-l).

Pre-conception paternal DDT exposure alters the sperm non-coding RNA content

To determine the impact of DDT exposure on paternal sperm, we performed RNA-seq analysis

to assess the small RNA patterns in mature cauda sperm of control and DDT-exposed males. The

distribution of the major small RNA species detected is in agreement with previous reports (YM

Chen Q, Cao Z, Li X, Zhang Y, Shi J, Feng GH, Peng H, Zhang X, Zhang Y, Qian J, Duan E,

Zhai Q, Zhou Q 2016; Sharma U 2016), with tRNA-derived fragments (tRFs) being the most

abundant sperm small RNA sub-type, followed by miRNAs and piRNAs in both groups (**Fig.**

5a). Interestingly, sperm of DDT-exposed males showed higher abundance of miRNAs (6.7%)

compare to controls (1.4%) (**Fig. 5a-b**). A total of 16 miRNAs were differentially expressed

(nine down- and seven up-regulated) in sperm of DDT-exposed mice compared to controls (**Fig.**

5c). Of the seven miRNAs upregulated in sperm of DDT-exposed males, miRNA-10b had the

highest surge, representing close to 60% of the overall miRNA increase (**Fig. 5d**).

To establish whether upregulation of miRNA-10b in sperm was, in fact, induced by DDT

exposure, we treated DDT-exposed mice with phenobarbital (PB), a drug known to promote

hepatic enzymes activation and accelerate DDT metabolism and excretion in humans (Davies et

al. 1969; Kwalick 1971) and animals (Alary et al. 1971; Lambert and Brodeur 1975, 1976).

Indeed, we observed an increase in hepatic phase I enzymes (P<0.001, **Fig.5e**) and a reduction in

DDT metabolites in males treated with both DDT and PB (**Table S4**). Further, compared to mice

exposed to DDT only, miRNA-10b was decreased in sperm of mice treated with both DDT and

PB with levels reversing to those observed in controls (P<0.05, **Fig. 5f**).

miRNA-10b embryonic injection replicates DDT daughters' phenotypes

miRNA-10b along with other members of the miRNA-10 family play critical roles in embryonic

development and cell differentiation through the regulation of Homeobox (*Hox*) genes (Tehler et

al. 2011). These genes regulate organogenesis including development of the mammary gland

(Carroll and Capecchi 2015; Chen and Capecchi 1999; Garin et al. 2006).

Next, we examined whether miRNA-10b was responsible for the phenotypes observed in DDT

offspring using embryonic (zygote) microinjections. While no major changes in mammary gland

morphology were detected (**Fig. S3**), female mice derived from the miR-10b zygote injections

(miR-10b females from now on) showed mammary tissue suppression of AMPK and increased

expression and activity of mTOR and down-stream effector, 4E-BP1, compared to controls (**Fig.**

6a, **Fig. S5**), a phenotype also observed in mammary glands of DDT daughters. More

importantly, we also found increased orthotopic (EO771 cells) mammary tumor development in

miR-10b females, which showed both increased tumor growth and higher tumor weights

compared to controls (P<0.05, P<0.01, **Fig. 6b-c**). Survival after tumor implantation was also

worse in miRNA-10b females (**Fig. 6d**, P<0.1).   Further, embryonic miR-10b injection

phenocopied the metabolic dysfunction of DDT daughters, with adult miR-10b females showing

hyperinsulinemia (P<0.01) compared to controls (**Fig. 6e**).

Discussion

Paternal environmental exposures have been shown to modulate offspring's health phenotypes

via sperm non-coding RNAs (YM Chen Q, Cao Z, Li X, Zhang Y, Shi J, Feng GH, Peng H,

Zhang X, Zhang Y, Qian J, Duan E, Zhai Q, Zhou Q 2016; Rodgers AB 2015; Sharma U 2016).

Here, we show that pre-conception paternal exposure to the pesticide DDT modulates the sperm

non-coding RNA load, particularly miRNAs, and programs female offspring's breast cancer

development in a pre-clinical animal model. We also found that miRNA-10b, the miRNA with

the highest increase in sperm of DDT-exposed fathers, can recapitulate phenotypes observed in

DDT daughters including alterations in the mammary tissue AMPK/mTOR axis and accelerated

breast tumor growth. These findings lend support to the notion that predisposition to diseases

such as cancer can be determined by factors other than genetic background and suggest that it

could be transmitted through sperm miRNAs. We have shown before that paternal obesity,

malnutrition or modulation of nutritional patterns can increase breast and other types of cancer in

the progeny (da Cruz et al. 2018; da Cruz et al. 2019; de Assis S 2012; Fontelles et al. 2016a).

This study further implies that environmentally-induced non-genetic or epigenetic predisposition

to cancer can occur and should be evaluated in human studies.

Previously published reports using the CHDS cohort showed that daughters of women exposed

to DDT in pregnancy have increased rates of breast cancer (Cohn BA 2015) and are more likely

to be diagnosed with advanced stage tumors. Likewise, we found that paternal DDT exposure in

the pre-conception window leads to increased breast cancer growth in daughters in a mouse

model. Our new observations and our previously published data (da Cruz et al. 2018; da Cruz et

al. 2019; Fontelles et al. 2016a; Fontelles et al. 2016b) indicate that paternal exposures are as

important as maternal ones in programming of breast and other cancers in offspring. This overlap

between the effects of maternal and paternal exposure on offspring's breast cancer phenotypes

suggest a common underlying mechanism that needs to be further investigated.

While no studies in humans have directly investigated the link between paternal DDT exposure

and breast cancer in their descendants, some evidence for a role of related paternal exposures

exists. For instance, an association between father's exposure to pesticides and a number of

childhood cancers in offspring has been reported (Bailey H 2014; Greenop KR 2013).

Epidemiologic studies show that paternal exposure to DDT and other pesticides is linked to low

birthweight (Robledo CA 2015) and our study supports those findings. Low birthweight is more

prevalent in minority populations (Lumpkins and Saint Onge 2017; Wang and Beydoun 2007)

which are more likely to be exposed to environmental toxicants such as DDT (CDC Upated

Tables-January 2019(2009); Daniels et al. 2018; Ruiz et al. 2018) and to develop more

aggressive breast tumors (Kohler et al. 2015).

It has been proposed that during the normal fertilization process, sperm small RNAs are

delivered to the oocyte and modulate the transcriptome during the first few cell divisions, setting

a signaling cascade that can impact embryonic development (Gapp and Miska 2016) and recent

reports support this view (YW Chen Q, Duan W 2016; Sharma U 2016). In line with that,

embryonic injection of miR-10b, the miRNA with the highest surge in sperm of DDT-exposed

males, replicates most of the major phenotypes observed in DDT daughters including higher

circulating levels of insulin, altered AMPK/mTOR signaling in mammary tissues and increased

tumor growth. While we have focused on miRNA-10b given its reported role in embryonic

development (Tehler et al. 2011), it is possible that the full phenotype observed in DDT

offspring reflects the interaction of that miRNA with other miRNAs and, likely, other non-

coding RNAs in sperm of DDT-exposed males. It is important to note, however, that other

paternal environmental exposures —including malnutrition and obesity— previously shown to

increase breast cancer development in daughters (da Cruz et al. 2018; Fontelles et al. 2016a;

Fontelles et al. 2016b) also lead to an increase in miRNA-10b levels in sperm (YM Chen Q, Cao

Z, Li X, Zhang Y, Shi J, Feng GH, Peng H, Zhang X, Zhang Y, Qian J, Duan E, Zhai Q, Zhou Q

2016; Cropley et al. 2016; da Cruz et al. 2018; Fontelles et al. 2016b), indicating that a common

mechanism underlying the effects of different paternal exposures on offspring phenotypes likely

exists and should be examined.

MicroRNA-10b along with other members of the miRNA-10 family play critical roles in

embryonic development and cell differentiation through the regulation of Homeobox (*Hox*)

genes (Tehler et al. 2011). These genes regulate organogenesis including development of the

mammary gland and other skin-derived tissues (Carroll and Capecchi 2015; Chen and Capecchi

1999; Garin et al. 2006). However, whether alterations in *Hox* genes in the embryonic mammary

gland play a mechanistic role in the increased breast cancer development in daughters of DDT-

exposed fathers needs to be examined in futures studies.

We previously showed that female offspring born small (due to paternal malnutrition) have re-

wiring of the nutrient sensing AMPK/mTOR pathway both in normal mammary tissues as well

as in tumors. Similarly, we found that DDT daughters, who are also born small, have alterations

in the mammary AMPK/mTOR axis, further suggesting a mutual mechanism between different

types of ancestral paternal exposures. Whether alterations in nutrient sensing mechanisms are

programmed by pre-conception exposure or are a consequence of the metabolic dysfunction

observed in DDT daughters remains to be determined. However, the same phenotype was

present in mammary glands of miRNA-10b females indicating that dysregulation of the

AMPK/mTOR axis is likely initiated through sperm miRNAs after fertilization but mechanisms

for how it impacts mammary development need to be further elucidated.

It also remains to be determined whether alterations in the AMPK/mTOR axis in mammary

tissues of DDT daughters and miR-10b females are functionally responsible for the observed

increase in tumor growth. In normal cells, proliferation is coupled to nutrient availability to

ensure proper cell growth and division (Viollet et al. 2009). The AMPK pathway functions as a

cellular energy sensor mechanism and provides a link between nutrient supplies and cell growth

and metabolism. This pathway controls mTOR activity and associated energy-consuming

biosynthetic activities such as protein synthesis (Shackelford and Shaw 2009). Not surprisingly,

suppression of AMPK and increased mTOR activity (due to loss of LKB1, the major AMPK

kinase) is responsible for the inheritable Peutz-Jeghers syndrome, in which patients have an 18-

fold increase in cancer risk, including breast cancer (Sebbagh et al. 2011). Alterations in

AMPK/mTOR axis have been shown to lead metabolic reprogramming in cancer cells (Faubert

et al. 2014; Mossmann et al. 2018; Saxton and Sabatini 2017) and our published studies confirm

those findings (da Cruz et al. 2018). Given its document role in cancer, the local mammary

changes in AMPK/mTOR signaling are likely mechanistic linked to accelerated tumor growth,

however, it will also be important to examined the role of systemic alterations in enhancing

breast cancer development in both DDT daughters and miRNA-10b females.

Although we clearly show that pre-conception paternal DDT exposure modulates the male

germline and programs female offspring's breast cancer development, our study has some

limitations. First, we only used one environmentally relevant dose of DDT in our investigations

and exposure was restricted to a short period of time in adulthood. In follow-up studies, it would

be important to examine whether other environmentally relevant doses of DDT would elicit the

same effects on the paternal germline and offspring's phenotypes. It would also be critical to

understand the impact of DDT exposure on sperm at different developmental windows and

possible differences stemming from longer exposure periods. Although embryonic injection of

miR-10b on its own replicated most of the effects of paternal DDT on offspring's mammary

cancer development, it will be important to test whether other miRNAs and small RNA subtypes

also play a role. Finally, due to age-specific requirements, we had to use different models of

breast cancer in studies involving DDT daughters (DMBA model) and miRNA-10b females

(orthotopic implantation). In both cases, we observed accelerated tumor growth, however, it is

likely that DMBA-induced primary tumors better mimic the three stages of cancer development

(initiation, promotion and progression) compared to orthotopic implantation models (Liu et al.

2015).

Conclusions: Our study shows that environmental toxicants such as DDT alter the male

germline non-coding RNAs and increase cancer susceptibility in the progeny. This could have

important implications for human health: According to published studies and the Center for

Disease Control and Prevention (CDC), minorities such as African Americans and Mexican-

Americans and recent immigrants have the highest circulating concentrations of this pesticide in

the U.S. (CDC 2009; Daniels et al. 2018; Ruiz et al. 2018). In addition, DDT is still used in

developing countries and Americans who could have been affected by paternal exposure to DDT

before its ban (Cohn BA 2015) are still alive today and reaching an age when the incidence of

breast and other cancers increases. While our study focused on DDT exposure, our findings are

likely to be relevant to other commonly used pesticides and flame retardants which, like DDT,

have anti-androgenic properties (Kharlyngdoh et al. 2018; Kortenkamp and Faust 2010; Stoker et

al. 2005).

Acknowledgments: We thank the following Lombardi Cancer Center Shared Resources (SR)

for their assistance: Animal Model SR, Histopathology & Tissue SR, Microscopy and Imaging

SR and Genomics & Epigenomics SR. We also thank Mr. Chip Hawkins and Dr. Roger Reeves

from the Johns Hopkins Transgenic Core Facility for technical support. Finally, we thank Dr.

Anni Warri from the University of Turku for insightful comments and suggestions to the

manuscript. This study was supported in part by The American Cancer Society (RSG-16-203-01-

NEC, Research Scholar Grant to S. de Assis), The Georgetown Environmental Initiative (pilot

fund to S. de Assis), and the National Institutes of Health (1P30-CA51008; Lombardi

Comprehensive Cancer Center Support Grant to Louis Weiner and CCSG pilot fund to S. de

Assis).

The small RNA-seq (small RNA) data has been deposited in GEO (Gene Expression Omnibus)

database with accession code pending.

Author contributions: S.D.A. conceived the study, oversaw the research and wrote the

manuscript; R.S. supervised the animal work and tissue collection, performed molecular analysis

and wrote the manuscript; H.C and C.C.F performed animal work. A.N. and A.K. processed

tissues and/or helped perform molecular analysis; M.I.C. assisted with animal studies; L.J.

performed RNA-seq data.

504
505

References

Alary JG, Guay P, Brodeur J. 1971. Effect of phenobarbital pretreatment on the metabolism of ddt in the rat and the bovine. Toxicol Appl Pharmacol 18:457-468.

Apostoli AJ, Skelhorne-Gross GE, Rubino RE, Peterson NT, Di Lena MA, Schneider MM, et al. 2014. Loss of pparγ expression in mammary secretory epithelial cells creates a pro-breast tumorigenic environment. Int J Cancer 134:1055-1066.

Bailey H FL, Infante-Rivard C, Glass DC, Miligi L, Dockerty JD, Lightfoot T, Clavel J, Roman E, Spector LG, Kaatsch P, Metayer C, Magnani C, Milne E, Polychronopoulou S, Simpson J, Rudant J, Sidi V, Rondelli R, Orsi L, Kang A, Petridou E, Schüz J. 2014. Parental occupational pesticide exposure and the risk of childhood leukemia in the offspring: Findings from the childhood leukemia international consortium. International Journal of Cancer 135:2157-2172.

Caronne BR FL, Habib N, Shea JM, Hart CE, Li R, Bock C, Li C, Gu H, Zamore PD, Meissner A, Weng Z, Hofmann HA, Friedman N, Rando OJ. 2010. Paternally induced transgenerational environment reprogramming of metabolic gene expression in mammals. Cell 143:1084-1096.

Carroll LS, Capecchi MR. 2015. Hoxc8 initiates an ectopic mammary program by regulating fgf10 and tbx3 expression and wnt/beta-catenin signaling. Development 142:4056-4067.

CDC. Upated Tables-January 2019(2009). The fourth national report on human exposure to environmental chemicals, 2009. (the Fourth Report, 2009).

Chen F, Capecchi MR. 1999. Paralogous mouse hox genes, hoxa9, hoxb9, and hoxd9, function together to control development of the mammary gland in response to pregnancy. Proc Natl Acad Sci U S A 96:541-546.

Chen Q YM, Cao Z, Li X, Zhang Y, Shi J, Feng GH, Peng H, Zhang X, Zhang Y, Qian J, Duan E, Zhai Q, Zhou Q. 2016. Sperm tsrnas contribute to intergenerational inheritance of an acquired metabolic disorder. Science 351:397-400.

Chen Q YW, Duan W. 2016. Epigenetic inheritance of acquired traits through sperm rnas and sperm rna modifications. Nature Reviews Genetics 17.

Clarke R, Ressom HW, Wang A, Xuan J, Liu MC, Gehan EA, et al. 2008. The properties of high-dimensional data spaces: Implications for exploring gene and protein expression data. Nat Rev Cancer 8:37-49.

Cohn BA, Wolff MS, Cirillo PM, Sholtz RI. 2007. Ddt and breast cancer in young women: New data on the significance of age at exposure. Environ Health Perspect 115:1406-1414.

Cohn BA, Cirillo PM, Terry MB. 2019. Ddt and breast cancer: Prospective study of induction time and susceptibility windows. J Natl Cancer Inst.

Cohn BA LMM, Krigbaum NY, Yeh G, Park JS, Zimmermann L, Cirillo PM. 2015. Ddt exposure in utero and breast cancer. Journal of Clinical Endocrinology and Metabolism 100:2865-2872.

Cropley JE, Eaton SA, Aiken A, Young PE, Giannoulatou E, Ho JWK, et al. 2016. Male-lineage transmission of an acquired metabolic phenotype induced by grand-paternal obesity. Mol Metab 5:699-708.

da Cruz RS, Carney EJ, Clarke J, Cao H, Cruz MI, Benitez C, et al. 2018. Paternal malnutrition programs breast cancer risk and tumor metabolism in offspring. Breast Cancer Res 20.

da Cruz RS, Clarke J, Curi AC, Al-Yawar A, Jin L, Baird A, et al. 2019. Parental obesity programs pancreatic cancer development in offspring. Endocr Relat Cancer.

Daniels SI, Chambers JC, Sanchez SS, La Merrill MA, Hubbard AE, Macherone A, et al. 2018. Elevated levels of organochlorine pesticides in south asian immigrants are associated with an

increased risk of diabetes. J Endocr Soc 2:832-841.
Davies JE, Edmundson WF, Carter CH, Barquet A. 1969. Effect of anticonvulsant drugs on
dicophane (d.D.T.) residues in man. Lancet 2:7-9.
de Assis S, Warri A, Cruz MI, Hilakivi-Clarke L. 2010. Changes in mammary gland morphology
and breast cancer risk in rats. J Vis Exp.
de Assis S KG, Hilakivi-Clarke L. 2006. High birth weight increases mammary tumorigenesis in
rats. International Journal of Cancer 119:1537-1546.
de Assis S WA, Cruz MI, Laja O, Tian Y, Zhang B, Wang Y, Huang T, Hilakivi-Clarke L. 2012.
High-fat or ethinyl-oestradiol intake during pregnancy increases mammary cancer risk in several
generations of offspring. Nature Communications 3.
de Oliveira Andrade F, Fontelles CC, Rosim MP, de Oliveira TF, de Melo Loureiro AP,
Mancini-Filho J, et al. 2014. Exposure to lard-based high-fat diet during fetal and lactation
periods modifies breast cancer susceptibility in adulthood in rats. J Nutr Biochem 25:613-622.
Elmore SA, Dixon D, Hailey JR, Harada T, Herbert RA, Maronpot RR, et al. 2016.
Recommendations from the inhand apoptosis/necrosis working group. Toxicol Pathol 44:173-
188.
Faubert B, Vincent EE, Griss T, Samborska B, Izreig S, Svensson RU, et al. 2014. Loss of the
tumor suppressor lkb1 promotes metabolic reprogramming of cancer cells via hif-1alpha. Proc
Natl Acad Sci U S A 111:2554-2559.
Fernandez-Twinn DS ES, Martin-Gronert MS, Terry-Adkins J, Wayman AP, Warner MJ, Luan
JA, Gusterson BA, Ozanne SE. 2010. Poor early growth and excessive adult calorie intake
independently and additively affect mitogenic signaling and increase mammary tumor
susceptibility. Carcinogenesis 31:1873-1881.
Fontelles CC, Carney E, Clarke J, Nguyen NM, Yin C, Jin L, et al. 2016a. Paternal overweight is
associated with increased breast cancer risk in daughters in a mouse model. Sci Rep 6:28602.
Fontelles CC, Guido LN, Rosim MP, Andrade Fde O, Jin L, Inchauspe J, et al. 2016b. Paternal
programming of breast cancer risk in daughters in a rat model: Opposing effects of animal- and
plant-based high-fat diets. Breast Cancer Res 18:71.
Gapp K, Miska EA. 2016. Trna fragments: Novel players in intergenerational inheritance. Cell
Res 26:395-396.
Gapp K JA, Sarkies P, Bohacek J, Pelzcar P, Prados J, Farinelli L, Miska E, Mansuy IM. 2014.
Implication of sperm rnas in transgenerational inheritence of the effects of early trauma in mice.
Nature Neuroscience 17:667-669.
Garin E, Lemieux M, Coulombe Y, Robinson GW, Jeannotte L. 2006. Stromal hoxa5 function
controls the growth and differentiation of mammary alveolar epithelium. Dev Dyn 235:1858-
1871.
Greenop KR PS, Bailey HD, Fritschi L, Attia J, Scott RJ, Glass DC, de Klerk NH, Alvaro F,
Armstrong BK, Milne E. 2013. Exposure to pesticides and the risk of childhood brain tumors.
Cancer Causes & Control 24:1269-1278.
Guido LN, Fontelles CC, Rosim MP, Pires VC, Cozzolino SM, Castro IA, et al. 2016. Paternal
selenium deficiency but not supplementation during preconception alters mammary gland
development and 7,12-dimethylbenz[a]anthracene-induced mammary carcinogenesis in female
rat offspring. Int J Cancer 139:1873-1882.
Kabasenche WP, Skinner MK. 2014. Ddt, epigenetic harm, and transgenerational environmental
justice. Environ Health 13:62.
Kawano M KH, Grandjean V, Kiani J, Rassoulzadegan M. 2012. Novel small noncoding rnas in

mouse spermatazoa, zygotes and early embryos. PLoS One 7.

Kharlyngdoh JB, Pradhan A, Olsson PE. 2018. Androgen receptor modulation following
combination exposure to brominated flame-retardants. Sci Rep 8:4843.

Knower KC TS, Leung YK, Ho SM, Clyne CD. 2014. Endocrine disruption of the epigenome: A
breast cancer link. Endorcrine Related Cancer 21:T33-55.

Kohler BA, Sherman RL, Howlader N, Jemal A, Ryerson AB, Henry KA, et al. 2015. Annual
report to the nation on the status of cancer, 1975-2011, featuring incidence of breast cancer
subtypes by race/ethnicity, poverty, and state. J Natl Cancer Inst 107:djv048.

Kortenkamp A, Faust M. 2010. Combined exposures to anti-androgenic chemicals: Steps
towards cumulative risk assessment. Int J Androl 33:463-474.

Kuiper GG, Lemmen JG, Carlsson B, Corton JC, Safe SH, van der Saag PT, et al. 1998.
Interaction of estrogenic chemicals and phytoestrogens with estrogen receptor beta.
Endocrinology 139:4252-4263.

Kwalick DS. 1971. Anticonvulsants and ddt residues. JAMA 215:120-121.

La Merrill M KE, Moshier E, Lindtner C, La Frano MR, Newman JW, Buettner C. 2014.
Perinatal exposure of mice to the pesticide ddt impairs energy expenditure and metabolism in
adult female offspring. PLoS One 9.

Lambert G, Brodeur J. 1975. [the effect of pretreatment with enzyme inducers on the acute
toxicity of ddt (1, 1-bis-(p-chlorophenyl)-2,2,2-trichloroethane) in rats]. Rev Can Biol 34:121-
129.

Lambert G, Brodeur J. 1976. [influence of various enzyme inductors and combinations of
inductors on ddt residue elimination in rats]. Rev Can Biol 35:33-41.

Liu Y, Yin T, Feng Y, Cona MM, Huang G, Liu J, et al. 2015. Mammalian models of chemically
induced primary malignancies exploitable for imaging-based preclinical theragnostic research.
Quant Imaging Med Surg 5:708-729.

Lumpkins CY, Saint Onge JM. 2017. Reducing low birth weight among african americans in the
midwest: A look at how faith-based organizations are poised to inform and influence health
communication on the developmental origins of health and disease (dohad). Healthcare (Basel)
5.

Mellemkjaer L OM, Sørensen HT, Thulstrup AM, Olsen J, Olsen JH. 2003. Birth weight and risk
of early-onset breast cancer (denmark). Cancer Causes Control 14:61-64.

MK JRS. 2007. Environmental epigenomics and disease susceptibility. Nature Review Genetics
8:253-262.

Mossmann D, Park S, Hall MN. 2018. Mtor signalling and cellular metabolism are mutual
determinants in cancer. Nat Rev Cancer 18:744-757.

Ng SF LR, Barres R, Owens JA, Morris MJ. 2010. Chronic high-fat diet in fathers programs β-
cell dysfunction in female rat offspring. Nature 467:963-966.

OJ R. 2012. Daddy issues: Paternal effects on phenotype. Cell 151:702-708.

Robledo CA YE, Mendola P, Sundaram R, Maisog J, Sweeney AM, Barr DB, Louis GMB.
2015. Preconception maternal and paternal exposure to persistent organic pollutants and birth
size: The life study. Environmental Health Perspectives 123:88-94.

Rodgers AB MC, Leu NA, Bale TL. 2015. Transgenerational epigenetic programming via sperm
microrna recapitulates effects of paternal stress. Proceedings of the National Academy of
Sciences of the United States of America 112:13699-13704.

Ruiz D, Becerra M, Jagai JS, Ard K, Sargis RM. 2018. Disparities in environmental exposures to
endocrine-disrupting chemicals and diabetes risk in vulnerable populations. Diabetes Care

41:193-205.

Sanderson M, Williams MA, Malone KE, Stanford JL, Emanuel I, White E, et al. 1996. Perinatal
factors and risk of breast cancer. Epidemiology 7:34-37.

Saxton RA, Sabatini DM. 2017. Mtor signaling in growth, metabolism, and disease. Cell
168:960-976.

SE MMF. 2013. Endocrine disruptors and the breast: Early life effects and later life disease.
Journal of Mammargy Gland Biology and Neoplasia 18:43-61.

Sebbagh M, Olschwang S, Santoni MJ, Borg JP. 2011. The lkb1 complex-ampk pathway: The
tree that hides the forest. Fam Cancer 10:415-424.

Shackelford DB, Shaw RJ. 2009. The lkb1-ampk pathway: Metabolism and growth control in
tumour suppression. Nat Rev Cancer 9:563-575.

Sharma U CC, Shea JM, Boskovic A, Derr AG, Bing XY, Belleannee C, Kucukural A, Serra
RW, Sun F, Song L, Caronne BR, Ricci EP, Li XZ, Fauquier L, Moore MJ, Sullivan R, Mello
CC, Garber M, Rando OJ. 2016. Biogenesis and function of trna fragments during sperm
maturation and fertilization in mammals. Science 351:391-396.

Skinner MK, Manikkam M, Tracey R, Guerrero-Bosagna C, Haque M, Nilsson EE. 2013.
Ancestral dichlorodiphenyltrichloroethane (ddt) exposure promotes epigenetic transgenerational
inheritance of obesity. BMC Med 11:228.

Stoker TE, Cooper RL, Lambright CS, Wilson VS, Furr J, Gray LE. 2005. In vivo and in vitro
anti-androgenic effects of de-71, a commercial polybrominated diphenyl ether (pbde) mixture.
Toxicol Appl Pharmacol 207:78-88.

Tehler D, Hoyland-Kroghsbo NM, Lund AH. 2011. The mir-10 microrna precursor family. RNA
Biol 8:728-734.

Trichopoulos D. 1990. Hypothesis: Does breast cancer originate in utero? Lancet 335:939-940.

Viollet B, Athea Y, Mounier R, Guigas B, Zarrinpashneh E, Horman S, et al. 2009. Ampk:
Lessons from transgenic and knockout animals. Front Biosci (Landmark Ed) 14:19-44.

Wang Y, Beydoun MA. 2007. The obesity epidemic in the united states--gender, age,
socioeconomic, racial/ethnic, and geographic characteristics: A systematic review and meta-
regression analysis. Epidemiol Rev 29:6-28.

Figure Legends

Fig-1. Paternal DDT exposure programs offspring's body weight. Body weight (grams) at (a) birth (n=8-17 litter/group), at (b) weaning and in (c) adulthood [longitudinal body weight from 6 to 30 weeks of age (n=25/group)] in control (CO) and DDT offspring. All data are mean ± SEM. $*p < 0.05$, $**p < 0.01$ by t-test (birthweight and weaning) or repeated measures ANOVA (longitudinal body weight).

Fig-2. Paternal DDT exposure programs metabolic dysfunction in daughters. (a) Serum insulin levels (n=6-7/group) in adult control (CO) and DDT female offspring; (b) Glucose tolerance test (GTT) in adult control and DDT female offspring (n=8/group). All data are mean ± SEM. $*p < 0.05$ by t-test (serum insulin levels) or repeated measures ANOVA (GTT).

Fig-3. Paternal DDT exposure programs breast cancer development in daughters. Carcinogen-induced mammary tumorigenesis in control (CO) and DDT female offspring: (a) Mammary tumor volume, (b) tumor weight (grams), (c) tumor incidence and (d) tumor latency (n=25/group). Tumor incidence is shown as percentage of animals with tumors. All other data are mean ± SEM. $*p < 0.05$, $**p < 0.01$ by t-test (tumor weight and tumor latency), repeated measures ANOVA (mammary tumor growth) or Kaplan-Meier analysis followed by log-rank test (tumor incidence).

Fig-4. Paternal DDT exposure programs the AMPK/mTOR signaling pathway in DDT daughters' mammary tissues. (a) Simplified schematic representation of the pathway; (b-c) Representative western blots showing protein expression (phosphorylated and total) levels of members of the AMPK/mTOR axis in (b) normal mammary tissues and in (c) tumors (n=6-7/group) of control (CO) and DDT female offspring. Protein expression levels were normalized by either β-actin or β-tubulin. Densitometry data is shown in Fig. S4.

Fig-5. DDT exposure reprograms the sperm non-coding RNA load. (a) Small RNA subtype distribution and (b) miRNA abundance in sperm of DDT-exposed and control (CO) males (n=4/group) assessed by RNA-seq; (c) Heat-map showing differentially expressed miRNAs in sperm from DDT-exposed and CO males; (d) miR-10b as a fraction of all upregulated miRNAs in sperm of DDT-exposed males; (e) Expression levels of Cyp2B10 assessed by q-PCR in liver tissues of CO and DDT-exposed male mice treated with phenobarbital (PB, 20mg/kg) or vehicle (n=4-5/group); (f) Expression levels of miRNA-10b, assessed by q-PCR, in sperm of control or DDT-exposed male mice treated with PB or vehicle (n=8-20/group). Expression levels of Cyp2B10 and miRNA-10b are relative to GAPDH and miR-26a, respectively. All data are mean ± SEM. $*p < 0.05$, $***p < 0.001$, $****p < 0.0001$ by two-way ANOVA.

Fig-6. Embryonic (zygote stage) injection of miRNA-10b recapitulate the phenotypes observed in DDT daughters. (a) Representative western blots showing protein expression (phosphorylated and total) levels of members of the AMPK/mTOR axis in normal mammary tissues of control (CO) and miRNA-10b females (Protein expression levels were normalized by β-actin,

n=5/group. Densitometry data is shown in Fig. S5); (b-d) Orthotopic tumor growth (EO771 cells, n=6-8/group) in adult CO and miRNA-10b females [(b) tumor volume, (c) tumor weight (grams) and (d) survival]. (e) Serum insulin levels in adult control (CO) and miR-10b females (n=6/group). Survival is shown as percentage of animals still alive at a given time; All other data are mean ± SEM. *p<0.05, **p<0.01 by t-test (insulin levels, tumor weights, repeated measures ANOVA (mammary tumor volume) or Kaplan-Meier analysis followed by log-rank test (survival).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

SUPPLEMENTARY INFORMATION

Supplementary Figures

Figure S1

Figure S2

Figure S3

Figure S4

Figure S5

Supplementary Figure Legends

Fig-S1. Experimental design: C57BL/6 male mice were exposed to DDT (1.7mg/kg body weight) by oral gavage for two weeks. DDT or control-vehicle (CO)-exposed males were mated with unexposed female mice to generate the offspring. To avoid litter-effect, pups were cross-fostered one day after birth. Pups from 2-3 dams were pooled and housed in a litter of 8-10 pups per nursing dam. Pups were weaned from mothers at 21 days of age.

Fig-S2. Cell proliferation is increased and apoptosis rates decreased in mammary tumors of DDT female offspring. Representative images and quantification (percentage of the positively stained nuclear area) of proliferation staining (ki67; a-c, arrows, n=6/group) and apoptosis (d-f, morphological assessment, H&E sections, arrows, n=14-17/group) in control (CO) and DDT tumors. All images captured at 40x. Values are mean ± SEM. *P<0.05 by t-test.

Fig-S3. No major changes in mammary gland morphology on PND50 in DDT or miRNA-10b female offspring. (a,b) Representative photomicrography of mammary gland whole mount (6.25x) and digitized image of the whole mount. (c,g) Number of TEBs; (d,h) Epithelial elongation (distance between lymph node and the end of the mammary epithelial tree); (e,i) Mammary epithelial area measurement (MEA); (f,j) Branching density values; (n=5-16/group) in control (CO), DDT or miRNA-10b females. All data are mean ± SEM. PND, post-natal day.

Fig-S4. Paternal DDT exposure programs the AMPK/mTOR signaling pathway in DDT offspring's mammary tissues. (a-l) Quantification of protein expression levels of members of the AMPK/mTOR axis in (a-f) normal mammary tissues and in (g-l) tumors (n=6-7/group) of control (CO) and DDT offspring. Protein expression levels were normalized by either β-actin or β-tubulin. Optical density of individual bands was quantified using Quantity-one software. Values are mean ± SEM. *p<0.05, **p<0.01, ***p<0.01 by t-test.

Fig-S5. Embryonic injection of miR-10b recapitulate the DDT-induced programming of the AMPK/mTOR signaling pathway in mammary tissues. (a-f) Quantification of protein expression levels of members of the AMPK/mTOR axis in normal mammary tissues (n=5/group). Protein expression levels were normalized by β-actin. Optical density of individual bands was quantified using Quantity-one software. Values are mean ± SEM. ***p<0.01, ****p<0.01 by t-test.

Supplementary Tables

Table S1: Number of offspring and number of contributing fathers per experiment.

Samples	Assay	Group	Father	Female
Mammary gland	Terminal End Buds	**CO**	2	12
		DDT	3	16
	Western Blot	**CO**	2	6
		DDT	2	6-7
Mammary tumor	Apoptosis	**CO**	6	14
		DDT	9	17
	Ki67	**CO**	5	6
		DDT	5	6
	Tumorigenesis	**CO**	5	25
		DDT	8	25
	Western Blot	**CO**	3	6-7
		DDT	4	6
Plasma	ELISA	**CO**	1	6
		DDT	3	7

Table S2 - List of Antibodies

Antibody	Application	Dilution	Description	Supplier
AMPKα	Western Blot	1:1000	Rabbit polyclonal	Cell Signaling #2532
Anti-Ki67	IHC	1:7	Rabbit monoclonal	BIOCARE #CRM325
mTOR	Western Blot	1:1000	Rabbit monoclonal	Cell Signaling#2983
Phospho-AMPKα	Western Blot	1:1000	Mouse monoclonal	Cell Signaling #2535
Phospho-mTOR	Western Blot	1:1000	Rabbit monoclonal	Cell Signaling#2971
Phospho-4E-BP1	Western Blot	1:1000	Rabbit monoclonal	Cell Signaling#2855
4E-BP1	Western Blot	1:1000	Rabbit monoclonal	Cell Signaling#9452
β-Actin	Western Blot	1:1000	Rabbit monoclonal	Cell Signaling #8457
β-Tubulin	Western Blot	1:1000	Mouse monoclonal	Thermo Fisher Scientific #MA5-16308
Anti-mouse IgG, HRP	Western Blot	1:1000	Goat	Cell Signaling # 7076S
Anti-rabbit IgG, HRP	Western Blot	1:1000	Horse	Cell Signaling # 7074S

Table S3. Gender distribution (proportion) and number of pups per litter in control and DDT groups

Group	Female	Male	Number of pups per liter (mean±SEM)
CO	0.53	0.47	8.7±0.37
DDT	0.55	0.45	8.2±0.29

Table S4 – Levels of DDT metabolites in liver of mice treated with vehicle-control, DDT or DDT+ phenobarbital (**PB**)

DDT metabolite	Group (concentration, ng/g)		
	CO	**DDT**	**DDT- PB**
o,p' DDE	ND	ND	ND
p,p DDE	ND	0.6	ND
o,p DDD	ND	ND	ND
p,p DDD	ND	0.3	ND

Levels of DDT metabolites in liver (n=3 pooled samples/group) of control (CO), DDT (1.7/mg/kg) and DDT+phenobarbital (1.7mg+20mg/kg; DDT-PB) treated male mice measured by GC-MS/MS 24h after the last treatment. ND, non-detectable (readings below assay detection limits and not distinguishable from the blank control).