Genetic Diversity and Population Structure in the Ryukyu Flying Fox Inferred from Remote Sampling in the Yaeyama Archipelago

Yuto Taki[1], Christian E. Vincenot[2,3*], Yu Sato[1,4] and Miho Inoue-Murayama[1,5*]

[1] Wildlife Research Center, Kyoto University, Kyoto, Japan

[2] Department of Social Informatics, Kyoto University, Kyoto, Japan

[3] Island Bat Research Group (IBRG), Yoshida-honmachi, Sakyo-ku, Kyoto, Japan

[4] Royal (Dick) School of Veterinary Studies and the Roslin Institute, University of Edinburgh, Edinburgh, United Kingdom

[5] Wildlife Genome Collaborative Research Group, National Institute for Environmental Studies, Tsukuba, Japan

*Correspondance:

E-mail: mmurayama@wrc.kyoto-u.ac.jp (MI-M)

vincenot@i.kyoto-u.ac.jp (CEV)

Keywords:

Inbreeding; microsatellite; mtDNA; Faeces; Pteropus dasymallus; Fruit bat

Running title:

Genetic diversity in Yaeyama flying fox

Abstract

The Ryukyu flying fox (*Pteropus dasymallus*) is distributed throughout the island chain spanning across southern Japan, Taiwan, and possibly the Philippines. Although *P. dasymallus* is listed as VU (vulnerable) in the IUCN Red List, only few genetic works have been conducted to support its conservation. In this study we analyzed 19 markers (mtDNA haplotypes and 18 microsatellite markers) to evaluate genetic diversity and investigate the genetic structure of this species.

mtDNA analysis was conducted with 142 DNA remote samples, mostly from feces, and wing tissues collected on eight islands (Miyako, Ishigaki, Kohama, Kuroshima, Hateruma, Taketomi, Iriomote, Yonaguni). 39 haplotypes were identified in 526bp of the control region, and haplotype network showed no clear genetic structure.

Microsatellite analysis was also conducted with 155 samples collected on six islands (Miyako, Ishigaki, Kohama, Taketomi, Iriomote, Yonaguni). It showed that the Yonaguni population exhibits low genetic diversity, high inbreeding, and clear genetic differentiation from other populations. Gene flow between Ishigaki and Miyako through small stepstone islands might be preventing inbreeding of the Miyako population.

We provide for the first time indirect proof of long-distance inter-island dispersal in the Ryukyu flying fox and revealed genetic diversity, gene flow and genetic differentiation among populations of the archipelago. These results will be useful for delineating conservation units and designing specific conservation policies for each island based on metapopulation genetic structure.

Introduction

The Chiroptera represent a speciose and highly threatened group of mammals [1]. In Japan, 37 of the 122 mammal species are bats, yet academic research on this order has been lacking and misdirected towards the least threatened species [2]. The country features significantly high endemicity with endemic bats most endangered while at the same time most poorly studied (ibid.).

Flying foxes, i.e. paleotropical fruit bats belonging to the *Pteropus* genus, serve important ecosystem functions as pollinators [3] and seed dispersers [4], and play a

disproportionally large ecological role in maintaining forest structure and biodiversity [5]. Their seed dispersing capacity relies on their ability to carry large seeds across remarkably large foraging ranges. For instance, the Mauritian flying fox *Pteropus* niger was reported to fly up to 92 km in a single night [6], while some other large *Pteropus* species have demonstrated an ability to migrate between islands [7]. This is particularly critical as many paleotropical islands have lost their megafauna [8], leaving only flying foxes to perform large seed dispersal [9]. Nonetheless, their service to forested insular ecosystems is easily disrupted by demographic declines, as flying foxes cease to function as seed dispersers long before they become rare [9]. This reckoning is all the more alarming as flying foxes are arguably the most endangered group of bats worldwide and are most threatened on islands [10]. The intense pressure that they face is best epitomized by the regular conflicts and mass fatalities observed in Australia [11-14] and the notorious mass culling campaigns inflicted to the remaining population of *P. niger* by the Mauritian government [10, 15-17].

The Ryukyu flying fox (*Pteropus dasymallus*) is distributed across the Ryukyu archipelago in Japan, two small islands of Taiwan, and possibly in the Philippines [18]. *P. dasymallus* mainly eats fruits, nectar, and sometimes leaves, and plays an important role in pollination and seed dispersal [19]. *P. dasymallus* is listed as Vulnerable in the IUCN Red List [20] and has been the object of serious conservation concerns, due to the recent discovery of unreported threat factors [21], conflicts with farmers and culling which have resulted in important fatality [18, 21-22]. Low likeability and lack of support for conservation further warrant increased attention [23].

Considering the largely insular distribution range of these bats, genetic information such as diversity, gene flow and genetic differentiation between islands is important for their conservation. Indeed, genetic analyses on another species, *Pteropus mariannus*, for instance revealed genetic structure and gene flow between islands, and suggested new subspecies classification and conservation units [24]. Yet, tissue sample collection is tedious in the case of solitary, tree-dwelling fruit bats. On the other hand, non-invasive DNA extraction from faeces also potentially poses technical challenges due to the small size and consistency of the faeces coupled with heavy concentration in phenols—a known PCR inhibitor [25]. Hence, to this date, few genetic studies have been conducted on *P. dasymallus* and no solid knowledge exists on long-range dispersal capabilities and

population structure. In this study, we try to remedy this situation by focusing on the Yaeyama archipelago, which hosts the most extensive part of the Ryukyu flying fox population and is inhabited by one allopatric subspecies, the Yaeyama flying fox *P.d. yayeyamae*. mtDNA haplotype and microsatellite analysis of *P. d. yayeyamae* were conducted based on remote DNA sampling using mostly feces and feeding marks. This work provided the first indirect evidence of gene flow between islands and revealed the genetic diversity and structure among the eight major islands of the chain.

Materials and methods

Samples

DNA was recovered from various direct and indirect sources, namely feces, tissue, feeding marks of fruits or flowers, blood and hair. Feces and fruit samples were collected from 2012 to 2019 (Fig 1) by selection of appropriate sites (mostly foraging sites, but also a few roosts) based on experience. We avoided picking several samples from the same sites, except on islands on which population density was so low that only few sites were available at the time of collection. Feces and fruit samples were stored in either silica beads or 99% EtOH. Tissue samples, on the other hand, were obtained on Ishigaki island using the standard procedure for bats, which involved capturing *P. dasymallus* individuals at night using mistnets and taking one 3mm wing biopsy from the plagiopatagium or between the 4th and 5th finger in the distal part of the chiropatagium, depending on wing condition (see detailed protocol used in [25, 30]).

In total, 135 samples from 8 islands (feces: $n = 95$, tissue: $n = 17$, fruits: $n = 23$) were used for mtDNA sequencing while 155 samples collected in 6 islands (feces: $n = 99$, tissue: $n = 8$, feeding marks: $n = 32$, hair: $n = 15$, blood: $n = 1$) were used for microsatellite analysis. The number of analyzed individuals for each island is shown in Table S1. DNA extraction was conducted with QIAGEN DNeasy Blood and Tissue Kit (QIAGEN) for tissue samples and QIAamp DNA Stool Mini Kit (QIAGEN) for fecal samples.

mtDNA haplotype analysis

A part of control region of mtDNA was amplified through PCR. ppM01F (5'-accagaaaaggggarcaacc-3') and ppmtCR-RS2(5'-caagcatcccccaaaaatta-3') [25] were used as primers and PCR System 9700 (GeneAmp) was used as a thermal cycler. The PCR conditions were: 95°C for 2 min; 40-45 cycles at 95°C for 30s, 50-55 °C for 30s, 74°C for 1 min; then a 10 min final extension at 74°C. PCR products were purified with High Pure PCR Product Purification Kit (Roche). After sequencing reaction, 526bp of control region was sequenced by ABI PRISM 3130xl Genetic Analyzer (Applied Biosystems). SNPs were detected through sequence alignment by MEGA7 [26]. Haplotypes were identified based on the SNPs. Phylogenetic analysis was conducted by MEGA7 [26] and the haplotype network [27] was constructed by PopART [28]. Also, haplotype diversity (h) and haplotype richness (hr) were calculated by Contrib [29].

Microsatellite analysis

18 microsatellite loci (Accession number: LC506191-LC506193, LC506193, LC506197-LC506198, LC506200-LC506202, LC506205-LC506209, LC506212-LC506213, LC506220, LC506223) which we developed specifically for *P. dasymallus* [30] were amplified through PCR using a PCR System 9700 Thermal Cycler (GeneAmp). Forward primers were synthesized with an M13 tag sequence (5'-GTTGTAAAACGACGGCCAGT-3') for fluorescent labeling. PCR was conducted in a final volume of 10 µl, containing 1 µl DNA, 5 µl Multiplex PCR Master Mix (QIAGEN), 0.2 µM of M13-tailed forward primer, reverse primer and a M13 fluorescent primer labeled with FAM, NED, or HEX, and 0.1 µg of T4 gene 32 Protein (Nippon Gene, Tokyo, JPN). The PCR conditions were: 94°C for 5 min; 45 cycles at 94°C for 30s, 60 °C for 45s, 72°C for 45s; then 8 cycles for M13 at 94 °C for 30s, 53 °C for 45s, 72°C for 45s, and a 10 min final extension at 72°C. Amplicon size was measured using an ABI PRISM 3130xl Genetic Analyzer (Applied Biosystems), and genotypes were scored by eye with Peak Scanner Software (Applied Biosystems)..

Genetic diversity analysis and Principal Coordinate Analysis (PCoA) were conducted by GenAlEx (version) [31-32]. Also, we investigated genetic structure with STRUCUTE [33]. Finally, *Fst* was calculated by GENEPOP on the Web [34-35] to check genetic differentiation and geneflow between islands. Taketomi was included in only PCoA

because the number of analyzed individuals was not enough for the other analyses.

Results

mtDNA haplotype analysis

45 SNPs were detected in 526 bp sequences (including 35 bp which is a part of neighboring translated gene) of 135 samples, and 39 haplotypes (RFF1~RFF39, Accession number: LC528174-LC528212) were defined. The SNPs of each haplotype are shown in Fig S1, and the number of samples for each haplotype on each island is shown in Table S2.

Haplotype network suggested that haplotypes from each island did not form any clusters and no clear genetic structure was detected (Fig S2). However, in the phylogenetic tree by maximum likelihood method, Miyako population seemed closer to Ishigaki population (Fig S3).

Haplotype distribution among islands revealed that 14 haplotypes out of 39 were shared between multiple islands (Fig 1). While some haplotypes (RFF14, RFF36) were widely shared from the east end to the west end of the distribution, others were only found in certain areas. RFF16 and RFF21 were only found in Iriomote and Yonaguni (the west area), while RFF7, RFF24, RFF30, RFF35, and RFF38 were only found in Miyako and Ishigaki (the east area).

The number of haplotypes (Nh), the number of unique haplotypes ($unique$), haplotype diversity (h), and haplotype richness (hr) of 4 islands are shown in Table 1. The h of Ishigaki and Iriomote were 0.96 and 1.00, while that of Miyako and Yonaguni were 0.69 and 0.61, respectively. Also, the hr of Ishigaki and Iriomote were 7.51 and 9.00, while that of Miyako and Yonaguni were 3.77 and 2.79, respectively. Genetic diversity was higher in Ishigaki and Iriomote than Miyako, and the diversity of Yonaguni was the lowest.

Microsatellite analysis

The number of analyzed individuals (n), observed heterozygosity (Ho), expected

heterozygosity (*He*), and inbreeding coefficient (*F*) of each island are shown in Table 2. *Ho* of Ishigaki, Kohama, Iriomote, Miyako were 0.666, 0.650, 0.631, 0.512, respectively. On the other hand, *Ho* of Yonaguni was 0.388 and *F* was 0.355. Yonaguni population had low genetic diversity and showed high level of inbreeding compared to the other islands. PCoA shows genetic distance between individuals, and there were 3 clusters: Yonaguni cluster, center cluster (Ishigaki, Taketomi, Kohama, Iriomote and a part of Miyako), and Miyako cluster (Fig 2).

In STRUCTURE, LnP(D) suggested that K=4 provides the best model (Fig S4), and the latter is shown in Fig S5. Ishigaki, Kohama, Iriomote and a part of Miyako populations had the same genetic composition, while the rest of Miyako and Yonaguni populations had each different genetic structure.

Fst between Yonaguni and the other islands were generally high (Table 3), and it shows Yonaguni population is genetically differentiated from the other islands' populations.

Discussion

Genetic diversity and its differentiations

Genetic diversity was high in populations of the central area (Ishigaki, Taketomi, Kohama, and Iriomote) and the lowest in Yonaguni population. Miyako population exhibited a medium level of genetic diversity. Comparing the *Ho* of each island (Table 2) to that of wild population of the notoriously endangered *Pteropus rodricensis* (*Ho* = 0.5718) [36], Yonaguni (*Ho* = 0.388) had much lower value, while that of Miyako (*Ho* = 0.512) was slightly lower. The other islands' were, however, higher (*Ho* = 0.631~0.666). Also, inbreeding coefficient was high in Yonaguni population. This indicates that genetic diversity has been getting low due to inbreeding depression in Yonaguni population.

In mtDNA haplotype analysis, no clear genetic differentiations between islands was detected. However, in microsatellite analysis, populations of 6 islands were genetically divided into 3 groups: Miyako group (Miyako), the Central group (Ishigaki, Taketomi, Kohama, and Iriomote), and Yonaguni group (Yonaguni). Populations in the central group had the same genetic composition, and Miyako group was partly differentiated from them.

Yonaguni group was highly differentiated from the other 2 groups.

Gene flow between islands

The main habitats of the Yaeyama flying fox are Ishigaki island and Iriomote island (Central group) and populations in these two have higher genetic diversity than other islands. Other islands of the Central group might be keeping their diversity because of the gene flow linking them with the Ishigaki and Iriomote populations.

The Miyako group is geographically far from the Central group, but there are some islands such as Tarama between Ishigaki and Miyako. There could be gene flow between the two groups through these islands, and that might be why Miyako group was not differentiated completely from the Central group (Fig 2, Table 3) and had a middle level of genetic diversity. Also, Miyako and Ishigaki had the lowest Fst value (0.0221). This could be just because of active gene flow between Ishigaki and Miyako through islands between them. However, it might also be explained by the establishment of the Miyako population through recent migration from Ishigaki, which would explain the high ratio of RFF30 in Miyako in mtDNA haplotype analysis, assuming that the founders had that haplotype. This is also supported by the mtDNA phylogenetic tree (Fig S3), in which the Miyako population seemed closer to Ishigaki population. Our dataset, however, does not allow to conclude on whether or not the colonization of Miyako effectively occurred recently and engendered a founder effect, or if the haplotype composition of Miyako is representative of a genetic drift resulting from strong population decline.

As regards Yonaguni, there is no steppingstone between this island and the Central group. In a research on the endangered Bonin flying fox *Pteropus pselaphon*, the other Japanese *Pteropus* species, genetic diversity of an isolated island was similar to that of Yonaguni [37]. Since Yonaguni lies far away from its closest islands, it can be considered geographically isolated and migration can safely be assumed to happen only on special occasions such as typhoons, which regularly sweep the region westwards. This is supported by our results, which show that Yonaguni genetically differentiated from the other populations, and that is causing low genetic diversity due to inbreeding depression.

Conservation implications

The observation of gene flow between islands provides the first indirect evidence that long-distance inter-island dispersal is occurring in the Ryukyu flying fox. This indicates that this species is capable of long flights and thus that individual activity may not be assumed to be limited to a small range. This confirms results of ongoing VHF and GPS tracking studies (Vincenot C.E., unpub. data). Furthermore, our results show that island populations may be connected through flight capability and thus generate demographic and genetic interdependence, which need to be resolved to inform conservation policies. Considering the results of our genetic research, the Yaeyama subspecies of the Ryukyu flying fox could be divided into 3 conservation units: Miyako unit (populations in Miyako islands), the central unit (populations in the center of distribution including Ishigaki, Taketomi, Kohama, and Iriomote populations), and Yonaguni unit (Yonaguni islands) (Fig 3). The central unit has many individuals and high genetic diversity, and gene flow between the central unit and Miyako unit might be preventing extinction of the Miyako unit. However, the Yonaguni unit is isolated from other populations and could be highly endangered. Recovery of genetic diversity is crucial for population viability of the Yonaguni unit. Particular care should be exerted to reduce pressure on this population by reducing threat factors (see [18,21]).

In conclusion, we offer the present results for consideration by the authorities and advocate for the design of conservation strategies, which are still lacking despite our previous calls for action. We propose that outer populations, such as the one on Yonaguni, be the object of focused attention and protection measures, as they seem most at risk of stochastic demographic collapse and inbreeding depression.

Ethical statements

Samples were collected under permission granted to C.E. Vincenot by the Japanese

Ministry of Environment (capture permits ref. 11-79, 11-105 and 11-62). The

experimental protocol for capture, handling and sample collection was also approved by

Kyoto University's Animal Experimentation Committee (ref. Inf-K15009, Inf-K17003, Inf-K19003).

Acknowledgments

We are grateful to Dr Anja Collazo (Island Bat Research Group, IBRG) for years of contribution in collecting samples and Hiromi Kobayashi for her helpful assistance in genotyping. This work was financially supported by KAKENHI No. 20H00420 for M.I.-M., and Leading graduate program in Primatology and Wildlife Science. Sample collection and initial genetics works were performed under KAKENHI No. 17K15054 and Pro Natura Japan support granted to C.E.V.

Note

Preliminary results of this project were presented at the Island Biology Conference in Hawaii (2014). More advanced versions of this work were presented at The International Symposium on Primatology and Wildlife Science (10~13th) (2018~2020), The Annual Meeting of the Japanese Society for DNA Polymorphism Research (27~28th) (2018~2019), The 66th Annual Meeting of the Ecological Society of Japan (2019), The 63rd PRIMATES Conference (2019), and The 14th International Conference on Environmental Enrichment (2019).

Authors Contributions

Yuto Taki: Data Curation, Formal Analysis, Investigation, Methodology, Software, Validation, Visualization, Writing

Christian E. Vincenot: Conceptualization, Funding Acquisition, Investigation, Methodology, Supervision, Writing

Yu Sato: Data Curation, Formal Analysis, Investigation, Methodology, Software, Writing

Miho Inoue-Murayama: Conceptualization, Funding Acquisition, Methodology, Resources, Supervision, Writing

References

1.   Voigt CC, Kingston T, editors. Bats in the Anthropocene: Conservation of Bats in a Changing World. 1st ed. Springer International Publishing; 2016.

2.   Preble J, Ohte N, Vincenot CE. In the Shadow of the Rising Sun: A Systematic Review of Japanese Bat Research and Conservation. Mammal Review, in press.

3.   Aziz SA, Clements GR, McConkey KR, Sritongchuay T, Pathil S, Yazid MN, Campos-Arceiz, A, Forget P, Bumrungsri, S. Pollination by the locally endangered island flying fox (*Pteropus hypomelanus*) enhances fruit production of the economically important durian (*Durio zibethinus*). Ecology and Evolution. 2017;7: 8670 - 8684.

4.   Oleksy R, Racey PA, Jones G. High-resolution GPS tracking reveals habitat selection and the potential for long-distance seed dispersal by Madagascan flying foxes *Pteropus rufus*. Global Ecology and Conservation. 2015;3:678-692.

5.   Florens FBV, Baider C, Marday V, Martin GMN, Zmanay Z, Oleksy R, Krivek G, Vincenot CE, Strasberg D, Kingston. Disproportionately large ecological role of a recently mass-culled flying fox in native forests of an oceanic island. Journal for Nature Conservation. 2017;40:85-93.

6.   Oleksy RZ, Ayady CL, Tatayah V, Jones C, Howey PW, Froidevaux JSP, Racey PA, Jones G. The movement ecology of the Mauritian flying fox (*Pteropus niger*): a long-term study using solar-powered GSM/GPS tags. Movement Ecology. 2019;7(12).

7.   Epstein JH, Olival KJ, Pulliam JRC, et al. *Pteropus vampyrus*, a hunted migratory species with a multinational home-range and a need for regional management. Journal of Applied Ecology. 2009;46(5):991-1002.

8.   Hansen DM, Galetti M. The Forgotten Megafauna. Science. 2009;324 (5923):42-43.

9.   McConkey KR, Drake DR. Low redundancy in seed dispersal within an island frugivore community. AoB PLANTS. 2015;7: plv088.

10. Vincenot C.E., Florens F.B.V., Kingston T. Can we protect island flying foxes? Science. 2017;355 (6332):1368–1370.

11. Westcott DA, Caley P, Heersink DK, McKeown A. A state-space modelling

approach to wildlife monitoring with application to flying-fox abundance. Scientific Reports. 2018;8:4038.

12. Tait J, Perotto-Baldivieso HL, McKeown A, Westcott DA. Are flying-foxes coming to town? Urbanisation of the spectacled flying-fox (*Pteropus conspicillatus*) in Australia. PLoS One. 2014;9(10):e109810.

13. Vincenot C, Petit S. Australia too casual with protection law. Nature. 2016;539:168.

14. Welbergen JA, Klose SM, Markus N, Eby P. Climate change and the effects of temperature extremes on Australian flying-foxes. Proc. R. Soc. B. 2007;275:419–425.

15. Florens FBV, Baider C. Mass-culling of a threatened island flying fox species failed to increase fruit growers' profits and revealed gaps to be addressed for effective conservation. Journal for Nature Conservation. 2019;47:58-64.

16. Florens FBV, Vincenot CE. Broader conservation strategies needed. Science. 2018;362(6413):409.

17. Nuwer R. These endangered bats are being killed by the thousands—here's why. National Geogrphic; 2019.

18. Vincenot CE, Collazo A.M., Russo D. The Ryukyu Flying Fox (*Pteropus dasymallus*) - A Review of Conservation Threats and Call for Reassessment. Mammalian Biology. 2017;83(4):71–77.

19. Lee YF, Takaso TY, Chiang YM, Nakanishi N, Tzeng HY, Yasuda K (2009) Variation in the nocturnal foraging distribution of and resource use by endangered Ryukyu flying-foxes (*Pteropus dasymallus*) on Iriomotejima, Japan. Contributions to Zoology, 78: 51-64

20. Vincenot, CE (2017) *Pteropus dasymallus*. The IUCN Red List of Threatened Species. 2017; e.T18722A22080614

21. Vincenot CE, Koyama L, Russo D. Near threatened? First report of unsuspected human-driven decline factors in the Ryukyu flying fox (*Pteropus dasymallus*) in Japan. Mamm Biol. 2015;80:273–277.

22. Weerach C, Shibata S, Vincenot CE. Amidst Nets and Typhoons – Conservation Implications of Bat-Farmer Conflicts on Okinawa Island. Forthcoming, under review.

23. Vincenot CE, Collazo AM, Wallmo K, Koyama L. Public awareness and perceptual

factors in the conservation of elusive species: The case of the endangered Ryukyu flying fox. Global Ecology and Conservation. 2015;3:526-540.

24. Brown VA, Brooke AP, Fordyce JA, McCracken GF. Genetic analysis of populations of the threatened bat *Pteropus mariannus*. Conservation Genetics. 2011;12: 933-941.

25. Vincenot CE, Collazo A, Kinoshita M, Koyama L. Good DNA from fruit bat guano? A Computational Framework for remote genetic sampling assessment. in preparation.

26. Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for bigger datasets. Molecular Biology and Evolution. 2016;33: 1870-1874.

27. Bandelt H, Forster P, Röhl A. Median-joining networks for inferring intraspecific phylogenies. Molecular Biology and Evolution. 1999;16: 37–48.

28. Leigh J, Bryant D. PopART: Full-feature software for haplotype network construction. Methods in Ecology and Evolution. 2015;6: 1110–1116.

29. Petit R, Mousadik A, Pons O. Identifying populations for conservation on the basis of genetic markers. Conservation Biology. 1998;12: 844-855.

30. Taki Y, Vincenot CE, Sato Y, Nakajima N, Inoue-Murayama M. Development of 34 microsatellite markers for the Ryukyu flying fox (*Pteropus dasymallus*) Conservation Genetics Resources. 2020; in press.

31. Peakall ROD, Smouse PE. GenAlEx 6: genetic analysis in excel. Population genetic software for teaching and research. Mol Ecol Notes. 2006;6:288–295.

32. Peakall R, Smouse PE. GenAlEx 6.5: genetic analysis in Excel. Population genetic software for teaching and research-an update. Bioinformatics. 2012;28: 2537-2539.

33. Pritchard JK, Stephens M, Donnelly P. Inference of Population Structure Using Multilocus Genotype Data. Genetics. 2000;155: 945–959.

34. Rousset F, Raymond M. Statistical analyses of population genetic data: new tools, old concepts. Torends Ecol Evol. 1997;12:313–317.

35. Rousset, F. Genepop'007: a complete reimplementation of the Genepop software for Windows and Linux. Mol. Ecol. Resources. 2008;8: 103-106.

36. O'Brien J, McCracken GF, Say L, Hayden TJ. Rodrigues fruit bats (*Pteropus rodricensis*, Megachiroptera: Pteropodidae) retain genetic diversity despite

population declines and founder events. Conserv Genet. 2007;8:1073.

37.   Okada A, Suzuki H, Inaba M, Horikoshi K, Shindo J. Genetic Structure and Cryptic Genealogy of the Bonin Flying Fox *Pteropus pselaphon* Revealed by Mitochondrial DNA and Microsatellite Markers. Acta Chiropterologica. 2014;16(1): 15-26.

Figure captions

Figure 1 mtDNA haplotype distribution. Ratio of the number of samples of each haplotype is shown for each island on the map. The sizes of circles represent the sample sizes. Haplotypes shared between islands are shown with the same colors. Source of the map was the Geospatial Information Authority of Japan.

Figure 2 Principal Coordinates (PCoA) based on microsatellite genotype. Each dot represents an individual and populations are shown with colors. Three clusters are shown with circles.

Figure 3 Suggested conservation units. Source of the map was the Geospatial Information Authority of Japan.

Figure S1 SNPs of each mtDNA haplotype. Base No shows positions in the 526bp sequence.

Figure S2 mtDNA haplotype network. Each haplotype is shown as a circle with ratio of the number of samples of each island. The sizes of circles represent the sample sizes.

Figure S3 Phylogenetic tree of mtDNA sequences by maximum likelihood method. Population and sample number are shown in each place in the tree.

Figure S4 Selection of K for STRUCTURE. LnP(D) shows the accuracy of the analysis for each K. K=4 is chosen as the best model.

Figure S5 The result of STRUCTURE at K=4. Each ancestral population is shown with colors.

Figure 1

Figure 2

Genetic diversity: middle

Isolation   Active gene flow

Genetic
diversity: low

Iriomote   Kohama

Yonaguni

Miyako

Some gene flow

Ishigaki

Extinction risk:
high

Taketomi

Hateruma   Genetic diversity: high

Kuroshima

20 km

Figure 3

Figure S1

Figure S2

Figure S3

Figure S4

Figure S5

Table 1 mtDNA haplotype diversity

Population	Nh	unique	h	hr
Miyako	12	1	0.69	3.77
Ishigaki	17	9	0.96	7.51
Iriomote	12	9	1.00	9.00
Yonaguni	6	1	0.61	2.79
Average	11.75	5	0.82	5.77

Nh: The number of haplotypes, *unique*: The number of unique haplotypes, *h*: haplotype diversity, *hr*: haplotype richness

Table 2 Genetic diversity of microsatellites

Ho: observed heterozygosity, *He*: expacted heterozygosity, *F*: inbreeding coefficient

Population	The number of analyzed individuals	*Ho*	*He*	*F*
Miyako	60	0.512	0.659	0.215
Ishigaki	24	0.666	0.688	0.042
Kohama	14	0.650	0.647	-0.020
Iriomote	23	0.631	0.710	0.114
Yonaguni	30	0.388	0.577	0.355

Table 3 *Fst* between populations

Population	Miyako	Ishigaki	Kohama	Iriomote
Ishigaki	0.0221			
Kohama	0.0577	0.0493		
Iriomote	0.0349	0.0296	0.0257	
Yonaguni	0.0665	0.0656	0.0740	0.0680

Table S1 Number of samples of each population used for analysis

Population	mtDNA n	Mocrosatellite n
Miyako	63	60
Ishigaki	25	24
Taketomi	1	4
Kuroshima	2	0
Kohama	3	14
Iriomote	12	23
Hateruma	4	0
Yonaguni	25	30
Total	135	155

Table S2 The number of samples of each mtDNA haplotype in each island

Haplotype	Miyako $n=63$	Ishigaki $n=25$	Taketomi $n=1$	Kuroshima $n=2$	*Kohama* $n=3$	Iriomote $n=12$	Hateruma $n=4$	Yonaguni $n=25$	Total $n=135$
RFF1	0	0	0	0	0	1	0	0	1
RFF2	0	1	0	0	0	0	0	0	1
RFF3	0	0	0	0	0	0	0	1	1
RFF4	0	1	0	0	0	0	0	0	1
RFF5	0	0	0	0	0	1	0	0	1
RFF6	0	1	0	0	0	0	0	0	1
RFF7	6	3	0	0	0	0	0	0	9
RFF8	0	1	0	0	0	0	0	0	1
RFF9	5	0	0	1	0	0	0	0	6
RFF10	0	0	1	0	0	0	0	0	1
RFF11	0	0	0	0	0	1	0	0	1
RFF12	1	0	0	0	0	0	0	0	1
RFF13	0	0	0	0	0	1	0	0	1
RFF14	3	3	0	0	0	0	0	1	7
RFF15	0	1	0	0	0	0	0	0	1
RFF16	0	0	0	0	0	1	0	2	3
RFF17	0	0	0	0	0	1	0	0	1
RFF18	0	2	0	0	0	0	0	0	2
RFF19	0	0	0	0	0	0	1	0	1
RFF20	0	0	0	0	1	0	0	0	1
RFF21	0	0	0	0	0	1	0	5	6
RFF22	1	0	0	1	0	0	0	0	2
RFF23	0	0	0	0	1	0	0	0	1
RFF24	2	2	0	0	0	0	0	0	4
RFF25	0	0	0	0	0	1	0	0	1
RFF26	0	0	0	0	0	1	0	0	1
RFF27	0	1	0	0	0	0	0	0	1
RFF28	0	0	0	0	0	1	0	0	1
RFF29	0	1	0	0	0	0	0	0	1
RFF30	34	1	0	0	0	0	0	0	35
RFF31	2	0	0	0	0	0	1	0	3
RFF32	0	0	0	0	0	0	1	0	1
RFF33	0	0	0	0	0	1	0	0	1
RFF34	1	0	0	0	0	0	1	0	2
RFF35	2	1	0	0	0	0	0	0	3
RFF36	5	3	0	0	0	1	0	1	10
RFF37	0	1	0	0	0	0	0	0	1
RFF38	1	1	0	0	0	0	0	0	2
RFF39	0	1	0	0	1	0	0	15	17