1

The role of UCP3 in brown adipose tissue mitochondrial bioenergetics is

complementary to that of UCP1

Christopher L. Riley[1], Edward M. Mills[2]

[1]Department of Cancer Biology, Dana Farber Cancer Institute

Boston, Massachusetts 02210 USA

[2]Department of Pharmacology and Toxicology, College of Pharmacy

The University of Texas at Austin, Austin, Texas 78712 USA

Abstract

Mitochondrial uncoupling protein 3 (UCP3) plays a complementary role in

uncoupling protein 1 (UCP1)-dependent uncoupling in mammalian

thermogenesis. Using mouse strains that lack UCP1 or UCP3, plus a

UCP1/UCP3 double knockout, we have previously demonstrated that UCP3 is

necessary for sustained heat production. However, how and where UCP3 is

essential for heat production remains unknown. Here we use brown adipocyte

mitochondria from mice lacking UCP1, UCP3, or UCP1 plus UCP3 (double

knockout) we found that in the absence of UCP1, UCP3 does not act as a free

fatty acid-inducible uncoupling protein; however, UCP3 is necessary for maximal

GDP-sensitive respiration. We additionally confirm that the loss of UCP1 is a

dominant regulator of mitochondrial respiratory chain activity, assembly, and

composition in brown adipose tissue. Our data suggest without UCP1, brown

adipose tissue is not a significant source of heat production, and that UCP3-

dependent heat generation serves a regulatory function outside of adipose

tissues. Together these findings suggest that in brown adipose tissue, UCP3's

thermogenic function is independent of its role as a *bona fide* uncoupling protein.

Introduction

Brown adipose tissue (BAT) mediates approximately 60% of adaptive non-

shivering thermogenesis in small mammals, and is an essential determinant of

the arousal from torpor in hibernators[1]. BAT's recently identified presence in a

significant percentage of adult humans, along with identification of brown-like

'beige' adipocytes in white adipose tissue, suggest that modulation of BAT

function holds great promise for the treatment of human metabolic disease

associated with overnutrition [2,3].

BAT mitochondria are unique in that they oxidize substrates to generate

heat, instead of ATP: this is a metabolic property that antagonizes obesity.

Mitochondrial uncoupling protein 1(UCP1) is the well-established thermogenic

mediator of BAT, and comprises approximately 20% of the BAT mitochondrial

proteome. Stimulated by fatty acids, UCP1 'uncouples' substrate oxidation from

ATP synthesis by physically transporting protons from the intermembrane space

back into the mitochondrial matrix[4].

While UCP1 remains unquestioned as the dominant mitochondrial

uncoupling protein, the close UCP1 homolog UCP3 (57% percent sequence

identity to UCP1) is also expressed in BAT, and exhibits uncoupling properties *in*

vitro as well as *in vivo*[5–7]. However, little is known about the function of UCP3

in BAT mitochondrial bioenergetics. UCP3 is expressed in skin[8], white and

brown adipose tissue, skeletal muscle, and heart [9]. Most physiological and

biochemical studies have examined the functions of UCP3 in skeletal muscle

mitochondria, where its role as a *bona fide* mitochondrial uncoupler is debated [10–12]. Its expression in skeletal muscle is highly correlated with fatty acid oxidation as well as with levels of reactive oxygen species. Nevertheless, UCP3 expression in BAT is differentially regulated, suggesting that UPC3 may function differently in BAT and skeletal muscle[13]. And in contrast to the earlier reports, Hilse *et al.* have demonstrated that the level of expression of UCP3 is two-fold higher in BAT mitochondria than in skeletal muscle.

Despite considerable interest about its role in metabolism in general, exactly what function UCP3 serves in murine BAT mitochondrial oxidation remains a mystery. We have established that UCP3 contributes significantly to the thermogenic responses to sympathomimetics and lipopolysaccharide *in vivo* [14,15]. Here we have sought to examine the bioenergetic role of UCP3 in mitochondrial function in BAT.

A recent report showed that UCP3 is expressed in BAT at higher levels that in SKM[9]. In combination with our findings *in vivo*, we hypothesized that UCP3 acts as a traditional uncoupler in BAT. The generation of a UCP1/UCP3 double knockout allowed us to characterize BAT mitochondrial uncoupling in the absence of UCP1 in BAT. Using this double knockout model, along with single UCP1 and UCP3 null mice, we found that the loss of UCP3 in brown adipose tissue decreases GDP-sensitive mitochondrial respiration in a UCP1-dependent manner, but does not significantly regulate fatty acid-inducible uncoupling. We also confirmed previous reports that UCP1 is necessary for maximal and

inducible respiration, and controls mitochondrial respiratory chain functions in

BAT.

Materials and Methods

Chemicals

All chemicals were purchased from Sigma Aldrich unless otherwise noted.

Animals

All mice were males between 8 and12 weeks of age at the time of sacrifice. They

were fed standard chow and water *ad libitum*, and maintained on a light dark

cycle of 12hr in a room maintained at 23 +/- 2^{o}C. Wild type, UCP1KO, UCP3KO,

and UCP1/3KO animals were on a C57B6/J background. UCP1KO mice were a

gift from Dr. Leslie Kozak (Maine Medical Research Institute, Scarborough, ME,

USA); UCP3KO were a gift from Mark Reitman, formerly of the National Institutes

of Health. UCPDK (UCP1/UCP3 double knockout animals) mice were generated

in house, by breeding heterozygous the UCP1KO mouse with the heterozygous

UCP3KO mouse. Every five generations, animals from all genotypes were

backcrossed with an in-house C57B6/J colony, to prevent genetic drift. Animals

were euthanized via isoflurane inhalation followed by cervical dislocation. The

University of Texas IACUC approved all animal experiments.

Mitochondrial isolation

Mitochondria were isolated as described in [16]. Briefly, intrascapular BAT was

harvested from 2-5 mice 8-12 week old male mice, rinsed in ice cold PBS, and

cleaned of surrounding white adipose tissue. BAT was then minced with fine

scissors, and homogenized in 250 mM sucrose. The homogenate was filtered

through a 40uM cell strainer and centrifuged at 8,500 x g. The fat layer and

supernant were removed; the pellet was resuspended in sucrose and

centrifuged at 800 x g. Resultant supernant was centrifuged, and the pellet was

resuspended in sucrose supplemented with 0.3% fatty acid free BSA, and

pelleted at 8,500 x g. The resulting pellet was resuspended in 200ul of 250mM

sucrose, and protein was measured with use of an BCA assay (Pierce). All steps

were performed at 4^{o}C, or on ice - unless otherwise noted. After preparation,

mitochondria were either used directly for oxygen consumption measurements,

or was frozen at -80^{o}C for further analysis.

Mitochondrial respiration

Isolated mitochondria (0.25-0.5mg/ml) in respiration buffer 0.6mL (0.1% BSA,

125mM sucrose, 20mM TES, 1mM EDTA, 4mM KP_i, 2mM $MgCl_2$ pH 7.2) were

added to a continuously mixed chamber kept at 37^{o}C (Hanstech); respiration was

monitored for 15-30 seconds, after which pyruvate (5mM) and malate (2.5mM),

GDP (1mM), ADP (1mM), Oligomycin (1.3ug/mL), and FCCP (0.5-2uM) were

added sequentially. In a separate run, titration of respiratory activity was induced

by oleic acid (50% EtOH) after mitochondrial respiration had been inhibited by

1mM GDP and 1.3ug/mL oligomycin. Oxygen consumption was measured every

second, until the experiment was completed. All mitochondrial respiratory chain

inhibitors were solubilized in 95% EtOH unless otherwise noted, and total EtOH

up to 0.1% was found to have no effect on respiration.

Mitochondrial complex activity assays

Mitochondrial complex assays were run essentially as described in [17]. Frozen

mitochondrial aliquots were thawed on ice, then freeze-thawed an additional two

times using a dry ice bath. Prepared mitochondria were used at a concentration

of 4ug/ml (CS, CI, CIV) or 5ug/ml (CII, CI-III, CII-CIII). Absorbance was read on a

Cary 40000 UV-Vis spectrometer at 37°C. Rates were calculated in Microsoft

Excel, using linear portions of the activity. All complex activities were normalized

to citrate synthase activity.

Western blotting

Fresh mitochondrial lysates were prepared and diluted (2ug/ul) in sample buffer,

and incubated for 5 minutes at 95°C. Mitochondrial proteins were loaded and run

on a Tris-glycine gel system and transferred to nitrocellulose membranes, which

were then incubated overnight with 5% milk in TBS-tween, followed by incubation

with primary antibodies, and then for an additional hour with secondary

antibodies, washed, and developed using ECL (Thermo-scientific).

High resolution Clear Native PAGE

Pelleted mitochondria (500ug) were resuspended in 50ul of buffer (50mM NaCl,

50 mM imidazole, 2mM 6-aminohexanoic acid, 1mM EDTA, pH 7) and solubilized

with digitonin (2g/g at 20%). Supernant was cleared by spinning at 20000 x g for

20 minutes. Glycerol was added (50% 2.5ul), and 10ul (~75ug) was loaded on a

3-12% Bis-Tris native gel. The Anode buffer was comprised of (25mM imidazole

pH 7) and the cathode buffer (50 mm Tricine, 7.5 mm imidazole, 0.01% DDM pH

7). Gels were run until the MW marker reached the end of the gel, then subjected

to sequential activity assays. <u>Complex I-II activity</u>: Gels were incubated for 10

minutes in Complex I buffer (2.5mg/ml nitrotetrazolium blue, 1 mg/ml NADH in

5mM Tris/HCl), transferred to 5mM Tris/HCl buffer, scanned, and then incubated

for an additional 20 minutes in Complex II buffer (2.5mg/ml nitrotetrazolium blue,

20mM Succinate, 2uM phenazine methanosulfate, 5mM Tris/HCl) and imaged.

<u>Complex III-IV activity</u>: Gels were incubated for 30 minutes in Complex III buffer

(.5mg/ml diaminobenzidine, 50mM sodium phosphate, pH 7.2), scanned imaged,

and placed back in Complex III buffer; cytochrome c was added to a 50uM final

concentration, gels were incubated for another 30 minutes, and imaged again.

Statistics

For all experiments a one-way ANOVA followed by a *Fisher LSD* test was used

to compare multiple genotypes, with significance set *a priori* at $p < .05$.

Results and Discussion

UCP1 is a dominant regulator of brown adipose tissue respiratory chain function,

expression, and assembly in BAT mitochondria.

To better understand how UCP1/3 affects mitochondrial oxidation, we

measured the activity of respiratory chains. We found that the loss of UCP1 in

the presence or absence of UCP3 decreased all respiratory complex activities

that were measured (Figure 1). The loss of UCP1 severely impacted complex I

activity in both UCP1 and UCPDK mice, with only ~20% of WT activity remaining

(In contrast, complex II activity is ~50% of WT activity). Complex IV activity was

also decreased in the absence of UCP1, to about 60% of WT activity. Notably,

complex IV activity was significantly increased in the UCP3KO mitochondria

(Figure 1C), and in agreement with the report by Liebig *et al.* [18].

To explore the decrease of respiratory chain activity, we determined

protein abundance via western blotting, and found that the loss of UCP1

decreased Ndufa9 (complex I), Uqcrc2 (complex III), and Coxiv (complex IV)

protein abundance in mitochondria (Figure 2A). Coxiv expression was also

decreased in the double knockout background. However, expression of Sdha

(complex II) was not affected, despite a reduction in its activity (Figure 1B, 2A).

Neither the expression of Oscp (complex V) nor of Mdh2(TCA cycle enzyme)

was changed in any of the genotypes (Figure 2). Because these enzymes only

represent a selected few of the many components of each respiratory complex,

we investigated how both UCP's influence the formation of respiratory chain

complex assembly in BAT. Using activity-based high resolution clear native

PAGE, we confirmed that in the absence of UCP1, complex I, complex III, and

complex IV are depleted and disorganized (Figure 2C), and complex III

occupancy was decreased in supercomplexes (Figure 3C). However, no

additional loss of mitochondrial complexes was detected in the UCPDK mice

when compared with mice in which only UCP1 was depleted (Figure 3C).

Together, these data unambiguously demonstrate that Ucp1 is necessary

for thermogenically competent brown adipose mitochondria. Our data strongly

support those of Kazak *et al.* We also expand their finding that Ucp1 is essential

for the functioning of the brown fat respiratory chain at 4C, to include mice that

are raised under standard housing conditions[19]. The loss of Ucp1 extensively

alters the BAT proteome, and highlight that UCP3 cannot compensate for the

loss of Upc1, even though Ucp3 is expressed in this model [9].

The discrepancies in Sdha protein abundance and complex II activity

between WT and UCP1-null animals points to the existence of a regulatory

function that does not allow electron transfer between the matrix-exposed Sdha-

Sdhb complex and the membrane-embedded Sdhc-Sdhd complex. Confirmation

of increased complex IV activity, despite similar protein abundance and complex

integrity in the UCP3-null mouse and the Ucp3-null hamster, as shown by

Liebig's work, suggests that this phenomenon reflects a mechanism wherein an

increase complex IV activity may compensate for the loss of Ucp3 function.

Loss of UCP3 decreases GDP sensitive respiration, but does not affect FFA-

induced respiration in BAT mitochondria.

As mitochondria are an integrated system, we used high-resolution

respirometry to further probe the bioenergetic function of UCP3 in brown adipose

mitochondria. The protocol outlined in Figure 4A utilizes continuous addition of

substrates and inhibitors, to elucidate GDP-sensitive (respiration after addition of

substrate minus respiration after GDP addition), ADP-linked (respiration after

addition of ADP minus respiration after oligomycin), maximal respiration (CCCP

induced) (Fig. 3A) [20]. In agreement with the complex activity data, loss of

UCP1 was found to be the dominant regulator of respiration (Fig. 3B-D). Notably,

the loss of UCP3 only (i.e., with UCP1 present) led to a decrease in GDP-

sensitive respiration (Fig. 3C). ADP-linked respiration was equal across all

genotypes, despite the significant decrease in total respiratory capacity (Fig. 3E).

We found no significant difference in free fatty acid-mediated uncoupling in the

presence or absence of UCP3(Fig. 3F).

The finding that ADP-linked respiratory capacity is the same between the

UCP1-null and WT mitochondria is critical: it suggests that the recently

discovered ATP-linked thermogenic pathways remain unaffected by the loss of

UCP1. This finding means that proposed futile cycles that require ATP production

(phospho-creatine and calcium) can run in the absence of UCP1 in BAT[21,22].

However, total respiratory capacity in this tissue is likely compromised, and heat

production would be severely impaired relative t WT animals.

Previous reports suggest that the role of Ucp3 in BAT is negligible. However,

those studies and the data presented in this report have critical differences.

Gong et al. used a mouse model on a mixed background that is not highly reliant on BAT thermogenesis[23]. Also, they employed succinate as the oxidative substrate - given that succinate is not compatible with maximal oxidation in BAT mitochondria, their results may have underestimated the function of UCP3[16]. Additionally, when both UCP3 expression and adrenergic tone in BAT were low, [16] they housed their mice at thermoneutrality [16]. And Liebig et al. used a mutant hamster with BAT-specific ablation of UCP3, and also found no significant effect on mitochondrial oxidation when respiration was driven with glycerol-3-phosphate (G3P). In contrast, here we have used pyruvate and malate, a complex I-linked substrate that requires the transport of both substrates across the inner membrane.

Discrepancies between our findings and earlier reports may be explained by UCP3's hypothesized transport function, through which it increases the availability of substrates pyruvate or malate during maximal UCP1 activation[11]. UCP2, which shares a 72% homology with UCP3, directly transports C4 metabolites, and controls glycolysis as well as glutamine oxidation in hepatocytes [24]. A recent study has demonstrated biochemical differences between UCP1 and UCP3 that point to differing transport roles[25]. In light of existing literature, our data suggest that UCP3 complements UCP1's uncoupling function, not as an uncoupler but as a solute transporter.

In summary, our works shows that UCP3 does have a role in mitochondrial oxidation in brown adipose tissue mitochondria. Future work will

investigate the possible transport functions of this protein and its role in substrate

oxidation more globally.

[1] D.O. Foster, M.L. Frydman, Tissue distribution of cold-induced

thermogenesis in conscious warm- or cold-acclimated rats reevaluated

from changes in tissue blood flow: the dominant role of brown adipose

tissue in the replacement of shivering by nonshivering thermogenesis.,

Can. J. Physiol. Pharmacol. 57 (1979) 257–70.

http://www.ncbi.nlm.nih.gov/pubmed/445227 (accessed November 27,

2017).

[2] A.M. Cypess, S. Lehman, G. Williams, I. Tal, D. Rodman, A.B. Goldfine,

F.C. Kuo, E.L. Palmer, Y.-H. Tseng, A. Doria, G.M. Kolodny, C.R. Kahn,

Identification and Importance of Brown Adipose Tissue in Adult Humans,

N. Engl. J. Med. 360 (2009) 1509–1517.

https://doi.org/10.1056/NEJMoa0810780.

[3] J.S. Flier, L.H. Underhill, J. Himms-Hagen, Thermogenesis in Brown

Adipose Tissue as an Energy Buffer, N. Engl. J. Med. 311 (1984) 1549–

1558. https://doi.org/10.1056/NEJM198412133112407.

[4] R.M. Locke, E. Rial, I.D. Scott, D.G. Nicholls, Fatty acids as acute

regulators of the proton conductance of hamster brown-fat mitochondria.,

Eur. J. Biochem. 129 (1982) 373–80.

http://www.ncbi.nlm.nih.gov/pubmed/6295765 (accessed November 18,

2017).

[5] K.S. Echtay, Q. Liu, T. Caskey, E. Winkler, K. Frischmuth, M.

Bienengräber, M. Klingenberg, Regulation of UCP3 by nucleotides is

different from regulation of UCP1., FEBS Lett. 450 (1999) 8–12.

http://www.ncbi.nlm.nih.gov/pubmed/10350047 (accessed November 18,

2017).

[6] D. Ricquier, F. Bouillaud, The uncoupling protein homologues: UCP1,

UCP2, UCP3, StUCP and AtUCP., Biochem. J. 345 Pt 2 (2000) 161–79.

http://www.ncbi.nlm.nih.gov/pubmed/10620491 (accessed November 16,

2017).

[7] K.S. Echtay, D. Roussel, J. St-Pierre, M.B. Jekabsons, S. Cadenas, J.A.

Stuart, J.A. Harper, S.J. Roebuck, A. Morrison, S. Pickering, J.C. Clapham,

M.D. Brand, Superoxide activates mitochondrial uncoupling proteins,

Nature. 415 (2002) 96–99. https://doi.org/10.1038/415096a.

[8] C.U. Lago, S.M. Nowinski, J.E. Rundhaug, M.E. Pfeiffer, K. Kiguchi, K.

Hirasaka, X. Yang, E.M. Abramson, S.B. Bratton, O. Rho, R. Colavitti, M.A.

Kenaston, T. Nikawa, C. Trempus, J. DiGiovanni, S.M. Fischer, E.M. Mills,

Mitochondrial respiratory uncoupling promotes keratinocyte differentiation

and blocks skin carcinogenesis, Oncogene. 31 (2012) 4725–4731.

https://doi.org/10.1038/onc.2011.630.

[9] K.E. Hilse, A. V Kalinovich, A. Rupprecht, A. Smorodchenko, U. Zeitz, K.

Staniek, R.G. Erben, E.E. Pohl, The expression of UCP3 directly correlates

to UCP1 abundance in brown adipose tissue., Biochim. Biophys. Acta.

1857 (2016) 72–8. https://doi.org/10.1016/j.bbabio.2015.10.011.

[10]   I.G. Shabalina, J. Hoeks, T. V. Kramarova, P. Schrauwen, B. Cannon, J. Nedergaard, Cold tolerance of UCP1-ablated mice: A skeletal muscle mitochondria switch toward lipid oxidation with marked UCP3 up-regulation not associated with increased basal, fatty acid- or ROS-induced uncoupling or enhanced GDP effects, Biochim. Biophys. Acta - Bioenerg. 1797 (2010) 968–980. https://doi.org/10.1016/j.bbabio.2010.02.033.

[11]   J. Mozo, G. Ferry, A. Studeny, C. Pecqueur, M. Rodriguez, J.A. Boutin, F. Bouillaud, Expression of UCP3 in CHO cells does not cause uncoupling, but controls mitochondrial activity in the presence of glucose, Biochem. J. 393 (2006) 431–439. https://doi.org/10.1042/BJ20050494.

[12]   A.J. Vidal-Puig, D. Grujic, C.-Y. Zhang, T. Hagen, O. Boss, Y. Ido, A. Szczepanik, J. Wade, V. Mootha, R. Cortright, D.M. Muoio, B.B. Lowell, Energy Metabolism in Uncoupling Protein 3 Gene Knockout Mice, J. Biol. Chem. 275 (2000) 16258–16266. https://doi.org/10.1074/jbc.M910179199.

[13]   F. Denjean, J. Lachuer, A. Géloën, F. Cohen-Adad, C. Moulin, H. Barré, C. Duchamp, Differential regulation of uncoupling protein-1, -2 and -3 gene expression by sympathetic innervation in brown adipose tissue of thermoneutral or cold-exposed rats., FEBS Lett. 444 (1999) 181–5. http://www.ncbi.nlm.nih.gov/pubmed/10050755 (accessed November 18, 2017).

[14]   C.L. Riley, C. Dao, M.A. Kenaston, L. Muto, S. Kohno, S.M. Nowinski, A.D. Solmonson, M. Pfeiffer, M.N. Sack, Z. Lu, G. Fiermonte, J.E. Sprague, E.M. Mills, The complementary and divergent roles of uncoupling proteins

1 and 3 in thermoregulation, J. Physiol. 594 (2016) 7455–7464.

https://doi.org/10.1113/JP272971.

[15] E.M. Mills, M.L. Banks, J.E. Sprague, T. Finkel, Pharmacology: uncoupling

the agony from ecstasy., Nature. 426 (2003) 403–4.

https://doi.org/10.1038/426403a.

[16] I.G. Shabalina, M. Ost, N. Petrovic, M. Vrbacky, J. Nedergaard, B. Cannon,

Uncoupling protein-1 is not leaky, Biochim. Biophys. Acta - Bioenerg. 1797

(2010) 773–784. https://doi.org/10.1016/j.bbabio.2010.04.007.

[17] M. Spinazzi, A. Casarin, V. Pertegato, L. Salviati, C. Angelini, Assessment

of mitochondrial respiratory chain enzymatic activities on tissues and

cultured cells, Nat. Protoc. 7 (2012) 1235–1246.

https://doi.org/10.1038/nprot.2012.058.

[18] M. Liebig, C. von Praun, G. Heldmaier, M. Klingenspor, Absence of UCP3

in Brown Adipose Tissue Does Not Impair Nonshivering Thermogenesis,

Physiol. Biochem. Zool. 77 (2004) 116–126.

https://doi.org/10.1086/381464.

[19] L. Kazak, E.T. Chouchani, I.G. Stavrovskaya, G.Z. Lu, M.P. Jedrychowski,

D.F. Egan, M. Kumari, X. Kong, B.K. Erickson, J. Szpyt, E.D. Rosen, M.P.

Murphy, B.S. Kristal, S.P. Gygi, B.M. Spiegelman, UCP1 deficiency causes

brown fat respiratory chain depletion and sensitizes mitochondria to

calcium overload-induced dysfunction, Proc. Natl. Acad. Sci. 114 (2017)

7981–7986. https://doi.org/10.1073/pnas.1705406114.

[20] A. Matthias, K.B. Ohlson, J.M. Fredriksson, A. Jacobsson, J. Nedergaard,

B. Cannon, Thermogenic responses in brown fat cells are fully UCP1-dependent. UCP2 or UCP3 do not substitute for UCP1 in adrenergically or fatty scid-induced thermogenesis., J. Biol. Chem. 275 (2000) 25073–81. https://doi.org/10.1074/jbc.M000547200.

[21] L. Kazak, E.T. Chouchani, M.P. Jedrychowski, B.K. Erickson, K. Shinoda, P. Cohen, R. Vetrivelan, G.Z. Lu, D. Laznik-Bogoslavski, S.C. Hasenfuss, S. Kajimura, S.P. Gygi, B.M. Spiegelman, A Creatine-Driven Substrate Cycle Enhances Energy Expenditure and Thermogenesis in Beige Fat, Cell. 163 (2015) 643–655. https://doi.org/10.1016/j.cell.2015.09.035.

[22] K. Ikeda, Q. Kang, T. Yoneshiro, J.P. Camporez, H. Maki, M. Homma, K. Shinoda, Y. Chen, X. Lu, P. Maretich, K. Tajima, K.M. Ajuwon, T. Soga, S. Kajimura, UCP1-independent signaling involving SERCA2b-mediated calcium cycling regulates beige fat thermogenesis and systemic glucose homeostasis, Nat. Med. 23 (2017) 1454–1465. https://doi.org/10.1038/nm.4429.

[23] D.-W. Gong, S. Monemdjou, O. Gavrilova, L.R. Leon, B. Marcus-Samuels, C.J. Chou, C. Everett, L.P. Kozak, C. Li, C. Deng, M.-E. Harper, M.L. Reitman, Lack of Obesity and Normal Response to Fasting and Thyroid Hormone in Mice Lacking Uncoupling Protein-3, J. Biol. Chem. 275 (2000) 16251–16257. https://doi.org/10.1074/jbc.M910177199.

[24] A. Vozza, G. Parisi, F. De Leonardis, F.M. Lasorsa, A. Castegna, D. Amorese, R. Marmo, V.M. Calcagnile, L. Palmieri, D. Ricquier, E. Paradies, P. Scarcia, F. Palmieri, F. Bouillaud, G. Fiermonte, UCP2 transports C4

metabolites out of mitochondria, regulating glucose and glutamine

oxidation, Proc. Natl. Acad. Sci. 111 (2014) 960–965.

https://doi.org/10.1073/pnas.1317400111.

[25] G. Macher, M. Koehler, A. Rupprecht, J. Kreiter, P. Hinterdorfer, E.E. Pohl,

Inhibition of mitochondrial UCP1 and UCP3 by purine nucleotides and

phosphate, Biochim. Biophys. Acta - Biomembr. (2017).

https://doi.org/10.1016/j.bbamem.2017.12.001.

Figure Legend

Figure 1. *Loss of UCP1 is the dominant regulator of mitochondrial respiratory chain enzyme activity*. (A) Complex I activity (B) Complex II activity and (C) Complex IV activity were each controlled for total mitochondria, using citrate synthase activity. Freeze-thawed mitochondria from WT (n=3) UCP1KO (n=3) UCP3KO (n=3) and UCPDK(n=4). *= $p< .05$, **=$p<.001$, ***= $p<.0001$, ****= $p<.00001$.

Figure 2. *Loss of UCP1 is the dominant regulator of respiratory chain protein expression and organization in brown adipose tissue mitochondria.* (A) Western blots of mitochondrial extracts from indicated genotypes. Five micrograms of mitochondrial protein from three separate mitochondrial preparations per genotype. (B) Densitometry from data in *A* was quantified using ImageJ, and was controlled for Mdh2 expression. N=3 per genotype. (C) Representative high resolution clear native PAGE analysis of mitochondria from indicated genotypes.

Figure 3. *Regulation of brown adipose tissue mitochondrial respiration by uncoupling proteins 1 and 3*. (A) Scheme for mitochondrial respiration assays, (B) Mitochondrial respiration from 5-6 different preparations per genotype, (C) GDP-sensitive respiration, (D) Maximal respiration, (E) ADP-linked respiration, and (F) Oleate induced respiration. (A-E) n= 5-6 per genotype *= $p< .05$, **=$p<.001$, ***= $p<.0001$, ****= $p<.00001$.

Figure 2

Figure 1

Figure 3