Unilateral vocal nerve resection alters neurogenesis in the avian song system in a

region-specific manner.

Jake V. Aronowitz[1†], Alice Perez[2†], Christopher O'Brien[1], Siaresh Aziz[1], Erica

Rodriguez[1], Kobi Wasner[1], Sissi Ribeiro[1], Dovounnae Green[1], Farhana Faruk[1], Carolyn

L. Pytte[1,2*]

[1] Psychology Department, Queens College, City University of New York, Flushing, NY

11367

[2] Psychology Department, Biology Department, The Graduate Center, City University of

New York, New York, NY 11016

[†]Equal contribution

*Corresponding author

Email: Carolyn.Pytte@QC.cuny.edu

Abstract

New neurons undergo a critical period soon after migration during which the behavior of

the animal may result in the survival or culling of these cells. In the songbird song

system, new neurons may be maintained in the song motor pathway with respect to

motor progression toward a target song--during juvenile song learning, seasonal song

restructuring, and experimentally manipulated song variability. However, it is not known

whether the quality of song per se, without progressive improvement, may also

influence new neuron survival. To test this idea, we experimentally altered song

acoustic structure by unilateral denervation of the syrinx. We found no effect of

aberrant song on numbers of new neurons in the HVC of the song motor pathway, a

loss of left-side dominance in new neurons in the auditory region caudomedial

nidopallium (NCM), and a bilateral decrease in new neurons in the basal ganglia

nucleus Area X. We propose new neuron survival may be determined in response to

behavioral feedback in accordance with the function of new neurons within a given brain

region. Studying the effects of singing behaviors on new neurons across multiple brain

regions that subserve singing may give rise to general rules underlying the regulation of

new neuron survival across taxa and brain regions more broadly.

Introduction

Most cell types that continue to be produced postnatally have predetermined lifespans, varying by tissue type, and largely independent of the experience of the cell. In healthy adults, epithelial cells survive less than 2 weeks (Spalding et al., 2005), white blood cells live for about 13-20 days, red blood cells survive 100-120 days (Krzyzanski et al., 2013), and liver hepatocytes turnover about every 200-300 days (Macdonald, 1961; Stocker and Heine, 1971). On the other hand, across species and brain regions, lifespans of neurons born in the postnatal brain are highly variable. More interestingly, lifespans differ not only by neuron type, but are influenced by the cell's activity and experience, setting them apart in this way from other continually regenerating cell types (Kirn et al., 1994; Kirn and Schwabl, 1997; Gheusi et al., 2000; Rochefort et al., 2002; Shors et al., 2002; Dupret et al., 2008; Belnoue et al., 2013; Liu and Nusslock, 2018; Larson et al., 2019).

Time-dependent changes in the expression of a number of transmembrane factors and intracellular signaling molecules in immature neurons seem to mediate the transition between cell survival and cell death, with different mosaics of receptors expressed throughout neuronal maturation in a cell-type specific manner (Pfisterer and Khodosevich, 2017). However, even after an early critical period of culling within the first few weeks of a neuron's life, microenvironmental and behavioral conditions continue to impact the neuronal lifespan (Kirn et al., 1999; Alvarez-Borda et al., 2004; Brenowitz and Larson, 2015; Shevchouk et al., 2017).

In the songbird, behavioral factors affecting new neuron survival have been studied most systematically in the nucleus HVC (Wilbrecht and Kirn, 2004; Brenowitz and

Larson, 2015), a sensorimotor region subserving song production that also responds to

song auditory stimuli (McCasland, 1987; George et al., 2005). Throughout the bird's life,

HVC receives new neurons that project to the premotor robust nucleus of the

arcopallium (RA), becoming part of the motor pathway for song production (HVC_{RA}). In

the canary, amount of singing positively correlates with new neuron survival (Li et al.,

2000; Alvarez-Borda and Nottebohm, 2002; Ball et al., 2004). This association is

mediated by BDNF and testosterone, both of which may directly increase new neuron

survival (Rasika et al., 1994; Rasika et al., 1999; Alvarez-Borda et al., 2004; Brenowitz

and Larson, 2015; Alward et al., 2016; Ball et al., 2020). Housing conditions also impact

the lifespan of both HVC_{RA} neurons and new neurons in the auditory region

caudomedial nidopallium (NCM), with more new neurons seen in zebra finches or

canaries housed in groups compared to singly housed birds (Lipkind et al., 2002;

Barnea et al., 2006; Walton et al., 2012; Shevchouk et al., 2017). Interestingly,

Shevchouk et al., (2017) show that new neuron proliferation and the subsequent

survival of new neurons can be independently affected by male- or female-paired

housing. Numerous behaviors may underlie this effect, including increased singing and

increased exposure to songs. Intriguingly, environmental factors of photo period and

temperature appear to influence new neuron numbers in the caudal nidopallium and

hippocampus in the non-seasonal zebra finch (Pozner et al., 2018).

Across taxa, a common modulator of neuronal lifespan is neural activity, with most

evidence indicating that activity confers survival (Alvarez-Borda and Nottebohm, 2002;

Ninkovic et al., 2007; Larson et al., 2013; Brenowitz and Larson, 2015; Pfisterer and

Khodosevich, 2017; Hall and Tropepe, 2018). However, neuronal activity has also

decreased new neuron survival of olfactory periglomerular cells under precise

conditions, perhaps fine-tuning olfactory sensitivity to odorant density (Khodosevich et

al., 2013). During brain development, neuronal activity conferring cell survival may

direct pathway maintenance and pruning thereby building system efficiency. Perhaps a

similar functional outcome occurs in adulthood. This has been proposed by Wilbrecht et

al. (2002) and (Wilbrecht and Kirn, 2004) in a model in which new HVC neurons

audition for a spot in the song production pathway and get the part when their

contribution improves song performance. Interestingly, this model considers not only the

amount of songs produced, but also the ongoing *quality* of the song. In this way, they

propose that the act of accurate progression in a goal directed behavior interacts with

song production in promoting new neuron retention (Wilbrecht and Kirn, 2004; Pytte et

al., 2008).

Consistent with this idea, experimentally altered song structure, by reversible

paralysis of the syringeal muscles, resulted in a positive correlation between numbers of

new neurons in HVC and song recovery (Pytte et al., 2011). However, the direction of

causality could not be determined. Increasingly accurate song production during song

recovery may have promoted the survival of new neurons or birds with higher numbers

of new neurons may have been better able to correct altered song structure (Pytte et al.,

2011). Song recovery, i.e., goal directed improvement in the behavior, was also

confounded by song quality per se. Songs that were the most aberrant recovered the

least. Therefore, although consistent with the model that behavioral improvement in

quality promotes new neuron survival, either song quality or song recovery may have

influenced new neuron survival in this study. Alternatively, perhaps new neurons in HVC

drove song recovery and song feedback was not a factor in neuronal survival.

To address this, we tested whether song quality without recovery impacts new

neuron survival. Here we produced an irreversibly aberrant song by unilaterally

denervating the syrinx in adult male zebra finches, resulting in stable, altered feedback

during song production. This allowed us to determine directly whether song quality may

impact numbers of new neurons, and also to broaden our investigation to song regions

outside of HVC.

In addition to HVC and NCM, the basal ganglia nucleus Area X processes song-

related feedback and continues to receive new neurons throughout adulthood

(Kosubek-Langer et al., 2017). Area X receives projections from a different population of

HVC neurons (HVC_X) than those that project to RA and is part of the anterior forebrain

pathway necessary for juvenile song learning (Sohrabji et al., 1990) and adult song

plasticity (Woolley et al., 2014). NCM affects HVC activity indirectly (Pawlisch and

Remage-Healey, 2015) and perhaps is a source of auditory input into the "song

system," the group of interconnected nuclei that subserve song behavior (Figure 1).

Given the functional and anatomical connections between HVC, NCM and Area X, we

explored effects of altered song feedback on new neurons across these regions.

We also speculated that differences in the effects of feedback on new neurons may

not only occur across brain regions, but also perhaps across the left and right

hemispheres of a given brain region. Tsoi et al. (2014) reported inter-hemispheric

differences whereby a greater number of new neurons are added to left NCM than right

NCM of adult zebra finches. Moreover, the degree of asymmetry was positively

correlated with the quality of song learning and the strength of neuronal memory for

songs (Tsoi et al., 2014). Improving our understanding of how new neuron survival is

regulated across brain regions and between hemispheres may inform strategies that

consider interconnected effects in a whole brain system in enhancing the survival of

new neurons post brain injury or disease (Mu and Gage, 2011; Sun, 2014; Winner and

Winkler, 2015) and in promoting brain health more generally.

Materials and Methods

Animals. All methods were approved by the Queens College Institutional Animal Care

and Use Committee (IACUC). Subjects were adult male zebra finches between 4 and

11 months of age. In zebra finches, only the males sing. We examined both the

recruitment and survival of new neurons using doublecortin (DCX) and

bromodeoxyuridine (BrdU) labeling, respectively. Different cohorts of birds were used in

labeling either DCX and BrdU, with overlapping individuals used within a label: DCX in

HVC (n = 20), Area X (n = 24), and NCM (n = 19); BrdU in HVC (n = 22), Area X (n =

40), and NCM (n = 22). Birds were hatched in either the Queens College or Wesleyan

University breeding colony and kept with their parents until 90 days of age. Thereafter,

birds were group housed, with experimental and control birds housed together in group

cages, within auditory and visual contact of both sexes throughout the study. Birds were

maintained on a 12:12 h light:dark schedule with food and water available *ad libitum*.

Song Recording and Analysis. Adult male zebra finches were temporarily housed

individually in sound attenuated chambers to record songs pre- and post-surgery. They

were recorded continuously for 2-4 days prior to surgery, and again 1-2 days after

surgery and then returned to their home cages. Recordings were made (Earthworks

precision audio SR20 Cardioid Microphone) using sound-activated Avisoft Recorder

(Avisoft Bioacoustics). Song bouts were edited to single motifs with Raven sound

analysis software (Cornell Lab of Ornithology). Changes in structure after surgery were

computed using Sound Analysis Pro (Tchernichovski et al., 2000). We used two

measures to assess song quality: (1) accuracy and (2) percentage of similarity (termed

here "similarity"). The accuracy score indicates the fidelity with which song elements

were reproduced by comparing 1 ms song segments between pre- and post-operative

motifs. Similarity indicates sameness in song structure at longer intervals (~50 ms)

(http://soundanalysispro.com/manual-1). We compared 10 pre-surgery and post-surgery

motif pairs for each bird and used the mean of these scores as a measure of

postoperative song change.

We were interested in whether the degree of global song change post TS-cut, or a

change in a particular song feature, corresponded to numbers of new neurons in

various brain regions. Therefore, we ran independent linear regressions between the

following measures of pre- to post-surgery song change and cell counts: accuracy,

similarity, pitch difference, FM difference, entropy difference, and goodness difference.

P values were corrected for multiple comparisons by adjusting the significance p value

to $0.05/6 = 0.0083$, using the Bonferroni correction. Uncorrected p values are reported

in the text along with the corrected p value threshold necessary for significance.

Singing rates were computed manually from continual voice-activated recordings

taken on days 2-4 post-surgery. Total songs were counted for each bird during a single

hour within two hours of lights coming on. Mean songs per hour were calculated.

Bromodeoxyuridine (BrdU) Injections and Tracheosyringeal Denervation. All birds

received intramuscular injections of 5-Bromo-2'-deoxyuridine (BrdU; 74 μg/g, pH 7.4,

Sigma) 3x/day for three days to label mitotically active cells. Surgeries were performed

21-22 days after the last BrdU injection, such that new neurons were 21-24 days old

when song feedback was altered. This time frame was selected in order to ensure new

neurons had migrated into the three regions of interest prior to nerve resection.

Birds were anesthetized with either a mixture of ketamine and xylazine (0.03-0.05

mg/g and 0.06 mg/g, respectively), or isoflurane in compressed air (3%; Henry Schein).

Using a surgical microscope (Zeiss Universal S3B), a ~3-mm rostro-caudal incision was

made < 1 mm lateral to the ventral midline to expose the tracheosyringeal nerve. A 2-4

mm section of either the left or right tracheosyringeal nerve (nXIIts, "TS") was resected

to prevent regrowth following surgery. Sham birds received all surgical manipulations

except nerve resection.

Histology. Birds experienced altered song for seven days and then were overdosed

with sodium pentobarbital (Euthasol) and transcardially perfused with 0.1 M phosphate

buffered saline (PBS, pH 7.4) followed by 4% paraformaldehyde (PFA; Sigma-Aldrich;

pH 7.4). Brains were post-fixed for 1 h in 4% PFA, rinsed in PBS for 3 h and embedded

in polyethylene glycol (MW = 1500; Polysciences). Six-ụm sagittal sections were cut on

a rotary microtome. The brain was oriented with the midline parallel to the blade edge.

The first complete section through the telencephalon was saved and subsequently

every sixth section of tissue was mounted onto Superfrost Plus + slides (VWR). All

series were stored at -20º C prior to processing using immunohistochemistry (IHC).

DCX Immunohistochemistry. Immunolabeling was performed using an anti-DCX goat polyclonal IgG (Santa Cruz Biotechnology; sc-8066, 1:150) or anti-DCX rabbit polyclonal IgG (Abcam; ab18723, 1:1000). Sections were brought to room temperature in TBS, followed by a 10-min wash in fresh TBS. Sections were then incubated for 30-min in a H_2O_2 solution (97% TBS, 1% methanol, 2% of 3% retail H_2O_2) to eliminate endogenous peroxidases. After three 5-min TBS rinses, sections were blocked with 5% normal horse serum (Jackson ImmunoResearch Laboratories Inc.) and 0.5% Triton X-100 (Sigma-Aldrich) in TBS for 30-min at room temperature, followed by exposure to anti-DCX polyclonal antibody in the same blocking buffer overnight. After three 5-min TBS rinses, sections were incubated for 3 h in biotinylated horse anti-goat (1:200, Vector Labs) or biotinylated goat anti-rabbit secondary antibody (1:200, Vector Labs) in TBS, rinsed again, and exposed for 1 h to an avidin-biotin complex (Vector Labs). Sections were rinsed in TBS and then reacted in a solution of 0.04% of 3,3'-diaminobenzidine tetrahydrochloride (DAB, Vector Labs) with nickel until the tissue changed color. Following three 5-min TBS rinses, sections were dehydrated in ethanols, delipidized in xylenes, and cover slipped with Krystalon (Millipore-Sigma).

BrdU/Hu Immunohistochemistry. Sections were brought to room temperature in 0.1 M phosphate buffer (PB; pH 7.4), then exposed to citrate buffer (pH 5.6 – 6.0) at 90-95°C for 10 min, followed by a 5-min PB wash (37°C), 3 min in a solution of 0.28 % pepsin in 400 ml 0.1 M HCl at 37°C, and three 5-min washes in PB at room temperature. Sections were blocked with 3% normal donkey serum (Jackson Labs) and 0.5 % Triton X-100 in PB for 1 h at room temperature, followed by 24-48 h exposure to sheep anti-BrdU (1:239, Capralogics) in block at 4°C. Sections were again rinsed in PB

and then incubated overnight in biotinylated donkey anti-sheep IgG in PB (1:200, Vector

Labs), followed by overnight incubation in streptavidin-conjugated Alexa 488 in PB

(1:800, ThermoFisher Scientific) for visualization of BrdU. The next day, sections were

washed with PB, blocked for one hour, and incubated overnight in mouse anti-Hu

primary antibody at 4°C (1:200 in 3% block; Invitrogen). After three 5-min PB rinses at

room temperature, tissue was exposed to donkey anti-mouse IgG conjugated to Cy-3 in

PB (1:80; EMD Millipore) for 1 h for visualization of Hu. Finally, sections were washed,

dehydrated with ethanols, delipidized with xylenes, and cover slipped with Krystalon

(Millipore-Sigma).

Microscopy. Data were collected without knowledge of bird identity, treatment, or brain

hemisphere. We had no predictions for potential hemispheric differences in new

neurons in HVC or Area X. Area measurements and cell counts for all regions were

performed using a computer-yoked microscope and mapping software (Olympus BX51;

Lucivid LED microprojection, Neurolucida, MicroBrightField, Inc.). The boundaries of all

brain regions were traced in 6-12 sections per hemisphere per bird. Boundaries for HVC

were established with dark-field optics based on neuropil density and contrast. Area X is

found ventral and anterior to the densely reflective oval shaped lateral magnocellular

nucleus of anterior nidopallium (lMAN) and is demarcated by a dense haze of terminals

and small cells. Area X and lMAN are separated by lamina pallio-subpallialis (LPS).

NCM was identified as in (Pytte et al., 2010). DCX[+] cells were visualized with bright

field light microscopy (Figure 2A,B). BrdU[+]/Hu[+] cells were visualized using fluorescein

isothiocyanate (FITC) and rhodamine filters and a dual FITC/rhodamine filter (Figure

2C,D,E). New neurons per square millimeter was calculated by dividing the numbers of

labeled cells by the total area sampled.

Statistical Analysis. Data are presented as means and SEMs unless otherwise

specified. Analyses were performed using one-way ANOVAs, two-factor ANOVAs with

repeated measures on hemisphere, and two-tailed *t*-tests for independent samples.

Tukey's HSD post-hoc tests were used following significant effects determined by

ANOVA. Following significant mixed ANOVAs, we performed one-way ANOVAs

between groups within individual hemispheres and then Holm's procedure with adjusted

p values for multiple tests. Pearson correlations were computed between cell counts

across brain regions or hemispheres. The lateralization index (LI) indicates the relative

number of new neurons between hemispheres, normalized for the bird's mean number

of new neurons per area sampled in both hemispheres (as in Tsoi et al., 2014). Higher

positive values indicate more new neurons in the left hemisphere relative to the right:

$$\frac{\text{Left neurons/mm}^2 - \text{Right neurons/mm}^2}{(\text{Left neurons} + \text{Right neurons})/(\text{Left area sampled} + \text{Right area sampled})}$$

Results

Song Analysis. Birds that underwent either left or right TS-cut had significantly lower

accuracy scores, a measure of song change post-denervation, than controls (One-way

ANOVA, df = 2, F = 13.95, p = 0.002, Figure 3A). Both left TS-cut and right TS-cut

scores were lower than those of controls (Tukey's post-Hoc test, p < .01) and there was

no difference between the two denervated groups (Tukey's post-Hoc test p > .05). On

the other hand, there was not a significant difference in pre-postop similarity scores

among groups (one way ANOVA, df = 2, F = 1.84, p = 0.22, Figure 3B), suggesting that

overall song structure was preserved in TS-cut birds despite degradation of fine

structure. Post-operative singing rates did not differ among groups (One-way ANOVA,

df = 2, F = 0.231, p = 0.797 data not shown).

Effect of TS-cut on New Neurons in HVC. There was no effect of treatment (df = 2, F

= 0.75, p = 0.49), hemisphere (F = 0.04, p = 0.84), or interaction (F = 0.29, p = 0.75) on

numbers of $BrdU^+/Hu^+$ cells in HVC that were 28-30 days old (two-factor ANOVA,

repeated measures on hemisphere, Figure 4A). We also quantified DCX+ cells, a

marker for immature neurons that is expressed for 2-3 weeks after mitosis in mammals

(Brown et al., 2003) and was calculated to be expressed in newborn neurons for a

similar time, approximately 20 days, in the canary (Balthazart, et al., 2008). The

morphology of DCX+ neurons can also be used to estimate neuronal age in this time

frame with fusiform, bipolar cells presumably with a migratory morphology being

younger, and round, multipolar cells being older (Figure 2A, B; Balthazart, et al., 2008).

We did not find treatment or hemisphere differences when fusiform and multipolar cell

counts were analyzed independently (p > 0.05 for all); therefore, we combined DCX+

cells. We found no effect of treatment (df = 2, F = 0.16, p = 0.85), hemisphere (df = 1, F

= 0.05, p = 0.83), or treatment by hemisphere interaction (df = 2, F = 0.39, p = 0.68) on

numbers of DCX+ cells in HVC (two-factor ANOVA, repeated measures on hemisphere,

Figure 4B). We also found no difference in new neuron numbers (neither BrdU+/Hu+,

nor DCX+) between hemispheres ipsilateral and contralateral to the TS nerve cut.

Unilateral TS-cuts had no effect on the lateralization index of BrdU+/Hu+ or DCX+

neurons in HVC (one-way ANOVA, df = 2, F < 0.01, p = 1.0 ; df = 2, F = 1.19, p = 0.33,

respectively, Figure 4C). We also found no relationship between the number BrdU+/Hu+

neurons or DCX+ cells in left HVC and right HVC of control animals (df = 5, r = 0.53, t =

1.38, p = 0.23; df = 5, r = 0.66, t = 1.1, p = 0.32, respectively, Figure 4D).

Effect of TS-cut on New Neurons in NCM. There was no main effect of treatment (F =

0.22, p = 0.80), hemisphere (F = 1.29, p = 0.27), or 3 way treatment by hemisphere

interaction (F = 0.83, p = 0.45) on numbers of BrdU+/Hu+ neurons in NCM (Figure 5A).

As in HVC, we found no difference among treatments or hemispheres when DCX+

fusiform and multipolar cells were analyzed separately; therefore, we combined these

categories. There was no main effect of treatment (F = 0.48, p = 0.63), hemisphere (F =

2.28, p = 0.15), or interaction on DCX+ cells in NCM (F = 0.14, p = 0.87, two-factor

ANOVA, repeated measures on hemisphere, Figure 5B).

Earlier work showed that in unmanipulated controls, there were more new neurons

in the left NCM than the right, and that this asymmetry was lost after a unilateral

tracheosyringeal nerve cut (Tsoi et al., 2014). Likewise, we found more BrdU[+]/Hu[+]

neurons in the left than the right NCM (paired sample t-test, t (11) = 2.26, p = 0.045) in

controls. Hemispheric asymmetry was not seen in either the left TS-cut or right TS-cut

group (paired t-test within group, p = 0.67, p = 0.74, respectively). Consistent with this

pattern, we found a significant difference in lateralization indices of BrdU+/Hu+ among

the three groups (df = 2, F = 3.92, p = 0.03). Interestingly, the lateralization index of only

the right TS-cut group differed from controls (HSD, p <0.05) whereas the lateralization

index of the left TS-cut group did not differ from that of controls or the right TS-cut group

(Figure 5C).

Unlike in HVC, in control birds, the numbers of BrdU+/Hu+ cells in left and right

NCM were positively correlated (df = 2, r = 0.69, t = 3.06, p = 0.01, Figure 5D) as in Tsoi

et al. (2014). There were no correlations between NCM hemispheres in either TS-cut

group (p = 0.45, left TS cut; p = 0.63, right TS-cut, data not shown). There were no

correlations in numbers of DCX+ cells between NCM hemispheres in any group (p >

0.05 for all, data not shown).

Effect of TS-cut on New Neurons in Area X. We found a significant difference in the

number of BrdU+/Hu+ neurons in Area X among the three treatment groups (two factor

repeated measures ANOVA, F = 7.47, p = 0.002, Figure 6A). There was no difference

between hemispheres (F = 0.25, p = 0.62) and no interaction (F = 1.16, p = 0.33).

Within the left hemisphere (one way ANOVA, df = 2, F = 3.31, p = 0.048), numbers of

BrdU+/Hu+ cells in both the left TS-cut and right TS-cut decreased compared to

controls (t = 2.05, p = 0.048; t = 2.39, p = 0.045, respectively). The same pattern was

found within the right hemisphere Area X (one way ANOVA, df = 2, F = 6.84, p = 0.003).

Compared to the right Area X of controls, there were fewer numbers of BrdU+/Hu+ cells

in the right Area X of both the left TS-cut and right TS-cut groups (t = 3.58, p = 0.002; t =

2.65, p = 0.012, respectively).

As in the other regions, we combined all DCX+ cells after finding no difference

within fusiform and round DCX+ cell types across treatments. There was no main effect

of treatment (2 factor mixed ANOVA, F = 0.07, p = 0.93), hemisphere (F = 1.48, p =

0.24), or interaction (F = 1.22, p = 0.32; Fig. 6B) on the number of DCX+ neurons in

Area X.

There was no significant difference among groups in the LI of BrdU+/Hu+ or DCX+

cells (one-way ANOVA, F = 0.2, p = 0.82; F = 0.20, p = 0.82, respectively, Figure 6C).

There were no correlations between the numbers of BrdU+/Hu+ neurons in the left and

right Area X of control animals (df = 10, r = 0.30, p = 0.35; Fig. 6D). However, DCX+

cells were correlated between hemispheres in control birds (df = 5, r = 0.92, p = 0.016,

Figure 6E) and in birds with a left side TS-cut (df = 7, r^2 = 0.85, p = 0.03), but not in birds

with a right side TS-cut (df = 6, r = 0.67, p = 0.27, data not shown).

Within Hemisphere Correlations in Numbers of New Neurons Between Adjacent

Regions. We were also interested in whether new neuron numbers may be linked

between functionally associated regions in control birds. However, we found no

correlations in numbers of BrdU+/Hu+ cells between NCM and HVC or between HVC

and Area X in either the left (df = 4, r = 0.69, p = 0.34; df = 4, r = 0.89, p = 0.063,

respectively) or right (df = 4, r = 0.69, p = 0.35; df = 5, r = 0.49, p = 0.61, respectively)

hemispheres. It is worth noting, however, that in each hemisphere removing a single

bird would have resulted in significant correlations between HVC and Area X and thus

larger sample sizes may find this to be the case. There were no correlations between

DCX+ cells within either hemisphere between NCM and HVC (left: df = 4, r = 0.48, p =

0.34; right: df = 4, r = 0.47, p = 0.35) or between HVC and Area X (left: df = 4, r = 0.79,

p = 0.063; right: df = 5, r = 0.23, p = 0.612, data not shown).

Discussion

This is a broad-strokes study, serving as a first pass in understanding whether the

quality of song structure affects new neuron survival and whether regulation of new

neuron survival is independent across brain region and hemisphere. We found that

irreversible disruption to song structure by unilateral denervation of the syrinx in adult

male zebra finches impacted the numbers of 28-30 day old neurons in a region and

hemisphere-specific manner. Altered song feedback had no effect on numbers of one

month old neurons in HVC, resulted in a loss of left-sided dominance of new neurons in

NCM consistent with Tsoi et al. (2014), and decreased neuron survival in both

hemispheres of Area X. These effects were not present in younger neurons expressing

DCX perhaps due to the lack of functional connectivity in this cohort at the time of

treatment.

HVC. The clearest example of a behavior that impacts new neurons in HVC is singing

(Li et al., 2000; Alvarez-Borda and Nottebohm, 2002; Ball et al., 2002; Alward et al.,

2016). However, functional implications of upregulating new neurons during singing are

not clear. In the non-seasonal zebra finch, new neurons appear to be added in surplus

of replacing dying cells as HVC neuronal packing density increases with increasing bird

age (Walton et al., 2012). Therefore, song production, and the increased density of

adult-formed neurons, may have the outcome of fine tuning motor stereotypy, or

improved resolution of fine motor gestures, increasing the precision of song structure

c.f.(Polomova et al., 2019). Consistent with the idea that new neurons contribute to

precision, song stereotypy continues to increase with age in adult zebra finches

(Brainard and Doupe, 2001; Pytte et al., 2007; James and Sakata, 2019). The proposal

that new neurons in the HVC-RA motor pathway increase the motor resolution for song

production borrows from the model that new neurons in the rodent hippocampus

contribute to memory resolution, providing improved pattern separation (Aimone et al.,

2011) and suggests a potential general rule of new neuron function in widely different

substrates.

With increased bird age, and thus increased packing density of HVC-RA neurons

(Walton et al., 2012), the motor pattern for song production is also increasingly resistant

to degradation after deafening (Lombardino and Nottebohm, 2000). Lombardino and

Nottebohm (2000) speculated that each time a bird sings, learning, or "motor

engrainment", cumulatively strengthens the memory of the song motor commands,

decreasing online reliance on auditory feedback. Combining these findings, perhaps the

act of singing, by upregulating new neurons, increases the substrate density for

encoding the motor pattern, resulting in greater motor pattern precision and durability.

This is consistent with the finding that more new neurons in HVC correspond to a more

stable song post-deafening (Pytte et al., 2012).

Independent of this model but compatible with it, it has been suggested that as

songs progress toward a target goal during development, seasonally, or with increased

stereotypy with age, feedback from continually improving song quality may actively

promote new neuron survival in HVC (Wilbrecht and Kirn, 2004; Kirn, 2010). Here we

examined whether altered song feedback, without song improvement, affected numbers

of new neurons. We found there was no group effect of the TS-cut on numbers of new

neurons in HVC, examining the left or right hemisphere independently and combined,

and also separately assessing the HVC hemisphere ipsilateral and contralateral to the

TS nerve cut. Thus the negative results reported here are consistent with the alternative

hypothesis that song improvement, i.e., progress toward a target, rather than altered

feedback per se is necessary for increased survival of new neurons in HVC (Pytte et al.,

2011).

NCM. NCM is thought to play a role in the recognition of conspecific songs (Chew et

al., 1996) and to encode at least one of potentially several distributed copies of a tutor's

song used in song learning (Solis and Doupe, 1997; Phan et al., 2006; Bolhuis and

Moorman, 2015). Neurons in NCM selectively respond to conspecific songs, the song of

the bird's tutor, as well as playback of the bird's own song (Grace et al., 2003; Hsu et

al., 2004). Moreover, NCM is active while a bird is singing, suggesting it may process

auditory feedback from song in real time (Jarvis et al., 1998). Motor recovery from song

distortion requires an intact NCM, although it is not clear whether this reflects access to

a stored representation of the bird's own song or that of the tutor (Canopoli et al., 2014).

NCM receives a robust influx of new neurons throughout adulthood, although very

little is known about the function of new neurons added. We know that new neurons are

upregulated in birds housed in groups (Lipkind et al., 2002; Barnea et al., 2006; Adar et

al., 2008a; Adar et al., 2008b) and are diminished in birds that have been deafened

(Pytte et al., 2010). Together, this suggests that activity in NCM promotes new neuron

survival; however, does not shed light on function. In a careful study of new neuron

survival across the rostral-caudal extent of NCM, Barnea et al. (2006) proposed an

interesting model describing how new neurons may encode information about the

calendar timepoint when an event (song stimulus) was experienced. More generally,

the continual addition of new neurons may preserve and/or improve the resolution of

stored song memories in either hemisphere of NCM, and perhaps enhance song

discrimination in general.

Our interest is in whether new neuron survival may be influenced by the degree to

which song feedback matches expected feedback, perhaps with respect to a stored

template of the bird's own song. In control birds, we found more new neurons in the left

than right NCM, consistent with Tsoi et al., (2014). This left-side lateralization in new

neurons was lost in the TS-cut bird. Although not statistically significant, the loss in

lateralization was due more to a decrease in numbers of new neurons in the left NCM

than an increase in new neurons in the right NCM. Thus, we hypothesize that a

template of the bird's own song may be lateralized to the left, rather than right, NCM.

Evidence for a lateralized response in NCM to the bird's own song is mixed.

Interestingly, estradiol synthesis in the left NCM but not the right NCM is necessary for a

bird's preference for his own song (Remage-Healey et al., 2010). In a behavioral study,

adult male zebra finches showed deficits in discriminating their own song from that of a

cage-mate when the left side ovoidalis, part of the ascending auditory pathway, was

lesioned and not when the right side was lesioned (Cynx et al., 1992). However,

selectivity for playback of the bird's own song was found to be lateralized toward the

right side in the midbrain dorsal lateral mesencephalic nucleus (MLd) the analog of the

mammalian inferior colliculus (Poirier et al., 2009). Like ovoidalis, MLd is part of the

unilateral ascending auditory pathway that eventually projects to NCM. Moreover, both

left and right NCM respond to playbacks of the bird's song (Soyman and Vicario, 2017).

In a different model, hemispheric asymmetry may be based on sound acoustic

features rather than categories of own song vs those of others (Van Ruijssevelt et al.,

2017). Playback of conspecific songs with filtered spectral structure increased BOLD

responses in the left NCM and slightly decreased BOLD responses in the right NCM

compared to nonmanipulated songs (Van Ruijssevelt et al., 2017). Because spectral

filtering removed frequency information from the song leaving primarily temporal

content, the authors interpreted this finding as indicating that the left NCM is specialized

for processing temporal information. They suggested that this asymmetry is normally

masked by the spectral content of whole songs (Van Ruijssevelt et al., 2017). On the

other hand, we suggest that perhaps the increased response in the left NCM reflects a

left-dominant sensitivity to aberrant information, resonating with our findings. The left

hemisphere of NCM has been shown to be sensitive to novel auditory experience more

broadly (Moorman et al., 2015; Yang and Vicario, 2015) consistent with the idea that

NCM (bilaterally) is sensitive to expectation (Lu and Vicario, 2017; Dong and Vicario,

2018). Thus perhaps it is the novelty of the distorted song, and not feedback about the

bird's own song *per se,* that underlies the observed decrease in new neurons in the left

NCM.

It is a consistent finding across studies that the magnitude of asymmetry in various

measures of NCM corresponds to performance in learning (Moorman and Nicol, 2015).

For instance, the degree of left side dominance in new neurons (Tsoi et al., 2014), and

left side dominance in activity during sleep (Moorman et al., 2015) predict accuracy of

song learning of the bird's own song. Left side dominance in NCM activity in response

to song playback is also correlated with rate of learning in a conspecific song

discrimination task (Bell et al., 2015). The results of the current study and that of Tsoi et

al., (2014) add that lateralization in new neurons is flexible and can be disrupted by the

bird's experience.

AREA X. We found significantly fewer new neurons in both hemispheres of Area X in

birds who received TS nerve cuts. Area X plays a role in juvenile song learning

(Sohrabji et al., 1990; Scharff and Nottebohm, 1991; Olveczky et al., 2005) and

modification of the song motor pattern in adults in response to social context, deafening,

and perturbations in feedback, although the mechanism by which it does so is debated

(Hessler and Doupe, 1999; Brainard and Doupe, 2000; Ali et al., 2013; Woolley et al.,

2014; Woolley and Kao, 2015; Woolley, 2016; Kojima et al., 2018) c.f. (Sanchez-

Valpuesta et al., 2019). In adult birds, the anterior forebrain pathway (AFP) more

generally mediates differences in song variability (Hessler and Doupe, 1999; Williams

and Mehta, 1999; Woolley et al., 2014).

As in HVC (Walton et al., 2012) new neurons are continually added to Area X,

increasing packing density with age, and are incorporated into functioning circuits

(Kosubek-Langer et al., 2017). Approximately 80% of new neurons express D1 and D2

receptors, become medium spiny neurons (MSN), and fire during singing (Kosubek-

Langer et al., 2017). New MSN are also innervated by glutamatergic synapses,

putatively from HVC and lMAN, by day 31 after birth dating (Kosubek-Langer et al.,

2017). Kosubek-Langer et al., (2017) suggest that the newly recruited MSNs fulfill the

same function as the existing older MSNs. These authors speculate that the continual

addition of MSNs results in relatively stronger inhibition by this increasing population on

the unchanged number of pallidal-like neurons. Downstream DLM then would diminish

excitation of LMAN and subsequently, of RA. Kosubek-Langer et al., (2017) note that

this model predicts diminishing AFP influence over the motor pathway as birds age,

consistent with behavioral studies (Lombardino and Nottebohm, 2000; Brainard and

Doupe, 2001; Pytte et al., 2007).

Song variability or degradation of acoustic structure can be induced experimentally

by distorting auditory feedback via tracheosyringeal nerve cut or deafening (Nordeen

and Nordeen, 1992; Williams and McKibben, 1992; Hough and Volman, 2002; Nordeen

and Nordeen, 2010). The AFP seems to mediate this degradation process, since

lesions of the AFP output nucleus LMAN prevent song deterioration after auditory

feedback distortion (Brainard and Doupe, 2000; Nordeen and Nordeen, 2010). In

addition, our results suggest that addition and/or survival of new neurons in Area X is

not immune to behavioral feedback and may be influenced by the quality of song

structure.

The mechanism by which altered song may result in decreased survival of new

neurons in Area X is not known, but may involve modulation of dopamine. Dopamine

interacts with mechanisms of new neuron survival (Appeltants et al., 2000; Cornil et al.,

2008; Leblois et al., 2010) and plays an important role in learning behaviors that are

reinforced by rewarding outcomes. Most of the song system nuclei receive

dopaminergic input (Lewis et al., 1981; Bottjer, 1993; Appeltants et al., 2000; Appeltants

et al., 2002). In particular, the substantia nigra pars compacta (SNc) and the ventral

tegmental area (VTA) send direct dopaminergic projections to Area X (Sasaki et al.,

2006). Through evaluation of predicted and actual auditory feedback, perhaps

dopaminergic signals modulate AFP output and influence song variability and learning.

Accurate song feedback may maintain dopamine activity and, in turn, new neuron

survival. Activation of dopaminergic neurons in Area X occurs in several contexts

(Scharff and Nottebohm, 1991; Kao and Brainard, 2006; Sasaki et al., 2006), including

in response to playback of the bird's own song (Gale and Perkel, 2010). It has

previously been demonstrated that singing induces dopamine release from SNc and

VTA into Area X (Sasaki et al., 2006). Likewise, it has been found that Area X may

make an indirect connection to SNc/VTA via a projection to ventral pallium (VP) (Gale et

al., 2008). Through this indirect pathway, the modulation of Area X output drives

changes in the firing rate of dopaminergic neurons in the SNc and VTA via the VP. This

relationship may suggest that Area X plays a role in feedback evaluation of song and

transmits this information to dopaminergic neurons. Perhaps this pathway results in

decreased dopamine signaling in the TS-cut birds with mismatched feedback and fewer

new neurons in Area X. Midbrain dopaminergic neurons send profuse projections to the

striatum, including Area X (Kitt and Brauth, 1986; Durstewitz et al., 1999), and much

sparser projections to the pallium, which includes HVC and NCM (Kitt and Brauth,

1986). Therefore, dopamine may be less influential in mediating new neuron survival in

HVC and NCM.

Numbers of New Neurons Between Regions. Few studies have made inter-regional

comparisons of adult neurogenesis in the mammalian brain. Notably, however, prenatal

stress has been shown to decrease neurogenesis in the mouse hippocampus in

adulthood, with no effect on neurogenesis in the olfactory bulb (Belnoue et al., 2013). In

the aging pigeon, the extent of new neuron decline is greater in the olfactory bulb than

in the hippocampus, again differentiating regulatory mechanisms between neurogenic

regions (Meskenaite et al., 2016). Rates of decline in new neurons in young adults also

differ between HVC and Area X (Pytte et al., 2007). Our results are consistent with the

idea that regulatory mechanisms operate independently across brain regions, even

those linked functionally or anatomically.

Taken together… Song learning by juveniles requires incremental improvement toward

a target motor behavior and requires both Area X and HVC, in concert with neuronal

tuning to the tutor and the bird's own song in NCM. Moreover, it occurs at a time when

new neuron incorporation is high. In one of many models of song learning, akin to

procedural motor learning more broadly, Area X and the AFP is thought to evaluate

feedback and influence subsequent motor commands produced by HVC, and the output

of feedback evaluation is thought to influence subsequent motor commands. Our

findings suggest that after the juvenile stage of song learning, the accumulation of new

neurons in Area X may correspond directly to the quality of song feedback in matching

to the target own-song template. On the other hand, new neurons in HVC are not culled

or maintained based on the output of a match between expected and received

feedback. New neurons in HVC may instead either 1) be influenced via feedback in

judging progress toward a goal of an ideal song or 2) are not affected by song feedback

at all, and instead new neurons may drive progress toward a goal song in a feedforward

direction only (Pytte et al., 2011). As in bilateral Area X, new neurons in left NCM may

be culled when song feedback no longer matches expected feedback of the bird's own

song. Thus, perhaps increasing the resolution of the neuronal substrate in the song

system and NCM by increasing new neurons serves to achieve increased song

stereotypy (Pytte et al., 2007) both by better resolution of song feedback evaluation

and/or by fine tuning motor output.

Conclusions. This study contributes to our understanding of whether mismatched

behavioral feedback affects new neuron recruitment and survival in brain regions that

subserve and monitor those behaviors. We found that altering song production via

unilateral tracheosyringeal denervation resulted in decreased 28-30 day old neurons in

Area X, loss of lateralization of neurons in NCM, and had no effect in HVC. This

indicates that the effects of syringeal denervation on neurogenesis vary depending both

on the region and hemisphere that receives new neurons.

Acknowledgements. John Kirn generously gifted birds for the study. Ben Koo

contributed to tissue processing and cell quantification.

This article is dedicated to John R. Kirn (1952-2019).

References

Adar E, Lotem A, Barnea A. 2008a. The effect of social environment on singing

behavior in the zebra finch (Taeniopygia guttata) and its implication for neuronal

recruitment. Behav Brain Res 187:178-184.

Adar E, Nottebohm F, Barnea A. 2008b. The relationship between nature of social

change, age, and position of new neurons and their survival in adult zebra finch

brain. J Neurosci 28:5394-5400.

Aimone JB, Deng W, Gage FH. 2011. Resolving new memories: a critical look at the

dentate gyrus, adult neurogenesis, and pattern separation. Neuron 70:589-596.

Ali F, Otchy TM, Pehlevan C, Fantana AL, Burak Y, Olveczky BP. 2013. The basal

ganglia is necessary for learning spectral, but not temporal, features of birdsong.

Neuron 80:494-506.

Alvarez-Borda B, Haripal B, Nottebohm F. 2004. Timing of brain-derived neurotrophic

factor exposure affects life expectancy of new neurons. Proc Natl Acad Sci U S A

101:3957-3961.

Alvarez-Borda B, Nottebohm F. 2002. Gonads and singing play separate, additive roles

in new neuron recruitment in adult canary brain. J Neurosci 22:8684-8690.

Alward BA, Madison FN, Parker SE, Balthazart J, Ball GF. 2016. Pleiotropic Control by

Testosterone of a Learned Vocal Behavior and Its Underlying

Neuroplasticity(1,2,3). eNeuro 3.

Appeltants D, Absil P, Balthazart J, Ball GF. 2000. Identification of the origin of

catecholaminergic inputs to HVc in canaries by retrograde tract tracing combined

with tyrosine hydroxylase immunocytochemistry. J Chem Neuroanat 18:117-133.

Appeltants D, Ball GF, Balthazart J. 2002. The origin of catecholaminergic inputs to the

song control nucleus RA in canaries. Neuroreport 13:649-653.

Ball GF, Auger CJ, Bernard DJ, Charlier TD, Sartor JJ, Riters LV, Balthazart J. 2004.

Seasonal plasticity in the song control system: multiple brain sites of steroid

hormone action and the importance of variation in song behavior. Ann N Y Acad

Sci 1016:586-610.

Ball GF, Madison FN, Balthazart J, Alward BA. 2020. How does testosterone act to

regulate a multifaceted adaptive response? Lessons from studies of the avian

song system. J Neuroendocrinol 32:e12793.

Ball GF, Riters LV, Balthazart J. 2002. Neuroendocrinology of song behavior and avian

brain plasticity: multiple sites of action of sex steroid hormones. Front

Neuroendocrinol 23:137-178.

Barnea A, Mishal A, Nottebohm F. 2006. Social and spatial changes induce multiple

survival regimes for new neurons in two regions of the adult brain: An anatomical

representation of time? Behav Brain Res 167:63-74.

Bell BA, Phan ML, Vicario DS. 2015. Neural responses in songbird forebrain reflect

learning rates, acquired salience, and stimulus novelty after auditory

discrimination training. J Neurophysiol 113:1480-1492.

Belnoue L, Grosjean N, Ladeveze E, Abrous DN, Koehl M. 2013. Prenatal stress

inhibits hippocampal neurogenesis but spares olfactory bulb neurogenesis. PLoS

One 8:e72972.

Bolhuis JJ, Moorman S. 2015. Birdsong memory and the brain: in search of the

template. Neurosci Biobehav Rev 50:41-55.

Bottjer SW. 1993. The distribution of tyrosine hydroxylase immunoreactivity in the brains

of male and female zebra finches. J Neurobiol 24:51-69.

Brainard MS, Doupe AJ. 2000. Auditory feedback in learning and maintenance of vocal

behaviour. Nat Rev Neurosci 1:31-40.

Brainard MS, Doupe AJ. 2001. Postlearning consolidation of birdsong: stabilizing effects

of age and anterior forebrain lesions. J Neurosci 21:2501-2517.

Brenowitz EA, Larson TA. 2015. Neurogenesis in the adult avian song-control system.

Cold Spring Harb Perspect Biol 7.

Canopoli A, Herbst JA, Hahnloser RH. 2014. A higher sensory brain region is involved

in reversing reinforcement-induced vocal changes in a songbird. J Neurosci

34:7018-7026.

Chew SJ, Vicario DS, Nottebohm F. 1996. A large-capacity memory system that

recognizes the calls and songs of individual birds. Proc Natl Acad Sci U S A

93:1950-1955.

Cornil CA, Castelino CB, Ball GF. 2008. Dopamine binds to alpha(2)-adrenergic

receptors in the song control system of zebra finches (Taeniopygia guttata). J

Chem Neuroanat 35:202-215.

Cynx J, Williams H, Nottebohm F. 1992. Hemispheric differences in avian song

discrimination. Proc Natl Acad Sci U S A 89:1372-1375.

Dong M, Vicario DS. 2018. Neural Correlate of Transition Violation and Deviance

Detection in the Songbird Auditory Forebrain. Front Syst Neurosci 12:46.

Dupret D, Revest JM, Koehl M, Ichas F, De Giorgi F, Costet P, Abrous DN, Piazza PV.

2008. Spatial relational memory requires hippocampal adult neurogenesis. PLoS

One 3:e1959.

Durstewitz D, Kroner S, Gunturkun O. 1999. The dopaminergic innervation of the avian

telencephalon. Prog Neurobiol 59:161-195.

Gale SD, Perkel DJ. 2010. A basal ganglia pathway drives selective auditory responses

in songbird dopaminergic neurons via disinhibition. J Neurosci 30:1027-1037.

Gale SD, Person AL, Perkel DJ. 2008. A novel basal ganglia pathway forms a loop

645   linking a vocal learning circuit with its dopaminergic input. J Comp Neurol

646   508:824-839.

647 George I, Cousillas H, Richard JP, Hausberger M. 2005. New insights into the auditory

648   processing of communicative signals in the HVC of awake songbirds.

649   Neuroscience 136:1-14.

650 Gheusi G, Cremer H, McLean H, Chazal G, Vincent JD, Lledo PM. 2000. Importance of

651   newly generated neurons in the adult olfactory bulb for odor discrimination. Proc

652   Natl Acad Sci U S A 97:1823-1828.

653 Grace JA, Amin N, Singh NC, Theunissen FE. 2003. Selectivity for conspecific song in

654   the zebra finch auditory forebrain. J Neurophysiol 89:472-487.

655 Hall ZJ, Tropepe V. 2018. Visual Experience Facilitates BDNF-Dependent Adaptive

656   Recruitment of New Neurons in the Postembryonic Optic Tectum. J Neurosci

657   38:2000-2014.

658 Hessler NA, Doupe AJ. 1999. Social context modulates singing-related neural activity in

659   the songbird forebrain. Nat Neurosci 2:209-211.

660 Hough GE, 2nd, Volman SF. 2002. Short-term and long-term effects of vocal distortion

661   on song maintenance in zebra finches. J Neurosci 22:1177-1186.

662 Hsu A, Woolley SM, Fremouw TE, Theunissen FE. 2004. Modulation power and phase

663   spectrum of natural sounds enhance neural encoding performed by single

664   auditory neurons. J Neurosci 24:9201-9211.

665 James LS, Sakata JT. 2019. Developmental modulation and predictability of age-

666   dependent vocal plasticity in adult zebra finches. Brain Res 1721:146336.

Jarvis ED, Scharff C, Grossman MR, Ramos JA, Nottebohm F. 1998. For whom the bird

sings: context-dependent gene expression. Neuron 21:775-788.

Kao MH, Brainard MS. 2006. Lesions of an avian basal ganglia circuit prevent context-

dependent changes to song variability. J Neurophysiol 96:1441-1455.

Khodosevich K, Lazarini F, von Engelhardt J, Kaneko H, Lledo PM, Monyer H. 2013.

Connective tissue growth factor regulates interneuron survival and information

processing in the olfactory bulb. Neuron 79:1136-1151.

Kirn J, O'Loughlin B, Kasparian S, Nottebohm F. 1994. Cell death and neuronal

recruitment in the high vocal center of adult male canaries are temporally related

to changes in song. Proc Natl Acad Sci U S A 91:7844-7848.

Kirn JR. 2010. The relationship of neurogenesis and growth of brain regions to song

learning. Brain Lang 115:29-44.

Kirn JR, Fishman Y, Sasportas K, Alvarez-Buylla A, Nottebohm F. 1999. Fate of new

neurons in adult canary high vocal center during the first 30 days after their

formation. J Comp Neurol 411:487-494.

Kirn JR, Schwabl H. 1997. Photoperiod regulation of neuron death in the adult canary. J

Neurobiol 33:223-231.

Kitt CA, Brauth SE. 1986. Telencephalic projections from midbrain and isthmal cell

groups in the pigeon. II. The nigral complex. J Comp Neurol 247:92-110.

Kojima S, Kao MH, Doupe AJ, Brainard MS. 2018. The Avian Basal Ganglia Are a

Source of Rapid Behavioral Variation That Enables Vocal Motor Exploration. J

Neurosci 38:9635-9647.

Kosubek-Langer J, Schulze L, Scharff C. 2017. Maturation, Behavioral Activation, and

Connectivity of Adult-Born Medium Spiny Neurons in a Striatal Song Nucleus.

Front Neurosci 11:323.

Krzyzanski W, Brier ME, Creed TM, Gaweda AE. 2013. Reticulocyte-based estimation

of red blood cell lifespan. Exp Hematol 41:817-822.

Larson TA, Thatra NM, Hou D, Hu RA, Brenowitz EA. 2019. Seasonal changes in

neuronal turnover in a forebrain nucleus in adult songbirds. J Comp Neurol

527:767-779.

Larson TA, Wang TW, Gale SD, Miller KE, Thatra NM, Caras ML, Perkel DJ, Brenowitz

EA. 2013. Postsynaptic neural activity regulates neuronal addition in the adult

avian song control system. Proc Natl Acad Sci U S A 110:16640-16644.

Leblois A, Wendel BJ, Perkel DJ. 2010. Striatal dopamine modulates basal ganglia

output and regulates social context-dependent behavioral variability through D1

receptors. J Neurosci 30:5730-5743.

Lewis JW, Ryan SM, Arnold AP, Butcher LL. 1981. Evidence for a catecholaminergic

projection to area X in the zebra finch. J Comp Neurol 196:347-354.

Li XC, Jarvis ED, Alvarez-Borda B, Lim DA, Nottebohm F. 2000. A relationship between

behavior, neurotrophin expression, and new neuron survival. Proc Natl Acad Sci

U S A 97:8584-8589.

Lipkind D, Nottebohm F, Rado R, Barnea A. 2002. Social change affects the survival of

new neurons in the forebrain of adult songbirds. Behav Brain Res 133:31-43.

Liu PZ, Nusslock R. 2018. Exercise-Mediated Neurogenesis in the Hippocampus via

BDNF. Front Neurosci 12:52.

Lombardino AJ, Nottebohm F. 2000. Age at deafening affects the stability of learned

713   song in adult male zebra finches. J Neurosci 20:5054-5064.

714   Lu K, Vicario DS. 2017. Familiar But Unexpected: Effects of Sound Context Statistics on

715   Auditory Responses in the Songbird Forebrain. J Neurosci 37:12006-12017.

716   Macdonald RA. 1961. "Lifespan" of liver cells. Autoradio-graphic study using tritiated

717   thymidine in normal, cirrhotic, and partially hepatectomized rats. Arch Intern Med

718   107:335-343.

719   McCasland JS. 1987. Neuronal control of bird song production. J Neurosci 7:23-39.

720   Moorman S, Gobes SM, van de Kamp FC, Zandbergen MA, Bolhuis JJ. 2015. Learning-

721   related brain hemispheric dominance in sleeping songbirds. Sci Rep 5:9041.

722   Moorman S, Nicol AU. 2015. Memory-related brain lateralisation in birds and humans.

723   Neurosci Biobehav Rev 50:86-102.

724   Mu Y, Gage FH. 2011. Adult hippocampal neurogenesis and its role in Alzheimer's

725   disease. Mol Neurodegener 6:85.

726   Ninkovic J, Mori T, Gotz M. 2007. Distinct modes of neuron addition in adult mouse

727   neurogenesis. J Neurosci 27:10906-10911.

728   Nordeen KW, Nordeen EJ. 1992. Auditory feedback is necessary for the maintenance of

729   stereotyped song in adult zebra finches. Behav Neural Biol 57:58-66.

730   Nordeen KW, Nordeen EJ. 2010. Deafening-induced vocal deterioration in adult

731   songbirds is reversed by disrupting a basal ganglia-forebrain circuit. J Neurosci

732   30:7392-7400.

733   Olveczky BP, Andalman AS, Fee MS. 2005. Vocal experimentation in the juvenile

734   songbird requires a basal ganglia circuit. PLoS Biol 3:e153.

Pawlisch BA, Remage-Healey L. 2015. Neuroestrogen signaling in the songbird

auditory cortex propagates into a sensorimotor network via an 'interface' nucleus.

Neuroscience 284:522-535.

Pfisterer U, Khodosevich K. 2017. Neuronal survival in the brain: neuron type-specific

mechanisms. Cell Death Dis 8:e2643.

Phan ML, Pytte CL, Vicario DS. 2006. Early auditory experience generates long-lasting

memories that may subserve vocal learning in songbirds. Proc Natl Acad Sci U S

A 103:1088-1093.

Poirier C, Boumans T, Verhoye M, Balthazart J, Van der Linden A. 2009. Own-song

recognition in the songbird auditory pathway: selectivity and lateralization. J

Neurosci 29:2252-2258.

Polomova J, Lukacova K, Bilcik B, Kubikova L. 2019. Is neurogenesis in two songbird

species related to their song sequence variability? Proc Biol Sci 286:20182872.

Pozner T, Vistoropsky Y, Moaraf S, Heiblum R, Barnea A. 2018. Questioning

Seasonality of Neuronal Plasticity in the Adult Avian Brain. Sci Rep 8:11289.

Pytte C, Yu YL, Wildstein S, George S, Kirn JR. 2011. Adult neuron addition to the

zebra finch song motor pathway correlates with the rate and extent of recovery

from botox-induced paralysis of the vocal muscles. J Neurosci 31:16958-16968.

Pytte CL, George S, Korman S, David E, Bogdan D, Kirn JR. 2012. Adult neurogenesis

is associated with the maintenance of a stereotyped, learned motor behavior. J

Neurosci 32:7052-7057.

Pytte CL, Gerson M, Miller J, Kirn JR. 2007. Increasing stereotypy in adult zebra finch

song correlates with a declining rate of adult neurogenesis. Dev Neurobiol

67:1699-1720.

Pytte CL, Parent C, Wildstein S, Varghese C, Oberlander S. 2010. Deafening

decreases neuronal incorporation in the zebra finch caudomedial nidopallium

(NCM). Behav Brain Res 211:141-147.

Rasika S, Alvarez-Buylla A, Nottebohm F. 1999. BDNF mediates the effects of

testosterone on the survival of new neurons in an adult brain. Neuron 22:53-62.

Rasika S, Nottebohm F, Alvarez-Buylla A. 1994. Testosterone increases the recruitment

and/or survival of new high vocal center neurons in adult female canaries. Proc

Natl Acad Sci U S A 91:7854-7858.

Remage-Healey L, Coleman MJ, Oyama RK, Schlinger BA. 2010. Brain estrogens

rapidly strengthen auditory encoding and guide song preference in a songbird.

Proc Natl Acad Sci U S A 107:3852-3857.

Rochefort C, Gheusi G, Vincent JD, Lledo PM. 2002. Enriched odor exposure increases

the number of newborn neurons in the adult olfactory bulb and improves odor

memory. J Neurosci 22:2679-2689.

Sanchez-Valpuesta M, Suzuki Y, Shibata Y, Toji N, Ji Y, Afrin N, Asogwa CN, Kojima I,

Mizuguchi D, Kojima S, Okanoya K, Okado H, Kobayashi K, Wada K. 2019.

Corticobasal ganglia projecting neurons are required for juvenile vocal learning

but not for adult vocal plasticity in songbirds. Proc Natl Acad Sci U S A

116:22833-22843.

Sasaki A, Sotnikova TD, Gainetdinov RR, Jarvis ED. 2006. Social context-dependent

singing-regulated dopamine. J Neurosci 26:9010-9014.

Scharff C, Nottebohm F. 1991. A comparative study of the behavioral deficits following

lesions of various parts of the zebra finch song system: implications for vocal

learning. J Neurosci 11:2896-2913.

Shevchouk OT, Ball GF, Cornil CA, Balthazart J. 2017. Studies of HVC Plasticity in

Adult Canaries Reveal Social Effects and Sex Differences as Well as Limitations

of Multiple Markers Available to Assess Adult Neurogenesis. PLoS One

12:e0170938.

Shors TJ, Townsend DA, Zhao M, Kozorovitskiy Y, Gould E. 2002. Neurogenesis may

relate to some but not all types of hippocampal-dependent learning.

Hippocampus 12:578-584.

Sohrabji F, Nordeen EJ, Nordeen KW. 1990. Selective impairment of song learning

following lesions of a forebrain nucleus in the juvenile zebra finch. Behav Neural

Biol 53:51-63.

Solis MM, Doupe AJ. 1997. Anterior forebrain neurons develop selectivity by an

intermediate stage of birdsong learning. J Neurosci 17:6447-6462.

Soyman E, Vicario DS. 2017. Principles of auditory processing differ between sensory

and premotor structures of the songbird forebrain. J Neurophysiol 117:1266-

1280.

Spalding KL, Bhardwaj RD, Buchholz BA, Druid H, Frisen J. 2005. Retrospective birth

dating of cells in humans. Cell 122:133-143.

Stocker E, Heine WD. 1971. [Proliferation and regeneration in liver and kidney of

juvenile rats. Autoradiographic studies after continuous infusion of 3H-thymidine

(author's transl)]. Verh Dtsch Ges Pathol 55:483-488.

Sun D. 2014. The potential of endogenous neurogenesis for brain repair and

regeneration following traumatic brain injury. Neural Regen Res 9:688-692.

Tchernichovski O, Nottebohm F, Ho CE, Pesaran B, Mitra PP. 2000. A procedure for an

automated measurement of song similarity. Anim Behav 59:1167-1176.

Tsoi SC, Aiya UV, Wasner KD, Phan ML, Pytte CL, Vicario DS. 2014. Hemispheric

asymmetry in new neurons in adulthood is associated with vocal learning and

auditory memory. PLoS One 9:e108929.

Van Ruijssevelt L, Washington SD, Hamaide J, Verhoye M, Keliris GA, Van der Linden

A. 2017. Song Processing in the Zebra Finch Auditory Forebrain Reflects

Asymmetric Sensitivity to Temporal and Spectral Structure. Front Neurosci

11:549.

Walton C, Pariser E, Nottebohm F. 2012. The zebra finch paradox: song is little

changed, but number of neurons doubles. J Neurosci 32:761-774.

Wilbrecht L, Kirn JR. 2004. Neuron addition and loss in the song system: regulation and

function. Ann N Y Acad Sci 1016:659-683.

Williams H, McKibben JR. 1992. Changes in stereotyped central motor patterns

controlling vocalization are induced by peripheral nerve injury. Behav Neural Biol

57:67-78.

Williams H, Mehta N. 1999. Changes in adult zebra finch song require a forebrain

nucleus that is not necessary for song production. J Neurobiol 39:14-28.

Winner B, Winkler J. 2015. Adult neurogenesis in neurodegenerative diseases. Cold

Spring Harb Perspect Biol 7:a021287.

Woolley SC. 2016. Social context differentially modulates activity of two interneuron

populations in an avian basal ganglia nucleus. J Neurophysiol 116:2831-2840.

Woolley SC, Kao MH. 2015. Variability in action: Contributions of a songbird cortical-

basal ganglia circuit to vocal motor learning and control. Neuroscience 296:39-

47.

Woolley SC, Rajan R, Joshua M, Doupe AJ. 2014. Emergence of context-dependent

variability across a basal ganglia network. Neuron 82:208-223.

Yang LM, Vicario DS. 2015. Exposure to a novel stimulus environment alters patterns of

lateralization in avian auditory cortex. Neuroscience 285:107-118.

Figure 1

Figure 1. **Diagram of the song system in a sagittal section, showing the vocal motor pathway (black solid lines) and the anterior forebrain pathway (dashed and white lines).** White lines show the feedback loop between the striatum, thalamus, nidopallium, and projection back to the striatum. AV = Nucleus avalanche; B = nucleus basorostralis; DM = dorsal medial nucleus; DLM = dorsal lateral nucleus of the medial thalamus; E = entopallium; LMAN = lateral magnocellular nucleus of the anterior nidopallium; LMO = lateral oval nucleus of the mesopallium; Nif = interfacial nucleus of the nidopallium; PAm = para-ambiguus; Ram = nucleus retroambiguus; Uva, nucleus uvaeformis; nXIIts = tracheosyringeal portion of the hypoglossal nucleus. Reproduced from Pytte et al., (2011); originally printed in Wilbrecht and Kirn (2004). Syrinx drawn by Franz Goller.

Figure 2

Figure 2. **Photomicrographs of doublecortin-positive neurons in HVC.** A. Fusiform, approximately 1-2 weeks old as indicated by its thin, elongated soma and unipolarity and B. Multipolar, approximately 2-3 weeks old as indicated by large, round soma and multiple processes (age estimates following Balthazart, et al., 2008). (C) Fluorescent markers used to identify 28-30 d old neurons in HVC, BrdU$^+$ nucleus visualized with FITC filter (D) Hu$^+$ neurons seen under rhodamine filter (E) BrdU$^+$/Hu$^+$ neuron showing colocalization of the two markers (rhodamine/FITC filter) in the same field of view. Scale bars A, B = 25 µm; C, D = 10 µm.

Figure 3

Figure 3. A. **Effects of TS-cuts on song structure.** A. Birds that underwent left (n = 5) or right (n = 4) TS-cut had accuracy scores that were significantly lower than those of controls (n = 2), demonstrating that unilateral tracheosyringeal nerve cut impacts the quality of song production. B. There were no significant differences in percentage similarity scores between pre-operative and post-operative songs of control (n = 2), left TS-cut (n = 5), and right TS-cut birds (n = 4). ** = p < .01

Figure 4

Figure 4. **Numbers of new neurons in HVC as a function of hemisphere and experimental condition.** A. Neither left (n = 7) nor right (n = 8) TS-cut affected the number of 28-30 d old BrdU+/Hu+ neurons in HVC compared to controls (n = 7). B. Neither left (n = 8) nor right (n = 5) TS-cut affected the number of ~1-3 week old DCX+ neurons in HVC compared with controls (n = 7). C. There were no differences in the Lateralization Index of BrdU+/Hu+ cells among treatment groups. D. There was no correlation between numbers of BrdU+/Hu+ cells between the left and right HVC.

Figure 5

Figure 5. **Numbers of new neurons in NCM as a function of hemisphere and experimental condition.** A. Left (n = 12) and right (n = 11) TS-cuts had no effect on the overall number of 28-30 d old BrdU+/Hu+ neurons in NCM. Control birds (n = 12) had more BrdU+/Hu+ neurons in left NCM than the right and this asymmetry was not seen in the left or right TS-cut group. B. Neither left (n = 7) nor right (n = 6) TS-cut

differed from controls (n = 7) in numbers of DCX+ neurons in NCM. C. There was a positive correlation between numbers of 28-30 d old neurons in left NCM and right NCM of control birds. * = p < 0.01.
Figure 6

Figure 6. **Numbers of new neurons in Area X as a function of hemisphere and experimental condition.** A. Left (n = 12) and right (n = 14) TS-cuts significantly decreased the number of 28-30 day old BrdU+/Hu+ neurons in Area X compared to controls (n = 12). B. Neither left (n = 9) nor right (n = 8) TS-cut groups

differed from controls (n = 7) in numbers of DCX+ neurons in Area X. C. Lateralization indices for BrdU+/Hu+ neurons in Area X. LIs did not differ between controls, left TS-cut, and right TS-cut conditions. D. There was no relationship between the numbers of BrdU+/Hu+ neurons in left versus right Area X in controls. E. We found a significant correlation between left and right Area X in numbers of DCX+ neurons in controls. ** = $p < .01$