Vocal convergence in the multi-level society of Guinea baboons

Julia Fischer[1,2,3*], Franziska Wegdell[1,3], Franziska Trede[1,4], Federica Dal Pesco[1,3] &

Kurt Hammerschmidt[1,3]

[1]*Cognitive Ethology Laboratory, German Primate Center, Kellnerweg 4, 37077 Göttingen,*

Germany

[2]*Department of Primate Cognition, Georg August University Göttingen, Göttingen, Germany*

[3]*Leibniz ScienceCampus Primate Cognition, Göttingen, Germany*

[4]*Primate Genetics Laboratory, German Primate Center, Göttingen, Germany*

Corresponding author: jfischer@dpz.eu

ORCIDs

Julia Fischer 0000-0002-5807-0074

Franziska Wegdell 0000-0002-3108-2999

Franziska Trede 0000-0003-3690-1006

Federica Dal Pesco 0000-0003-2326-1185

Kurt Hammerschmidt 0000-0002-3430-2993

ABSTRACT

The extent to which nonhuman primate vocalizations are amenable to modification

through experience is relevant for understanding the substrate from which human

speech evolved. One road to investigate the flexibility in vocal production has been to

study differences in vocal behaviour between different social groups. We examined the

vocal behaviour of Guinea baboons, *Papio papio*, ranging in the Niokolo Koba National

Park in Senegal. Guinea baboons live in a multi-level society, with parties nested within

gangs. We investigated whether the acoustic structure of grunts of 30 male baboons of

five gangs differed in relation to social level and genetic relatedness. Males in this

species are philopatric, resulting in increased male relatedness within gangs and

parties. Grunts from members of the same gang were more similar to each other than

across gangs (N = 435 dyads), but for parties within gangs we found no evidence for

higher similarity (N = 169 dyads). Acoustic similarity did not correlate with genetic

relatedness. Our study provides evidence for acoustic convergence in male Guinea

baboon grunts; the observed nonlinear relationship between social level and acoustic

similarity may reflect the limits of the extent to which vocal accommodation is possible,

or even advantageous.

KEY WORDS: baboon, *Papio papio*, primate communication, relatedness, vocal

learning

BACKGROUND

The structure of nonhuman primate vocalisations appears to be highly conserved [1].

Nevertheless, modifications in vocal output in relation to experience appear to be possible

within species-specific constraints [2–4], which spurred debates about whether or not

nonhuman primates show evidence for vocal learning [5,6]. To overcome the futile

arguments about whether or not modification in vocal output should be taken as evidence for

imitation, we recently proposed a framework that aims to distinguish between different

mechanisms that may contribute to acoustic variation as a result of auditory experience [7];

see also [8] for similar considerations about vocal learning more generally.

Of particular relevance in the present context are two of the mechanisms, namely

'specific auditory facilitation' and 'learning from success'. Specific auditory facilitation is

based on the idea of a "common coding" framework, according to which perception and

action are represented by the same system [9,10]. So, listening to specific calls may increase

the likelihood of producing corresponding call variants by the listener. Variation in the

exposure to specific variants should translate into corresponding variation in acoustic

similarity. In other words, subjects that interact more frequently should be more similar to

each other than those that interact less frequently. Acoustic similarity may also result from

genetic relatedness, however. For instance, subjects that are highly related may also have a

similar morphology of the vocal production apparatus [11]. Before conclusions about the role

of experience can be drawn, it is therefore necessary to check whether potential acoustic

variation between individuals can (also) be explained by genetic distance.

We here set out to test the idea of auditory facilitation by comparing the acoustic

variation in male Guinea baboon, *Papio papio,* grunts. Guinea baboons are an intriguing

model to examine the influence of auditory experience and social group membership, as they

live in a nested multi-level society with male philopatry [12]. Males associate with specific

females and young in 'units' [13]. Around four to five such units form 'parties', and 2-3 parties

team up to form a 'gang' [14]. Males are spatially highly tolerant and show low levels of

aggression [15]. They also form differentiated and stable relationships with other males [16].

During affiliative interactions with other group members, males produce low frequency tonal

grunts (Fig. 1). Such grunts have been shown to increase the likelihood of a subsequent

affiliative interaction or infant handling [17].

If specific auditory facilitation affects the structure of calls, subjects that interact

frequently with one another should produce calls that are more similar to each other. In other

words, members of the same party should have the greatest similarity, and members of the

same gang should produce calls that are more similar to each other than to calls produced

by members of another gang. If genetic relatedness affects vocal structure, dyads that are

more highly related should reveal greater acoustic similarity.

METHODS

We obtained recordings of grunts from a total of 30 male baboons between 2010 and 2016.

Thirteen of the males were members of the "Mare" gang and 14 of the "Simenti" gang, while

3 males resided in three different neighbouring gangs ("R", "N", and "O"). The Mare gang

comprised two parties of 6 and 7 males each; the Simenti gang comprised two parties of 5

and 9 males each.

The vocalizations were recorded using Marantz PMD 661 recorders (D &M

Professional, Longford, U.K.) with Sennheiser directional microphones (K6 power module +

ME66 recording head; Sennheiser, Wedemark, Germany) equipped with Rycote windshields

(RycoteWindjammer, Rycote Ltd., Stroud, U.K.). We used Avisoft SASLab Pro (Avisoft

Bioacoustic, Berlin, Germany) to check the recording quality, and to label and extract grunts

with sufficient quality and low background noise. We only used calls recorded at a maximum

distance of 3 m.

We reduced the sampling frequency from 44.1 to 5 and 1 kHz to obtain an

appropriate frequency resolution for the estimation of acoustic features, and calculated two

1024 pt Fast Fourier Transformation (FFT), one resulting in a frequency range of 2500 Hz

(frequency resolution 5 Hz, temporal resolution 6.4ms) and a second FFT resulting in a

frequency range of 500 Hz (frequency resolution of 1 Hz, and a temporal resolution of

16 ms). The resulting frequency time spectra were analyzed with a custom software program

LMA 2019, which allows a visual control of the accuracy of parameter estimation [18,19]. We

focused on the following acoustic features: fundamental frequency (F0), peak frequency,

distribution of frequency amplitudes and noisiness expressed as Wiener entropy (see

Supplementary Material Table S1). In total, we included 780 grunts in the acoustic analysis.

On average, we used 26 calls per subject in the analysis (range: 5-127). The Mare and

Simenti gang males were represented by 390 and 366 grunts, respectively; the other three

gangs by 7-10 grunts.

To quantify the acoustic similarity, we used a discriminant function analysis with

which we obtained a pairwise F-value for the acoustic distance for each dyad. This approach

has been applied in different studies examining relationships between acoustic structure and

genetic or geographic distance [20–22]. To determine the F-values, we entered all 82

acoustic features into a stepwise discriminant function (DFA) with subject identity as

grouping variable. The selection criterion for acoustic features to enter the discriminant

function was P_{in} = 0.05 and to remove P_{out} = 0.1. The DFA used 33 acoustic features for the

individual discrimination. F-values were calculated using IBM SPSS version 26.0 (IBM,

Armonk, NY). To assess the classification result of the individual discrimination of male

grunts, we performed a permuted DFA [23].

We extracted DNA from fecal samples using the First-DNA all tissue kit (Genial®).

We characterized genetic variation by assessing the individual allele variation on 24

polymorphic autosomal microsatellite markers. The 24 markers were amplified using the

Multiplex PCR Kit (QIAGEN) and fluorescent-labelled primers. PCR products were separated

and detected through capillary gel electrophoresis on an ABI 3130xL Genetic Analyzer

(Applied Biosystems®, USA). Microsatellite allele sizes were evaluated using Gene Mapper

5 (Applied Biosystems®). One locus (D1s548) showed signs of null alleles and significant

deviations from Hardy-Weinberg equilibrium and was therefore excluded, resulting in a total

of 23 loci included in the relatedness estimation (calculated with MICRO-CHECKER - version

2.2.3 [24] and the PopGenReport R package - version 3.0.0 [25], respectively). We used the

R package "related version 1.0" [26,27] to estimate relatedness using R version 3.4.4 and

RStudio version 1.1.456. The Queller-Goodnight estimator [28] showed the best performance

and was thus chosen in the present analysis. This estimator ranges from -1 to 1. Negative

values indicate that dyads are less related than on average, while positive values indicate

that they are more highly related than on average (see [16] for a detailed description of the

analysis).

These and the following statistical analyses were conducted in the R environment

(version 3.6.2; [29]), using the RStudio interface (version 1.1.447; [30]). We used a mantel

matrix correlation test (package "vegan" version 2.5.6) to test the correlation between

acoustic and genetic variation. To test the effect of membership to a given social level, we

applied categorical mantel tests, using "Same gang membership" (Yes/No) as the categorical

predictor variable, and QG or F (transformed as ln(1+F)) as the continuous variable. This

analysis was based on 435 dyads. To study the effect of party membership, we only

considered dyads that belonged to the same gang (N = 169) and used a restricted

permutation approach where males were permuted between parties within gangs. We used

1,000 permutations in all analyses, except the one for the variation between parties within

gangs, where we used 10,000 permutations. The raw data and the code for statistical

analysis are deposited at

https://osf.io/h7q5r/?view_only=e24fb7b53e7a4c57a1fe67db5d2452ab

RESULTS

Confirming previous analyses, males were more highly related within gangs than between

gangs (P = 0.009, N = 435), though the average (mean ± se) relatedness did not differ much

within gangs (QG estimator: –0.003 ± 0.011, N = 169 dyads) and between gangs (–0.054 ±

0.014, N = 266 dyads). Within gangs, males in the same party were more highly related on

average (QG estimator: 0.024 ±0.024) than males that were not members of the same party

(-0.029 ± 0.015, P = 0.035).

Grunts could be assigned to the correct individual significantly more frequently than

by chance, with an average correct assignment of 38.1 % (chance level 3.3%, pDFA, P <

0.001). Across the five gangs, the acoustic similarity did not correlate with genetic similarity (r

= -0.002, P = 0.497, Fig. S1). Yet, grunts of males within gangs were more similar to each

other than between gangs (categorical Mantel test, P = 0.001). The variation in acoustic

dissimilarity for within and between gangs is shown in Fig. 2A. Grunts of dyads in the same

gang had a (mean ± se) dissimilarity of F = 1.08 ± 0.02, while grunts of dyads in different

gangs had a mean dissimilarity of F = 1.25 ± 0.02. We found no evidence that grunts from

males within parties were more similar to each other than between parties in the same gang

(Fig. 2B, P = 0.6).

DISCUSSION

The structure of male grunts varied clearly between members of different gangs, but not

between members of parties within a gang. Males in the same gang were also more highly

related to one another, but this did not seem to account for the acoustic variation between

gangs, as evidenced by the lack of correlation between genetic and acoustic similarity.

Interaction frequency did not appear to account for the emergence of acoustic similarity

either, as grunts of males in the same party were not acoustically more similar to each other

than grunts of males in different parties in the same gang. This was somewhat surprising,

given that in a previous study, dyads within parties were observed within 100 m proximity

65% of the instances, while dyads of different parties were found in 100 m proximity in 24%

of the instances. Dyads of different gangs were observed within 100 m only 1 % of the time

[14]. Moreover, grunts occur almost exclusively within members of the same party [17].

The absence of a high degree of similarity within parties discounts the idea of a linear

relationship between auditory experience and vocal production. One possible explanation for

the observed nonlinear relationship between social level and acoustic similarity may be that

there are limits to the extent to which vocal accommodation is possible, or even

advantageous. Indeed, there may also be selective pressure to maintain individual identity.

Since the acoustic structure of grunts is relatively simple, this may additionally place a barrier

on vocal convergence.

Alternatively, gang membership may have a special social value that promotes the

adjustment of calls to that particular level. With the present data and in the absence of

experimental manipulation, we are not able to distinguish between these two hypotheses,

although the latter seems rather unlikely. Either way, it would be illuminating to track the

vocal output of males that transfer between different levels, but since males are philopatric

and dispersal is female-biased, such transfers are rare and rather unpredictable. We

therefore have no data to test to which degree and over which time course males would

adjust the structure of their calls.

The degree of variation in grunts between different social levels is relatively small

though and it is not clear whether subjects pay attention to this variation. In a previous study

[31], we tested male responses to the playbacks of grunts of males that share the same

home range as the study males ("neighbours") vs. to grunts of males living 50 km away

("strangers"). As control, we played back the grunts of males from the own gang.

Surprisingly, males responded strongly only to the grunts from males of the own gang, but

largely ignored neighbour or stranger males' calls (see also [32]). In principle, these

responses could be explained by recognition of the males' individual characteristics. Yet, it

might also be the case that males recognize the 'sound' of their own gang. Playbacks

presenting artificially created grunts bearing the own gang's characteristics vs. another

gang's characteristics would be needed to test this conjecture.

Taken together, we find evidence for a moderate degree of vocal convergence in

Guinea baboons. The magnitude of the change is difficult to compare to those of other

studies on nonhuman primates [3–5], given the differences in methodological approaches,

but broadly appears to be in a similar range. Indeed, all these studies reported minor

variation within the species-specific range. What does such group-specific variation tell us

about the substrate for the evolution of speech? Perhaps not so much, as the observed

variation could simply be a by-product of auditory experience, rather than volitional imitation,

as in the case of human speech. We therefore iterate that simpler explanations for vocal

plasticity need to be ruled out before we can safely claim to have identified precursors to

human vocal learning [see 7].

CONCLUSION

While this study provides evidence for acoustic convergence in male Guinea baboon grunts,

it also highlights that there are limits to the extent to which vocal accommodation is possible,

or even advantageous. More generally, our findings add to the body of evidence that within

species-specific constraints, subtle and (potentially) meaningful variation can be found in

nonhuman primate vocalizations. This variation does not compare to the open-ended

possibility of vocal imitation found in human speech, however.

Acknowledgments

We thank the Diréction des Parcs Nationaux (DPN) and Ministère de l'Environnement et de

la Protéction de la Nature (MEPN) de la République du Sénégal for permission to work in the

Niokolo Koba National Park. Particular thanks go to Mallé Guye for essential support, and

the field assistants and all team members of the CRP Simenti for long-term data collection.

Roger Mundry provided invaluable statistical advice. This research was supported by the

Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) - Project number

254142454 / GRK 2070, by the Leibniz ScienceCampus Primate Cognition funded by the

Leibniz Association, and by the Christian-Vogel Fonds of the Gesellschaft für Primatologie.

246   Author contributions

JF, KH and FW conceived the study, FW collected the data, FW and KH conducted the

acoustic analyses, FDP and FT conducted the genetic analysis, JF and KH did the statistical

analysis and prepared the figures, JF wrote the manuscript with input from all authors.

Figure Legends

Figure 1. Grunt examples from four different males. Frequency (kHz) on the y-axis, time (s)

on the x-axis. The spectrogram was created using Avisoft-SASLab Pro 5.2 (1.024 pt FFT,

sampling frequency: 11 kHz, time resolution: 2.9 ms, Flat Top window).

Figure 2. Acoustic variation between dyads that belong to (A) different gangs or the same

gang, and (B) different parties or the same party within a gang. Boxplots with medians and

interquartile range. Whiskers show values within 1.5 times of the inter-quartile range. Dots

represent dyadic values.

References

1.   Wegdell F, Hammerschmidt K, Fischer J. 2019 Conserved alarm calls but rapid

auditory learning in monkey responses to novel flying objects. *Nat. Ecol. Evol.* **3**,

1039–1042. (doi:10.1038/s41559-019-0903-5)

2.   Fischer J, Hage SR. 2019 Primate vocalization as a model for human speech: Scopes

and limits. In *Human Language: From Genes and Brains to Behavior* (ed P Hagoort),

pp. 639–656. Cambridge MA: MIT Press.

3.   Mitani JC, Hasegawa T, Gros-Louis J, Marler P, Byrne RW. 1992 Dialects in wild

chimpanzees? *Am. J. Primatol.* **27**, 233–243. (doi:10.1002/ajp.1350270402)

4.   Crockford C, Herbinger I, Vigilant L, Boesch C. 2004 Wild chimpanzees produce

group-specific calls: A case for vocal learning? *Ethology* **110**, 221–243.

(doi:10.1111/j.1439-0310.2004.00968.x)

5.   Watson SK, Townsend SW, Schel AM, Wilke C, Wallace EK, Cheng L, West V,

Slocombe KE. 2015 Vocal learning in the functionally referential food grunts of

chimpanzees. *Curr. Biol.* **25**, 495–499. (doi:10.1016/j.cub.2014.12.032)

6.   Fischer J, Wheeler BC, Higham JP. 2015 Is there any evidence for vocal learning in

chimpanzee food calls? *Curr. Biol.* **25**, R1028–R1029. (doi:10.1016/j.cub.2015.09.010)

7.   Fischer J, Hammerschmidt K. 2020 Towards a new taxonomy of primate vocal

production learning. *Philos. Trans. R. Soc. B Biol. Sci.* **375**, 20190045.

(doi:10.1098/rstb.2019.0045)

8.   Arriaga G, Jarvis ED. 2013 Mouse vocal communication system: Are ultrasounds

learned or innate? *Brain Lang.* **124**, 96–116. (doi:10.1016/j.bandl.2012.10.002)

9.   Prinz W, Neumann O. 1990 A common-coding approach to perception and action. In

Relationships between perception and action: Current approaches, pp. 167–201.

Berlin & New York: Springer.

10.   Jeannerod M. 1994 The representing brain: Neural correlates of motor intention and

imagery. *Behav. Brain Sci.* **17**, 187–202. (doi:10.1017/S0140525X00034026)

11.   Hauser MD. 1992 Articulatory and social factors influence the acoustic structure of

rhesus monkey vocalizations: A learned mode of production? *J. Acoust. Soc. Am.* **91**,

2175–2179. (doi:10.1121/1.403676)

12. Fischer J *et al.* 2017 Charting the neglected West: The social system of Guinea

baboons. *Am. J. Phys. Anthropol.* **162**, 15–31. (doi:10.1002/ajpa.23144)

13. Goffe AS, Zinner D, Fischer J. 2016 Sex and friendship in a multilevel society:

Behavioural patterns and associations between female and male Guinea baboons.

Behav. Ecol. Sociobiol. **70**, 323–336. (doi:10.1007/s00265-015-2050-6)

14. Patzelt A, Kopp GH, Ndao I, Kalbitzer U, Zinner D, Fischer J. 2014 Male tolerance and

male-male bonds in a multilevel primate society. *Proc. Natl. Acad. Sci. U. S. A.* **111**,

14740–14745. (doi:10.1073/pnas.1405811111)

15. Kalbitzer U, Heistermann M, Cheney DL, Seyfarth RM, Fischer J. 2015 Social

behavior and patterns of testosterone and glucocorticoid levels differ between male

chacma and Guinea baboons. *Horm. Behav.* **75**, 100–110.

(doi:10.1016/j.yhbeh.2015.08.013)

16. Dal Pesco F. 2019 Dynamics and fitness benefits of male-male sociality in wild Guinea

baboons (Papio papio). Georg-August-University Göttingen. See

http://hdl.handle.net/21.11130/00-1735-0000-0005-134F-E.

17. Faraut L, Siviter H, Pesco FD, Fischer J. 2019 How life in a tolerant society affects the

usage of grunts: evidence from female and male Guinea baboons. *Anim. Behav.* **153**,

83–93. (doi:10.1016/j.anbehav.2019.05.003)

18. Schrader L, Hammerschmidt K. 1997 Computer-aided analysis of acoustic parameters

in animal vocalizations: A multi-parametric approach. *Bioacoustics* **7**, 247–265.

(doi:10.1080/09524622.1997.9753338)

19. Fischer J, Noser R, Hammerschmidt K. 2013 Bioacoustic field research: A primer to

acoustic analyses and playback experiments with primates. *Am. J. Primatol.* **75**, 643–

663. (doi:10.1002/ajp.22153)

20. Naguib M, Hammerschmidt K, Wirth J. 2001 Microgeographic variation, habitat effects

and individual signature cues in calls of chiffchaffs *Phylloscopus collybita canarensis*.

Ethology **107**, 341–355. (doi:10.1046/j.1439-0310.2001.00669.x)

21.   Thinh VN, Hallam C, Roos C, Hammerschmidt K. 2011 Concordance between vocal

and genetic diversity in crested gibbons. *BMC Evol. Biol.* **11**, 36. (doi:10.1186/1471-

2148-11-36)

22.   Meyer D, Hodges JK, Rinaldi D, Wijaya A, Roos C, Hammerschmidt K. 2012 Acoustic

structure of male loud-calls support molecular phylogeny of Sumatran and Javanese

leaf monkeys (genus *Presbytis*). *BMC Evol. Biol.* **12**, 16. (doi:10.1186/1471-2148-12-

16)

23.   Mundry R, Sommer C. 2007 Discriminant function analysis with nonindependent data:

consequences and an alternative. *Anim. Behav.* **74**, 965–976.

(doi:10.1016/j.anbehav.2006.12.028)

24.   Van Oosterhout C, Hutchinson WF, Wills DPM, Shipley P. 2004 MICRO-CHECKER:

Software for identifying and correcting genotyping errors in microsatellite data. *Mol.*

Ecol. Notes (doi:10.1111/j.1471-8286.2004.00684.x)

25.   Gruber B, Adamack AT. 2015 landgenreport: A new R function to simplify landscape

genetic analysis using resistance surface layers. *Mol. Ecol. Resour.*

(doi:10.1111/1755-0998.12381)

26.   Wang J. 2011 Coancestry: a program for simulating, estimating and analysing

relatedness and inbreeding coefficients. *Mol. Ecol. Resour.* **11**, 141–145.

(doi:10.1111/j.1755-0998.2010.02885.x)

27.   Pew J, Muir PH, Wang J, Frasier TR. 2015 related: An R package for analysing

pairwise relatedness from codominant molecular markers. *Mol. Ecol. Resour.* **15**, 557–

561. (doi:10.1111/1755-0998.12323)

28.   Queller DC, Goodnight KF. 1989 Estimating relatedness using genetic markers.

Evolution (N. Y). **43**, 258–275. (doi:10.1111/j.1558-5646.1989.tb04226.x)

29.   R Core Team. 2019 R: A Language and Environment for Statistical Computing.

30.   RStudio Team. 2018 RStudio: Integrated Development for R.

31.   Maciej P, Patzelt A, Ndao I, Hammerschmidt K, Fischer J. 2013 Social monitoring in a

multilevel society: A playback study with male Guinea baboons. *Behav. Ecol.*

Sociobiol. **67**, 61–68. (doi:10.1007/s00265-012-1425-1)

32.   Faraut L, Fischer J. 2019 How life in a tolerant society affects the attention to social

information in baboons. *Anim. Behav.* **152**, 11–17.

(doi:10.1016/j.anbehav.2019.04.004)