The consequences of size-selective fishing mortality for larval production and

sustainable yield in species with obligate male care

Authors:

Holly K. Kindsvater[1], Kim Tallaksen Halvorsen[2], Tonje Knutsen Sørdalen[3,4], Suzanne H.

Alonzo[5]

Affiliations:

[1]Dept. of Fish and Wildlife Conservation, Virginia Polytechnic Institute and State

University, Blacksburg, VA 24060, USA

[2]Institute of Marine Research, Austevoll Research Station, Austevoll, Norway

[3]Institute of Marine Research, Flødevigen Research Station, His, Norway

[4]Centre for Coastal Research (CCR), University of Agder, 4604 Kristiansand, Norway

[5]Department of Ecology and Evolutionary Biology, University of California, Santa Cruz,

CA, 95060, USA

Abstract

Size-based harvest limits or gear regulations are often used to manage fishing mortality

and ensure the spawning biomass of females is sufficiently protected. Yet, management

interactions with species' mating systems that affect fishery sustainability and yield are rarely

considered. For species with obligate male care of eggs, it is possible that size-specific harvest of

males will decrease larval production. In order to examine how size-based management practices

interact with mating systems, we modeled fisheries of two species with obligate male care,

Symphodus melops (corkwing wrasse) and *Ophiodon elongatus* (lingcod) under two management

scenarios, a minimum size limit and a harvest slot limit. We simulated the population dynamics,

larval production, and yield to the fishery under a range of fishing mortalities. We also modeled

size-dependent male care to determine its interaction with management. In both species, the slot

limit decreased yield by less than 12% at low fishing mortalities; at higher mortalities,

individuals rarely survived to outgrow the slot and spawning potential decreased substantially

relative to unfished levels. The spawning potential decreased less when managed with a slot limit

if we included a positive feedback between male size, care, and hatching success, but the benefit

of implementing the slot depended both on the relative proportions of males and females selected

by the fishery and on our assumptions regarding male size and care. This work highlights that the

effects of size- and sex-selective fisheries management can be nuanced and produce counter-

intuitive results.

Keywords

Fisheries, slot harvest limit, minimum size limit, population model, male care, sexual selection,

wrasse, lingcod

Introduction

The sustainability of a fishery depends on the intensity of fishing pressure, the selectivity of the fishery for different sizes, ages, or sexes, and natural variation in population age and size structure arising from the species' biology (Lowerre-Barbieri et al., 2017). Sustainable fisheries management aims to maximize yield to the fishery, either in numbers or in biomass, over the long term, by controlling total fishing mortality, the selectivity of the fishery, or both. To achieve this aim, management must also account for annual changes in population abundance and age structure due to fishing, natural environmental variation, and biological responses of the fished species to these processes (Halliday & Pinhorn, 2009; Zhou et al., 2010). A large body of theoretical and empirical literature has shown that recruitment to the fishery (i.e., larval or juvenile survival) is related to spawning biomass and contemporary environmental conditions (Browman, 2014; Munch, Giron-Nava, & Sugihara, 2018; Pritt, Roseman, & O'Brien, 2014), but that over longer time scales, generation time, body size, and age at maturation are positively correlated with a stock's risk of overexploitation (Hutchings & Kuparinen, 2017; Hutchings & Reynolds, 2004; Kindsvater, Mangel, Reynolds, & Dulvy, 2016).

The potential effects of fishing on mating system dynamics and recruitment success have rarely been incorporated into management, despite evidence fishing can affect these processes (Alonzo & Mangel, 2004; Kendall & Quinn, 2013; Rowe & Hutchings, 2003; Sørdalen et al., 2018). The effects of fishing on mating systems may be most important when fishing targets a specific size in sexually dimorphic species (Carroll & Lowerre-Barbieri, 2019; H.K. Kindsvater, Reynolds, Sadovy de Mitcheson, & Mangel, 2017; Sørdalen et al., 2018). In such cases, differential mortality between the sexes and altered sex ratios can increase the likelihood of egg

or sperm limitation (depending on which sex is larger; Gosselin, Sainte-Marie, & Bernatchez,

2005; Hines et al., 2003). While the importance of large females to egg production and larval

survival is well acknowledged (Arlinghaus, Matsumura, & Dieckmann, 2010; Birkeland &

Dayton, 2005; Hixon, Johnson, & Sogard, 2014), the reproductive role of (large) males is often

ignored. However, males and sperm may also be limited (Alonzo *et al.* 2008; Sato 2012), body

size can have positive paternal effects (Uusi-Heikkilä, Kuparinen, Wolter, Meinelt, &

Arlinghaus, 2012), and theory predicts reduced reproductive rates when females are choosy but

males with preferred phenotypes are scarce (Møller & Legendre, 2001).

The interaction between fishing and obligate male care, in which territorial males must

defend eggs laid in nests to ensure eggs successfully hatch, has rarely been studied despite the

fact that several commercially and recreationally important species display this behavior (Table

1; Halvorsen et al. 2017; King and Withler 2005; Sutter et al., 2012). For these species, size-

selective fishing could decrease larval production by limiting the availability of nests or the

quality of male care, by selecting for smaller males. Smaller males may have smaller nests, fewer

nests or provide less effective care (Cargnelli & Neff, 2006; Ingebrigt Uglem & Rosenqvist,

2002; Wiegmann & Baylis, 1995). Moreover, the most effective guarding males could be the

largest, most aggressive males, so that the traits that correlate with the most effective parental

care could also increase vulnerability to fishing (Andersen, Marty, & Arlinghaus, 2018; Sutter et

al., 2012).

Despite growing recognition of the importance of maintaining diverse size structure in

fished populations, there is little theory predicting how size-selective fishing will interact with

reproductive behavior and influence population productivity. Conventional wisdom suggests

size-selective management (e.g., minimum size limits), designed to allow individuals to mature

and reproduce successfully before being fished, will ensure sustainable yields in the long term (

Birkeland, & Dayton, 2005; Froese, 2004;). More recently, protection of the largest, oldest

individuals has been recognized as a desirable management outcome if large individuals play

important roles ecologically or contribute high quality larvae to the population (Ahrens, Allen,

Walters, & Arlinghaus, 2020; Hixon, Johnson, & Sogard, 2014). A maximum size limit, in

addition to a minimum size limit, often referred to as a harvest slot, has been suggested to

prevent sex-selective harvest when one sex grows faster than the other (Halvorsen et al., 2016;

Morson, Munroe, Harner, & Marshall, 2017). Slot limits have been increasingly implemented in

recreational fisheries as a means of ensuring that fishing's effects on stock size, age, or sex

structure are balanced, or to maintain a suitable number of large fish available to anglers (Gwinn

et al., 2015). Slot limits have been advocated in fisheries for species in which mating success

and egg survival also depend on body size as in species with obligate paternal care (Halvorsen et

al., 2017). However, when considering the effectiveness of a minimum size limit vs. a slot limit,

it can be unclear how each management action will interact with the species' mating system and

sex-specific growth differences to affect fishery sustainability and yield. For example, in species

with paternal care, it is unclear whether it is more important for management actions to protect

females (typically the gamete-limiting sex) or males to ensure a sustainable larval supply.

To explore these alternatives, we use simulations of age-, size- and sex-structured

population dynamics to understand how fishing will interact with natural variation in male and

female size structure. We focus on comparing the effects of different size-selective management

scenarios in species with obligate male care, but different mating systems and different sex-

specific growth patterns. With a model of age-, size- and sex-structured population dynamics, we

compare the effects of a fishery selecting predominantly the large nesting males of a species with

multiple male phenotypes (*Symphodus melops,* the corkwing wrasse), with a fishery for a species

where females are the larger sex, and both males and females are caught (*Ophiodon elongatus*,

lingcod). Both of these species have obligate male care of eggs laid in benthic nests. We

specifically evaluate the consequences of alternative management tactics (a minimum size limit

vs. a slot limit) on the production of eggs and recruits by calculating the Spawning Potential

Ratio (SPR) and yield in numbers. By tracking male and female numbers and sizes in the model,

we are able to quantify the effects of management on both egg production and the availability of

paternal care. In comparing these species, we aim to gain general insights into the way that

species biology and fishery selectivity interact.

As fishing removes males of a specific size, the average size of a nesting male is

expected to decrease, because fewer males will survive to the largest age and size class. Whether

fishing affects larval production in species with male care depends on the consequences of size-

and sex-selective fishing for the egg density in each nest, and the extent to which egg

survivorship depends on density. In fishes, male care is often regarded as shareable, although

there is some evidence for density-dependent egg survival within a nest (Klug, Lindstrom, & St.

Mary, 2006). In addition, male size has shown to be positively correlated to the intensity of care

and nest survival (Suski & Ridgway, 2007; Wiegmann & Baylis, 1995). Here, we develop a new

metric of the availability of care *per egg* and quantify how that is expected to change with the

impacts of fishing on nest size and number, as well as on egg production. This is the first

theoretical investigation of the interplay between fishing and larval production in species with

alternative mating tactics and obligate male care.

Study species

We modelled two contrasting fisheries for species with obligate male care: the

commercial fishery for live *S. melops* in southern and western Norway and the recreational

fishery for *O. elongatus* in the northwestern USA. These are among the two best-studied marine

fisheries for species with parental care, and each fishery has a history of management that

includes minimum size limits and slot limits.

Corkwing wrasse

S. melops, the corkwing wrasse, has two male mating tactics, with large nesting males

providing care for eggs and sneaker males that steal fertilizations during spawning events (Potts,

1974; I. Uglem, Rosenqvist, & Wasslavik, 2000). Male and female corkwing wrasse have

different growth and maturation rates, as do the two male life-history pathways (territorial

nesting males and sneakers (Halvorsen et al., 2016; Uglem, Rosenqvist, & Wasslavik, 2000).

Sneaker males mature after one year. Nesting males and females mature between two and three

years, and can live for up to nine years.

The parameters for growth and maturation rates in our model were estimated from the

data published in Halvorsen et al. (2016), combined with more recent samples collected in 2017..

Growth and maturation rates as a function of age are plotted in Figure 1 (a,c); functions are in

Table 2; Parameter estimates are in Table 3. We assume mass-at-age $W(a)$ is a cubic function of

length, and we estimate egg production as a function of mass following the estimates in Chalaris

(2011). Like many batch spawners, corkwing wrasse are income breeders, but very little is

known about female re-mating rates (Stiver et al., 2018; Ingebrigt, Uglem & Rosenqvist, 2002)

and the number of times individual females spawn each season is unknown. While we do not

have direct information on natural mortality rates for each life-history type of corkwing wrasse,

estimates of total mortality (natural mortality and fishing mortality) can be estimated from the

frequency of each age class by life-history type in the catch data (Halvorsen *unpublished data*).

The commercial fishery for live corkwing wrasse has grown rapidly since 2010 as the wrasse are

used in louse biocontrol in salmon aquaculture ((Blanco Gonzalez et al., 2019; Halvorsen et al.,

2017). The fishery is size-selective, as the largest individuals fare better in salmon pens, and

since nesting males grow faster and mature older than females and sneaker males, their life

history is less protected by the current minimum size regulations. To account for this, a slot

harvest limit - where only fish between 120 and 170 mm can be retained – has been proposed to

protect the largest males in response to scientific concerns about the effects of fishing on the

mating system (Fig 1a; Halvorsen et al., 2017). In England, corkwing wrasse is managed by slot

sizes in two districts (IFCA 2020).

Lingcod

We compared the effect of fisheries management on the wrasse system with a species

with a contrasting life history (Figure 1, b,d), *O. elongatus,* lingcod. The lingcod is the only

extant member of the family Hexigrammidae and is native to western North America. Lingcod

are large, long-lived fish, although the ages of maturity of the nesting male wrasse and lingcod

are both between two and three years (Cass, Beamish, McFarlane, & Canada. Department of

Fisheries and Oceans., 1990; King & Withler, 2005). Lingcod females are capable of growing to

over two meters long and they live, on average, six years longer than males, with lower natural

mortality. As in the corkwing wrasse, males will defend eggs laid by one or more females in

benthic habitat for a period of 6 to 8 weeks, depending on water temperature.

182    Estimates of growth, maturation, and mortality rates, as well as estimates of mass-

183 specific egg production by females, and initial size at recruitment, were reported in Jagielo and

184 Wallace (2005), an international stock assessment that encompassed populations from California

185 to the Salish Sea. While there is undoubtedly some regional variation in growth and body size,

186 these growth and maturation values are representative of the species and are useful when

187 examining the effect of the contrasting life histories and mating systems of lingcod and corkwing

188 wrasse (Figure 1). There are two notable differences in lingcod growth patterns and mating

189 system that differ from the corkwing wrasse. First, female lingcod are the larger sex, and spawn

190 with only one male each season. Second, while smaller males are known to steal fertilizations by

191 sneaking during spawning events, it is likely that male mating tactics change ontogenetically as

192 males grow, and are not separate life history pathways (King and Withler 2005).

193    Lingcod have been fished commercially and recreationally for decades and were

194 considered to be overfished by the late 20th century due to intense fishing in the 1980s and

195 1990s (Haltuch et al., 2017; Jagielo & Wallace, 2005). Management has varied among Alaska,

196 British Columbia, Washington, Oregon, and California according to the history of the fishery and

197 natural latitudinal variation in growth and body size. Current management practices are a mix of

198 area closures, bag limits, minimum-size limits, and slot limits in different parts of the range. In

199 Washington State, recreational fishers are regulated by a slot limit of 26-36 inches (Fig 1b). Due

200 to the implementation of precautionary management measures in recent years, lingcod

201 populations are rebuilding and appear to be stable throughout much of the range (Haltuch et al.,

202 2017).

Methods

We developed a deterministic model of equilibrium population dynamics (Table 2),

which is closely related to the size-and age- structured model commonly used in fisheries stock

assessments (Mangel 2006). However, our model differs in several important ways. First, we

allowed for the sexes to have different growth rates. We also addressed the two growth patterns

of the different male life history strategies (nesting male and sneaker) for the corkwing wrasse

population. Second, we modeled differences in natural mortality and fishery susceptibility for

each life-history type. Third, we considered the relationship between the male size structure,

paternal care, and larval production. Figure 2 shows a general schematic of the model. With this

model, we can evaluate the potential consequences of fishing for population productivity, given

different assumptions regarding how male size structure affects the availability of paternal care

in nests and the resulting consequences for the production of larvae.

Male size and paternal care

We assume the effectiveness of paternal care in a nest – which is essential for the

production of viable larvae in both species – increases with the size of nesting males in both

corkwing wrasse and lingcod. In both species, larger males have larger nests and nest multiple

times in a season or defend multiple nests at once (King & Withler, 2005b; Potts, 1985; Ingebrigt

Uglem & Rosenqvist, 2002). It is also possible that large males are more effective at nest defense

than small nesting males, as is the case for smallmouth bass (Wiegmann & Baylis, 1995).

However, the exact relationship between egg number and larval numbers as a function of male

size and nest size is unknown for these and most other species. Per-capita survival of eggs in the

nest could be a decreasing function of egg density and nest size (i.e., larval production could be a

density-dependent function of egg number due to crowding). Both males and females may be

adapted to avoid spawning under conditions of nest crowding, but if the number of nests is

limited, they might choose to spawn anyway. As there are a range of possibilities, as a starting

point, we chose a simple relationship between male size and available nest care to represent both

male nest size and number of nests. We developed a Male Size Index, which we represent as θ,

representing the probability that a territorial male of a given length is mature, and the average

length of territorial males, weighted by the abundance of each age class (Table 2). We modeled

available nest care as a power function, such that care $C = \theta^p$. We varied p from one (linear) to

1.25 (slightly concave up). We used the nest care function to calculate the per-egg availability of

care in a nest as the ratio of available care C to total egg production each year. Both care

availability and egg production decreased with fishing, but the selectivity of fishing pressure for

each sex determined whether *care-per-egg* decreased or increased. We characterize this change

as the Care Ratio, which we represent as *R:* the care-per-egg in the fished population, relative to

the care-per-egg in the unfished population (Table 2).

Using this new metric *R,* we evaluated the consequences of a feedback from fishing on

larval production, the total number of eggs that hatch successfully year, due to a change in care-

per-egg. We assumed larval hatching success changed proportionally according to the Care Ratio

in this subset of our analyses. For example, if available care per egg decreased with fishing

mortality relative to the unfished population (i.e., *R < 1*), we assumed this also decreased larval

hatching success by the same fraction. In this way we estimated the indirect effect of changes in

per-egg availability of care after fishing removed some males and females. Our goal was to

compare the net effect of these changes to male and female demography on yield and larval

production, with and without the potential feedback between care availability and hatching

success. In every scenario, we calculated expected yield and larval production, sometimes

referred to as spawning potential. The latter is correlated with a population's capacity to buffer

stochastic environmental variation (Kindsvater et al., 2016; O'Farrell & Botsford, 2005).

Recruitment and population dynamics of both species

Births and deaths in our simulated population were determined by the demographic

composition of males and females. The maturation rate and mortality rate of each sex of each

species determines the number of mature individuals in each age class (a) alive at time $N(t)$, if

they are mature $p_m(a)$, and the relationship between fecundity and body mass $W(a)$. Egg

production $E(t)$ depends on the number and size of mature females, and the mass-fecundity

relationship for each species (Table 2; parameters are given in Table 3). In scenarios with no

indirect effect of care availability, egg production and larval production $P(t)$ were perfectly

correlated, $E(t) = P(t)$. As illustrated in Figure 2, with a feedback incorporating care

availability, larval production decreased according the care ratio, $E(t) = RP(t)$. For all

scenarios, we assumed the population dynamics are regulated by density-dependence in

recruitment to the year one age class. In other words, larval survival from hatching to recruitment

depends on larval density, but adult survival and growth are independent of density. We assumed

the recruitment function followed the Beverton-Holt equation (Mangel, 2006). The Beverton-

Holt relationship specifies the maximum probability that a larva survives to recruitment α (i.e.,

the productivity of the population at low density) and a metric of the strength of density

dependence β. These parameters arise from specific environmental conditions that are difficult to

measure, so they are often estimated from stock-recruitment relationships that relate recruitment

to spawning stock biomass (Mangel, 2006). In species lacking stock-recruitment relationships,

such as the corkwing wrasse, we can characterize the effects of fishing on a population relative

to an arbitrary unfished population. For comparison, we completed the same analysis on lingcod.

Importantly, as long as recruitment overfishing is not occurring, our assumptions regarding

density-dependent recruitment (i.e., the Beverton-Holt stock recruitment curve) do not have a

strong effect on relative yield and relative spawning potential (sometimes called the Spawning

Potential Ratio, SPR). To make sure this assumption is valid for the results presented here, we

used a range of values for α and β and checked to make sure they did not influence on our

conclusions.

After individuals recruit to the population model after 1 year at size L_R (in cm), they

grow and reproduce each year according to the specific growth, maturation, and mortality rates

reported for each species (Table 3, Figure 2). Individuals live to at most A_{max} years. Given the

balance of birth and death rates, the population will equilibrate at a steady state population

biomass. We checked that our simulated populations all reached this equilibrium within 100

years, at which time we "fished" our populations and made sure the fished population reached a

new steady state. The selectivity of fishing mortality varied according to the management

scenario we simulated. We evaluated a range of fishing mortality coefficients (F in Tables 2 and

3) for each scenario. For each species, we compared a minimum size limit with a slot size limit,

based on size limits currently used in management. We also compared scenarios with and

without a feedback between care and larval production. This factorial comparison allowed us to

tease apart the effects of management on the indirect effects of fishing on yield and larval

production. Doing this for both species revealed how the details of each life history interacted

with management decisions.

Results

For each factorial combination, we calculated yield to the fishery in numbers of fish,

rather than biomass. To evaluate spawning potential, we compared the lifetime egg production of

individuals in the fished population to those in the unfished, which is a method of calculating the

Spawning Potential Ratio (SPR; e.g. Kindsvater et al., 2016). This reference point is commonly

used to understand the recovery potential of a population, and conversely, its risk of

overexploitation (O'Farrell & Botsford, 2005). Finally, we calculated the Care Ratio R: the care

per egg available in the fished population, relative to the unfished population (Table 2).

Corkwing wrasse

Figure 3 shows that implementing the harvest slot limit led to a small decrease in yield to

the wrasse fishery. A decrease in yield after the implementation of the slot is expected, because

the largest individuals are protected from fishing. Natural mortality is high enough in corkwing

wrasse that very few individuals in our model survived to grow beyond the maximum size limit,

especially under strong fishing mortality. Therefore, the percent change in yield between the

minimum size limit and the slot was greatest at low fishing mortality (Fig. 3c). This implies that

for species with high natural mortality, the benefits of a slot limit will be countered by any

corresponding change in fishing mortality on size classes within the slot. In other words, if the

harvest rate of size classes within the slot increases as fishers respond to its implementation, it is

possible the benefits of the slot will be undetectable.

In our model, yield did not change in the scenarios including the feedback between male

size structure and larval production. While the slot limit increased care capacity by decreasing

the number of large males captured, the minimum size limit was sufficient to protect egg

production enough that recruitment to the adult population (at age one) was not affected. This is

a consequence of our use of a deterministic Beverton-Holt recruitment function. Because stock

recruitment relationships vary according to inter-annual environmental fluctuations, the result

that yield is stable under high size-selective fishing mortality is an oversimplification of reality

and could lead to overconfidence in a fishery's sustainability. Therefore, we focus on the

Spawning Potential Ratio (a proxy for larval production) as an indicator of the ability of the

population to buffer environmental stochasticity.

The dashed lines in each panel of Figure 4 show the SPR (squares) and the Care Ratio R

(circles), which reflects the care capacity of the population. We plot R even in the scenarios

where there is no indirect effect of care capacity on egg survival, to better illustrate the

interaction between spawning potential and R. As expected, spawning potential is lower at all

levels of fishing mortality when egg survival to the larval stage decreases with the availability of

care. In other words, fishing has a stronger negative effect on larval supply when we assume

there is an effect of care capacity on larval production. However, implementing a harvest slot

mitigates this effect (compare SPR in Fig. 4c and 4d). Additionally, the efficacy of the slot in

increasing larval supply (i.e., the difference in squared dashed-lines in 4d to 4b) depends on the

shape parameter p, which determines nonlinearity of the relationship between male size and care

capacity. At higher values of p ($1 < p < 2$), protecting the largest males with a slot limit has an

outsized effect on the population's care capacity and increases the SPR in Fig 4d, especially at

low levels of fishing mortality when the slot is most effective (not shown).

Lingcod

As in the wrasse, the yield to the fishery (in number of fish) decreased with the

implementation of a harvest slot limit (Fig. 5), but did not vary with an effect of care capacity on

larval production. Again, we attribute this stability in catch to our choice of the Beverton-Holt

stock recruitment function and the fact the minimum size limit is sufficient to prevent

recruitment overfishing in our deterministic model. As in the wrasse, yield differences arising

from the two management scenarios was greatest at low fishing mortality; at high rates of fishing

mortality yield did not differ substantially between the minimum size and slot limit scenarios

(Fig. 5), as few individuals survived to outgrow the maximum size limit.

Lingcod females are larger than males, so under a minimum size limit, the main effect of

fishing was to strongly reduce the spawning potential of females (Fig. 6a, dashed line with

squares). Including a feedback with care capacity mitigated this decrease in SPR (Fig. 6c, dashed

line with squares). This pattern held in both management scenarios, but the increase due to care

was more dramatic in the minimum size limit scenario (Fig. 6a and 6c). This is because in

lingcod, the availability of care increased *above* unfished levels after fishing, because egg

production decreased faster than nests. The increase in care-per-egg is reflected in a Care Ratio

greater than one in all fishing scenarios (Fig. 6). This effect was less dramatic with slot limit, as

these only protected the female population; males did not grow beyond the upper size limit (Fig.

1b). For this reason, the shape parameter p, representing the degree to which the largest males

contributed disproportionately to total care capacity, did not affect the results in Fig. 6.

Discussion

Our study is part of a growing body of literature evaluating the potential for size-selective

management strategies to balance yield, spawning potential, and long-term fishery sustainability

(Gwinn et al., 2015; Kindsvater et al., 2017, Ahrens et al., 2020). The influence of male size

structure on population productivity in fished species with obligate male care has been

recognized as an important aspect of effective management, but left un-explored (Halvorsen et

al., 2016; Stubberud et al., 2019). Here, we modelled fisheries for two such species to investigate

the effects of different size-selective management strategies on yield, spawning potential, and

care. By using a simulation approach, we were able to investigate the possible importance of an

indirect effect (feedback) of fishing on per-egg care, which depends on both male and female

size structure. We found that for both species, without any assumptions about how male care

influences larval production, the slot limit was most effective at protecting larval production

when fishing mortality was low. Under high fishing mortality, in the steady state, very few

individuals survived beyond the maximum size limit. This result might change if fishing

decreases natural mortality, which is possible if the mechanisms of adult mortality are density

dependent. This is a plausible mechanism for compensatory productivity, especially for species

with high natural mortality (Andersen, Jacobsen, Jansen, & Beyer, 2017; Rose, Cowan,

Winemiller, Myers, & Hilborn, 2001). Furthermore, our results indicate that slot limits alone

could be insufficient to prevent overexploitation. In both species, implementing a slot limit led to

a difference in yield of 5% or less once fishing mortality (F) exceeded 0.4 (Fig. 3c, Fig. 5c),

while spawning potential was reduced to 60% or less of unfished levels in the wrasse when we

assumed a decrease in care availability due to fishing. Our results suggest the growth patterns

and mating systems of both species may increase their vulnerability to overfishing, for different

reasons, and despite management measures.

Corkwing wrasse

The benefit of the slot limit to larval production - relative to populations managed with a

minimum size limit - could only be detected in the corkwing wrasse simulations when care was

reduced by fishing in a way that reduced the proportion of surviving larvae by the fraction R.

This is because, under fishing, no females grew large enough to benefit from the maximum size

limit, so the slot did not affect egg production. The details regarding how care affects larval

survival are important. First, we assumed per-capita egg survival decreased linearly with nest

availability, i.e. there were no non-linear effects (such as increased survival at low density). This

assumption may not hold for all species with obligate male care, but in *S. ocellatus,* a congeneric

wrasse species, male care is largely sharable (i.e., we don't expect a decrease in egg survival at

high densities) and at low egg densities, males are more likely to abandon nests (Alonzo 2004).

Second, the more nonlinear the relationship between the Male Size Index and care capacity, the

more effective the slot limit was at increasing larval production. We do not know how much

more effective larger males are in providing care, but in the corkwing wrasse, large males have

larger nests and start nesting earlier, which could allow them to care for more eggs and complete

more nesting cycles in a spawning season, suggesting that they do provide an outsized

contribution to care (Ingebrigt, Uglem & Rosenqvist, 2002). Finally, in corkwing wrasse, size-

selective fishing has two impacts: disproportionate removal of nesting males, and decreasing the

amount of care available for eggs. We did not model the effect of male removal on fertilization

success, because this species also has sneaker males capable of fertilizing eggs, so when some

nesting males are present, sperm limitation is unlikely. However, in the long term, any increase

in the mating success of these smaller males may be offset by changes in the inter- and intra-

sexual dynamics of males and females, as the fitness of sneaker males also depends on the

frequency of nesting males and the choosiness of females.

Lingcod

By contrast, for lingcod, the slot limit was effective in increasing larval production

(relative to that expected under a minimum size limit), even without a feedback between nest

availability and care capacity. However, the fishery selectively removed a greater proportion of

the female spawning population than the male population. As a result, this species remained

vulnerable to recruitment overfishing, even with a slot limit in place, if total fishing mortality

was not controlled. This result is likely to be the case for any species where females are much

larger than males and fishing is strongly size-selective (regardless of whether they have male

care). For this reason, our assumptions about the potential indirect effects of fishing on male

care were less important than the direct effect of fishing on egg production. For lingcod,

spawning potential was reduced to 40% or less of unfished levels once fishing mortality

exceeded 0.4, unless we assumed there was a "rescue" effect due to the increase in availability of

per-egg care (Fig. 6). Currently we do not know whether increased care availability has any such

compensatory effect, or if it does, whether the magnitude we considered is realistic. However,

our result suggests the use of bag limits, in addition to slot limits, have been important in

allowing lingcod stocks to rebuild.

In our model, recruitment depended on a deterministic Beverton-Holt relationship

between larval production and survival. Therefore, our model results regarding yield were not

sensitive to our assumptions about feedbacks from male care (Fig. 3; Fig. 5), despite differences

in larval production (SPR) arising under different management and harvest scenarios (Fig 4, Fig.

6). In reality, we expect that the relationship between larval supply and yield will fluctuate

according to environmental conditions, though detecting this relationship is notoriously tricky

(Munch et al., 2018). By using the Beverton-Holt relationship, we assumed there was no

compensatory density-dependence (as there would be if we used a Ricker function). We chose

the Beverton-Holt because it is a more conservative model of recruitment dynamics (i.e., fishing

does not increase population productivity as it would under a Ricker model). We also have no

evidence for cannibalism at high population densities, or for interference in egg survival at high

adult population densities, which have been suggested to drive Ricker-like dynamics in some

species. If we had used a Ricker function, the relative increase in larval production expected after

implementing a slot size limit would be offset by the decline in recruitment success at high larval

densities.

When males of different sizes compete for fertilizations, as in species large fighter males

and smaller sneaker males, it is unclear how adaptation to size-selective fishing will affect the

distribution of male sizes and population productivity (DeFilippo et al., 2019; Kendall et al.,

2014; Kendall & Quinn, 2013). Some theory predicts that sexual selection will reinforce

adaptation to fishing when males are under directional sexual selection, but fishing increases

variance in male reproductive success (Hutchings & Rowe, 2008). There is some evidence from

sockeye salmon, however, that the removal of large males by fishing elevates the relative fitness

of secondary males (in salmon, known as "jacks"), and increases the frequencies of smaller

males in subsequent generations (DeFilippo et al., 2019). In other species with multiple male

mating tactics, this may not be the case. For example, in a wrasse with paternal care, high

sneaking rates have been found to decrease the willingness of nesting males and females to mate

(Alonzo & Heckman, 2010; Alonzo & Warner, 1999). In this case, it is possible that sexual

selection will actually decrease the success of the remaining (unfished) nesting males in the short

term, because the density of small sneaker males at remaining nests will increase. The frequency-

dependent selection that is maintaining the stable polymorphism in male mating tactic may be

disrupted, and the potential interactions with fishing-induced selection on maturation and growth

is unknown. One possibility is that by increasing sneaker density at nests, fishing could indirectly

hamper population productivity by decreasing the number of eggs that are spawned, even if

females are not fished directly – an undesirable outcome of size-selective management.

In species that do not have secondary males, such as lingcod, it is possible that

fertilization rates could be a limiting factor in fisheries that target large males (Alonzo and

Mangel 2004). Several studies support the assumption of a positive correlation between male

body size and reproductive success in fishes with male care (Bose et al., 2018; Cargnelli & Neff,

2006; Wiegmann & Baylis, 1995). Among fisheries targeting fishes with male care, some are

managed by harvest slots, reflecting that the importance of large males (and females) is

acknowledged (Table 1). Several of these species are also protogynous hermaphrodites. For these

species, management that protects the largest individuals may be necessary to prevent depletion

of terminal males to avoid sperm limitation, and can also help to buffer fisheries-induced

reduction in size- at sex-change (Alonzo et al., 2008; Kindsvater, Reynolds, Sadovy de

Mitcheson, & Mangel, 2017; Sato et al., 2018).

Although harvest slots may seem to be an intuitive measure to balance fishing mortality

between sexes, we show that management protecting large individuals of any sex is likely to

have small effects if natural mortality is relatively high. That does not mean that large males

should be ignored, but rather that management strategies are carefully evaluated and effects

monitored. Our results are consistent with three rules of thumb to promote the sustainability of

fisheries for species with paternal care: (i) control fishing mortality (e.g. implement quotas or

bag limits); (ii) allow both males and females to spawn at least once before fished (e.g. enforce

minimum size limits or gear modifications) and (iii) reduce or restrict fishing during the nesting

period which can directly affect productivity if guarding males are fished (Froese, 2004;

Overzee & Rijinsdorp, 2015; Suski 2003.). A spawning season closure has been implemented in

the Norwegian wrasse fishery in recent years, but wrasse can be fished during the spawning

season in Sweden (Faust, Halvorsen, Andersen, Knutsen, & André, 2018). Lastly, in order to

better predict the consequences of harvesting fish with male care, dedicated field studies are

needed under realistic conditions. For example, how does disproportional removal of large males

affect competition and social hierarchies among remaining males, and to what extent does

reduced availability of nests affect female mating decisions and offspring survival?

In summary, our results highlight the need to exert caution when managing fishes with

obligate male care, and that slot size limits are no silver bullet for ensuring long term viability if

overall fishing mortality is high and not controlled. Furthermore, the evolutionary outcomes of

size-selective fishing for the relative fitness of the alternative reproductive tactics deserves

further scientific investigation. Despite these caveats, our results suggest that the current

management strategies used for both corkwing wrasse in Norway and lingcod in western North

America, have likely benefited the sustainability and rebuilding rates of the respective fisheries.

Acknowledgements

This work was supported by NSF DEB-1556779 to HKK, funding from Virginia Tech to

HKK, NSF IOS-1655297 to SHA and funding from the University of California Santa Cruz to

SHA.

References

Ahrens, R., Allen MS, Walters C, & Arlinghaus R. (2020). Saving large fish through harvest slots outperforms the classical minimum-length limit when the aim is to achieve multiple harvest and catch-related fisheries objectives. *Fish and Fisheries*. In press. https://doi.org/10.1111/faf.12442

Alonzo, S. H., & Heckman, K. L. (2010). The unexpected but understandable dynamics of mating, paternity and paternal care in the ocellated wrasse. *Proceedings of the Royal Society B: Biological Sciences*, 277(1678), 115–122.

Alonzo, S. H., Ish, T., Key, M., MacCall, A. D., & Mangel, M. (2008). The importance of incorporating protogynous sex change into stock assessments. *Bulletin of Marine Science*, 83(1), 163–179.

Alonzo, S. H., & Mangel, M. (2004). The effects of size-selective fisheries on the stock dynamics of and sperm limitation in sex-changing fish. *Fishery Bulletin*. 102:1-13

Alonzo, S. H., & Warner, R. R. (1999). A trade-off generated by sexual conflict: Mediterranean wrasse males refuse present mates to increase future success. *Behavioral Ecology*, 10(1), 105–111.

Andersen, K. H., Jacobsen, N. S., Jansen, T., & Beyer, J. E. (2017). When in life does density dependence occur in fish populations? *Fish and Fisheries*, 18(4), 656–667. https://doi.org/10.1111/faf.12195

Andersen, K. H., Marty, L., & Arlinghaus, R. (2018). Evolution of boldness and life history in response to selective harvesting. *Canadian Journal of Fisheries and Aquatic Sciences*, 75(2), 271–281. https://doi.org/10.1139/cjfas-2016-0350

Arlinghaus, R., Matsumura, S., & Dieckmann, U. (2010). The conservation and fishery benefits of protecting large pike (*Esox lucius* L.) by harvest regulations in recreational fishing. *Biological Conservation*, 143(6), 1444–1459. https://doi.org/10.1016/j.biocon.2010.03.020

Birkeland, C., & Dayton, P. K. (2005). The importance in fishery management of leaving the big ones. *Trends in Ecology & Evolution*, 20(7), 356–358. https://doi.org/10.1016/j.tree.2005.03.015

Blanco Gonzalez, E., Espeland, S. H., Jentoft, S., Hansen, M. M., Robalo, J. I., Stenseth, N. C., & Jorde, P. E. (2019). Interbreeding between local and translocated populations of a cleaner fish in an experimental mesocosm predicts risk of disrupted local adaptation. *Ecology and Evolution*, 9(11), 6665–6677. https://doi.org/10.1002/ece3.5246

549 Bose, A. P. H., Cogliati, K. M., Luymes, N., Bass, A. H., Marchaterre, M. A., Sisneros, J. A., …
550 Balshine, S. (2018). Phenotypic traits and resource quality as factors affecting male reproductive
551 success in a toadfish. *Behavioral Ecology*,29(2) 496–507. https://doi.org/10.1093/beheco/ary002
552

553 Browman, H. I. (2014). Commemorating 100 years since Hjort's 1914 treatise on fluctuations in
554 the great fisheries of northern Europe: Where we have been, where we are, and where we are
555 going. *ICES Journal of Marine Science*, Vol. 71, pp. 1989–1992.
556 https://doi.org/10.1093/icesjms/fsu159
557

558 Cargnelli, L. M., & Neff, B. D. (2006). Condition-dependent nesting in bluegill sunfish *Lepomis*
559 *macrochirus*. *Journal of Animal Ecology*, 75(3), 627–633. https://doi.org/10.1111/j.1365-
560 2656.2006.01083.x
561

562 Carroll, J., & Lowerre-Barbieri, S. (2019). Interactions of dimorphic growth, reproductive
563 behavior, and a size-regulated fishery: a case study using spotted seatrout *Cynoscion nebulosus.*
564 *Marine Ecology Progress Series*, 608, 233–245. https://doi.org/10.3354/meps12795
565

566 Cass, A. J., Beamish, R. J., McFarlane, G. A. (1990). Lingcod (*Ophiodon elongatus*). *Can. Spec.*
567 *Pub. Fish. Aquat. Sci*. 109:40.
568

569 Chalaris, A. (2011). Preliminary study of fecundity, sperm characteristics, behaviour and
570 hormonal induction in native wrasse species. Master's of Science thesis, University of Stirling.
571

572 DeFilippo, L. B., Schindler, D. E., Ohlberger, J., Schaberg, K. L., Foster, M. B., Ruhl, D., &
573 Punt, A. E. (2019). Recruitment variation disrupts the stability of alternative life histories in an
574 exploited salmon population. *Evolutionary Applications*, 12(2), 214–229.
575 https://doi.org/10.1111/eva.12709
576

577 Enberg, K., Jørgensen, C., Dunlop, E. S., Heino, M., & Dieckmann, U. (2009). Implications of
578 fisheries-induced evolution for stock rebuilding and recovery. *Evolutionary Applications*, 2(3),
579 394–414. https://doi.org/10.1111/j.1752-4571.2009.00077.x
580

581 Faust, E., Halvorsen, K. T., Andersen, P., Knutsen, H., & André, C. (2018). Cleaner fish escape
582 salmon farms and hybridize with local wrasse populations. *Royal Society Open Science*.
583 https://doi.org/10.1098/rsos.171752
584

585 Froese, R. (2004). Keep it simple: Three indicators to deal with overfishing. *Fish and Fisheries*,
586 5(1), 86–91. https://doi.org/10.1111/j.1467-2979.2004.00144.x
587

588 Gosselin, T., Sainte-Marie, B., & Bernatchez, L. (2005). Geographic variation of multiple
589 paternity in the American lobster, *Homarus americanus. Molecular Ecology*.
590 https://doi.org/10.1111/j.1365-294X.2005.02498.x
591

592 Gwinn, D. C., Allen, M. S., Johnston, F. D., Brown, P., Todd, C. R., & Arlinghaus, R. (2015).
593 Rethinking length-based fisheries regulations: The value of protecting old and large fish with
594 harvest slots. *Fish and Fisheries*, 16(2), 259–281. https://doi.org/10.1111/faf.12053
595
596 Halliday, R. G., & Pinhorn, A. T. (2009). The roles of fishing and environmental change in the
597 decline of Northwest Atlantic groundfish populations in the early 1990s. *Fisheries Research,*
598 97(3), 163–182. https://doi.org/10.1016/j.fishres.2009.02.004
599
600 Haltuch, M. A., Wallace, J., Akselrud, C. A., Nowlis, J., Barnett, L. A. K., Valero, J. L., … Lam,
601 L. .2017 Lingcod Stock Assessment. Pacific Fisheries Management Council, Portland, OR.
602 Available from https://www.pcouncil.org/stock-assessments-and-fishery-evaluation-safe-
603 documents/
604
605 Halvorsen, K. T., Larsen, T., Sørdalen, T. K., Vøllestad, L. A., Knutsen, H., & Olsen, E. M.
606 (2017). Impact of harvesting cleaner fish for salmonid aquaculture assessed from replicated
607 coastal marine protected areas. *Marine Biology Research*, 13(4), 359–369.
608 https://doi.org/10.1080/17451000.2016.1262042
609
610 Halvorsen, K. T., Sørdalen, T. K., Durif, C., Knutsen, H., Olsen, E. M., Skiftesvik, A. B., …
611 Vøllestad, L. A. (2016). Male-biased sexual size dimorphism in the nest building corkwing
612 wrasse (Symphodus melops): implications for a size regulated fishery . *ICES Journal of Marine*
613 *Science*, 73(10), 2586–2594. https://doi.org/10.1093/icesjms/fsw135
614
615 Halvorsen, K. T., Sørdalen, T. K., Vøllestad, L. A., Skiftesvik, A. B., Espeland, S. H., & Olsen,
616 E. M. (2017). Sex-and size-selective harvesting of corkwing wrasse (Symphodus melops)-a
617 cleaner fish used in salmonid aquaculture. *ICES Journal of Marine Science*, 74(3), 660–669.
618 https://doi.org/10.1093/icesjms/fsw221
619
620 Hines, A. H., Jivoff, P. R., Bushmann, P. J., Van Montfrans, J., Reed, S. A., Wolcott, D. L., &
621 Wolcott, T. G. (2003). Evidence for sperm limitation in the blue crab, *Callinectes sapidus*.
622 *Bulletin of Marine Science.* 72(2): 287–310.
623
624 Hixon, M. A., Johnson, D. W., & Sogard, S. M. (2014). BOFFFFs: On the importance of
625 conserving old-growth age structure in fishery populations. *ICES Journal of Marine Science,*
626 71(8), 2171–2185. https://doi.org/10.1093/icesjms/fst200
627
628 Hutchings, J. A., & Kuparinen, A. (2017). Empirical links between natural mortality and
629 recovery in marine fishes. *Proceedings of the Royal Society, Series B*, 284(1856), 20170693.
630 https://doi.org/10.1098/rspb.2017.0693
631
632 Hutchings, J. A., & Reynolds, J. D. (2004). Marine Fish Population Collapses: Consequences for
633 Recovery and Extinction Risk. *BioScience* 54, 297–309
634   https://academic.oup.com/bioscience/article-abstract/54/4/297/284117
635

Hutchings, J. A., & Rowe, S. (2008). Consequences of sexual selection for fisheries-induced evolution: an exploratory analysis. *Evolutionary Applications*, 1(1), 129–136. https://doi.org/10.1111/j.1752-4571.2007.00009.x

IFCA (2020). Inshore Fisheries and Conservation Authority. Wrasse Fishery Guidance. Available at https://secure.toolkitfiles.co.uk/clients/25364/sitedata/files/Wrasse-Guidance.pdf

Jagielo, T. H., & Wallace, F. R. (2005). Assessment of Lingcod (*Ophiodon elongatus*) for the Pacific Fishery Management Council. Montesano, WA.

Kendall, N. W., Dieckmann, U., Heino, M., Punt, A. E., & Quinn, T. P. (2014). Evolution of age and length at maturation of Alaskan salmon under size-selective harvest. *Evolutionary Applications*, 7(2), 313–322. https://doi.org/10.1111/eva.12123

Kendall, N. W., & Quinn, T. P. (2013). Size-selective fishing affects sex ratios and the opportunity for sexual selection in Alaskan sockeye salmon *Oncorhynchus nerka. Oikos*, 122(3), 411–420. https://doi.org/10.1111/j.1600-0706.2012.20319.x

Kindsvater, H. K., Mangel, M., Reynolds, J. D., & Dulvy, N. K. (2016). Ten principles from evolutionary ecology essential for effective marine conservation. *Ecology and Evolution*, 6(7). https://doi.org/10.1002/ece3.2012

Kindsvater, H. K., Reynolds, J. D., Sadovy de Mitcheson, Y., & Mangel, M. (2017). Selectivity matters: Rules of thumb for management of plate-sized, sex-changing fish in the live reef food fish trade. *Fish and Fisheries*, 18(5). https://doi.org/10.1111/faf.12208

King, J. R., & Withler, R. E. (2005). Male nest site fidelity and female serial polyandry in lingcod (Ophiodon elongatus, Hexagrammidae). *Molecular Ecology*, 14(2), 653–660. https://doi.org/10.1111/j.1365-294X.2005.02438.x

Klug, H., Lindstrom, K., & St. Mary, C. M. (2006). Parents benefit from eating offspring: density-dependent egg survivorship compensates for filial cannibalism. *Evolution* 60(10), 2087-2095.
.
Lowerre-Barbieri, S., DeCelles, G., Pepin, P., Catalán, I. A., Muhling, B., Erisman, B., … Paris, C. B. (2017). Reproductive resilience: a paradigm shift in understanding spawner-recruit systems in exploited marine fish. *Fish and Fisher*ies, 18(2), 285–312. https://doi.org/10.1111/faf.12180

Mangel, Marc. (2006). The Theoretical Biologist's Toolbox : Quantitative Methods for Ecology and Evolutionary Biology. Cambridge University Press.

Morson, J. M., Munroe, D., Harner, R., & Marshall, R. (2017). Evaluating the potential for a sex-balanced harvest approach in the recreational summer flounder fishery. *North American Journal of Fisheries Management,* 37(6), 1231–1242. https://doi.org/10.1080/02755947.2017.1362490

682 Munch, S. B., Giron-Nava, A., & Sugihara, G. (2018). Nonlinear dynamics and noise in fisheries
683 recruitment: A global meta-analysis. *Fish and Fisheries*, 19(6), 964–973.
684 https://doi.org/10.1111/faf.12304
685
686 O'Farrell, M. R., & Botsford, L. W. (2005). Estimation of change in lifetime egg production
687 from length frequency data. *Canadian Journal of Fisheries and Aquatic Sciences*, 62(7), 1626–
688 1639. https://doi.org/10.1139/f05-064
689
690 Overzee, H. Van, & Rijinsdorp, A. (2015). Effects of fishing during the spawning period:
691 implications for sustainable management. *Reviews in Fish Biology and Fisheries*, 25(1), 65–83.
692 https://doi.org/10.1007/s11160-014-9370-x
693
694 Potts, G. W. (1974). The colouration and its behavioural significance in the corkwing wrasse,
695 Crenilabrus Melops. *Journal of the Marine Biological Association of the United Kingdom*, 94,
696 925–938. https://doi.org/10.1017/S0025315400057659
697
698 Potts, G. W. (1985). The nest structure of the corkwing wrasse, Crenilabrus Melops (Labridae:
699 Teleostei). *Journal of the Marine Biological Association of the United Kingdom*, 65(02), 531–
700 546. https://doi.org/10.1017/S002531540005058X
701
702 Pritt, J. J., Roseman, E. F., & O'Brien, T. P. (2014). Mechanisms driving recruitment variability
703 in fish: Comparisons between the Laurentian Great Lakes and marine systems. *ICES Journal of
704 Marine Science*, 71(8), 2252–2267. https://doi.org/10.1093/icesjms/fsu080
705
706 Rose, K. A., Cowan, J. H., Winemiller, K. O., Myers, R. A., & Hilborn, R. (2001).
707 Compensatory density dependence in fish populations: Importance, controversy, understanding
708 and prognosis. *Fish and Fisheries*, 2(4), 293–327. https://doi.org/10.1046/j.1467-
709 2960.2001.00056.x
710
711 Rowe, S., & Hutchings, J. A. (2003). Mating systems and the conservation of commercially
712 exploited marine fish. *Trends in Ecology & Evolution*, 18(11), 567–572.
713 https://doi.org/10.1016/J.TREE.2003.09.004
714
715 Sato, T. (2012). Impacts of large male-selective harvesting on reproduction: illustration with
716 large decapod crustacean resources. *Aqua-BioScience Monographs*, 5(3), 67–102.
717 https://doi.org/10.5047/absm.2012.00503.0067
718
719 Sato, T., Kobayashi, M., Kurihara, T., Takebe, T., Hirai, N., Okuzawa, K., … Teruya, K. (2018).
720 Impact of fishing losses of males on the reproductive output of the large protogynous fish,
721 Choerodon schoenleinii. *Marine Ecology Progress Series*, 602, 225–235.
722 https://doi.org/10.3354/meps12675
723
724 Sørdalen, T. K., Halvorsen, K. T., Harrison, H. B., Ellis, C. D., Vøllestad, L. A., Knutsen, H., …
725 Olsen, E. M. (2018). Harvesting changes mating behaviour in European lobster. *Evolutionary
726 Applications*, 11(6), 963–977. https://doi.org/10.1111/eva.12611
727

728 Stiver, K. A., Kindsvater, H. K., Tamburello, N., Heckman, K. L., Klein, J., & Alonzo, S. H.
729 (2018). Intentional multiple mating by females in a species where sneak fertilization circumvents
730 female choice for parental males. *Journal of Fish Biology*, 93(2), 324–333.
731 https://doi.org/10.1111/jfb.13766
732
733 Stubberud, M. W., Vindenes, Y., Vøllestad, L. A., Winfield, I. J., Stenseth, N. C., & Langangen,
734 Ø. (2019). Effects of size- and sex-selective harvesting: An integral projection model approach.
735 *Ecology and Evolution*, 9(22), 12556–12570. https://doi.org/10.1002/ece3.5719
736
737 Suski, C.D. (2003). The effect of catch-and-release angling on the parental care behavior of male
738 smallmouth bass. *Transactions of the American Fisheries Society*, 132, 210–218.
739
740 Suski, C. D., & Ridgway, M. S. (2007). Climate and body size influence nest survival in a fish
741 with parental care. *Journal of Animal Ecology*, 76(4), 730–739. https://doi.org/10.1111/j.1365-
742 2656.2007.01242.x
743
744 Sutter, D. A. H., Suski, C. D., Philipp, D. P., Klefoth, T., Wahl, D. H., Kersten, P., …
745 Arlinghaus, R. (2012). Recreational fishing selectively captures individuals with the highest
746 fitness potential. *Proceedings of the National Academy of Sciences of the United States of
747 America*, 109(51), 20960–20965. https://doi.org/10.1073/pnas.1212536109
748
749 Uglem, I., Rosenqvist, G., & Wasslavik, H. S. (2000). Phenotypic variation between dimorphic
750 males in corkwing wrasse. *Journal of Fish Biology*, 57(1), 1–14. https://doi.org/10.1111/j.1095-
751 8649.2000.tb00771.x
752
753 Uglem, Ingebrigt, & Rosenqvist, G. (2002). Nest building and mating in relation to male size in
754 corkwing wrasse, *Symphodus melops*. *Environmental Biology of Fishes* 63, 17-25.
755
756 Uusi-Heikkilä, S., Kuparinen, A., Wolter, C., Meinelt, T., & Arlinghaus, R. (2012). Paternal
757 body size affects reproductive success in laboratory-held zebrafish (*Danio rerio*). *Environmental
758 Biology of Fishes,* 93(4), 461–474. https://doi.org/10.1007/s10641-011-9937-5
759
760 Wiegmann, D. D., & Baylis, J. R. (1995). Male body size and paternal behaviour in smallmouth
761 bass, Micropterus dolomieui (Pisces: Centrarchidae). *Animal Behaviour*, 50(6), 1543–1555.
762 https://doi.org/10.1016/0003-3472(95)80010-7
763
764 Zhou, S., Smith, A. D. M., Punt, A. E., Richardson, A. J., Gibbs, M., Fulton, E. A., … Sainsbury,
765 K. (2010). Ecosystem-based fisheries management requires a change to the selective fishing
766 philosophy. *Proceedings of the National Academy of Sciences of the United States of America,*
767 Vol. 107, pp. 9485–9489. https://doi.org/10.1073/pnas.0912771107

Figure captions

Figure 1. Length-at-age functions and age-specific maturation rates for Corkwing wrasse (left column) and lingcod (right column). For the wrasse, estimates of growth (panel a) and maturation rates (panel b) are available for each life-history pathway in the western Norway stock (females: black line; nesting males: blue line; sneaker males: red line). For lingcod, sex-specific estimates (females: black; males: blue) are available from stocks in Washington State. Note female lingcod are the larger sex. Parameters for each function are given in Table 3.

Figure 2. Generic schematic of the population model and where the reproductive behavior operates. The model assumes that recruitment to the adult population happens at after one year of age for both species. After that, they experience constant annual mortality, which varies according to sex and male tactic. Maturation rates for each sex and life history vary according to age and/or size, depending on how they were estimated empirically (Table 3). Female fecundity depends on her mass, which is a function of her age in this model. In the scenarios where we included an effect (feedback) of male care, the probability an egg hatched and becomes a planktonic larva is proportional to the Care Ratio R, based on the average size of territorial mature males, weighted by their frequency in a stable age distribution under fishing, relative to their frequency in the unfished population (Table 2).

Figure 3. Spawning potential and care capacity for each of the factorial combinations of corkwing wrasse simulations. We modeled fisheries with a minimum size limit (left column) or a slot limit (right column), each with (bottom row) and without (top row) a feedback between care

and larval production, across a range of fishing mortalities. Blue square-dashed lines represent

the SPR - relative larval production - at each level of fishing mortality. In each plot, red circle-

dashed lines represent the Care Ratio, R. For corkwing wrasse, R was bounded between 0 and 1,

so the right y-axis scale represents both dashed lines.

Figure 4. Yield of corkwing wrasse, in numbers of fish, for the minimum size limit (panel a) and

the slot limit (panel b). We calculate the percent difference in yield under the two management

scenarios (panel c). Yield is always greater under the minimum size limit.

Figure 5. Spawning potential and care capacity for each of the factorial combinations of lingcod

simulations. Again, we modeled fisheries with a minimum size limit (left column) or a slot limit

(right column), each with (bottom row) and without (top row) a feedback between care and larval

production, across a range of fishing mortalities. Blue square-dashed lines represent the SPR -

relative larval production - at each level of fishing mortality. In each plot, red circle-dashed lines

represent the Care Ratio R. For lingcod, R increased above 1 (with no fishing), ranging from 3-

20 under the highest level of fishing mortality F. The right y-axis scale (red) indicates this

change, which was greatest under a minimum size limit (panel a), as egg production was

severely depleted in this scenario.

Figure 6. Yield of lingcod, in numbers of fish, for the minimum size limit (panel a) and the slot

limit (panel b). We calculate the percent difference in yield under the two management scenarios

(panel c). Yield is always greater under the minimum size limit, but that difference narrows as

the population becomes overfished.

Table 1. Fished species with male care.

Species	Larger sex	Alternative male tactic	Hermaphroditism	Habitat	Fishery	Management includes slot limit (some populations)
Ballan wrasse (*Labrus bergylta*)	Male	No	Protogynous	Marine	Commercial and recreational	Yes
Cuckoo wrasse (*Labrus mixtus*)	Male	Yes	Protogynous	Marine	Commercial	Yes
Corkwing wrasse (*Symphodus melops*)	Male	Yes	No	Marine	Commercial	Yes
Black bream (*Spondyliosoma cantharus*)	Male	No	Protogynous	Marine	Recreational	No
Bluegill sunfish (*Lepomis macrochirus*)	Male	Yes	No	Fresh water	Recreational	No
Largemouth bass (*Micropterus salmoides*)	Monomorphic	No	No	Fresh water	Recreational	Yes
Smallmouth bass (*Micropterus dolomieu*)	Monomorphic	No	No	Fresh water	Recreational	Yes
Atlantic wolffish (*Anarhichas lupus*)	Monomorphic	No	No	Marine	Commercial	No
Butterfly peacock bass (*Cichla ocellaris*)*	Monomorphic	No	No	Fresh water	Recreational	No
Lingcod (*Ophiodon elongatus*)	Female	No	No	Marine	Commercial and recreational	Yes
Lumpfish (*Cyclopterus lumpus*)	Female	No	No	Marine	Commercial	No

*Note Butterfly peacock bass has biparental care.

Table 2.

Process	Equation	Interpretation
Von Bertalanffy growth function	$L(i, a+1) = L(i,a)e^{-k} + L_{\infty,i}(1 - e^{-k_i})$.	$L_{\infty,i}$ and k_i determine size-at-age L of each life history type (in discrete time t)
Maturation probability function (wrasse nesting males)	$p_m(i = nm, a) = \dfrac{1}{1 + e^{(-\gamma + \rho L(i=nm,a))}}$	For $i = nm$, wrasse nesting males, maturation is based on length L and logistic parameters γ and ρ. All females and sneaker males mature in years 2 and 1, respectively, which we denote as a_{100}
Maturation probability function (lingcod)	$p_m(i, a) = \dfrac{1}{1 + e^{-q(a - a_{mat,i})}}$	For lingcod $a_{mat,i}$ is the age at which 50% of individuals mature; q determines the steepness ogive.
Body mass	$W(i,a) = \nu_i L(i,a)^{\omega_i}$	ν and ω are scale and shape parameters for each life history type i
Natural mortality	$M(i,a)$	M assumed constant for all ages after age 1
Fishing mortality	$F(a) = \begin{cases} 0 & L(a) < \text{ min size limit} \\ F & \text{min size limit } < L(a) < \text{max size limit} \end{cases}$	Depends on fishing mortality rate F and selectivity (size limit or harvest slot limit)
Total annual egg production	$E(t) = \sum_a N(i=f,a,t)p_m(i=f,a)bW(i=f,a)^c$	Where $i = f$ is the female population. Mass and fecundity depend on length, which depends on age. Mass-specific fecundity parameters are b and c
Density dependent recruitment	$N_0(t+1) = \dfrac{\alpha P(t)}{1 + \beta P(t)}$	Beverton-Holt recruitment function
Spawning Potential Ratio	$SPR = \dfrac{\sum_a^{a_{max}} P(a)e^{-M(i=f,a)-F(a)}}{\sum_a^{a_{max}} N(i=f,a,t)p_m(a)bW(i=f,a)\,e^{-M(i=f,a)}}$	Egg production in the fished population relative to the unfished population. Note that the maximum ages in each population may differ
Male Size Index	$\theta = mean\left(\dfrac{L(a)p_m(a)}{\bar{N}_T}\right)$	θ is the mean probability a territorial male is mature given its length, weighted by the number of territorial males of that size alive in the steady state population
Care Ratio	$R = \dfrac{\theta_{fished}^p \Big/ \sum_a^{a_{max}} P(a)e^{-M(i=f,a)-F(a)}}{\theta_{unfished}^p \Big/ \sum_a^{a_{max}} N(i=f,a,t)p_m(a)bW(i=f,a)\,e^{-M(i=f,a)}}$	The parameter p shapes nonlinearity of the male size-care relationship (1 = linear). R is the proportional change in care-per-egg expected after fishing removes some males and females.
Population dynamics through time	$N(i, a+1, t+1) = \begin{cases} N_0(i,t) & if\ a = 0 \\ N(i,a,t)\,e^{-(M(a)+F(a))} & if\ a > 0 \end{cases}$	$N(i,a,t)$ is the number of individuals of each life history type i at age a alive at time t. $N_0(i,t)$ is made up of equal numbers of males and females. Male tactics (for the wrasse) occur in equal proportions at birth

	Table 3. Simulation parameters					
N	Initial population size (assumed 50:50 sex ratio, 25% of males assumed to be sneakers)	1000				
T_{max}	Number of years in our population simulation.	200				
$T_{fishing}$	Time when fishing mortality begins	100				
		Corkwing wrasse parameters			**Lingcod parameters**	
		Females	**Territorial males**	**Sneaker males**	**Females**	**Territorial males**
$M(a)$	Natural mortality rate (assumed constant)	0.54	0.54	0.40	0.18	0.32
A_{max}	Maximum age (years)	8	8	9	20	14
L_{inf}	Asymptotic size in von Bertalanffy growth function	177.5 mm	199.72 mm	180.8 mm	112.8 cm	81.87 cm
k	Growth coefficient in von Bertalanffy growth function	0.2972	0.3994	0.2625	0.145	0.223
L	Length-at-age; L(1) is reported here as the initial size at recruitment	81.6 mm	93.1 mm	78.3 mm	35 cm	34 cm
ν	Scale coefficient relating length to body mass (in kg)	6.6×10^{-7}	9.3×10^{-7}	8.5×10^{-7}	1.7×10^{-3}	4.0×10^{-3}
ω	Shape exponent relating length to body mass	3.176	3.089	3.115	3.40	3.21
a_{100}	Age at maturation (wrasse females/sneaker males)	2	-	1	-	-
γ	Coefficient of length-based maturation function for wrasse nesting males	-	18.5836	-	-	-
ρ	Shape coefficient of length-based maturation function for wrasse nesting males	-	0.13361	-	-	-
a_{mat}	Age when 50% of lingcod are mature	-	-	-	1.129	1.124
q	Steepness of age-based maturation function for lingcod	-	-	-	3.814	3.233
b	Mass-specific fecundity coefficient (mass in kg)	2.95×10^{5}	-	-	2.82×10^{-4}	-
c	Mass-specific fecundity exponent	1			3.0011	-
	Recruitment parameters					
α	Steepness parameter of the Beverton-Holt recruitment function	0.05			0.1	
β	Parameter of the Beverton-Holt recruitment function	8×10^{-6}			1×10^{-5}	
	Exploitation parameters					
-	Minimum size limit (mm)	120			26 inches (66.04 cm)	
-	Maximum size limit (slot)	170			36 inches (91.44 cm)	
F	Fishing mortality rate	Ranged from 0 to 0.7				

a) **Corkwing wrasse**

b) **Lingcod**

c)

d)

Corkwing wrasse

a) Minimum size limit   b) Slot size limit   c) Percent decrease in yield with slot limit

Fishing mortality

Corkwing wrasse

Lingcod

a) Minimum size limit

b) Slot size limit

c) Percent decrease in yield with slot limit

Fishing mortality

Lingcod