Synergism between microRNA-124 and ELAVL3 drives neuronal gene

upregulation during direct neuronal reprogramming of human fibroblasts

Ya-Lin Lu[1,2,3], Yangjian Liu[1,2], and Andrew S. Yoo[1,2*]

[1]Department of Developmental Biology, Washington University School of Medicine, St. Louis,

MO 63110, USA

[2]Center for Regenerative Medicine, Washington University School of Medicine, St. Louis, MO

63110, USA

[3]Program in Developmental, Regenerative and Stem Cell Biology, Washington University

School of Medicine, St. Louis, MO 63110, USA

*Correspondence: yooa@wustl.edu

Summary

Human neurons generated by direct conversion of adult fibroblasts can serve instrumental tools

for modeling diseases as well as a platform to study neurogenic processes. Neuron-enriched

microRNAs (miRNAs), miR-9/9* and miR-124 (miR-9/9*-124), direct cell fate switching of

human fibroblasts to neurons when ectopically expressed. A critical component of the miRNA-

mediated conversion process is the role of miR-9/9*-124 targeting and repressing anti-

neurogenic genes. However, how these miRNAs function beyond the onset of the neuronal

program remains unclear. Here, we performed the Argonaute (AGO) HITS-CLIP analysis to map

direct target transcripts of miR-9/9*-124 when cells start adopting the neuronal fate during the

conversion (reprogramming) process. Surprisingly, we found that miR-124, in particular, targets

a suite of genes associated with neuronal differentiation and function, and these targets appeared

to be upregulated by miR-124, as knocking down miR-124 function diminished the expression of

these targets. A critical example of the upregulated miR-124 target genes is *PTBP2*, a neuron-

enriched RNA-binding protein known to be repressed by its homolog PTBP1. We elected to

focus on *PTBP2* to gain insights into the mechanism underlying the activity of miR-124 as an

inducer of its target genes. Paradoxically, both *PTBP1* and *PTBP2* are repressive targets of miR-

124 in non-neuronal cells, yet PTBP2 selectively escapes the repression and becomes enriched in

neurons. By interrogating 3'UTR sequences of *PTBP1* and *PTBP2*, we provide evidence that the

switching of miR-124 activity to an activator is determined by the sequence availability at the

target transcript that allows the binding of neuronal RNA-binding proteins, in particular

ELAVL3 for select upregulation of PTBP2. ELAVL3 binds and interacts with AGO at PTBP2

3'UTR, yet ELAVL3 binding alone without miR-124 sites is insufficient to trigger the

upregulation of PTBP2. Knocking down ELAVL3 expression during neuronal reprogramming

leads to PTBP2 downregulation and impairs neuronal conversion, highlighting the concerted

interplay between miR-124 and ELAVL3 that drives neuronal fate acquisition. By elucidating

how miR-124 promotes PTBP2 induction, our study lends support to the underexplored role of

miRNAs in promoting target gene expression in neurons.

Introduction

MiR-9/9* and miR-124 (miR-9/9*-124) are efficient reprogramming effectors that, when

ectopically expressed in human adult fibroblasts (HAFs), induce an extensive reconfiguration of

the chromatin accessibility landscape leading to the erasure of fibroblast fate and activation of

the neuronal program (Abernathy et al. 2017). The conversion process by miR-9/9*-124 shares

similarities to molecular cascades underlying neurogenesis during neural development such as

the downregulation of REST, a well-established transcription repressor of neuronal genes (Ballas

et al., 2005; Lee et al., 2018; Schoenherr and Anderson, 1995) and switching of homologous

chromatin modifiers from non-neuronal to neuronal counterparts including DNMT3B to

DNMT3A, subunits of BAF/BRM-associated factor (BAF) complexes, and TOP2A to TOP2B

(Abernathy et al. 2017; Lee et al. 2018; Lessard et al. 2007; Staahl et al. 2013; Watanabe et al.

1994; Yoo et al. 2009; Tsutsui et al. 2001). Although the direct repression by brain-enriched

miRNAs on non-neuronal targets, for instance, *REST* (Packer et al., 2008; Visvanathan et al.,

2007), *ACTL6A* (Yoo et al., 2009), and *PTBP1* (Makeyev et al., 2007) for the induction of

neuronal counterparts offer a classic representation of miRNAs acting as negative regulators of

downstream target genes, how these miRNAs function in the midst of neuronal milieu remains

unknown.

The robustness of miRNA-based neuronal reprogramming allows for the investigation of

the pathways downstream of miR-9/9*-124 that promote neuronal identity (Abernathy et al.

2017). In the present study, we employed high-throughput sequencing of RNA isolated by

crosslinking immunoprecipitation (HITS-CLIP) of Argonaute (AGO) during miR-9/9*-124-

mediated neuronal reprogramming of human fibroblasts to map miR-9/9*-124-target interactions

at the onset of neuronal fate acquisition. Unexpectedly, we identified a suite of neuronal genes

enriched with AGO binding, in particular, corresponding to the binding sites of miR-124. The

upregulation of these neuronal genes required miR-124, suggesting that miR-124 not only

function as a repressor, but also as an effector to promote neuronal gene expression. Although

the ability of miRNAs as a positive effector of downstream target genes has been implicated

before (Truesdell et al., 2012; Vasudevan et al., 2007; Vasudevan and Steitz, 2007), little is

known about the molecular mechanism that governs a miRNA's activity as an activator in

neurons.

Among the genes identified to be targeted and upregulated by miR-124 was *PTBP2*, a

neuron-enriched RNA-binding protein (RBP). We elected to further focus on *PTBP2* to

understand the mechanism underlying miR-124 as an activator because the miR-124-PTBP2 axis

presented intriguing questions. A non-neuronal homolog *PTBP1* (Boutz et al., 2007) and *PTBP2*

both contain miR-124 target sites at their 3'UTRs (Makeyev et al., 2007; Xue et al., 2016). Why

is there selective repression on *PTBP1* while *PTBP2* escapes the repression by miR-124 in

neurons? The current view on the initial activation of *PTBP2* points to the activity of PTBP1 as a

repressor of *PTBP2* that is relieved upon miR-124 expression at the onset of neuronal

differentiation (Makeyev et al., 2007). However, the mechanism by which *PTBP2* expression

persists while being targeted by miR-124 remains unknown. In this study, we describe a series of

experiments to provide evidence for an RNA-binding protein, ELAVL3, induced during

neuronal conversion, directly acting on *PTBP2* 3'UTR to mediate the mode of miR-124 as an

activator, and the synergism between miR-124 and ELAVL3 that drives the neuronal conversion

of human fibroblasts.

Results

AGO binds both downregulated and upregulated genes during miR-9/9*-124-mediated

direct neuronal reprogramming of human fibroblasts

We have previously shown that ectopically expressing miR-9/9*-124 in human adult fibroblasts

(HAFs) induced a neuronal state characterized by the appearance of neuronal markers (such as

MAP2, TUBB3, NCAM, and SNAP25) (Figure 1A) and electrical excitability (Abernathy et al.

2017). To infer the role of miR-9/9*-124 as converting cells acquire the neuronal identity, we

carried out AGO HITS-CLIP after two weeks into reprogramming, a time point when neuronal

genes start being activated (Abernathy et al. 2017), to identify transcripts bound with AGO

loaded with miR-9/9* or miR-124 over the non-specific miRNA (miR-NS) control (Log$_2$FC $\geq$ 2;

adj.P-value < 0.05). We then compared these hits to the list of differentially expressed genes

(DEGs) (by RNA-seq analysis; -2 $\leq$ Log$_2$FC $\geq$ 2; adj.P-value < 0.05) at day 20 of miRNA-

induced neurons (miNs) (Abernathy et al. 2017). As expected, we found target transcripts that

were downregulated consistent with the repressive mode of miRNAs (Figure 1B, 123 genes

labeled as green dots). Some of these are known targets of miR-124, such as *SOX9* (Farrell et al.

2011), *TEAD1* (Mucaj et al., 2015; Sun et al., 2015), and *VAMP3* (Chen et al. 2019).

Interestingly, we also discovered 301 unique gene transcripts with enriched AGO binding, which

were upregulated in day 20 miNs (Figure 1B, red dots). By gene ontology (GO) analysis, these

upregulated genes were associated with various neuronal processes such as synaptic transmission

and neuronal differentiation (red) in contrast to downregulated target genes (green) (Figure 1C).

Moreover, when examined against the time-course transcriptome analysis during neuronal

reprogramming (Abernathy et al. 2017), the expression of the identified neuronal target genes

showed continuous upregulation during the later time point of neuronal conversion (Figure 1D).

For example, transcripts of neuronal genes such as *MAP2*, *PTBP2*, and *SLC4A8* that were highly

expressed in day 20 miNs harbored AGO binding containing (Figure 1E). To determine the

fraction of upregulated neuronal genes that are likely actual targets of miR-9/9* and/or miR-124,

we extracted the AGO-enriched sequences and predicted the duplex formation by either miR-

9/9* and/or miR-124 through RNAhybrid (a maximum free energy threshold of -20 kcal/mol)

(Rehmsmeier, 2004). Of the 301 upregulated DEGs in miNs bound by AGO (Figure 1B), 261

(~86%) gene transcripts are predicted to contain miR-9/9* and/or miR-124 binding sites (Figure

1F). These transcripts harboring miR-9/9*-124 sites are also associated with various neuronal

GO terms such as in neuronal differentiation and synaptic processes (Figure 1F) as they include a

selection of neuronal genes identified earlier (Figures D-E). Of these upregulated miR-9/9*-124

target genes (in which approximately a half harboring target sites for both miR-9/9* and miR-

124), more than 90% of the total genes (240 genes) contain miR-124 target sequence, while less

than 60% of the genes (147 genes) contain miR-9/9* target sites (Figure 1G). These results

collectively demonstrate that the AGO-loaded transcripts differentially (up or down) respond to

miR-9/9*-124 during the neuronal conversion.

MiR-124 target genes are upregulated during neuronal conversion

As a larger fraction of the upregulated genes are predicted to contain miR-124 target sequences

(Figure 1G), we further tested if the upregulated genes were *bona fida* miR-124 targets by

knocking down miR-124 expression through the use of a tough decoy to inhibit the miRNA

activity (Bak et al., 2013; Haraguchi et al., 2009). The effect of the lentivirus-based tough decoy

for miR-124 (TuD-miR-124) was monitored after transduction by following the concurrent RFP

reporter expression and measuring the mature miR-124 level. TuD-miR-124 yielded more than

60% reduction of miR-124 expression in comparison to the control, non-specific miRNA tough

decoy (TuD-miR-NS) (Figure S1A-S1B). We then performed RNA-seq analysis on day 20 miNs

treated with either TuD-miR-NS or TuD-miR-124 to identify genes that fail to be upregulated

with TuD-miR-124 (Figure 2A, S1C) ($Log_2FC \leq 0.5$; adj.P-value < 0.01). When compared with

upregulated DEGs in day 20 miNs that were also enriched for AGO binding (Figure 1), we

identified 118 genes that failed to be upregulated upon miR-124 reduction (Figure 2A, Table S1).

With GO analysis, these identified genes were associated with neuronal terms such as neuronal

development and synaptic activity (Figure 2B).

To further analyze the miR-124 and target interactions, we looked into the potential

duplex formations between miR-124 and the identified AGO HITS-CLIP peak regions using

RNAhybrid (a maximum free energy threshold of -20 kcal/mol) (Rehmsmeier, 2004). From the

3'UTR sequences of the 118 genes, we identified a spectrum of miR-124 and target 3'UTR

duplex configurations from the canonical 2-8 seed base pairing to non-canonical base pairing

starting at position 3 and 4 (Figures 2C, S1D, and S2). These examples include *MAP2* 3'UTR, in

which miR-124 base pairing is predicted to start at position 3, while miR-124 target sequences

on *PTBP2* 3'UTR at the two peaks are predicted to start at position 1 and position 2, respectively

(Figure 2C). Interestingly, across the predicted miR-124:target duplex configurations, we

observed consistent auxiliary 2-3 base pairing at the 3' end of miR-124 (Figure S1D).

Our results so far support the notion that miR-124 can target and promote the expression

of select neuronal genes when cells acquire the neuronal fate during neuronal conversion. We

wondered if the active mode of miR-124 on neuronal genes can be recapitulated in non-neuronal

cell types. We cloned the 3'UTRs of the identified neuronal target genes into a luciferase

reporter construct and transfected into a non-neuronal cell type, HEK293T (a human embryonic

kidney cell line), with a miR-9/9*-124 expression construct. After 48 hours post-transfection, we

found that unlike miRNA-induced neurons, miR-9/9*-124, instead, led to the repression of the

luciferase signal in HEK293T cells in comparison to the control miR-NS, while the non-target

control, *PGK1* 3'UTR was unaffected (Figure 2D). This result suggests that additional

determinants available in neuronal cells may be in play to govern the activity of miR-124 as a

positive regulator while simultaneously functioning to repress non-neuronal targets (Figure 2E).

PTBP1 and PTBP2 3'UTR as targets of miR-124

To further dissect the mechanism underlying the dual modes of miR-124 on its targets, we

elected to focus on *PTBP2* from the list of our identified neuronal targets because i) *PTBP2* and

its non-neuronal homolog*, PTBP1*, both contain miR-124 sites in their 3'UTRs (Figure 3A), and

ii) both *PTBP1* and *PTBP2* 3'UTRs are targeted and repressed by miR-124 in HEK293T cells

(Figure 3A), and iii) contrastingly, as observed in the human brain, the neuronal conversion

establishes the mutually exclusive expression between *PTBP1* and *PTBP2* (Figure 3B).

Therefore, the PTB homologs represent an ideal example to investigate how two closely related

targets respond differently to miR-124.

The 3'UTRs of human *PTBP1* and *PTBP2* contain two predicted target sites for miR-124

(yellow bar) and one site for miR-9 (blue bar) (Figure 3A). Luciferase reporter assays in the non-

neuronal HEK293T cells showed that miR-9/9*-124 or miR-124 alone (but not miR-9/9* alone)

was sufficient to repress both *PTBP1* and *PTBP2* through their 3'UTRs indicating that miR-124

is the primary miRNA targeting both PTB 3'UTRs (Figure 3A). Moreover, mutating the two

miR-124 sites in *PTBP2* 3'UTR (s1 and s2) rendered the 3'UTR insensitive to miR-124 (Figure

S3A). While our results pointing to the ability of miR-124 to target and repress 3'UTRs of both

PTBP1 and *PTBP2* in cell lines were consistent with previous findings (Makeyev et al., 2007;

Xue et al., 2016), miRNA-mediated neuronal conversion of HAFs establishes the mutually

exclusive expression of PTBP1 and PTBP2 (Figure 3B), as *in vivo* neurons (Boutz et al., 2007).

Because *PTBP1* and *PTBP2* are regulated oppositely in miNs, we asked whether the

repressive activity of miR-124 on *PTBP2* 3'UTR could be reversed with a prolonged neurogenic

input by miR-9/9*-124 in HEK293T cells. We expressed a destabilized EGFP reporter

containing *PTBP1* 3'UTR, *PTBP2* 3'UTR, or control 3'UTR (CTL) that lacks a 3'UTR in

HEK293T cells with the miR-9/9*-124 expression construct (Figure SB). Analogous to PTBP2

upregulation during neuronal reprogramming of HAFs, we observed the selective increase in

EGFP expression with *PTBP2* 3'UTR within ten days of miR-9/9*-124 expression, contrasting

the continued repression of EGFP with *PTBP1* 3'UTR (Figures S3B-C). These results suggest

that with the prolonged neurogenic input, *PTBP2* 3'UTR responds to miR-124 differentially

from when measured after 48 hours with miRNA expression in HEK293T cells (Figure 3A).

MiR-124 accentuates PTBP2 expression beyond the induction mediated by the

downregulation of PTBP1

Previous studies have shown that during development, PTBP1 destabilizes *PTBP2* transcript in

non-neuronal cells by alternative splicing, and at the onset of neurogenesis, miR-124 directly

represses *PTBP1* resulting in PTBP2 induction (Boutz et al., 2007; Makeyev et al., 2007). Upon

miR-9/9*-124 expression in HAFs, we observed the concomitant downregulation and

upregulation of endogenous PTBP1 and PTBP2, respectively. As expected, the initial PTBP2

upregulation was abrogated when *PTBP1* cDNA was overexpressed (Figure 3C), supporting the

role of PTBP1 reduction as the initiation step of PTBP2 expression. We then sought to stratify

the contribution of PTBP1 repression alone versus the input of miRNAs to the overall PTBP2

level. When PTBP2 protein levels were compared between PTBP1 knockdown with a short

hairpin RNA (shRNA) and miR-9/9*-124 expression, we found that miR-9/9*-124 enhanced the

PTBP2 level by nearly a two-fold increase over the PTBP1 knockdown alone condition (Figure

3D). In fact, PTBP2 was more upregulated with miR-9/9*-124 despite the more pronounced

reduction of PTBP1 with shRNA (Figure 3D), demonstrating that miR-9/9*-124 accentuate

PTBP2 expression beyond the level induced by PTBP1 downregulation.

Functional significance of PTBP2 expression for neuronal conversion

To probe the functional importance of PTBP2 expression, we performed the knockdown

experiments of PTBP2 with an shRNA during neuronal conversion. PTBP2 shRNA completely

impaired the induction of neurons marked by the loss of MAP2 expression, a neuronal marker

(Figure 3E). The effect of shRNA was specific to PTBP2 knockdown as supplementing *PTBP2*

cDNA rescued the reprogramming defect (Figure 3E). These findings were somewhat surprising

as the results from a previous study indicated that reduction of PTBP2 by shRNA was found to

promote maturation of reprogrammed neurons (Xue et al. 2016). It is not clear why shRNA-

versus miRNA-based reprogramming approaches lead to these conflicting results. Our results,

however, clearly demonstrate the essential role of PTBP2 at the onset of neuronal conversion.

Furthermore, knocking down PTBP2 in primary cultured human neurons resulted in increased

cell death as measured with SYTOX assay (Figure S4A-B), consistent with results from previous

studies that demonstrated the essential function of PTBP2 in primary neurons (Li et al., 2014).

To further gain insights into PTBP2 function during neuronal conversion, we performed

Human Clariom D Assay in HAFs expressing the non-specific control miR-NS, miR-9/9*-124,

or miR-9/9*-124 with PTBP2 shRNA to identify genes whose expression and alternative splicing

patterns were affected by the reduction of PTBP2. By two weeks, miR-9/9*-124-expressing cells

showed significant downregulation of *PTBP1* and other non-neuronal fibroblast genes (*FBN1*

and *S100A4*), and upregulation of *PTBP2* and neuronal genes (*NEFM,* and *SNAP25*) in

comparison to the control miR-NS expression (FC $\geq$ 1.5, ANOVA p-value < 0.05) (Figure S4C)

(Abernathy et al., 2017). To examine the alternative splicing events mediated by PTBP2 during

neuronal reprogramming, differentially spliced events were compared between PTBP2 shRNA

and control shRNA (shCTL) conditions in the miR-9/9*-124 expression background. We found

that reducing PTBP2 led changes in splicing events (-2 $\leq$ splicing index $\geq$ 2, ANOVA p-value <

0.05) indicated by red (positive splicing index) and green dots (negative splicing index) (Figure

S4D). This analysis identified some of the known splicing targets of PTBP2, such as the exon

skipping or exclusion in *UNC13B*, *DLG4*, and *CADM3*, and inclusion in *DNM1* and *SMARCC2*

transcripts (Figure S4D) (Li et al., 2014; Licatalosi et al., 2012; Vuong et al., 2016; Zheng et al.,

2012). Comparing genes upregulated in response to miR-9/9*-124 to genes associated with

PTBP2-mediated alternative splicing (differential splicing events between miR-9/9*-124 shCTL

and shPTBP2 conditions), identified 1183 differentially spliced transcripts of genes involved in

processes such as neuronal differentiation and signaling (Figure S4E) (Table S2). Altogether, our

results support the role of PTBP2 as a crucial regulator of the neuronal program during the

neuronal conversion of HAFs.

Differential sequence composition between PTBP1 and PTBP2 3'UTRs

We sought to identify effectors that promote miR-124 function in enhancing PTBP2 expression.

RNA-binding proteins (RBPs) have been shown to interact with miRNA-loaded RISC

complexes to modulate target gene expression (Iadevaia and Gerber, 2015; Jiang and Coller,

2012; Plass et al., 2017). We ran the sequence of *PTBP2* 3'UTR under three RBP motif

prediction databases, including RBPDB (Cook et al., 2011), RBPmap (Paz et al., 2014), and

beRBP (Yu et al., 2018) (Figure S5A). Across the three databases, two RBPs, ELAVL1 and

PUM2, were consistently predicted to bind to *PTBP2* 3'UTR (Figure S5A). As PTBP2 is

neuronal-enriched, we focused on the family of RBPs in which ELAVL1 belongs to, the

ELAVL-family proteins known to include neuron-specific homologs. Both the expression of

ELAVL1 and Pumilio family RBPs, *PUM2* or its homolog, *PUM1*, are not specific to neurons

(Lin et al., 2018; Okano and Darnell, 1997; Spassov and Jurecic, 2002). *ELAVL1* is known to be

ubiquitously expressed while the other three family members, *ELAVL2*, *ELAVL3*, and *ELAVL4*

(collectively referred to as neuronal *ELAVLs*, *nELAVLs*), are neuron-specific (Okano and

Darnell, 1997). As *PTBP2* is a neuronal gene, we reasoned that nELAVLs would be concurrently

enriched and bind to *PTBP2* 3'UTR as ELAVLs share RNA-binding domains (Okano and

Darnell, 1997). We examined the HITS-CLIP data of nELAVLs in the human brain (Scheckel et

al., 2016) and found nELAVLs to be highly enriched at *PTBP2* 3'UTR, in contrast to *PTBP1*

3'UTR (Figure S5B-C). This selective binding is likely due to the presence of AU-rich elements

(AREs) within *PTBP2* 3'UTR since ELAVLs have been shown to bind to AREs (Ince-Dunn et

al., 2012; Scheckel et al., 2016), and *PTBP1* 3'UTR lacks the ARE (Figure S5B). Moreover, the

binding of nELAVLs mapped to the ARE directly upstream of the first miR-124 site (s1; seed,

highlighted in red) within the 3'UTR of *PTBP2* (Figure S5C).

To examine if nELAVLs are induced during neuronal conversion, we measured the

expression of ELAVL-family binding proteins as well as other brain-enriched RBPs, including

NOVA-, RBFOX-family proteins, and SSRM4 by qPCR. We found that miNs express the

neuronal RBPs similarly to the human brain, such as the selective upregulation of nELAVLs, in

contrast to the ubiquitous expression of *ELAVL1* (Figure 4A) (Okano and Darnell, 1997). To

determine if nELAVL induction occurs concurrently with PTBP2 upregulation, we assessed

ELAVL expression at multiple time points during the neuronal conversion. We found that the

transcriptional activation of nELAVLs, with *ELAVL3* being the most robust one (blue), aligned

with the upregulation of *PTBP2* (black) by nine days into reprogramming (Figure 4B).

MiRNA-mediated PTBP2 induction requires ELAVL3 binding at PTBP2 3'UTR

We then tested whether ARE within *PTBP2* 3'UTR would serve as a sequence that binds

ELAVL proteins by expressing 3'UTRs of *PTBP1*, *PTBP2,* or *PTBP2* without ARE

(PTBP2ΔARE 3'UTR) with individual FLAG-tagged ELAVLs (ELAVL1-4) in HEK293T cells

(Figure 4C). RNA-immunoprecipitation (RIP) of FLAG-ELAVLs with FLAG antibody,

followed by qPCR for detecting the loaded 3'UTRs using primers specific for either *PTBP1* or

PTBP2 3'UTRs displayed significant enrichment for *PTBP2* 3'UTR with nELAVL (ELAVL2, 3,

and 4) pull-downs, while the binding of ELAVL1 to *PTBP2* 3'UTR was minimal. We could not

detect significant enrichment for *PTBP1* 3'UTR for any of the ELAVLs (Figure 4C).

Importantly, deleting ARE in *PTBP2* 3'UTR (PTBP2ΔARE 3'UTR) abolished the binding of

nELAVL proteins to *PTBP2* 3'UTR (Figure 4C) indicating that the ARE within *PTBP2* 3'UTR

serves to recruit nELAVLs.

We asked whether adding nELAVLs to the non-neuronal context of HEK293T cells,

thereby reconstituting nELAVLs that become available during neuronal conversion, would

alleviate miR-124-mediated repression of PTBP2 and enhance PTBP2 expression instead.

Adding individual ELAVLs to control *PGK1* 3'UTR had no effect on luminescence (relative

luminescence of ~ 1; *PGK1* 3'UTR histograms) (Figure 4D). Luciferase activities with *PTBP1*

3'UTR remained repressed upon miR-9/9*-124 expression compared to the control miR-NS

irrespective of the ELAVL addition (relative luminescence of < 1; *PTBP1* 3'UTR histograms)

(Figure 4D). MiR-9/9*-124 led to the repression *PTBP2* 3'UTR in the absence of ELAVLs

(black, relative luminescence ratio < 1; *PTBP2* 3'UTR histograms) (Figure 4D). However,

adding ELAVL2 (green) and ELAVL3 (blue) significantly alleviated miR-9/9*-124-mediated

repression on *PTBP2* 3'UTR, with especially ELAVL3 (blue) having the most significant effect

on elevating the luciferase activity in comparison to the control construct (Figure 4D). To

examine the requirement of ARE, we repeated the ELAVL addition experiments using the

luciferase cassette with *PTBP2* 3'UTR lacking the ARE sequence (*PTBP2ΔARE*). Deleting ARE

abolished the effect of ELAVL3 (blue) on *PTBP2* (*PTBP2ΔARE* 3'UTR histograms), which

stayed repressed (Figure 4D), demonstrating the direct role of ELAVL3 at ARE of *PTBP2*

3'UTR in modifying the function miR-124.

Selective activity of ELAVL3 on PTBP2 3'UTR is dependent on the hinge region

ELAVL1-4 members exhibit high sequence homology across all three functional RNA

recognition motifs (RRMs) except for the non-conserved spacer region (also referred to as hinge

region) flanked by RRM2 and RRM3 (Hinman et al., 2013; Okano and Darnell, 1997). To better

understand the specificity of ELAVL3 on *PTBP2* 3'UTR regulation, we mutagenized the hinge

region of ELAVL1 and ELAVL3 by deleting or swapping the hinge region (H) between

ELAVL1 and ELAVL3. Deleting the H in *ELAVL3* (*ELAVL3ΔH*) (light blue) abrogated the

alleviating effect on *PTBP2* 3'UTR repression, whereas no effect was observed with *PGK1*

3'UTR (Figure 4E, *PGK1* 3'UTR histograms). Moreover, replacing the *ELAVL3* H with *ELAVL1*

H (*ELAVL3-E1H*) (dark blue) still led to the failure of alleviating the *PTBP2* 3'UTR repression

(Figure 4E), and none of the *ELAVL3* variants (wild-type and mutants) had any effect on

PTBP2ΔARE 3'UTR (Figure 4E, *PTBP2ΔARE* 3'UTR histograms). These results indicate that

the specificity of ELAVL3 to *PTBP2* 3'UTR is mediated by the *ELAVL3* H region. This notion

is further supported by the increase in the luminescence readout of *PTBP2* 3'UTR in HEK293T

cells when *ELAVL1* H is replaced by *ELAVL3* H (*ELAVL1-E3H*) (dark red) compared to wild-

type *ELAVL1* (red) (Figure 4F, *PTBP2* 3'UTR histograms). Our results altogether uncover the

specificity of ELAVL3 in promoting PTBP2 expression.

Synergism between nELAVL and AGO requires miR-124 site in PTBP2 3'UTR

We further tested whether the miR-124 sites within *PTBP2* 3'UTR would also be critical for

mediating the PTBP2 upregulation with ELAVL3. By mutating the miR-124 seed-match

sequences within *PTBP2* 3'UTR (*PTBP2Δs1s2* 3'UTR), adding ELAVLs failed to enhance the

luciferase activity over the control (CTL, black; relative luminescence ratio unchanged;

PTBP2Δs1s2 3'UTR histograms) (Figure 5A). This result is in contrast to wild-type *PTBP2*

3'UTR where ELAVL3 can enhance luciferase activity in the presence of miR-124 target sites

(Figures 4D, 5A). Interestingly, the lack of the increased luminescence with ELAVL3 addition

was not due to the failure of ELAVL binding to *PTBP2Δs1s2* 3'UTR because qPCR analysis

with nELAVL-RIP showed persistent binding of ELAVLs to *PTBP2Δs1s2* 3'UTR (Figure 5B).

These results altogether suggest the requirement of both ARE and miR-124 sites in *PTBP2*

3'UTR for PTBP2 upregulation (Figure 5C).

ELAVL3 promotes PTBP2 expression during neuronal reprogramming

To further examine if ELAVL3 would be critical for inducing PTBP2 expression during the

neuronal conversion of HAFs, we performed ELAVL3 knockdown by shRNA (shELAVL3) to

assess PTBP2 expression and neuronal reprogramming. Knocking down ELAVL3 resulted in the

significant downregulation of PTBP2 expression as determined by immunostaining, qPCR, and

immunoblotting analyses, and impairment of the conversion process (Figures 6A-C). This

knockdown effect was specific for ELAVL3 downregulation as PTBP2 expression and neuronal

fate acquisition could be rescued by overexpressing *ELAVL3* cDNA with shELAV3 (Figures 6A-

B). It is noteworthy that reducing the function of other nELAVLs (ELAVL2 and 4) had only a

minor effect on PTBP2 expression (Figures 6B-C), highlighting the role of ELAVL3 as a

primary driver for PTBP2 upregulation with miR-124.

MiR-124 and nELAVL interaction for other neuronal transcripts

To gain insights into whether the synergism of miR-124 and ELAVLs for miRNA-mediated

upregulation would be unique for PTBP2 or a broader mechanism to other neuronal genes, we

carried out a comparative analysis between multiple datasets: i) the previously identified DEGs

upregulated in response to miR-9/9*-124 at day 20 into reprogramming (Abernathy et al. 2017),

ii) DEGs bound by AGO HITS-CLIP (Figure 1), and iii) DEGs responsive to miR-124 tough

decoy (Figure 2), to iv) genes bound by nELAVLs in the human brain (Scheckel et al., 2016)

(Figure 6D). These multiple comparisons identified 54 genes, including *PTBP2* (Table S3),

enriched for neuronal biological terms generally involved in synaptic processes (Figure 6D).

Examples of these identified genes include *MAP2*, *SEMA6A, RCAN2, DOCK3*, *SCN1A*, and

CAMK1D, some of which some we showed as targets of miR-124 (Figure 2, Table S3). Although

the additional identified targets remain to be further dissected for the functional meaning

underlying the upregulation, our findings pointing to these transcripts harboring both AGO-miR-

124 and nELAVL binding sites suggest that the synergistic activities of miR-124 and nELAVL

for promoting gene expression may extend to other genes important for neuronal differentiation

and function (Figure 6E).

Discussion

In the present study, we uncovered the role of miR-124 in the upregulation of genes associated

with neuronal differentiation and function during the neuronal conversion of human fibroblasts.

This finding provides insights into the function of miRNAs in addition to their canonical role as

a repressor of downstream target genes. Of the bound AGO transcripts are *bona fida* miR-124

target genes in which the 3'UTRs can be repressed in a non-neuronal context, but reverses upon

neuronal induction. By examining the mutually exclusive regulation of *PTBP1* and *PTBP2*, we

reveal how miR-124 plays a bifunctional role depending on the sequence composition at the

3'UTR, the availability of neuronal ELAVLs, and the interaction with AGOs to mediate the

switching of PTB homolog expression during the neuronal conversion of HAFs. Although we

focus specifically on the interplay between miR-124 and ELAVL3 for PTBP2 upregulation,

future studies should also examine if other RBPs can also synergize with AGOs for such target

gene regulation. For example, FXR1, an RBP, has been shown in previous studies to interact

with AGOs to facilitate gene expression in non-neuronal cells (Truesdell et al., 2012; Vasudevan

et al., 2007; Vasudevan and Steitz, 2007).

The selective role of ELAVL3, and not ubiquitous ELAVL1 or neuronal-specific

ELAVL2 and ELAVL4, in mediating PTBP2 induction in neurons highlights the functional

specificity of ELAVL family members in neurons. Like ELAVL3, ELAVL1 binds AREs and has

been shown to interact with RISC components (Kim et al., 2009; Vasudevan and Steitz, 2007).

Our results, however, clearly defines the functional specificity inherent in ELAVL3 that cannot

be replaced by ELAVL1, especially for regulating PTBP2 expression with miR-124. We found

that the specificity is, at least in part, driven by the hinge region inferred by the region-swapping

experiments between ELAVL1 and ELAVL3. As there is no known ARE recognition unique to

each ELAVL family member, it is likely that interactors associating with the hinge region may

be regulating ELAVL specificity and targeting (Fujiwara et al., 2012; Hinman et al., 2013).

To investigate if the synergism of miR-124 and nELAVLs for transcript stabilization and

activation can be generalized to other neuronal transcripts beyond *PTBP2*, we examined the

existing nELAVL HITS-CLIP dataset and uncovered that miR-124- and nELAVL-mediated

upregulation of target transcripts may not be a unique occurrence to *PTBP2*, but likely an

overlooked mechanism that maintains gene expression in neurons. Interestingly, as we identified

numerous neuronal genes to be targets of miR-124, in addition to *PTBP2*, such as *MAP2,*

RCAN2, and SEMA6A, we argue that miR-124 may be critical for enhancing the overall neuronal

program. This notion also indicates that miR-9/9*-124 is highly neurogenic as these miRNAs not

only allow for the conversion of HAFs into neurons (Abernathy et al. 2017; Yoo et al. 2011), but

also enhanced the expression of neuronal markers, such as *MAP2*, when overexpressed during

neuronal differentiation of human pluripotent stem cell-derived neurons (Ishikawa et al., 2020;

Sun et al., 2016). Although further tests are required to see if all the identified transcripts

targeted by both miR-124 and nELAVLs share the same mechanism as PTBP2, our results from

miR124-TuD and its effect on other neuronal genes lend support to the general role of miR-124

as a positive regulator of some its target genes in neurons. Future experiments taking a closer

look at RNA structure, motif proximity with RBPs, and additional interacting proteins will

provide further mechanistic insights to how a single miRNA can simultaneously repress and

activate different transcripts in a cell context-dependent manner.

Our study offers insights into the bifunctional mode of miR-124 conferring miRNAs as a

reprogramming effector that can contribute to the neuronal program by upregulating specific

neuronal genes. As miRNAs have been typically checked for their targets in non-neuronal cell

lines, it is plausible that the dual-mode of miRNAs may not be unique to miR-124 in neurons,

but instead utilized and altered in other cellular contexts with the help of other cell type-specific

RBPs. The governance of miRNA activity by the sequences within 3'UTR highlights the role of

cell type-specific RBPs as core regulators of gene expression.

Materials and Methods

Cell culture

Primary human fibroblasts used in this study was from a 22-year-old female (GM02171, NIGMS

Coriell Institute for Medical Research) while human neonatal fibroblasts (ScienCell, 2310) was

used exclusively for the HITS-CLIP experiment. Primary human neurons were obtained

commercially (ScienCell, 1520) with gender and age of the source undisclosed. Fibroblasts were

maintained in high glucose Dulbecco's Modified Eagle Medium (Gibco, 11960044) containing

10% FBS (Gibco, 10437028), MEM non-essential amino acids (Gibco, 11140050), sodium

pyruvate (Gibco, 11360070), GlutaMAX (Gibco, 35050061), HEPES (Gibco, 15630080),

penicillin-streptomycin (Gibco, 15130122), and 2-mercaptoethanol (Gibco, 21985023) at 37°C.

Human neurons were maintained in neuronal media (NM; ScienCell, 1521) at 37°C.

Lenti-X 293T (Clontech, 632180) cells were maintained in high glucose Dulbecco's

Modified Eagle Medium (Gibco, 11960044) containing 10% FBS (Gibco, 10437028), MEM

non-essential amino acids (Gibco, 11140050), sodium pyruvate (Gibco, 11360070), GlutaMAX

(Gibco, 35050061), HEPES (Gibco, 15630080), penicillin-streptomycin (Gibco, 15130122), and

2-mercaptoethanol (Gibco, 21985023) at 37°C.

Primary mouse cortical neurons were maintained in Neurobasal-A medium (Gibco,

10888022), B-27 plus supplement (Gibco, A3582801), GlutaMAX (Gibco, 35050061), and

penicillin-streptomycin (Gibco, 15130122) at 37°C.

MiR-9/9*-124-mediated neuronal conversion

To initiate reprogramming, doxycycline-inducible pT-BclXL-miR-9/9*-124 (Addgene, 60857)

and reverse tetracycline-controlled transactivator rtTA (Addgene, 66810) lentivirus with 8

ug/mL polybrene (Sigma, H9268) was added to a plate of confluent fibroblasts and spinfected at

37°C for 30 min at 1,000xG. Full media change with 1 μg/mL doxycycline (DOX; Sigma-

Aldrich, D9891) occurred the following day. Two days following transduction, cells underwent

another media change supplemented with DOX and respective antibiotics (Puromycin, Life

Technologies, A11138-03; Blasticidin S HCl, Life Technologies, A11139-03). Five days after

transduction, cells were plated onto poly-l-ornithine (Sigma-Aldrich, P4957), fibronectin

(Sigma-Aldrich, F4759), and laminin (Sigma-Aldrich, L2020) coated coverslips or onto 10cm^2

Primaria plates (Corning, 353803) followed by full media switch to neuronal media (NM;

ScienCell, 1521) the next day supplemented with 200 μM dibutyl-cyclic AMP (cAMP; Sigma-

Aldrich, D0627), 1 mM valproic acid (VPA; Sigma Aldrich, P4543), 10 ng/mL human BDNF

(PeproTech, 450-02), 10 ng/mL human NT-3 (Peprotech, 450-03), 1 μM retinoic acid (RA;

Sigma-Aldrich, R2625), RevitaCell supplement (Gibco, A2644501), and antibiotics. DOX was

supplemented every 2 days while half media changes occurred every 4 days until day 30.

Plasmids and cloning

For luciferase assay, full length 3'UTR of target transcripts were cloned and ligated into

pmirGLO vector. For mutagenizing miR-124 target sites, QuikChange XL site-directed

mutagenesis kit (Agilent, 200516) was used according to the manufacturer's protocol. Using the

same UTR sequences, 3'UTR was attached immediately downstream of a destablized EGFP

reporter and subcloned into lentiviral vector. Sequences for shRNAs were synthesized through

Integrated DNA Technologies, annealed, and ligated into the pLKO.1 vector (Addgene, 8453 or

26655). Overexpression vectors of either lentiviral (N106 or N174) or mammalian expression

(pcDNA3.1+, Invtrogen, V79020) was cut using NotI cut site (NEB, R0189) for insert ligation.

Tough decoy for miR-124 was synthesized (GeneScript) and subcloned into pLemir vector using

MluI (NEB, R0198) and NotI sites.

Lentivirus production

Lentivirus was produced as previously described (Richner et al., 2015). Briefly, 1.5 µg pMD2.G,

4.5 µg psPAX2, 6 µg of plasmid in lentiviral backbone, 600 µl Opti-MEM (Life Technologies,

31985) and 48 µl of 2 mg/mL polyethyleneimine (PEI; Polysciences, 24765) were mixed and

transfected into Lenti-X 293T (Clontech, 632180) plated at 6×10^6 cells per 10 cm^2 dish. Media

was changed the following day, and viral supernatant was collected, filtered and spun at

70,000xG for 2 hr at 4°C two days later. The viral pellet collected per 10 cm^2 dish was

resuspended in 1 mL PBS.

Immunostaining analysis

Cells were fixed with 4% paraformaldehyde (PFA; Electron Microscopy Sciences, 15710) for 20

mins at room temperature (RT) followed by three washes with PBS. Cells were permeabilized

and blocked in 0.3% TritonX-100, 2% normal goat serum (NGS; Jackson ImmunoResearch

Laboratories, 005-000-121) and 5% bovine serum albumin (BSA; Sigma-Aldrich, A7906) in

PBS for 1 hr at RT prior to incubation with primary antibodies overnight at 4°C. After three

washes with PBS, cells were incubated with respective secondary antibodies for 1 hr at RT.

Coverslips were mounted onto coverslides with ProLong Gold antifade reagent (Invitrogen,

P36934) for imaging using Leica SP5X white light laser confocal system with Leica Application

Suite (LAS) Advanced Fluorescence. See Supplemental Table S4 for a list of antibodies used.

SYTOX assay

488 SYTOX assay was performed as previously described (Victor et al., 2018). Briefly, 0.1 µM

489 SYTOX gene nucleic acid stain (Invitrogen, S7020) and 1 µl/mL of Hoeschst 33342 (Thermo

490 Scientific, 66249) were added into cell medium. Samples were incubated for at least 15 mins in

491 37°C prior to imaging. Images were taken using Leica DMI 4000B inverted microscope with

492 Leica Application Suite (LAS) Advanced Fluorescence.

Luciferase assay

HEK 293 cells plated in 96-well plate were transfected with 100 ng of pSilencer-miRNA, 100 ng

of pmirGLO containing 3'UTR of interest, and PEI (Polysciences, 24765) with Opti-MEM (Life

Technologies, 31985). Forty-eight hours after transfection, luciferase activity was assayed using

Dual-Glo luciferase assay system (Promega, E2920) according to the manufacturer's protocol

using Synergy H1 Hybrid plate reader (BioTek). Luciferase activity was obtained by normalizing

firefly luminescence to renilla luminescence (luciferase activity = firefly/renilla) followed by

normalizing to respective pSilencer-miR-NS control.

Flow cytometry

Destabilized EGFP reporter with or without 3'UTR of interest was transduced into HEK 293

cells to establish a stable reporter containing cell line with Blasticidin S HCl (Life Technologies,

A11139-03) selection. The day following transduction with miR-9/9*-124 lentivirus, media was

changed with the addition of DOX (Sigma-Aldrich, D9891). At day 3, media change was

supplemented with DOX and puromycin (Life Technologies, A11138-03). DOX was

supplemented every 2 days following transduction. At day 10, cells were imaged, and collected

for flow cytometry. Briefly, cells were collected in PBS and incubated with propidium iodide

(PI; Sigma-Aldrich, P4861) on ice until ready. Using FACSCalibur (BD Biosciences), all PI-

negative cell population was obtained for the gating of GFP-negative and -positive cell

population.

Quantitative reverse transcription PCR

Total RNA of cells was extracted using TRIzol Reagent (Invitrogen, 15596026). Reverse

transcription was performed using SuperScript III first strand synthesis system for RT-PCR

(Invitrogen, 18080-051) according to the manufacturer's protocol from fibroblasts,

reprogrammed neurons, and human brain total RNA (Invitrogen, AM7962). Quantitative PCR

was performed using SYBR Green PCR master mix (Applied Biosystems, 4309155) and

StepOnePlus Real-Time PCR system (Applied Biosystems, 4376600) according to the

manufacturer's protocol against target genes.

HITS-CLIP

AGO HITS-CLIP was performed on cells after 2 weeks into reprogramming of miR-NS or miR-

9/9*-124-expressing neonatal fibroblasts (ScienCell, 2310). Cells were harvested, UV-

crosslinked, lysed, and processed according to Moore et al. 2014. Briefly, cross-linked cells were

lysed and treated with RQ1 DNase (Promega, M6101) and RNaseA (Thermo). Complex

containing AGO-miRNA-mRNA were immunoprecipitated overnight at 4°C with pan-AGO

antibody. The immunoprecipitated complex was radio-labelled and extracted after running on

NuPAGE gel (Thermo). RNA from the 130kDa band was extracted for sequencing using TruSeq

Small RNA Library Preparation Kits (San Diego, CA). Samples were sequences using Illumina

HiSeq 2500 platform at the Genome Technology Access Center (GTAC) at Washington

University School of Medicine, St. Louis.

MiRNA Tough Decoy

Total RNA was extracted from day 20 cells expressing miR-9/9*-124 + TuD-miR-NS, and miR-

9/9*-124 + TuD-miR-124 using TRIzol Reagent (Invitrogen, 15596026) in combination with

RNeasy micro kit (Qiagen, 74004). RNA quality (RIN > 9.6) was determined with 2100

Bioanalyzer (Agilent) and samples underwent low input Takara-Clontech SMARTer kit (Takara,

639490) library preparation. Samples were sequenced using NovaSeq S4 and processed at the

Genome Technology Access Center (GTAC) at Washington University School of Medicine, St.

Louis.

Human Clariom D Microarray

For Human Clariom D Array (Affymetrix), total RNA was extracted from day 14 cells

expressing miR-NS, miR-9/9*-124 + shCTL, and miR-9/9*-124 + shPTBP2 using TRIzol

Reagent in combination with RNeasy mini kit (Qiagen, 74104). RNA quality (RIN > 9.6) was

determined with 2100 Bioanalyzer and samples (biological duplicates each) underwent

amplification and hybridization according to manufacturer's protocol by GTAC at Washington

University School of Medicine, St. Louis.

Immunoblot analysis

Cells were lysed with sonication (Diagenode, UCD-200) in RIPA buffer (Thermo Scientific,

89900) supplemented with protease inhibitor cocktail tablet (Roche, 04693132001). Protein

concentration of cleared lysate was measured using Pierce BCA protein assay kit (Thermo

Scientific, 23227) and read with Synergy H1 Hybrid plate reader (BioTek). Lysate and sample

buffer (Life Technologies, NP0008) were boiled, separated with Bis-Tris gels, and transferred to

nitrocellulose membrane (GE Healthcare Life Sciences, 10600006). Membrane was blocked with

5% milk for 1 hr at RT and incubated with primary antibody overnight at 4°C. After three

washes of TBST (1X TBS and 0.1% Tween-20), the membrane was incubated with respective

horseradish peroxidase-conjugated antibody for 1 hr at RT followed by three washes with TBST.

Blots were developed with ECL system (Thermo Scientific, 34580) and imaged or developed

onto film. See Supplemental Table S3 for a list of antibodies used.

Immunoprecipitation analysis

Cells were lysed with sonication (Diagenode, UCD-200) in IP buffer (20 mM HEPES, 150 mM

NaCl, 10% glycerol, 5 mM EDTA, 1% Triton X-100) supplemented with protease inhibitor

cocktail tablet (Roche, 04693132001). Cleared lysate was incubated with anti-FLAG M2

magnetic beads (Sigma-Aldrich, M8823) overnight with rotation at 4°C. The beads were washed

three times with IP buffer and bound proteins were boiled and eluted with sample buffer (Life

Technologies, NP0008), separated with Bis-Tris gels, and immunoblotted.

RNA-IP

Following immunoprecipitation with anti-FLAG magnetic beads (Sigma-Aldrich, M8823) or

anti-V5 antibody (Invitrogen, MA5-15253) coupled to Dynabeads protein G (Life Technologies,

NP10003D), beads were resuspended in 90 ul of IP buffer with proteinase K (NEB, P8107S) for

30 mins at 37°C. To extract RNA, 1 mL of TRIzol Reagent (Invitrogen, 15596026) was added to

the bead slurry. Final RNA was DNase I (Invitrogen, 18068015) treated prior to RT-qPCR.

GO enrichment analysis

Gene ontology analyses were performed using Metascape (Tripathi et al., 2015) with minimum

overlap of 3, P value cutoff of 0.01, and minimum enrichment of 1.5. Entire gene list was used

for each GO analysis.

RNAhybrid miRNA-target duplex analysis

To predict the presence of miR-124-3p binding sites at HITS-CLIP peaks, peak sequences were

extracted by on the genomic coordinates and processed through RNAhybrid (Rehmsmeier, 2004)

against miR-124-3p miRNA sequence with at least a free energy threshold of -20 kcal/mol.

RBP binding analysis

RNA-binding protein prediction database/software, RBPDB (Cook et al., 2011)), RBPmap (Paz

et al., 2014), and beRBP (Yu et al., 2018) for *PTBP2* 3'UTR sequence were used. Default

criteria were selected for all three software for non-bias prediction of any human RBP motifs.

Data Analyses

597   AGO HITS-CLIP reads for miR-NS and miR-9/9*-124-expressing cells at least two

weeks into reprogramming were trimmed and aligned to human genome hg38 using STAR with

default parameters. Aligned reads at CDS and UTRs were binned at a continuously 100 bp bins.

Two separate sets of bins were generated of which the two sets are offset by 50 bp in order to

capture all alignments. Aligned reads were quantified using featureCounts and differentially

present bins or peaks were determined using DESeq2 with the following criteria: adj. p-value <

0.05 and LogFC of $\leq$ -2 and $\geq$ 2. HITS-CLIP datasets will be publicly available.

604   RNA-seq for day 20 miNs expressing either CTL TuD-miR-NS or TuD-miR-124 was

605   analyzed by GTAC's RNA-seq pipeline. Briefly, reads were aligned to hg38 with STAR and

606   processed through EdgeR to obtain differentially expressed genes with adj. p-value of < 0.05.

607   RNA-seq dataset consisting of TuD-miR-NS and TuD-miR-124 at day 20 will be publicly

608   available.

609   Human Clariom D Array data was analyzed using manufacturer's software, Expression

Console followed by Transcriptome Analysis Console (TAC). For gene level analysis, genes

comparing miR-9/9*-124 vs miR-NS linear fold change of $\geq$ 1.5 and ANOVA P < 0.05 were

considered to upregulated in neuronal conditions. For splicing analysis, splicing events with

linear splicing index $\leq$ -2 and $\geq$ 2, ANOVA P < 0.05 were considered significant splicing events.

Human Clariom D array dataset consisting of miR-NS, miR-9/9*-124 with shCTL, and miR-

9/9*-124 with shPTBP2 at day 14 will be publicly available.

Author Contributions

Y.L.L. conducted experiments shown in Figures 2-5 and associated supplementary Figures 1, 3-

6. Y.J.L. conducted experiments shown in Figure 1 and supplementary Figure 2. A.S.Y.

supervised the study. Y.L.L. and A.S.Y. wrote the manuscript.

Acknowledgements

We thank the GTAC at Washington University for the sequencing service and support, and flow

Cytometry Core at Washington University for equipment use. We thank We thank P. Gontarz,

for the bioinformatics analysis of AGO HITS-CLIP, D. Annamalai and A. H. Kim for the help

with tough decoy designs, and K. Cates and L. Capano for the helpful suggestions with the

manuscript. Y.L.L. is supported by the LIFENAD Fellowship. This study was supported through

awards and funds to A.S.Y by the Andrew B. and Virginia C. Craig Faculty Fellowship

endowment, NIH Director's Innovator Award (DP2NS083372), Presidential Early Career Award

for Scientists and Engineers (PECASE), and NIH (RF1AG056296 and R01NS107488).

Declaration of Interests

The authors declare no competing interest.

References

Abernathy DG, Kim WK, McCoy MJ, Lake AM, Ouwenga R, Lee SW, Xing X, Li D, Lee HJ,
    Heuckeroth RO, Dougherty JD, Wang T, Yoo AS. 2017a. MicroRNAs Induce a
    Permissive Chromatin Environment that Enables Neuronal Subtype-Specific
    Reprogramming of Adult Human Fibroblasts. *Cell Stem Cell* **21**:332-348.e9.
    doi:10.1016/j.stem.2017.08.002

Bak RO, Hollensen AK, Primo MN, Sorensen CD, Mikkelsen JG. 2013. Potent microRNA
suppression by RNA Pol II-transcribed "Tough Decoy" inhibitors. *RNA* **19**:280–293.
doi:10.1261/rna.034850.112
Ballas N, Grunseich C, Lu DD, Speh JC, Mandel G. 2005. REST and Its Corepressors Mediate
Plasticity of Neuronal Gene Chromatin throughout Neurogenesis. *Cell* **121**:645–657.
doi:10.1016/j.cell.2005.03.013
Boutz PL, Stoilov P, Li Q, Lin C-H, Chawla G, Ostrow K, Shiue L, Ares M, Black DL. 2007. A
post-transcriptional regulatory switch in polypyrimidine tract-binding proteins
reprograms alternative splicing in developing neurons. *Genes & Development* **21**:1636–
1652. doi:10.1101/gad.1558107
Chen Y, Sun J, Chen W, Wu G, Wang Y, Zhu K, Wang J. 2019. miR-124/VAMP3 is a novel
therapeutic target for mitigation of surgical trauma-induced microglial activation. *Sig*
Transduct Target Ther **4**:27. doi:10.1038/s41392-019-0061-x
Cook KB, Kazan H, Zuberi K, Morris Q, Hughes TR. 2011. RBPDB: a database of RNA-
binding specificities. *Nucleic Acids Research* **39**:D301–D308. doi:10.1093/nar/gkq1069
Farrell BC, Power EM, Dermott KWM. 2011. Developmentally regulated expression of Sox9
and microRNAs 124,128 and 23 in neuroepithelial stem cells in the developing spinal
cord. *International Journal of Developmental Neuroscience* **29**:31–36.
doi:10.1016/j.ijdevneu.2010.10.001
Fujiwara T, Fukao A, Sasano Y, Matsuzaki H, Kikkawa U, Imataka H, Inoue K, Endo S,
Sonenberg N, Thoma C, Sakamoto H. 2012. Functional and direct interaction between
the RNA binding protein HuD and active Akt1. *Nucleic Acids Research* **40**:1944–1953.
doi:10.1093/nar/gkr979
Haraguchi T, Ozaki Y, Iba H. 2009. Vectors expressing efficient RNA decoys achieve the long-
term suppression of specific microRNA activity in mammalian cells. *Nucleic Acids*
Research **37**:e43–e43. doi:10.1093/nar/gkp040
Hinman MN, Zhou H-L, Sharma A, Lou H. 2013. All three RNA recognition motifs and the
hinge region of HuC play distinct roles in the regulation of alternative splicing. *Nucleic*
Acids Research **41**:5049–5061. doi:10.1093/nar/gkt166
Iadevaia V, Gerber A. 2015. Combinatorial Control of mRNA Fates by RNA-Binding Proteins
and Non-Coding RNAs. *Biomolecules* **5**:2207–2222. doi:10.3390/biom5042207
Ince-Dunn G, Okano HJ, Jensen KB, Park W-Y, Zhong R, Ule J, Mele A, Fak JJ, Yang C, Zhang
C, Yoo J, Herre M, Okano H, Noebels JL, Darnell RB. 2012. Neuronal Elav-like (Hu)
Proteins Regulate RNA Splicing and Abundance to Control Glutamate Levels and
Neuronal Excitability. *Neuron* **75**:1067–1080. doi:10.1016/j.neuron.2012.07.009
Ishikawa M, Aoyama T, Shibata S, Sone T, Miyoshi H, Watanabe H, Nakamura M, Morota S,
Uchino H, Yoo AS, Okano H. 2020. miRNA-Based Rapid Differentiation of Purified
Neurons from hPSCs Advancestowards Quick Screening for Neuronal Disease
Phenotypes In Vitro. *Cells* **9**:532. doi:10.3390/cells9030532
Jiang P, Coller H. 2012. Functional Interactions Between microRNAs and RNA Binding
Proteins. *MicroRNA e* **1**:70–79. doi:10.2174/2211536611201010070
Kim HH, Kuwano Y, Srikantan S, Lee EK, Martindale JL, Gorospe M. 2009. HuR recruits let-
7/RISC to repress c-Myc expression. *Genes & Development* **23**:1743–1748.
doi:10.1101/gad.1812509
Lee SW, Oh YM, Lu Y-L, Kim WK, Yoo AS. 2018. MicroRNAs Overcome Cell Fate Barrier by
Reducing EZH2-Controlled REST Stability during Neuronal Conversion of Human Adult
Fibroblasts. *Developmental Cell* **46**:73-84.e7. doi:10.1016/j.devcel.2018.06.007

Lessard J, Wu JI, Ranish JA, Wan M, Winslow MM, Staahl BT, Wu H, Aebersold R, Graef IA,
Crabtree GR. 2007. An Essential Switch in Subunit Composition of a Chromatin
Remodeling Complex during Neural Development. *Neuron* **55**:201–215.
doi:10.1016/j.neuron.2007.06.019
Li Q, Zheng S, Han A, Lin C-H, Stoilov P, Fu X-D, Black DL. 2014. The splicing regulator
PTBP2 controls a program of embryonic splicing required for neuronal maturation. *eLife*
3. doi:10.7554/eLife.01201
Licatalosi DD, Yano M, Fak JJ, Mele A, Grabinski SE, Zhang C, Darnell RB. 2012. Ptbp2
represses adult-specific splicing to regulate the generation of neuronal precursors in the
embryonic brain. *Genes & Development* **26**:1626–1642. doi:10.1101/gad.191338.112
Lin K, Zhang S, Shi Q, Zhu M, Gao L, Xia W, Geng B, Zheng Z, Xu EY. 2018. Essential
requirement of mammalian *Pumilio* family in embryonic development. *MBoC* **29**:2922–
2932. doi:10.1091/mbc.E18-06-0369
Makeyev EV, Zhang J, Carrasco MA, Maniatis T. 2007. The MicroRNA miR-124 Promotes
Neuronal Differentiation by Triggering Brain-Specific Alternative Pre-mRNA Splicing.
Molecular Cell **27**:435–448. doi:10.1016/j.molcel.2007.07.015
Moore MJ, Zhang C, Gantman EC, Mele A, Darnell JC, Darnell RB. 2014. Mapping Argonaute
and conventional RNA-binding protein interactions with RNA at single-nucleotide
resolution using HITS-CLIP and CIMS analysis. *Nat Protoc* **9**:263–293.
doi:10.1038/nprot.2014.012
Mucaj V, Lee SS, Skuli N, Giannoukos DN, Qiu B, Eisinger-Mathason TSK, Nakazawa MS,
Shay JES, Gopal PP, Venneti S, Lal P, Minn AJ, Simon MC, Mathew LK. 2015.
MicroRNA-124 expression counteracts pro-survival stress responses in glioblastoma.
Oncogene **34**:2204–2214. doi:10.1038/onc.2014.168
Okano HJ, Darnell RB. 1997. A Hierarchy of Hu RNA Binding Proteins in Developing and
Adult Neurons. *The Journal of Neuroscience* **17**:3024–3037.
doi:10.1523/JNEUROSCI.17-09-03024.1997
Packer AN, Xing Y, Harper SQ, Jones L, Davidson BL. 2008. The Bifunctional microRNA miR-
9/miR-9* Regulates REST and CoREST and Is Downregulated in Huntington's Disease.
Journal of Neuroscience **28**:14341–14346. doi:10.1523/JNEUROSCI.2390-08.2008
Paz I, Kosti I, Ares M, Cline M, Mandel-Gutfreund Y. 2014. RBPmap: a web server for mapping
binding sites of RNA-binding proteins. *Nucleic Acids Research* **42**:W361–W367.
doi:10.1093/nar/gku406
Plass M, Rasmussen SH, Krogh A. 2017. Highly accessible AU-rich regions in 3' untranslated
regions are hotspots for binding of regulatory factors. *PLOS Computational Biology*
13:e1005460. doi:10.1371/journal.pcbi.1005460
Rehmsmeier M. 2004. Fast and effective prediction of microRNA/target duplexes. *RNA*
10:1507–1517. doi:10.1261/rna.5248604
Richner M, Victor MB, Liu Y, Abernathy D, Yoo AS. 2015. MicroRNA-based conversion of
human fibroblasts into striatal medium spiny neurons. *Nature Protocols* **10**:1543–1555.
doi:10.1038/nprot.2015.102
Scheckel C, Drapeau E, Frias MA, Park CY, Fak J, Zucker-Scharff I, Kou Y, Haroutunian V,
Ma'ayan A, Buxbaum JD, Darnell RB. 2016. Regulatory consequences of neuronal
ELAV-like protein binding to coding and non-coding RNAs in human brain. *eLife* **5**.
doi:10.7554/eLife.10421

Schoenherr C, Anderson D. 1995. The neuron-restrictive silencer factor (NRSF): a coordinate
repressor of multiple neuron-specific genes. *Science* **267**:1360–1363.
doi:10.1126/science.7871435
Spassov DS, Jurecic R. 2002. Cloning and comparative sequence analysis of PUM1 and PUM2
genes, human members of the Pumilio family of RNA-binding proteinsq 10.
Staahl BT, Tang J, Wu W, Sun A, Gitler AD, Yoo AS, Crabtree GR. 2013. Kinetic Analysis of
npBAF to nBAF Switching Reveals Exchange of SS18 with CREST and Integration with
Neural Developmental Pathways. *Journal of Neuroscience* **33**:10348–10361.
doi:10.1523/JNEUROSCI.1258-13.2013
Sun AX, Yuan Q, Tan S, Xiao Y, Wang D, Khoo ATT, Sani L, Tran H-D, Kim P, Chiew YS,
Lee KJ, Yen Y-C, Ng HH, Lim B, Je HS. 2016. Direct Induction and Functional
Maturation of Forebrain GABAergic Neurons from Human Pluripotent Stem Cells. *Cell
Reports* **16**:1942–1953. doi:10.1016/j.celrep.2016.07.035
Sun Y, Luo Z-M, Guo X-M, Su D-F, Liu X. 2015. An updated role of microRNA-124 in central
nervous system disorders: a review. *Front Cell Neurosci* **9**. doi:10.3389/fncel.2015.00193
Tripathi S, Pohl MO, Zhou Y, Rodriguez-Frandsen A, Wang G, Stein DA, Moulton HM,
DeJesus P, Che J, Mulder LCF, Yángüez E, Andenmatten D, Pache L, Manicassamy B,
Albrecht RA, Gonzalez MG, Nguyen Q, Brass A, Elledge S, White M, Shapira S,
Hacohen N, Karlas A, Meyer TF, Shales M, Gatorano A, Johnson JR, Jang G, Johnson T,
Verschueren E, Sanders D, Krogan N, Shaw M, König R, Stertz S, García-Sastre A,
Chanda SK. 2015. Meta- and Orthogonal Integration of Influenza "OMICs" Data Defines
a Role for UBR4 in Virus Budding. *Cell Host & Microbe* **18**:723–735.
doi:10.1016/j.chom.2015.11.002
Truesdell SS, Mortensen RD, Seo M, Schroeder JC, Lee JH, LeTonqueze O, Vasudevan S. 2012.
MicroRNA-mediated mRNA Translation Activation in Quiescent Cells and Oocytes
Involves Recruitment of a Nuclear microRNP. *Scientific Reports* **2**.
doi:10.1038/srep00842
Tsutsui Ken, Tsutsui Kimiko, Sano K, Kikuchi A, Tokunaga A. 2001. Involvement of DNA
Topoisomerase IIβ in Neuronal Differentiation. *Journal of Biological Chemistry*
276:5769–5778. doi:10.1074/jbc.M008517200
Vasudevan S, Steitz JA. 2007. AU-Rich-Element-Mediated Upregulation of Translation by
FXR1 and Argonaute 2. *Cell* **128**:1105–1118. doi:10.1016/j.cell.2007.01.038
Vasudevan S, Tong Y, Steitz JA. 2007. Switching from Repression to Activation: MicroRNAs
Can Up-Regulate Translation. *Science* **318**:1931–1934. doi:10.1126/science.1149460
Victor MB, Richner M, Olsen HE, Lee SW, Monteys AM, Ma C, Huh CJ, Zhang B, Davidson
BL, Yang XW, Yoo AS. 2018. Striatal neurons directly converted from Huntington's
disease patient fibroblasts recapitulate age-associated disease phenotypes. *Nature
Neuroscience*. doi:10.1038/s41593-018-0075-7
Visvanathan J, Lee S, Lee B, Lee JW, Lee S-K. 2007. The microRNA miR-124 antagonizes the
anti-neural REST/SCP1 pathway during embryonic CNS development. *Genes &
Development* **21**:744–749. doi:10.1101/gad.1519107
Vuong JK, Lin C-H, Zhang M, Chen L, Black DL, Zheng S. 2016. PTBP1 and PTBP2 Serve
Both Specific and Redundant Functions in Neuronal Pre-mRNA Splicing. *Cell Reports*
17:2766–2775. doi:10.1016/j.celrep.2016.11.034
Watanabe M, Tsutsui Ken, Tsutsui Kimiko, Inoue Y. 1994. Differential expressions of the
topoisomerase IIa and IIb mRNAs in developing rat brain. *Neuroscience Research*
19:51–57.

Xue Y, Qian H, Hu J, Zhou B, Zhou Y, Hu X, Karakhanyan A, Pang Z, Fu X-D. 2016.
Sequential regulatory loops as key gatekeepers for neuronal reprogramming in human
cells. *Nature Neuroscience* **19**:807–815. doi:10.1038/nn.4297
Yoo AS, Staahl BT, Chen L, Crabtree GR. 2009. MicroRNA-mediated switching of chromatin-
remodelling complexes in neural development. *Nature* **460**:642–646.
doi:10.1038/nature08139
Yoo AS, Sun AX, Li L, Shcheglovitov A, Portmann T, Li Y, Lee-Messer C, Dolmetsch RE,
Tsien RW, Crabtree GR. 2011. MicroRNA-mediated conversion of human fibroblasts to
neurons. *Nature* **476**:228–231. doi:10.1038/nature10323
Yu H, Wang J, Sheng Q, Liu Q, Shyr Y. 2018. beRBP: binding estimation for human RNA-
binding proteins. *Nucleic Acids Research*. doi:10.1093/nar/gky1294
Zheng S, Gray EE, Chawla G, Porse BT, O'Dell TJ, Black DL. 2012. PSD-95 is post-
transcriptionally repressed during early neural development by PTBP1 and PTBP2.
Nature Neuroscience **15**:381–388. doi:10.1038/nn.3026
794

Figure 1. Enrichment of AGO binding on transcripts of neuronal genes upregulated during miR-9/9*-124-mediated neuronal conversion

(A) Examples of miR-9/9*-124-mediated direct reprogramming of human adult dermal fibroblasts (HAFs) into neurons (miNs). Photographs show immunostaining of the fibroblast marker (S100A4) and neuronal markers (MAP2, TUBB3, NCAM, and SNAP25). Scale bar = 20 μm.

(B) A volcano plot of differentially expressed genes at day 20 miNs and enrichment of AGO binding in response to miR-9/9*-124 expression identified by AGO HITS-CLIP analysis. Red dots, day 20 mRNA expression Log2FC ≥ 2; adj.P-value < 0.05, AGO binding Log2FC ≥ 2, adj.P-value < 0.05. Green dots, day 20 mRNA expression Log2FC ≤ 2; adj.P-value < 0.05, AGO binding Log2FC ≥ 2, adj.P-value < 0.05.

(C) Top biological GO terms of upregulated (red) and downregulated (green) DEGs in day 20 miNs differentially bound by AGO.

(D) Time course heatmap of select upregulated DEGs enriched for AGO binding with miR-9/9*-124 expression.

(E) Track views of AGO CLIP-seq tracks (top) and RNA-seq tracks (bottom) for gene examples showing the enrichment of AGO binding miR-9/9*-124 expression (over control miR-NS) and increased transcript levels at 3'UTRs.

(F) Of the 301 unique upregulated DEGs identified in (A), 261 are predicted to harbor miR-9/9* and/or miR-124 sites at the AGO-enriched regions through RNAhybrid prediction. The graph indicates top biological GO terms associated with the 261 upregulated DEGs containing miR-9/9* and/or miR-124 target sites.

(G) Breakdown of the 261 upregulated DEGs in (F) based on common or specific targets of miR-9/9* and/or miR-124.

Figure 2. Identification of miR-124 target genes that fail to be upregulated upon the inhibition of miR-124 during miRNA-mediated neuronal conversion

(A) Left, miR-124 activity is reduced through the use of tough decoy (TuD). By overlapping with upregulated DEGs bound by AGO (Figure 1), 118 genes were identified as genes that fail to be increased upon the reduction of miR-124. Right, a heatmap of z-scores of the 118 genes from RNA-seq comparing miNs between TuD-miR-NS and TuD-miR-124 treatments.

(B) Top biological GO terms associated with the 118 genes identified in (A).

(C) Predicted binding of miR-124 to the 3'UTR sequences of neuronal target genes, for example, *MAP2* and *PTBP2*, according to RNAhybrid prediction at the highlighted AGO HITS-CLIP peaks.

(D) Luciferase assays in HEK293T cells of upregulated neuronal target genes of miR-124 selected from A). In the non-neuronal context of HEK293T cells, 3'UTRs from the neuronal genes are targeted and repressed, instead, by miR-9/9*-124. Luminescence measured after 48 hrs of transfection and normalized to miR-NS control of each condition. Data are represented by mean ± SEM from three independent experiments (from left, ** P = 0.0094, P = 0.0011; *** P < 0.001; * P = 0.048, 0.015, 0.012; *** P < 0.001).

(E) A diagram of the observed phenomenon in which miR-9/9*-124 can promote neuronal identity by simultaneously targeting both non-neuronal genes for repression while promoting the expression of neuronal genes during neuronal conversion.

Figure 3. MiR-124 targets both PTBs, but differentially regulate PTB expression during neuronal conversion.
(A) Top, a schematic diagram of PTB 3'UTRs with miR-124 (yellow) and miR-9 (blue) target sites. S1 and s2 refer to the two conserved miR-124 target sites on PTBP2 3'UTR. Bottom, luciferase assays with luminescence measured after 48 hrs of transfection and normalized to miR-NS control in each condition. Data are represented by mean ± SEM from four independent experiments. Two-way ANOVA followed by Dunnett's test (from left, PTBP1 3'UTR ** P = 0.0043, 0.0034; PTBP2 3'UTR * P = 0.0355, ** P = 0.0016)
(B) Top, A heatmap of gene expression assessed by qPCR in starting HAFs, day 30 miRNA-induced neurons (miNs), and human brain RNA. Bottom, PTB switching is recapitulated during the miRNA-mediated direct conversion of HAFs into neurons. HAFs and day 30 miNs immunostained for PTBP1 and PTBP2, along with a fibroblast marker, S100A4, and a pan-neuronal marker, TUBB3. Scale bar = 50 μm.
(C) Initial induction of PTBP2 by the reduction of PTBP1 by either shRNA or miRNAs during miRNA-mediated neuronal reprogramming. PTBP1 overexpression in the miR-9/9*-124 expression background suppresses the PTBP2 induction.
(D) Left, an immunoblot showing that PTBP1 knockdown in HAFs resulted in induction of PTBP2, but PTBP2 expression becomes more pronounced in the presence of miR-9/9*-124 compared to PTBP1 knowdown only. Right, quantification of PTBP2 band intensity as relative fold changes compared to PTBP1 knockdown alone. Data were normalized to HSP90 from four independent experiments. The plots were represented in mean ± SEM. One-way ANOVA followed by Tukey's test (from top *** P = 0.0002, 0.0004; ns P = 0.8925).
(E) Top left, a schematic diagram of the experimental procedure. Right, photographs of day 30 miNs treated with CTL shRNA, PTBP2 shRNA, or PTBP2 shRNA with PTBP2 cDNA. Cells were immunostained for PTBP2 and MAP2. Bottom left, quantification of the percentage of MAP2-positive cells with two or more neurite processes over the total number of DAPI-positive cells. Scale bar = 50 μm. Data are represented as mean ± SEM. One-way ANOVA followed by Tukey's test (*** P < 0.0001; ** P 0.0016). shCTL n = MAP2 356/482; shPTBP2 n = MAP2 31/239; shPTBP2+PTBP2 n = MAP2 225/454.
(F) A model of differential miR-124 activity on PTB 3'UTRs during neuronal conversion.

Figure 4. MiRNA-mediated PTBP2 upregulation requires nELAVL binding at PTBP2 3'UTR

(A) An expression heatmap of neuronal-enriched RBPs determined by qPCR in starting HAFs, day 20 miNs, and human brain RNA.

(B) Time course qPCR assays of PTB and ELAVL transcripts during neuronal reprogramming.

(C) Left, a diagram of luciferase constructs containing different PTB 3'UTRs: PTBP1 3'UTR, and PTBP2 3'UTR with and without (PTBP2ΔARE) the AU-rich element (ARE) used for RIP and luciferase assays. Right, RIP of individual FLAG-tagged ELAVLs 48 hrs after transfection. Enrichment normalized to input determined by qPCR for PTBP1 or PTBP2 3'UTR. Data are represented by mean ± SEM from three independent experiments. Two-way ANOVA followed by Dunnett's test (from left, PTBP2 3'UTR *** P = 0.0001, ** P = 0.0086, 0.0042).

(D) Luciferase assays with the addition of individual ELAVLs in the luciferase constructs containing the control PGK1, PTBP1, PTBP2 or PTBP2ΔARE 3'UTR. Luminescence was measured 48 hrs after the transfection and normalized miR-9/9*-124 to miR-NS control of each condition. Data are represented by mean ± SEM from at least three independent experiments. Two-way ANOVA followed by Dunnett's test (* P = 0.0312, *** P = 0.0001).

(E) Left, a schematic diagrm of ELAVL3 hinge mutants. Right, luciferase assays with the addition of wild-type ELAVL3 or ELAVL3 hinge mutants (ELAVL3Δ H: ELAVL3 hinge deletion; ELAVL3-E1H: ELAVL3 with ELAVL1 hinge). Luminescence was measured 48 hrs after transfection and normalized miR-9/9*-124 to miR-NS control of each condition. Data are represented by mean ± SEM from at least four independent experiments. Two-way ANOVA followed by Tukey's test (from left ** P = 0.0020, * P = 0.0435, * P = 0.0286).

(F) Left, a schematic diagram of ELAVL1 hinge mutants. Right, luciferase assays with the addition of wild-type ELAVL1 or ELAVL1 hinge mutants (ELAVL1 ΔH: ELAVL1 hinge deletion; ELAVL1-E3H: ELAVL1 with ELAVL3 hinge). Luminescence was measured 48 hrs after transfection and normalized miR-9/9*-124 to miR-NS control of each condition. Data represented in mean ± SEM from at least four independent experiments. Two-way ANOVA followed by Tukey's test (** P = 0.0069, * P = 0.0260).

Figure 5. Synergism of nELAVL and miR-124 on PTBP2 3'UTR
(A) Top, a diagram of luciferase constructs containing PTBP2 3'UTR with or without miR-124 sites (PTBP2Δs1s2). Bottom, luciferase assays with the addition of individual ELAVLs in PGK1 control, PTBP2, or PTBP2Δs1s2 3'UTR. Luminescence was measured 48 hrs after transfection and normalized miR-9/9*-124 to miR-NS control of each condition. Data are represented by mean ± SEM from at least three independent experiments.Two-way ANOVA followed by Dunnett's test (* P = 0.0312, *** P = 0.0001).
(B) RIP of individual FLAG-tagged ELAVLs in PTBP2Δs1s2 3'UTR 48 hrs after transfection. Enrichment normalized to input determined through qPCR against PTBP2 3'UTR. Data are represented by mean ± SEM from three independent experiments. Two-way ANOVA followed by Dunnett's test (from left, PTBP2Δs1s2 3'UTR ** P = 0.0018, *** P = 0.0001, *** P = 0.0001).
(C) A schematic summary of nELAVL and RISC activities on PTB 3'UTRs.

Figure 6. PTBP2 upregulation during reprogramming requires ELAVL3

(A) Left, day 30 miNs with shRNA knockdown against CTL, ELAVL3, and ELAVL3 with ELAVL3 cDNA rescue. Cells were immunostained for MAP2 and PTBP2. Scale bar 50 = μm. Right, quantification of the percentage of MAP2-positive cells over the total number of DAPI-positive cells with two or more neurites (left). Data represented as mean ± SEM. One-way ANOVA followed by Tukey's test (from left, *** P < 0.0001, *** P < 0.0001). shCTL n = MAP2 803/1060; shELAVL3 n = MAP2 67/1052; shELAVL3 + ELAVL3 n = MAP2 431/617.

(B) Left, immunoblot analysis of PTBP2 in day14 miNs with knockdown against CTL, ELAVL2, ELAVL3, and ELAVL4. Right, quantification of the PTBP2 band intensity as a relative fold change compared to shCTL normalized to HSP90 in four independent experiments. Data are represented by mean ± SEM. One-way ANOVA followed by Dunnett's test (* P = 0.0422).

(C) The relative quantity of PTBP2 transcript upon individual nELAVL knockdown compared to CTL at day14 miNs determined by qPCR. Data are represented by mean ± SEM from 3 independent experiments. One-way ANOVA followed by Dunnett's test (*** P = 0.0001).

(D) Left, a Venn diagram of day 20 upregulated DEGs bound by AGO identified by AGO HITS-CLIP in miNs overlapped with nELAVL-bound targets in the human brain. Right, top biological GO terms associated with the 54 genes that are predicted to be upregulated in miNs and targeted by both AGO and nELAVLs (left).

(E) A model of miR-9/9*-124 activity in simultaneously repressing non-neuronal targets while promoting the expression of select neuronal targets, such as PTBP2, for the successful acquisition of neuronal identity during the neuronal conversion of HAFs.