Persistent Epigenetic Reprogramming of Sweet Taste by Diet

Anoumid Vaziri[1, 2], Morteza Khabiri[3,†], Brendan T. Genaw[5], Christina E. May[2, 6], Peter L. Freddolino[3, 4],

Monica Dus[1, 2, 5, 6]*

[1] The Molecular, Cellular and Developmental Biology Graduate Program, The University of Michigan,

Ann Arbor, MI 49109

[2] Department of Molecular, Cellular and Developmental Biology, College of Literature, Science, and the

Arts, The University of Michigan, Ann Arbor, MI 49109

[3] Department of Biological Chemistry, The University of Michigan, Ann Arbor, MI 48109, USA

[4] Department of Computational Medicine and Bioinformatics, The University of Michigan, Ann Arbor, MI

48109, USA

[5] Program in Biology, College of Literature, Science, and the Arts, The University of Michigan, Ann

Arbor, MI, 48109, USA

[6] The Neuroscience Graduate Program, The University of Michigan, Ann Arbor, MI 49109

† Current address: Department of Biological Sciences, Quinnipiac University, Hamden, CT 06518

* Correspondence: mdus@umich.edu

Abstract

Diets rich in sugar, salt, and fat alter taste perception and food intake, leading to obesity and metabolic

disorders, but the molecular mechanisms through which this occurs are unknown. Here we show that in

response to a high sugar diet, the epigenetic regulator Polycomb Repressive Complex 2.1 (PRC2.1)

persistently reprograms the sensory neurons of *D. melanogaster* flies to reduce sweet sensation and

promote obesity. In animals fed high sugar, the binding of PRC2.1 to the chromatin of the sweet gustatory

neurons is redistributed to repress a developmental transcriptional network that modulates the

responsiveness of these cells to sweet stimuli, reducing sweet sensation. Importantly, half of these

transcriptional changes persist despite returning the animals to a control diet, causing a permanent decrease

in sweet taste. Our results uncover a new epigenetic mechanism that, in response to the dietary

environment, regulates neural plasticity and feeding behavior to promote obesity.

Introduction

Diets high in processed foods promote higher calorie intake and weight gain, increasing the risk for chronic and metabolic diseases (*1*). How these foods cause overconsumption, however, is still unclear. Processed foods are high in salt, and fat, which we are genetically programmed to like because of their high caloric-density (*2, 3*). Interestingly, evidence is emerging that the levels of salt, sugar, and fat in diets can alter taste sensation in humans (*4–7*), raising the question of whether these sensory changes may influence food intake, obesity, and metabolic disease (*8, 9*). This idea is supported by a number of recent animal studies which found changes in taste, neural responses, and food preferences in rodents fed high nutrient diets (*10–17*). However, due to the complexity of the mammalian taste system and the lack of genetic tools, we know next to nothing about the molecular mechanisms through which diet composition affects taste sensation and obesity. Thus, studies in genetically tractable model organisms could help shed light on this question and help us define evidence-based strategies to curb the prevalence of obesity and metabolic disease, which currently affects billions of people worldwide.

We recently found that high dietary sugar dulls the responses of the *D. melanogaster* taste neurons to sweet stimuli, causing higher food intake and weight gain, arguing that the effects of diet on taste are conserved (*18, 19*). In this manuscript we exploited the exquisite genetics tools of the fly and the relative simplicity of its sensory system to uncover the mechanisms through which high levels of dietary sugar reshape the sensory neurons to promote weight gain and obesity. We report that the Polycomb Repressive Complex 2.1 (PRC2.1), a chromatin silencing complex conserved from plants to humans (*20*), tunes the activity of the sweet sensory neurons and taste sensation in response to the food environment by repressing a transcriptional program that shapes the synaptic, signaling, and metabolic properties of these cells. Interestingly, this diet-dependent transcriptional remodeling persisted even when animals were returned to the control diet, leading to lasting changes in sweet taste sensation that depended on the constitutive activity of PRC2.1. Together our findings suggest that diet composition activates epigenetic mechanisms that reprogram sensory responses to food; this sensory reprogramming determines the perception of future stimuli, leading to long-lasting alterations in behavior that increase the risk for obesity and metabolic disease.

Results

PRC2.1 modulates sweet taste in response to the food environment

Drosophila melanogaster flies fed high dietary sugar experience lower sweet taste sensation as a result of the decreased responsiveness of the sweet sensory neurons to sugar stimuli (*18*). Given the importance of sensory cues to control eating, and recent data that diet also impacts taste in mammals (*10–15, 18*), we set out to identify the molecular mechanisms through which the food environment shapes sensory responses. Since sweet taste deficits develop within 2-3 days upon exposure to the high sugar diet and independently of weight gain (*18*), we reasoned that gene regulatory mechanisms may be involved in modulating the responses of the sensory neurons. To test this hypothesis, we conducted a screen for gene regulatory and epigenetic factors necessary for the sweet taste defects caused by a high sugar diet. To do this, we fed control (*w1118cs*) and mutant flies a control diet (CD, ~5% sucrose) or a diet supplemented with 30% sucrose (sugar diet, SD) for 7 days and then tested their ability to taste using the Proboscis Extension Response (*21*). This behavioral assay measures taste responses by quantifying the amount of proboscis extension (0= no extension, 0.5=partial extension, 1= full extension) when the fly labellum – where the dendrites and cell bodies of the taste neurons are located (Fig. S1A) – is stimulated with three different concentrations of sucrose (30%, 10%, 5%); this generates a taste curve where flies respond more intensely to higher sugar stimuli (Fig. 1B, gray circles). Flies fed a sugar diet show a marked decrease in PER compared to control diet flies (Fig. 1B, gray squares); however, mutants for the core Polycomb Repressive Complex 2 (PRC2) – which includes the histone 3 lysine 27 (H3K27) methyltransferase *Enhancer of Zeste* (*E(z)*), and the obligate accessory factors *Suppressor of zeste 12,* (*Su(z)12*) and *extra sex combs,* (*esc*) (Fig. 1A) – had largely the same proboscis extension response (PER) on a control and sugar diet (Fig. 1B, right, red shades). To confirm the role of PRC2 in taste sensation deficits, we supplemented the control and sugar diet with EED226, a PRC2 inhibitor (herein referred to as EEDi) that destabilizes the core complex by binding to the tri-methyl H3K27 (H3K27me3) binding pocket of EED (the homologue of *esc* in *M. musculus*) (*22*). While animals fed a sugar diet plus vehicle (10% DMSO) experienced lower PER, those fed a SD+EEDi retained normal sweet taste responses (Fig. 1C), consistent with results from the PRC2 mutants. Thus, mutations and inhibition of PRC2 prevents the blunting of sweet taste that occurs in the high sugar food environment.

In flies PRC2 forms two main subcomplexes, PRC2.1 and PRC2.2, which contain distinct

accessory factors that can influence the targeting of the core complex to the chromatin (*23*). Mutations in

the *Polycomb-like* (*Pcl*) gene, the accessory factor to PRC2.1, phenocopied PRC2 mutants and prevented

sweet taste deficits in flies fed a sugar diet (Fig. 1D). In contrast, flies with deficits in the PRC2.2-

members *Jumonji, AT rich interactive domain 2* (*Jarid2*) and *jing* still showed a blunting of sweet taste

responses in flies fed a sugar diet (Fig. S1B). Interestingly, members of the Polycomb Repressive Complex

1 (PRC1) and the recruiter complex PhoRC were also not required for the taste changes in responses to a

sugar diet (Fig. S1C and Fig. S1D-E). Thus, the PRC2.1 complex is necessary for the sensory changes that

occur in the high sugar environment.

We next asked if PRC2.1 is required specifically in the sweet sensory neurons to decrease their

responses to sweet stimuli on the sugar diet. To do this, we used the GAL4/UAS system to knock down

Pcl in the sweet taste neurons using the *Gustatory receptor 5a* GAL4 driver, *Gr5a-GAL4,* which labels

~60 cells in the proboscis of adult flies (*24*); we selected *Pcl* to narrow the effect to the PRC2.1 complex.

Knockdown of *Pcl* in *Gr5a+* neurons using two independent RNAi transgenes (50% knockdown

efficiency, Fig. S2A) prevented sweet taste deficits in animals fed a sugar diet (Fig. 1E, and Fig. S2B). *Pcl*

knockdown, however, had no effect on sweet taste on a control diet (Fig. S2C), in accordance with the

observation that *E(z)* and *Pcl* mutants have no effect on taste on a control diet (Fig. 1B, C) and suggesting

that these phenotypes are uncovered only by the high sugar food environment.

Since *Pcl* is thought to target the PRC2 core complex to chromatin (*23*), we hypothesized that its

overexpression may be sufficient to induce sweet taste deficits even in the absence of a high sugar food

environment. Indeed, overexpression of *Pcl* specifically in the *Gr5a+* neurons induced sweet taste deficits

in flies fed a control diet compared to transgenic controls (Fig. 1F). The effects of *Pcl* overexpression were

abolished by treatment with the PRC2 inhibitor EEDi (Fig. 1G), arguing that *Pcl* causes sweet taste deficits

entirely through the action of PRC2 and not through some yet unidentified mechanism. Importantly, *Pcl*

overexpression had no effect on the number of *Gr5a+* neurons in the proboscis (Fig. S2D), and so the taste

deficits cannot be attributed to a decrease in the number of cells. To exclude the possibility that the effects

of manipulating *Pcl* on sweet taste were developmental, we used the temperature sensitive *tubulin-GAL80ts*

transgene to limit expression of *UAS-Pcl* and *PclRNAi-1* only in adult flies. Switching the flies to the non-

permissive temperature and diet 4 days post eclosion, resulted in the same effects on sweet taste as using

the *Gr5a-GAL4* alone (Fig. S2E). Together, these experiments establish that the PRC2.1 complex is

required cell-autonomously in the *Gr5a+* neurons to mediate the effects of a high sugar diet on sweet

taste.

***Pcl* mutant flies have normal sensory responses and are resistant to diet-induced obesity**

120   Flies on a high sugar diet have lower sweet taste because the neural responses of the taste neurons

to sweet stimuli are decreased (*18*). Since *Pcl* mutants have identical taste on a control and sugar diet, we

hypothesized that the responses of the sensory neurons to sucrose stimulation should also be similar. To

test this, we expressed the genetically encoded presynaptic calcium indicator *UAS-GCaMP6s-Brp-*

mCherry (25) in the sweet sensing neurons and measured their *in vivo* responses to stimulation of the

proboscis with 20% sucrose in *Pcl* mutant animals (Fig. 2A). Indeed, the responses to sucrose stimulation

were identical in *Pcl* mutant flies fed a control diet and sugar diet (Fig. 2B), matching the behavioral data

(Fig. 1); importantly, this rescue was not due to an increase in the number of sweet taste cells (Fig. 2C).

128   We previously showed that restoration of sweet taste neuron activity in flies fed high dietary sugar

protected them from diet-induced obesity (*18*). Since *Pcl* mutants abolished the deficits in neural and

behavioral responses to sweetness in animals fed a high sugar diet, we anticipated that they should also

prevent a diet-dependent increase in triglycerides. Indeed, sugar-diet flies with knockdown of *Pcl* in the

Gr5a+ neurons remained as lean as animals on a control diet (Fig. 2D), while triglycerides increased in

control flies fed a sugar diet (Fig. 2D). Importantly, there was no difference in the levels of triglycerides

between control and *Pcl* knockdown flies fed a control diet (Fig. 2D). Together, these data suggest that, in

response to the food environment, *Pcl* modulates the responsiveness of the sweet gustatory neurons to

promote diet-induced obesity.

***Pcl* chromatin occupancy is redistributed in the high sugar environment**

139   Our experiments show that PRC2.1 plays a critical role in the neural activity, behavior, and the

metabolic state of animals exposed to the high sugar food environment. To identify the molecular

mechanisms underlying these phenotypes, we measured the chromatin occupancy of *Pcl* in the ~60 *Gr5a+*

neurons using Targeted DNA Adenine Methyltransferase Identification (TaDa) (Fig. 3A) (*26-28*). To do

this we generated *Dam::Pcl* (*UAS-LT3-Dam::Pcl*) transgenic flies and compared them to *Dam*-only flies

(*UAS-LT3-Dam*) to control for non-specific methylation by the freely diffusing *Dam* protein (*26*) (Fig. 3A)

and to obtain a measure of chromatin accessibility *in vivo* (CATaDa) (*29*). To specifically profile *Pcl*

binding to chromatin in the sweet sensory neurons and limit the induction of *Dam*, we expressed the

Dam::Pcl and *Dam* transgenes in combination with *Gr5a-GAL4*;*tubulin-GAL80ts*. To induce the

expression of each UAS transgene we shifted the flies to the permissive temperature (28ºC) for 20 hours

after they had been exposed to a control or sugar diet for 3 days (Fig. 3A). We selected this time point

because we previously showed that sweet taste defects developed within 3 days of exposure to the sugar

diet (*18*).

Most of the variation in the biological replicates of *Dam::Pcl* normalized to *Dam* alone (see

Methods) was due to diet (Fig. S3A), consistent with high Pearson correlations within each dietary

condition (Fig. S3B). Further, the chromatin accessibility profile of *Dam* at the *Gr5a* sweet taste receptor

gene promoter was high, while at the *Gr66a* bitter taste receptor promoter –which is only expressed in

bitter cells, closely located near the sweet cells– was low (Fig. 3B), suggesting that the transgenes were

appropriately targeted to the sweet taste neurons and that the limited induction controlled for background

DNA methylation.

We first analyzed *Pcl* chromatin occupancy in the *Gr5a+* neurons of flies on a control diet by

comparing our data to a previous study that annotated five major chromatin types in *D. melanogaster* using

a similar technique (DNA Adenine Methyltransferase Identification, Dam-ID) (*30*). *Pcl* targets were

enriched in Polycomb chromatin (blue), compared to other chromatin types (green and black = repressive;

red and yellow= active) (Fig. 3C); for example, *Pcl* occupancy was high and *Dam* accessibility low at two

known Polycomb blue chromatin clusters, (Fig. S3E), while the opposite was true for regions in other

chromatin types (Fig. S3F, red and yellow) (*30*). We next asked whether *Pcl* was enriched at Polycomb

Response Elements (PREs), cis-regulatory sequences to which Polycomb Group Proteins bind in *D.*

melanogaster (*31,32*). Using a recently developed tool that predicts PRE regions genome-wide (*33*), we

found that *Pcl* was present in these regions (Fig. 3D, gray line), with an enrichment for intergenic (3.2-

fold enriched, $p<0.001$, Monte Carlo permutation test) and enhancer PREs (2.9-fold enriched, $p<0.001$,

Monte Carlo permutation test).

171   To determine changes in *Pcl* occupancy with diet, we compared *Pcl* chromatin binding between

flies fed a control and sugar diet. While ~70% of the overall *Pcl* peaks were shared between the control

(CD) and sugar diet (SD) (Fig. S3C, D), we found more *Pcl* at PREs on a sugar diet compared to a control

diet (Fig. 3D, purple line). Interestingly, chromatin accessibility at both *Pcl* peaks (Fig. 3E) and PREs (Fig.

3F) was decreased in the sugar diet condition. Our analysis also showed that differentially bound *Pcl* peaks

had a 3.3-fold enrichment of overlap for enhancer-type PREs ($p<0.001$, Monte Carlo permutation

test). Further examination of the differentially bound genes, revealed a redistribution in *Pcl* occupancy

(Fig. S3G), with a similar number of genes with higher (group 1) and lower (group 2) *Pcl* binding on a

sugar diet compared to the control diet (Fig. 3G). Using iPAGE for pathway enrichment analysis (*34*) we

found that most of the genes differentially bound by *Pcl* were transcription factors with both promoter and

enhancer binding. Notably, transcription factors implicated in axon target recognition and nervous system

development showed an enrichment in *Pcl* binding on a sugar diet, while those involved in proboscis

development and feeding behavior had both an increase and decrease in *Pcl* occupancy on a sugar diet

(Fig. 3H, for full iPAGE analysis see Fig. S4). While the large majority of genes differentially bound by

Pcl were in the gene regulation category (80%), the pathway enrichment analysis also uncovered a few

metabolism Gene Ontology (GO) terms (Fig. 3H and Fig. S4). In summary, we found that PRC2.1 targeted

transcription factors involved in neuronal processes and development in the *Gr5a+* neurons, and its

chromatin occupancy was redistributed at these loci in the high sugar diet environment. This redistribution

could result in changes in the expression of these transcription factors and their targets, and in turn, affect

the responsiveness of the sensory neurons and taste sensation.

PRC2.1 sculpts the transcriptional responses of the *Gr5a+* neurons in response to diet

193   To test the hypothesis that redistribution of PRC2.1 chromatin occupancy alters the physiology of

the sweet sensing neurons by changing gene expression, we used Translating mRNA Affinity Purification

(TRAP) (*35*) to isolate mRNAs associated with the ribosomes of these cells. (Fig. 4A). To capture the

dynamics of this process, we collected samples from age-matched *Gr5a>Rpl3-3XFLAG* flies fed a sugar

diet for 3 and 7 days (Fig. S5A). We first verified that this technique selected for mRNAs in the *Gr5a+*

neurons alone by quantifying the normalized read counts (*Gr5a+*/input) for three sweet taste receptor

genes *Gr5a*, *Gr64f*, and *Gr64a*, which are expressed in cells labeled by the *Gr5a*-GAL4. Indeed, these

transcripts were enriched in the *Gr5a+* fraction compared to the input (Fig. S5B), while the opposite was

true for the bitter receptor gene *Gr66a*, which is expressed only in the bitter sensing neurons in the taste

sensilla (*36*) (Fig. S5B).

Notably, we observed a large negative skew in gene expression in the *Gr5a+* neurons of flies fed a

sugar diet for 3 (SD3, mint; compared to the control diet) and 7 days (SD7, teal; compared to the control

diet) (Fig. 4B-C, -0.8 and -1.7 skew), with ~68% and 87% of differentially expressed genes (DEG, each

compared to control diet, Wald test, $q< 0.1$) showing negative log$_2$ fold changes (l$_2$fc) respectively (Fig.

4B-C, SD3 and SD7), consistent with the idea that a repressive gene regulatory mechanism is involved in

this process. Overall, we found ~800 transcripts differentially expressed at each time point compared to the

control diet, while ~190 changed at both time points (Fig. 4D, Venn diagram, Wald test, $q< 0.1$). Gene

Ontology (GO) analysis (*34*) revealed that these genes were part of biological pathways involved in 3

broad categories: neural function/signaling, metabolism, and gene regulation (Fig. S6 and Fig. S7). GO

terms for neuron-specific processes, such as dendritic membrane, sensory perception of chemical stimulus,

and presynaptic/vesicle transport, were enriched at both timepoints (Fig. S6 and Fig. S7), suggesting that

PRC2.1 may alter the physiology of the sensory neurons through these pathways in response to a high

sugar environment. Interestingly, flies fed a sugar diet for 7 days also showed changes in GO terms linked

to neurodevelopmental processes, such as asymmetric neuroblast division and neuron projection

morphogenesis, indicating that longer exposure to the diet led to additional alterations in neural function,

which may explain the worsening of sweet taste sensation at the later time point (*18*). GO terms associated

with metabolic changes were present in higher numbers in flies fed a sugar diet for 7 days, consistent with

the findings that longer exposure to the high sugar diet leads to higher fat accumulation (*18*). Finally, we

observed changes in "regulatory" GO terms such as transcription factor and corepressor, consistent with

the redistribution of *Pcl* chromatin occupancy in response to a high sugar diet. Thus, consumption of a

high sugar diet altered neural, regulatory, and metabolic genes in the *Gr5a+* cells.

To determine the role of PRC2.1 in these changes, we performed the transcriptional profiling

experiments in the $Gr5a+$ neurons of Pcl_{c429} mutant animals fed a control diet and sugar diet for 7 days

(CD and SD7) (Fig. S5C). Strikingly, the Pcl mutation abolished the negative skew (Fig. S5D) and largely

nullified the effects of the high sugar diet environment on gene expression. Specifically, of the genes

repressed by a sugar diet (Fig. 4D, heatmap) 32% had a positive $\log_2$fold change (Wald test, $q< 0.1$) and

76% were unchanged ($q<0.1$, practical equivalence test using a null hypothesis of a change of at least 1.5-

fold; see Methods for details) between Pcl mutants fed a control and sugar diet. This effect was reflected in

the GO analysis where terms changed by a high sugar diet in Pcl wild-type animals, such as dendritic

membrane, sensory perception of chemical stimuli, synapse, and carbohydrate metabolic process, showed

opposite trends in $\log_2$ fold changes in Pcl mutants (Fig. S8). Thus, Pcl mutations abolished nearly all the

gene expression changes induced by a high sugar diet consistent with their effects on behavior (PER,

Fig.1), neural function (*in vivo* calcium imaging, Fig. 2), and metabolism (triglycerides, Fig. 2). Together,

these findings support the hypothesis that PRC2.1 tunes taste sensation to the food environment by

influencing the expression of genes involved in the physiology of the sensory neurons.

PRC2.1 represses a transcriptional program required for sweet taste

We discovered that a high sugar diet environment repressed gene expression in the sweet sensory

neurons, and that Pcl mutations almost entirely abolished this effect. This, together with the discovery that

Pcl binding primarily changed at the enhancers of transcription factor genes on a sugar diet, suggests that

Pcl redistribution may affect the expression of transcription factors that, in turn, control genes responsible

for the overall responsiveness of these sensory neurons to sweetness. This idea was supported by the

observation that Pcl-bound genes had lower expression levels than those not bound by it in the $Gr5a+$

neurons (Fig. 4E), with many genes showing high binding and low expression ($\log_2$tpm <2, dark purple),

while others having higher mRNA read counts ($\log_2$tpm >5, light purple) (Fig. S5E). To test this

hypothesis, we looked for transcription factors that were differentially bound by Pcl directly and that

showed changes in gene expression on a sugar diet (Fig. S5F). This analysis revealed 5 transcription

factors: four of these were activators – *GATAe* (Zn finger)*, nubbin/pdm* (*nub*, POU homeobox)*, Ptx1*

(paired-domain homeobox), and *caudal* (*cad*, hox-like homeobox)– which had higher Pcl binding (Fig.

5A) and lower mRNA levels on a sugar diet (Fig. 4F). The fifth transcription factor was the suppressor

scarecrow (*Scro*, NK-like homeobox), which had lower *Pcl* binding (Fig. 5A) and higher mRNAs levels

on a sugar diet (Fig. 4F). Notably, mutations in *Pcl* reversed the effects of a high sugar diet on the

expression of these 5 genes, suggesting that the binding of *Pcl* modulates their mRNA levels (Fig. 4F).

Interestingly, *cad, Ptx1, GATAe,* and *nub* were enriched in the *Gr5a+* neurons compared to the input

(heads), while *Scro* was depleted in these cells (Fig. S9A). To test the effects of these five transcription

factors on sweet taste, we manipulated their expression to mimic the direction of change on a high sugar

diet. Knockdown of *cad, Ptx1, GATAe,* or *nub*, and overexpression of *Scro* in the *Gr5a+* neurons of flies

fed a control diet led to a decrease in sweet taste sensation (Fig. 5B) comparable to that experienced by

animals on a sugar diet (Fig. 1B). Thus, *cad, Ptx1, GATAe,* and *nub* are direct targets of PRC2.1 and

necessary for sweet taste, while overexpression of *Scro* is sufficient to decrease it. However,

overexpression of each activator alone and knockdown of *Scro* was not sufficient to rescue sweet taste in

flies fed a sugar diet (Fig. S9B).

Given that the 4 activators are required for sweet taste sensation, we reasoned that they may

control the expression of genes important for the proper function of the *Gr5a+* neurons and normal sweet

taste. To identify candidate target genes, we tiled the entire *D. melanogaster* genome using the motifs for

each of these 5 transcription factors, converted the hits for each transcription factor to z-scores, and

determined candidate regulatory targets based on estimates of the z-score threshold for binding in each

case (Fig. S9C; see Methods for details). We then flagged as "targets" the set of genes that had a putative

binding site (exceeding our transcription factors-specific z-score cutoff) within a 2 kb region upstream of

the annotated ORF start (Fig. S9D-H), and examined their expression pattern in the *Gr5a+* neurons of flies

on a control and sugar diet. This analysis revealed 658 genes that were collectively regulated by these five

transcription factors. Targets of the transcriptional activators Cad, Ptx1, GATAe and Nub– which *Pcl*

repressed on a sugar diet– showed negative log2fold changes on a sugar diet (Fig. 5C, SD7, teal) that

reverted in the *Pcl* mutants (Fig. 5C, pink). Conversely, targets of the transcriptional repressor Scro –

which was released from *Pcl* binding and had higher mRNA levels on a sugar diet – showed negative

log2fold changes on a sugar diet (Fig. 5C, SD7, teal) that reverted in *Pcl* mutants (Fig. 5C, pink).

Strikingly, the 658 putative targets of these 5 transcription factors accounted for nearly all the genes

changed by a high sugar diet (Fig. 4), suggesting that by directly modulating the expression of *Ptx1, cad,*

GATAe, nub, and *Scro*, PRC2.1 could influence the neural state of the *Gr5a+* neurons via their targets.

When we examined whether these targets were regulated by more than one of the 5 transcription factors

targeted by *Pcl*, we found significant overlap among the regulons of all of the 4 transcriptional activators

with the exception of *Ptx1-nub* (Fisher's exact test, FDR-corrected $p<0.000001$) (Fig. 5D), suggesting that

they could direct the expression of a common set of genes. Further, a portion of *Scro* targets also

overlapped with the regulons of each of the activators (Fisher's exact test, $q<0.000001$), indicating that

Scro could further drive the negative skew in gene expression on a high sugar diet beyond that caused by

the direct binding of PRC2.1 to the activators. Transcription factors that share common targets are often

part of feed-forward loops, where they regulate one another and themselves to ensure stability of gene

expression. Indeed, we found that *GATAe* had binding sites in the promoters of all four regulators

considered here (*cad*, *Scro*, *Ptx1*, *nub*), in addition to binding its own promoter in an auto-regulatory loop

(Fig. 5E). Furthermore, Cad also targeted itself, *nub* was one of Ptx1 targets, and Scro regulated both *cad*

and *GATAe*, forming a negative feedback loop with the latter (Fig. 5D). Thus, the 5 transcription factors

differentially bound by PRC2.1 on a high sugar diet, form a hub that regulates the physiology of the

Gr5a+ neurons. To gain a deeper understanding of which aspects of physiology were changed, we used

pathway enrichment analysis on the regulons for each transcription factor. GATAe targets, which comprise

a large number of the genes regulated by the 4 other transcription factors, were enriched for GO terms

involved in synaptic assembly and growth, terminal bouton, neural projection morphogenesis, and protein

kinase regulation (summarized in Fig. 5D, and Supplementary File 5). In contrast, Ptx1 targets were

enriched in GO terms implicated in cyclic AMP signaling, detection of chemical stimuli, and

morphogenesis (summarized in Fig. 5D, and Supplementary File 5), Cad targets showed enrichments in

adenylate cyclase activity, sensory perception, and neuropeptide signaling (summarized in Fig. 5D, and

Supplementary File 5), and Nub targets in calcium signaling and nucleosome (summarized in Fig. 5D, and

Supplementary File 5). The targets of the repressor Scro showed enrichment in both neural and metabolic

GO terms such as olfactory behavior and carbohydrate metabolic process (summarized in Fig. 5D, and

Supplementary File 5).

To test the possibility that these targets form a functional network, we used STRING (*37*) and

found a significant number of interactions above the expected number (protein-protein interaction

enrichment of $p < 1.0e-16$) suggesting that the targets are indeed part of a functional and biologically

connected network in the *Gr5a+* neurons. We then used a subset of the neural targets to build a second

network to identify candidate target genes that may play a direct role in neural physiology and sweet taste.

This network showed strong interactions between genes involved in synaptic organization and signal

transduction and their connection with the upstream regulators (Fig. S14A). We chose two genes at the

edges of the network, which are less likely to have redundant functions, the *Adenylyl Cyclase X D* (*ACXD*)

gene (*38*) and the *Activity Regulated Cytoskeleton Associated Protein 1* (*Arc1*) (*39*), which are involved in

the sensing of stimuli and in synaptic plasticity, respectively. Knockdown and mutations of *Arc1* or *ACXD*

in the *Gr5a+* cells of flies on a control diet led to a significant decrease in sweet taste responses compared

to the transgenic controls (Fig. S14B-D). Together, these lines of evidence suggest that PRC2.1 mediates

the effects of a high sugar diet on sweet taste by directly controlling the expression of a transcription

program required for sweet taste.

The persistent phenotypic memory of the food environment is dependent on PRC2.1

The cellular fates created by Polycomb Group proteins are inherited as stable memories across cell

divisions to ensure phenotypic stability even in the absence of the triggering stimuli (*40-44*). We therefore

asked if the neural and behavioral state created by PRC2.1 in the high sugar diet environment was

maintained when files were moved to the control diet after eating a sugar diet for 7 days (SD>CD) (Fig.

6A). We found that these animals had lower sweet taste, similar to that of age-matched flies fed a sugar

diet for 7 days (Fig.6B, SD>CD compared to CD>SD). However, their triglyceride levels were similar to

those of control diet flies (Fig. S14E), suggesting that while fat storage was reversible when flies only had

access to the "healthy" control diet, sweet taste deficits persisted.

To understand how this phenotype compares to that of the control diet flies at the molecular level,

we conducted TRAP of the *Gr5a+* neurons of flies in the SD>CD and CD>CD conditions. mRNAs from

flies on a SD>CD showed a strong negative skew in $\log_2$fold changes compared to the control diet group

(Fig. 6C, -2.02), reminiscent of the skew we observed in flies fed a sugar diet (Fig. 4C). Furthermore, we

observed that 47% (310/658) of genes in the transcriptional network repressed by PRC2.1 on a sugar diet

were still decreased in SD>CD flies (Fig. 6D). Interestingly, the SD>CD animals clustered with the sugar

diet 7 (SD7) group compared to sugar diet 3 (SD3) and *Pcl* mutants fed a sugar diet (Fig. 6D). Thus, at the

molecular level, half of the neural state established by dietary sugar via PRC2.1 persisted. To test the

hypothesis that PRC2.1 plays an active role in the maintenance of this neural state in the absence of the

sugar diet, we inhibited PRC2 activity during the "recovery" diet using an EEDi inhibitor

(SD>CD+EEDi). Remarkably, these animals showed a restoration of wild-type sweet taste (Fig. 6E, green

triangles) compared to flies fed the recovery diet without EEDi supplementation (Fig. 6B, SD>CD).

Together, these data indicate that the sensory neurons retain a transcriptional and phenotypic memory of

the sugar diet environment that leads to long lasting behavioral deficits, and that PRC2.1 is constitutively

required for the persistence of this state.

Discussion

In this study we set out to understand how diet composition alters the gustatory system to promote

food intake and weight gain. Specifically, we took advantage of the simple sensory system of *D.*

melanogaster and its exquisite genetic and neural tools to identify the molecular mechanisms through

which diet composition changes neural state, physiology, and behavior. Here we show that the decrease in

sweet taste sensation that flies experience after chronic exposure to a high sugar diet is caused by the cell-

autonomous action of the Polycomb Repressive Complex 2.1 in the sweet gustatory neurons. Mutations

and pharmacological inhibition of PRC2.1 blocked the effects of the food environment on neural activity,

behavior, and obesity. While we do not exclude the possibility that PRC1 and PhoRC may also be

involved, we found that mutations or knockdown in these complexes had no effect on taste. In the high

sugar food environment, PRC2.1 chromatin occupancy was redistributed, leading to the repression of

transcription factors, neural, signaling, and metabolic genes that decreased the responsiveness of the

Gr5a+ neurons and the fly's sensory experience. However, we discovered that PRC2.1 did not directly

bind to neuronal genes in these cells and that, instead, it targeted transcription factors involved in sensory

neuron development, synaptic function, and axon targeting. Specifically, on a high sugar diet *Pcl* binding

was increased at the *cad*, *GATAe*, *nub/pdm*, *Ptx1* loci and decreased at the *Scro* locus, with corresponding

changes in the mRNA levels of these genes (Fig. 6F, model). Together, the decrease in the mRNA levels

of the 4 activators (*cad*, *GATAe*, *nub/pdm*, *Ptx1*) and the increase in the repressor (*Scro*), led to the

repression of genes implicated in synaptic function, signal transduction, and metabolism. Interestingly, we

uncovered several feed-forward regulatory loops among the transcriptional regulators, suggesting that this

hub could ensure the stability of the neural state unless the change in environment is chronic. This

transcriptional network controlled by these regulators was necessary for sweet taste, because knockdown

of the four activators and a few of their targets, as well as overexpression of *Scro* (Fig. 5B and Fig. S10B-

D), resulted in a decrease in sweet taste on a control diet.

Interestingly, several of the transcription factors we identified – *Ptx1*, *Scro*, and *nub/pdm* – have

been shown to control the proper branching, synaptic connectivity, and function of sensory neurons (*45–*

51), while others (*cad*, *nub/pdm*) play a role in neuroblast development (*52, 53*); PRC2 also functions as a

competence factor in neural proliferation, differentiation and sensory neurons (*45, 52, 54*). Importantly, we

found that the 4 activators that are repressed by *Pcl* in the high sugar condition are enriched in the *Gr5a+*

cells on a control diet, while *Scro* is depleted. Thus, the transcriptional network of ~658 genes controlled

by this developmental transcription hub is likely to define the intrinsic properties of the sweet sensing

neurons. Indeed, we observed that many of the target genes involved in signaling, synaptic function, and

cell adhesion such as the kinase *haspin*, the adenylate cyclase *ACXD*, *sytalpha*, *Arc1*, tetraspanin, jonan,

and innexin proteins among others, were part of highly interconnected network, which we speculate, could

affect the sweet gustatory neurons at both the functional and connectivity levels. Since we did not observe

a change in the expression of the sweet taste receptors, or the misexpression of other taste receptors (*36*),

our data are not consistent with a complete loss of identity of the *Gr5a+* neurons. Instead, we propose that

PRC2.1 tunes the sweet sensory neurons to the dietary environment by altering a transcriptional network

that controls the intrinsic properties of these cells, especially those involved in signal transduction,

connectivity, synaptic function, and metabolism. Studies that test the effects of this network on the

connectivity, morphology, and signal transduction of the sweet sensory neurons will shed light on how

exactly the transcriptional remodeling caused by PRC2.1 impacts the *Gr5a+* cells. The question of how

exactly PRC2.1 senses the changes in the food environment also remains open. Recent studies suggest that

the activity of Polycomb Group proteins is directly and indirectly linked to cellular metabolism, including

kinase signaling cascades, GlcNAcylation, and the availability of cofactors for histone modifications (*55–*

59). Our experiments show that PRC2.1 chromatin occupancy changes depending on the dietary

environment, and provides a new model to tackle this question. Finally, dysregulation of Polycomb-

associated chromatin has been found in mice and humans with diet-induced obesity (*60–62*), suggesting

that the mechanisms we discovered here could also underlie the chemosensory alterations reported in

mammals.

More broadly our work opens up the exciting possibility that PRC2 may modulate neural plasticity

in response to environmental conditions. Despite its central role in development and maintenance of neural

identity, studies have not directly linked PRC2.1 to neural plasticity. However, in other post-mitotic cells

such as muscle, Polycomb Group Proteins are known to reshape transcriptional programs according to

environmental stressors, such as oxidative stress, injury, temperature, and light (*55, 56*). Our findings

advance the conceptual understanding of the role of Polycomb Group Proteins in the nervous system and

argue that PRC2 could regulate "neural states" and metaplasticity in response to environment stimuli

beyond diet. Using neuroepigenetic mechanisms like those employed by Polycomb Group Proteins to tune

neural states to external conditions could provide several advantages compared to the medley of other

cellular, receptor, or synaptic plasticity based mechanisms. Specifically, it would allow cells to 1)

orchestrate a coordinated response, 2) create a memory of the environment, and 3) buffer small

fluctuations until a substantial challenge is perceived. It is particularly fascinating to think about the

molecular mechanisms through which these neural states may be established. The need of neurons to

constantly maintain their identity may mean that environmental signals like the extent of sensory

stimulation could alter the expression of developmental gene batteries and affect neural states (*63*). Indeed,

it has been speculated that some forms of plasticity may re-engage developmental programs that specify

the intrinsic properties of neurons (*64, 65*). Here we observed that the regulators of the transcriptional

network we uncovered function in sensory neuron development and are enriched in the *Gr5a+* cells. Thus,

it could be a hallmark of neuroepigenetic plasticity to exploit developmental programs, linking the known

role of PRC2 in establishing cell fates with this newly discovered function in modulating cell states.

Incidentally, engaging developmental programs could be the reason why some environments and

experiences leave a memory that leads to the persistent expression of the phenotype beyond the presence

of the triggering stimulus, as these could target neural connectivity and set synaptic weights thresholds. Defining the circuit-specific changes of the pioneering studies on maternal care, addiction, and learning and memory (*66–69*) would put this hypothesis to test. Here we found that the changes in taste sensation and half of the sugar diet neural state set by PRC2.1 remained even after animals were moved back to the control diet for a week. A limitation of our study is that due to the small number of *Gr5a+* neurons and their anatomically inaccessible location, we were not able to measure the identity of the molecular memory in these cells alone. However, we saw that the phenotypic memory of the high sugar food environment was dependent on the constitutive action of PRC2.1. Based on other studies showing that the H3K27 methyl mark acts as a molecular memory during development (*40, 41*), we speculate that this is likely to be the molecular signal in the *Gr5a+* cells too.

In conclusion, we show that PRC2.1 mediates the effects of dietary sugar on sweet taste by establishing persistent alterations in the taste neurons that remain as a phenotypic and neural memory of the previous food environment. We speculate that this memory may lock animals into patterns of feeding behavior that become maladaptive and promote obesity. Thus, diet can induce lasting molecular alterations that restrict the behavioral plasticity of animals and impact disease risk. Since the content of sugar in processed foods is similar to that we fed flies and the function of Polycomb Group Proteins is conserved from plants to humans (*20*), our work is broadly relevant to understanding the effects of processed food on the mammalian taste system and its impact on food intake and a whole range of diet-related conditions and diseases that affect billions of people around the globe.

Materials and Methods

Fly Husbandry and Strains:

All flies were grown and maintained on cornmeal food (Bloomington Food B recipe) at 25°C and 45%–55% humidity under a 12:12 hour light-dark cycle (ZT0 at 9 AM). Male flies were collected under CO2 anesthesia 1-3 days after eclosion and maintained in a vial that housed 35-40 flies. Flies were acclimated to their new vial environment for an additional 2 days. For all experiments, flies were changed to fresh food vials every other day.

For all dietary manipulations, the following compounds were mixed into standard cornmeal food (Bloomington Food B recipe) (0.58 calories per gram) by melting, mixing, and pouring new vials as in (*70*) and (*71*). For the 30% sugar diet (1.41 calories per gram) Domino granulated sugar w/v was added. For the EED226 inhibitor diet (AxonMedchem), EED226 was solubilized in 10% DMSO and added to control or 30% sugar diet at a total concentration of 8 uM.

For genetic manipulations the GAL4/UAS system was used to express transgenes of interest in *Gustatory receptor 5a Gr5a-GAL4*. For each GAL4/UAS cross, transgenic controls were made by crossing the *w1118CS* (gift from A. Simon) to GAL4 or UAS flies, sex-matched to those used in the GAL4/UAS cross. The following fly lines are used in this paper: *Gr5a-GAL4/Cyo* (gift from K. Scott, University of California, Berkeley), *nsyb-GAL4/Cyo* (gift from J. Simpson, University of California, Santa Barbara), *Gr64f-GAL4/Cyo* (gift from H. Amrein, University of Texas, A&M), *UAS-GCaMP6s-Brp-mCherry* (BDSC #77131), *Tubulin-GAL80ts* (BDSC #7018), *UAS-Ogt-RNAi* and *UAS-Ogt-GFP/Cyo* (gift from C. Lehner, University of Zurich), *UAS-Pcl* (FlyORF #F001897), *UAS-Pcl-RNAi* (VDRC #v220046, BDSC #33945), *UAS-Rpl3-3XFLAG* (gift from D. Dickman, University of Southern California), *PclT1/CKG* (gift from G. Cavalli, Université de Montpellier), *Pclc241/Cyo*, *E(z)c249/TM3*, *Su(z)12c253*, *Escc289* (gift from N. Liu, Chinese Academy of Sciences), *Pc1/TM1* (BDSC #1728), *ph-d 401ph-p602/FM7* (BDSC #5444), *Psch27/Cyo* (BDSC #5547), *UAS-LT3-Dam* (gift from A.H Brand, University of Cambridge), *UAS-LT3-Dam::PCL* (made in this study), *UAS-Arc1-RNAi* (BDSC #25954), *Arcesm13* (BDSC #37530), *Arc1esm18* (BDSC #37531), *UAS-Arc1* (BDSC #37532), *UAS-Ptx1-RNAi* (VDRC #107785), *UAS-cad-RNAi* (BDSC #34702), *UAS-GATAe-RNAi* (BDSC #33748), *UAS-nub* (FlyORF #F000147), *UAS-Scro-RNAi*

(BDSC #29387), *UAS-Ptx1* (FlyORF #F003469), *UAS-Scro* (FlyORF F003427), *UAS-nub-RNAi* (BDSC

#55305), *UAS-cad* (FlyORF #F000471) *UAS-Jarid2-RNAi* (BDSC #32891, #26184), *UAS-ACXD-RNAi*

(BDSC #62871, #35589), *UAS-jing-RNAi* (BDSC #27084), *UAS-Sfmbt-RNAi* (BDSC #32473).

Proboscis Extension Response:

Male flies were food deprived for 18-24 hours in a vial with a Kimwipe dampened with 2 mL of milliQ-

filtered deionized (milliQ DI) water. Proboscis extension response (PER) was carried out as described in

(*21*).

Proboscis Immunofluorescence:

Probosces were dissected in 1xPBS and fixed in 4% PFA, mounted in FocusClear (CelExplorer) on

coverslips. Cell bodies were imaged using a FV1200 Olympus confocal with a 20x objective. Cells were

counted using Imaris Image analysis software.

Triglyceride Measurements:

Total triglycerides normalized to total protein were measured as described in *(72)*. Briefly, two flies per

biological replicate were homogenized in lysis buffer (140 mM NaCl, 50 mM Tris-HCl pH 7.4, 0.1%

Triton-X) containing protease inhibitor (Thermo Scientific). Lysate extract was used to determine protein

and triglyceride concentrations using Pierce BCA assay (Thermo Scientific, abs 562 nm) and Triglycerides

LiquiColor test (Stanbio, abs 500 nm), respectively.

Calcium Imaging:

Male flies expressing *GCaMP6s-Brp-mCherry* (*25*) in the sweet sensing neurons were food deprived for

18-24 hours. The flies for live imaging were prepared similar to LeDue et al. (*73*). Briefly, flies were fixed

to a custom-printed plastic slide with paraffin wax and the proboscis waxed to an extended position. Distal

leg segments were removed to prevent tarsal interference with labellar stimulation. To image the SEZ, a

sugarless artificial hemolymph solution filled the well surrounding the head. Subsequently, the dorsal

cuticle between the eyes was removed by microdissection to expose the brain. Each fly proboscis was

tested with milliQ water before stimulating with 20% sucrose dissolved in milliQ water. Stimulus (a piece

of Kimwipe soaked in tastant and held with forceps) delivery to the proboscis was manual and timed to

coincide with the 100th recording sample of each time series. Imaging was carried out using an upright

confocal microscope (Olympus, FluoView 1200 BX61WI), a 20x water-immersion objective and laser

excitation at 488 and 543 nm. Recordings were made at 4 Hz (512 x 512 pixels). Plane of interest was kept

to the most ventral neuropil regions innervated by the sweet sensing neurons.

RNA Extraction and Quantitative RT-PCR:

For all RNA extractions used for qPCR, fly heads from 10-20 flies were dissected into Trizol (Ambion)

and homogenized with plastic pestles. RNA was extracted by phenol chloroform (Ambion), and

precipitated by isopropanol with Glycoblue Coprecipitant (Invitrogen). RNA pellet was washed as needed

with 75% ethanol and subsequently eluted in nuclease free water and treated with DNase I, according to

manufacturer's instructions (Turbo DNA-free DNA removal kit, Ambion). All steps were carried out in

RNase free conditions, and RNA was stored at -80C until further processing.

Complementary DNA was synthesized by Superscript III (Invitrogen) reverse transcriptase with the

addition of Ribolock RNase inhibitor (Thermo Fisher Scientific). qPCR reactions were carried out using

Power SYBR Green PCR master mix (Applied Biosystems) based on manufacturer's instructions. Primers

were added at a 2.5 uM concentration. All reactions were run on a 96-well plate on the StepOnePLus Real-

Time PCR System (Applied Biosystems) and quantifications were made relative to the reference gene

Ribosomal protein 49 (Rp49). Primer sequences are listed below and were tested for efficiency prior to the

qPCR experiment:

Gene	Sequence
Rp49 Forward	ATGCTAAGCTGTCGCACAAA
Rp49 Reverse	ACTTCTTGAATCCGGTGGGC

Pcl Forward	ACCCATGAAAATGCAACAGCC
Pcl Reverse	TTATCCGCATGGTAGGGCAA
Pcl_Gibson_Forward	ctctgaagaggatctgGCCGGCGCaATGATGAACAACCATTTTCACTTGCAAC
Pcl_Gibson_Reverse	agtaaggttccttcacaaagatcctTCAATTTCCAAGCAATCCAATCGCC

Affinity purification of ribosome associated mRNA (TRAP):

300 heads (10,000 *Gr5a+* cells) per biological replicate were collected using pre-chilled sieves in liquid

nitrogen on dry ice. Frozen heads were then lysed as previously described (*35*). From the lysate total RNA

was extracted by TRIzol LS Reagent (ThermoFisher scientific, 10296010) as input. The remainder of the

lysate was incubated with Dynabeads protein G (ThermoFisher scientific, 10004D) to preclear samples for

2 hours and subsequently incubated with Dynabeads protein G coated with an anti-Flag antibody (Sigma,

F1804). The lysate-beads mixture was incubated at 4°C with rotation for 2 hours, then. RNA was extracted

from ribosomes bound to the beads by TRIzol Reagent (*35*).

Targeted DNA Adenine Methyltransferase Identification (TaDa):

For the *UAS-LT3-Dam::Pcl* construct, the coding region of the *Pcl* gene was amplified from the *pCRE-*

NDAM-Myc-DO69-Pcl (gift from Bas Van Steensel, The Netherlands Cancer Institute) with primers listed

above and assembled into the *UAS-LT3-DAM* plasmid (gift from Andrea Brand, University of Cambridge)

using the NEBuilder® HiFi DNA Assembly kit based on manufacturer's instructions (NEB). Transgenic

animals were validated by RT-PCR for correct insert. These lines were crossed to *Gr5a-GAL4;tubulin-*

GAL80ts. All animals were raised and maintained at 20 °C. Expression of *Dam::Pcl* and *Dam* were induced

at 28 °C for 18-20 hours. For all experiments 300 heads of males and female flies were collected per

replicate on dry ice by sieving. DNA was extracted from frozen heads following kit instructions

(Invitrogen). For identification of methylated regions purified DNA was digested by DpnI followed by

PCR purification of digested sequences. TaDa adaptors were ligated by T4 DNA ligase (NEB). Adapter

ligated DNA was PCR amplified according to protocol (*28*), and subsequently purified. Purified DNA was

540 digested with DpnII followed by sonication to yield fragments averaging 200-300bp. TaDa adaptors were

541 removed from sonicated DNA by digestion. Sonicated DNA is used for library preparation (*28*).

Library Preparation for TRAP and TaDa:

544 RNA sequencing libraries were generated using the Ovation SoLo RNA-Seq System for *Drosophila*

545 (Nugen, 0502-96). All reactions included integrated HL-dsDNase treatment (ArcticZymes, Cat. #70800-

546 201). DNA sequencing libraries were generated using the Takara ThruPlex kit (cat #022818) using 3ng

547 input and 10 cycles of PCR. All libraries were sequenced on the Illumina NextSeq platform (High-output

548 kit v2 75 cycles) at the University of Michigan core facility.

High Throughput RNA-seq Analysis

Fastq files were assessed for quality using FastQC (*74*). Reads with a quality score below 30 were

discarded. Sequencing reads were aligned by STAR (*75*) to dmel-all-chromosomes of the dm6 genome

downloaded from Ensemble, and gene counts were obtained by HTseq (*76*). Count files were used as input

to call differential RNA abundance by DESeq2 (*77*). All pairwise comparisons were made to the control

diet of the corresponding genotypes, such that sugar diet three days and sugar diet seven days were

compared to the age matched control diet group. In *Pcl* mutant experiments, the pairwise comparison was

made between sugar diet and control diet within the age-matched *Pcl$_{c429}$* genotype group. A cutoff of

*q*val<0.1 was used to call differentially expressed genes. Skew in log$_2$fold changes was measured using the

R package Skewness (e1071). RNAseq data visualization was carried out in R studio using ggplot2 and the

following packages, pheatmap (*78*), Venneuler (*79*), and EnhancedVolcano (*80*). To cluster columns and

rows in pheatmap "Ward.D " clustering was used.

High Throughput TaDa and CATaDa Analysis

Fastq files were assessed for quality using FastQC (*74*). Reads with a quality score below 30 were

discarded. The damidseq_pipeline was used to align, extend, and generate log2 ratio files (*Dam::Pcl/Dam*)

in GATC resolution as described previously (*81*). In short, the pipeline uses Bowtie2 (*82*) to align reads to

dmel-all-chromosomes of the dm6 genome downloaded from Ensemble, followed by read extension to 300

bp (or to the closest GATC, whichever is first). Bam output is used to generate the ratio file in bedgraph

format. Bedgraph files were converted to bigwig and visualized in the UCSD genome browser. Correlation

coefficients and PCA plot between biological replicates were computed by multibigwigSummary and

plotCorrelation in deepTools (*83*). Fold Change traces for SD/CD of $\log_2$(*Dam::Pcl/Dam*) were generated

by calculating the mean coverage profile of all replicates for each condition and subsequently calculating

fold change between the sugar diet and control diet condition with deepTools bigwigCompare (*83*). Peaks

were identified from $\log_2$(*Dam::Pcl/Dam*) ratio files using find_peaks (FDR<0.01) (*81*). To do this, the

binding intensity thresholds are identified from the dataset, the dataset is then shuffled randomly, and the

frequency of consecutive regions (i.e. GATC fragments or bins) with a score greater than the threshold is

calculated. The FDR is the observed / expected for a number of consecutive fragments above a given

threshold. Association of genes to peaks was made using the peaks2genes script (*81*) and dm6 genome

annotations. Overlapping intervals or nearby intervals were merged into a single interval using MergeBED

in Bedtools (*84*). Intervals common in all replicate peak files were identified by Multiple Intersect in

Bedtools (*84*). DiffBind was used to determine differentially bound sites on peak files based on differences

in read intensities (*85*).

For CATaDa experiments, all analyses were performed similar to those of TaDa with the exceptions that

1) *Dam* only profiles were not normalized as ratios but shown as non-normalized binding profiles, 2) *Dam*

only coverage plots were generated by converting bam files to bigwig files normalized to 1x dm6 genome,

and 3) peaks were called using MACS2 call peaks on alignment files without building the shifting model

with an of FDR<0.05 (*86*).

To determine the proportion of genes that fit within the various chromatin domain subtypes, we first

matched *Dam::Pcl/Dam* targets to coordinates identified by Filion et al and then determined their gene

count in each chromatin subtype (observed) compared to the whole genome (expected).

iPAGE Analysis

All Gene Ontology (GO) term enrichment analysis was performed using the iPAGE package (*34*), using

gene-GO term associations extracted from the Flybase dmel 6.08 2015_05 release (*87*). For analysis of

TRAP data, iPAGE was run in continuous mode, with $\log_2$fold changes divided into seven equally

populated bins. For analysis of TaDa and TF regulatory targets, iPAGE was run in discrete mode, using the

groups specified for each calculation. For all discrete calculations, independence filtering was deactivated

due to the less informative available signal. All other iPAGE settings were left at their default values. All

shown GO terms pass the significance tests for overall information described in (*34*); in addition, for each

term, individual bins showing especially strong contributions (*p*-value such that a Benjamini-Hochberg

FDR (*88*) calculated across that row, yields $q<0.05$) are highlighted with a strong black box.

PREdictor

Identification of predicted PRE sites was performed exactly as described in (*33*), using the dm6 genome.

As suggested in (*33*), we use a threshold confidence score of 0.8 to identify the PREs used in the present

analysis. A complete list of predicted PREs, with accompanying confidence scores, is shown in

Supplementary File 1. Enrichments of overlap between different PRE classes and *Pcl* occupancy locations

were calculated by comparing the observed overlap frequency with the overlaps for 1,000 random

shuffling of the binding/differential binding peak locations (calculated using bedtools 2.17.0 (*84*)).

Calculation of regulatory targets of transcription factors

To identify likely targets of each transcription factor of interest, we drew upon the transcription factor

binding site calculations described in (*33*), in which the motif of each transcription factor was scanned

along every base pair of the *D. melanogaster* dm6 genome using FIMO (*89*), and the base pair-wise

binding results converted to robust z-scores. For each TF, we then considered its regulon to consist of all

genes with at least one binding site with z-score above a TF-specific threshold within 2 kb upstream of the

beginning of the gene. We identified TF-specific thresholds by manual inspection of a plot of the average

expression changes between conditions vs. threshold, aiming to identify a point of maximum information

content relative to noise (see Supplemental Figure 9 for the plots used to identify TF-specific z-score

thresholds). Once the set of targets (regulon) for each factor was identified, we tested for significant

enrichment or depletion of overlaps between the regulons using Fisher's exact test, reporting Benjamini-

Hochberg false discovery rates (FDRs) (*88*). All calculated odds ratios were positive, indicating enriched

overlaps between the regulons.

STRING network analysis

To develop a functional network between the candidate neural targets of cad, Ptx1, nub, GATAe and Scro, we uploaded genes categorized into neural/signaling GO terms based on DAVID functional annotations (*90, 91*) to the STRING database search (*37*). Genes were clustered by their reported protein-protein interactions and corresponding confidence scores (*37*) and plotted in Cytoscape (v3.7.1) (*92*). In this network edges do not represent direct protein-protein interaction but rather represent a functional interaction. For network see Supplementary File 6.

Data Analysis and Statistics

Statistical tests, sample size, and *p* or *q* values are listed in each figure legend. Data were evaluated for normality and appropriate statistical tests applied if data were not normally distributed, all the tests, biological samples, and the *p* and *q* values are listed in the figure legends and specific analysis under each methods session. Because the inferential value of a failure to reject the null hypothesis in frequentist statistical approaches is limited, for all RNA-seq expression datasets, we coupled our standard differential expression with a test for whether each gene could be flagged as 'significantly not different'. Defining a region of practical equivalence (ROPE) as a change of no more than 1.5-fold in either direction, we tested the null hypothesis of at least a 1.5-fold change for each gene, using the gene-wise estimates of the standard error in $\log_2$fold change (reported by Deseq2) and the assumption that the actual $\log_2$fold changes are normally distributed. Rejection of the null hypothesis in this test is taken as positive evidence that the gene's expression is not changed substantially between the conditions of interest. Python code for the practical equivalence test can be found on Github as calc_sig_unchanged.py. All data in the figures are shown as Mean ± SEM, **** $p < 0.0001$, *** $p < 0.001$, ** $p < 0.01$, *$p < 0.05$ unless otherwise indicated.

References

1. K. D. Hall, A. Ayuketah, R. Brychta, H. Cai, T. Cassimatis, K. Y. Chen, S. T. Chung, E. Costa, A. Courville, V. Darcey, L. A. Fletcher, C. G. Forde, A. M. Gharib, J. Guo, R. Howard, P. V. Joseph, S. McGehee, R. Ouwerkerk, K. Raisinger, I. Rozga, M. Stagliano, M. Walter, P. J. Walter, S. Yang, M. Zhou, Ultra-Processed Diets Cause Excess Calorie Intake and Weight Gain: An Inpatient Randomized Controlled Trial of Ad Libitum Food Intake. Cell Metabolism. 30 (2019), pp. 67–77.e3.

2. G. K. Beauchamp, P. Jiang, Comparative biology of taste: Insights into mechanism and function. Flavour. 4, 9 (2015).

3. K. C. Berridge, M. L. Kringelbach, Pleasure systems in the brain. Neuron. 86, 646–664 (2015).

4. M. Bertino, G. K. Beauchamp, K. Engelman, Long-term reduction in dietary sodium alters the taste of salt. Am. J. Clin. Nutr. 36, 1134–1144 (1982).

5. G. K. Beauchamp, M. Bertino, D. Burke, K. Engelman, Experimental sodium depletion and salt taste in normal human volunteers. Am. J. Clin. Nutr. 51, 881–889 (1990).

6. J. E. Stewart, R. S. J. Keast, Recent fat intake modulates fat taste sensitivity in lean and overweight subjects. Int. J. Obes. 36, 834–842 (2012).

7. P. M. Wise, L. Nattress, L. J. Flammer, G. K. Beauchamp, Reduced dietary intake of simple sugars alters perceived sweet taste intensity but not perceived pleasantness. Am. J. Clin. Nutr. 103, 50–60 (2016).

8. D. Liu, N. Archer, K. Duesing, G. Hannan, R. Keast, Mechanism of fat taste perception: Association with diet and obesity. Prog. Lipid Res. 63, 41–49 (2016).

9. A. Wittekind, K. Higgins, L. McGale, C. Schwartz, N. S. Stamataki, G. K. Beauchamp, A. Bonnema, P. Dussort, S. Gibson, C. de Graaf, J. C. G. Halford, C. F. M. Marsaux, R. D. Mattes, J. McLaughlin, D. J. Mela, S. Nicklaus, P. J. Rogers, I. A. Macdonald, A workshop on "Dietary Sweetness—Is It an Issue?" Int. J. Obes. 42, 934 (2018).

10. A. Kaufman, E. Choo, A. Koh, R. Dando, Inflammation arising from obesity reduces taste bud abundance and inhibits renewal. PLoS Biol. 16, e2001959 (2018).

11. M. S. Weiss, A. Hajnal, K. Czaja, P. M. Di Lorenzo, Taste Responses in the Nucleus of the Solitary Tract of Awake Obese Rats Are Blunted Compared With Those in Lean Rats. Front. Integr. Neurosci. 13, 35 (2019).

12. A. B. Maliphol, D. J. Garth, K. F. Medler, Diet-induced obesity reduces the responsiveness of the peripheral taste receptor cells. PLoS One. 8, e79403 (2013).

13. Z. C. Ahart, L. E. Martin, B. R. Kemp, D. D. Banik, S. G. E. Roberts, A. M. Torregrossa, K. Medler, Differential effects of diet and weight on taste responses in diet-induced obese mice. bioRxiv (2019), p. 564211.

14. K. Chen, J. Yan, Y. Suo, J. Li, Q. Wang, B. Lv, Nutritional status alters saccharin intake and sweet receptor mRNA expression in rat taste buds. Brain Res. 1325, 53–62 (2010).

15. L. P. McCluskey, L. He, G. Dong, R. Harris, Chronic exposure to liquid sucrose and dry sucrose diet have differential effects on peripheral taste responses in female rats. Appetite. 145, 104499 (2020).

16. J. M. Crow, Obesity: insensitive issue. Nature. 486, S12–3 (2012).

17. F. Sartor, L. F. Donaldson, D. A. Markland, H. Loveday, M. J. Jackson, H.-P. Kubis, Taste perception and implicit attitude toward sweet related to body mass index and soft drink supplementation. Appetite. 57, 237–246 (2011).

18.   C. E. May, A. Vaziri, Y. Q. Lin, O. Grushko, M. Khabiri, Q.-P. Wang, K. J. Holme, S. D. Pletcher,
P. L. Freddolino, G. G. Neely, M. Dus, High Dietary Sugar Reshapes Sweet Taste to Promote
Feeding Behavior in Drosophila melanogaster. Cell Rep. 27, 1675–1685.e7 (2019).

19.   C. E. May, J. Rosander, J. Gottfried, E. Dennis, M. Dus, Dietary sugar inhibits satiation by
decreasing the central processing of sweet taste. bioRxiv (2019) (available at
https://www.biorxiv.org/content/10.1101/2019.12.16.877613v1.abstract).

20.   B. Schuettengruber, H.-M. Bourbon, L. Di Croce, G. Cavalli, Genome Regulation by Polycomb
and Trithorax: 70 Years and Counting. Cell. 171, 34–57 (2017).

21.   T. Shiraiwa, J. R. Carlson, Proboscis extension response (PER) assay in Drosophila. J. Vis. Exp.,
193 (2007).

22.   W. Qi, K. Zhao, J. Gu, Y. Huang, Y. Wang, H. Zhang, M. Zhang, J. Zhang, Z. Yu, L. Li, L. Teng,
S. Chuai, C. Zhang, M. Zhao, H. Chan, Z. Chen, D. Fang, Q. Fei, L. Feng, L. Feng, Y. Gao, H. Ge,
X. Ge, G. Li, A. Lingel, Y. Lin, Y. Liu, F. Luo, M. Shi, L. Wang, Z. Wang, Y. Yu, J. Zeng, C.
Zeng, L. Zhang, Q. Zhang, S. Zhou, C. Oyang, P. Atadja, E. Li, An allosteric PRC2 inhibitor
targeting the H3K27me3 binding pocket of EED. Nat. Chem. Biol. 13, 381–388 (2017).

23.   A. Laugesen, J. W. Højfeldt, K. Helin, Molecular Mechanisms Directing PRC2 Recruitment and
H3K27 Methylation. Mol. Cell. 74, 8–18 (2019).

24.   S. Chyb, A. Dahanukar, A. Wickens, J. R. Carlson, Drosophila Gr5a encodes a taste receptor tuned
to trehalose. Proc. Natl. Acad. Sci. U. S. A. 100 Suppl 2, 14526–14530 (2003).

25.   B. Kiragasi, J. Wondolowski, Y. Li, D. K. Dickman, A Presynaptic Glutamate Receptor Subunit
Confers Robustness to Neurotransmission and Homeostatic Potentiation. Cell Rep. 19, 2694–2706
(2017).

26.   B. van Steensel, S. Henikoff, Identification of in vivo DNA targets of chromatin proteins using
tethered dam methyltransferase. Nat. Biotechnol. 18, 424–428 (2000).

27.   T. D. Southall, K. S. Gold, B. Egger, C. M. Davidson, E. E. Caygill, O. J. Marshall, A. H. Brand,
Cell-type-specific profiling of gene expression and chromatin binding without cell isolation:
assaying RNA Pol II occupancy in neural stem cells. Dev. Cell. 26, 101–112 (2013).

28.   O. J. Marshall, T. D. Southall, S. W. Cheetham, A. H. Brand, Cell-type-specific profiling of
protein–DNA interactions without cell isolation using targeted DamID with next-generation
sequencing. Nat. Protoc. 11, 1586–1598 (2016).

29.   G. N. Aughey, A. E. Gomez, J. Thomson, H. Yin, T. D. Southhall, CATaDa reveals global
remodelling of chromatin accessibility during stem cell differentiation in vivo. eLife, 7 (2018).

30.   G. J. Filion, J. G. van Bemmel, U. Braunschweig, W. Talhout, J. Kind, L. D. Ward, W. Brugman,
I. J. de Castro, R. M. Kerkhoven, H. J. Bussemaker, B. van Steensel, Systematic protein location
mapping reveals five principal chromatin types in Drosophila cells. Cell. 143, 212–224 (2010).

31.   C. S. Chan, L. Rastelli, V. Pirrotta, A Polycomb response element in the Ubx gene that determines
an epigenetically inherited state of repression. EMBO J. 13, 2553–2564 (1994).

32.   J. Müller, J. A. Kassis, Polycomb response elements and targeting of Polycomb group proteins in
Drosophila. Curr. Opin. Genet. Dev. 16, 476–484 (2006).

33.   M. Khabiri, P. L. Freddolino, Genome-wide Prediction of Potential Polycomb Response Elements
and their Functions. bioRxiv (2019), p. 516500.

34.   H. Goodarzi, O. Elemento, S. Tavazoie, Revealing global regulatory perturbations across human
cancers. Mol. Cell. 36, 900–911 (2009).

35.   X. Chen, D. Dickman, Development of a tissue-specific ribosome profiling approach in
Drosophila enables genome-wide evaluation of translational adaptations. PLoS Genet. 13,
e1007117 (2017).

36. K. Scott, Gustatory Processing in Drosophila melanogaster. Annu. Rev. Entomol. 63, 15–30
(2018).

37. D. Szklarczyk, A. L. Gable, D. Lyon, A. Junge, S. Wyder, J. Huerta-Cepas, M. Simonovic, N. T.
Doncheva, J. H. Morris, P. Bork, L. J. Jensen, C. von Mering, STRING v11: protein–protein
association networks with increased coverage, supporting functional discovery in genome-wide
experimental datasets. Nucleic Acids Res. 47, D607–D613 (2019).

38. K. Ueno, Y. Kidokoro, Adenylyl cyclase encoded by AC78C participates in sugar perception in
Drosophila melanogaster. Eur. J. Neurosci. 28, 1956–1966 (2008).

39. J. Ashley, B. Cordy, D. Lucia, L. G. Fradkin, V. Budnik, T. Thomson, Retrovirus-like Gag Protein
Arc1 Binds RNA and Traffics across Synaptic Boutons. Cell. 172, 262–274.e11 (2018).

40. F. Laprell, K. Finkl, J. Müller, Propagation of Polycomb-repressed chromatin requires sequence-
specific recruitment to DNA. Science. 356, 85–88 (2017).

41. R. T. Coleman, G. Struhl, Causal role for inheritance of H3K27me3 in maintaining the OFF state
of a Drosophila HOX gene. Science. 356 (2017), doi:10.1126/science.aai8236.

42. P. N. C. B. Audergon, S. Catania, A. Kagansky, P. Tong, M. Shukla, A. L. Pidoux, R. C. Allshire,
Epigenetics. Restricted epigenetic inheritance of H3K9 methylation. Science. 348, 132–135
(2015).

43. K. Ragunathan, G. Jih, D. Moazed, Epigenetics. Epigenetic inheritance uncoupled from sequence-
specific recruitment. Science. 348, 1258699 (2015).

44. L. J. Gaydos, W. Wang, S. Strome, Gene repression. H3K27me and PRC2 transmit a memory of
repression across generations and during development. Science. 345, 1515–1518 (2014).

45. J. Z. Parrish, K. Emoto, L. Y. Jan, Y. N. Jan, Polycomb genes interact with the tumor suppressor
genes hippo and warts in the maintenance of Drosophila sensory neuron dendrites. Genes Dev. 21,
956–972 (2007).

46. J. Z. Parrish, M. D. Kim, L. Y. Jan, Y. N. Jan, Genome-wide analyses identify transcription factors
required for proper morphogenesis of Drosophila sensory neuron dendrites. Genes Dev. 20, 820–
835 (2006).

47. E. P. R. Iyer, S. C. Iyer, L. Sullivan, D. Wang, R. Meduri, L. L. Graybeal, D. N. Cox, Functional
Genomic Analyses of Two Morphologically Distinct Classes of Drosophila Sensory Neurons:
Post-Mitotic Roles of Transcription Factors in Dendritic Patterning. PLoS ONE. 8 (2013), p.
e72434.

48. G. Vorbrüggen, R. Constien, O. Zilian, E. A. Wimmer, G. Dowe, H. Taubert, M. Noll, H. Jäckle,
Embryonic expression and characterization of a Ptx1 homolog in Drosophila. Mech. Dev. 68, 139–
147 (1997).

49. S. Zaffran, G. Das, M. Frasch, The NK-2 homeobox gene scarecrow (scro) is expressed in
pharynx, ventral nerve cord and brain of Drosophila embryos. Mech. Dev. 94, 237–241 (2000).

50. M. M. Corty, J. Tam, W. B. Grueber, Dendritic diversification through transcription factor-
mediated suppression of alternative morphologies. Development. 143, 1351–1362 (2016).

51. C. J. Neumann, S. M. Cohen, Boundary formation in Drosophila wing: Notch activity attenuated
by the POU protein Nubbin. Science. 281, 409–413 (1998).

52. C. Q. Doe, Temporal Patterning in the Drosophila CNS. Annu. Rev. Cell Dev. Biol. 33, 219–240
(2017).

53. M. Kohwi, C. Q. Doe, Temporal fate specification and neural progenitor competence during
development. Nat. Rev. Neurosci. 14, 823–838 (2013).

54. S. Bahrampour, C. Jonsson, S. Thor, Brain expansion promoted by polycomb-mediated anterior
enhancement of a neural stem cell proliferation program. PLoS Biol. 17, e3000163 (2019).

55.　F. Marasca, B. Bodega, V. Orlando, How Polycomb-Mediated Cell Memory Deals With a
Changing Environment: Variations in PcG complexes and proteins assortment convey plasticity to
epigenetic regulation as a response to environment. Bioessays. 40, e1700137 (2018).

56.　A. Kolybaba, A.-K. Classen, Sensing cellular states—signaling to chromatin pathways targeting
Polycomb and Trithorax group function. Cell Tissue Res. 356, 477–493 (2014).

57.　M. C. Gambetta, K. Oktaba, J. Müller, Essential role of the glycosyltransferase sxc/Ogt in
polycomb repression. Science. 325, 93–96 (2009).

58.　M. C. Gambetta, J. Müller, O-GlcNAcylation prevents aggregation of the Polycomb group
repressor polyhomeotic. Dev. Cell. 31, 629–639 (2014).

59.　C.-S. Chu, P.-W. Lo, Y.-H. Yeh, P.-H. Hsu, S.-H. Peng, Y.-C. Teng, M.-L. Kang, C.-H. Wong, L.-
J. Juan, O-GlcNAcylation regulates EZH2 protein stability and function. Proc. Natl. Acad. Sci. U.
S. A. 111, 1355–1360 (2014).

60.　A. Öst, A. Lempradl, E. Casas, M. Weigert, T. Tiko, M. Deniz, L. Pantano, U. Boenisch, P. M.
Itskov, M. Stoeckius, M. Ruf, N. Rajewsky, G. Reuter, N. Iovino, C. Ribeiro, M. Alenius, S.
Heyne, T. Vavouri, J. A. Pospisilik, Paternal diet defines offspring chromatin state and
intergenerational obesity. Cell. 159, 1352–1364 (2014).

61.　T. T.-H. Lu, S. Heyne, E. Dror, E. Casas, L. Leonhardt, T. Boenke, C.-H. Yang, Sagar, L.
Arrigoni, K. Dalgaard, R. Teperino, L. Enders, M. Selvaraj, M. Ruf, S. J. Raja, H. Xie, U.
Boenisch, S. H. Orkin, F. C. Lynn, B. G. Hoffman, D. Grün, T. Vavouri, A. M. Lempradl, J. A.
Pospisilik, The Polycomb-Dependent Epigenome Controls β Cell Dysfunction, Dedifferentiation,
and Diabetes. Cell Metab. 27, 1294–1308.e7 (2018).

62.　K. Dalgaard, K. Landgraf, S. Heyne, A. Lempradl, J. Longinotto, K. Gossens, M. Ruf, M.
Orthofer, R. Strogantsev, M. Selvaraj, T. T.-H. Lu, E. Casas, R. Teperino, M. A. Surani, I.
Zvetkova, D. Rimmington, Y. C. L. Tung, B. Lam, R. Larder, G. S. H. Yeo, S. O'Rahilly, T.
Vavouri, E. Whitelaw, J. M. Penninger, T. Jenuwein, C.-L. Cheung, A. C. Ferguson-Smith, A. P.
Coll, A. Körner, J. A. Pospisilik, Trim28 Haploinsufficiency Triggers Bi-stable Epigenetic
Obesity. Cell. 164, 353–364 (2016).

63.　E. S. Deneris, O. Hobert, Maintenance of postmitotic neuronal cell identity. Nat. Neurosci. 17,
899–907 (2014).

64.　J. Z. Parrish, C. C. Kim, L. Tang, S. Bergquist, T. Wang, J. L. Derisi, L. Y. Jan, Y. N. Jan, G. W.
Davis, Krüppel mediates the selective rebalancing of ion channel expression. Neuron. 82, 537–544
(2014).

65.　E. Marder, A. A. Prinz, Modeling stability in neuron and network function: the role of activity in
homeostasis. Bioessays. 24, 1145–1154 (2002).

66.　J. D. Sweatt, The emerging field of neuroepigenetics. Neuron. 80, 624–632 (2013).

67.　E. J. Nestler, C. Lüscher, The Molecular Basis of Drug Addiction: Linking Epigenetic to Synaptic
and Circuit Mechanisms. Neuron. 102, 48–59 (2019).

68.　C. Dulac, Brain function and chromatin plasticity. Nature. 465, 728–735 (2010).

69.　F. A. Champagne, Epigenetic mechanisms and the transgenerational effects of maternal care.
Front. Neuroendocrinol. 29, 386–397 (2008).

70.　L. P. Musselman, R. P. Kühnlein, Drosophila as a model to study obesity and metabolic disease. J.
Exp. Biol. 221 (2018), doi:10.1242/jeb.163881.

71.　J. Na, L. P. Musselman, J. Pendse, T. J. Baranski, R. Bodmer, K. Ocorr, R. Cagan, A Drosophila
Model of High Sugar Diet-Induced Cardiomyopathy. PLoS Genetics. 9 (2013), p. e1003175.

72.　J. M. Tennessen, W. E. Barry, J. Cox, C. S. Thummel, Methods for studying metabolism in
Drosophila. Methods. 68, 105–115 (2014).

73. E. E. LeDue, Y.-C. Chen, A. Y. Jung, A. Dahanukar, M. D. Gordon, Pharyngeal sense organs
drive robust sugar consumption in Drosophila. Nature Communications. 6 (2015), ,
doi:10.1038/ncomms7667.

74. S. Andrews, Others, FastQC: a quality control tool for high throughput sequence data (2010).

75. A. Dobin, C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson, T. R.
Gingeras, STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 29, 15–21 (2013).

76. S. Anders, P. T. Pyl, W. Huber, HTSeq--a Python framework to work with high-throughput
sequencing data. Bioinformatics. 31, 166–169 (2015).

77. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. Genome Biol. 15, 550 (2014).

78. R. Kolde, Pheatmap: pretty heatmaps. R package version (2012) (available at
https://scholar.google.ca/scholar?cluster=14492249782051319874,4384187057944727023,160593
88166350612695,16769163128501098493,5741623497242498138&hl=en&as_sdt=0,5&sciodt=0,
5).

79. L. Wilkinson, Exact and approximate area-proportional circular Venn and Euler diagrams. IEEE
Trans. Vis. Comput. Graph. 18, 321–331 (2012).

80. K. Blighe, EnhancedVolcano: Publication-ready volcano plots with enhanced colouring and
labeling. R package version 1.2. 0 (2019).

81. O. J. Marshall, A. H. Brand, damidseq_pipeline: an automated pipeline for processing DamID
sequencing datasets. Bioinformatics. 31, 3371–3373 (2015).

82. B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. Nat. Methods. 9, 357–
359 (2012).

83. F. Ramírez, D. P. Ryan, B. Grüning, V. Bhardwaj, F. Kilpert, A. S. Richter, S. Heyne, F. Dündar,
T. Manke, deepTools2: a next generation web server for deep-sequencing data analysis. Nucleic
Acids Res. 44, W160–5 (2016).

84. A. R. Quinlan, I. M. Hall, BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics. 26, 841–842 (2010).

85. R. Stark, G. Brown, Others, DiffBind: differential binding analysis of ChIP-Seq peak data. R
package version. 100, 4–3 (2011).

86. J. Feng, T. Liu, B. Qin, Y. Zhang, X. S. Liu, Identifying ChIP-seq enrichment using MACS. Nat.
Protoc. 7, 1728–1740 (2012).

87. J. Thurmond, J. L. Goodman, V. B. Strelets, H. Attrill, L. S. Gramates, S. J. Marygold, B. B.
Matthews, G. Millburn, G. Antonazzo, V. Trovisco, T. C. Kaufman, B. R. Calvi, FlyBase
Consortium, FlyBase 2.0: the next generation. Nucleic Acids Res. 47, D759–D765 (2019).

88. Y. Benjamini, Y. Hochberg, Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. J. R. Stat. Soc. Series B Stat. Methodol. 57, 289–300 (1995).

89. C. E. Grant, T. L. Bailey, W. S. Noble, FIMO: scanning for occurrences of a given motif.
Bioinformatics. 27, 1017–1018 (2011).

90. D. W. Huang, B. T. Sherman, R. A. Lempicki, Systematic and integrative analysis of large gene
lists using DAVID Bioinformatics Resources. Nature Protoc. 2009; 4 (1): 44--57 (2008).

91. D. W. Huang, B. T. Sherman, R. A. Lempicki, Bioinformatics enrichment tools: paths toward the
comprehensive functional analysis of large gene lists. Nucleic Acids Res. 37, 1–13 (2009).
92. P. Shannon, A. Markiel, O. Ozier, N. S. Baliga, J. T. Wang, D. Ramage, N. Amin, B.
Schwikowski, T. Ideker, Cytoscape: a software environment for integrated models of biomolecular
interaction networks. Genome Res. 13, 2498–2504 (2003).

Acknowledgements: We thank Sundeep Kalantry, Josie Clowney, and Giacomo Cavalli for helpful

comments and discussions. We also thank the University of Indiana at Bloomington, the VDRC, the

FLYORF stock collections, and all the investigators who kindly shared fly lines with us. We also thank

Daniel Wilinski for help with data analysis, Dion Dickman and Xun Chen for helpful comments on

riboTRAP, and Margarita Brovkina (E.J. Clowney lab) for helpful discussions on TaDa analysis. Matt

Burkhard assisted with analysis of expression changes for different Gene Ontology terms. Julia Kuhn

designed some of the graphics for the manuscript. This work was funded by NIH R00 DK-97141 and NIH

1DP2DK-113750, the Klingenstein-Simons Fellowship in the Neurosciences, and the Rita Allen

Foundation (to M.D.) and NIH R35 GM-128637 (to P.L.F). **Author contributions:** A.V performed all

experiments and analyzed RNAseq, TaDa, and CATaDa datasets, with the exception of *in vivo* calcium

imaging. B.T.G helped with PER and triglyceride measurements. C.E.M performed *in vivo* calcium

imaging. M.K and P.L.F developed the PREdictor, calculated regulatory targets of transcription factors,

ran and analyzed Gene Ontology term enrichment analysis, provided statistical consultation, and tested for

practical significance. M.D oversought the project and secured funding. A.V and M.D designed the

experiments, wrote the manuscript, and prepared the figures with input from all authors. **Competing**

interests: The authors declare no competing interests. **Data availability:** All high throughput sequencing

data files can be found on Gene Expression Omnibus GSE146245.

A

PRC2.1

E(z)

Pcl

Esc   Su(z)12

PRC2

B

w1118^{cs} *E(z)*^{c249} *Su(z)12*^{c253} *esc*^{c289}

PER

1.0

0.5

0.0

○ CD
□ SD

****

****

***

30 10 5   30 10 5   30 10 5   30 10 5

[%Sucrose]

C

+*vehicle*   +*EEDi*

PER

1.0

0.5

0.0

****

****

○ CD
□ SD

30 10 5   30 10 5

[%Sucrose]

D

w1118^{cs} *Pcl*^{c429} *Pcl*^{T1}

PER

1.0

0.5

0.0

****

****

○ CD
□ SD

30 10 5   30 10 5   30 10 5

[%Sucrose]

E

PER

1.0

0.5

0.0

Gr5a-GAL4/+
UAS-Pcl^{RNAi-1}*/+*
Gr5a>Pcl^{RNAi-1}

**
****

****
****

□ SD

30 10 5

[%Sucrose]

F

PER

1.0

0.5

0.0

Gr5a-GAL4/+
UAS-Pcl/+
Gr5a>Pcl

**

****
*

****
****

○ CD

30 10 5

[%Sucrose]

G

+*vehicle*   +*EEDi*

PER

1.0

0.5

0.0

Gr5a-GAL4/+
UAS-Pcl/+
Gr5a>Pcl

*

****
****

○ CD

30 10 5   30 10 5

[%Sucrose]

Fig. 1. PRC2.1 modulates sweet taste in response to the food environment.

(A) Schematic of *Drosophila melanogaster* PRC2 core complex consisting of *Enhancer of zeste* (*E(z))*,

Suppressor of zeste (*Su(z)12)*, *extra sex combs (esc*) and the accessory protein *Polycomb-like* (*Pcl*). **(B-G)**

Taste responses (y axis) to stimulation of the labellum with 30%, 10% and 5% sucrose (x axis) of age-

matched males **(B)** $w1118_{cs}$ (gray), $E(z)_{c249/+}$ (dark red), $Su(z)12_{c253/+}$ (fuschia), and $esc_{c289/+}$ (red) flies on a

control (circle) or sugar (square) diet for 7 days. n = 34-68; **(C)** $w1118_{cs}$ flies on a control (circle) or sugar

(square) diet supplemented with vehicle (10% DMSO) or 8 μM EEDi for 7 days. n = 32-43; **(D)** $w1118_{cs}$

(gray), $Pcl_{c429/+}$ (pink) and $Pcl_{T1/+}$ (pink) flies on a control (circle) or sugar (square) diet for 7 days. n = 36-

82; **(E)** $Gr5a>Pcl_{RNAi-1}$ (pink), and parental transgenic controls (gray, crossed to $w1118_{cs}$) flies on a sugar

diet for 7 days. n = 42-63; **(F)** $Gr5a>Pcl$ (pink), and parental transgenic controls (gray, crossed to $w1118_{cs}$)

flies on a control diet for 7 days. n = 36-61; **(G)** $Gr5a>Pcl$ flies on a control diet supplemented with

vehicle (10% DMSO) or 8 μM EEDi for 7 days. n = 30-35. Data are shown as mean ± SEM, Kruskal-

Wallis Dunn's multiple comparisons, comparisons to control diet within genotype groups. **** $p <$

0.0001, *** $p < 0.001$, ** $p < 0.01$, *$p < 0.05$ for all panels in this figure.

Fig. 2. *Pcl* mutant flies have normal sensory responses and are resistant to diet-induced obesity.

(A) Schematic showing the setup for *in vivo* calcium imaging: the fly proboscis is stimulated with 20%

sucrose while recording from the subesophageal zone (SEZ) which contains the presynaptic terminals of

the sweet taste neurons (shown here labeled with *synaptotagmin::GFP*). **(B)** Average $\Delta F/F_0$ calcium

response traces (left panel) and peak % $\Delta F/F_0$ (right panel) responses to sucrose stimulation of the

proboscis (arrow) in age-matched male *Gr64f>GCaMP6s-Bruchpilot-mCherry;Pcl$_{c429}$* flies fed a control

(gray, circle) or sugar (pink, square) diet for 7 days, n = 32-34. **(C)** Quantification of GFP-labeled cells in

the labella of *Gr64f;CD8-GFP* flies crossed to *w1118$_{cs}$* (as control, dark gray) or *Gr64f;CD8-*

GFP>Pcl$_{RNAi}$ (pink) on a control (circle) or sugar (square) diet for 7 days. n=5-16 flies, Kruskal-Wallis

Dunn's multiple comparisons, comparison to control diet of each genotype, no significance. **(D)**

Triglyceride levels normalized to protein in *Gr5a>Pcl$_{RNAi}$* (pink) and parental transgenic control flies (dark

gray, crossed to *w1118$_{cs}$*) fed a control (circle) or sugar (square) diet for 7 days. n=8, two-way ANOVA

with Sidak's multiple comparisons test, comparisons to control diet of each genotype. All data are shown

as mean ± SEM, $****p < 0.0001$, $***p < 0.001$, $**p < 0.01$, and $*p < 0.05$ for all panels unless indicated.

Fig. 3. *Pcl* chromatin occupancy is redistributed in the high sugar environment.

(A) Schematic of Targeted Dam-ID (TaDa) of *Dam::Pcl* and *Dam* (CATaDa, top panel) and of TaDa

induction paradigm (bottom panel). Age-matched *Gr5a;tubulin-GAL80>UAS-LT3-Dam::Pcl* and

Gr5a;tubulin-GAL80>UAS-LT3-Dam flies were placed on a control (gray) or sugar (purple) diet for 3

days at 20-21°C, then switched to 28°C between day 3-4 to induce *Dam* expression (arrow) (bottom panel).

(B) CATaDa chromatin coverage (chromatin accessibility) normalized to 1x dm6 genome (reads per

genome coverage) of the three biological replicates from control diet flies within a 5 kb window at the

sweet gustatory receptor *Gr5a* and the bitter gustatory receptor *Gr66a*. **(C)** Fold change of the proportion

of genes allocated to the five chromatin states (assigned in color) according to their transcription start site

(TSS), normalized to the expected proportion across the whole genome (see Methods). **(D)** Mean

$\log_2$(*Dam::Pcl/Dam*) chromatin binding coverage for a 10 kb window centered on predicted PREs on a

control (gray) and sugar (purple) diet. **(E-F)** Mean CATaDa chromatin coverage normalized to 1x dm6

genome for a 10 kb window centered on **(E)** *Pcl* peaks and **(F)** predicted PREs on a control (gray) and

sugar (yellow) diet. **(G)** Box plots represent the median and variance of normalized *Dam::Pcl/Dam* reads

for genes differentially bound by *Pcl* on a control (gray) and sugar (purple) diet. Genes are placed into

groups with higher (Group 1) or lower (Group 2) *Pcl* chromatin binding in the sugar diet environment. **(H)**

GO terms associated with differentially bound genes by *Dam::Pcl/Dam* identified by iPAGE, text boxes

represent the GO term category, metabolism (orange) and regulatory (lavender) (for the full iPAGE list see

Supplemental Figure 4). For all panels peak FDR<0.01 (see Methods).

Fig. 4. PRC2.1 sculpts the transcriptional responses of the *Gr5a+* neurons in response to diet.

(A) Schematic depiction of the TRAP technique used to profile the changes in the *Gr5a+* neurons. **(B-C)** Volcano plots representing changes in gene expression in the *Gr5a+* neurons of age matched male *Gr5a>UAS-Rpl3-3XFLAG* flies on a sugar diet for 3 days (SD3, mint) and 7 days (SD7, teal). n=2-3 replicates per condition (10,000 *Gr5a+* cells per replicate). Non-significant genes are shown in black and genes with $q<0.1$ (Wald test) are in mint or teal for SD3 and SD7 respectively, comparison is to the control diet group. **(D)** Venn diagram of differentially expressed genes (DEGs) at SD3/CD (mint), SD7/CD (teal), and the overlap between SD3/SD7 (Wald test, $q<0.1$). Heatmaps show $\log_2$ fold changes ($l_2$fc) for DEGs in each condition specified in Venn diagram (left columns in heatmap, SD3, SD7, and SD3+SD7, all comparisons to control diet) and *Pcl$_{c429}$* mutant flies (right column in heatmap, *Pcl$_{c429}$ SD7* comparison to control diet); $l_2$fc range from purple (high) to gold (low). **(E)** Normalized read counts (Tpm+1) from TRAP for *Pcl* bound (pink) and not bound genes (gray) on a control diet. Box plots represent the median and variance, *p*-value = $3.196 \times 10_{-6}$, two-tailed t-test. **(F)** $\log_2$ fold changes for *scarecrow* (*Scro*), *Paired-like homeobox domain 1* (*Ptx1*), *caudal* (*cad*), *GATAe,* and *nubbin* (*nub*) in control SD7/CD, SD3/CD flies, and *Pcl$_{c429}$*SD7/CD mutant flies. L_2fc ranges from purple (high) to gold (low).

Fig. 5. PRC2.1 represses a transcriptional program required for sweet taste.

(A) $Log_2(Dam::Pcl/Dam)$ chromatin binding coverage tracks of flies on a control (black) and sugar

(purple) diet within a 50 kb window at the *cad*, *Ptx1*, *GATAe*, *nub*, and *Scro* genes visualized with the

UCSD genome browser. The three biological replicates for each condition are superimposed. Turquoise

traces are SD/CD fold changes. Peaks are shown as black boxes below respective traces ($q<0.01$) and

genes are shown in dense format to include all potential isoforms. **(B)** Taste responses (y axis) to

stimulation of the labellum with 30%, 10% and 5% sucrose (x axis) in age-matched males of

Gr5a>cad_{RNAi} (gold), *Gr5a>Ptx1_{RNAi}* (blue), *Gr5a>GATAe_{RNAi}* (green), *Gr5a>nub_{RNAi}* (lavender),

Gr5a>Scro (salmon), and parental transgenic control (gray, crossed to $w1118_{cs}$) flies on a control diet for 7

days. n = 30-54, Kruskal-Wallis Dunn's multiple comparisons, comparisons to both transgenic controls.

All data shown as mean ± SEM, $****p < 0.0001$, $***p < 0.001$, $**p < 0.01$, and $*p < 0.05$. **(C)** Box plots

represent the median and variance of the log$_2$ fold changes (l$_2$fc) for candidate gene targets of *cad*, *Ptx1*,

GATAe, *nub*, and *Scro* (for threshold cutoff score see Supplemental Figure 9) in control flies at SD7 (teal),

and *Pcl_{c429}* mutants at SD7 (pink). **(D)** Venn diagram representation of the overlap of the candidate gene

targets of *cad* (gold), *Ptx1* (blue), *GATAe* (green), *nub* (lavender), and *Scro* (salmon). **(E)** A transcriptional

loop between *cad* (gold), *Ptx1* (blue), *GATAe* (green), *nub* (lavender), and *Scro* (salmon) mediated by *Pcl*.

Green arrows represent activation and red bars represent inhibition. GO terms associated with each

transcriptional regulator identified by iPAGE are listed. Text boxes represent the GO term category,

metabolism (orange), regulatory (lavender) and neural/signal (blue) (for the full iPAGE list see

Supplemental File 5).

Fig. 6. The persistent phenotypic memory of the food environment is dependent on PRC2.1.

(A) Diagram representing dietary manipulations, CD, control diet (gray), SD, sugar diet (purple), and >

represents the switch in diet. **(B)** Taste responses (y axis) to stimulation of the labellum with 30%, 10%

and 5% sucrose (x axis) of age-matched males $w1118_{cs}$ flies on a control (CD>CD, circle), control to sugar

(CD>SD, triangle), and sugar to control (SD>CD, diamond) diet as shown in (A) n =57-64, Kruskal-Wallis

Dunn's multiple comparisons, comparisons to control diet. All data shown as mean ± SEM, $****p <$

0.0001, $***p < 0.001$, $**p < 0.01$, and $*p < 0.05$ in this panel. **(C)** Volcano plot with the changes in gene

expression in the $Gr5a+$ neurons of age-matched male $Gr5a>UAS-Rpl3-3XFLAG$ flies on a sugar to

control (SD>CD, green) diet compared to the control diet group (CD>CD, Wald test, $q<0.1$, ns is non-

significant). n=2 replicates per condition (~10,000 $Gr5a+$ cells per replicate). **(D)** Heatmap showing how

the differentially expressed genes identified in Figure 5C are changing in the SD>CD condition (310/658,

48% Wald test, $q<0.1$) compared to the other conditions SD3, SD7, and $Pclc_{429}$SD7 (column headings, all

comparisons to control diet). Log2fold changes for these genes were determined by comparing each group

to its own control (see Methods). L2fc ranges from purple (high) to gold (low). **(E)** Taste responses (y axis)

to stimulation of the labellum with 30%, 10% and 5% sucrose (x axis) of age-matched males $w1118_{cs}$ flies

on a control (circle), control to sugar (triangle), and sugar to control (diamond, green) diet for 14 days.

EEDi was supplemented to the diets at the day 7 transition (see diagram in figure). n =34-46, Kruskal-

Wallis Dunn's multiple comparisons, comparisons to control diet. All data shown as mean ± SEM, $****p$

< 0.0001, $***p < 0.001$, $**p < 0.01$, and $*p < 0.05$ in this panel. **(F)** Model of molecular changes in the

$Gr5a+$ neurons on a control (top panel) and sugar (bottom panel) diet, showing the redistribution in

PRC2.1 binding, and the effects on the transcription of the regulators (green and red arrows), and neural

responses to sweet stimuli.

Vaziri et al.
Supplement 1

Fig. S1. The PRC1 and PhoRC complex are not required for sugar diet mediated sweet taste defects.

(A) The fly proboscis contains the sweet sensory neurons soma (green) with dendrites extending into the

hair sensilla and their axons projecting onto the subesophageal zone (SEZ, delimited in yellow) of the fly

brain. **(B-E)** Taste responses (y axis) to stimulation of the labellum with 30%, 10% and 5% sucrose (x

axis) of age-matched males of **(B)** $Gr5a>Jarid2_{RNAi-1}$ (dark blue), $Gr5a>Jarid_{RNAi-2}$ (light blue), and

$Gr5a>jing_{RNAi-1}$ (orange) and parental transgenic control (gray, crossed to $w1118_{cs}$) flies on a sugar diet for

7 days. n = 25-34, Kruskal-Wallis Dunn's multiple comparisons, comparisons to transgenic controls. **(C)**

$w1118_{cs}$, ph-$d_{[401]/+}ph$-$p_{[602]/+}$, $Psc_{27/+}$,and $Pc_{1/+}$ flies on a control (circle) or sugar (square) diet for 7 days.

n = 30-42, Kruskal-Wallis Dunn's multiple comparisons, comparisons to control diet within each genotype

group. **(D)** $w1118_{cs}$, $phol_{[81A]/+}$ flies on a control (circle) or sugar (square) diet for 7 days. n = 32-34,

Kruskal-Wallis Dunn's multiple comparisons, comparisons to control diet within each genotype group. **(E)**

$Gr5a>Sfmbt_{RNAi}$ (dark pink) and parental transgenic control (gray, crossed to $w1118_{cs}$) flies on a sugar diet

for 7 days. n = 26-46, Kruskal-Wallis Dunn's multiple comparisons, comparisons to transgenic controls.

All data shown as mean ± SEM, ∗∗∗∗$p < 0.0001$, ∗∗∗$p < 0.001$, ∗∗$p < 0.01$, and ∗$p < 0.05$ for all panels

unless indicated.

Fig. S2. The effects of *Pcl* on sweet taste are independent of changes in cell number or development.

(A) Fold change of *Pcl* mRNA levels in the heads of flies with and without pan-neuronal (*nsyb-GAL4*) *Pcl*

knockdown with two independent RNAi lines (dark gray, plum, and pink respectively) measured by qPCR

and normalized to the gene *Rp49*. n=6, Kruskal-Wallis Dunn's multiple comparisons, compared to GAL4

transgenic control (crossed to *w1118cs*). **(B-C)** Taste responses (y axis) to stimulation of the labellum with

30%, 10% and 5% sucrose (x axis) of age-matched males of **(B)** *Gr5a>PclRNAi-2* and parental transgenic

control (gray, crossed to *w1118cs*) flies on a sugar diet for 7 days. n = 42-51, Kruskal-Wallis Dunn's

multiple comparisons, comparisons to transgenic controls. **(C)** *Gr5a>PclRNAi-1* and parental transgenic

control (gray, crossed to *w1118cs*) flies on a control diet for 7 days. n =30-35, Kruskal-Wallis Dunn's

multiple comparisons, comparisons to transgenic controls. **(D)** Quantification of the number of sweet taste

GFP-labeled cells in the labella of *Gr64f;CD8-GFP* flies crossed to *w1118cs* (as control, gray) or

Gr64f;CD8-GFP>Pcl (pink) on a control (circle) or sugar (square) diet for 7 days. n=5-16 probosces, no

significance, Kruskal-Wallis Dunn's multiple comparisons, comparison to control diet of each genotype.

(E) Taste responses (y axis) to stimulation of the labellum with 30%, 10% and 5% sucrose (x axis) of age-

matched males of *Gr5a;tubulin-GAL80ts>PclRNAi-1*, *Gr5a;tubulin-GAL80ts>Pcl,* and parental transgenic

control (gray, crossed to *w1118cs*) flies on a control (circle, right) or sugar (square, left) diet for 7 days. n =

34-49, Kruskal-Wallis Dunn's multiple comparisons, comparisons to transgenic control. All data shown as

mean ± SEM, ∗∗∗∗$p < 0.0001$, ∗∗∗$p < 0.001$, ∗∗$p < 0.01$, and ∗$p < 0.05$ for all panels unless indicated.

1038

Fig. S3. Chromatin occupancy and accessibility analysis of *Pcl* in the *Gr5a+* neurons.

(A) Principal component analysis of normalized $\log_2$(*Dam::Pcl/Dam*) flies on a control (CD, gray) or

sugar (SD, purple) diet. **(B)** The three biological replicates of $\log_2$(*Dam::Pcl/Dam*) on a control (CD, gray)

or sugar (SD, purple) diet are shown, with high Pearson correlation coefficients within each dietary

condition, and low correlation coefficients between dietary conditions. Scale shows degree of correlation.

(C-D) Left panel: overlap of $\log_2$(*Dam::Pcl/Dam)* chromatin binding peaks between a control (gray) and

sugar (purple) diet (find_peaks, $q<0.01$). Right panel: overlap of CATaDa chromatin peaks between a

control (gray) and sugar (yellow) diet (MACS2, $q<0.05$). **(E-F)** $\log_2$*(Dam::Pcl/Dam*) chromatin binding

coverage at *Ultrabithorax* (*Ubx*), *abdominal A* (*abd-A*), *Abdominal B* (*Abd-B*), *polyhomeotic distal* (*ph-d*)

and *polyhomeotic proximal* (*ph-p*) (Top panel, E), and *Tat interactive protein 60* (*Tip60*) and *Suppressor of*

variegation 3-9 (*Su(var)3-9*) (Top panel, F). Bottom panel: CATaDa chromatin coverage (chromatin

accessibility) of the three biological replicates at the loci mentioned above. **(G)** Volcano plot showing the

differential chromatin binding of *Dam::Pcl* on a sugar diet compared to a control diet, (Wald test, $q<0.05$,

purple), non-significant peaks are shown in black (ns).

1053

 Fig. S4. Pathway enrichment analysis of *Pcl* chromatin targets in the *Gr5a+* neurons.

 iPAGE identification of pathways depleted (left) or enriched (right) compared to background gene list

 (middle) from the differentially bound chromatin peaks by *Dam::Pcl* flies on a sugar diet compared to a

 control diet. Scale represents over-representation (orange) or under-representation (purple) of genes within

 a specific bin for the corresponding GO term. Black outlined boxes represent *q*<0.05.

Fig. S5. PRC2.1 mediates the transcriptional responses of the *Gr5a+* neurons to diet.

(A) Principal component analysis of age matched male *Gr5a>UAS-Rpl3-3XFLAG* flies on a control diet

(CD, gray), sugar diet for three days (SD3, mint), and sugar diet for seven days (SD7, teal). **(B)**

Normalized read counts for *Gustatory receptor Gr66a* (bitter), *Gustatory receptor Gr64f, Gustatory*

receptor Gr5a, and *Gustatory receptor Gr64a* (all sweet) from the input fraction (dark green box) and

Gr5a+ fraction (light green box) specific conditions. Both samples also include the dietary conditions (not

specified in figure), hues of purple from light to dark represent low to high expression. **(C)** Principal

component analysis of age matched male *Gr5a;Pcl$_{c429}$>UAS-Rpl3-3XFLAG* flies on a control diet (CD,

gray) and sugar diet for seven days (SD7, pink). **(D)** Volcano plot with changes in gene expression in the

Gr5a+ neurons of age matched male *Gr5a;Pcl$_{c429}$>UAS-Rpl3-3XFLAG* flies on a SD for 7 days

(*Pcl$_{c429}$*SD7, pink) compared to the control diet condition in the same genotype (Wald test, $q<0.1$), n=3

replicates per condition (~10,000 *Gr5a+* cells per replicate). **(E)** Normalized reads (Tpm+1) from TRAP

for genes (x-axis) bound by *Pcl* and normalized to $\log_2$(*Dam::Pcl/Dam*) reads at these genes (y-axis) on a

control diet. Scale represents gene density (light to dark purple). **(F)** Fold change of *Dam::Pcl* chromatin

binding for genes differentially bound on sugar diet compared to control diet (y-axis) ($q<0.05$) and their

respective differentially expressed $\log_2$ fold changes on a sugar diet compared to control diet (x-axis)

($q<0.1$). Genes are shown as circles and colored based on GO term category, metabolism (orange) and

regulatory (lavender).

Fig. S6. Pathway enrichment analysis of the *Gr5a+* neurons in flies fed a sugar diet for 3 days.

iPAGE identification of pathways enriched in age-matched male *Gr5a>UAS-Rpl3-3XFLAG* flies on a

sugar diet for three days compared to a control diet. Bins (top) show the range of $\log_2$ fold changes for

genes within their corresponding GO terms. Scale represents over-representation (orange) or under-

representation (purple) of genes within a specific bin for the corresponding GO term. Black outlined boxes

represent *q*<0.05.

2.55
-4.12

SD7/CD

nucleolus, GO:0005730
G-protein coupled amine receptor activity, GO:0008227
cGMP biosynthetic process, GO:0006182
acetylcholine-activated cation-selective channel activity, GO:0004889
voltage-gated calcium channel complex, GO:0005891
snRNA pseudouridylation guide activity, GO:0030565
voltage-gated potassium channel complex, GO:0008076
gamma-tubulin binding, GO:0043015
positive regulation of glial cell proliferation, GO:0060252
G-protein coupled receptor activity, GO:0004930
sodium channel regulator activity, GO:0017080
sensory perception of chemical stimulus, GO:0007606
deactivation of rhodopsin mediated signaling, GO:0016059
histone H3 acetylation, GO:0043966
dendrite, GO:0030425
vesicle-mediated transport, GO:0016192
neuromuscular junction, GO:0031594
kinesin binding, GO:0019894
precatalytic spliceosome, GO:0071011
guanyl-nucleotide exchange factor activity, GO:0005085
GTPase activator activity, GO:0005096
circadian rhythm, GO:0007623
DNA repair, GO:0006281
mitotic sister chromatid separation, GO:0051306
neuron projection morphogenesis, GO:0048812
asymmetric neuroblast division, GO:0055059
RNA processing, GO:0006396
compound eye development, GO:0048749
protein complex, GO:0043234
muscle organ development, GO:0007517
TORC1 signaling, GO:0038202
transcription corepressor activity, GO:0003714
positive regulation of JNK cascade, GO:0046330
nuclear pore organization, GO:0006999
nucleosome, GO:0000786
cytosolic ribosome, GO:0022626
regulation of protein kinase activity, GO:0045859
chitin-based cuticle development, GO:0040003
mitochondrial large ribosomal subunit, GO:0005762
NADH dehydrogenase activity, GO:0003954
ligand-gated ion channel activity, GO:0015276
endomembrane system, GO:0012505
pyridoxal phosphate binding, GO:0030170
proteasome-mediated ubiquitin-dependent process, GO:0043161
glutathione transferase activity, GO:0004364
metabolic process, GO:0008152
plasma membrane proton-transporting V-type ATPase complex, GO:0033181
carbohydrate metabolic process, GO:0005975
UDP-glucose metabolic process, GO:0006011
transporter activity, GO:0005215
glycolytic process, GO:0006096
serine-type endopeptidase inhibitor activity, GO:0004867
dynein complex, GO:0030286
transmembrane transporter activity, GO:0022857
aminopeptidase activity, GO:0004177
secondary active organic cation transporter activity, GO:0008513
iron ion binding, GO:0005506
alkaline phosphatase activity, GO:0004035
defense response to Gram-positive bacterium, GO:0050830
metallocarboxypeptidase activity, GO:0004181
ketohexokinase activity, GO:0004454
hexose metabolic process, GO:0019318
alpha-mannosidase activity, GO:0004559
lipase activity, GO:0016298
sperm storage, GO:0046693
glucose transmembrane transporter activity, GO:0005355
larval serum protein complex, GO:0005616

Under-representation   -5   Over-representation

1085

Fig. S7. Pathway enrichment analysis of the *Gr5a+* neurons in flies fed a high sugar diet for 7 days.

iPAGE identification of pathways enriched in age-matched male *Gr5a>UAS-Rpl3-3XFLAG* flies on a

sugar diet for seven days compared to a control diet. Bins (top) show the range of $\log_2$ fold changes for

genes within their corresponding GO terms. Scale represents over-representation (orange) or under-

representation (purple) of genes within a specific bin for the corresponding GO term. Black outlined boxes

represent $q<0.05$.

Pcl^c429 SD7/CD

dynein complex, GO:0030286
sperm competition, GO:0046692
protein targeting to mitochondrion, GO:0006626
procollagen-proline 4-dioxygenase activity, GO:0004656
male meiosis, GO:0007140
protein polyglutamylation, GO:0018095
fructokinase activity, GO:0008865
fatty acid elongase activity, GO:0009922
nucleotide catabolic process, GO:0009166
aminopeptidase activity, GO:0004177
spermatogenesis, exchange of chromosomal proteins, GO:0035093
proteasome-mediated ubiquitin-dependent process, GO:0043161
catalytic step 2 spliceosome, GO:0071013
ribonucleoprotein complex assembly, GO:0022618
transcription factor TFIID complex, GO:0005669
AAG codon-amino acid adaptor activity, GO:0033444
AUG codon-amino acid adaptor activity, GO:0033436
nucleosome, GO:0000786
cytosolic ribosome, GO:0022626
rRNA 2'-O-methylation, GO:0000451
regulation of protein kinase activity, GO:0045859
dendrite membrane, GO:0032590
UAC codon-amino acid adaptor activity, GO:0033410
AUU codon-amino acid adaptor activity, GO:0033433
AAC codon-amino acid adaptor activity, GO:0033442
GAG codon-amino acid adaptor activity, GO:0033460
CCU codon-amino acid adaptor activity, GO:0033421
GUG codon-amino acid adaptor activity, GO:0033452
chitin-based cuticle development, GO:0040003
taste receptor activity, GO:0008527
compound eye development, GO:0048749
sensory perception of chemical stimulus, GO:0007606
behavioral response to ethanol, GO:0048149
chorion, GO:0042600
larval serum protein complex, GO:0005616
detection of chemical stimulus, GO:0009593
structural constituent of peritrophic membrane, GO:0016490
cellular response to starvation, GO:0009267
signal transduction, GO:0007165
synapse organization, GO:0050808
chromatin silencing, GO:0006342
vesicle-mediated transport, GO:0016192
embryonic development via the syncytial blastoderm, GO:0001700
neurotransmitter transport, GO:0006836
carbohydrate metabolic process, GO:0005975
male courtship behavior, GO:0045433
ribosome, GO:0005840
myofibril assembly, GO:0030239
positive regulation of neuron death, GO:1901216
translational initiation, GO:0006413
phototransduction, GO:0007602
septate junction, GO:0005918
mitochondrial respiratory chain complex IV, GO:0005751
endomembrane system, GO:0012505
neuropeptide receptor activity, GO:0008188
cell adhesion, GO:0007155

2.61
-0.49

Over-representation
5
5
Under-representation

1092

Fig. S8. Pathway enrichment analysis of the *Gr5a+* neurons in *Pclc429* mutant flies fed a high sugar

diet for 7 days.

iPAGE identification of pathways enriched in age-matched male *Gr5a;Pclc429>UAS-Rpl3-3XFLAG* flies

on a sugar diet for seven days compared to a control diet. Bins (top) show the range of $\log_2$ fold changes

for genes within their corresponding GO terms. Scale represents over-representation (orange) or under-

representation (purple) of genes within a specific bin for the corresponding GO term. Black outlined boxes

represent *q*<0.05.

1100

Fig. S9. Identification of the transcriptional program mediated by PRC2.1 on a sugar diet.

(A) Normalized read counts for *Scro*, *cad*, *nub*, *GATAe*, and *Ptx1* in the input fraction (dark green box) and the *Gr5a+* fraction (light green box) from *Gr5a>UAS-Rpl3-3XFLAG* age-matched male flies on a control diet; hues of purple from light to dark show low to high expression. **(B)** Taste responses (y axis) to stimulation of the labellum with 30%, 10% and 5% sucrose (x axis) of age-matched males of *Gr5a>cad (gold), Gr5a>Ptx1 (blue), Gr5a>nub (lavender), Gr5a>Scro_{RNAi}* (coral), and parental transgenic control (gray, crossed to *w1118_{cs}*) flies on a sugar diet for 7 days. n = 31-51, Kruskal-Wallis Dunn's multiple comparisons, comparisons to transgenic controls. **(C)** Schematic of how the targets of the 5 transcription factors in Figure 5 were identified: the DNA binding motifs for each transcription factor were scanned along every base pair of the *D. melanogaster* genome using FIMO, and the basepair-wise binding results converted to robust z-scores. **(D-H)** Threshold scores used to identify the regulons of each transcription factor. Y-axis represents the averaged log_2 fold change of the predicted targets based on the cutoff score, respectively.

1114

Fig. S10. PRC2.1 modulates the synaptic and neural properties of the *Gr5a+* neurons on a sugar

diet.

(A) Functional network created with STRING featuring PRC2.1, *cad*, *Ptx1*, *GATAe*, *nub*, *Scro,* and their

predicted targets (gray) from Figure 5C that can be classified into the GO term category neural/signaling.

Each node represents a gene. Edges indicate functional protein-protein interactions, not direct interactions.

Edges represented with dashed lines are interactions identified in this study. **(B-D)** Taste responses (y axis)

to stimulation of the labellum with 30%, 10% and 5% sucrose (x axis) of age-matched males of **(B)**

w1118cs, *Arc1esm18*, and *Arc1esm13* flies on a control diet for 7 days. n = 38-45, Kruskal-Wallis Dunn's

multiple comparisons, comparisons to transgenic controls. **(C)** *Gr5a>Arc1RNAi (turquoise),* and parental

transgenic control (gray, crossed to *w1118cs*) flies on a control diet for 7 days. n = 28-47, Kruskal-Wallis

Dunn's multiple comparisons, comparison to control diet. **(D)** *Gr5a>ACXDRNAi-1* (dark purple) *and*

Gr5a>ACXDRNAi-2 (light purple)*,* and parental transgenic controls (gray, crossed to *w1118cs*) flies on a

control diet for 7 days. n = 28-47, Kruskal-Wallis Dunn's multiple comparisons, comparison to control

diet. **(E)** Triglyceride levels normalized to protein in *w1118cs* flies fed a control (gray, circle) or sugar

(gray, square) diet for 14 days, and flies fed a SD>CD reversal diet for a total of 14 days (7 days in each

diet). n=16, two-way ANOVA with Sidak's multiple comparisons test, comparisons to control diet of each

genotype. All data are shown as mean ± SEM, $****p < 0.0001$, $***p < 0.001$, $**p < 0.01$, and $*p < 0.05$

for all panels unless indicated.

Supplementary Files

Supplementary file 1:

PRE prediction calls for the dmel6.08 genome. Relating to Figure 3.

Supplementary file 2:

Differential binding analysis of *Dam::Pcl/Dam* on a sugar diet compared to control diet.

Supplementary file 3:

Differential expression analysis of TRAP experiments for sugar diet 3 (SD3), sugar diet 7 (SD7), *Pcl$_{c429}$*

SD7 (all comparisons to control diet) and CD>CD and SD>CD.

Supplementary file 4:

Transcription factor regulatory target calls relating to Figures 5.

Supplementary file 5:

iPAGE identification of pathways over-represented or under-represented in the predicted gene targets of

GATAe, cad, nub, Ptx1 and *Scro.*

Supplementary file 6:

STRING interaction network relating to Supplement Figure 14.

Supplementary file 7:

Gene table for heatmap relating to Figure 6D.