Ion transport mechanisms for smoke inhalation injured airway epithelial barrier

Jianjun Chang[1,2], Zaixing Chen[1&], Runzhen Zhao[1], Hong-Guang Nie[3*], Hong-Long Ji[1,4*]

[1]Department of Cellular and Molecular Biology, University of Texas Health Science Center at Tyler, 11937 US Hwy 271, Tyler Texas 75708, USA; [2]Institute of Health Sciences, China Medical University, Shenyang 110122, Liaoning, China; [3]Department of Stem Cells and Regenerative Medicine, College of Basic Medical Science, China Medical University, Shenyang 110122, Liaoning, China; [4]Texas Lung Injury Institute, University of Texas Health Science Center at Tyler, Tyler Texas 75708, USA

e-mail: jianjun_chang@yeah.net, zaixingchen@sina.com, runzhen.zhao@uthct.edu, hgnie@cmu.edu.cn, james.ji@uthct.edu

[&]Present address: School of Pharmacy, China Medical University, Shenyang 110122, Liaoning, China

[*]Corresponding author.

Acknowledgments: This work was supported by the grants from the National Institute of Health (NIH HL134828), and the National Natural Science Foundation of China (NSFC 81670010).

Word count: Title 10, Abstract 248, Main text word count 6715

Acrolein

CFTR

Cl⁻

ENaC

Na⁺

2K⁺

Na/K-ATPase

3Na⁺

Apical

Basolateral

Graphical Headlights

- Thermal stress and acrolein are two essential determinants for smoke-inhalation injury, impairing airway epithelial barrier.

- Transcellular ion transport pathways via the ENaC, CFTR, and Na/K-ATPase are interrupted by both thermal stress and acrolein, one of the most potent smoke toxins.

- Heat and acrolein damage the integrity of the airway epithelium through suppressing and relocating the tight junctions.

Abstract Smoke inhalation injury is the leading cause of death in firefighters and victims. Inhaled hot air and toxic smoke are the predominant hazards to the respiratory epithelium. We aimed to analyze the effects of thermal stress and smoke aldehyde on the permeability of the airway epithelial barrier. Transepithelial resistance (R_{TE}) and short-circuit current (I_{SC}) of mouse tracheal epithelial monolayers were digitized by an Ussing chamber setup. Zonula occludens-1 tight junctions were visualized under confocal microscopy. A cell viability test and fluorescein isothiocyanate-dextran assay were performed. Thermal stress (40°C) decreased R_{TE} in a two-phase manner. Meanwhile, thermal stress increased I_{SC} followed by its decline. Na^+ depletion, amiloride (an inhibitor for epithelial Na^+ channels [ENaCs]), ouabain (a blocker for Na^+/K^+-ATPase) and CFTRinh-172 (a blocker of cystic fibrosis transmembrane regulator [CFTR]) altered the responses of R_{TE} and I_{SC} to thermal stress. Steady-state 40°C increased activity of ENaCs, Na^+/K^+-ATPase, and CFTR. Acrolein, one of the main oxidative unsaturated aldehydes in fire smoke, eliminated R_{TE} and I_{SC}. Na^+ depletion, amiloride, ouabain, and CFTRinh-172 suppressed acrolein-sensitive I_{SC}, but showed activating effects on acrolein-sensitive R_{TE}. Thermal stress or acrolein disrupted zonula occludens-1 tight junctions, increased fluorescein isothiocyanate-dextran permeability but did not cause cell death or detachment. The synergistic effects of thermal stress and acrolein exacerbated the damage to monolayers. In conclusion, the paracellular pathway mediated by the tight junctions and the transcellular pathway mediated by active and passive ion transport pathways contribute to impairment of the airway epithelial barrier caused by thermal stress and acrolein.

Keywords: thermal stress; acrolein; tracheal epithelial monolayers; ion transport; tight junctions

1   Introduction

Fire is a common disaster in urban and rural populations. Smoke is a complicated mixture of hot air, toxic chemicals, and suspended particles (Fitzgerald and Flood 2006). Inhalation of harmful smoke causes direct impairment of respiratory function and is the leading cause of fire-related death (Haponik 1993). Inhaled hot air and toxic chemicals in smoke from fire impair the respiratory tract and lungs.

The airway epithelium is the most important defense barrier in the respiratory tract (Wang et al. 2008). Approximately 80% of the cells are ciliated in the normal respiratory tract. These cells form a tight airway epithelial layer, which can effectively prevent various external attacks (Hou et al. 2019). Destruction of the integrity of the airway epithelial barrier can be an important indicator of occurrence and the development of many respiratory diseases (Wang et al. 2007). Under pathological conditions, the integrity of the airway epithelial barrier is disrupted and its permeability is increased. Permeability of a tight airway epithelium is determined by paracellular and transcellular pathways. Paracellular pathway is predominantly regulated by tight junctions and lateral intercellular space. Transcellular permeability is predominantly controlled by active and passive ion transport pathways. Apical epithelial Na^+ channels (ENaCs) and cystic fibrosis transmembrane regulator (CFTR), as well as basolateral Na^+/K^+-ATPase, are mainly responsible for Na^+/Cl^- transport in the respiratory epithelium (Chang et al. 2018; Wang et al. 2007). In respiratory epithelium, passive ion transport mediated by ENaCs is an important determinant of transcellular permeability, and active Na^+/K^+-ATPase provides the driving force for passive ion transport systems.

A previous study showed the mats of ciliated cells in rat tracheal epithelium post-thermal injury (Dubick et al. 2002). Thermal stress regulates ENaCs in alveolar epithelial cells (Howard et al. 2013) and CFTR is implicated in thermal stress-induced signaling events in bronchial epithelial cells (Dong et al. 2015). Additionally, Na^+/K^+-ATPase is heat-sensitive in rabbit esophageal epithelium (Tobey et al. 1999). However, whether thermal stress regulates permeability of the airway epithelial barrier through the paracellular pathway (mediated by tight junctions) and the transcellular pathway (mediated by ion channels and transporters) in primary airway epithelial cells remains unclear.

Smoke aldehydes are released from the burning of wood and paper products, as well as cotton fabrics. Formaldehyde and acrolein

are the two main gaseous α, β unsaturated aldehydes in fire and cigarette smoke (Reinhardt and Ottmar 2004; Anthony et al. 2007; Alwis et al. 2015). Inhalation of formaldehyde and acrolein results in gene induction, inflammation, cell apoptosis, and necrosis in the respiratory epithelium (Bein and Leikauf 2011; Meacher and Menzel 1999). Our previous studies showed that formaldehyde and crotonaldehyde inhibited the activity and expression of ENaCs in alveolar epithelial cells (Cui et al. 2016; Li et al. 2017). Acrolein led to irritation of the airway and impaired fluid homeostasis (Borchers et al. 1998; Romet-Haddad et al. 1992; Roux et al. 2002), as well as downregulated CFTR activity in respiratory epithelial cells (Alexander et al. 2012). Acrolein is used to prepare animal models for acute lung injury and lung edema (Hales et al. 1988). However, the pathological role of acrolein in the presence of thermal stress in the smoke-injured airway epithelial barrier has not been studied.

Therefore, this study aimed to investigate the effects of thermal stress or/and acrolein on permeability of the airway epithelial barrier. We recorded the transepithelial resistance (R_{TE}) and short-circuit current (I_{SC}) in primary mouse tracheal epithelial (MTE) monolayers in an Ussing chamber setup. To further determine the underlying transcellular mechanisms, inhibitors for ion channels and transporters (i.e., amiloride, ouabain, and CFTRinh-172) were applied to MTE monolayers. To examine the role of the paracellular pathway, the tight junctions were visualized under confocal microscopy, and the fluorescein isothiocyanate (FITC)-dextran assay was performed.

Materials and methods

Animals

Forty healthy wild-type C57BL/6 mice (males, 20; females, 20), 8-12 weeks old (mean: 10.86 ± 0.28 weeks) and weighed 18-25 g (mean: 20.03 ± 0.30 g), were purchased from the Jackson Laboratory and the Laboratory Animal Center of China Medical University. Mice were housed in a pathogen-free facility, the husbandry condition of which included a suitable light/dark cycle, temperature, drinking water, and food. All experiments were performed according to the guidelines and regulations of the Animal Care and Use Ethics Committee, and all protocols were approved by the University of Texas Health Science Center at Tyler and China Medical University. The reference numbers of ethics approval are IACUC 611 and SCXK (Liao) 2018-0001.

Isolation and culture of MTE cells

Pooled MTE cells were isolated from C57BL/6 mice and cultured according to our previous study (Chen et al. 2014). Briefly, tracheas of anesthetized mice were removed, and cleaned tracheas were incubated in Dulbecco's modified eagle medium (cat# 30-2002, American Type Culture Collection, USA) containing 0.1% protease XIV (cat# P5147, Sigma, USA), 0.01% DNase (cat #DN25, Sigma-Aldrich, St. Louis, MO), and 1% fetal bovine serum (cat# 26140-087, Gibco, USA) at 4°C for 24 h. MTE cells were seeded onto 6.5-mm diameter, collagen IV (cat# 3410-010-01, Trevigen, Gaithersburg, MD)-coated transwell inserts (cat# 3413, Corning-Costar, Lowell, MA) at a density of $3.0 \times 10^5/cm^2$. These cells were grown in a 1:1 mixture of Ham's F-12 medium (cat# 11765-054, Invitrogen, Camarillo, CA) and 3T3 fibroblast preconditioned Dulbecco's modified eagle medium supplemented with insulin (10 μg/ml, cat# I1882, Sigma-Aldrich, St. Louis, MO), hydrocortisone (1 μM, cat# H0396, Sigma-Aldrich, St. Louis, MO), endothelial cell growth supplement (3.75 μg/ml, cat# E0760, Sigma-Aldrich, St. Louis, MO), epidermal growth factor (25 ng/ml, cat# E4127, Sigma-Aldrich, St. Louis, MO), triiodothyronine (30 nM, cat# T6397, Sigma-Aldrich, St. Louis, MO), iron-saturated transferrin (5 μg/ml, cat# T1283, Sigma-Aldrich, St. Louis, MO), cholera toxin (10 ng/ml, cat# C8052, Sigma-Aldrich, St. Louis, MO), and dexamethasone (250 nM, cat# D2915, Sigma-Aldrich, St. Louis, MO). MTE cells from one mouse were seeded onto three to four transwell inserts and cultured for up to 12 days. Polarized monolayers with a reading of R_{TE} >1000 Ω by an epithelial voltohmmeter (WPI, Sarasota, FL) were used. Monolayers from the same batch of mice were randomly allocated to control and treated groups.

Preparation of human bronchial epithelial monolayers

Human bronchial epithelial (HBE) cells were purchased from American Type Culture Collection. Cells were seeded in plastic T-75 flasks and grown in keratinocyte culture medium (cat# 2101, Sciencell, China) supplemented with keratinocyte growth factor (1%). The culture medium was changed every 48 h until 90% confluent. HBE cells were seeded (10^6 cells/cm^2) onto collagen IV-coated transwell inserts and grown in keratinocyte culture medium supplemented with keratinocyte growth factor (1%), insulin-transferrin-selenium solution, (1.5%, cat# 41400045, Gibco, USA) and dexamethasone (1 μM, cat# D2915, Sigma-Aldrich, St. Louis, MO). Polarized monolayers at 12-day post-seeding were used. At all stages of culture, cells were maintained at 37°C in 5% CO_2 in an air incubator.

Preparation and administration of ion transport regulators

Amiloride (an inhibitor for ENaCs, cat# A4562, Sigma-Aldrich, St. Louis, MO) was reconstituted in an apical bath solution containing 50% dimethyl sulfoxide (DMSO). CFTRinh-172 (an inhibitor for CFTR, cat# C2992, Sigma-Aldrich, St. Louis, MO) was reconstituted in DMSO. Ouabain (an inhibitor for Na^+/K^+-ATPase, cat# O3125, Sigma-Aldrich, St. Louis, MO) was reconstituted in a basolateral bath solution. Reconstituted amiloride (100 μM) or CFTRinh-172 (20 μM) was added to the apical bath solution of MTE monolayers. Reconstituted ouabain (1 mM) was added to the basolateral bath solution. Vehicle solution of an equal volume was added to the corresponding bath solution as the control.

Concurrent measurements of I_{SC} and R_{TE}

Measurement of I_{SC} and R_{TE} of MTE or HBE monolayers was performed as we described previously (Han et al. 2010; Nie et al. 2009; Li et al. 2017). In brief, monolayers were mounted into Ussing chambers (Physiologic Instruments, San Diego, CA) and *bathed in* saline solution containing (in mM) 120 NaCl, 25 $NaHCO_3$, 3.3 KH_2PO_4, 0.83 K_2HPO_4, 1.2 $CaCl_2$, 1.2 $MgCl_2$, 10 HEPES, 10 D-mannitol (apical), and 10 D-glucose (basolateral). The saline bath solution was bubbled continuously with a gas mixture of 95% O_2 and 5% CO_2. Transepithelial potential was short-circuited to 0 mV, and R_{TE} and I_{SC} levels were measured with an epithelial voltage clamp (VCC-MC8, Physiologic Instruments, San Diego, CA). A 10-mV pulse of 1 s duration was imposed every 10 s to monitor R_{TE} levels. When R_{TE} and I_{SC} levels were stable for at least 10 min, monolayers were treated as follows: 1) amiloride (100 μM) was added to the apical bath solution; 2) CFTRinh-172 (20 μM) was added to the apical bath solution; 3) ouabain (1 mM) was applied to the basolateral bath solution; and 4) saline bath solution of both apical and basolateral chambers was replaced with Na^+-free bath solution or Cl^--free bath solution. The same volume of vehicle solution was added to corresponding chambers, which was used as the control of amiloride, CFTRinh-172, or ouabain. The composition of the Na^+-free bath solution contained (in mM) 120 N-methyl-D-glucamine-Cl, 25 $KHCO_3$, 3.3 KH_2PO_4, 0.83 K_2HPO_4, 1.2 $CaCl_2$, 1.2 $MgCl_2$, 10 HEPES, 10 D-mannitol (apical), and 10 D-glucose (basolateral). The Cl^--free bath solution contained (in mM) 120 sodium gluconate, 25 $NaHCO_3$, 3.3 KH_2PO_4, 0.83 K_2HPO_4, 1.2 $Ca(NO_3)_2$, 1.2 $MgSO_4$, 10 HEPES, 10 D-mannitol (apical), and 10 D-glucose (basolateral). Before replacement of Na^+-free or Cl^--free bath solution, recording of R_{TE} and I_{SC}

in the Ussing chamber system was stopped, and then saline bath solution was aspirated from the apical and basolateral chambers. Warm

saline (as the control), Na^+-free bath solution, or Cl^--free bath solution was then added to the chambers. Finally, recording in the Ussing

chamber system was restored.

Application of thermal stress or acrolein to monolayers

All experiments of thermal stress or acrolein application were simultaneously performed with recording in the Ussing chamber system.

In the experiments for studying the effects of thermal stress on R_{TE} and I_{SC}, and the underlying mechanisms, monolayers were mounted

into the chambers in which the temperature was 37°C. The temperature of bath solution of the apical and basolateral chambers was

gradually elevated to the designed degree (40°C) from 37°C. A bath circulator (Thermo Fisher Scientific, CA) was used to regulate the

temperature of the chambers. The bath circulator was set to the desired temperature. Water was pumped from the bath of the circulator

and continuously recirculated around the chambers, thereby transferring heat to the bath solution in the chambers. Additionally,

thermometers in the apical and basolateral bath solution of the chambers were used to monitor and confirm the temperature setting. In

the experiments for investigating the effects of thermal stress on ion transport system, the bath solution in the chambers was first set to

the desired temperature (40°C), and then monolayers were mounted into chambers to achieve instant exposure of thermal stress. In the

experiments of acrolein exposure, a range of working concentrations from 5 to 500 μM was accumulatively pipetted into the apical bath

solution. Acrolein vapor was minimized without gas bubbling of the bath solution, and the bath solution was moderately blown up to 10

times to ensure even distribution of acrolein. To study the synergistic effects of acrolein and thermal stress, monolayers were mounted

into chambers in which the temperature was already 40°C or 37°C (as the control). When R_{TE} and I_{SC} traces were stable, a series of

concentrations of acrolein were added to the apical bath solution.

Immunofluorescent staining and imaging of MTE monolayers

MTE monolayers that were *bathed in* saline solution in Ussing chambers were exposed to 40°C (120 min) or acrolein (500 μM). For

synergistic treatment of thermal stress and acrolein, MTE monolayers were *bathed in* saline solution (40°C or 37°C), and then 500 μM

acrolein was added to the apical bath solution. After this treatment, monolayers were prepared for staining. Monolayers were fixed in

phosphate-buffered saline containing 4% paraformaldehyde for 15 min and washed with phosphate-buffered saline. Monolayers were

permeabilized with 0.5% Triton X-100 and blocked with a blocking buffer containing 10% goat serum (cat# G9023, Sigma-Aldrich, St.

Louis, MO) and 1% bovine serum albumin (cat# A3803, Sigma-Aldrich, St. Louis, MO) for 1 h. Monolayers were incubated with rabbit

anti-mouse zonula occludens-1 (ZO-1) antibody (1:50; cat# 61-7300, Invitrogen, Camarillo, CA) or rabbit anti-mouse occludin antibody

(1:200; cat# 40-4700, Invitrogen, Camarillo, CA) in blocking buffer overnight at 4°C. Cells were incubated with Alexa Fluor 488-

labeled goat anti-rabbit IgG (1:1000, cat# 111-545-045 Jackson Immuno Research Laboratories, Carlsbad, CA) and Hoechst (1:1000,

cat# 561908, BD Biosciences, San Jose, CA) and then mounted with Vectashield mounting media (cat# H1000, Vector Laboratories,

Burlingame, CA). Images were obtained with a Zeiss LSM 510 confocal microscope (Carl Zeiss AG, Germany). A series of optical

sections were collected at 1-μm intervals in the Z-axis. Images and quantification of fluorescence intensity were analyzed with ImageJ

(NIH).

Cell counting kit-8 assay

Cell viability was quantified by the cell counting kit-8 assay (cat# C0037, Beyotime, Jiangsu, China). Culture medium containing 10%

cell counting kit-8 solution was added apically to MTE monolayers at post-treatment of 37°C (control), 40°C (120 min), acrolein (500

μM), or 40°C combined with acrolein. After 2 h of incubation at 37°C, culture medium in transwell inserts was collected. Absorbance

was measured at a wavelength of 450 nm.

FITC‑dextran assay

To measure dextran permeability, FITC-dextran (4 kDa, 1 mg/ml, cat# FD4, Sigma-Aldrich, St. Louis, MO) was added to the transwell

inserts. Aliquots were withdrawn from the lower chambers after 4 h and assayed for fluorescence at 530 nm.

Statistical analysis

Data are expressed as the mean ± SEM. Normality and homoscedasticity tests were performed by the Shapiro-Wilk and Levene tests.

The real power of the sample size with alpha = 0.05 is shown in the figure legends. The student's two-tailed t-test was used for comparing

two groups with parametric data. For comparison of multiple groups, we performed one-way analysis of variance followed by

Bonferroni's test for all groups in the experiments. For non-parametric data, the Mann-Whitney U test was used. Differences were

considered significant when the *P* value was < 0.05. The $\tau_{1/2}$ value was computed by fitting R_{TE} raw data with the ExpDec1 function

($y = y_0 + Ae^{-x/t}$), where y is R_{TE}, x is time, y_0 is offset; and A is amplitude. The dose-response curve for acrolein was generated

by fitting the raw data with the Hill equation ($y = START + (END - START)\frac{x^n}{k^n + x^n}$), where y is the normalized R_{TE}/I_{SC} value, x is

the logarithmic concentration of acrolein, *START* and *END* are initial and ending values of R_{TE}, respectively, k is the K_i value, and n is

the total number of x. In thermal stress-treated R_{TE}/I_{SC} traces, $\triangle R_{TE}$ was defined as the difference between the initial R_{TE} and end of

R_{TE} within one phase (phase 1 [P1] or phase 2 [P2]) or the entire procedure, and $\triangle I_{SC}$ was defined as the difference between the basal

I_{SC} and the peak I_{SC} at P1 or the difference between the peak I_{SC} and end of I_{SC} at P2. In acrolein-treated R_{TE}/I_{SC} traces, $\triangle R_{TE}/\triangle I_{SC}$ was

defined as the difference between basal R_{TE}/I_{SC} and acrolein-resistant R_{TE}/I_{SC}. ENaCs, CFTR, or Na^+/K^+-ATPase activity was determined

as the difference between the basal I_{SC} and corresponding specific inhibitor-resistant I_{SC}. Statistical analysis was performed with Origin

Pro 8.0 (OriginLab Corp. MA).

Results

Thermal stress alters R_{TE} and I_{SC} in MTE and HBE monolayers

The airway epithelial barrier is the primary line of defense against external attacks. Under pathogenic conditions, the integrity of this

barrier is damaged and permeability is increased. We postulated that thermal stress would regulate permeability of the airway epithelium.

In our study, basal R_{TE} and I_{SC} in polarized MTE monolayers were recorded at 37°C. To investigate the effect of thermal stress, the

temperature of the bath solution was elevated from 37°C. R_{TE} and I_{SC} were eliminated in minutes when the temperature of the bath

solution was elevated to 44°C or 42°C (Fig. S1). Therefore, 40°C was selected for the temperature of thermal stress. R_{TE} was decreased

when the temperature of the bath solution was raised to 40°C from 37°C. Moreover, the entire process was divided into two stages of

P1 and P2 (Fig. 1a). At each phase, R_{TE} was significantly decreased in a time-dependent manner (Fig. 1c). We used $\tau_{1/2}$ to reflect the

declining rate of R_{TE} within one phase, which was computed by fitting R_{TE} raw data with the ExpDec1 function. The $\tau_{1/2}$ at P2 was

significantly higher than that at P1 (Fig. 1e, $P < 0.001$). We defined P1 and P2 of I_{SC} using the R_{TE} time course. In contrast, I_{SC}, which

was recorded in parallel with R_{TE}, was increased at P1 and decreased at P2 (Fig. 1b and 1d, all $P < 0.01$). To overcome the difference

between species, we recorded R_{TE} and I_{SC} in HBE monolayers during thermal stress. We found that R_{TE} of HBE monolayers was

decreased by thermal stress, which was similar to the result from MTE monolayers (Fig. 1f and 1h, $P = 0.077$). However, I_{SC} of HBE

monolayers was increased to a high and stable level by thermal stress (Fig. 1g and 1i, $^*P < 0.05$). Our data suggest that R_{TE} and I_{SC} of

primary MTE monolayers respond to thermal stress in a phase-dependent, diverse manner.

Na^+ transport mediates thermal stress-induced bioelectric changes

In respiratory epithelium, Na^+ transport is mainly regulated by ENaCs and Na^+/K^+-ATPase (Chang et al. 2018). Passive ion transport

across tight epithelial monolayers includes ion channels (i.e., ENaCs) and transporters, which are an important determinant of

transcellular permeability. Active Na^+/K^+-ATPase provides the driving force for the passive ion transport systems. To examine the role

of epithelial Na^+ transport in thermal stress-altered R_{TE} and I_{SC}, Na^+-free bath solution, amiloride, or ouabain was applied to MTE

monolayers *bathed in* Ussing chambers at 37°C followed by elevation to 40°C. At 37°C, Na^+-free bath solution and amiloride, but not

ouabain, caused an increment in R_{TE} (Fig. 2a, black line). A drop in I_{SC} was seen at this time (Fig. 2b, black line). During thermal stress,

R_{TE} was decreased in a two-phase manner in the presence of amiloride, which was similar to the result from the saline group. In contrast,

a continuous decrease in R_{TE} was observed in the presence of Na^+-free bath solution or ouabain, which could not be divided into two

significant phases (Fig. 2a, red line). I_{SC} at P1 and P2 was observed in the presence of Na^+-free bath solution or amiloride (Fig. 2b, red

line). We computed thermal stress-sensitive R_{TE} at P1 and P2 ($\triangle R_{TE}$, in Fig. 2c), and found that amiloride markedly increased thermal

stress-sensitive R_{TE} at P1 ($P < 0.01$), but there were no significant effects at P2 ($P = 0.56$). Thermal stress caused the continuous reduction

of R_{TE} in the presence of the Na^+-free bath solution and ouabain, so we calculated thermal stress-sensitive R_{TE} in the entire procedure

($\triangle R_{TE}$ in Fig. 2d), and found that Na^+-free bath solution markedly increased thermal stress-sensitive R_{TE} and ouabain significantly

decreased thermal stress-sensitive R_{TE} in the entire of procedure (all $P < 0.05$). We calculated thermal stress-sensitive I_{SC} at P1 and P2

($\triangle I_{SC}$, in Fig. 2e and 2f), and found that thermal stress-sensitive I_{SC} at P1 was significantly decreased in the presence of Na^+-free bath

7   solution, amiloride, or ouabain (Fig. 2e, all $P < 0.05$). Additionally, Na^+-free bath solution and amiloride markedly decreased thermal

8   stress-sensitive I_{SC} at P2 (Fig. 2f, $P < 0.001$). Because we could not observe the obvious I_{SC} P2 in the presence of ouabain, we did not

9   calculate thermal stress-sensitive I_{SC} at P2 in the presence of ouabain. These data indicate that changes in R_{TE} and I_{SC} caused by thermal

stress may be predominately determined by transcellular Na^+ transport powered by ENaCs and Na^+/K^+-ATPase. Interestingly, blockade

of Na^+/K^+-ATPase showed lower thermal stress-sensitive I_{SC} compared with Na^+-free bath solution. The potential mechanisms of this

finding may be the temperature-dependent Na^+/K^+-ATPase and others, including ENaCs, CFTR, and K^+ channels (KCs). We attempted

to further confirm the effects of thermal stress on permeabilized half monolayers. However, neither R_{TE} nor I_{SC} levels were stable enough

to perform further analysis (Fig. S2).

CFTR is required for thermal stress-induced alteration

We next examined the contribution of epithelial Cl^- transport to thermal stress-induced alterations in R_{TE} and I_{SC} of MTE monolayers.

Transcellular Cl^- transport via apical CFTR comprises majority of transcellular anion flux (Sheppard and Welsh 1999; Fuller and Benos

1992). At 37°C, Cl^--free bath solution or CFTRinh-172 was applied to block overall epithelial Cl^- transport or CFTR. Blockade of CFTR

led to an increment in R_{TE} and a drop in I_{SC} (Fig. 3a and 3b, black line). However, Cl^--free bath solution caused a transient increment,

followed by a slow decline to 0.24 ± 0.03 k$\Omega \times cm^2$ in R_{TE} and almost zero in I_{SC} (Fig. S3). Therefore, we could not perform further

analysis. During thermal stress, R_{TE} and I_{SC} were changed in a two-phase manner in the presence of CFTRinh-172, which was similar

to the saline group (Fig. 3a and 3b, red line). CFTRinh-172 significantly increased thermal stress-sensitive R_{TE} for both phases (Fig. 3c,

all $P < 0.01$). CFTRinh-172 significantly decreased thermal stress-sensitive I_{SC} for both phases (Fig. 3d and 3e, $P < 0.05$). These

findings suggest that CFTR plays a role in changes in R_{TE} and I_{SC} caused by thermal stress.

Steady-state elevated temperature increases transcellular ion channel activity

To confirm the two-phase alteration in I_{SC} following a gradual rise in bath temperature, we alternatively mounted MTE monolayers to

Ussing chambers that were already set at 40°C. Interestingly, elevated and stable total I_{SC} levels were observed under this steady-state,

elevated temperature condition, which was different from the two-phase manner when the temperature was gradually elevated (Fig. 4a,

4c, and 4e, basal I_{SC} of the black line and red line). These observations suggested that the time course of thermal stress determined the

severity of impaired transcellular ion transport in MTE monolayers. To inhibit the activity of ion channels and transporters, including

ENaCs, CFTR, and Na^+/K^+-ATPase, their specific inhibitors, including amiloride, CFTRinh-172, and ouabain, were added to a bath

solution at 37°C or 40°C. I_{SC} was inhibited by these specific inhibitors at 37°C and 40°C (Fig. 4a, 4c, and 4e). We computed total I_{SC},

specific inhibitor-resistant I_{SC}, and the difference between these two I_{SC} (Figs. 4b, 4d, and 4f). The difference between total I_{SC} and

specific inhibitor-resistant I_{SC} reflected the activity of corresponding ion channels or transporters, and we found that thermal stress

increased the activity of ENaCs, CFTR and Na^+/K^+-ATPase (all $P < 0.05$). We also computed the basal R_{TE} value and post-inhibitor

value applying at 37°C or 40°C (Fig. 4g). The mean basal R_{TE} level at 40°C (0.57 ± 0.05 k$\Omega \times cm^2$) was significantly less than that at

37°C (1.08 ± 0.06 k$\Omega \times cm^2$) ($P < 0.001$). An increase in R_{TE} was observed in the presence of amiloride and CFTRinh-172 (all $P < 0.05$),

but not ouabain (all $P > 0.05$), at 37°C and 40°C. These data suggest that a steady-state elevated temperature increases the activity of

ENaCs, CFTR, and Na^+/K^+-ATPase.

Acrolein downregulates R_{TE} and I_{SC}

Our previous studies showed that formaldehyde and crotonaldehyde regulated ENaCs in alveolar epithelial cells (Cui et al. 2016; Li et

al. 2017). Acrolein is an industrial chemical of high toxicity and a toxic combustion product (Stevens and Maier 2008), and

concentrations of which can vary widely in different fire and tobacco smoke. One kilogram of different types of wood can produce

0.374-2.35 mM of acrolein (Faroon et al. 2008). In the case of burning of tobacco, the acrolein concentrations are 34-503 μM when

smoke from a single cigarette is bubbled through 10 ml of buffered saline (Burcham et al. 2010). To examine the effects of acrolein on

the permeability of MTE monolayers, acrolein (approximately 500 μM) was applied to monolayers bathed at 37°C and at a steady-state

of 40°C. A total of 500 μM acrolein decreased R_{TE} and I_{SC} at 37°C and 40°C (Fig. 5a and 5b). To compare the difference in effects of

acrolein at 37°C and 40°C, the dose-response curve for acrolein was generated by fitting raw data points with the Hill equation. By

fitting raw R_{TE} data, we found that the K_i value for acrolein was 120.22 μM at 37°C and 79.43 μM at 40°C (Fig. 5c). By fitting raw I_{SC}

data, we found that the K_i value for acrolein was 125.89 μM at 37°C and 77.62 μM at 40°C (Fig. 5d). In HBE monolayers, a reduction

of R_{TE} and I_{SC} caused by 500 μM acrolein was observed, which was similar to the result from MTE monolayers (Fig. 5e and 5f). In the

R_{TE} dose-response curve of HBE monolayers, the K_i value for acrolein was 83.18 μM at 37°C and 97.73 μM at 40°C (Fig. 5g). In the

I_{SC} dose-response curve of HBE monolayers, the K_i value for acrolein was 125.89 μM at 37°C and 112.21 μM at 40°C (Fig. 5h). Our

data suggest that acrolein impairs airway epithelial permeability in an atypical concentration-dependent manner and an elevated

temperature exacerbates this damaging process.

Na^+ transport participates in acrolein-induced reduction of R_{TE} and I_{SC}

We examined the role of epithelial Na^+ transport on the decrease in R_{TE} and I_{SC} by acrolein in MTE monolayers. At the basal level of

37°C, Na^+-free bath solution, amiloride, or ouabain was used to block overall epithelial Na^+ transport, ENaCs or Na^+/K^+-ATPase. After

pre-inhibition of these Na^+ transport pathways, 500 μM acrolein was added to the apical bath solution, and changes in R_{TE} and I_{SC} caused

by acrolein were observed. At 37°C, Na^+-free bath solution and amiloride, but not ouabain, caused an increment in R_{TE} (Fig. 6a, black

line). Additionally, a drop in I_{SC} induced by Na^+-free bath solution, amiloride, and ouabain, was observed (Fig. 6b, black line). During

exposure of 500 μM acrolein, R_{TE} was decreased in the presence of Na^+-free bath solution, amiloride, and ouabain (Fig. 6a, blue line).

Acrolein continuously decreased I_{SC} in the presence of amiloride but showed no obvious effect in the presence of Na^+-free bath solution

and ouabain (Fig. 6b, blue line). We calculated acrolein-sensitive R_{TE} ($\triangle R_{TE}$, Fig. 6c), and found that Na^+-free bath solution, amiloride,

and ouabain significantly increased acrolein-sensitive R_{TE} ($P < 0.05$). Acrolein-sensitive I_{SC} ($\triangle I_{SC}$, Fig. 6d) was also computed, and it

was significantly decreased by Na^+-free bath solution, amiloride, and ouabain (all $P < 0.01$). These data suggest that changes in R_{TE} and

I_{SC} induced by acrolein may be related to transcellular pathways that are mediated by ENaCs and Na^+/K^+-ATPase.

CFTR mediates the acrolein-induced reduction in R_{TE} and I_{SC}

We further examined the role of CFTR in the decrease in R_{TE} and I_{SC} by acrolein in MTE monolayers. At basal level of 37°C, CFTRinh-

0 172 was used to inhibit CFTR, and then 500 μM acrolein was added to the monolayers. Changes in R_{TE} and I_{SC} caused by acrolein were

1 observed. At the basal level, CFTRinh-172 increased R_{TE} (Fig. 7a, black line) and decreased I_{SC} (Fig. 7b, black line). During exposure

2 of acrolein, R_{TE} and I_{SC} were decreased (Figs. 7a and 7b, blue line). CFTRinh-172 significantly increased acrolein-sensitive R_{TE} (Fig.

7c, $P < 0.01$) and markedly decreased acrolein-sensitive I_{SC} (Fig. 7d, $P < 0.001$). These findings suggest that CFTR may participate in the reduction of R_{TE} and I_{SC} caused by acrolein.

Thermal stress and acrolein impair tight junctions

Our findings suggested that the transcellular pathway is required for the effects of thermal stress and acrolein on permeability of MTE monolayers. Therefore, we next examined if thermal stress and acrolein affect the paracellular pathway. We observed the structure and distribution of the tight junction protein ZO-1 in MTE monolayers. In MTE monolayers, ZO-1 is one of the most widely studied scaffold proteins and is uniformly expressed after an air-liquid interface is achieved (Umeda et al. 2006). Confocal microscopy showed that thermal stress disrupted the integrity of ZO-1 (Fig. 8e) compared with the control group (Fig. 8b). Damage to the structure of ZO-1 was also observed in acrolein-treated monolayers (Fig. 8h). Moreover, synergistic treatment of thermal stress and acrolein further exacerbated the damage to ZO-1 (Fig. 8k) compared with thermal stress (Fig. 8e) or acrolein group (Fig. 8h). We quantified fluorescence intensity and it was consistent with that from the representative confocal images (Fig. 8m). To determine the effects of thermal stress and acrolein on cell detachment and death of MTE monolayers, we tested cell number and viability of monolayers post-thermal stress and after acrolein. We found no significant cell detachment or death under these conditions (Fig. S4, all $P > 0.05$). We examined expression of occludin and claudins, but their expression levels were insufficient for visualization (Fig. S6). The FITC-dextran assay was performed to further determine the effects of thermal stress and acrolein on paracellular permeability of MTE monolayers. We found that thermal stress and acrolein significantly increased the permeability of MTE monolayers to FITC-dextran (Fig. S6p, $P < 0.001$).

Discussion

Smoke inhalation injury is the main cause of death during a fire, and it is an important risk factor for increasing morbidity and mortality (You et al. 2014). Pathogenic factors in inhalation injury, such as heat and toxic chemicals, cause direct injury to the airway epithelial barrier. In this study, we aimed to investigate the effects of thermal stress and acrolein on permeability of the airway epithelial barrier. MTE cells were isolated from mouse trachea and used to culture the model of the air-liquid interface (MTE monolayers). This

in vitro model mainly contained goblet and cilia cells and imitated the physiological state of the normal airway epithelium (Hou et al. 2019). MTE monolayers are suitable for studying respiratory diseases, and they have been applied in toxicology, respiratory tract infection, ion transport, cell carcinogenesis, and other experiments (Davidson et al. 2000; Horani et al. 2013). In our study, R_{TE} and I_{SC} were markedly changed in thermal stress or acrolein-treated MTE monolayers. The changes in R_{TE} and I_{SC} caused by thermal stress or acrolein were related to the transcellular pathway, which was mediated by ENaCs, CFTR, and Na^+/K^+-ATPase. Additionally, thermal stress or acrolein regulated the paracellular pathway, which was mediated by tight junctions. These findings suggest the potential roles of the transcellular and paracellular pathways in thermal stress or acrolein in damaging the airway epithelial barrier. Our findings may provide new directions for the understanding, diagnosis, and treatment of smoke inhalation injury.

Transcellular and paracellular pathways are two essential determinants of R_{TE} and I_{SC}. In the current study, during thermal stress, R_{TE} and I_{SC} of MTE monolayers were changed in a two-phase (P1 and P2) manner. I_{SC} was increased at P1, which could have been due to enhanced ion transport by thermal stress. Concurrently, increased permeability led to a reduction in R_{TE}. The reverse change in R_{TE} and I_{SC} indicated that monolayers at P1 still had normal physiological characteristics. Subsequently, R_{TE} and I_{SC} were decreased at P2. These findings could have resulted from damaged gap and tight junctions, which were shown by confocal images. Another explanation for our findings is ion diffusion across MTE monolayers upon depletion of ATP. Therefore, we speculate that effects caused by thermal stress are reversible at P1 and irreversible at P2. A previous study provided the basis for our speculation (Tobey et al. 1999) in which increased I_{SC} and decreased R_{TE} were observed during application of heat to the esophageal epithelium, and recovery was observed when the temperature returned to 37°C. Cell death and detachment cause increased permeability of monolayers. However, our data from cell counting and viability ruled out this possibility. To overcome the difference between species, we cultured HBE monolayers and monitored R_{TE} and I_{SC}. We found that R_{TE} of HBE monolayers was decreased by thermal stress, which was similar to the result from MTE monolayers. In contrast, thermal stress increased I_{SC} of HBE monolayers to a high and stable level, which was similar to P1 in MTE monolayers. However, we could not monitor the decreased trace of I_{SC} (i.e., P2 in MTE monolayers). These findings indicated that HBE monolayers showed lower thermal sensitivity and higher thermal resistance than primary MTE monolayers.

7   Various time courses of thermal stress caused obvious different regulation in I_{SC} in our study. We observed an elevated I_{SC} trace at

8   P1 and a declined trace at P2 when the temperature was gradually raised to 40°C from 37°C. However, under the condition of a steady-

9   state at 40°C, an elevated and stable I_{SC} trace was visualized instead of two phases. In a previous study, researchers found elevated stable

I_{SC} levels of rabbit esophageal epithelium in an Ussing chamber system in which the temperature had already been set to 49°C (Tobey

et al. 1999). These divergent responses in I_{SC} and R_{TE} to different time courses of thermal stress could be due to the response of the

individual transport system to thermal stress. Host cell adaption to environmental stress could explain the diverse observations between

the two procedures of thermal stress.

The paracellular and transcellular pathways are vital for establishing or dissipating ion concentration gradients and thus are important

for determining the ionic composition of the apical compartment and net volume flow (Flynn et al. 2009). Therefore, both pathways

work in concert and are functionally matched to meet the transport requirements of the specific tissue. The paracellular pathway is

mainly formed by tight junctions, and this pathway is located near the apical side of the cells and the lateral intercellular space (Tsukita

and Furuse 2002; Van Itallie and Anderson 2006). Tight junctions form the functional and structural boundary that separates apical and

basolateral compartments and also determine the ion transport properties of the paracellular pathway. Tight junctions are composed of

a complex of proteins that determine ion selectivity and conductance of the paracellular pathway (Farquhar and Palade 1963). ZO-1 is

one of the most widely studied tight junction protein (Umeda et al. 2006) and is uniformly expressed after the air-liquid interface is

achieved (Kuroishi et al. 2009). We found that thermal stress or acrolein decreased ZO-1 expression and disrupted its tight structure,

which indicated that thermal stress or acrolein regulated the paracellular ion transport pathway. Inhibitors (e.g., Amiloride, ouabain and

CFTRinh-172) can only block ion channels and transporters that belong to the transcellular pathway and inhibit the function of ion

transport. Therefore, inhibitors for ENaCs, CFTR, and Na^{+}/K^{+}-ATPase might not significantly affect these proteins. In our study, the

FITC-dextran flux assay showed increased FITC-dextran permeability in thermal stress or acrolein-treated MTE monolayers. Thermal

stress or acrolein not only damaged the complete structure of the tight junction but also increased its permeability.

In the current study, when adding acrolein to an apical bath solution, the bath solution was blown up to 10 times to ensure even

distribution of acrolein instead of gas bubbling. Maintaining bubbling of a bath solution facilitates loss of acrolein. This situation was supported by our results for comparing two experimental conditions with and without air bubbling (Fig. S5). A total of 500 μM acrolein adversely affected the integrity of the airway epithelial barrier, which was supported by a reduction in R_{TE} and I_{SC}, a disrupted ZO-1 structure, as shown by confocal images, and the increased FITC-dextran permeability in MTE monolayers. Therefore, 500 μM acrolein caused irreversible damage to the airway epithelial barrier. Through a fitted R_{TE}/I_{SC} dose-response curve for acrolein in MTE monolayers, we found that the K_i value for acrolein was lower at 40°C than at 37°C. Therefore, the same dose of acrolein caused more serious damage to R_{TE}/I_{SC} under thermal stress conditions compared with 37°C. Moreover, the damage to the integrity and permeability of tight junctions caused by acrolein was more severe at 40°C than at 37°C. The synergistic effects of thermal stress and acrolein induced further impairment to the airway epithelial barrier. Our previous studies indicated that oxidative aldehydes, including formaldehyde and crotonaldehyde, inhibited activity of ENaCs by causing activation of reactive oxygen species (ROS) in alveolar epithelium cell monolayers (Cui et al. 2016; Li et al. 2017). Modulation of antioxidant enzymes affected the thermal sensitivity of respiratory cells. Additionally, lowering superoxide dismutase enzyme levels resulted in a significant reduction in thermal resistance (Omar et al. 1987). However, overexpression of manganese superoxide dismutase by stable transfection provided cellular resistance against the cytotoxic effect of hyperthermia (Li and Oberley 1997; Kuninaka et al. 2000). Therefore, the synergistic effects of thermal stress and acrolein in the present study may have resulted from increased thermal sensitivity to cells, which was caused by acrolein-activated oxidative signaling.

As mentioned above, damage to the airway epithelium due to acrolein mainly results from its downstream signal via cellular oxidative stress, such as glutathione depletion, and subsequently ROS stimulation (Wang et al. 2009). In addition to ENaCs, Na^+/K^+-ATPase, and CFTR, some other proteins, including KCs, the Ca^{2+}-activated Cl^- channels (CaCCs), SCL26A9, and $Na^+/K^+/2Cl^-$ (NKCC), play roles in ion transport in the respiratory epithelium. Additionally, these proteins may be regulated by acrolein by induction of oxidative stress (Londino et al. 2017). As the major Ca^{2+}-activated KCs of ciliated cells, $K_{Ca}1.1$/KCNMA1 plays a role in maintaining fluid secretion and is affected by oxidative stress (Manzanares et al. 2011; Kis et al. 2016; Hermann et al. 2015). Moreover, exposure to

tobacco smoke inhibits $K_{Ca}1.1$/KCNMA1 and results in a reduction in airway surface liquid in human bronchial epithelial cells (Sailland et al. 2017). Anoctamin 6 functions as a CaCCs, and a Ca^{2+}-dependent phospholipid scramblase, which are stimulated by an increase in ROS and subsequent peroxidation of membrane lipids (Schreiber and Ousingsawat 2018; Scudieri et al. 2015). NKCC is located on the basolateral membrane of the tracheal epithelium and is an important conduit for Cl^- entry in the liquid-transporting epithelium (Gillie et al. 2001). A previous study reported that NKCC activation may contribute to the protective system against ROS-mediated damage to the airway epithelium (Matsuno et al. 2008). These studies may provide proof of acrolein damaging KCs, CaCCs, and NKCC in the airway epithelium.

In addition to the effects of smoke inhalation on the tracheal epithelium, we also found interactions between different ion transport systems. Apical and basolateral ion transport systems play diverse roles in the relationship between R_{TE} and I_{SC}. In our study, blockade of Na^+ entry or Cl^- secretion via apical ENaCs and CFTR, respectively, caused a reciprocal change in R_{TE} and I_{SC}. In contrast, inhibition of basolateral Na^+/K^+-ATPase resulted in a reduction in R_{TE} and I_{SC}. Additionally, a Cl^--free solution showed a similar effect as ouabain in inhibiting I_{SC}. This finding indicates the potential dependence of Na^+/K^+-ATPase on anion transport or interactions between vertical Cl^- and K^+ transporters. This interesting similar effect can be excluded for the associations between Cl^- and Na^+ transporters, because a Na^+-free solution led to an opposite effect on R_{TE} and I_{SC}. Our study showed that the apical ENaCs and CFTR modulated R_{TE}.

There are some limitations to this study, which need to be addressed in the future. Although the MTE *in vitro* model imitates the physiological state and complies with the morphological characteristics of the normal airway epithelium, there are still some differences from the *in vivo* environment. However, an *in vivo* model does not allow us to precisely measure R_{TE} and I_{SC}. Future studies need to focus on the effects of thermal stress or acrolein on inflammatory signal transduction pathways at the molecular level. In addition to the oxidative stress pathway, the mitogen-activated protein kinase (MAPK) pathway appears to play a role in regulating the airway epithelial barrier (Huang et al. 2016; Lee et al. 2018). A previous study suggested that MAPK Hog1 phosphorylation was activated to a peak at approximately 5 min after heat shock, and phosphorylated Hog1 declined to the basal level at 30 min (Dunayevich et al. 2018). Researchers reached a similar conclusion in another study, where they found that heat-activated MAPK signaling was elevated to a peak

at 5 min, and activation was weakened or even disappeared over time (Dong et al. 2015). In the present study, application of thermal stress to MTE monolayers caused a bi-phasic change, which indicated a sophisticated condition and a long time course. An *in vivo* model can show higher sensitivity and resistance to thermal stress than an *in vitro* model. Therefore, we ultimately to perform an *in vivo* study of thermal stress or acrolein-induced MAPK signaling.

Thermal stress and acrolein are the two main pathogenic factors of smoke inhalation injury. Our study suggests that these two factors damage the integrity of the airway epithelial barrier. The underlying mechanisms are related to the transcelluar pathway (mediated by ion channels and transporters) and the paracellular pathway (mediated by tight junctions). Morever, to the best of our knowledge, we have shown for the first time the synergistic effects of thermal stress and acrolein on impairment of the airway epithelial barrier.

Authors' contributions Hong-Long Ji conceived study, designed experiments, edited manuscript, and approved submission. Jianjun Chang, Zaixing Chen, and Hong-Guang Nie performed experiments, analyzed data, and plotted graphs. Jianjun Chang, Runzhen Zhao, Hong-Guang Nie, and Zaixing Chen prepared manuscript.

Compliance with ethical standards

Conflict of interest No conflicts of interest, financial or otherwise, are declared by the authors.

Abbreviations ENaCs, epithelial Na^+ channels; CFTR, cystic fibrosis transmembrane regulator; R_{TE}, transepithelial resistance; I_{SC}, short-circuit current; MTE, mouse tracheal epithelial; FITC, fluorescein isothiocyanate; HBE, human bronchial epithelial; DMSO, dimethyl sulfoxide; ZO-1, zonula occludens-1; P1, phase 1; P2, phase 2; ASI, amiloride-sensitive I_{SC}; KCs, K^+ channels; ROS, reactive oxygen species; CaCCs, Ca^{2+}-activated Cl^- channels; NKCC, $Na^+/K^+/2Cl^-$; MAPK, mitogen-activated protein kinase.

8 References

Alexander NS, Blount A, Zhang S, Skinner D, Hicks SB, Chestnut M, et al. Cystic fibrosis transmembrane conductance regulator modulation by the tobacco smoke toxin acrolein. Laryngoscope. 2012;122:1193-7.

Alwis KU, deCastro BR, Morrow JC, Blount BC. Acrolein Exposure in U.S. Tobacco Smokers and Non-Tobacco Users: NHANES 2005-2006. Environ Health Perspect. 2015;123:1302-8.

Anthony TR, Joggerst P, James L, Burgess JL, Leonard SS, Shogren ES. Method development study for APR cartridge evaluation in fire overhaul exposures. Ann Occup Hyg. 2007;51:703-16.

Bein K, Leikauf GD. Acrolein - a pulmonary hazard. Mol Nutr Food Res. 2011;55:1342-60.

Borchers MT, Wert SE, Leikauf GD. Acrolein-induced MUC5ac expression in rat airways. Am J Physiol. 1998;274:L573-81.

Burcham PC, Raso A, Thompson CA. Intermediate filament carbonylation during acute acrolein toxicity in A549 lung cells: functional consequences, chaperone redistribution, and protection by bisulfite. Antioxid Redox Signal. 2010;12(3):337-47.

Chang J, Ding Y, Zhou Z, Nie HG, Ji HL. Transepithelial Fluid and Salt Re-Absorption Regulated by cGK2 Signals. Int J Mol Sci. 2018;19:E881.

Chen Z, Zhao R, Zhao M, Liang X, Bhattarai D, Dhiman R, et al. Regulation of epithelial sodium channels in urokinase plasminogen activator deficiency. Am J Physiol Lung Cell Mol Physiol. 2014;307:L609-17.

Cui Y, Li H, Wu S, Zhao R, Du D, Ding Y, et al. Formaldehyde impairs transepithelial sodium transport. Sci Rep. 2016;6:35857.

Davidson DJ, Kilanowski FM, Randell SH, Sheppard DN, Dorin JR. A primary culture model of differentiated murine tracheal epithelium. Am J Physiol Lung Cell Mol Physiol. 2000;279(4):L766-78.

Dong ZW, Chen J, Ruan YC, Zhou T, Chen Y, Chen Y, et al. CFTR-regulated MAPK/NF-kappaB signaling in pulmonary inflammation in thermal inhalation injury. Sci Rep. 2015;5:15946

Dubick MA, Carden SC, Jordan BS, Langlinais PC, Mozingo DW. Indices of antioxidant status in rats subjected to wood smoke inhalation and/or thermal injury. Toxicology. 2002;176(1-2):145-57.

Dunayevich P, Baltanas R, Clemente JA, Couto A, Sapochnik D, Vasen G, et al. Heat-stress triggers MAPK crosstalk to turn on the hyperosmotic response pathway. Sci Rep. 2018;8(1):15168.

Faroon O, Roney N, Taylor J, Ashizawa A, Lumpkin MH, Plewak DJ. Acrolein environmental levels and potential for human exposure. Toxicol Ind Health. 2008;24(8):543-64.

Farquhar MG, Palade GE. Junctional complexes in various epithelia. The Journal of cell biology. 1963;17:375-412. doi:10.1083/jcb.17.2.375.

Farquhar MG, Palade GE. Junctional complexes in various epithelia. The Journal of cell biology. 1963;17:375-412.

Fitzgerald KT, Flood AA. Smoke inhalation. Clin Tech Small Anim Prac. 2006;21:205-14.

Flynn AN, Itani OA, Moninger TO, Welsh MJ. Acute regulation of tight junction ion selectivity in human airway epithelia. Proc Natl Acad Sci U S A. 2009;106:3591-6.

Fuller CM, Benos DJ. CFTR! Am J Physiol. 1992;263:C267-86.

Gillie DJ, Pace AJ, Coakley RJ, Koller BH, Barker PM. Liquid and ion transport by fetal airway and lung epithelia of mice deficient in sodium-potassium-2-chloride transporter. Am J Respir Cell Mol Biol. 2001;25(1):14-20.

Hales CA, Barkin PW, Jung W, Trautman E, Lamborghini D, Herrig N, et al. Synthetic smoke with acrolein but not HCl produces pulmonary edema. J Appl Physiol (1985). 1988;64:1121-33.

Han DY, Nie HG, Gu X, Nayak RC, Su XF, Fu J, et al. K+ channel openers restore verapamil-inhibited lung fluid resolution and transepithelial ion transport. Respir Res. 2010;11:65.

Haponik EF. Clinical smoke inhalation injury: pulmonary effects. Occup Med. 1993;8:430-68.

Hermann A, Sitdikova GF, Weiger TM. Oxidative Stress and Maxi Calcium-Activated Potassium (BK) Channels. Biomolecules. 2015;5(3):1870-911.

Horani A, Dickinson JD, Brody SL. Applications of mouse airway epithelial cell culture for asthma research. Methods Mol Biol.

2013;1032:91-107.

Hou Y, Cui Y, Zhou Z, Liu H, Zhang H, Ding Y, et al. Upregulation of the WNK4 Signaling Pathway Inhibits Epithelial Sodium Channels of Mouse Tracheal Epithelial Cells After Influenza A Infection. Front Pharmacol. 2019;10:12.

Howard M, Roux J, Iles KE, Miyazawa B, Christiaans S, Anjum N, et al. Activation of the heat shock response attenuates the interleukin 1beta-mediated inhibition of the amiloride-sensitive alveolar epithelial ion transport. Shock. 2013;39:189-96.

Huang W, Zhao H, Dong H, Wu Y, Yao L, Zou F, et al. High-mobility group box 1 impairs airway epithelial barrier function through the activation of the RAGE/ERK pathway. Int J Mol Med. 2016;37(5):1189-98.

Kis A, Krick S, Baumlin N, Salathe M. Airway Hydration, Apical K(+) Secretion, and the Large-Conductance, Ca(2+)-activated and Voltage-dependent Potassium (BK) Channel. Ann Am Thorac Soc. 2016;13 Suppl 2:S163-8.

Kuninaka S, Ichinose Y, Koja K, Toh Y. Suppression of manganese superoxide dismutase augments sensitivity to radiation, hyperthermia and doxorubicin in colon cancer cell lines by inducing apoptosis. Br J Cancer. 2000;83(7):928-34.

Kuroishi S, Suda T, Fujisawa T, Ide K, Inui N, Nakamura Y, et al. Epithelial-mesenchymal transition induced by transforming growth factor-beta1 in mouse tracheal epithelial cells. Respirology. 2009;14:828-37.

Lee KE, Jee HM, Hong JY, Kim MN, Oh MS, Kim YS, et al. German Cockroach Extract Induces Matrix Metalloproteinase-1 Expression, Leading to Tight Junction Disruption in Human Airway Epithelial Cells. Yunsei Med J. 2018;59(10):1222-31.

Li JJ, Oberley LW. Overexpression of manganese-containing superoxide dismutase confers resistance to the cytotoxicity of tumor necrosis factor alpha and/or hyperthermia. Cancer Res. 1997;57(10):1991-8.

Li Y, Chang J, Cui Y, Zhao R, Ding Y, Hou Y, et al. Novel mechanisms for crotonaldehyde-induced lung edema. Oncotarget. 2017;8:83509-22.

Londino JD, Lazrak A, Collawn JF, Bebok Z, Harrod KS, Matalon S. Influenza virus infection alters ion channel function of airway and alveolar cells: mechanisms and physiological sequelae. Am J Physiol Lung Cell Mol Physiol. 2017;313(5):L845-l58.

Manzanares D, Gonzalez C, Ivonnet P, Chen RS, Valencia-Gattas M, Conner GE, et al. Functional apical large conductance, Ca2+-activated, and voltage-dependent K+ channels are required for maintenance of airway surface liquid volume. J Biol Chem. 2011;286(22):19830-9.

Matsuno T, Ito Y, Ohashi T, Morise M, Takeda N, Shimokata K, et al. Dual pathway activated by tert-butyl hydroperoxide in human airway anion secretion. J Pharmacol Exp Ther. 2008;327(2):453-64.

Meacher DM, Menzel DB. Glutathione depletion in lung cells by low-molecular-weight aldehydes. Cell Biol Toxicol. 1999;15:163-71.

Nie HG, Chen L, Han DY, Li J, Song WF, Wei SP, et al. Regulation of epithelial sodium channels by cGMP/PKGII. J Physiol. 2009;587:2663-76

Omar RA, Yano S, Kikkawa Y. Antioxidant enzymes and survival of normal and simian virus 40-transformed mouse embryo cells after hyperthermia. Cancer Res. 1987;47(13):3473-6.

Reinhardt TE, Ottmar RD. Baseline measurements of smoke exposure among wildland firefighters. J Occup Environ Hyg. 2004;1:593-606.

Romet-Haddad S, Marano F, Blanquart C, Baeza-Squiban A. Tracheal epithelium in culture: a model for toxicity testing of inhaled molecules. Cell Biol Toxicol. 1992;8:141-50.

Roux E, Ouedraogo N, Hyvelin JM, Savineau JP, Marthan R. In vitro effect of air pollutants on human bronchi. Cell Biol Toxicol. 2002;18:289-99.

Sailland J, Grosche A, Baumlin N, Dennis JS, Schmid A, Krick S, et al. Role of Smad3 and p38 Signalling in Cigarette Smoke-induced CFTR and BK dysfunction in Primary Human Bronchial Airway Epithelial Cells. Sci Rep. 2017;7(1):10506.

Schreiber R, Ousingsawat J. Regulation of TMEM16A/ANO1 and TMEM16F/ANO6 ion currents and phospholipid scrambling by Ca(2+) and plasma membrane lipid. J Physiol. 2018;596(2):217-29.

Scudieri P, Caci E, Venturini A, Sondo E, Pianigiani G, Marchetti C, et al. Ion channel and lipid scramblase activity associated with expression of TMEM16F/ANO6 isoforms. J Physiol. 2015;593(17):3829-48.

Sheppard DN, Welsh MJ. Structure and function of the CFTR chloride channel. Physiol Rev. 1999;79:S23-45.

Stevens JF, Maier CS. Acrolein: sources, metabolism, and biomolecular interactions relevant to human health and disease. Mol Nutr Food Res. 2008;52:7-25.

Tobey NA, Sikka D, Marten E, Caymaz-Bor C, Hosseini SS, Orlando RC. Effect of heat stress on rabbit esophageal epithelium. Am J Physiol. 1999;276:G1322-30.

Tsukita S, Furuse M. Claudin-based barrier in simple and stratified cellular sheets. Curr Opin Cell Biol. 2002;14:531-6.

Umeda K, Ikenouchi J, Katahira-Tayama S, Furuse K, Sasaki H, Nakayama M, et al. ZO-1 and ZO-2 independently determine where claudins are polymerized in tight-junction strand formation. Cell. 2006;126:741-54.

Van Itallie CM, Anderson JM. Claudins and epithelial paracellular transport. Annu Rev Physiol. 2006;68:403-29.

Wang T, Liu Y, Chen L, Wang X, Hu XR, Feng YL, et al. Effect of sildenafil on acrolein-induced airway inflammation and mucus production in rats. European Respir J. 2009;33(5):1122-32.

Wang X, Adler KB, Erjefalt J, Bai C. Airway epithelial dysfunction in the development of acute lung injury and acute respiratory distress syndrome. Expert Rev Respir Med. 2007;1:149-55.

Wang Y, Bai C, Li K, Adler KB, Wang X. Role of airway epithelial cells in development of asthma and allergic rhinitis. Respir Med. 2008;102:949-55.

You K, Yang HT, Kym D, Yoon J, HaejunYim, Cho YS, et al. Inhalation injury in burn patients: establishing the link between diagnosis and prognosis. Burns. 2014;40:1470-5.

Figure Legends

Fig. 1 Thermal stress alternates bioelectric features in mouse tracheal epithelial (MTE) and human bronchial epithelial (HBE)

monolayers. (a) Representative transepithelial resistance (R_{TE}) trace in MTE monolayer mounted on an Ussing chamber setup. R_{TE} was

decreased in a two-phase manner (P1 and P2 were labeled) during the temperature of the bath solution was raised to 40°C (red line)

from 37°C (black line). When the temperature rose to 38.5°C, the changes in R_{TE} and I_{SC} were observed, and it took 4.7 min for the

course from 37°C to 40°C. **(b)** Short-circuit current (I_{SC}) trace recorded simultaneously in the same MTE monolayer. P1 and P2 of I_{SC}

were defined using the R_{TE} time course. I_{SC} was increased at P1 and decreased at P2. **(c)** Average R_{TE} levels of MTE monolayers at

different time points during thermal stress. Student's t-test. $^{***}P < 0.001$. $n = 20$. **(d)** Average I_{SC} levels of MTE monolayers. Student's

t-test. $^{***}P < 0.001$. $n = 20$. **(e)** Time for reducing half of the total R_{TE} ($\tau_{1/2}$). The $\tau_{1/2}$ value was computed by fitting R_{TE} raw data with

the ExpDec1 function. ($y_0 = 0.36 \pm 0.04$, $A = 0.87 \pm 0.04$ [P1]; $y_0 = -0.87 \pm 0.11$, $A = 3.22 \pm 0.36$ [P2]). Student's t-test. $^{***}P < 0.001$. n

$= 12$. **(f)** Representative R_{TE} trace in HBE monolayer. R_{TE} was decreased during the temperature of the bath solution was elevated to

40°C (red line) from 37°C (black line). **(g)** I_{SC} trace recorded simultaneously in the same HBE monolayer. I_{SC} was increased to a high

and stable level during thermal stress. **(h)** Average R_{TE} levels of HBE monolayers under the conditions of 37°C and 40°C. Student's t-

test. NS, no significance. $n = 6$. **(i)** Average I_{SC} levels of HBE monolayers. Student's t-test. $^{*}P < 0.05$. $n = 6$.

Fig. 2 Na^{+} transport mediates thermal stress-induced bioelectric changes in MTE monolayers. (a, b) Representative R_{TE} and I_{SC}

traces in the presence of saline bath solution (control group, solid line), Na^{+}-free bath solution (dashed lines), amiloride (Amil, 100 μM,

dotted lines), or ouabain (1 mM, dashed-dotted lines). At 37°C (black lines), Na^{+}-free bath solution, amiloride or ouabain was applied

at the time pointed by the arrow. And the temperature of the bath solution was elevated to 40°C (red lines). When the temperature rose

to 38.5°C, the changes in R_{TE} and I_{SC} were observed. **(c)** Average thermal stress-sensitive R_{TE} levels ($\triangle R_{TE}$, the difference between the

0 initial R_{TE} and ending R_{TE} within one phase [P1 or P2]). Mann-Whitney U test and Student's t-test. $^{**}P < 0.01$. NS, no significance. $n =$

1 27. **(d)** Average thermal stress-sensitive R_{TE} levels ($\triangle R_{TE}$, the difference between the initial R_{TE} and ending R_{TE} during the entire process

of thermal stress). Student's t-test. $^*P < 0.05$ and $^{**}P < 0.01$. $n = 17$. **(e)** Average thermal stress-sensitive I_{SC} levels at P1 ($\triangle I_{SC}$, the

difference between the basal I_{SC} and the peak I_{SC} at P1). Student's t-test. $^*P < 0.05$ and $^{***}P < 0.001$. $n = 29$. **(f)** Average thermal stress-

sensitive I_{SC} levels at P2 ($\triangle I_{SC}$, the difference between the peak I_{SC} and the ending I_{SC} at P2). Student's t-test. $^{***}P < 0.001$. $n = 20$.

Fig. 3 CFTR is involved in thermal stress-induced bioelectric changes in MTE monolayers (a, b) Representative R_{TE} and I_{SC} traces

in the presence of saline bath solution (control group, solid line) and CFTRinh-172 (CFTRinh, 20 μM, dashed line). Segments of traces

recorded at different bath temperatures were shown as black (37°C) and red lines (40°C). Arrows indicated the time to add CFTRinh-

172. When the temperature rose to 38.5°C, the obvious bioelectric changes were observed. **(c)** Average thermal stress-sensitive R_{TE}

levels ($\triangle R_{TE}$, the difference between the initial R_{TE} and ending R_{TE} within one phase [P1 or P2]). Student's t-test. $^{**}P < 0.01$ and $^{***}P <$

0.001. $n = 18$. **(d)** Average thermal stress-sensitive I_{SC} levels at P1 ($\triangle I_{SC}$, the difference between the basal I_{SC} and the peak I_{SC} at P1).

Student's t-test. $^*P < 0.05$. $n = 9$. **(e)** Average thermal stress-sensitive I_{SC} levels at P2 ($\triangle I_{SC}$, the difference between the peak I_{SC} and the

ending I_{SC} at P2). Student's t-test. $^*P < 0.05$. $n = 9$.

Fig. 4. Activation of ENaCs, CFTR, and Na^+-K^+-ATPase of MTE monolayers by thermal stress. (a, c, and e) Representative I_{SC}

traces before and after the addition of amiloride (100 μM), CFTRinh (20 μM), or ouabain (1 mM) at 37°C (black lines) and 40°C (red

lines). Arrows indicated the time to add inhibitors. **(b, d and f)** Average I_{SC} levels. Total I_{SC} meant the basal I_{SC} level; +Amil I_{SC},

+CFTRinh I_{SC}, and +Ouabain I_{SC} meant inhibitor-resistant I_{SC} level; ASI (amiloride-sensitive I_{SC}), CFTRinh I_{SC} and Ouabain I_{SC}

respectively reflected the activity of each ion channel and was calculated by the difference between the basal I_{SC} and inhibitor-resistant

I_{SC}. Student's t-test and Mann-Whitney U test. $^*P < 0.05$ and $^{**}P < 0.01$. NS, no significance. $n = 18$. **(g)** Average R_{TE} levels. Student's t-

test and Mann-Whitney U test. $^*P < 0.05$ and $^{***}P < 0.001$. NS, no significance. $n = 36$.

Fig. 5 Acrolein impairs bioelectric features in MTE and HBE monolayers. (a, b) Representative R_{TE} and I_{SC} traces in the presence

of 5, 50, and 500 µM acrolein at 37°C (black lines) and 40°C (red lines) in MTE monolayers. Acrolein (ACR, 5, 50, and 500 µM) was

pipetted to the bath as indicated by arrows. 500 µM acrolein decreased R_{TE} and I_{SC}. **(c, d)** Dose-response curve for acrolein in MTE

monolayers. Normalized R_{TE}/I_{SC} points were fitted with the Hill equation (*START* = 1, *END* = 0, *n* = 22). The K_i value of R_{TE} for acrolein

was 120.22 µM at 37°C and 79.43 µM at 40°C; The K_i value of I_{SC} for acrolein was 125.89 µM at 37°C and 77.62 µM at 40°C. **(e, f)**

Representative R_{TE} and I_{SC} traces of HBE monolayers in the presence of 5, 50, and 500 µM acrolein under conditions of 37°C (black

line) and 40°C (red line). Arrows indicated the time to add acrolein (ACR, 5, 50, and 500 µM). R_{TE} and I_{SC} were decreased by 500 µM

acrolein. **(g, h)** Dose-response curve for acrolein in HBE monolayers. Normalized R_{TE}/I_{SC} points were fitted by the Hill equation (*START*

= 1, *END* = 0, *n* = 2). The K_i value of R_{TE} for acrolein was 83.18 µM at 37°C and 97.73 µM at 40°C, respectively; the K_i value of I_{SC} for

acrolein was 125.89 µM at 37°C and 112.21 µM at 40°C, respectively.

Fig. 6 Na$^+$ transport is involved in acrolein-impaired bioelectric features in MTE monolayers. (a, b) Representative R_{TE} and I_{SC}

traces in the presence of saline bath solution (control group, solid lines), Na$^+$-free bath solution (dashed lines), amiloride (Amil, 100 µM,

dotted lines), or ouabain (1 mM, dashed-dotted line). At basal level (black lines), Na$^+$-free bath solution, amiloride or ouabain was

applied at the time pointed by the arrow. And then 500 µM acrolein (ACR, blue lines) was added. **(c)** Average acrolein-sensitive R_{TE}

levels ($\triangle R_{TE}$, the difference between basal R_{TE} and acrolein-resistant R_{TE}). Mann-Whitney U test. $^*P < 0.05$. *n* = 16. **(d)** Average acrolein-

sensitive I_{SC} levels ($\triangle I_{SC}$, the difference between basal I_{SC} and acrolein-resistant I_{SC}). Student's t-test. $^{**}P < 0.01$ and $^{***}P < 0.001$. *n* =

16.

Fig. 7 CFTR mediates acrolein-impaired bioelectric features in MTE monolayers. (a, b) Representative R_{TE} and I_{SC} traces in the

presence of saline bath solution (control group, solid lines) and CFTRinh-172 (20 µM, dashed lines). Arrows showed the time for adding

CFTRinh-172. Then monolayers were exposed to 500 µM acrolein (ACR, blue lines). **(c)** Average acrolein-sensitive R_{TE} levels ($\triangle R_{TE}$,

the difference between basal R_{TE} and acrolein-resistant R_{TE}). Student's t-test. $^{**}P < 0.01$. *n* = 11. **(d)** Average acrolein-sensitive I_{SC} levels

6   ($\triangle I_{SC}$, the difference between basal I_{SC} and acrolein-resistant I_{SC}). Student's t-test. $^{***}P < 0.001$. $n = 11$.

8   **Fig. 8 The synergistic effects of thermal stress and acrolein on ZO-1 tight junction in MTE monolayers.** Laser scanning confocol

9   imaging for ZO-1 protein in MTE monolayers treated by 37°C **(a-c)**, 40°C for 120 min **(d-f)**, 500 μM acrolein **(g-i)**, and combination of

0   40°C and 500 μM acrolein **(j-l)**. Nucleus was labeled with hoechst (blue), and tight junction was labeled with ZO-1 (green). Scale bar:

1   50 μm. **(m)** Quantification of ZO-1 fluorescence intensity. Student's t-test. $^{*}P < 0.05$, $^{***}P < 0.001$, $^{\&}P < 0.05$ and $^{\#}P < 0.05$. $n = 12$.

Supplementary materals

Ion transport mechanisms for smoke inhalation injured airway epithelial barrier

Jianjun Chang, Zaixing Chen[&], Runzhen Zhao, Hong-Guang Nie, Hong-Long Ji

Supplemental Figure LegendsFig. S1 Extreme thermal stress disrupts bioelectric features in minutes in mouse tracheal epithelial (MTE) monolayers. (a) Representative transepithelial resistance (R_{TE}) trace at 37°C, 42°C, and 44°C. R_{TE} was eliminated to zero at 44°C. **(b)** Short-circuit current (I_{SC}) trace recorded simultaneously of the same monolayer. At 44°C, I_{SC} was decreased followed by a jump, probably because of the damage to tight monolayer. **(c)** Representative R_{TE} trace at 37°C, 40°C, and 42°C. R_{TE} was eliminated to zero at 42°C. **(d)** I_{SC} trace of the same monolayer. I_{SC} was decreased at 42°C and then shot up and out of the scale in minutes probably due to impaired monolayer.

Fig. S2 Long process of apical and basolateral-permeabilizing destabilizes R_{TE} and I_{SC} measurements in MTE monolayers. (a) Representative R_{TE} traces of apical-permeabilized MTE monolayers. After long time of apical permeabilizing process, R_{TE} was decreased near zero. **(b)** I_{SC} traces monitored synchronously of the same monolayers. I_{SC} was reduced followed by a jump. **(c)** Representative R_{TE} traces of basolateral-permeabilized MTE monolayers. R_{TE} was almost decreased to zero. **(d)** I_{SC} traces of the same monolayers. I_{SC} was decreased and then out of the scale upwardly probably due to the damage to tight monolayers.

Fig. S3 Cl-free bath solution eliminates R_{TE} and I_{SC} levels in MTE monolayers. (a) Representative R_{TE} trace in the presence of saline bath solution (control group, black line) and Cl-free bath solution (green line). Cl-free bath solution caused the transient increment followed by the slow decline of R_{TE}. **(b)** I_{SC} trace recorded synchronously of the same monolayer. **(c)** Average R_{TE} levels. R_{TE} was eliminated to 0.24 ± 0.03 k$\Omega \times$ cm^2 compared with the control group (0.91 ± 0.05 k$\Omega \times$ cm^2), probably due to impaired tight monolayers. Student's t-test. [***]$P < 0.001$. $n = 8$. **(d)** Average I_{SC} levels. Student's t-test. [***]$P < 0.001$. $n = 8$.

Fig. S4 Cell count and viability in MTE monolayers. (a) Cell number of monolayers post the exposure of thermal stress (120 min) and acrolein (500 μM). Ten randomly selected images across the monolayers from at least 3 independent experiments were captured and counted for total cells with DAPI staining using a cell count plug-in of the ImageJ. No significant difference was observed among each group. Thermal stress and acrolein could not induce significant cell detachment of monolayers. NS, no significance. $n = 12$. **(b)** Cell viability of monolayers treated by thermal stress (120 min) and acrolein (500 μM). At the end of treatment, cell viability was measured with the cell counting kit-8 assay. Thermal stress and acrolein failed to cause obvious cell death of monolayers. NS, no significance. $n = 12$.

Fig. S5 Comparison of the effects of acrolein on I_{SC} and R_{TE} in bubbled and unbubbled MTE monolayers. (a) Representative R_{TE} trace when unbubbling (black line) or bubbling (red line). **(b)** I_{SC} trace recorded simultaneously of the same monolayers. Keeping bubbling the bath solution facilitated the loss of acrolein and mitigated the effects of acrolein on R_{TE} and I_{SC}.

Fig. S6 Thermal stress and acrolein impair occludin tight junction of MTE monolayers. Laser scanning confocol imaging for occludin tight junction of MTE monolayers treated by 37°C **(a-c)**, 40°C for 120 min **(d-f)**, 500 μM acrolein **(g-i)**, and the combination of 40°C and 500 μM acrolein **(j-l)**. Nuclei were labeled with hoechst (blue), and tight junctions were labeled with occludin (green). Scale bar: 10 μm. **(m-o)** Immunolocalization of claudin tight junction of normal MTE monolayer. Nuclei were labeled with hoechst (blue), and tight junctions were labeled with claudin (green). The expression of claudin was insufficient to be visualized. Scale bar: 50 μm. **(p)** Fluorescein isothiocyanate-dextran (4 kDa) permeability assay of MTE monolayers. MTE monolayers were treated by 37°C, 40°C for 120 min, 500 μM acrolein, and the combination of 40°C and 500 μM acrolein. Thermal stress and acrolein increased the permeability of fluorescein isothiocyanate-dextran. Student's t-test. [***]$P < 0.001$. $n = 14$.

Fig. S1

Fig. S2

Fig. S3

Fig. S4

Fig. S5

Fig. S6

Supplemental data table 1. Parameters when R_{TE} raw data were fitted by the ExpDec1 function

Group	P1								P2			
Sample number	1	2	3	4	5	6	7	8	1	2	3	4
y_0	0.34	0.46	0.52	0.097	0.30	0.35	0.36	0.42	-0.89	-1.08	-0.97	-0.55
A	0.92	0.97	0.96	0.87	0.87	0.83	0.91	0.64	3.13	3.88	3.62	2.26