PprA interacts with replication proteins and affects their physicochemical properties required
for replication initiation in *Deinococcus radiodurans*

Ganesh K. Maurya [$,1, 2] and Hari S. Misra *[1, 2]

[1]Molecular Biology Division, Bhabha Atomic Research Centre, Mumbai 400085. India
[2]Homi Bhabha National Institute, Mumbai, 400094. India

*Address for Correspondence

Dr. H. S. Misra

Molecular Biology Division

Bhabha Atomic Research Centre

Mumbai- 400085

Tel: 91-22-25593821

Fax: 91-22-25505151 (EN: 13238)

Email: hsmisra@barc.gov.in
16

17 $ Present Address: Mahila Mahavidyalaya, Banaras Hindu University, Varanasi-221005.
18 India.

Abstract

The deletion mutant of *pprA,* a gene encoding pleiotropic functions in radioresistant bacterium *Deinococcus radiodurans,* showed an increased genomic content and ploidy in chromosome I and chromosome II. We identified *oriC* in chromosome I *(oriCI)* and demonstrated the sequence specific interaction of deinococcal DnaA (drDnaA) with *oriCI.* drDnaA and drDnaB showed ATPase activity while drDnaB catalyzed $5'{\rightarrow}3'$ dsDNA helicase activity. These proteins showed both homotypic and heterotypic interactions. The roles of C-terminal domain of drDnaA in *oriCI* binding and its stimulation of ATPase activity were demonstrated. Notably, PprA showed ~2 times higher affinity to drDnaA as compared to drDnaB and attenuated both homotypic and heterotypic interactions of these proteins. Interestingly, the ATPase activity of drDnaA but not drDnaB was inhibited in presence of PprA. These results suggested that PprA influences the physicochemical properties of drDnaA and drDnaB that are required for initiation of DNA replication at *oriCI* site in this bacterium.

Introduction

The <u>origin of replication</u> in bacterial chromosome (*oriC*) is a discrete locus that contains AT rich conserved DNA motifs and the varying number of 9 mer repeats of non-palindromic sequences called DnaA boxes. These boxes are recognized by a replication initiator protein named DnaA, followed by the assembly of replication initiation complex at *oriC* (Messer, 2002; Mott and Berger, 2007). Mechanisms underlying replication initiation have been characterised in large number of bacteria harbouring limited copies of single circular chromosome as inheritable genetic material (Zakrzewska-Czerwińska et al., 2007; Leonard and Grimwade, 2015). It has been shown in *E. coli* that DnaA-ATP oligomer binds to DnaA boxes at *oriC* and unwinds the adjacent AT rich region. Subsequently, a hexameric complex of replicative helicase DnaB and its loader DnaC ($DnaB_6$-$DnaC_6$) is recruited to the unwound region in *oriC* resulting to the formation of pre-priming complex (Skarstad and Katayama, 2013; Chodavarapu and Kaguni, 2016). This provides the site for binding of primase and activation of various events required for the progression of replication complex comprised of DNA polymerase III holoenzyme. DnaB hexameric ring translocates bidirectionally in order to unwind the parental duplex DNA while 3' end of the primer is extended by polymerase complex (McHenry, 2011; Yao and O'Donnell, 2010). In *E. coli* it has been shown that the initiation of DNA replication at *oriC* site is tightly regulated and the next round of *oriC* mediated DNA replication must wait till the *oriC* of newly replicated daughter chromosomes is fully methylated. Therefore, under normal growth conditions, the number of copies of primary chromosome per cell is expected to be less than 2 (Skarstad and Katayama, 2013). Recently, the bacteria with multiple copies of multipartite genome system have been reported. Notably, most of them are either parasite to some forms of life or exhibits super tolerance to abiotic stresses (Misra et al, 2018). The ploidy of chromosomes in these bacteria allowed us to revisit the principle of *oriC* regulation as known in bacteria containing less than

2 copies of circular chromosome per cell. Mechanisms underlying regulation of *oriC* function

in multipartite genome harbouring bacteria have not been studied in detail. In case of *Vibrio*

cholera, which harbours two chromosomes viz; chromosome I (Chr I) and chromosome II

(Chr II), the Chr I replicates similar to *E. coli* chromosome while Chr II follows replication

mechanism similar to low copy number P1 and F plasmids (Egan et al., 2005; Egan and

Waldor, 2003; Jha et al., 2012; Ramachandran et al., 2017; Koch et al., 2010; Fournes et al.,

2018).

Deinococcus radiodurans is one of the most radiation resistant bacteria, characterized for its

extraordinary ability to withstand the lethal effects of DNA damaging agents including

radiation and desiccation (Zahradka et al., 2006; Cox and Battista, 2005; Slade and Radman,

2011; Misra et al., 2013). It also harbours the multipartite genome system comprised of two

chromosomes (Chr I (2,648,638bp) and ChrII (412,348bp) and a megaplasmid (177,466bp)

and plasmid (45,704 bp) (White et al., 1999). Interestingly, each of these genome elements is

found in multiple copies per cell (Hansen, 1978). Chr I of *D. radiodurans* encodes putative

DnaA (DR_0002) and DnaB (DR_0549) (hereafter named drDnaA and drDnaB, respectively)

while chromosome II encodes PprA (DR_A0346), which has been characterized for various

functions (Narumi et al., 2004, Kota et al., 2014, 2016; Adachi et al, 2014). Recently, the

extended structure of PprA has been reported where a possibility of it acting as protein

scaffold (Adachi et al., 2019), which might explain PprA roles in large number of molecular

events. Here, for the first time, we report the functional characterization of chromosome

replication initiation proteins drDnaA and drDnaB in *D. radiodurans* and demonstrated that

PprA a pleiotropic protein involved in radioresistance plays a role in the regulation of DNA

replication. We showed that drDnaA interacts with *origin of replication* (*oriCI*) in a sequence

specific manner. drDnaA is characterized as an *oriCI* responsive ATPase while drDnaB as

ATP dependent 5′→3′ dsDNA helicase with higher affinity for ssDNA as compared to

dsDNA. Interestingly, PprA interacted with drDnaA at relatively higher affinity than drDnaB

and inhibited both homotypic and heterotypic interactions of these proteins. Further, PprA

downregulated the ATPase activity of drDnaA but showed no effect on ATPase and helicase

activities of drDnaB. These results suggested that drDnaA and drDnaB confers the desired

functions for the initiation of replication at *oriCI* in *D. radiodurans*. Furthermore, the

interference of PprA in the physicochemical properties of these replication proteins and an

increase in copy numbers of both primary and secondary chromosomes in absence of *pprA*

has suggested its involvement in the regulation of chromosomal replication in this bacterium.

Results

The *pprA* deletion affects genomic content in *D. radiodurans*

Earlier, the regulatory role of PprA in cell division and genome maintenance has been

demonstrated (Devigne et al., 2013; Kota et al., 2014; Devigne at al., 2016; Kota et al., 2016).

The DNA content and the copy number of genome elements in *pprA* deletion mutant were

compared with wild type *D. radiodurans*. The total DNA content in mutant cells (~8.41 ±

1.05 fg per cell) was found to be ~3 fold higher than wild type cells (~3.05 ± 0.47 fg per cell)

(Fig. 1A). Similarly, the DAPI fluorescence in Δ*pprA* mutant was ~2 fold higher as compared

to wild type (Fig. 1B). The cell scan analysis of DAPI stained cells showed that ~80 % Δ*pprA*

cells have ~2 folds higher DAPI fluorescence as compared to wild type cells grown

identically (Fig. 1C). When we checked the copy number of each replicon, the average copy

number of Chr I and Chr II was ~2.5 to 3-fold higher in Δ*pprA* mutant as compared to wild

type. For instance, the copy number of Chr I was ~ 8 in wild type while it was ~18 in Δ*pprA*

mutant. The copy number of Chr II had also increased from ~7 in wild type to ~17 per cell in

Δ*pprA* mutant (Fig. 1D). Interestingly, the copy number of megaplasmid and small plasmid

did not change. This increase in genomic content and chromosomal copy number indicated a

strong possibility of continued DNA replication in the absence of PprA. Thus, a possible role

of PprA in the regulation of DNA replication is suggested in *D. radiodurans*.

PprA interacts with DnaA and DnaB of *D. radiodurans*

In order to get the mechanistic insights on the regulatory role of PprA in DNA replication, the

physical and functional interaction of PprA with drDnaA and drDnaB proteins were

separately monitored. *E. coli* BTH101 (*cyaA*⁻) co-expressing T18 tagged PprA and T25

tagged drDnaA or drDnaB showed reconstitution of active CyaA leading to the expression of

β-galactosidase activity (Fig. 2A). The reconstitution of active CyaA from its T18 and T25

domains would be possible only when target proteins tagged with these domains interact.

PprA interaction with drDnaA and drDnaB was further confirmed using surface plasmon

resonance. For that recombinant PprA protein was immobilized on a bare gold sensor chip

and the purified recombinant drDnaA or drDnaB (Fig. S2) was passed over the immobilised

PprA in the mobile phase. Results showed a concentration-dependent increase in SPR signals

in case of drDnaA or drDnaB (Fig. 2 B,C). The dissociation constant (Kd) for drDnaA was

$5.41 \times 10^{-7} \pm 1.8 \times 10^{-8}$ [M] while it was $9.71 \times 10^{-7} \pm 1.1 \times 10^{-8}$ [M] for drDnaB. These

indicated that PprA interacts with drDnaA with ~2-fold higher affinity than drDnaB. We have

also tested the interaction of C-terminal deletion mutant of DnaA (DnaΔCt) with PprA in

surrogate *E. coli* using co-immunoprecipitation, as described in methodology. We found that

truncation of C-terminus from drDnaA has not affected its interaction with PprA (Fig. 2D).

This suggests a possibility of N-terminal and/or middle region of drDnaA could be the region

of interaction with PprA protein in this bacterium.

The interaction of PprA with drDnaA or drDnaB was also monitored in *D. radiodurans*. For

that, the total proteins of *D. radiodurans* cells expressing T18 tagged drDnaA or drDnaB on

plasmid and PprA on chromosome were precipitated using antibodies against PprA, and

interaction of PprA with drDnaA / drDnaB partners if any, were detected using T18

antibodies. PprA interaction with drDnaA and drDnaB was confirmed in *D. radiodurans*

(Fig. 2E). These results suggested that PprA interacts with both drDnaA and drDnaB in *D.*

radiodurans, and that might help PprA to regulate the functions of these proteins in this

bacterium.

The chromosome I of *D. radiodurans* contains putative *oriCI*

The putative origin of replication of chromosome I in *D. radiodurans* R1 was predicted using

DOriC database (DoriC accession number – ORI10010007) (Luo and Gao, 2019). The 1183

to 1903 bp upstream to dr*dnaA* gene (DR_0002) was searched for consensus sequences of 9

mer DnaA-boxes using WebLogo online tool (Crooks et al., 2004). The 1273 to 1772 bp

(~500 bp) region upstream to dr*dnaA* coding sequence contains 13 DnaA-boxes of 9 mer

each and was named as putative *oriCI*. These DnaA boxes are distributed in random fashion

on both sense (4 DnaA boxes) and antisense (9 DnaA boxes) strands of chromosome I (Fig.

3). Unlike *E. coli oriC*, the structure of *oriCI* in *D. radiodurans* is complex. For instance, it is

comparatively longer than the *oriC* of *E. coli* and contains many 'non-perfect' DnaA-boxes.

The *oriCI* has 46.2 % GC content indicating that it is an AT rich region. When we analysed

this region for typical *E. coli* type DnaA-boxes, 8 out of 13 DnaA-boxes were perfect like

DnaA-boxes (TTATCCACA) of *oriC* in *E. coli* (Messer, 2002). The other 5 boxes differ

from *E. coli* type by single nucleotide (Fig. 3) but were like the DnaA-boxes in the

chromosome of other bacteria like *Cyanothece* 51142, *T. thermophilus* and *B. subtilis* (Huang

et al., 2015; Schaper et al., 2000; Moriya et al., 1988).

A web logo of these DnaA-boxes has generated a consensus sequence of
T(A/G)TA(T)TCCACA. These DnaA-boxes are distributed in random fashion on both sense
(4 DnaA-boxes) and antisense (9 DnaA-boxes) strands of chromosome I (Fig. 3). This
suggested that *oriCI* in chromosome I of *D. radiodurans* is largely similar to *oriC* of *E.coli.*
166

drDnaA is a sequence specific *oriCI* binding protein

The recombinant drDnaA was purified from recombinant *E. coli* (Fig.S1A) and its *oriCI*

binding activity was checked by electrophoretic mobility shift assays (EMSA). *oriCI*

containing 13 repeats of DnaA-boxes was generated by PCR amplification and radiolabelled

using [^{32}P]γ-ATP. The DNA substrate was incubated with purified recombinant drDnaA and

the nucleoprotein complex was separated on native poly-acrylamide gel electrophoresis

(PAGE). drDnaA showed sequence-specific interaction with *oriCI in vitro* as this interaction

was not affected by addition of 50-fold excess of cold non-specific DNA (Fig. 4A, B).

Interestingly, the affinity of drDnaA to *oriCI* (Kd = 1.68 ± 0.23 μM) increased by ~ 8-fold in

the presence of ATP (Kd = 0.243 ± 0.02 μM) (Fig. 4C, D). We further checked the affinity of

DnaA with different number of DnaA boxes in *oriCI* in the presence of ATP and found that

as the number of DnaA boxes reduces, the affinity of DrDnaA also reduces (Table 1). . Like

E. coli DnaA, drDnaA has also shown specific affinity for single DnaA box though the

affinity with perfect DnaA box (TTATCCACA; Kd = 2.88 ± 0.28 μM) is ~2 fold higher than

non-perfect type DnaA boxes like TTTTCCACA or GTATCCACA (Kd = 4.21 ± 0.15 μM or

4.36 ± 0.18 μM, respectively) (Fig. 2 G-I).These results showed that drDnaA is an *oriCI*

binding protein and its interaction with *oriCI* is stimulated in the presence of ATP.

drDnaA binds *oriCI* through its C-terminal and N-terminal domain associated ATPase

activity is stimulated by *oriCI in vitro*

Since drDnaA showed relatively higher affinity for *oriCI* in the presence of ATP (Fig. 4C),

the ability of drDnaA for ATP hydrolysis and the effect of *oriCI* on its ATPase function was

further investigated. We observed that drDnaA can hydrolyse [^{32}P] αATP into [^{32}P] αADP

(Fig 5A) and this activity was stimulated in the presence of *oriCI* (Fig. 5B, 5C). Interestingly,

the ATPase activity of drDnaA was not affected in the presence of non-specific dsDNA (Fig.

S3). The helix-turn-helix motif in domain IV at the C-terminal of DnaA has been reported for

sequence specificity in many bacteria (Roth and Messer, 1995; Sutton and Kaguni, 1997;

Majka et al., 1999; Blaesing et al., 2000; Messer et al., 1999; Fujikawa et al., 2003).

Therefore, the C-terminal (domain IV) truncation was created in drDnaA (DnaAΔCt) (Fig

6A) and its binding to *oriCI* was checked in the presence of ATP. DnaAΔCt failed to bind

oriCI (Fig. 6B). Although, this truncation did not affect ATPase activity of drDnaA (Fig 6C),

as expected the *oriCI* stimulation of ATPase activity was not observed in DnaAΔCt protein

(Fig 6D, 6E). These results suggested that drDnaA is an *oriCI* binding ATPase and its C-

terminal domain IV is essential for recognition to *oriCI* sequence and thereby the ATPase

activity stimulation by this cognate *cis* element.

drDnaB is a ssDNA responsive ATPase and an ATP dependent 5′→3′ dsDNA helicase

drDnaB showed non-specific DNA binding activity with *oriCI* as chase with non-specific

dsDNA has significantly competed out *oriCI* from drDnaB binding. The affinity of drDnaB

to *oriCI* (Kd = 8.89±0.92 µM) was not affected by ATP (Kd = 8.23±0.41 µM). However,

drDnaB affinity to ssDNA (Kd= 3.11±0.11 µM) was nearly ~3 times higher than dsDNA

(*oriCI*) (Kd= 8.89±0.92 µM). drDnaB affinity to ssDNA was further increased by 2 folds in

the presence of ATP (Kd = 1.42±0.05 µM) (Fig. 7). These results suggested that drDnaB

prefers binding to ssDNA over dsDNA in ATP independent manner, and addition of ATP

improves its affinity for ssDNA but not to dsDNA. drDnaB could hydrolyse [^{32}P]-αATP to

$[^{32}P]$-αADP (Fig 8A), which was stimulated in the presence of ssDNA (Fig. 8B, 8C). Similar

observations were reported earlier for other DnaB homologs (Biswas et al., 2009; Zhang et

al., 2014). DnaBs are characterized as the replicative helicases and play essential role in *oriC*

mediated DNA replication in other bacteria (LeBowitz and McMacke, 1986; Patel and Picha,

2000; Zawilak, et al., 2001). Therefore, the helicase activity of recombinant drDnaB was

checked on dsDNA having either 3′ or 5′ overhangs in the presence and absence of ATP and

products were monitored in gel and FRET based assays. Results showed that drDnaB

unwinds dsDNA with 5′ overhang only and not with 3′ overhang (Compare Fig. 9A and 9E

with 9C and 9F). This 5′ → 3′ dsDNA helicase activity required the hydrolysable ATP as no

DNA unwinding was observed when ATP was replaced with ATP-γ-S (Fig 9B). The similar

results were obtained in FRET assays with FAM as donor and BHQ as acceptor (Fig 9D).

Results showed that the fluorescence has increased in the presence of protein and ATP but

not with ATP-γ-S (Fig 9E and 9F). These results suggested that drDnaB is an ATP dependent

5′ → 3′ dsDNA helicase, which largely agrees with the characteristics of DnaBs of other

bacteria (Soultanas and Wigley, 2002; Soni et al., 2003; Nitharwal et al., 2012; Zhang et al.,

2014).

drDnaA and drDnaB show homotypic and heterotypic interactions

The possible interactions of drDnaA and drDnaB proteins were monitored using bacterial

two-hybrid (BACTH) system and co-immunoprecipitation *ex-vivo* and *in vivo*. Both N-

terminal and C-terminal fusions of drDnaA and drDnaB with T18 and T25 BACTH tags were

expressed in *E. coli* BTH101 (*cyaA⁻*) (Fig S4), and the expression of β-galactosidase upon

reconstitution of active CyaA from its T18 and T25 domains through interactions of target

proteins was monitored. The *E. coli* expressing drDnaA and drDnaB tagged with T18 and

T25 through their C-terminals showed both homotypic and heterotypic interactions and the

expression of β-galactosidase activity was observed in both spot and liquid assay (Fig. 10).

Some of these interactions were further confirmed by immunoprecipitation in *E. coli*

BTH101 as described in method. The co-IP results from surrogate *E. coli* agreed with the

results obtained from BACTH analysis (Fig 10 B,C). For *in vivo* interaction studies, the *D.*

radiodurans cells were co-expressed drDnaA or drDnaB with T18 tag and / or polyhistidine

tag on respective plasmids. Total proteins from these cells were immunoprecipitated using

polyhis antibodies and the probable interacting partner was detected using anti-T18

antibodies. The results obtained were similar to that of BACTH and co-IP analysis in

surrogate *E. coli* BTH101 expressing these proteins in different combination (Fig. 10 D). The

results indicated that both these proteins interact through N-terminal domains as reported for

their homologs (Seitz et al., 2000; Messer, 2002; Matthew and Simmons, 2019). Since, C-

terminal domain of drDnaA is required for *oriCI* binding but not for both homotypic and

heterotypic interaction with drDnaA and drDnaB, the interaction of these proteins seems to

be independent of drDnaA interaction to *oriCI*. This suggested that the N-terminal and or

middle region of these proteins seems to involve in oligomerization. This was further

confirmed when DnaAΔCt did not affect its interaction with full length drDnaA and drDnaB

(Fig. 10 E). Further mutational analysis of both deinococcal DnaA and DnaB would require

to exactly map the interaction domains or residues between these proteins.

PprA modulates *in vitro* function of drDnaA but not drDnaB

The ATP hydrolysis is the key feature of bacterial DnaA and DnaB proteins characterized till

date, and this activity is required for the initiation and progression of *oriC* mediated

replication. drDnaA and drDnaB proteins exhibit ATPase activity *in vitro* (Fig. 5, 8).

Therefore, the effect of PprA *per se* on ATPase activity of cognate drDnaA and drDnaB

proteins was checked *in vitro*. Interestingly, PprA did not hydrolyse ATP into ADP by itself.

However, the ATPase activity of drDnaA was significantly reduced in the presence of

equimolar concentration of PprA irrespective of the order of addition of these proteins in

reaction mixture (Fig. 11 A, B).There was no effect of PprA on either ATPase activity (Fig

11C & 11D) or 5′→3′ dsDNA helicase activity (Fig 12) of drDnaB *in vitro*. Since, PprA did

not affect drDnaB function *in vitro*, the possibility of ATP titration by PprA and if that

becomes the limiting factor for ATPase activity of drDnaA was nearly ruled out. Since the

ATPase activity of DnaA is required for *oriC* dependent replication initiation in bacteria

(Messer, 2002; Chodavarapu & Kaguni, 2016), the suppression of ATPase activity of

drDnaA by PprA seems to be an important step in *oriCI* regulation in *D. radiodurans*.

Further, drDnaB is found to be a non-specific DNA binding protein with its preference to

ssDNA while PprA prefers dsDNA. Therefore, if these differences have accounted to the

significant effect of PprA on drDnaA and nearly no effect on drDnaB function cannot be

ruled out.

PprA negatively regulates drDnaA and drDnaB interaction in surrogate *E. coli*

Since, PprA interact with drDnaA or drDnaB with different affinities, the possibility of PprA

affecting drDnaA and drDnaB interactions was hypothesized and tested. The *E. coli* BTH101

cells co-expressing drDnaA and drDnaB interacting combination was co-expressed with

PprA from pSpecpprA (Rajpurohit and Misra, 2013) and the expression of β-galactosidase

activity was monitored. We observed that the expression of β-galactosidase activity arising

from the established interaction of drDnaA and drDnaB (Fig 2) was reduced in the presence

of PprA as compared to control without PprA (Fig. 13A-C). This effect of PprA was

observed on homotypic and heterotypic interactions of both these replication proteins. These

results suggested that PprA interferes with the oligomerization of drDnaA and drDnaB

proteins of *D. radiodurans* as depicted in Fig 13D. In different bacteria, it has been reported

that oligomerization of both DnaA and DnaB followed by their interaction would be required

for faithful DNA replication initiation and elongation. Any defect in oligomerization or

interaction of these proteins affects replication initiation process. Therefore, PprA

interference in the oligomerization of drDnaA and drDnaB could be a crucial step in the

regulation of replication initiation process in *D. radiodurans*.

Discussion

Molecular basis of *oriCI* regulation has been studied mostly in bacteria that harbour limited

copies of single circular chromosome as genetic materials. Very limited information exists on

the regulation of DNA replication in bacteria containing multiple chromosomes and ploid

genome. In *Vibrio cholerae,* a multipartite genome harbouring bacterium, Chr I replication is

initiated by DnaA protein while Chr II replication in initiated by RctB protein that binds and

unwinds an array of repeats present in *oriCII* of Chr II (Egan et al., 2005; Jha et al., 2012;

Ramachandaran et al., 2017; Bruhn et al., 2018). Since Chr I is larger than Chr II in *V.*

cholerae, the temporal regulation of replication initiation could help in synchronization of

replication termination of both the chromosomes before the onset of cell division (Rasmussen

et al., 2007, Val et al., 2016; Val et al, 2014).

D. radiodurans, is another bacterium that harbours polyploid multipartite genome comprising

of 2 chromosomes and 2 plasmids (White et al., 1999). Molecular studies on the regulation of

DNA replication and control of ploidy of genome elements particularly chromosomes have

not been reported yet in this bacterium. Here, for the first time, we have brought forth some

evidence to suggest that primary chromosome (chromosome I) encodes the functional DNA

replication initiation proteins drDnaA and drDnaB and confers a *cis element* comprised of

both *E. coli* type typical DnaA-boxes and non-perfect DnaA-boxes between *drdnaA* and

drdnaN loci. Based on location on chromosome and structure with 13 AT rich DnaA-boxes, 2

GATC regions and 2 AT rich clusters (12 bp and 20 bp) between 2^{nd} and 3^{rd} DnaA box

(White et al., 1999; Luo and Gao, 2018), the cis element was predicted to be *oriC* in

chromosome I (*oriCI*) (Fig. 3). Further, we demonstrated that recombinant drDnaA binds

specifically to *oriCI* while drDnaB binds preferentially and non-specifically to ssDNA over

dsDNA. The presence of ATP has significantly increased the affinity of drDnaA toward *ori*

CI by ~ 8 folds (Table 1). Further, the affinity of drDnaA has been reduced with decreasing

number of DnaA boxes in *oriCI* (Table 1) and its affinity for non-perfect single DnaA box

viz. TTTTCCACA or GTATCCACA was found to be extremely low. The ATPase activity

of drDnaA and drDnaB was stimulated by *oriCI* and ssDNA, respectively. Both these

proteins undergo homotypic and heterotypic oligomerization *in vitro*. Similar characteristics

of DnaA and its interaction with cognate *oriC* have been reported in other bacteria including

B. subtilis (Yoshikawa & Wake, 1993)*, M. tuberculosis* (Zawilak et al., 2004)*, T.*

thermophilus (Schaper et al., 2000)*, Streptomyces coelicolor* and *Spiroplasma citri* (Richter

et al., 1998). Like other eubacteria, the C-terminal domain IV of drDnaA is found to be

essential for binding with its cognate *oriCI*. The presence of ATPase activity in drDnaA and

drDnaB proteins and its stimulation by *oriCI* and ssDNA, respectively, as well as ATP

dependent dsDNA helicase function in drDnaB are some of the typical characteristics known

for these proteins in other bacteria, and thus make them potentially important for regulation

of *oriCI* functions. The DnaBs or its homologs from different bacteria have shown different

polarity in dsDNA helicase functions (Caspi et al., 2001; Naqvi et al., 2003). drDnaB shows

ATP dependent unwinding of duplex DNA in $5' \rightarrow 3'$ polarity, which is like the DnaB of *E.*

coli (LeBowitz & McMacken, 1986)*, H. pylori* (Soni et al., 2003), *M. tuberculosis* (Zhang et

al., 2014) and DnaB like helicase in *Bacillus anthracis* (Naqvi et al., 2003). Initiation of DNA

replication requires the oligomerisation of DnaA and DnaB at the origin of replication site

(Messer, 2002; Zawilak-Pawlik, et al., 2017; Matthew and Simmons, 2019). drDnaA and

drDnaB undergo both homotypic and heterotypic oligomerization and interact with *oriCI*

region and ssDNA, respectively. These findings together suggest that the 13 AT-rich DnaA-

boxes upstream to dr*dna*A is most likely the *origin of replication* in chromosome I, and

drDnaA and drDnaB together constitute the replication initiation complex with *oriCI* of *D.*

radiodurans.

Various mechanisms have been suggested for the regulation of DNA replication. For

instance, in *E. coli* the initiation of DNA replication is regulated by differential affinity of

DnaA and SeqA proteins to methylated and hemi-methylated *oriC* and, regulated inactivation

of DnaA (RIDA) by Hda (Skarstad and Katayama, 2013). CtrA involvement in the regulation

of DNA replication in *Caulobacter crescentus* (Spencer et al., 2009) and *CrtS* protein

interaction with methylated DNA in case of *V. cholerae* (Fournes et al., 2018) have been

reported. Involvement of genome segregation proteins like Soj and Spo0J in case of *Bacillus*

subtilis (Ogura et al., 2003; Murray and Errington, 2008; Scholefield et al., 2011) and ParA

and ParB in the regulation of secondary genome copy number in *D. radiodurans* (Maurya et

al., 2019a, 2019b) have also been suggested. In case of *oriC* mediated initiation of DNA

replication, DnaA binding at *oriC* site triggers the recruitment of (DnaB)6-(DnaC)6

hexameric replicative helicase in several bacteria (Chodavarapu and Kaguni, 2016). Once the

helicase is bound to *oriC*, the dissociation of DnaC from DnaB is required for subsequent

round of DNA replication. Like many other bacteria, the *D. radiodurans* genome also does

not encode DnaC homologue and therefore, the regulation of DNA replication initiation by

drDnaA and drDnaB would be either DnaC independent or it will involve some other proteins

in the system.

Earlier, it has been shown that ATP hydrolysis by DnaA is required for melting of AT rich

region in *oriC* followed by initiation of replication (Messer, 2002; Chodavarapu & Kaguni,

2016). The oligomerisation of both DnaA and DnaB is required for their sequential loading at

oriC, unwinding of DNA duplex and generation of replication fork, respectively (Arai and

Kornberg, 1981; Chodavarapu & Kaguni, 2016; Patel and Picha, 2000). Here, we report that

PprA regulates the macromolecular interactions between drDnaA and drDnaB proteins that

would be needed for drDnaA to initiate replication. Further, we observed that the genomic

content and ploidy levels of *D. radiodurans* cells lacking PprA has increased significantly as

compared to wild type (Fig 1). How does PprA regulate genomic content and ploidy levels in

D. radiodurans is not clear. However, the higher affinity of PprA to drDnaA (Kd= 5.41 X 10^{-}

^{7} ± 1.8 X 10^{-8} [M]) as compared to drDnaB (Kd= 9.71 X 10^{-7} ± 1.1 X 10^{-8} [M]) (Fig 2B, C),

inhibition of ATPase activity of drDnaA by PprA while no effect on ATPase and dsDNA

helicase function of drDnaB *in vitro* (Fig 11, 12) and the loss of drDnaA and drDnaB

interaction in the presence of PprA (Fig 13) are some of the results that indicate the

important role of PprA in the regulation of function of replication initiation proteins.

Therefore, the absence of PprA if has allowed replication to continue at least one cycle and

resulted to an increase in DNA content *per cell* could be speculated. Regulation of DNA

replication by a cell cycle regulatory protein competing for binding to *oriC* site has also been

suggested (Quoin et al., 1998). Therefore, a possibility of PprA competing with drDnaA

binding to *oriCI* was checked and found that PprA does bind to *oriCI* but it was non-specific

(Fig S5). These results together supported the role of PprA in regulation of DNA replication

perhaps by modulating the activities of replication initiation proteins in *D. radiodurans*.

In conclusion, here we have characterized all the components like *oriCI*, drDnaA and drDnaB

involved in *oriC* mediated initiation of bacterial chromosome replication from a multipartite

genome harbouring radioresistant bacterium *D. radiodurans*. All these components showed

canonical functions required for initiation of chromosomal DNA replication as reported in

other bacteria. We found PprA, a protein with pleiotropic functions in this bacterium,

regulates the physicochemical properties of these proteins that are required for their functions

in *oriC* mediated chromosome replication. While further studies would be required to get the

deeper understanding on the role of PprA in regulation of DNA replication in *D. radiodurans*

and if it plays a check point regulator of *oriC* mediated DNA replication during DSB repair.

The available results that include the increased genomic content per cell in Δ*pprA* mutant and

PprA inhibition of drDnaA and drDnaB interaction as well as drDnaA's ATPase activity

together clearly suggest the role of PprA in the initiation of chromosome replication in *D.*

radiodurans. How this activity of PprA is regulated under different growth conditions and if

phosphorylation of PprA as reported earlier (Rajpurohit et al, 2013) provides a regulatory

mechanism in the arrest of DNA replication in quiescent cells engaged in DSB repair during

post irradiation recovery of this bacterium would be investigated independently.

Materials and Methods

Bacterial strains and plasmids

All the bacterial strains and plasmids and oligonucleotides used in this study are given in

Table S1 and Table S2, respectively. *D. radiodurans* R1 (ATCC13939) was grown in TGY

(Bactotryptone (1%), Glucose (0.1%) and Yeast extract (0.5%)) medium at 32°C (Schaefer et

al 2000). *E. coli* strain NovaBlue was used for cloning and maintenance of all the plasmids

while *E. coli* strain BL21(DE3) pLysS was used for the expression of recombinant proteins.

E. coli strain BTH 101 was used for Bacterial Two-Hybrid System (BACTH) based protein-

protein interaction studies. Standard protocols for all recombinant techniques were used as

described in (Green and Sambrook, 2012). Molecular biology grade chemicals and enzymes

were procured from Merck Inc and New England Biolabs, USA. Radiolabeled nucleotides

were purchased from Department of Atomic Energy-Board of Radiation and Isotope

Technology (DAE-BRIT), India.

Cloning, expression and purification of proteins

The dr*dnaA* (DR_0002) and dr*dnaB* (DR_0549) genes were PCR amplified from the genomic

DNA of *D. radiodurans* R1 using pETdnaAF & pETdnaAR, and pETdnaBF & pETdnaBR

primers, respectively. The PCR products were cloned in pET28a (+) plasmid at *Bam*HI and

*Eco*RI sites to yield pETDnaA and pETDnaB plasmids (Table S1). The recombinant

plasmids were sequenced, and correctness of inserts was ascertained. The C-terminal (domain

IV) truncation of drDnaA was made using pETDAΔCtR primer along with pETdnaAF (Table

S2), and resulting plasmid was named pETDAΔCt (Table S1). The recombinant proteins

were purified by nickel affinity chromatography as described earlier (Maurya et al., 2019a).

Fractions showing more than 95% purity were pooled and dialyzed in buffer A containing

200 mM NaCl and further purified from HiTrap Heparin HP affinity columns (GE Healthcare

Life sciences) using a linear gradient of NaCl. Different fractions were analyzed on SDS-

PAGE and fractions containing desired protein were pooled and precipitated with 30% w/v

ammonium sulphate at 8°C. The precipitate was dissolved in R-buffer (20 mM Tris-HCl pH

7.6, 0.1 mM EDTA, 0.5 mM DTT) containing 1 M NaCl. After centrifugation at 16,000 rpm

for 30 minutes, supernatant containing soluble proteins were processed for gel filtration

chromatography. The purified protein was dialyzed in dialysis buffer containing 20 mM Tris-

HCl pH 7.6, 200 mM NaCl, 50% glycerol, 1 mM $MgCl_2$, 0.5 mM DTT and 1 mM PMSF.

Similarly, PprA was expressed from pETpprA and purified as described in (Kota and Misra,

2006) followed by gel filtration as described above.

Protein DNA interaction studies

The recombinant proteins interaction with different forms of DNA was studied using

electrophoretic mobility shift assay (EMSA) as described earlier (Maurya et al., 2019a). In

brief, the 500 bp *oriCI* region (1272-1771) containing 13 repeats of 9 mer of DnaA-boxes

(T(A/G)TA(T)TCCACA) was PCR amplified using OriIFw and OriIRw primers (Table S1

Fig. 1) and gel purified. In addition, different fragments of *oriCI* with 11, 7, 3 and 1 repeat(s)

were also either PCR amplified or chemically synthesised and annealed. DNA substrates

were labeled with [γ-^{32}P] ATP using T4 polynucleotide kinase. Approximately 30 nM labeled

substrate was incubated with different concentrations of recombinant drDnaA in a reaction

buffer B containing 50 mM Tris-HCl (pH 8.0), 75mM KCl, 5mM MgSO$_4$ and 0.1mM DTT at

37°C for 15 min. The reaction was performed in the presence and absence of 1 mM ATP. For

the competition assay, a saturating concentration of protein was incubated with DNA

substrate before different concentrations of non-specific competitor DNA of similar length

(regions of *ftsZ* gene in Table S1) was added and further incubated as per experimental

requirements. Similarly, oligonucleotide 99F (Das and Misra, 2012) was radiolabeled at 5'

end and used as ssDNA substrate for interaction with drDnaB. The interaction of drDnaB

with *oriCI* (dsDNA) was monitored in absence and presence of 1 mM ATP as described for

drDnaA above. The reaction mixtures were separated on 6-8 % native PAGE gels, the gels

were dried, and autoradiograms were developed on X-ray films. The band intensity of

unbound and bound fraction was quantified densitometrically and computed using Image J

2.0 software. The fraction of DNA bound to the protein was plotted against the protein

concentrations by using GraphPad Prism6 and the *Kd* values for the curve fitting of

individual plots were determined as described in (Maurya et al., 2019a).

DNA helicase activity assay

The unwinding of double strand DNA substrates (with 5' overhang as well as 3' overhang)

by drDnaB (2µM) in the presence of different combinations of 1 mM ATP and 2 µM PprA

was monitored using fluorescent resonance energy transfer (FRET) approach using 96-well

plates as described in (Khairnar et al., 2019). In brief, the substrates used were FAM and

BHQ labelled complementary oligonucleotides (Table S1). For preparation of overhang

dsDNA, the oligonucleotides of complementary strands producing 5'overhang or 3' overhang

were mixed, heated at 95 □C for 10 minutes and then slowly annealed by switching off the

heating block for overnight. The helicase assays were performed in volume of 100 µL in

helicase buffer (25 mM Tris pH 7.6, 10 mM NaCl, 1 mM MgCl$_2$, 100 µM DTT, 100 µg/ml

BSA, 25 mM KCl and 2 % glycerol) at 37 □C. To avoid re-annealing of unwound DNA,

excess of DNA trap complementary to the BHQ labelled DNA was added to each reaction

mixture. The reaction was excited at 490 nm and emission was recorded at 520 nm. The

fluorescence signals were monitored at an interval of 30 seconds using microplate reader

(Biotek Synergy H1).

Similarly, gel-based DNA helicase activity assay of drDnaB was carried out as descried in

(Khairnar et al., 2019). In brief, the 3' overhang substrate was made by annealing the [^{32}P]

radiolabelled 3OV99F (5' TTTTTGCGGTTCATATGGAATTCC3') with 99F

(5'GGAATTCCAATGAACCGCAAAACCGCAAAAACCGTACCGA3') as described in

(Das and Misra, 2012). Similarly, 5' overhang substrate was generated by annealing of

radiolabelled 5OV99F (5'TCGGTACGGTTTTTGCGGTTCATA3') with 99F

oligonucleotides. The annealed substrates were incubated with 2 µM drDnaB with or without

2 µM PprA proteins in the presence and absence of 1 mM ATP or ATP-γ-S in helicase assay

buffer at 37˚C. Aliquots were taken at different time interval and reactions were stopped with

stop solution (20 % glycerol, 20 mM EDTA, 125µg Proteinase K, 0.2 % SDS, 50 mM KCl)

at 37 □C for 10 minutes. The products were separated on 8 % denaturing PAGE, dried and

exposed to X-ray films. The autoradiograms were developed and documented.

Bacterial Two-Hybrid (BACTH) system assay

drDnaA and drDnaB interaction was monitored in the absence and presence of PprA using

bacterial two-hybrid system (BACTH), described in (Maurya et al., 2016, 2018). In brief, the

coding sequence of DR_0002 (drDnaA) was cloned at *Bam*HI and *Eco*RI sites in pUT18,

pUT18C and pKT25 plasmids to yield pUT18DA, pUT18CDA & pKTDA, respectively.

Similarly, coding sequences of DR_0549 (drDnaB) was cloned at *Kpn*I and *Eco*RI sites in

pUT18, pUT18C and pKT25 plasmids to yield pUT18DB, pUT18CDB & pKTDB,

respectively. The construction of pKNTDA and pKNTDB plasmids has been described in

(Maurya et al., 2019b) and pUTpprA in (Kota et al., 2014). *E. coli* strain BTH101 (*cyaA*⁻)

was co-transformed with these plasmids in different combinations and expression of β

galactosidase was monitored as described earlier. *E.coli* BTH101 harbouring only vectors

were used as negative control while those harbouring pUTEFA and pKNTEFZ (Modi and

Misra, 2014) were used as positive control. For monitoring the effect of PprA on drDnaA and

drDnaB interactions, the PprA was expressed on pSpecpprA (Rajpurohit and Misra, 2013)

while p11559 vector was used for negative control. Expression of β-galactosidase as an

indication of protein-protein interaction was monitored using spot assay and in liquid culture

and β-galactosidase activity was calculated in Miller units as described in (Battesti et al.,

2012).

Co-immunoprecipitation assay

Protein-protein interactions in surrogate *E. coli* (BTH101) and *D. radiodurans* were

monitored by co-immunoprecipitation as described in (Maurya et al 2016; Maurya et al.,

2018). In brief, the total proteins of the recombinant BTH101 cells co-expressing drDnaA

and drDnaB proteins on BACTH plasmids (Table S1) were immunoprecipitated using

polyclonal antibodies against T25 and counterpart was detected using T18 antibodies. PprA's

effect on drDnaA and drDnaB interaction was studied by *in trans* expression of PprA on

pSpecpprA. Signals were detected using anti-mouse secondary antibodies conjugated with

alkaline phosphatase using BCIP/NBT substrates (Roche Biochemical, Mannheim).

Similarly, for monitoring drDnaA, drDnaAΔCt and drDnaB in *D. radiodurans* by co-

immunoprecipitation, the coding sequences of drDnaA, drDnaAΔCt and drDnaB tagged with

polyhistidine were PCR amplified using pETHisFw and pETHisRw primers and cloned in

pRADgro plasmid at *Apa*I and *Xba*I sites to yield pRADhisDA, pRADhisDACt and

pRADhisDB, respectively (Table S1). In addition, the coding sequences of T18 tagged

drDnaA and drDnaB were PCR amplified using BTHF(pv) and BTHR(pv) primers from

pUT18DAand pUT18DB plasmids and cloned in p11559 plasmid at *Nde*I and *Xho*I sites to

yield pV18DA and pV18DB, respectively (Table S1). These plasmids were co-transformed in

different combinations in *D. radiodurans* and induced with 5 mM IPTG as requried. The cell-

free extracts of *D. radiodurans* expressing drDnaA, drDnaAΔCt and drDnaB in different

combinations were prepared and immunoprecipitated using polyhistidine antibodies as

described earlier (Maurya et al., 2019a). The immunoprecipitates were purified using Protein

G Immunoprecipitation Kit (Cat. No. IP50, Sigma-Aldrich Inc.) and were separated on SDS-

PAGE for western blotting using monoclonal antibodies against T18 as described above. To

show interaction of PprA with drDnaA or drDnaB, wild type cells of *D. radiodurans*

expressing native PprA were transformed with pV18DA or pV18DB and expression of T18

tagged drDnaA or drDnaB was induced with 5 mM IPTG. The cell free extracts of these

transformants were prepared and immunoprecipitated using Anti-PprA antibodies. The

purified immunoprecipitates were processed for western blotting using T18 antibodies as

described above.

Surface Plasmon Resonance

Interaction of drDnaA and drDnaB with PprA was also investigated using surface plasmon

resonance (SPR; Autolab Esprit, Netherland) as described in (Maurya et al., 2018). For this,

20 μM PprA was immobilized on a bare gold sensor chip using EDC-NHS chemistry as

described in user manual at 20°C which results about 200 response units in running buffer

[20 mM Tris (pH 7.6)]. The different concentrations (4 – 20 μM) of recombinant drDnaA or

drDnaB were used in the mobile phase and incubated with 1 mM ATP and 1 mM $MgCl_2$ and

passed over the PprA-bound sensor chip in one channel. Reaction buffer containing 20 mM

Tris (pH 7.6), 1 mM ATP and 1 mM $MgCl_2$ was used as buffer control to flow from another

channel over immobilized PprA. The response units for each concentration of proteins were

recorded and normalized with buffer control. Further, data were processed using the inbuilt

Autolab kinetic evaluation software (V5.4) to find dissociation constant and plotted after

curve smoothening using GraphPad Prizm6 software.

Thin Layer Chromatography for ATPase assay

ATPase activity of drDnaA, drDnaAΔCt and drDnaB was measured as the release of [^{32}P]

αADP from [^{32}P] αATP using Thin Layer Chromatography (TLC) as described earlier

(Maurya et al., 2019a). In brief, different concentrations (0 - 3 μM) of drDnaA or drDnaB

were mixed with 30 nM [^{32}P] αATP in the absence and presence of 0.2 pmol dsDNA or 0.1

pmol ssDNA in 10 μl containing buffer B, respectively. The reaction mixture was incubated

at 37$^{°}$C for 30 min. For monitoring the effect of PprA on ATPase activity, 2 μM PprA was

used with 2 μM drDnaA or 2 μM drDnaB as described above. The reaction mixture was

stopped at different time period (0 - 30 min) with 10 mM EDTA solution and 1 μl of it was

spotted on PEI-Cellulose F^{+} TLC sheet. The spots were air-dried, and products were

separated on a solid support in a buffer system containing 0.75 M KH_2PO_4 / H_3PO_4 (pH 3.5).

The TLC sheets were air dried, exposed to X-ray film and autoradiograms were developed.

Spot intensities of samples were determined by densitometry using Image J 2.0 software. The

percentage ratio of ADP to ATP was calculated and plotted using the GraphPad Prizm6

software.

Fluorescence microscopy

The *D. radiodurans* R1 (WT) and its Δ*pprA* mutant cells were grown till the exponential

phase, and the equal number of cells were incubated with 2.5µg 4',6-diamidino-2-

phenylindole dihydrochloride (DAPI) per / ~10^8 cells for nucleoid staining. These cells were

washed twice in phosphate buffer saline and then mounted on slides coated with 0.8%

agarose. Cells were imaged in DIC channel (automatic exposure) and DAPI channel (50

milliseconds exposure) under identical conditions for both WT and Δ*pprA* mutant using an

Olympus IX83 inverted fluorescence microscope equipped with an Olympus DP80 CCD

monochrome camera. Large number of cells (~150) from both WT and Δ*pprA* mutant were

processed and fluorescence intensity of DAPI stained nucleoid was measured using 'intensity

profile' tool in cellSens1.16 software installed with microscope. Mean fluorescence intensity

was plotted against sample type using GraphPad Prizm6 software. In addition, relative

frequency distribution (%) was also plotted against fluorescence intensity using GraphPad

Prizm6 software.

Determination of ploidy in wild type and Δ*pprA* mutant

The amount of DNA and copy number of all four genome elements like chromosome I,

chromosome II, megaplasmid and plasmid per cell of wild type and Δ*pprA* mutant was

determined as described earlier (Maurya et al., 2019a). In brief, the exponentially growing

wild type and Δ*pprA* cells were adjusted at optical density at 600 nm and their cell numbers

were determined using a Neubauer cell counter. The collected cells were washed in PBS

followed by 70 % ethanol wash and lysed in solution containing 10 mM Tris pH 7.6, 1 mM

EDTA, 4 mg / ml lysozyme. The cells were incubated at 37°C and cell debris was removed

by centrifugation at 10000 rpm for 5 min. The integrity of isolated genomic DNA was

confirmed by 0.8% agarose gel electrophoresis. DNA content was measured as OD260nm

and the genomic copy number was determined using quantitative real-time PCR as described

in (Breuert et al., 2006). For quantification of genome copy number, two different genes

were taken per replicon showing similar PCR efficiency. For examples, the *ftsZ* and *ftsE* for

chromosome I, DR_A0155 and DR_A0002 for chromosome II, DR_B003 and DR_B0076

for megaplasmid and DR_C001 and DR_C018 for small plasmid (Table S2). PCR efficiency

of each gene was analysed and was found to be > 96% for each (data not shown). We have

followed the Minimum Information for Publication of Quantitative Real-Time PCR

Experiments (MIQE) guidelines (Bustin et al., 2009) for qPCR using Roche Light cycler and

the cycle threshold (Cp) values were determined. The experiment was performed using three

independent biological replicates for each sample. The Cp value for each gene of respective

replicon was compared with a dilution series of a PCR product of known concentration, as a

standard (Fig S1). The copy number of each replicon by both genes per cell was determined

using the cell number present at the time of cell lysis. An average of copy number from two

genes per replicon in WT and Δ*pprA* mutant was calculated and represented along with bio-

statistical analysis.

Acknowledgements: Authors are grateful to Prof. V Nagaraja and Prof. D N Rao, Indian

Institute of Science, Bangalore, for their intellectual comments on the findings and other

valuable suggestions. We thank Ms Shruti Mishra and Dr Sheetal Uppal and Dr C Rajani

Kant for their technical comments on the manuscript. GKM and NP are grateful to DAE and

DST, Govt of India, for DAE and INSPIRE-research fellowships, respectively.

Competing Interests: Authors have no financial or non-financial competing interests.

References

1. Adachi, M., Hirayama, H., Shimizu, R., Satoh, K., Narumi, I. & Kuroki, R. (2014).

Interaction of double-stranded DNA with polymerized PprA protein from

Deinococcus radiodurans. Protein Science 23, 1349-1358.

2. Adachi, M., Shimizu, R., Shibazaki, C., Satoh, K., Fujiwara, S., Arai, S., Narumi, I. &

Kuroki, R. (2019). Extended structure of pleiotropic DNA repair-promoting protein

PprA from *Deinococcus radiodurans*. FASEB J 33, 3647-3658.

3. Arai, K.I., & Kornberg, A. (1981). Mechanism of dnaB protein action. II. ATP

hydrolysis by dnaB protein dependent on single-or double-stranded DNA. J Biol.

Chem. 256, 5253-5259.

4. Battesti, A. & Bouveret, E. (2012). The bacterial two-hybrid system based on

adenylate cyclase reconstitution in *Escherichia coli*. Methods 58, 325-334.

5. Biswas, E. E., Barnes, M. H., Moir, D. T., & Biswas, S. B. (2009). An essential DnaB

helicase of *Bacillus anthracis*: identification, characterization, and mechanism of

action. *J. Bacteriol*, *191*(1), 249-260.

6. Blaesing F., Weigel, C., & Messer, W. (2000). Analysis of the DNA

binding domain of *Escherichia coli* DnaA protein. Mol Microbiol

36, 557-569.

7. Breuert, S., Allers, T., Spohn, G., & Soppa, J. (2006). Regulated

polyploidy in halophilic archaea. PLoS One 1, e92

8. Bruhn, M., Schindler, D., Kemter, F.S., Wiley, M.R., Chase, K., Koroleva, G.I.,

Palacios, G., Sozhamannan, S. & Waldminghaus, T. (2018). Functionality of two

origins of replication in *Vibrio cholerae* strains with a single chromosome. Front

Microbiol 9, 2932.

9. Bustin, S.A., Benes, V., Garson, J.A., Hellemans, J., Huggett, J., Kubista, M.,

Mueller, R., Nolan, T., Pfaffl, M.W., Shipley, G.L. & Vandesompele, J. (2009). The

MIQE guidelines: minimum information for publication of quantitative real-time PCR

experiments. Clin. Chem. 55, 611-622

10. Caspi, R., Pacek, M., Consiglieri, G., Helinski, D. R., Toukdarian, A., & Konieczny,

I. (2001). A broad host range replicon with different requirements for replication

initiation in three bacterial species. EMBO Journal, 20, 3262-3271.

11. Chodavarapu, S., & Kaguni, J. M. (2016). Replication initiation in bacteria. In *The*

Enzymes (Vol. 39, pp. 1-30). Academic Press.

12. Cox MM, Battista JR. 2005. *Deinococcus radiodurans* - the consummate survivor.

Nat Rev Microbiol 3, 882-892.

13. Crooks, G. E., Hon, G., Chandonia, J. M., & Brenner, S. E. (2004). WebLogo: a

sequence logo generator. Genome Res. 14(6), 1188-1190.

14. Das, A.D., & Misra, H.S. (2012). DR2417, a hypothetical protein characterized as a

novel β-CASP family nuclease in radiation resistant bacterium, *Deinococcus*

radiodurans. Biochim Biophys Acta (GS) 1820,1052-1061.

15. Devigne, A., Mersaoui, S., Bouthier-de-la-Tour, C., Sommer, S., & Servant, P.

(2013). The PprA protein is required for accurate cell division of γ-irradiated

Deinococcus radiodurans bacterium. DNA Repair 12**,** 265-272.

16. Devigne, A., Guérin, P., Lisboa, J., Quevillon-Cheruel, S., Armengaud, J., Sommer,

S, Bouthier de la Tour C, and Servant, P. (2016). PprA protein is involved in

chromosome segregation via its physical and functional interaction with DNA gyrase

in irradiated *Deinococcus radiodurans* bacteria. mSphere, 1, e00036-15.

17. Egan, E. S., and Waldor, M. K. (2003). Distinct replication requirements for the

643   two *Vibrio cholerae* chromosomes. Cell 114, 521–530.

644   18. Egan, E. S., Fogel, M. A., and Waldor, M. K. (2005). Divided genomes: negotiating

645   the cell cycle in prokaryotes with multiple chromosomes. Mol. Microbiol. 56, 1129–

646   1138.

647   19. Fournes, F., Val, M. E., Skovgaard, O., and Mazel, D. (2018). Replicate once per cell

648   cycle: replication control of secondary chromosomes. Front. Microbiol. 9, 1833.

649   20. Fujikawa, N., Kurumizaka, H., Nureki, O., Terada, T., Shirouzu, M., Katayama, T., &

650   Yokoyama, S. (2003). Structural basis of replication origin recognition by the DnaA

651   protein. Nucleic Acids Research, 31, 2077-2086.

652   21. Green, M.R., & Sambrook, J. Molecular cloning : a laboratory manual. Cold Spring

653   Harbor, N.Y: Cold Spring Harbor Laboratory Press. 2012.

654   22. Hansen, M.T. (1978). Multiplicity of genome equivalents in the radiation-resistant

655   bacterium *Micrococcus radiodurans*. J Bacteriol 134, 71-75.

656   23. Huang, H., Song, C. C., Yang, Z. L., Dong, Y., Hu, Y. Z., & Gao, F. (2015).

657   Identification of the replication origins from *cyanothece* ATCC 51142 and their

658   interactions with the DnaA protein: from in silico to in vitro studies. Front

659   Microbiol. *6*, 1370.

660   24. Jha, J. K., Baek, J. H., Venkova-Canova, T., and Chattoraj, D. K. (2012).

661   Chromosome dynamics in multichromosome bacteria. Biochim. Biophys. Acta 1819,

662   826–829.

663   25. Karimova, G., Pidoux, J., Ullmann, A., and Ladant, D. (1998) A bacterial two-hybrid

664   system based on a reconstituted signal transduction pathway. *Proc. Nat. Acad. Sci.*

665   *USA* **95**, 5752-5756.

26. Khairnar, N.P., Maurya, G.K., Pandey, N., Das, A., Misra, H.S. (2019). DrRecQ

regulates guanine quadruplex DNA structure dynamics and its impact on

radioresistance in *Deinococcus radiodurans*. Mol Microbiol. 112, 854-865.

27. Koch, B., Ma, X., and Lobner-Olesen, A. (2010). Replication of *Vibrio*

cholerae chromosome I in *Escherichia coli*: dependence on dam methylation. J.

Bacteriol. 192, 3903–3914.

28. Kota, S., Misra, H.S. (2006). PprA: A protein implicated in radioresistance of

Deinococcus radiodurans stimulates catalase activity in *Escherichia coli*. Appl

Microbiol Biotechnol. 72,790-796.

29. Kota, S., Charaka, V.K., Ringgaard, S., Waldor, M.K., & Misra, H.S. (2014). PprA

Contributes to *Deinococcus radiodurans* Resistance to Nalidixic Acid, Genome

Maintenance after DNA Damage and Interacts with Deinococcal Topoisomerases.

PLoS One 9, e85288.

30. Kota, S., Rajpurohit, Y. S., Charaka, V. K., Satoh, K., Narumi, I., & Misra, H. S.

(2016). DNA gyrase of *Deinococcus radiodurans* is characterized as type II bacterial

topoisomerase and its activity is differentially regulated by PprA in

vitro. Extremophiles, 20, 195-205.

31. LeBowitz, J.H. & McMacken, R. (1986). The *Escherichia coli* dnaB

replication protein is a DNA helicase. J. Biol. Chem., 261, 4738-4748.

32. Leonard, A.C, & Grimwade, J.E. (2015). The orisome: structure and function. Front

Microbiol. 6,545

33. Luo, H., & Gao, F. (2018). DoriC 10.0: an updated database of replication origins in

prokaryotic genomes including chromosomes and plasmids. Nucleic Acids Research,

47, D74–D77.

34. Majka, J., Jakimowicz, D., Messer, W., Schrempf, H., Lisowski, M.
and Zakrzewska-Czerwinska, J. (1999) Interactions of the *Streptomyces lividans*
initiator protein DnaA with its target. Eur. J. Biochem.
260, 325-335.

35. Matthews, L. A., & Simmons, L. A. (2019). Cryptic protein interactions regulate
DNA replication initiation. Mol Microbiol 111, 118-130.

36. Maurya, G.K., Modi, K., and Misra, H.S. (2016). Divisome and segrosome
components of *Deinococcus radiodurans* interact through cell division regulatory
proteins. Microbiology (UK) 162, 1321-1334.

37. Maurya, G. K., Modi, K., Banerjee, M., Chaudhary, R., Rajpurohit, Y. S., and Misra,
H. S. (2018) Phosphorylation of FtsZ and FtsA by a DNA Damage-Responsive
Ser/Thr Protein Kinase Affects Their Functional Interactions in *Deinococcus*
radiodurans. mSphere 3, 4.

38. Maurya, G. K., Kota, S., Kumar, N. N., Tewari, R., & Misra, H. S. (2019a). ParA
proteins of secondary genome elements crosstalk and regulate radioresistance through
genome copy number reduction in *Deinococcus radiodurans*. Biochem J. 476, 909-
930.

39. Maurya, G. K., Kota, S., & Misra, H. S. (2019b). Characterisation of ParB encoded on
multipartite genome in *Deinococcus radiodurans* and their roles in
radioresistance. Microbiol Res 223, 22-32.

40. McHenry, C.S. (2011) Bacterial replicases and related polymerases. Curr Opin Chem
Biol. 15, 587–594.

41. Messer,W., Blaesing,F., Majka,J., Nardmann,J., Schaper,S., Schmidt,A.,
Seitx,H., Speck,C., Tungler,D., Wegrzyn,G., Weigel,C., Welzeck,M. and

Zakrzewska-Czerwinska, J. (1999). Functional domains of DnaA proteins. Biochimie 81, 819-825.

42. Messer, W. (2002). The bacterial replication initiator DnaA. DnaA and oriC, the bacterial mode to initiate DNA replication. FEMS Microbiol Rev 26, 355-374.

43. Misra, H.S., Khairnar, N.P., Kota S., Shrivastava, S., Joshi, V.P., and Apte S.K. (2006) An Exonuclease I sensitive DNA repair pathway in *Deinococcus radiodurans*: a major determinant of radiation resistance. *Mol. Microbiol.* **59**,1308-1316.

44. Misra, H.S., Rajpurohit, Y.S., & Kota, S. (2013). Physiological and molecular basis of extreme radioresistance in *Deinococcus radiodurans*. Curr. Sci. 104,194-205.

45. Misra, H. S., Maurya, G. K., Kota, S., & Charaka, V.K. (2018) Maintenance of multipartite genome system and its functional significance in bacteria. J. Genet. 97, 1013–1038.

46. Modi, K.M., Misra, H.S. (2014). Dr-FtsA, an actin homologue in *Deinococcus radiodurans* differentially affects Dr-FtsZ and Ec-FtsZ functions *in vitro*. PLoS ONE 9, e115918.

47. Moriya, S., Fukuoka, T., Ogasawara, N., & Yoshikawa, H. (1988). Regulation of initiation of the chromosomal replication by DnaA-boxes in the origin region of the *Bacillus subtilis* chromosome. *EMBO J 7*, 2911-2917.

48. Mott, M.L. & Berger J.M. (2007). DNA replication initiation: mechanisms and regulation in bacteria. Nat Rev Microbiol. 5, 343-354.

49. Murray, H., & Errington, J. (2008). Dynamic control of the DNA replication initiation protein DnaA by Soj/ParA. Cell 135, 74–84.

50. Naqvi, A., Tinsley, E. & Khan, S.A. (2003). Purification and characterization of the PcrA helicase of *Bacillus anthracis*. J Bacteriol 185, 6633–6639.

51. Narumi, I., Satoh, K., Cui, S., Funayama, T., Kitayama, S., & Watanabe, H. (2004).

PprA: a novel protein from *Deinococcus radiodurans* that stimulates DNA

ligation. Mol Microbiol 54, 278-285.

52. Nitharwal, R. G., Verma, V., Subbarao, N., Dasgupta, S., Choudhury, N. R., & Dhar,

S. K. (2012). DNA binding activity of *Helicobacter pylori* DnaB helicase: the role of

the N-terminal domain in modulating DNA binding activities. FEBS J, 279, 234-250.

53. Ogura, Y., Ogasawara, N., Harry, E.J., Moriya, S. (2003). Increasing the ratio of Soj

to Spo0J promotes replication initiation in *Bacillus subtilis*. J Bacteriol 185, 6316–

6324.

54. Patel, S. S., & Picha, K. M. (2000). Structure and function of hexameric

helicases. Ann Rev Biochem 69, 651-697.

55. Quon, K.C., Yang, B., Domian, I.J., Shapiro, L., & Marczynski, G.T.

(1998). Negative control of bacterial DNA replication by a cell cycle regulatory

protein that binds at the chromosome origin. Proc Natl Acad Sci 95, 120–125.

56. Rajpurohit, Y.S., & Misra, H.S. (2013). Structure-function study of deinococcal

serine/threonine protein kinase implicates its kinase activity and DNA repair protein

phosphorylation roles in radioresistance of *Deinococcus radiodurans*. Int J Biochem

Cell Biol 45, 2541-2552.

57. Ramachandran, R., Jha, J., Paulsson, J., and Chattoraj, D. (2017). Random versus cell

cycle-regulated replication initiation in bacteria: insights from studying *Vibrio*

cholerae chromosome 2. Microbiol. Mol. Biol. Rev. 81, e00033-16.

58. Rasmussen, T., Jensen, R. B., and Skovgaard, O. (2007). The two chromosomes

of *Vibrio cholerae* are initiated at different time points in the cell cycle. EMBO J. 26,

3124–3131.

59. Richter, S., Hess, W. R., Krause, M. & Messer, W. (1998). Unique organization of the

dnaA region from *Prochlorococcus marinus* CCMP1375, a marine cyanobacterium.

Mol. Gen. Genet. 257, 534-541.

60. Roth, A. and Messer, W. (1995) The DNA binding domain of the

initiator protein DnaA. EMBO J. 14, 2106-2111.

61. Schäefer, M., Schmitz, C., Facius, R., Horneck, G., Milow, B., Funken, K.H., &

Ortner, J. (2000). Systematic Study of Parameters Influencing the Action of Rose

Bengal with Visible Light on Bacterial Cells: Comparison Between the Biological

Effect and Singlet-Oxygen Production. Photochem Photobiol 71, 514-523.

62. Schaper, S., Nardmann, J., Lu «der, G., Lurz, R., Speck, C., &

Messer, W. (2000). Identification of the chromosomal replication

origin from *Thermus thermophilus* and its interaction with the replication initiator

DnaA. J Mol Biol 299, 655-665.

63. Scholefield, G., Whiting, R., Errington, J., & Murray, H. 2011. Spo0J regulates the

oligomeric state of Soj to trigger its switch from an activator to an inhibitor of DNA

replication initiation. Mol Microbiol 79, 1089–1100.

64. Seitz, H., Weigel, C., & Messer, W. (2000). The interaction domains of the DnaA and

DnaB replication proteins of *Escherichia coli*. Mol. Microbiol., *37*(5), 1270-1279.

65. Skarstad, K., & Katayama, T. (2013). Regulating DNA Replication in Bacteria. Cold

Spring Harb Perspect Biol. 5, a012922.

66. Slade, D., & Radman, M. (2011). Oxidative stress resistance in *Deinococcus*

radiodurans. Microbiol Mol Biol Rev 75, 133-191.

67. Soni, R. K., Mehra, P., Choudhury, N. R., Mukhopadhyay, G., & Dhar, S. K. (2003).

Functional characterization of *Helicobacter pylori* DnaB helicase. *Nucleic Acids*

Research, *31*(23), 6828-6840.

68. Soultanas, P., & Wigley, D. B. (2002). Site-directed mutagenesis reveals roles for
conserved amino acid residues in the hexameric DNA helicase DnaB from *Bacillus*
stearothermophilus. Nucleic Acids Res 30, 4051-4060.
790

791   69. Spencer, W., Siam, R., Ouimet, M-C, Bastedo, D.P., & Marczynski, G.T. (2009).
792   CtrA, a Global Response Regulator, Uses a Distinct Second Category of Weak DNA
793   Binding Sites for Cell Cycle Transcription Control in *Caulobacter crescentus*. J Bact
         191, 5458-5470.

70. Sutton, M.D. and Kaguni, J.M. (1997). The *Escherichia coli* dnaA gene:
four functional domains. J Mol Biol 12, 546-561.

71. Val, M. E., Soler-Bistue, A., Bland, M. J., & Mazel, D. (2014). Management of
multipartite genomes: the *Vibrio cholerae* model. Curr Opin Microbiol 22, 120–126.

72. Val, M. E., Marbouty, M., de Lemos Martins, F., Kennedy, S. P., Kemble, H., Bland,
M. J., Possoz, C., Koszul, R., & Skovgaard, O. (2016). A checkpoint control
orchestrates the replication of the two chromosomes of *Vibrio cholerae*. *Sci.*
Adv. 2:e1501914.

73. White, O., Eisen, J.A., Heidelberg, J.F., Hickey, E.K., Peterson, J.D., Dodson, R.J.,
Haft, D.H., Gwinn, M.L., Nelson, W.C., Richardson, D.L., Moffat, K.S., Qin, H.,
Jiang, L., Pamphile, W., Crosby, M., Shen, M., Vamathevan, J.J., Lam, P., McDonald,
L., Utterback, T., Zalewski, C., Makarova, K.S., Aravind, L., Daly, M.J., Minton,
K.W., Fleischmann, R., Ketchum. K.A, Nelson, K.E., Salzberg, S., Smith, H.O.,
Venter, J.C., & Fraser, C.M. (1999). Genome sequence of the radioresistant bacterium
Deinococcus radiodurans R1. Science 286, 1571–1577.

74. Yao, N.Y., & O'Donnell, M. (2010). SnapShot: the replisome. Cell. 141, 1088, e1081.

75. Yoshikawa, H. & Wake, R.G. (1993). Initiation and termination of
chromosome replication. In: *Bacillus subtilis* and Other Gram-Positive Bacteria:

Biochemistry, Physiology, and Molecular Genetics

(Sonenshein, A.L., Hoch, J.A. and Losick, R., Eds.), pp. 507-528.

ASM, Washington, DC.

76. Zahradka, K., Slade, D., Bailone, A., Sommer, S., Averbeck, D., Petranovic, M.,

Lindner, A.B., & Radman, M. (2006). Reassembly of shattered chromosomes in

Deinococcus radiodurans. Nature 443, 569–573.

77. Zakrzewska-Czerwińska, J., Jakimowicz, D., Zawilak-Pawlik, A., & Messer, W.

(2007). Regulation of the initiation of chromosomal replication in bacteria. FEMS

Microbiol. Rev. 31, 378–387.

78. Zawilak, A., Cebrat, S., Mackiewicz, P., Krol-Hulewicz, A., Jakimowicz, D., Messer,

W., Gosciniak, G. & Zakrzewska-Czerwinska, J. (2001). Identification of a putative

chromosomal replication origin from *Helicobacter pylori* and its interaction with the

initiator protein DnaA. Nucleic Acids Res. 29, 2251-2259

79. Zawilak, A., Agnieszka, K. O. I. S., Konopa, G., Smulczyk-Krawczyszyn, A., &

Zakrzewska-Czerwińska, J. (2004). *Mycobacterium tuberculosis* DnaA initiator

protein: purification and DNA-binding requirements. *Biochemical Journal*, *382*(1),

247-252.

80. Zawilak-Pawlik, A., Nowaczyk, M., & Zakrzewska-Czerwińska, J. (2017). The role

of the N-terminal domains of bacterial initiator DnaA in the assembly and regulation

of the bacterial replication initiation complex. *Genes*, *8*(5), 136.

81. Zhang, H., Zhang, Z., Yang, J., & He, Z. G. (2014). Functional characterization of

DnaB helicase and its modulation by single-stranded DNA binding protein in

Mycobacterium tuberculosis. FEBS J 281, 1256-1266.

Table 1. Dissociation constant (Kd) of drDnaA with *oriI* and its repeat variants with different
number of DnaA boxes was measured in the presence and absence of ATP.

Number of DnaA boxes	ATP	Dissociation constant (Kd) (µM)
13 (TTATCCACA)$_{13}$	-	1.78±0.23
13 (TTATCCACA)$_{13}$	+	0.243±0.021
11(TTATCCACA)$_{11}$	+	0.241±0.01
7(TTATCCACA)$_7$	+	0.373±0.01
3(TTATCCACA)$_3$	+	1.08±0.12
1 (TTATCCACA)	+	2.88±0.28
1 (TTTTCCACA) non-perfect repeat	+	4.21±0.15
1 (GTATCCACA) non-perfect repeat	+	4.36±0.18

Figures

Fig. 1. Change in genomic content in *pprA* deletion mutant of *D. radiodurans*. *D. radiodurans* R1 (WT) and its *pprA* mutant (Δ*pprA*) were grown to mid-logarithmic phase. The amount of DNA per cell was measured spectrophotometrically (A) and through DAPI staining of nucleoid (B). The change in DAPI fluorescence in wild type and Δ*pprA* mutant cells was scanned in ~150 cells and relative frequency of cells containing different amount of DAPI stained nucleoid was estimated (C). Similarly, wild type and mutant populations were determined for the copy number of chromosome I (ChrI), chromosome II (ChrII), megaplasmid (Mp) and small plasmid (Sp) (D). Data presented in A, B and D are mean ± SD (n=9). Data given in C are from the wild type cell population (n=148) and mutant population (n=159).

Fig 2. PprA interaction with drDnaA and drDnaB of *D. radiodurans*. *E. coli* BTH 101 cells co-expressing drDnaA-C18 (DA-C18), drDnaA-C25 (DA-C25), drDnaB-C25 (DB-C25), PprA-C18 in different combinations were detected for expression of β-galactosidase in spot test and in liquid (A). The bacterial two hybrid system vectors pUT18 and pKNT25 were used as negative control while *E. coli* FtsA (EcFtsA) and FtsZ (EcFtsZ) were used a positive control. Recombinant PprA interactions with recombinant drDnaA (B) and drDnaB (C) of *D. radiodurans* were studied by surface plasmon resonance. *Ex vivo* interaction of PprA expressing from PprA-C18 and C-terminal truncated DnaA with histidine tag (HisAΔCt) was monitored in surrogate *E. coli* (D). *In vivo* interaction of native PprA expressing on chromosome and drDnaA (DA-C18), and drDnaB (DB-C18) on plasmids was monitored in *D. radiodurans* R1. Total proteins were precipitated with antibodies against polyhistidine (D) and PprA (E). Perspective interacting partners were detected using antibodies against T18 domain of CyaA. Sizes of fusions were compared with molecular weight marker (M). C in panel E indicates plasmid control.

Fig. 3. Organization of putative origin of replication in chromosome I (*oriCI*) of *D. radiodurans.* A *cis* element region (1273-1772) located between *dnaN* and *dnaA* in chromosome I of *D. radiodurans* was analyzed for putative *oriC* signature. This fragment is found containing 13 copies of conserved DnaA boxes, 2 GATC motif and 2 AT rich regions and thus predicted to be *oriCI in D. radiodurans*.

Fig. 4. Recombinant drDnaA interaction with putative *oriCI*. The *oriCI* (*oriI*) DNA was
radiolabeled and incubated with increasing concentration of recombinant drDnaA protein in
the absence (A, B) and presence (C, D) of 1mM ATP. The nucleoprotein complexes were
analyzed on native PAGE. The 5 (1), 10 (2), 20 (3) and 50 (4) fold higher molar
concentration of non-specific dsDNA (NS-DNA) was added into reaction mixture and
analyzed on non-denaturing PAGE. Band intensity of free form and bound form of DNA was
estimated densitometrically and plotted as mean ± SD (n=3) of bound fraction in percentage.

A   B   C

Fig. 5. ATPase activity in recombinant drDnaA. An increasing concentration (0.5, 1, 1.5, 2.0

and 3 µM) of recombinant drDnaA (DnaA) was incubated with $[^{32}P]$-αATP (αATP) in the

absence (A) and presence (B) of *oriCI* (*oriI*)and generation of $[^{32}P]$-αADP (αADP) product

was detected on TLC after autoradiography. Spot intensity was quantified densitometrically

and percent of ADP/ATP ratios were plotted as a function of drDnaA concentration (C).

Results were analyzed using student t-test and significant difference in data sets with p values

of 0.05 or less is shown as (*).

Fig. 6. Role of C-terminal putative helix turn helix motifs containing domain in drDnaA function. The 99 amino acids of drDnaA were removed from its C-terminal and resulting DnaAΔCt derivative was generated (A). An increasing concentration of recombinant protein was incubated with radiolabeled *oriCI* (*oriI*) as described in Fig 4A. Products were analysed on native PAGE and autoradiogram was developed (B). Similarly, 2 μM concentration of DnaAΔCt was incubated for different time points with [^{32}P]-αATP in the absence (C) and presence (D) of *oriCI* (*oriI*). Products were separated on TLC and analyzed as described in Fig 5. The percentage of ADP/ATP ratios were plotted as a function of time (E). Results were analyzed using student t-test and significant difference in data sets with p values of 0.05 or less is shown as (*).

Fig 7

Fig. 7. DNA binding activity of DnaB of *D. radiodurans* (drDnaB). An increasing concentration (μM) of recombinant purified drDnaB (DnaB) was incubated with radiolebeled *oriCI* (*oriI*) taken as dsDNA (A, B) and radiolabeled ssDNA (C and D) in the absence (A and C) and presence (B and D) of ATP. Products were analyzed on non-denaturing PAGE. Autoradiograms were developed, band intensity of free and bound form of DNA was estimated densitometrically from autoradiogram A (E), B (F), C (G) and D (H), and the mean ± SD (n=3) of percent bound fraction was plotted.

Fig. 8. ATPase activity of purified recombinant drDnaB. An increasing concentration of
drDnaB (DnaB) was incubated with $[^{32}P]$-αATP (αATP) in the absence (A) and presence (B)
of ssDNA and generation of $[^{32}P]$-αADP (αADP) product was detected on TLC. Spot
intensity was quantified densitometrically and the percent of ADP/ATP ratios were plotted as
mean ± SD (n=3). Results were analyzed using student t test and significant difference in data
set with p values of 0.05 or less is marked with (*).

1005

1006

1007

1008

1009

Fig. 9. Helicase activity assay of recombinant drDnaB. The purified recombinant drDnaB (DnaB) (2 μM) was incubated with radiolabeled dsDNA having 5' overhang (A) and 3' overhang (B) in the presence and absence of ATP. The role of ATP hydrolysis on helicase activity was monitored with dsDNA with 5' overhang in the presence and absence of ATP and ATP-γ-S for different time points (C). Products were analyzed on denaturing PAGE and autoradiograms were developed. The helicase activity was also monitored using FRET assay on substrate containing FAM at 5' end and BHQ at 3' end of dsDNA (D). Loss of FRET was monitored as gain of FAM emission with the substrate having 5' overhang (E) and 3' overhang (F) as a function of time in the presence and absence of ATP and ATP-γ-S. Data represents the sets of the reproducible experiment repeated 3 times.

1020

1021

1022

1023

Fig 10. Interaction between drDnaA and drDnaB proteins. The translation fusions of drDnaA

and drDnaB with T18/T25 tags at N-terminal (N18/N25-DA and N18/N25-DB) and at C

terminal (DA-C18/C25 and DB-C18/C25) were expressed in *E. coli* BTH 101 cells in

different combination. The expression of β-galactosidase was checked in spot test and in

liquid culture (A). Total proteins from *E. coli* expressing some of these combinations were

immunoprecipitated with T25 antibodies and interacting partner were detected using T18

antibodies (B and C). The pUT18 (T18) and pKNT25 (T25) vector were used as negative

control while *E. coli* FtsA (EcFtsA) and FtsZ (EcFtsZ) were used a positive control. *In vivo*

interaction of drDnaA (DA-C18), DnaA with histidine tag (HisDA), drDnaB (DB-C18),

drDnaB with histidine tag (HisDB) and C-terminal truncated drDnaA with histidine tag (His-

DAΔct) was monitored in *D. radiodurans* R1 (WT) expressing these recombinant proteins in

different combinations on the plasmids. Total proteins were precipitated with polyhis

antibodies and perspective interacting partners were detected using antibodies against T18

domain of CyaA.

1038

1039

1040

1041

Fig. 11. Effect of recombinant PprA on ATPase activity of recombinant drDnaA and drDnaB.

Recombinant PprA was incubated with drDnaA (DnaA) (A, B) and drDnaB (DnaB) (C, D)

separately. ATP hydrolysis was monitored using [^{32}P]-αATP (αATP) at different time

interval and generation of [^{32}P]-αADP (αADP) product was estimated. Percentage of ADP to

ATP ratios were plotted as the mean ± SD (n=3) (B, D). Data sets were analyzed by student t

test and significant difference if any is given as the p values less than 0.01 and 0.001 were

denoted as (**) and (***), respectively and non-significance is marked as (ns).

1049

1050

1051

1052

Fig 12. Effect of purified PprA on helicase activity of drDnaB. Recombinant drDnaB (DnaB)

was incubated with radiolabeled dsDNA having 5' overhangs for different time interval in the

presence of different combinations of PprA and ATP. The products were analyzed on

denaturing PAGE (A). Similarly, FRET substrate of dsDNA with 5' overhang (Fig 9D) was

incubated with drDnaB (DnaB) in different combinations as described in A. Helicase activity

was monitored as the loss of FRET and gain of FAM emission with time (B). Data given are

representative of the reproducible experiments repeated 2 times.

1060

1061

1062

Fig 13. Influence of PprA on drDnaA and drDnaB interaction in surrogate *E.coli*. *E. coli*
BTH 101 cells co-expressing C-terminal fusions of T18/T25 tags with drDnaA (DA-C18,
DA-C25), DnaB (DB-C25, DB-C18) and PprA in different combinations. The expression of
β-galactosidase as an indication of two proteins interaction was monitored in spot test and in
liquid assay (A). The pUT18 and pKNT25 vectors were used as negative control while *E. coli*
FtsA (EcFtsA) and FtsZ (EcFtsZ) were used a positive control. Total proteins of *E. coli*
expressing some of these combinations were immunoprecipitated using T25 antibodies and
interacting partners were detected by T18 antibodies (B, C). In Fig. B and C, the upper panels
are immunoblots while lower panels are corresponding SDS-PAGE gel. PprA influence on
interaction of these proteins is summarised (D).

1073

1074