Age-dependent selection on MHC class 2 variation in a free-living ruminant

Wei Huang[1], Kara L Dicks, Jarrod D Hadfield, Susan E Johnston, Keith T Ballingall, Josephine M Pemberton

Abstract

Genes within the major histocompatibility complex (MHC) are the most variable identified in vertebrates. Pathogen-mediated selection (PMS) is believed to be the main force maintaining diversity at MHC class I and II genes, but it has proven hard to demonstrate the exact PMS regime that is acting in natural populations. Demonstrating contemporary selection on MHC alleles is not trivial, and previous work has been constrained by limited genetic tools, low sample sizes and short time scales and has sometimes involved anticonservative statistical approaches. Here, we use appropriate statistical approaches to examine associations between MHC variation and several fitness measurements including total fitness (lifetime breeding success) and five fitness components, in 3400 wild Soay sheep (*Ovis aries*) monitored over their lifetimes between 1989 and 2012. We found haplotypes C and D were associated with decreased and increased male total fitness respectively. In terms of fitness components, juvenile survival was positively associated with haplotype divergence. Of the eight MHC haplotypes (A-H), haplotypes C and F were associated with decreased adult male breeding success and decreased adult female life span respectively. Consistent with the increased male total fitness, haplotype D, which is the rarest, has increased in frequency throughout the study period. Our results suggest that contemporary balancing selection is acting on MHC class II genes in Soay sheep and that different selection mechanisms are acting between juveniles and adults.

Introduction

The vertebrate immune system has evolved to protect the host from infection through a variety of innate and adaptive immune mechanisms. Many of the proteins that control the specificity of immune function are the products of genes that are amongst the most diverse within vertebrate populations. Such diversity is believed to have evolved in response to pathogen diversity. Investigating the mechanisms shaping the diversity of immune genes has been a major focus in evolutionary biology (1). Genes located within the major histocompatibility complex (MHC) encode proteins that play important roles in the

---

[1] To whom correspondence should be addressed. Email: wei.huang@ed.ac.uk

vertebrate immune system and have been intensively studied across many species,
including model organisms (2-4). MHC molecules, the products of MHC class I and class II
genes, present peptide fragments to antigen-specific receptors on T cells to induce adaptive
immune responses. MHC class I molecules present peptides derived from intracellular
pathogens such as viruses to CD8 +ve T cells, while MHC class II molecules present peptides
derived from extracellular pathogens such as bacteria and parasites to CD4 +ve T cells (5).
40
MHC class I and II genes are the most polymorphic loci identified in vertebrate populations
with hundreds of alleles described in many species (6, 7). Pathogen-mediated balancing
selection is thought to be the principal driver maintaining such high levels of allelic diversity
(8-11). However, the specific selection mechanisms which act to maintain high levels of
MHC allelic diversity are not clear (3, 12, 13). Three models have been proposed to explain
how PMS acts to maintain such diversity:
47
1) *Heterozygote advantage*: individuals which are more heterozygous across the MHC can
respond to a greater variety of pathogens. As a result, variation at MHC loci is selected
and maintained (14). In addition, *divergent allele advantage* has been proposed as an
extension to heterozygote advantage. Here, individuals with a greater functional
divergence between alleles should have a selective advantage because their MHC
molecules can bind a broader range of antigens (15).
54
2) *Rare-allele advantage*: there is strong selection on pathogens to evade immune
protection provided by the most common host MHC alleles. Therefore, rare alleles have
a selective advantage. However, the advantage disappears as the frequency of a
protective MHC allele increases and is absent at equilibrium. The arms race between
hosts and pathogens sets up a cyclical process causing MHC alleles and pathogens to
fluctuate, meaning that MHC diversity is maintained by negative frequency-dependent
selection (16-18). It should be noted that since rare alleles appear most frequently as
heterozygotes, this can manifest as heterozygote advantage for specific alleles under
some circumstances.
64
3) *Fluctuating selection*: a changing environment results in variation in the abundance of
different pathogens in space and time, which generates directional selection in different
host subpopulations and/or at different time points. Thus, different alleles will be
favoured in different subpopulations or at different times, meaning MHC diversity is
maintained as a consequence (19, 20). Unlike rare-allele advantage the change in
selection coefficients in time and/or space is not driven by coevolution in the pathogens.
71
To test each selection model we can examine associations between MHC diversity and
parasite presence or parasite load; different patterns of associations can be explained as a
result of different selection mechanisms (13). A number of studies on MHC genes in natural

populations have examined associations between MHC variation and parasites, often with
conflicting results. Some studies found that MHC heterozygosity or certain MHC alleles
confer resistance to parasite infection (21-23), while other studies found no association
between MHC alleles and parasite load (24). However, such studies may be limited by
incomplete data on the diversity and abundance of the parasite populations present and
parasites that strongly select for MHC diversity may have been missed. As a consequence,
quantifying relationships between MHC genes and pathogens may not have been able to
explicitly elucidate which selection mechanisms were operating, although some
experimental studies have managed to demonstrate a clear mechanism by examining
associations between MHC and pathogens (25, 26). An alternative means to test selection
on MHC genes would be to examine MHC-dependent fitness patterns (4, 27).

As with studies examining associations between MHC and parasites, MHC-fitness
associations can also indicate different selection mechanisms. Under heterozygote
advantage, we predict a positive association between MHC heterozygosity and fitness.
Under rare-allele advantage, if MHC alleles are not in equilibrium, we may detect selection
favouring specific rare alleles and probably also disfavouring specific common alleles. Under
fluctuating selection, we would also observe specific alleles being favoured or disfavoured
by selection. But unlike rare-allele advantage, this should occur regardless of their frequency.
However, it is hard to distinguish rare-allele advantage and fluctuating selection by
examining MHC-fitness associations (13). Because measuring fitness is generally challenging
except in some long-term individual-based studies of natural populations, only a few studies
have tested associations between fitness and MHC variation in the wild, again with mixed
results. First, some MHC studies in wild populations have found a positive association
between MHC heterozygosity and fitness. For instance, the number of MHC alleles was
found to be positively associated with increased apparent survival (return rate from initial
year of capture) in common yellowthroats *(Geothlypis trichas)* (28). Similarly, Banks et al.
(29) found an association between MHC heterozygosity and survival in a natural population
of mountain brushtail possums *(Trichosurus cunninghami)*. Second, some studies have
reported associations between fitness and specific MHC alleles or supertypes (clusters of
MHC alleles with similar physicochemical properties at their antigen-binding sites). For
example, Lukasch et al. (30) found that a specific MHC allele was associated with higher
survival of house sparrow fledglings *(Passer domesticus)*. Another study of great tits *(Parus*
major) using mark-recapture data identified three MHC supertypes associated with fitness
components, one with increased survival, one with increased lifetime reproductive success
and another one with decreased lifetime reproductive success (31). Kloch et al. (32) also
found that two MHC alleles from different phylogenetic clusters were associated with
winter survival in root voles *(Microtus oeconomus)*. Finally, there are some studies where no
associations are detected. In collared flycatchers (Ficedula albicollis), no association
between MHC variation and lifetime reproductive success was found (24). Similarly in house

sparrows (*Passer domesticus*), no significant associations between MHC variation and
survival were identified (33).
A complication in studies of fitness association is that associations may differ between
individuals of different sex, age or nutritional status, presumably due to variation in their
ability to counter the threat from pathogens. For example, in a population of black-legged
kittiwake (*Rissa tridactyla*), the association between MHC diversity and fitness was only
found in second-hatched female chicks suggesting sex and hatching order could modulate
MHC-fitness association (34). In a previous study of Seychelles warbler (*Acrocephalus*
sechellensis), a positive association with MHC diversity was only found in juvenile survival
but not in adult survival (35). However, the impact of age on MHC-fitness associations has
been rarely studied because of the limitation of study length and scale in wild populations.
Thus, longitudinal studies with fitness measurement through all life stages of study
individuals are required to study MHC-dependent fitness effects.
128
Apart from the difficulty in measuring fitness to study selection on MHC genes, many
previous studies of selection on MHC genes are limited by the depth and quality of the
genetic data available. Within a typical mammalian MHC region, multiple closely linked and
duplicated genes are present which are inherited together as a haplotype. With the
development of next generation sequencing (NGS) technologies, we are now able to cost-
effectively genotype large numbers of samples (36, 37). However, in species where the MHC
region is poorly characterised such as in fish or passerine birds, it is difficult to assign a
sequence to an individual locus or identify if a locus is functional. Without locus-specific
genotyping, multiple loci may co-amplify, which will impact subsequent testing for
associations between MHC diversity and fitness. For example, heterozygote advantage
cannot be demonstrated without knowing that alleles are alternatives at the same locus.
Even if locus-specific genotyping is possible, many studies have focused on a single MHC
locus and ignored linkage with other MHC loci (29, 32, 38). Locus-specific genotyping and
characterisation of haplotypes are desirable when testing for associations between fitness
and MHC diversity in wild populations.
144
Lastly, statistical analyses of selection on MHC variation could be more precise for
controlling confounders than is often deployed. Measures of parasite load and fitness are
quantitative traits likely to be determined by both environmental factors and multiple
genetic factors, including MHC variation. As such, it is possible that studies of contemporary
selection may confound MHC similarity with genetic relatedness, so an MHC effect could be
due to variation elsewhere in the genome rather than the MHC itself. Selection on MHC
genes in natural populations has mostly been assessed in long-term individual-based studies
which include a lot of genetically related individuals (31, 35). To our knowledge, few MHC
studies in natural populations have been conducted in a quantitative genetic framework. In
addition, given the ubiquity of inbreeding depression, any analysis of selection should
include an estimate of individual inbreeding to ensure MHC effects are not confounded with

genome-wide inbreeding effects. However, only a few previous studies, e.g. (39, 40) have accounted for inbreeding.

Here, we present the first investigation of contemporary selection acting on the MHC in a free-living population that overcomes the issues raised above. The Soay sheep *(Ovis aries)* population on the island of Hirta, St Kilda is one of the most intensively studied wild animal populations in the world (41). Since 1985, individual Soay sheep living in the Village Bay study area have been followed from birth, through all breeding attempts, until death. In addition, using a combination of observation and SNP genotypes, a multigenerational pedigree for the population has been assembled which includes annual and lifetime measures of fitness for both sexes (42). Previous work characterised diversity at the MHC class II *DRB1* gene using an internal microsatellite and identified associations between MHC genotype, fitness and parasite infection (43, 44). However, the only fitness measure studied was juvenile survival. Another study using the same genetic dataset of MHC-linked microsatellite alleles found higher temporal and spatial variation in these loci than in putatively neutral microsatellite loci in other parts of the genome, suggesting fluctuating selection on MHC genes (45). Recently, using sequencing-based genotyping of functional MHC class II DR and DQ loci we have shown that eight MHC haplotypes are segregating in the study population (46). Using a panel of 13 SNPs, MHC class II diplotypes (i.e. haplotype complement) were characterised for 5349 individuals sampled between 1989 and 2012 (47). Diplotypes were in Hardy-Weinberg equilibrium both within cohorts (cite Kara BioRxiv MS) and within the population alive each year (47). Here, we use this data to investigate associations between MHC variation and fitness. We test for associations between (1) MHC variation and total fitness, defined as lifetime breeding success of each individual and (2) MHC variation and five fitness components. We fit MHC variation including divergence, heterozygosity and individual haplotypes within quantitative genetic models in order to distinguish the effect of MHC haplotypes from genome-wide effects. Finally (3), we perform gene-drop simulations to examine whether haplotype frequencies have changed more than expected from a neutral process.

Methods

Study system

Soay sheep have lived virtually unmanaged on the island of Soay, in the St. Kilda archipelago, for thousands of years. In 1932, 107 Soay sheep were introduced to the larger neighbouring island of Hirta and have been living there unmanaged since. A previous study demonstrated that an introgression event between Soay sheep and a more modern breed occurred approximately 150 years ago (48). From 1985, a long-term study has been conducted on the sheep resident in the Village Bay area of Hirta to investigate ecological and evolutionary questions (41). The vast majority of lambs are born in April or May of each year. At their first

live capture, lambs are ear-tagged to enable life-long identification and discs of ear tissue
removed in the tagging process are used for DNA extraction. Parentage is inferred for each
individual using a subset of 315 unlinked SNPs derived from the Illumina Ovine 50K SNP
array, on which most individuals alive since 1990 have been genotyped (49). In addition, the
R package Sequoia (50) was used to cluster half-siblings which share a parent that has not
been genotyped. In cases where no SNP genotypes were available, a small number of
parentage inferences were made using field observations (for mothers) or a previous
microsatellite genotype approach (51). The pedigree used here includes 7447 individuals,
with 7014 paternal links and 6229 maternal links. This study investigated associations
between MHC variation and fitness measurements defined as follows:

• Total fitness, the number of offspring an individual produced over its lifetime;
• Juvenile survival (JS), a binary variable indicating whether a lamb survived from birth
until May 1st of the following year.
• Adult annual survival (adult AS), a binary variable indicating whether an adult
survived or not in a given calendar year; an adult is an animal that survived to age 1.
• Adult life span (adult LS), the age of an adult when it died;
• Adult annual breeding success (adult ABS), the number of offspring an adult
produced in a given calendar year;
• Adult lifetime breeding success (adult LBS), the number of offspring an adult
produced over its lifetime;

MHC Class IIa Haplotyping

The genetic data used in this study were obtained from a previous study (46, 47). In that
study the seven expressed loci (*DRB1*, *DQA1*, *DQA2*, *DQA2-like*, *DQB1*, *DQB2* and *DQB2-like*)
within the MHC class IIa region were characterised in 118 selected Soay sheep using
sequencing-based genotyping. A total of eight MHC haplotypes were identified and named
A to H. These haplotypes were confirmed in an additional 94 individuals that were unrelated
based on the pedigree. A panel of 13 SNPs located in the region of MHC class IIa haplotypes
was then selected to impute the eight MHC class IIa haplotypes in 5951 Soay sheep
genotyped using Kompetitive Allele-specific PCR (KASP). This panel included 11 SNPs known
to be variable in Soays based on data from the Illumina Ovine infinium HD chip and two
other SNPs identified within the *DQA1* gene (52). After quality control, the diplotypes of
5349 individuals that lived in the study area between 1985 and 2012 were identified. As
cohort effects were fitted in the statistical analyses, we excluded animals born before 1989
since too few were genotyped to be representative of their birth cohort. We also removed
the cohort born in 2010 which had a high genotyping failure rate in the KASP genotyping
assay, all animals which died as foetuses and any individuals subjected to experimental
treatments which may have affected their survival or breeding performance (e.g.
anthelmintic boluses). After these exclusions, a total of 3440 individuals (1750 males and

1650 females) remained for statistical modelling of total fitness and JS (Table 1). The
frequency of the rarest haplotype (D) was 3.02%. With the exception of animals
homozygous for haplotype D (N=5), moderate numbers of individuals (N>20) of each
diplotype are represented (Supplementary 1). For each individual successfully diplotyped,
MHC divergence was measured by the proportion of the amino acid sequence that differed
between the two MHC haplotypes (p-distance)(53) (Supplementary 2).
241

242 Statistical analysis

We used generalized linear mixed models (GLMMs) to study the associations between MHC
haplotypes and fitness measurements. The primary models were animal models (AMs) that
include a covariance structure proportional to the pedigree relatedness matrix, but we also
repeated all tests without pedigree information. In null models, we fitted a set of fixed and
random effects for each fitness measurement (Table 1) including genomic inbreeding (F_{grm})
as a fixed effect, as previous studies demonstrated inbreeding depression in this population
(54, 55). None of the fitness measurements in our study had a normal distribution, so we
used Markov chain Monte Carlo techniques implemented in MCMCglmm v 2.28 (56) to run
models.
252

To examine whether there were associations between MHC variation and fitness
measurements, we fitted MHC heterozygosity, MHC divergence and individual MHC
haplotypes simultaneously into the null model. Each haplotype was fitted as dosage (0, 1 or
2 copies) as a fixed effect in all the models. We performed a Wald test using the posterior
mean and the posterior covariance matrix to test whether MHC haplotypes explained
variation in fitness. In these models, Haplotype H was treated as a reference so the results
for individual haplotypes from our models are relative to haplotype H. When a Wald test for
haplotype differences was significant, we conducted an additional analysis (Supplementary
4) comparing the estimate for each haplotype with the mean of the estimates for all other
haplotypes, to identify individual haplotype effects.
263

Finally, we used models with different error distributions to combat the heterogeneity of
ontogenies and distributions of fitness measurements (Supplementary 3). We used probit
regression to run a unisex model for JS. As the ontogenies and distributions of total fitness
and adult fitness components were different between the two sexes, we used separate
models for males and females. For both males and females, we used zero-inflated Poisson
models for total fitness, probit regression for adult AS and Poisson regression for adult LS
and LBS. As females can only have up to three offspring each year while males had between
0 and 24, we used multiple category probit (ordinal) regression for female ABS and Poisson
regression for male ABS. All the models were run for 200,000 iterations in R v3.5.2 (57).
273

274 Table 1. Fixed and random effects fitted in null model and sample sizes, for each fitness
275 measurement. Sample sizes are shown by number of individuals (records).
276

Fitness measurement	Fixed effects			Random effects			Sample sizes	
	Litter Size	Age (years, quadratic)	Inbreeding ($F_{grm)}$	Birth Year	Year of Fitness Measure	ID	Female	Male
Total fitness			×	×			1650	1750
JS	×		×	×			1650	1750
Adult AS		×	×	×	×	×	551 (3093)	529 (1610)
Adult LS			×	×			574	545
Adult ABS		×	×	×	×	×	574 (3667)	545 (2155)
Adult LBS			×	×			574	545

Gene-drop analysis

We performed gene-drop simulations with a custom script in R v3.5.2 (57) to model the
expected frequency change of MHC haplotypes under genetic drift given the pedigree
structure. Hereafter, the first four birth cohorts (1989 to 1992) are defined as the "founder"
cohorts, whereas all subsequent cohorts are defined as "simulated" cohorts (1993 to 2012).
A total of 5000 simulations were conducted as follows. Individuals in the founder cohorts
were assigned their observed diplotypes; founder cohort individuals with unknown
diplotypes were assigned one by either: (a) sampling a haplotype from each parent
assuming Mendelian segregation; or (b) sampling a haplotype with the probability of the
observed haplotype frequencies in the same birth cohort when one or more parents were
unknown. In the simulated cohorts, each individual diplotype was sampled using
approaches (a) and (b) above. By this method, simulated diplotypes were generated for
each individual in the pedigree born after 1992. Using these data with the record of birth
year and death year of each individual, we compared the observed and simulated changes
in MHC haplotype frequency for the standing population (all the individuals living in a single
year), using a linear regression with year (1993 to 2012) as the predictor variable. In cases
where the observed slope fell within the top or bottom 2.5% of simulated slopes, the allele
frequency was deemed to have been subject to directional selection.
296

Results

Associations between MHC and fitness measurement

The results of models with and without pedigree information were largely similar (Supplementary 5 to 7). Here we report the results of animal models, i.e. including the pedigree as a random effect.

Figure 1. Associations between MHC and fitness measurements in Soay sheep derived from animal models. (A) Associations between MHC divergence and fitness measurements in juveniles (circle), females (squares) and males (triangles). Bars represent the 95% credibility intervals and those not overlapped with zero represent significant results. (B) Associations between MHC haplotypes and total fitness in females and males. The solid line represents the model intercept at haplotype H and posterior means and credible intervals for each haplotype are plotted relative to H. Dashed lines indicate the average posterior mean of all eight haplotypes. Circles indicate the Wald test was not significant, other symbols indicate it was significant. Haplotype effects that were significantly different from the mean are shown as squares with non-significant ones show as triangles. (C and D) Associations between MHC haplotypes and adult fitness components in females (C) and males (D). Symbols and lines as for B.

317 Table 2. Results (p value) of Wald tests for animal models testing for differences between
318 haplotypes in fitness measures (d.f.=7). Bold numbers show significant results (p <0.05).

Sex	Fitness measurement	p value
/	JS	0.13
Female	Total fitness	0.45
	AS	0.073
	LS	**0.028**
	ABS	0.92
	LBS	0.12
Male	**Total fitness**	**0.0034**
	AS	0.39
	LS	0.55
	ABS	**0.019**
	LBS	**0.039**

Total fitness

321 There was no association between MHC heterozygosity or MHC divergence and total fitness
322 in either sex (Figure 1A). In the Wald test for haplotype effects a significant association was
323 found for males, but not for females (Table 2). Haplotype C was associated with decreased
324 male total fitness while haplotype D was associated with increased total fitness (Figure 1B).

Juvenile survival

327 We found a positive association between MHC divergence and JS (Figure 1A). Considering
328 haplotype effects, the Wald test was not significant (Table 2).

Adult annual survival

331 There was no association between MHC heterozygosity or MHC divergence and adult annual
332 survival (AS) in either sex (Figure 1A). The Wald test for haplotype effects in AS was not
333 significant for either sex (Table 2).

Adult life span

336 There was no association between MHC heterozygosity or MHC divergence and adult life
337 span (LS) in either sex (Figure 1A). In the Wald test for haplotype effects a significant
338 association was found for females, but not for males (Table 2). Haplotype F was associated
339 with decreased adult female life span (Figure 1C).

Annual breeding success

342 There was no association between MHC heterozygosity or MHC divergence and annual
343 breeding success (ABS) in either sex (Figure 1A). In the Wald test for haplotype effects a

344 significant association was found for males, but not for females (Table 2). Haplotype C was
345 associated with decreased adult male annual breeding success (Figure 1D).

Lifetime breeding success

348 We found no association between MHC heterozygosity or MHC divergence and adult
349 lifetime breeding success (LBS) in either sex (Figure 1A). In the Wald test for haplotype
350 effects a significant association was found for males, but not for females (Table 2).
351 Haplotype C were associated with decreased adult male lifetime breeding success (Figure
352 1D).

Models without fitting the pedigree

The results obtained from models not including the pedigree were generally consistent with
those above for associations between MHC heterozygosity/divergence and fitness
measurements (Supplementary 5). For associations between individual MHC haplotypes and
fitness measurements, estimates were mostly similar, but there were minor variations in
terms of significance of individual haplotypes (Table 1, Supplementary 5). One difference
was that we that found haplotype D was associated with increased female adult LS in
models without pedigree fitted (Supplementary 5).

Gene-drop analysis

All haplotype frequency changes were within expectation due to drift alone except for
haplotype D (Figure 2). Of the 5000 simulations for Haplotype D, only 97 simulated
regression slopes were greater than the slope generated from real data (p=0.0194).
Therefore, this haplotype increased in frequency more than predicted by chance.

Figure 2. Results of the gene drop analysis for each haplotype. The black lines represent the

result of simulations (N=5000) and the red lines represent the observed frequency of each

haplotype in the Soay sheep population over the same time interval.

Discussion

In this study we investigated associations between MHC class II diversity (heterozygosity,

divergence and specific haplotypes) and fitness measurements including total fitness and

five fitness components in a sample of individually-monitored free-living ruminants using

generalized linear mixed models. We fitted MHC divergence, MHC heterozygosity and

individual MHC haplotype effects together to test whether there is divergent

allele/heterozygote advantage, rare-allele advantage or fluctuation selection. For all models,

we included genetic relatedness to eliminate potential false positive results due to other

genomic regions in common in related individuals and a genome inbreeding estimator to

control genome-wide heterozygosity and the effect of inbreeding depression. We found a

negative association between haplotype C and male total fitness and a positive association

between haplotype D and male total fitness. We found MHC divergence was associated with

increased JS. We also identified a negative association between haplotype C and adult male

ABS and LBS and a negative association between haplotype F and adult female LS. Finally,

we found that the rarest haplotype, D, increased significantly in frequency over the 23 year

study period.

In relation to the various hypotheses for the evolution of MHC diversity, our study supports

multiple hypotheses. In wild populations, evidence of MHC heterozygote fitness advantage

has been reported in some previous studies (28, 29, 35) but not in others (24, 58). In
addition, some studies have found evidence of divergent allele advantage, over and above
heterozygote advantage, for example (40). In this study we found evidence for the divergent
allele advantage hypothesis as MHC divergence was positively associated with JS
independent of MHC heterozygosity. In addition, selection favouring or disfavouring
particular haplotypes, as is likely under negative frequency-dependent selection when not in
equilibrium or by the fluctuating selection hypothesis, has been reported in previous studies
(31, 44). In this study, we also found evidence for negative frequency-dependent selection
or fluctuating selection in that specific MHC haplotypes were associated with fitness
measurements including total fitness, adult female LS and adult male ABS and LBS. Although
haplotype D is rare, favoured and increasing, suggestive of rare allele advantage, our results
are consistent with both rare allele advantage and fluctuating selection andwe are not
capable of differentiating them at this time (13).
407
In this study, we found that MHC-fitness associations varied among different fitness
measurements. We found evidence of divergent allele advantage in juvenile survival, but did
not observe any evidence of these effects in total fitness and adult fitness components.
These results are consistent with a previous study of Seychelles warbler that found a
positive association between MHC diversity and juvenile but not adult survival (35). In Soay
sheep, parasite infection intensity (measured as strongyle egg count) (59) and mortality is
highest in the juvenile period (60) and parasite infection intensity is also negatively
correlated with juvenile survival (61). Therefore, lambs with divergent MHC haplotypes may
be better able to respond to a wider range of pathogen antigens, increasing their chances of
surviving the harsh juvenile period. However, as more MHC homozygote lambs die in their
first year, selection for divergent MHC constitution may no longer be evident in adults due
to selective disappearance. Alternatively, the lack of selection for MHC heterozygosity or
divergence in adult sheep could also result from maturation of the immune system or a
change in the parasite community (59, 62). In terms of specific haplotypes, we did not find
any association between specific MHC haplotypes and juvenile survival, but we did detect
associations with total fitness and adult fitness components which could be a reflection of
the heterogeneity of parasite infection and the development of immunity to parasites
acquired between the juvenile and adult periods.
426
Although many studies in wild populations have identified associations between specific
MHC alleles and fitness, few studies have observed an allele frequency change in response
to selection. One exception is a study of the great reed warbler (*Acrocephalus arundinaceus*)
that identified significantly higher variation in MHC allele frequencies between cohorts than
at neutral loci, suggesting that fluctuating selection may be acting on MHC variation (63). In
a previous experimental study, MHC alleles providing resistance to the respective specific
parasite increased in frequency in the next host generation (64). If MHC-linked fitness
effects are strong enough, we should be able to see changes in haplotype frequency over

time in Soay sheep. In our study, we found haplotype C and D were associated with
decreased and increased male total fitness respectively. In addition, the point estimates of
haplotype C and D for female total fitness are, if anything, in the same direction (though not
significant). Consistent with this, the gene drop analysis showed that haplotype D has
increased significantly through the study period (p=0.0194). However, the frequency of
haplotype C did not change more than expected by chance. A possible reason for this
difference lies in the effects on adult life span (see Figure 1B, supplementary figure s5.2).
Haplotype C was not significantly associated with adult life span in either sex, whereas
haplotype D was associated with adult female life span and the point estimates of haplotype
D for male adult life span was in the same direction though not significant. If individuals
carrying haplotype D remain in the population longer, as this suggests, then they contribute
to the significant increase in haplotype frequency in standing population each year. To our
knowledge, our study is the first demonstrating that the frequency of a MHC haplotype
conferring selective advantage also increased through the study period in a wild population.
449
Several aspects of our analysis improve on earlier MHC-fitness association studies.
Benefiting from MHC haplotype data generated by locus-specific genotyping, this is one of
the few studies using dosage models to precisely examine whether there is evidence for
heterozygote advantage and divergent allele advantage which are caused by non-additive
genetic effects (65, 66). Indeed, we demonstrated signature of divergent haplotype
advantage in JS. In addition, to our knowledge, this is the first study to use animal models in
a study of association between MHC and fitness measurements in wild populations. As the
AM accounts for relatedness, we predicted that it would be more conservative and show
fewer associations than GLMMs that did not account for relatedness. In fact, we found the
results to be largely consistent between the two models. This may be due to the generally
low heritability of the fitness traits (Supplementary 9). However, for traits which have a
higher heritability, the results of animal models will be more conservative as they would
exclude the false positive effects caused by genomic regions other than the MHC.
463
Long term individual-based studies like the Soay sheep project also enable repeat
measurements of the strength and direction of selection over time (67, 68). In a previous
study, three MHC-linked microsatellite alleles were found to be associated with survival in
Soay sheep using individuals from 1985 to 1994 (44). Microsatellite alleles OLADRB 257 and
205 were associated with decreased juvenile and yearling survival respectively whereas the
OLADRB 263 allele was associated with increased yearling survival. Some microsatellite
alleles can be linked directly to the MHC class II haplotypes studied here: OLADRB allele 257
is linked to haplotype B (Supplementary 10). Thus, we might expect some consistency
between the results from the previous study and the current one. However, in our analysis
of juvenile survival, we did not find any evidence for haplotype-specific effects (Wald tests
not significant; Table 2). The inconsistency between the previous and current study is
probably due to differences in genotyping method, sample size and statistical methods

including fitting inbreeding. In addition, alleles OLADRB 205 and 263 are associated with
multiple haplotypes, and we have studied annual survival rather than yearling survival,
which may also contribute to a failure to repeat the results.

Here we have identified a number of associations between MHC diversity and fitness
measurements which are likely due to PMS. We would therefore expect to observe similar
associations between MHC class II variation and measures of immunity to key parasites. In a
previous study using the same MHC haplotype dataset, we examined associations between
MHC class II variation and five phenotypic traits including strongyle faecal egg count, FEC
and titres of *Teladorsagia circumcincta* specific immunoglobulin isotypes IgA, IgE and IgG.
Although FEC is negatively correlated with fitness, no significant association between MHC
variation and FEC was identified. In contrast, MHC divergence was found to be positively
associated with lamb IgA titres. Specific haplotypes were also associated with antibody titres
including a positive association between haplotype C and adult IgE and a positive
association between haplotype D and IgE. However, no association was identifiedwith
haplotype F that might match the female adult LS result found here. Our failure to identify
MHC-FEC associations may result from the nature of this measure, its relationship with
parasite burden and immunity to gastrointestinal nematode parasites. Other
microorganisms, pathogenic or benign, are likely be involved in selection on MHC genes,
including gut microbiota (69). Further studies examining associations between MHC
diversity and the range of pathogens each animal is exposed to will help to clarify the nature
of the selective forces acting to maintain MHC diversity in Soay sheep.

In summary, we used well-characterized MHC class II haplotypes to investigate MHC-linked
fitness effects in Soay sheep. To our knowledge, our dataset, with more than 3000
individuals, is the largest yet used to study selection on MHC variation in a wild population.
Our results support the existence of contemporary selection on MHC class II variation in
Soay sheep, when genetic relatedness and inbreeding are controlled for. However, the
selection mechanisms differs between juveniles and adults. We observed that the frequency
of a rare MHC haplotype with a selective advantage has increased significantly during the
study period. Our study highlights the importance of investigating selection on MHC genes
across the whole lifespan and suggests different selection mechanisms on MHC variation
could act simultaneously in a natural population.

Acknowledgments

We thank the National Trust for Scotland for permission to work on St. Kilda and QinetiQ for
logistics and support. We thank J. Pilkington, I. Stevenson and many volunteers who have
helped with fieldwork on the island and all those who have contributed to keeping the
project going over many years. W. Huang was supported by Edinburgh Global research
scholarship. K. Ballingall received funding from the European Union's Horizon 2020 research

and innovation programme under grant agreement no. 731014KB (VetBioNet) and
acknowledges the support received from the Scottish Government's strategic research
programme. The long-term project on St Kilda, the KASP genotyping and the SNP array
genotyping were funded by the UK Natural Environment Research Council and the European
Research Council.

References

1.   Obbard DJ, Welch JJ, Kim KW, & Jiggins FM (2009) Quantifying Adaptive Evolution in the
Drosophila Immune System. *Plos Genet* 5(10).
2.   Acevedo-Whitehouse K & Cunningham AA (2006) Is MHC enough for understanding wildlife
immunogenetics? *Trends Ecol Evol* 21(8):433-438.
3.   Bernatchez L & Landry C (2003) MHC studies in nonmodel vertebrates: what have we
learned about natural selection in 15 years? *Journal of Evolutionary Biology* 16(3):363-377.
4.   Piertney SB & Oliver MK (2006) The evolutionary ecology of the major histocompatibility
complex. *Heredity* 96(1):7-21.
5.   Garrigan D & Hedrick PW (2003) Detecting adaptive molecular polymorphism, lessons from
the MHC. *Am J Hum Genet* 73(5):375-375.
6.   Apanius V, Penn D, Slev PR, Ruff LR, & Potts WK (1997) The nature of selection on the major
histocompatibility complex. *Crit Rev Immunol* 17(2):179-224.
7.   Maccari G*, et al.* (2017) IPD-MHC 2.0: an improved inter-species database for the study of
the major histocompatibility complex. *Nucleic Acids Res* 45(D1):D860-D864.
8.   Oppelt C, Starkloff A, Rausch P, Von Holst D, & Rodel HG (2010) Major histocompatibility
complex variation and age-specific endoparasite load in subadult European rabbits.
Molecular ecology 19(19):4155-4167.
9.   Radwan J, Kuduk K, Levy E, LeBas N, & Babik W (2014) Parasite load and MHC diversity in
undisturbed and agriculturally modified habitats of the ornate dragon lizard. *Molecular*
ecology 23(24):5966-5978.
10. Oliver MK, Telfer S, & Piertney SB (2009) Major histocompatibility complex (MHC)
heterozygote superiority to natural multi-parasite infections in the water vole (Arvicola
terrestris). *Proceedings. Biological sciences* 276(1659):1119-1128.
11. Westerdahl H, Asghar M, Hasselquist D, & Bensch S (2012) Quantitative disease resistance:
to better understand parasite-mediated selection on major histocompatibility complex.
Proceedings. Biological sciences 279(1728):577-584.
12. Jordan WC & Bruford MW (1998) New perspectives on mate choice and the MHC. *Heredity*
81:239-245.
13. Spurgin LG & Richardson DS (2010) How pathogens drive genetic diversity: MHC,
mechanisms and misunderstandings. *P R Soc B* 277(1684):979-988.
14. Takahata N & Nei M (1990) Allelic Genealogy under Overdominant and Frequency-
Dependent Selection and Polymorphism of Major Histocompatibility Complex Loci. *Genetics*
124(4):967-978.
15. Wakeland EK*, et al.* (1990) Ancestral polymorphisms of MHC class II genes: divergent allele
advantage. *Immunol Res* 9(2):115-122.
16. Hughes AL & Nei M (1988) Pattern of Nucleotide Substitution at Major Histocompatibility
Complex Class-I Loci Reveals Overdominant Selection. *Nature* 335(6186):167-170.
17. Ejsmond MJ & Radwan J (2015) Red Queen Processes Drive Positive Selection on Major
Histocompatibility Complex (MHC) Genes. *Plos Comput Biol* 11(11).
18. Borghans JAM, Beltman JB, & De Boer RJ (2004) MHC polymorphism under host-pathogen
coevolution. *Immunogenetics* 55(11):732-739.

19. Hedrick PW (2002) Pathogen resistance and genetic variation at MHC loci. *Evolution;*
international journal of organic evolution 56(10):1902-1908.
20. Hill AVS, *et al.* (1991) Common West African Hla Antigens Are Associated with Protection
from Severe Malaria. *Nature* 352(6336):595-600.
21. Rivero-de Aguilar J, *et al.* (2016) MHC-I provides both quantitative resistance and
susceptibility to blood parasites in blue tits in the wild. *J Avian Biol* 47(5):669-677.
22. Westerdahl H, Stjernman M, Raberg L, Lannefors M, & Nilsson JA (2013) MHC-I Affects
Infection Intensity but Not Infection Status with a Frequent Avian Malaria Parasite in Blue
Tits. *Plos One* 8(8).
23. Westerdahl H, *et al.* (2005) Associations between malaria and MHC genes in a migratory
songbird. *P R Soc B* 272(1571):1511-1518.
24. Radwan J, *et al.* (2012) MHC diversity, malaria and lifetime reproductive success in collared
flycatchers. *Molecular ecology* 21(10):2469-2479.
25. Bolnick DI & Stutz WE (2017) Frequency dependence limits divergent evolution by favouring
rare immigrants over residents. *Nature* 546(7657):285-+.
26. Phillips KP, *et al.* (2018) Immunogenetic novelty confers a selective advantage in host-
pathogen coevolution. *Proceedings of the National Academy of Sciences of the United States*
of America 115(7):1552-1557.
27. Sutton JT, Nakagawa S, Robertson BC, & Jamieson IG (2011) Disentangling the roles of
natural selection and genetic drift in shaping variation at MHC immunity genes. *Mol Ecol*
20(21):4408-4420.
28. Dunn PO, Bollmer JL, Freeman-Gallant CR, & Whittingham LA (2013) Mhc Variation Is
Related to a Sexually Selected Ornament, Survival, and Parasite Resistance in Common
Yellowthroats. *Evolution* 67(3):679-687.
29. Banks SC, Dubach J, Viggers KL, & Lindenmayer DB (2010) Adult survival and microsatellite
diversity in possums: effects of major histocompatibility complex-linked microsatellite
diversity but not multilocus inbreeding estimators. *Oecologia* 162(2):359-370.
30. Lukasch B, *et al.* (2017) Major histocompatibility complex genes partly explain early survival
in house sparrows. *Sci Rep-Uk* 7.
31. Sepil I, Lachish S, & Sheldon BC (2013) Mhc-linked survival and lifetime reproductive success
in a wild population of great tits. *Molecular ecology* 22(2):384-396.
32. Kloch A, Baran K, Buczek M, Konarzewski M, & Radwan J (2013) MHC influences infection
with parasites and winter survival in the root vole Microtus oeconomus. *Evol Ecol* 27(3):635-
653.
33. Karlsson M, *et al.* (2015) House sparrowPasser domesticussurvival is not associated with
MHC-I diversity, but possibly with specific MHC-I alleles. *Journal of Avian Biology* 46(2):167-
174.
34. Pineaux M, *et al.* (2020) Sex and hatching order modulate the association between MHC-II
diversity and fitness in early-life stages of a wild seabird. *Mol Ecol* 29(17):3316-3329.
35. Brouwer L, *et al.* (2010) MHC-dependent survival in a wild population: evidence for hidden
genetic benefits gained through extra-pair fertilizations. *Molecular ecology* 19(16):3444-
3455.
36. Babik W (2010) Methods for MHC genotyping in non-model vertebrates. *Mol Ecol Resour*
10(2):237-251.
37. Babik W, Taberlet P, Ejsmond MJ, & Radwan J (2009) New generation sequencers as a tool
for genotyping of highly polymorphic multilocus MHC system. *Mol Ecol Resour* 9(3):713-719.
38. Osborne AJ, *et al.* (2015) Heterozygote advantage at MHC DRB may influence response to
infectious disease epizootics. *Molecular ecology* 24(7):1419-1432.
39. Thoss M, Ilmonen P, Musolf K, & Penn DJ (2011) Major histocompatibility complex
heterozygosity enhances reproductive success. *Molecular ecology* 20(7):1546-1557.

40. Lenz TL, Mueller B, Trillmich F, & Wolf JBW (2013) Divergent allele advantage at MHC-DRB
through direct and maternal genotypic effects and its consequences for allele pool
composition and mating. *P R Soc B* 280(1762).
41. Clutton-Brock T & Pemberton JM (2004) Soay sheep - Dynamics and selection in an island
population. *Cambridge University Press*.
42. Berenos C*, et al.* (2015) Heterogeneity of genetic architecture of body size traits in a free-
living population. *Molecular ecology* 24(8):1810-1830.
43. Paterson S (1998) Evidence for balancing selection at the major histocompatibility complex
in a free-living ruminant. *J Hered* 89(4):289-294.
44. Paterson S, Wilson K, & Pemberton JM (1998) Major histocompatibility complex variation
associated with juvenile survival and parasite resistance in a large unmanaged ungulate
population (Ovis aries L.). *Proceedings of the National Academy of Sciences of the United
States of America* 95(7):3714-3719.
45. Charbonnel N & Pemberton J (2005) A long-term genetic survey of an ungulate population
reveals balancing selection acting on MHC through spatial and temporal fluctuations in
selection. *Heredity* 95(5):377-388.
46. Dicks KL, Pemberton JM, & Ballingall KT (2019) Characterisation of major histocompatibility
complex class IIa haplotypes in an island sheep population. *Immunogenetics* 71(5-6):383-393.
47. Dicks K (2017) Unravelling major histocompatibility complex diversity in the Soay Sheep of
St.Kilda. *University of Edinburgh*.
48. Feulner PGD*, et al.* (2013) Introgression and the fate of domesticated genes in a wild
mammal population. *Molecular ecology* 22(16):4210-4221.
49. Berenos C, Ellis PA, Pilkington JG, & Pemberton JM (2014) Estimating quantitative genetic
parameters in wild populations: a comparison of pedigree and genomic approaches. *Mol
Ecol* 23(14):3434-3451.
50. Huisman J (2017) Pedigree reconstruction from SNP data: parentage assignment, sibship
clustering and beyond. *Molecular Ecology Resources* 17(5):1009-1024.
51. Morrissey MB*, et al.* (2012) The prediction of adaptive evolution: empirical application of the
secondary theorem of selection and comparison to the breeder's equation. *Evolution;
international journal of organic evolution* 66(8):2399-2410.
52. Dicks KL, Pemberton JM, Ballingall KT, & Johnston SE (2020) Haplotyping MHC class IIa by
high throughput screening in an isolated sheep population. *bioRxiv*:2020.2007.2020.212225.
53. Henikoff S (1996) Scores for sequence searches and alignments. *Curr Opin Struc Biol*
6(3):353-360.
54. Berenos C, Ellis PA, Pilkington JG, & Pemberton JM (2016) Genomic analysis reveals
depression due to both individual and maternal inbreeding in a free-living mammal
population. *Mol Ecol* 25(13):3152-3168.
55. Stoffel MA, Johnston SE, Pilkington JG, & Pemberton JM (2020) Genetic architecture and
lifetime dynamics of inbreeding depression in a wild mammal.
bioRxiv:2020.2005.2027.118877.
56. Hadfield JD (2010) MCMC Methods for Multi-Response Generalized Linear Mixed Models:
The MCMCglmm R Package. *J Stat Softw* 33(2):1-22.
57. R Core Team (2013) R: A language and environment for statistical computing. .
58. Sepil I, Lachish S, Hinks AE, & Sheldon BC (2013) Mhc supertypes confer both qualitative and
quantitative resistance to avian malaria infections in a wild bird population. *P R Soc B*
280(1759).
59. Craig BH, Pilkington JG, & Pemberton JM (2006) Gastrointestinal nematode species burdens
and host mortality in a feral sheep population. *Parasitology* 133:485-496.
60. Clutton-Brock TH & Pemberton JM (2004) *Soay sheep: dynamics and selection in an island
population* (Cambridge University Press).

61. Hayward AD, *et al.* (2011) Natural selection on a measure of parasite resistance varies across
ages and environmental conditions in a wild mammal. *J Evolution Biol* 24(8):1664-1676.
62. Sparks AM, *et al.* (2018) Natural Selection on Antihelminth Antibodies in a Wild Mammal
Population. *Am Nat* 192(6):745-760.
63. Westerdahl H, Hansson B, Bensch S, & Hasselquist D (2004) Between-year variation of MHC
allele frequencies in great reed warblers: selection or drift? *J Evol Biol* 17(3):485-492.
64. Eizaguirre C, Lenz TL, Kalbe M, & Milinski M (2012) Rapid and adaptive evolution of MHC
genes under parasite selection in experimental vertebrate populations. *Nature*
communications 3:621.
65. Hu XL, *et al.* (2015) Additive and interaction effects at three amino acid positions in HLA-DQ
and HLA-DR molecules drive type 1 diabetes risk. *Nature genetics* 47(8):898-+.
66. Lenz TL, *et al.* (2015) Widespread non-additive and interaction effects within HLA loci
modulate the risk of autoimmune diseases. *Nature genetics* 47(9):1085-+.
67. Siepielski AM, DiBattista JD, & Carlson SM (2009) It's about time: the temporal dynamics of
phenotypic selection in the wild. *Ecol Lett* 12(11):1261-1276.
68. Siepielski AM, DiBattista JD, Evans JA, & Carlson SM (2011) Differences in the temporal
dynamics of phenotypic selection among fitness components in the wild. *P R Soc B*
278(1711):1572-1580.
69. Bolnick DI, *et al.* (2014) Major Histocompatibility Complex class IIb polymorphism influences
gut microbiota composition and diversity. *Molecular ecology* 23(19):4831-4845.

684
685