A rare MHC haplotype confers selective advantage in a free-living
ruminant

Wei Huang, Kara L Dicks, Jarrod D Hadfield, Susan E Johnston, Keith T Ballingall, Josephine M Pemberton

Abstract

The major histocompatibility complex (MHC) is the most variable gene family known in vertebrates. Parasite-mediated selection (PMS) is
believed to be the main force maintaining diversity at MHC genes, but it has proven hard to demonstrate the exact PMS regime that is acting
in natural populations. Demonstrating contemporary selection on MHC alleles is not trivial, and previous work has been constrained by limited
genetic tools, low sample sizes, short time scales and anticonservative statistical approaches. Here, we use a conservative statistical approach
to examine associations between MHC genes and fitness components, using haplotypes of expressed MHC class II genes in a large sample of
Soay sheep (*Ovis aries*) monitored over their lifetimes between 1989 and 2012. Of the eight MHC haplotypes (A-H) identified, we found that
haplotype C was associated with decreased male breeding success, haplotype D was associated with increased female life span, and haplotype
F was associated with decreased female life span. Consistent with the increased lifespan in females, haplotype D has increased in frequency
throughout the study period. Our results suggest the existence of contemporary balancing selection on MHC class II genes in Soay sheep as
might be expected under the hypotheses of rare allele advantage or fluctuating selection and do not support an effect of heterozygous
advantage.

Introduction

The vertebrate immune system has evolved to protect the host from infection through a variety of innate and adaptive immune mechanisms. Many of the proteins that control the specificity of immune function are the products of genes that are amongst the most diverse within vertebrate populations. Such diversity is believed to have evolved in response to pathogen diversity. Investigating the mechanisms shaping the diversity of immune genes has been a major focus in evolutionary biology (1). Genes located within the major histocompatibility complex (MHC) encode proteins that play important roles in the vertebrate immune system and have been intensively studied across many species, including model organisms (2-4). MHC molecules, the products of MHC class I and class II genes, present peptide fragments to antigen-specific receptors on T cells to induce adaptive immune responses. MHC class I molecules present peptides derived from intracellular pathogens such as viruses to CD8 +ve T cells; while MHC class II molecules present peptides derived from extracellular pathogens such as bacteria and parasites to CD4 +ve T cells (5).

MHC class I and II genes are the most polymorphic loci identified in vertebrate populations with hundreds of alleles described in many species (6). Pathogen-mediated balancing selection is thought to be the principal driver maintaining such high levels of allelic diversity (7-10). However, the specific selection mechanisms which act to maintain high levels of MHC allelic diversity are not clear (3, 11, 12). Three models have been proposed to explain how PMS acts to maintain such diversity:

1) *Heterozygote advantage*: individuals which are more heterozygous across the MHC can respond to a greater variety of pathogens. i.e. there is a consistent advantage to heterozygosity regardless of the alleles involved. As a result, variation at MHC loci is selected and maintained (13).

2) *Rare-allele advantage*: there is strong selection on pathogens to evade immune protection provided by the most common host MHC alleles. Therefore, rare alleles have a selective advantage. However, the advantage disappears as the frequency of a protective MHC allele increases and is absent at equilibrium. The arms race between hosts and pathogens sets up a cyclical process causing MHC alleles and pathogens to fluctuate, meaning that MHC diversity is maintained by negative frequency-dependent selection (14). It should be noted that

43 since rare alleles appear most frequently as heterozygotes, this can manifest as heterozygote advantage for specific alleles under some
44 circumstances.
45
46 3) *Fluctuating selection*: a changing environment results in variation in the abundance of different pathogens in space and time, which
47   generates directional selection in different host subpopulations and/or at different time points. Thus, different alleles will be favoured in
48   different subpopulations or at different times, meaning MHC diversity is maintained as a consequence (15, 16). Unlike rare-allele
49   advantage the change in selection coefficients in time and/or space is not driven by coevolution in the pathogens.
50
To test each selection model we can examine associations between MHC diversity and parasite presence or parasite load; different patterns of
associations can be explained as a result of different selection mechanisms. An association between MHC heterozygosity and reduced parasite
load could be viewed as evidence of heterozygous advantage, while if MHC alleles or haplotypes are correlated with protection against a
particular parasite species, rare-allele advantage or fluctuating selection could be the mechanism (12). A number of studies on MHC genes in
natural populations have examined associations between MHC variation and parasites, often with conflicting results. Some studies found that
MHC heterozygosity or certain MHC alleles confer resistance to parasite infection (17-19), while another study found no association between
MHC alleles and parasite load (20). However, prior to the development of high throughput sequencing technologies, such studies were
probably limited by incomplete data on the diversity and abundance of the parasite populations present. As a consequence, parasites that
strongly select for specific MHC alleles or haplotypes may have been missed. Therefore, quantifying relationships between MHC alleles and
pathogens may not have been able to reliably elucidate which selective mechanisms were operating. An alternative means to test for selection
on MHC genes would be to examine MHC-dependent fitness patterns (4, 21).
62
As with studies examining associations between MHC and parasites, MHC-fitness associations can also indicate different selection mechanisms.
Under heterozygote advantage, heterozygotes are always fitter regardless of the alleles they carry, so we predict a positive association
between MHC heterozygosity and fitness. Under rare-allele advantage, if MHC alleles have reached equilibrium frequency, no association
between MHC alleles and fitness should be detectable. However, when MHC alleles are not in equilibrium, we may detect selection favouring
specific rare alleles and probably also disfavouring specific common alleles. Under fluctuating selection, we would also observe specific alleles
being favoured or disfavoured by selection. But unlike rare-allele advantage, this should occur regardless of their frequency. However, it is

hard to distinguish rare-allele advantage and fluctuation selection by examining MHC-fitness associations (12). Because measuring fitness is
generally challenging except in some long-term individual-based studies of natural populations, only a few studies have tested associations
between fitness and MHC variation in the wild. Moreover, most such studies have only tested associations between MHC and a single fitness
component, again with mixed results. First, some MHC studies in wild populations have found a positive association between MHC
heterozygosity and fitness. For instance, the number of MHC alleles was found to be positively associated with increased apparent survival
(return rate from initial year of capture) in common yellowthroats *(Geothlypis trichas)* (22). Similarly, Banks et al. (23) found there was an
association between MHC heterozygosity and survival in a natural population of mountain brushtail possums *(Trichosurus cunninghami)*.
Second, some studies have reported associations between fitness and specific MHC alleles or supertypes (clusters of MHC alleles with similar
physicochemical properties at their antigen-binding sites). For example, Lukasch et al. (24) found that a specific MHC allele was associated with
higher survival of house sparrow fledglings *(Passer domesticus)*. Another study of great tits *(Parus major)* using mark-recapture data identified
three MHC supertypes associated with fitness components, one with increased survival, one with increased lifetime reproductive success and
another one with decreased lifetime reproductive success (25). Kloch et al. (26) also found that two MHC alleles from different clusters were
associated with winter survival in root voles *(Microtus oeconomus)*. Finally, there are some studies where no associations are detected. In
collared flycatcher (*Ficedula albicollis*), no association between MHC genes and lifetime reproductive success were found (20). Similar in
house sparrow (*Passer domesticus*), no significant association between MHC genes and survival were identified (27).
84
Many studies of selection on the MHC are also limited by the depth and quality of the genetic data available. Within a typical mammalian MHC
region, multiple closely linked and duplicated genes are present which are inherited together as a haplotype. With the development of next
generation sequencing (NGS) technologies, we are now able to cost-effectively genotype large numbers of samples (28, 29). However, in
species where the MHC region is poorly characterised such as in fish or passerine birds, it is difficult to assign a sequence to an individual locus
or identify if a locus is functional. Without locus-specific genotyping, multiple loci may co-amplify, which will impact subsequent testing for
associations between MHC diversity and fitness. Even if locus-specific genotyping is possible, most studies have focused on a single locus and
ignored linkage with other loci (23, 26, 30). Locus-specific genotyping and characterisation of haplotypes is desirable when testing for
associations between fitness and MHC diversity in wild populations.
93

Lastly, statistical analyses of selection on MHC genes require greater sophistication than is often deployed. Measures of parasite load and
fitness are quantitative traits likely to be determined by both environmental factors and multiple genetic factors, including MHC genes. As such,
it is possible that many studies of contemporary selection confound MHC similarity with genetic relatedness, so an MHC effect could be due to
loci elsewhere in the genome rather than the MHC itself. Selection on MHC genes in natural populations has mostly been assessed in long-
term individual-based studies which include a lot of genetically related individuals (25, 31). To our knowledge, no MHC study in a natural
population has been conducted in a quantitative genetic framework and thus they are anti-conservative. Similarly, given the ubiquity of
inbreeding depression, any analysis of selection on MHC testing for heterozygote advantage should include an estimate of individual
inbreeding to ensure MHC effects are not confounded with genome-wide inbreeding effects.
102
Here, we present the first investigation of contemporary selection acting on the MHC in a free-living population that aims to overcome the
issues raised above. The Soay sheep (*Ovis aries*) population on the island of Hirta, St Kilda is one of the most intensively studied wild animal
populations in the world (32). Since 1985, individual Soay sheep living in the Village Bay study area have been followed from birth, through all
breeding attempts, until death. In addition, using a combination of observation and SNP genotypes, a multigenerational pedigree for the
population has been assembled which includes annual and lifetime measures of fitness for both sexes (33). Previous work characterised
diversity at the MHC class II locus *DRB1* using an internal microsatellite and identified associations between MHC genotype, fitness and
parasite infection (34, 35). However, the only fitness measure studied was juvenile survival. Another study using the same genetic dataset of
MHC-linked microsatellite alleles found higher temporal and spatial variation in these loci than in putatively neutral microsatellite loci in other
parts of the genome, suggesting fluctuating selection on MHC genes (36). Recently, using sequencing-based genotyping of functional MHC
class II DR and DQ loci we have shown that eight MHC haplotypes are segregating in the study population (37). Using a panel of 13 SNPs, MHC
class II haplotypes were characterised for 5349 individuals sampled between 1989 and 2012 (38). While diplotypes were consistently in Hardy-
Weinberg equilibrium, haplotype frequencies are more even than expected by chance across all life history stages and standing population
samples (Ewens-Watterson test), which is consistent with balancing selection (38). Here, we use this haplotype data to investigate associations
between (1) MHC heterozygosity and fitness and (2) individual MHC haplotypes and fitness. We fit MHC haplotype within quantitative genetic
models in order to distinguish the effect of MHC haplotypes from genome-wide effects. In addition (3), we perform gene-drop simulations to
examine whether haplotype frequencies have changed more than expected from a neutral process.

Methods

Study system

Soay sheep have lived virtually unmanaged on the island of Soay, in the St. Kilda archipelago, for thousands of years. In 1932, 107 Soay sheep were introduced to the larger neighbouring island of Hirta and have been living there unmanaged since. From 1985, a long-term study has been conducted on the sheep resident in the Village Bay area of Hirta to investigate ecological and evolutionary questions (32). The vast majority of lambs are born in April or May of each year. At their first live capture, lambs are ear-tagged to enable life-long identification and discs of ear tissue removed in the tagging process are used for DNA extraction. Parentage is inferred for each individual using a subset of 315 unlinked SNPs derived from the Illumina Ovine 50K SNP array, on which most individuals alive since 1990 have been genotyped (39). In addition, the R package Sequoia (40) was used to cluster half-siblings which share a parent that has not been genotyped. In cases where no SNP genotypes were available, a small number of parentage inferences were made using field observations (for mothers) or a previous microsatellite genotype approach (41). The pedigree used here includes 7447 individuals, with 7014 paternal links and 6229 maternal links. This study investigated associations between MHC haplotypes and five fitness components defined as follows:

- Juvenile survival (JS), a binary variable indicating whether a lamb survived from birth until May 1st of the following year.
- Annual Survival (AS), a binary variable indicating whether an individual survived or not in a given calendar year;
- Life-span (LS), the age of an individual when it died;
- Annual Breeding Success (ABS), the number of offspring an individual produced in a given calendar year;
- Lifetime Breeding Success (LBS), the number of offspring an individual produced over its lifetime;

Analyses of LBS, ABS, LS and AS only included sheep that survived the juvenile period.

MHC Class IIa Haplotyping

The genetic data used in this study were obtained from a previous study (37, 38). In that study the seven expressed loci (*DRB1*, *DQA1*, *DQA2*, *DQA2-like*, *DQB1*, *DQB2* and *DQB2-like*) within the MHC class IIa region were characterised in 118 selected Soay sheep using sequencing-based genotyping. A total of eight MHC haplotypes were identified and named A to H. These haplotypes were confirmed in an additional 94

individuals that were unrelated based on the pedigree. A panel of 13 SNPs located in the region of MHC class IIa haplotypes was then selected
to impute the eight MHC class IIa haplotypes in 6752 Soay sheep genotyped using Kompetitive Allele-specific PCR (KASP). This panel included
11 SNPs known to be variable in Soays based on data from the Illumina Ovine infinium HD chip (42) and two other SNPs identified within *DQA1*
gene (42). After quality control, the diplotypes of 5349 individuals that lived in the study area between 1985 and 2012 were identified. As we
investigate cohort effects in the statistical analyses, we excluded animals born before 1989 since too few were genotyped to be representative
of their birth cohort; we also removed the cohort born in 2010 which had a high genotyping failure rate in the KASP genotyping assay. We also
excluded any individuals previously subjected to experimental treatments which may have affected their survival or breeding performance (e.g.
anthelmintic boluses). After these exclusions, a total of 3440 individuals (1750 males and 1650 females) remained for statistical modelling of JS
(Table 1). The frequency of rarest haplotype (D) was 3.02%. With the exception of animals homozygous for haplotype D (N=5), moderate
numbers of individuals (N>20) of each diplotype (i.e. haplotype complement) are represented (Supplementary 1).
154

Statistical analysis

We used generalized linear mixed models (GLMMs) to study the associations between MHC haplotypes and fitness components. The primary
models were animal models (AMs) including a covariance structure proportional to the pedigree relatedness matrix, but we also repeated all
tests without pedigree information. In null models, we fitted a list of fixed and random effects for each fitness component (Table 1) including
genomic inbreeding (F_{grm}) as a fixed effect, as a previous study demonstrated inbreeding depression in this population (43). None of the five
fitness components in our study had a normal distribution, so we used Markov chain Monte Carlo techniques implemented in MCMCglmm v
2.28 (44) in R v3.5.2 (45) to run models. As the fitness components have different ontogenies and distributions between males and females
(Supplementary 2), separate models were built for males and females. For both males and females, we used probit regression for JS and AS
and we used Poisson regression for LS and LBS. As females can only have up to two offspring each year while males had between 0 and 24, we
used multiple category probit regression for female ABS and Poisson regression for male ABS. To examine whether there were associations
between MHC genes and fitness components, we fitted all haplotypes simultaneously into the null model. Each haplotype was fitted either as
a count (0, 1 or 2 copies) in the dosage models or as 1/0 in the presence/absence models. These two models capture the extreme scenarios of
pure additive gene action and complete dominance respectively. We included MHC heterozygosity as a fixed effect in both dosage and
presence/absence models, which assumes a common dominance deviation across MHC diplotypes. We performed a Wald test using the
posterior mean and posterior covariance matrix to test whether haplotypes explained variation in fitness. In these models, Haplotype H was

treated as a reference and dropped in our results so the results for individual haplotypes from our models are relative to haplotype H.

Therefore, when a Wald test for haplotype differences was significant, we conducted an additional analysis (Supplementary 3) comparing the

estimate for each haplotype with the mean of the estimates for all other haplotypes to identify individual haplotype effects.

Table 1. Fixed and random effects fitted in null model and sample sizes of each fitness component. Sample sizes are shown by number of

individuals (records).

Fitness component	Fixed effects			Random effects			Sample sizes	
	Litter Size	Age (years, quadratic)	Inbreeding ($F_{grm)}$	Birth Year	Year of Fitness Measure	ID	Female	Male
JS	×		×	×			1650	1750
AS		×	×	×	×	×	551 (3093)	529 (1610)
LS			×	×			574	545
ABS		×	×	×	×	×	574 (3667)	545 (2155)
LBS			×	×			574	545

Gene-drop analysis

We performed gene-drop simulations with a custom script in R v3.5.2 (45) to model the expected frequency change of MHC haplotypes under

genetic drift given the pedigree structure. Hereafter, the first four birth cohorts (1989 to 1992) are defined as the "founder" cohorts, whereas

all subsequent cohorts are defined as "simulated" cohorts (1993 to 2012). A total of 10,000 simulations were conducted as follows. Individuals

in the founder cohorts were assigned their observed diplotypes; founder cohort individuals with unknown diplotypes were assigned one by

either: (a) sampling a haplotype from each parent assuming Mendelian segregation; or (b) sampling a haplotype with the probability of the

observed haplotype frequencies in the same birth cohort when one or more parents were unknown. In the simulated cohorts, each individual

diplotype was sampled using approaches (a) and (b) above. By this method, simulated diplotypes were generated for each individual in the

pedigree born after 1992. Using these data, we compared the observed and simulated changes in MHC haplotype frequency for individuals

with known diplotypes, using a linear regression with cohort year (1993 to 2012) as the predictor variable. In cases where the observed slope

fell within the top or bottom 2.5% of simulated slopes, the allele frequency was deemed to have been subject to directional selection.

Results

Associations between MHC and fitness components

192 The results of models with and without pedigree information were largely similar (Supplementary 4 to 6). Here we report the results of animal

193 models, i.e. including the pedigree as a random effect.

195
196 Figure 1. Associations between MHC heterozygosity and fitness components in Soay sheep in the dosage and presence/absence animal models
197 in females (circles) and males (triangles). Bars represent the 95% credibility intervals and those not overlapped with zero represent significant
198 results.
199
200
201

Table 2. Results (p value) of Wald tests for animal models (d.f.=7). Bold numbers show significant results (p <0.05).

Fitness component	Female		Male	
	Dosage model	Presence/absence model	Dosage model	Presence/absence model
JS	0.47	0.35	0.46	0.49
AS	0.068	0.066	0.43	0.56
LS	**0.039**	**0.046**	0.54	0.62
ABS	0.99	0.98	**0.024**	0.069
LBS	0.16	0.16	**0.044**	0.071

206
Figure 2. Associations between MHC haplotypes and fitness components in Soay sheep derived from animal models in females (A) and males
(B). Only fitness traits in which Wald tests were significant in at least either dosage or presence/absence model are shown here. Bars represent
the 95% credibility intervals. The solid lines represent the model intercept at haplotype H and posterior means and credible intervals for each
haplotype are plotted relative to H. Dashed lines indicate the average posterior mean of all eight haplotypes. Circles showed that Wald test
was not significant for this model. For models that Wald tests were significant, haplotype effects that were significantly different from the
mean are presented in squares with non-significant ones presented in triangles.

Juvenile survival

215 We found a positive association between MHC heterozygosity and female JS in the presence/absence model (Figure 1). Considering haplotype
216 effects, the Wald test was not significant in either the dosage or presence/absence models for either sex (Table 2, Supplementary Figure 4.2).

Annual survival

219 There was no association between MHC heterozygosity and annual survival (AS) in either sex (Figure 1). The Wald test for haplotype effects in
220 AS was not significant in either the dosage or presence/absence models for either sex (Table 2, Supplementary Figure 4.2).

Lifespan

There was a negative association between MHC heterozygosity and male life span (LS) in the dosage model and a negative association
between MHC heterozygosity and female life span (LS) in the presence/absence model (Figure 1). In the Wald test for haplotype effects a
significant association was found for female LS in both dosage model and presence/absence models, but neither test was significant for males
(Table 2, Figure 2A). Haplotype D was associated with increased female life span in the presence/absence model. Haplotype H was associated
with decreased female life span in the dosage model while haplotype F was associated with decreased female life span in both the dosage and
presence/absence models (Figure 2A).

Annual breeding success

231 There was no association between MHC heterozygosity and annual breeding success (ABS) in either sex (Figure 1). In the Wald test for
232 haplotype effects, a significant association was found for male ABS in the dosage but not the presence /absence model; neither test was
233 significant in female ABS (Table 2). Haplotype C was associated with decreased male ABS in the dosage model (Figure 2B).

Lifetime breeding success

236 There was a negative association between MHC heterozygosity and male lifetime breeding success (LBS) in the dosage model (Figure 1). The
237 Wald test for haplotype effects was significant in male LBS but only in the dosage model; neither test was significant in female LBS (Table 2).
238 Haplotype C was associated with decreased male LBS in the dosage model (Figure 2B).
239
240 The results obtained from models not including the pedigree were largely consistent with those above for associations between MHC
241 heterozygosity and fitness components (Supplementary 4). For associations between individual MHC haplotypes and fitness components,
242 estimates were largely similar but there were minor variations in terms of significance of Wald tests (Table 1, Supplementary 4). The Wald test
243 was no longer significant for male LBS in the dosage model but the Wald test was significant for male ABS in the presence/absence model.
244 Haplotype C was still associated with decreased male ABS in the presence/absence model (Supplementary 4).
245
246

247 # Gene-drop analysis

248
249 All haplotype frequency changes were within expectation due to drift alone except for haplotype D (Figure 3, Supplementary 7). Of the 10000
250 simulations for Haplotype D, only 272 simulated regression slopes were greater than the slope generated from real data (p=0.028). However,
251 this haplotype frequency change of haplotype was not significant for a two-tailed test.
252

254 Figure 3. Results of the gene drop analysis for each haplotype. The black lines represent the result of simulations (N=10000) and the red lines

255 represent the observed frequency of each haplotype in the Soay sheep population over the same time interval.

Discussion

In this study we investigated associations between MHC class II genes (heterozygosity and specific haplotypes) and five fitness components in a sample of individually-monitored free-living ruminants using generalized linear mixed models. We fitted individual MHC haplotype effects as both dosage and presence/absence and included MHC heterozygosity in all models. In addition, we included genetic relatedness to eliminate potential false positive results due to other genomic regions in common in related individuals. We found some associations between MHC heterozygosity and fitness components but the associations differed somewhat between dosage and presence/absence models. MHC heterozygosity was associated with increased female juvenile survival and decreased female LS in the presence/absence but not the dosage models; and MHC heterozygosity was associated with decreased male LS and LBS in the dosage models but not the presence/absence models. We also identified statistically significant associations between specific MHC class II haplotypes and three fitness components: female LS, male ABS and male LBS. The associations between specific MHC haplotypes and female LS were identified in both dosage and presence/absence models while associations in male ABS and LBS were only identified in the dosage models.

As outlined above, evidence of MHC heterozygote fitness advantage has been reported in some previous studies (22, 23, 31) although other studies did not support this hypothesised mechanism. In addition, theoretical work has questioned the role of heterozygote advantage in maintaining MHC diversity and suggested that heterozygote advantage alone is not sufficient to account for the high degree of MHC polymorphism (46). In this study we only found a positive association between MHC heterozygosity and female JS in the presence/absence model which was not reflected in the dosage model. We did not find any positive associations between MHC heterozygosity and any adult fitness component. Indeed, we found that MHC heterozygosity was associated with decreased male LS and male LBS in the dosage model as well as decreased female LS in the presence/absence model which is contradictory to a fixed advantage for MHC heterozygosity. Overall, it seems unlikely that allelic diversity in MHC class II genes in Soay sheep is maintained by heterozygous advantage.

If heterozygote advantage does not explain the maintenance of MHC diversity then selection favouring or disfavouring particular haplotypes, as is likely under negative frequency-dependent selection when not in equilibrium or by the fluctuating selection hypothesis may be present (25, 35). Indeed, we found significant associations between specific MHC haplotypes and three adult fitness components. Haplotype D was associated with increased female life span while haplotypes F and H were associated with decreased female life span. Haplotype C was

associated with reduced male annual breeding success and lifetime breeding success. If these MHC-linked fitness effects are strong enough
and not in conflict with other effects, we should be able to see changes in haplotype frequency over time in response. Examination of effects
of MHC-fitness associations on both sexes suggests haplotype D often had positive coefficients in adult fitness components in both sexes, but
the patterns for the other haplotypes showing significant associations in specific fitness components are much less clear (Supplementary 4).
Consistent with this, the gene drop analysis showed that haplotype D has increased through the study period, although the change was not
significant for a two-tailed test (p=0.028). Taken together, modelling of fitness component associations and gene-drop simulations suggest
there is contemporary selection on MHC genes in Soay sheep, mainly comprising selection favouring haplotype D.
288
Regarding the specific mechanism of selection, one possibility is that the MHC class II genes in Soay sheep are under rare allele advantage
caused by an arms race between hosts and parasites. Previous theoretical work has suggested that frequency-dependent selection resulting
from host-parasite coevolution could favor novel alleles and is important in maintaining MHC polymorphism (47, 48). A recent empirical study
using guppies (*Poecilia reticulata*) and their common ectoparasite in controlled infections showed that novel MHC variants could reduce the
infection regardless of individual MHC variability and genetic background, which provides direct evidence of rare allele advantage in shaping
MHC diversity (49). Under rare allele advantage, fitness differences and frequency changes are only expected when the system is not in
equilibrium. If haplotypes are rarer than their equilibrium frequency, they will confer an advantage in fitness. Haplotype D is the rarest
haplotype in the Soay population with a very low haplotype frequency at the beginning of our study (only 7 individuals in the founder cohorts
1989-1992). Therefore, positive selection on haplotype D could come from rare-allele advantage (Figure 3).
298
However, our results are also consistent with the fluctuating selection hypothesis. Previous theoretical work showed that diversity of MHC
genes could be maintained by temporally fluctuating selection on its own in the absence of heterozygote and rare allele advantage if the
resistance alleles were dominant (15) although few empirical studies have found signatures of temporal fluctuation selection in a single
population. A study of the great reed warbler (*Acrocephalus arundinaceus*) identified significantly higher variation in MHC allele frequencies
between cohorts than at neutral loci suggesting that fluctuating selection may be acting on MHC genes (50). In our study, haplotype D may
confer the best resistance to the current parasite community on St Kilda under the fluctuating selection hypothesis.
305

306 Our results are thus consistent with both rare allele advantage and fluctuating selection and we are not capable of differentiating them. The
307 key differences between these two mechanisms are first, that fluctuating selection requires a changing external environment while rare allele
308 advantage is caused by an arms race between host and parasites, and second, that selection is directional for fluctuating selection while for
309 rare allele advantage selection is cyclical. However, not only are these differences hard to identify in real time, but also they are not mutually
310 exclusive, meaning that it is very hard to differentiate rare allele advantage and fluctuating selection in wild populations (12).
311
Several aspects of our analysis are novel in studies of MHC-fitness associations. To our knowledge, this is the first study which uses animal
models and includes a genomic inbreeding estimator in a study of association between MHC and fitness components in wild populations. As
the AM accounts for relatedness, we predicted that it would be more conservative and show fewer associations than GLMMs without
accounting for relatedness. In fact, we found the results to be largely consistent between the two models. This may be due to the generally
low heritability of the fitness traits (Supplementary 8). However, for traits which have a higher heritability, the results of animal models will be
more conservative as they would exclude the false positive effects caused by genomic regions other than the MHC. We are also the first to
examine MHC-fitness associations using both dosage models and presence/absence models in a wild population. We found that dosage
models and presence/absence models were generally consistent in terms of estimates (Figure 2). One possible explanation for this is that for
most haplotypes, homozygotes were relatively rare compared with heterozygotes. Thus, in some cases, there may not have been sufficient
power to differentiate dosage and presence/absence models.
322
Long term individual-based studies like the Soay sheep project also enable repeat measurements of the strength and direction of selection
over time (51, 52). In a previous study, three MHC-linked microsatellite alleles were found to be associated with survival in Soay sheep using
individuals from 1985 to 1994 (35). Microsatellite alleles OLADRB 257 and 205 were associated with decreased juvenile and yearling survival
respectively whereas the OLADRB 263 allele was associated with increased yearling survival. These microsatellite alleles can be linked to the
MHC class II haplotypes studied here: OLADRB allele 257 is linked to haplotype B (Supplementary 9). However in our analysis of juvenile
survival, we did not find any evidence for haplotype differences (Wald tests not significant; Table 2) although estimates for haplotype B in
juvenile survival are consistently negative in females. The inconsistency between the previous and current study is probably due to differences
in sample size and statistical methods. Additionally, as alleles 205 and 263 are associated with multiple haplotypes, and we have studied
annual survival rather than yearling survival, it is hard to draw further comparisons.

In summary, we used well-characterized MHC class II haplotypes to investigate MHC-linked fitness effects in Soay sheep. To our knowledge,

our dataset, with more than 3000 individuals, is the largest yet used to study selection on MHC genes in a wild population. Our results support

the existence of contemporary selection on MHC class II genes in Soay sheep and suggest rare allele advantage or fluctuating selection is the

mechanism in shaping MHC diversity in Soay sheep rather than heterozygous advantage.

Acknowledgments

We thank the National Trust for Scotland for permission to work on St. Kilda and QinetiQ for logistics and support. We thank J. Pilkington, I.

Stevenson and many volunteers who have helped with fieldwork on the island and all those who have contributed to keeping the project going

over many years. W. Huang was supported by Edinburgh Global research scholarship. K. Ballingall received funding from the European Union's

Horizon 2020 research and innovation programme under grant agreement no. 731014KB (VetBioNet) and acknowledges the support received

from the Scottish Government's strategic research programme. The long-term project on St Kilda, the KASP genotyping and the SNP array

genotyping were funded by the UK Natural Environment Research Council and the European Research Council.

References

1.   Obbard DJ, Welch JJ, Kim KW, & Jiggins FM (2009) Quantifying Adaptive Evolution in the Drosophila Immune System. *Plos Genet* 5(10).

2.   Acevedo-Whitehouse K & Cunningham AA (2006) Is MHC enough for understanding wildlife immunogenetics? *Trends Ecol Evol* 21(8):433-438.

3.   Bernatchez L & Landry C (2003) MHC studies in nonmodel vertebrates: what have we learned about natural selection in 15 years? *Journal of*

Evolutionary Biology 16(3):363-377.

4.   Piertney SB & Oliver MK (2006) The evolutionary ecology of the major histocompatibility complex. *Heredity* 96(1):7-21.

5.   Garrigan D & Hedrick PW (2003) Detecting adaptive molecular polymorphism, lessons from the MHC. *Am J Hum Genet* 73(5):375-375.

6.   Apanius V, Penn D, Slev PR, Ruff LR, & Potts WK (1997) The nature of selection on the major histocompatibility complex. *Crit Rev Immunol*

17(2):179-224.

7.   Oppelt C, Starkloff A, Rausch P, Von Holst D, & Rodel HG (2010) Major histocompatibility complex variation and age-specific endoparasite load in

subadult European rabbits. *Molecular ecology* 19(19):4155-4167.

8.   Radwan J, Kuduk K, Levy E, LeBas N, & Babik W (2014) Parasite load and MHC diversity in undisturbed and agriculturally modified habitats of the
ornate dragon lizard. *Molecular ecology* 23(24):5966-5978.
9.   Oliver MK, Telfer S, & Piertney SB (2009) Major histocompatibility complex (MHC) heterozygote superiority to natural multi-parasite infections in
the water vole (Arvicola terrestris). *Proceedings. Biological sciences* 276(1659):1119-1128.
10.   Westerdahl H, Asghar M, Hasselquist D, & Bensch S (2012) Quantitative disease resistance: to better understand parasite-mediated selection on
major histocompatibility complex. *Proceedings. Biological sciences* 279(1728):577-584.
11.   Jordan WC & Bruford MW (1998) New perspectives on mate choice and the MHC. *Heredity* 81:239-245.
12.   Spurgin LG & Richardson DS (2010) How pathogens drive genetic diversity: MHC, mechanisms and misunderstandings. *P R Soc B* 277(1684):979-988.
13.   Takahata N & Nei M (1990) Allelic Genealogy under Overdominant and Frequency-Dependent Selection and Polymorphism of Major
Histocompatibility Complex Loci. *Genetics* 124(4):967-978.
14.   Hughes AL & Nei M (1988) Pattern of Nucleotide Substitution at Major Histocompatibility Complex Class-I Loci Reveals Overdominant Selection.
Nature 335(6186):167-170.
15.   Hedrick PW (2002) Pathogen resistance and genetic variation at MHC loci. *Evolution; international journal of organic evolution* 56(10):1902-1908.
16.   Hill AVS, *et al.* (1991) Common West African Hla Antigens Are Associated with Protection from Severe Malaria. *Nature* 352(6336):595-600.
17.   Rivero-de Aguilar J, *et al.* (2016) MHC-I provides both quantitative resistance and susceptibility to blood parasites in blue tits in the wild. *J Avian Biol*
47(5):669-677.
18.   Westerdahl H, Stjernman M, Raberg L, Lannefors M, & Nilsson JA (2013) MHC-I Affects Infection Intensity but Not Infection Status with a Frequent
Avian Malaria Parasite in Blue Tits. *Plos One* 8(8).
19.   Westerdahl H, *et al.* (2005) Associations between malaria and MHC genes in a migratory songbird. *P R Soc B* 272(1571):1511-1518.
20.   Radwan J, *et al.* (2012) MHC diversity, malaria and lifetime reproductive success in collared flycatchers. *Molecular ecology* 21(10):2469-2479.
21.   Sutton JT, Nakagawa S, Robertson BC, & Jamieson IG (2011) Disentangling the roles of natural selection and genetic drift in shaping variation at
MHC immunity genes. *Mol Ecol* 20(21):4408-4420.
22.   Dunn PO, Bollmer JL, Freeman-Gallant CR, & Whittingham LA (2013) Mhc Variation Is Related to a Sexually Selected Ornament, Survival, and
Parasite Resistance in Common Yellowthroats. *Evolution; international journal of organic evolution* 67(3):679-687.
23.   Banks SC, Dubach J, Viggers KL, & Lindenmayer DB (2010) Adult survival and microsatellite diversity in possums: effects of major histocompatibility
complex-linked microsatellite diversity but not multilocus inbreeding estimators. *Oecologia* 162(2):359-370.
24.   Lukasch B, *et al.* (2017) Major histocompatibility complex genes partly explain early survival in house sparrows. *Sci Rep-Uk* 7.
25.   Sepil I, Lachish S, & Sheldon BC (2013) Mhc-linked survival and lifetime reproductive success in a wild population of great tits. *Molecular ecology*
22(2):384-396.
26.   Kloch A, Baran K, Buczek M, Konarzewski M, & Radwan J (2013) MHC influences infection with parasites and winter survival in the root vole
Microtus oeconomus. *Evol Ecol* 27(3):635-653.
27.   Karlsson M, *et al.* (2015) House sparrowPasser domesticussurvival is not associated with MHC-I diversity, but possibly with specific MHC-I alleles.
Journal of Avian Biology 46(2):167-174.

28. Babik W (2010) Methods for MHC genotyping in non-model vertebrates. *Mol Ecol Resour* 10(2):237-251.
29. Babik W, Taberlet P, Ejsmond MJ, & Radwan J (2009) New generation sequencers as a tool for genotyping of highly polymorphic multilocus MHC
system. *Mol Ecol Resour* 9(3):713-719.
30. Osborne AJ*, et al.* (2015) Heterozygote advantage at MHC DRB may influence response to infectious disease epizootics. *Molecular ecology*
24(7):1419-1432.
31. Brouwer L*, et al.* (2010) MHC-dependent survival in a wild population: evidence for hidden genetic benefits gained through extra-pair fertilizations.
Molecular ecology 19(16):3444-3455.
32. Clutton-Brock T & Pemberton JM (2004) Soay sheep - Dynamics and selection in an island population. *Cambridge University Press*.
33. Berenos C*, et al.* (2015) Heterogeneity of genetic architecture of body size traits in a free-living population. *Molecular ecology* 24(8):1810-1830.
34. Paterson S (1998) Evidence for balancing selection at the major histocompatibility complex in a free-living ruminant. *J Hered* 89(4):289-294.
35. Paterson S, Wilson K, & Pemberton JM (1998) Major histocompatibility complex variation associated with juvenile survival and parasite resistance in
a large unmanaged ungulate population (Ovis aries L.). *Proceedings of the National Academy of Sciences of the United States of America* 95(7):3714-
3719.
36. Charbonnel N & Pemberton J (2005) A long-term genetic survey of an ungulate population reveals balancing selection acting on MHC through
spatial and temporal fluctuations in selection. *Heredity* 95(5):377-388.
37. Dicks KL, Pemberton JM, & Ballingall KT (2019) Characterisation of major histocompatibility complex class IIa haplotypes in an island sheep
population. *Immunogenetics* 71(5-6):383-393.
38. Dicks K (2017) Unravelling major histocompatibility complex diversity in the Soay Sheep of St.Kilda. *University of Edinburgh*.
39. Berenos C, Ellis PA, Pilkington JG, & Pemberton JM (2014) Estimating quantitative genetic parameters in wild populations: a comparison of pedigree
and genomic approaches. *Mol Ecol* 23(14):3434-3451.
40. Huisman J (2017) Pedigree reconstruction from SNP data: parentage assignment, sibship clustering and beyond. *Molecular Ecology Resources*
17(5):1009-1024.
41. Morrissey MB*, et al.* (2012) The prediction of adaptive evolution: empirical application of the secondary theorem of selection and comparison to
the breeder's equation. *Evolution; international journal of organic evolution* 66(8):2399-2410.
42. Johnston SE, Berenos C, Slate J, & Pemberton JM (2016) Conserved Genetic Architecture Underlying Individual Recombination Rate Variation in a
Wild Population of Soay Sheep (Ovis aries). *Genetics* 203(1):583-+.
43. Berenos C, Ellis PA, Pilkington JG, & Pemberton JM (2016) Genomic analysis reveals depression due to both individual and maternal inbreeding in a
free-living mammal population. *Mol Ecol* 25(13):3152-3168.
44. Hadfield JD (2010) MCMC Methods for Multi-Response Generalized Linear Mixed Models: The MCMCglmm R Package. *J Stat Softw* 33(2):1-22.
45. R Core Team (2013) R: A language and environment for statistical computing. .
46. De Boer RJ, Borghans JAM, van Boven M, Kesmir C, & Weissing FJ (2004) Heterozygote advantage fails to explain the high degree of polymorphism
of the MHC. *Immunogenetics* 55(11):725-731.

47.   Ejsmond MJ & Radwan J (2015) Red Queen Processes Drive Positive Selection on Major Histocompatibility Complex (MHC) Genes. *Plos Comput Biol*
11(11).
48.   Borghans JAM, Beltman JB, & De Boer RJ (2004) MHC polymorphism under host-pathogen coevolution. *Immunogenetics* 55(11):732-739.
49.   Phillips KP, *et al.* (2018) Immunogenetic novelty confers a selective advantage in host-pathogen coevolution. *Proceedings of the National Academy*
of Sciences of the United States of America 115(7):1552-1557.
50.   Westerdahl H, Hansson B, Bensch S, & Hasselquist D (2004) Between-year variation of MHC allele frequencies in great reed warblers: selection or
drift? *J Evol Biol* 17(3):485-492.
51.   Siepielski AM, DiBattista JD, & Carlson SM (2009) It's about time: the temporal dynamics of phenotypic selection in the wild. *Ecol Lett* 12(11):1261-
1276.
52.   Siepielski AM, DiBattista JD, Evans JA, & Carlson SM (2011) Differences in the temporal dynamics of phenotypic selection among fitness components
in the wild. *P R Soc B* 278(1711):1572-1580.