Negative selection may cause grossly altered but broadly stable karyotypes in metastatic colorectal cancer

William Cross*[1,2], Salpie Nowinski*[1,3], George D Cresswell*[3,23], Maximilian Mossner*[1,3], Abhirup Banerjee*[1], Bingxin Lu*[4,5], Marc Williams[1,6], Georgios Vlachogiannis[3], Laura Gay[1], Ann-Marie Baker[1,3], Christopher Kimberley[1], Freddie Whiting[1,3], Hayley Belnoue-Davis[11], Pierre Martinez[1,7], Maria Traki[1], Viola Walther[1], Kane Smith[1,3], Javier Fernandez-Mateos[3], Erika Yara[3], Erica Oliveira[3], Salvatore Milite[22], Giulio Caravagna[3], Chela James[22], George Elia[1], Alison Berner[1,2], Ryan Changho Choi[1,8,9,], Pradeep Ramagiri[10], Ritika Chauhan[10], Nik Matthews[10], Jamie Murphy[11], Anthony Antoniou[8], Susan Clark[8,11], Miriam Mitchison[12], Jo-Anne Chin Aleong[13], Enric Domingo[14], Inmaculada Spiteri[3], Stuart AC McDonald[1], Darryl Shibata[15], Miangela M Lacle[16], Lai Mun Wang[17,18], Morgan Moorghen[8], Ian PM Tomlinson[19], Marco Novelli[2,12], Marnix Jansen[2,12], Alan Watson[20], Nicholas A Wright[1], John Bridgewater[2,12], Manuel Rodriguez-Justo[2,12], Chris P Barnes[4,5], Hemant Kocher[13,21], Simon J Leedham[17], Andrea Sottoriva^[22,3], Trevor A Graham^[1,3]

* These authors contributed equally to this manuscript.
^ For correspondence: andrea.sottoriva@fht.org and trevor.graham@icr.ac.uk

Centre for Genomics and Computational Biology, Barts Cancer Institute, Barts and the London School of Medicine and Dentistry, Queen Mary University of London, Charterhouse Sq, London, UK.
UCL Cancer Institute, UCL, 72 Huntley St, Bloomsbury, London, UK
Centre for Evolution and Cancer, The Institute of Cancer Research, London, UK.
Department of Cell and Developmental Biology, University College London, London, UK
UCL Genetics Institute, University College London, London, UK
Computational Oncology, Department of Epidemiology and Biostatistics, Memorial Sloan Kettering Cancer Center, 321 East 61st St., New York, NY 10065, USA
Univ Lyon, Université Claude Bernard Lyon 1, INSERM 1052, CNRS 5286, Centre Léon Bérard, Cancer Research Center of Lyon, Lyon, 69008, France
St Mark's Hospital, Harrow, London, UK
St. George Hospital, Sydney, Australia
Tumour Profiling Unit (TPU), Institute of Cancer Research, London, UK
Department of Surgery and Cancer, Imperial College London, London, UK
UCL Hospitals NHS Trust, London, UK
Barts Health NHS Trust, The Royal London Hospital, Whitechapel, London, UK.
Department of Oncology, University of Oxford, Oxford
Department of Pathology, University of Southern California Keck School of Medicine, Los Angeles, CA 90033, USA
Department of Pathology, University Medical Center Utrecht, Utrecht, Netherlands
Wellcome Trust Centre Human Genetics, University of Oxford, Roosevelt Drive, Oxford, UK
Ludwig Institute, University of Oxford, Oxford, UK
Cancer Research UK Edinburgh Centre, MRC Institute of Genetics and Molecular Medicine, University of Edinburgh, Western General Hospital, Edinburgh, UK
Department of Gastroenterology, Barts Health, Whipps Cross Hospital, Leytonstone, London, UK.
Centre for Tumour Biology, Barts Cancer Institute
Computational Biology Research Centre, Human Technopole, Milan, Italy
St. Anna Children's Cancer Research Institute, Vienna, Austria

Contributions:
Study conception and design: AS & TG
Funding and supervision: SL, HK, NW, AS, TG
Manuscript writing: WC, MM, SN, GC, MW, AS, TG

Bioinformatics: WC, SN, GC, MW, PM, GC, TG, SM, CJ
Mathematical modelling: BL, MW, AS, CB, TG
Experimental work: MM, LG, AB, CK, MT, VW, KS, SA, RC, HD, IS, NM
Provision of patient materials: SA, AB, JM, AA, SC, JC, ED, LM, IT, MMo, MMi, ML, AW, SM, MN, MRJ,
JB, HK, SL
Histopathology: MRJ, ML, MJ, NW, MN, LM, MM
58
59
Abstract

Aneuploidy, the loss and gain of whole and part chromosomes, is near-ubiquitous in cancer genomes
and likely defines cancer cell biology. However, the temporal evolutionary dynamics that select for
aneuploidy remain uncharacterised. Here we perform longitudinal genomic analysis of 755 samples
from a total of 167 patients with colorectal-derived neoplastic lesions that represent distinct stages
of tumour evolution through metastasis and treatment. Adenomas typically had few copy number
alterations (CNAs) and most were subclonal, whereas cancers had many clonal CNAs, suggesting that
progression goes through a CNA bottleneck. Individual CRC glands from the same tumour typically
had very similar karyotypes, despite evidence of ongoing instability at the cell level in patient
tumours, cell lines and organoids. CNAs in metastatic lesions sampled from liver and other organs,
after chemotherapy or targeted therapies, and in late recurrences were typically similar to the
primary tumour. Mathematical modelling and statistical inference indicated that these data are
consistent with the action of negative selection on CNAs that 'traps' cancer cell genomes on a fitness
peak defined by the specific pattern of chromosomal aberrations. These data suggest that the initial
progression of colorectal cancer requires the traversal of a rugged fitness landscape and subsequent
CNA evolution, including metastatic dissemination and therapeutic resistance, is constrained by
negative selection.

77
78

Introduction

Most cancer genomes are aneuploid[1]. This is the result of gains and losses of whole or part chromosomes which are attained through a variety of mechanisms that include chromosome mis-segregation, aberrant double strand break repair, and genome doubling[2]. In colorectal cancer (CRC), approximately 85% of cancers are classified as chromosomally unstable (CIN) by virtue of having an abnormal karyotype and aneuploidy[3]. The remaining ~15% of cancers exhibit genetic instability at the mutational level and are classified as hypermutant, but exhibit limited chromosome aberrations.

Aneuploidy likely has a causal role in cancer evolution. First, the observed ubiquity of aneuploidy in cancers and the prevalence of recurrent chromosome copy number alterations (CNAs) is suggestive that some CNAs are positively selected[4,5]. Second, aneuploidy is predictive of progression to cancer in premalignant diseases such as Barrett oesophagus[6] and ulcerative colitis[7]. Third, in established cancers, individual CNAs have prognostic value over-and-above single nucleotide alterations in key cancer driver genes[8]. Fourth, ploidy has been shown to have prognostic value in many types of established cancer[9], and notably within-tumour heterogeneity of CNAs, as principally measured by the number of lineages with distinct karyotypes, is prognostic pan-cancer[10]. In lung cancer, the diversity of CNAs but not single nucleotide variants (SNVs) has prognostic value[11].

Aneuploidy should also be subject to negative selection because the loss or gain of large genomic regions will affect the expression of thousands of genes and regulatory elements, some of which are likely to adversely affect tumour cell viability[12]. Moreover, it is likely that some chromosomal conformations prevent correct chromosomes segregation, resulting in cell death. Chromosomal-scale alterations are therefore expected to result in trade-offs between the dosages of positively- and negatively-selected elements[13], potentially leading to an apparently stable aneuploid genome. Chromosomal changes may also have a buffering effect against deleterious point mutations[14]. In such an evolutionary regimen, most alterations are either neutral or negatively selected[15]. Indeed, studies show that the burden of CNAs is non-monotonically related with prognosis across cancer types[10] (first observed in breast cancer[16]). As such, cancers carrying an intermediate level of alterations ('just right' aneuploidy) are associated with worse prognosis than cancers with lower or higher levels of aneuploidy. Observing negative selection may be complicated by the fact that less fit variants are effectively removed from the population and do not come to clinical prominence.

The spatial-temporal evolutionary dynamics that produce the aneuploidy observed in cancer genomes remain poorly understood, in part because longitudinal measurement is often not clinically-feasible and is compounded by the challenge of reconstructing ancestral history of chromosome gains and losses[17]. CRC presents a unique opportunity to track clonal evolution over space and time: pre-cancerous lesions (colorectal adenomas, CRAs) are occasionally 'caught in the act' of transformation and are found to contain a small focus of cancer (early CRCs). CRC frequently metastasizes to the liver, and repeated hepatic metastatectomy can provide a source of longitudinally sampled tumour material amenable for molecular analysis.

Results

Benign-to-malignant transformation through a CNA bottleneck

Here we investigated the evolution of aneuploidy in a large cross-sectional and longitudinally followed cohort of adenomas and cancers. Previously we observed that the degree of aneuploidy was much higher in a small cohort of (malignant) colorectal cancers than (benign) adenomas[18]. To validate this observation, we performed multi-region spatial sampling and genome-wide copy number alteration analysis (by shallow whole genome sequencing [sWGS] or SNP-array) in n=139 archival colorectal lesions (19 adenomas and 81 stage II/III CRCs, termed "regular CRCs"), and 39 early CRCs of which 23/39 (59%) contained residual adenoma (Fig 1A). In early CRCs, adenoma and

carcinoma regions were separately subsampled (minimum 2 regions per neoplasm). The percentage
of genome altered (PGA) by a CNA was lowest in adenomas and highest in regular CRCs (8.5% and
27.2% respectively, p=8.4x10^{-14}) and early CRCs (25.3%; Fig 1B). Gains were more prevalent than
losses in adenomas (p=0.0052; Fig S1A) whereas they occurred at similar frequency relative to
baseline ploidy in cancers (p=0.75, p=0.32, respectively; Fig S1A). Which CNA events tended to be
clonal or subclonal depended on tumour stage (Fig S1B).

136
Figure 1: Aneuploidy increases through colorectal cancer progression, and clonal diversity decreases. (A)
Heatmap showing copy number alterations (CNAs) in adenomas (orange; top), early cancers (green; middle)
and established cancers (blue; bottom). Thin black lines delineate cases, within black lines are multiple regions
of each case. Genome doubling (GD) status is shown on the left of the heatmap. Red=gain, blue=loss. (B)
Percentage genome altered (PGA) by adenomas, early cancers and then later stage cancers. PGA increases
through progression. (C) The proportion of CNAs that are subclonal decreases through progression from
adenomas to carcinomas. (D) Phase space of PGA versus proportion of subclonal CNAs that are subclonal.
Adenomas (orange) have low PGA but high CNA subclonality, whereas CRCs (blue) have high PGA but low CNA
subclonality. Early cancers (green) are more disparate in PGA and CNA subclonality. (E) The proportion of CNAs
that are subclonal is not significantly associated with overall survival in CRCs (Kaplan Meier analysis, groups
split on the median proportion of subclonal CNAs across the cohort). (F) Frequency of genome doubling (GD) by
tumour stage. GD is very rare in adenomas, more common in later stage cancers where it is usually found to be
clonal. (G) The proportion of CNAs that are subclonal by tumour stage and genome doubling status. Clonal
genome doubling is associated with a lower proportion of subclonal CNAs.

151
CNA subclonal diversity (the proportion of total altered genome not altered in all samples from a
case; Methods) was greatest in adenomas and low in regular CRCs (p=0.0016; Fig 1C; Table S1; Table

S2). Different measures to quantify CNA subclonality were highly correlated (Fig S2A). This suggests
that progression from benign to malignant tumours goes through an aneuploidy-divergence 'phase
space' from low-aneuploidy/high-diversity to high-aneuploidy/low-diversity (Fig 1D), indicating a
selective genetic bottleneck acting on CNAs. Consistently, within the early CRCs, the fraction of
subclonal CNAs was higher between the adenoma components than between cancer components
(p=1.2x10^-6; Fig S3A). Moreover, subclonal CNA fraction in cancer components of early CRCs was
significantly lower than in regular CRCs (p=0.046; Fig S3A; Fig S3B).
161
We trained a classifier to infer the presence of genome doubling (GD) from shallow WGS data
(Methods; Fig S4). GD was present in only a single adenoma and was clonal (Fig 1F). In regular CRCs,
GD was more common (16/81, 18%) and was almost always clonal (15/16, 94%; Fig 1F). Within early
CRCs, no adenoma component had GD, whereas a third of the early cancers (13/39) had evidence of
GD (n=13/39; Fisher's test p=0.0011) and typically this was clonal (11/13, 85%). These observations
suggest that GD occurs at, or shortly prior to, the time of invasion. The proportion of subclonal CNAs
was lower in GD than non-GD CRCs (p=0.00084; Fig 1F) and divergent CNAs were significantly
shorter in length (p=0.015; Fig S5A). Therefore, following GD the aneuploid genome accrues little
further alteration within primary CRCs, suggesting the process of genome doubling produces a fit
genotype which drives passage through the CNA bottleneck.
172
We investigated the relationship between CNA diversity and clinical outcome in regular CRCs. The
diversity of pre-existing adaptive phenotypes in a population should correlate with population
evolvability, and consequently more diverse populations are more likely to harbour an individual
pre-adapted to a new selective pressure. Accordingly, intra-tumour clonal diversity has been
observed as a pan-cancer prognostic biomarker[10] and it is prognostic in the premalignant lesion
Barrett's Oesophagus[19]. However, in our cohort, the proportion of subclonal CNAs was not
associated with overall survival (comparison of cases above and below median proportion of
subclonal CNAs ; p=0.37; Fig 1E; Table S3) Consequently, these data suggested that intra-tumour
CNA diversity did not strongly reflect the heterogeneity of adaptive phenotypes in regular CRC, and
was consistent with prior observations of a lack of positive selection experienced by CRC[20,21]. The
presence of GD was also not associated with overall survival (p=0.56, Fig S5C).
184
Limited CNA diversity at single gland resolution in CRCs
Previous reports using fluorescence *in situ* hybridisation (FISH) to measure CNAs at single cell
resolution found widespread chromosomal instability in colorectal cancer cell lines[22] and primary
tumours[20]. To investigate CNA heterogeneity at higher resolution we exploited the fact that well-to-
moderately differentiated CRCs are made up of a collection of glands, each composed of a few
thousand cells recently derived from a common ancestor[20,23]. Glands can reasonably be considered
as units of selection in CRC, as gland fission likely expands the cancer cell population[20]. We extracted
a total of 307 individual colorectal cancer glands from the notional 'left' and 'right' opposite sides of
an additional 6 regular CRCs (Fig 2A, Table S3) and performed high-depth exome sequencing (59
glands) or shallow whole genome sequencing (sWGS; 248 glands). In addition, we also analysed at
least two bulk samples composed of numerous glands from each tumour.
196
At gland level (individual clone) resolution the pattern of CNAs was quite similar within cancers (Fig
2B, Table S4-5). In all cases, a 'core karyotype' was evident across the glands from that case, with a
small numbers of glands showing additional subclonal CNAs (mean CNAs per gland = 7, mean clonal
CNAs=6.5). We note that intra-tumour CNA heterogeneity was detected, but we emphasise the
clonality of most CNAs. Zooming in further, single cell analysis by FISH in four additional cases (Fig
S6) and single cell sequencing in case C274 (reported in Bollen *et al.*[24]) confirmed ongoing instability
at the level of individual cancer cells within glands.
204

205
Figure 2: Lack of CNA heterogeneity at the level of individual colorectal cancer glands. (A) (left column)
Heatmaps showing CNAs in individual glands in each of six CRCs. Glands were sequenced from arbitrary
opposite sides of each CRC as indicated by the heatmap on the right-hand side. Red hue colours indicate gains,
blue losses. (centre column) Phylogenetic trees reconstructed from individual gland single nucleotide alteration
calls from exome sequenced glands. Trees are balanced (comparison with Yule model). Note the trunk length
is not shown to scale, number above trunk represent number of clonal SNVs. Branches are coloured by notional
tumour side where the gland was collected. HI=homoplasy index. (right column) Results of Bayesian inference
to determine whether the observed CNA diversity is best supported by a model with or without stabilising
selection. Model selection by deviance information criteria (DIC) suggests that the stabilising selection model is
better supported for four patients (10 simulation replicates used for each boxplot). (B) Schematic showing
shape of fitness landscape used in the computational simulations of stabilising selection. The model parameter
determines the steepness of the single fitness peak in the landscape, at position defined by the above
heatmap illustrating the optimal CNA pattern. (C) Heatmaps showing CNAs of the randomly sampled glands in
one simulation as function of selective strength when CNA rate is fixed (0.2). The simulations were based on
data from patient C319, and 77 glands were sampled when the population size reached 100000. (D) Genetic
divergence (top) and percentage of the genome altered (bottom) as a function of CNA rate and selection
strength (10 replicates for each case, the same simulation settings as in C). Strong selection on an initially
optimal clone supresses diversity and CNA accumulation and higher mutation rates lead to more diversity and
more CNA accumulation. The box plots show the median (centre), 1st (lower hinge), and 3rd (upper hinge)

227
Single nucleotide variants (SNVs) were called from individual gland exome sequencing data and were
used to reconstruct phylogenetic trees (Methods; Fig 2C, Table S6). Trees were balanced (ancestral
nodes have equal numbers of offspring; Yule Model; p>0.05 for all cases) suggestive of a lack of
stringent subclonal positive selection, in accordance with previous reports[20,21]. Individual glands
from MSS tumours contained on average 156 SNVs (range: 68-232) which equated to 40 more SNVs
than the most recent common ancestor (MRCA) cell of the cancer (371 SNVs for hypermutant cases).
We calculated the SNV accumulation rate per year using only C>T mutations in the four trinucleotide
contexts associated with COSMIC mutational signature 1 (ACG, CCG, GCG, TCG) which are known to
accrue at a constant rate in CRC[25]. This analysis produced an estimate of the time to the MRCA of 24
years (range: 11-30 years). Despite this elapsed time, we observed little subclonal accrual of CNAs
within the cancers.
239
Mathematical modelling negative selection on CNAs
We hypothesised that the distinct pattern of CNAs observed in each CRC represented a (local)
optimum in the fitness landscape, with negative (purifying) selection suppressing diversity at gland
and bulk levels. To test this hypothesis, we developed a mathematical model of CNA accrual during
cancer growth and the evolutionary dynamics due to the action of negative selection on the CNAs.
Individuals in the model ('glands') had variable fitness ($s = \frac{b_i - d_i}{b_o - d_o} - 1$) defined by the gland's
karyotype according to the relation:
247
248
$$b_i - d_i = \frac{b_o - d_o}{1 + \alpha * d(G_i, G_o)},$$
249
where b_i (d_i) was the birth (death) rate of gland i, G_i was the karyotype of gland i, G_o was a pre-
specified 'optimal' karyotype that maximised fitness, b_o (d_o) was the "optimal" birth (death) rate,
and the function $d(G_i, G_o)$ measured the genetic distance between the two karyotypes. A gland's
karyotype was represented by a vector of length C, where C was the number of bins used in our
empirical data analysis, with each entry representing the integer copy number of each locus in the
gland's (abstracted) genome. The number of CNAs followed Poisson distribution at mutation rate μ
per gland division, where the size of each CNA (in the unit of bins) was sampled from the sizes of
observed CNAs in our data. A gland's fitness was then inversely proportional to the absolute
distance between a gland's karyotype and the maximum fitness karyotype (G_o); the parameter α
scaled the strength of stabilising selection experienced by non-optimal karyotypes, with $\alpha = 0$
defining a "flat" landscape where all karyotypes have the same fitness, $s = s_{max}$ (Fig 2B).
261
Stochastic simulations showed that ongoing mutation within a growing tumour in the absence of
negative selection ($\alpha = 0$) generated high CNA heterogeneity (Fig 2C&D). A growing tumour that
initially occupied the fitness maxima (karyotype of the first cancer gland = G_o) experienced negative
selection which suppressed CNA diversity, with stronger negative selection (greater α) more
severely limiting diversity (Fig 2C&D), and higher mutation rates generating more diversity (Fig 2D).
Thus, modelling suggested that negative selection for an optimal CNA karyotype was a plausible
explanation of the apparent stability of the aneuploid genome observed in CRC glands.
269
Our previous single cell sequencing data, derived from cells extracted from colorectal glands from
the same tumours analysed in this study[24], provided an empirical measurement of the lower limit of
the CNA rate at $\mu \approx 0.1$ new CNAs per gland division. The dynamics of cells within glands affect the
scaling of per-cell to per-gland mutation rates in a complex fashion dependent on the mechanism of
selection upon cell/gland karyotypes; for simplicity we neglected to consider these dynamics and

took prior distributions that set per-gland mutation rates the same as cellular rates. We used
Bayesian model selection to compare the fit of models with negative selection versus no selection
(purely neutral evolution) on the 6 cancers subjected to gland-by-gland analysis. In 4/6 cancers the
model with stabilising selection better represented the data than the model without stabilising
selection (Fig 2A), and it was not possible to distinguish between the models in 1/6 cases. The model
rests on a number of assumptions, and the goodness of these assumptions (see Discussion)
determines the strength of the conclusion that can be drawn from these analyses. Nevertheless,
modelling provided some quantitative evidence that the low levels of CNAs diversity in primary CRCs
was frequently due to the acting of stabilising selection upon a core karyotype.
284
Ongoing CNA accrural in CRC model systems
Our model of negative selection presumes that CNA events were constantly generated during
tumour development and were purged from the population by negative selection, leading to
minimal CNA heterogeneity. An alternative hypothesis is that CNA alteration rates are very low most
of the time, possibly with occasional bursts of high CNA rates that produced the observed highly-
altered genomes. We note this is not necessarily the same as *punctuated evolution*, which does not
require bursts of mutation rates because phenotypes develop in peripherally isolated populations in
separated environmental niches[26]. High ongoing CNA rates in cancer and specifically in colorectal
malignancies, rather than a burst of CAN events, has been demonstrated previously[24], and was
supported by single cell assays in our dataset (Fig S6 and ref [24]).
295
We sought to provide further evidence of an ongoing high CNA rate in CRC. We generated single cell
DNA sequencing data from SW620 colon cancer cells (Fig S7A, Table S7). Using 91 high quality cells
(see Methods) we performed CNA calling and phylogenetic analyses. This highlighted that individual
cells had ongoing chromosomal instability because the cells had generated multiple new CNAs since
their MRCA, in addition to a number of shared (clonal) CNAs in the population (Fig 3A). The new
CNAs were spread across the genome, and phylogenetic analysis showed that cell-unique (tip) CNAs
were enriched for chromosome 11 losses. This may indicate that the copy number gain state (CN =
3) in chromosome 11 is unstable and frequently reversed in SW620 cells, or could indicate selection
for this loss in our cell culture conditions.
305
306 We next performed single cell DNA sequencing in a patient-derived colorectal cancer organoid[27],
307 which also showed ongoing CNA generation in individual cells that were spread across the entire
308 genome (Fig 3C-D, Fig S7B, Table S7).
309

310
Figure 3: Ongoing copy number alterations in model systems. *(A) Schematic of the method used to*
obtain and sequence single-cells from cultured colorectal cell-line SW620 and patient derived
organoid 3994-1117. Cell line SW620 underwent single-cell dispensing using the cellenONE system,
followed by direct library prep (DLP) while the 10x Genomics Chromium was used to prepare single
cell libraries from the patient derived organoid S994-1117. (B) Heatmap showing copy number
alterations (CNAs) in single-cells from colorectal cell-line SW620 and (C) from patient derived
organoid 3994-1117. Red=gain, blue=loss. (D) Phylogenetic tree constructed using CNAs in the CRC
cell line SW620. (E) Frequency of copy number gains and losses at the tips of the SW620 phylogenetic
tree as in 3G. (F) Phylogenetic tree constructed using CNAs in the CRC organoid 3994-117. (G)
Frequency of copy number gains and losses at the tips of the 3994-117 phylogenetic tree.
321
322

323 ***Negative selection on CNAs continues through metastatic dissemination and treatment***
324 We then investigated how CNAs evolved during metastatic spread and treatment. Colonisation of a
325 new tissue and the application of therapy represent new selective pressures that could change the
326 core karyotype observed in primary CRCs. Even in the absence of positive selection, bottlenecks
327 during metastatic colonisation combined with the strong effects of genetic drift in an initial small
328 population, could lead to gross differences between primary tumour and metastatic deposits (i.e.
329 founder effect). CRC frequently metastasises to the liver and less frequently to other organs,
330 including the lung. We collected tissue from 23 primary sites and 68 metastatic lesions from 22
331 patients, with the material collected longitudinally from a mode of 2 time-points (range 1-5 time
332 points). Five patients remained untreated throughout their studied time course (an additional three
333 patients received no treatment between sampled timepoints), while other patients had experienced
334 a diverse range of treatment regimens (Table S8). Using sWGS, we measured inferred absolute copy
335 number of CNAs for each tumour deposit (total of 168 distinct regions analysed) and called GD as
336 described above. Comparing metastatic lesions to our previous primary CRC cohort (Fig 1), we
337 observed that CNA diversity only fractionally increased in the metastatic cohort (Fig S8A). The
338 frequency of GD was significantly higher in metastatic cases (18/22 [82%] had GD) versus regular
339 CRCs (16/81 [20%] cases GD; p<0.0001; Fig S8B), and was comparable to previous reports of GD in
340 metastatic CRC[28]. GD was clonal in two-thirds of metastatic cases with GD (12/18; 66%).
341
342 We examined how CNAs evolved over space and time by comparing patterns of CNAs within and
343 between metastatic lesions and to the primary tumour. Additionally, we performed phylogenetic
344 analysis per patient in the cohort based on CNAs using MEDICC2[29] (Fig S9). We observed several
345 examples of the maintenance of a core abnormal karyotype in the primary tumour wherein the
346 pattern of CNAs did not greatly change in metastatic lesions through space, time, metastasis to
347 other tissues and through chemotherapy and/or targeted treatment (Fig 3A-D, Fig S10-12, Table S9).
348 In nearly all cases there were some CNA differences between lesions in the same patient, and we do
349 not discount that these differences may be important for tumour cell biology.
350
351 We measured the proportion of the aberrant genome that was subclonal in pairwise sample
352 comparisons within each patient, the mean of which we term divergence, in addition to counting
353 phylogenetic events. Patient LM0027 displayed little divergence in karyotype between the primary
354 tumour and a metastasis taken a year later in the liver, with no treatment in-between (Fig 3A;
355 divergence of 0.26 [0.18-0.31], mean absolute PGA difference 5.4 percentage points, mean number
356 of phylogenetic events 11 [4-17]). In patient LM0071, despite 23 months between the primary colon
357 and metastatic liver samples 3-months of oxaliplatin and capecitabine chemotherapy, maintenance
358 of a core karyotype was observed (Fig 3A; divergence 0.17 [0.03-0.22], mean absolute PGA
359 difference 1.6pp, mean number of events 5.1 [2-7]). LM0029 was followed for 28 months from
360 primary diagnosis, through hepatic metastatectomies, one pulmonary metastatectomy, and three
361 challenges of chemotherapy in combination with anti-epidermal growth factor monoclonal antibody
362 cetuximab. Again, the core karyotype was broadly maintained throughout, with the largest
363 difference seen in the lung metastases which lacked a gain of chromosome 2 (Fig 4A, divergence
364 between consecutive sampling points of 0.19 [0.13-0.25], 0.34 [0.19-0.60], 0.26 [0.12-0.52], 0.38
365 respectively, mean absolute PGA differences 4.6pp, 6.1pp, 6.2pp and 18pp respectively, mean
366 number of events, 33 [30-36], 27 [14-38], 25 [13-40], 25 [25-25], respectively). Similarly, in patient
367 LM0139, resection of the primary CRC at baseline was followed by hepatic metastatectomy at 23
368 months and pulmonary metastatectomy at 72 months. Despite 6 years of elapsed evolution and
369 metastasis to two separate organs, the final karyotype sampled in the lung had a very similar pattern
370 of CNAs to the initial karyotype observed in the primary, demonstrating marked stability across time
371 and space (divergence between first and last timepoint 0.28 [0.18-0.38], mean absolute PGA
372 difference 12pp, mean number of events 36 [35-36]; Fig S13A).

373
Figure 4: Stability of CNAs across space, time, tissue and treatment in colorectal cancer. (A)
Longitudinal assessment of CNAs in primary colorectal cancers and metachronous metastatic lesions
in liver and lung showed similar CNA profiles were observed over space time, tissue and through
various treatment courses. Timelines of four representative metastatic patients are shown.
Schematics show organ where cancer deposit was sampled. Genome-wide view of CNAs with gains
(red) and losses (blue) relative to baseline ploidy. Proportion of Aberrant Genome Subclonal (PoAGS)
and number of events are displayed for each comparison presented sequentially. To the right of each
timeline is the corresponding phylogenetic tree as described in Fig S9. (B) Mean percent genome
altered (PGA) averaged across all lesions at the sampling time by time since baseline (removal of the
primary CRC). The PGA remained broadly constant over time in each patient (x-axis is days and is on
a log scale). (C) The pairwise difference in CNAs between regions of the primary tumours was not
significantly different to the pairwise difference between regions of paired metastatic lesions. (D)
Across timepoint genetic divergence was not significantly different after treatment with various
chemotherapy and/or targeted agents. (E) GD was common in metastatic primary tumours and
metastatic deposits and was sometimes present in some but not all of the metastatic lesions from a
patient (called subclonal GD). (F) An example case (LM0032) where GD was subclonal in the primary
but clonal in the metastasis. The greater burden of CNAs is evident in the GD samples. (G) Frequency
of copy number gains (red) and losses (losses) present on the clonal branches in the set of
phylogenetic trees created from the cohort. Over-representation of some CNAs is evident. (H)

Frequency of copy number gains and losses at the tips of the phylogenetic trees (i.e. unique events).
CNAs are uniformly spread across the genome.
395
Despite these examples of relative stability, we observed some cases where CNA patterns diverged
markedly between lesions. For example, the metastatic lesions in the ovary and omentum sampled
at the second timepoint in LM0018, taken only 7 months after the sampling of the primary, showed
the highest between timepoint divergence in the cohort (divergence 0.62 [0.55-0.7], mean absolute
PGA difference 27pp, mean number of events 42 [32-52], Fig S10). The patient received FOLFOX in
between sampling points. These samples were our only examples of ovary and omentum spread but
raise the possibility that CNAs may change substantially to enable colonisation of these organs. We
also highlight case LM0011 which also showed a large increase in divergence between the two
timepoints in the colon and liver taken 16 months apart despite receiving no treatment in the
interval (divergence 0.53 [0.49-0.62], mean absolute PGA difference 21pp, mean number of events
52 [40-67], Fig S10).
407
Alterations in the proportion of genome altered (PGA) were generally low in sequential timepoints
indicating aneuploidy stability (mean PGA increased by 3.4pp on average for consecutive timepoints
in the cohort; Fig 4B). Within the same time point, CNA differences between metastatic regions was
not significantly different from CNA differences between primary regions (Fig 4C; p=0.2, linear mixed
model, Fig S13B) and divergence within regions of primary colorectal samples was consistent with
our analysis of regular CRCs (divergence 0.24). Overall, CNA divergence was not significantly
different in on-treatment intervals versus off-treatment intervals (p=0.19, linear mixed model, Fig
3D).
416
Subclonal GD was inferred in six patients (LM0018, LM0019, LM0029, LM0032, LM0056, LM0124)
and in these cases GD was enriched in metastatic samples versus primaries (Fig 4E). For example, the
metastatic samples in LM0032 were GD whereas one of the two primary samples was not,
suggesting GD was subclonal at the primary site and potentially led to the metastatic clone (Fig 4F).
In LM0056 and LM0124 all primary samples were classified as non-GD, yet all bar one metastasis
sample in each patient was considered GD, indicating the GD likely originated after metastasis. Only
a single liver metastasis region was called as GD in LM0019. In primary CRCs, GD tumours had less
diverse CNAs. The reason for this difference could be related to elapsed time: in GD primaries could
be fit and grow rapidly with little elapsed time for new CNA evolution despite an intrinsically higher
mutation rate, whereas in metastasis sufficient time elapses for new CNAs to accrue.
427
Based on the phylogenetic structure inferred for all 22 patients, we separated CNA events into three
categories: clonal (events preceding the MRCA of the tumour samples), intermediate (events shared
between samples but not preceding the MRCA) and tip (events unique to each sample). Clonal CNAs
had occurred early in the cancer's development, and so selection had had time to operate on these
events. In comparison, tip CNAs had occurred more recently and so may not yet have been assorted
by selection. The distribution of clonal events across the genome showed some CNAs were far more
common than others consistent with the typical pattern for CRC[30]. Events such as 8q, 13 and 20 gain,
as well as 8p and 17p loss were enriched only in clonal alterations (Fig 4G). Contrastingly, the events
at the tips of the phylogenetic trees (Fig 4H) showed a strikingly uniform distribution across the
genome suggesting an ongoing accrual of CNAs at a constant rate across the genome, prior to
assortment by selection.
439
440
Discussion
Our study reveals the dynamics of CNA during the evolution of colorectal cancer. We observe large
changes in CNAs during the progression from benign to malignant disease, but thereafter the cancer

karyotype appears quite stable, and typically remains relatively unchanged over many years,
through metastasis to distant organs, and despite treatment. Taken together, these data suggested
a model whereby strong positive selection for an abnormal karyotype occurs during transformation
from benign to malignant disease, and that thereafter the karyotype evolution is constrained by
negative selection acting at the level of individual cells including in metastasis and during treatment.
This is a distinct mechanism to transient bursts in the CNA rate as has been previously suggested to
occur in breast cancer evolution[31]. We note that our finding of pervasive negative selection at CNA
level is congruent with our previous observation of "Big Bang" effectively-neutral evolution in
CRCs[20,21], because negative selection acts to remove deleterious variants before they can expand to
detectable levels and so only neutrally evolving lineages remain. Positively selected variants, both
SNVs and CNAs, are clonal in the tumour and part of the 'trunk' of the phylogenetic tree but do not
drive subclone expansions.
456
We propose an altered "core karyotype" is necessary for the initiation of malignancy, and that the
same karyotype is also sufficient for subsequent metastatic spread. These evolutionary dynamics are
consistent with the observation of early metastasis in CRC[32], whereby the genome of the initial
primary tumour is competent for metastasis. Given the inter-tumour heterogeneity in patterns of
CNAs, we suggest that the fitness landscape traversed by CNAs is 'rugged', with local fitness peaks
located at normal-diploid and at multiple aberrant-genome positions, and that fitness of any
individual karyotype is likely to be somewhat patient-specific due to the background of germline and
somatic variants and tumour microenvironment composition. There could be multiple local maxima
close together in genotype space – or similarly new CNAs could be deleterious or neutral – our
analysis is insensitive to detect these dynamics. Our data do suggest that fitness peaks are likely
sufficiently distant in karyotype space to prevent easy transition between peaks, and potentially that
the selective landscape of CNAs differs between benign and malignant disease.
469
Given the potentially defining role of CNAs in CRC evolution, we hypothesise that the fixed pattern
of CNAs in any individual CRC likely defines the biology of that tumour, and consequently is a major
determinant of patient prognosis and treatment response. That the karyotype is relatively fixed
through treatment is surprising, as presumably the effectiveness of therapy is dependent on it acting
as a novel and strong selective pressure. We had expected this new selective pressure to change the
fitness landscape and as a result cause wholesale change in the CNAs observed post-therapy. It may
be that the fitness peak of maximal treatment-resistance is too far from that occupied by an
untreated tumour and so is highly unlikely to evolve. However, we emphasise that our analysis has
taken a genome-wide average view, and adaption could be driven by individual and/or small
CNAs[8,33] for which our analysis is insensitive to detect. Nevertheless, an alternative explanation is
that chemotherapy response in CRC may be entirely plastic[34], and so does not involve the clonal
selection of individual lineages – a phenomenon that may be common across cancer types[35]. In CRC,
previous work is suggestive that plasticity is established very early in CRC[35] and is consistent with
reports of early metastatic spread[32].
484
We reiterate that there are typically clear genetic differences between matched primaries and mets,
and we do not rule out that these differences could contribute importantly to tumour biology: we
recognise that our analysis is insensitive to detect a driving role for such events. We acknowledge
that our study is limited by its principal reliance upon bulk sampling, including our use of individual
tumour glands, and stronger evidence for or against negative selection on CNAs will derive from
large-scale single cell sequencing analyses. Our modelling relies on a number of critical but untested
assumptions, such as simple scaling of mutation rates between tumour cells and glands, steady
division rates of disseminated tumour cells, uncertain estimates of the age of metastases, and
monoclonal origins of metastases. All these assumptions can alter the expectation of the degree of

genetic divergence between and within lesions. Appropriately addressing these assumptions is a
major undertaking and should be a priority for future work.
496
497 Collectively, out study demonstrates a plausible role for negative selection acting on the karyotype
498 to define the genomes of colorectal cancers.
499

Methods
501 Full methods are provided as supplementary material.
502

Acknowledgments
MJ, AS and TG acknowledge funding from Cancer Research UK (A22745, A22909 and A19771
respectively). TG, AS and SL are funded by the Wellcome Trust (202778/Z/16/Z, 202778/B/16/Z and
206314/Z/17/Z respectively) and via a Wellcome Trust award to the Centre for Evolution and Cancer
at the ICR (105104/Z/14/Z). SL and LMW acknowledge funding from the National Institute for Health
Research (NIHR) Oxford Biomedical Research Centre. DS, AS and TG received funding for this project
from the NIH via the Cancer Systems Biology Consortium U54 scheme (CA217376). TG received pilot
funding from the Barts Charity. AB was supported by Barts Health NHS Trust and Barts Charity grant.
ED is supported by the S:CORT consortium which is funded by a grant from the Medical Research
Council and Cancer Research UK. JB is in part funded by the UCLH/UCL Biomedical Research Centre.
MJ, MN and MRJ receive funding support from Cancer Research UK UCL Experimental Cancer
Medicine Centre and the Department of Health's National Institute for Health Research Biomedical
Research Centres funding scheme (Reference: C12125/ A15576). The research leading to these
results has received funding from the People Programme (Marie Curie Actions) of the European
Union's Seventh Framework Programme (FP7/2007-2013) under REA grant agreement n° 608765
(MM). This research was supported by the National Institute for Health Research (NIHR) Biomedical
Research Centre based at Guy's and St Thomas' NHS Foundation Trust and King's College London
who performed some of the genome sequencing, and we note that the views expressed are those of
the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health. We are
grateful for the support of the QMUL Genome Centre (Charles Mein) and Institute of Cancer
Research Tumour Profiling Unit for support with genomic assays. A small project grant from the UK
Bowel and Cancer Research (BACR) Charity to TG and SL pump primed this research. Organ likeness
in Fig 3A, 3F and Fig S6, S10-12 were based on images in Servier Medical Art (licensed under a
Creative Commons Attribution 3.0 Unported License).

Conflicts of interest statement
529 TG and AMB are named as coinventors on a patent application that describe a method for TCR
530 sequencing (GB2305655.9), and TG is named on a method to measure evolutionary dynamics in
531 cancers using DNA methylation (GB2317139.0). TG has received an honorarium from Genentech.
532

References
1.   Taylor, A. M. *et al.* Genomic and Functional Approaches to Understanding Cancer
Aneuploidy. *Cancer Cell* **33**, 676-689.e3 (2018).
2.   Sansregret, L., Vanhaesebroeck, B. & Swanton, C. Determinants and clinical
implications of chromosomal instability in cancer. *Nat. Rev. Clin. Oncol.* **15**, 139–150
(2018).
3.   Muzny, D. M. *et al.* Comprehensive molecular characterization of human colon and
rectal cancer. *Nature* **487**, 330–337 (2012).
4.   Watkins, T. B. K. *et al.* Pervasive chromosomal instability and karyotype order in
tumour evolution. *Nature* **587**, 126–132 (2020).
5.   Mermel, C. H. *et al.* GISTIC2.0 facilitates sensitive and confident localization of the

targets of focal somatic copy-number alteration in human cancers. *Genome Biol.* **12**,
R41 (2011).

6. Li, X. *et al.* Assessment of Esophageal Adenocarcinoma Risk Using Somatic
547 Chromosome Alterations in Longitudinal Samples in Barrett's Esophagus. *Cancer*
548 *Prev. Res. (Phila).* **8**, 845–856 (2015).

7. Rubin, C. E. *et al.* DNA aneuploidy in colonic biopsies predicts future development of
550 dysplasia in ulcerative colitis. *Gastroenterology* **103**, 1611–1620 (1992).

8. Smith, J. C. & Sheltzer, J. M. Systematic identification of mutations and copy number
552 alterations associated with cancer patient prognosis. *Elife* **7**, (2018).

9. Hieronymus, H. *et al.* Tumor copy number alteration burden is a pan-cancer
554 prognostic factor associated with recurrence and death. *Elife* **7**, (2018).

10. Andor, N. *et al.* Pan-cancer analysis of the extent and consequences of intratumor
556 heterogeneity. *Nat. Med.* **22**, 105–113 (2016).

11. Jamal-Hanjani, M. *et al.* Tracking the Evolution of Non-Small-Cell Lung Cancer. *N.*
558 *Engl. J. Med.* **376**, 2109–2121 (2017).

12. Sheltzer, J. M. & Amon, A. The aneuploidy paradox: costs and benefits of an incorrect
560 karyotype. *Trends Genet.* **27**, 446–453 (2011).

13. Li, L. *et al.* Cancer-causing karyotypes: chromosomal equilibria between destabilizing
562 aneuploidy and stabilizing selection for oncogenic function. *Cancer Genet. Cytogenet.*
563 **188**, 1–25 (2009).

14. López, S. *et al.* Interplay between whole-genome doubling and the accumulation of
565 deleterious alterations in cancer evolution. *Nat. Genet.* **52**, 283–293 (2020).

15. Kimura, M. Evolutionary Rate at the Molecular Level. *Nature* **217**, 624–626 (1968).

16. Birkbak, N. J. *et al.* Paradoxical relationship between chromosomal instability and
568 survival outcome in cancer. *Cancer Res.* **71**, 3447–3452 (2011).

17. Schwartz, R. & Schäffer, A. A. The evolution of tumour phylogenetics: principles and
570 practice. *Nat. Rev. Genet.* **18**, 213–229 (2017).

18. Cross, W. *et al.* The evolutionary landscape of colorectal tumorigenesis. *Nat. Ecol.*
572 *Evol.* **2**, 1661–1672 (2018).

19. Maley, C. C. *et al.* Genetic clonal diversity predicts progression to esophageal
574 adenocarcinoma. *Nat. Genet.* **38**, 468–473 (2006).

20. Sottoriva, A. *et al.* A Big Bang model of human colorectal tumor growth. *Nat. Genet.*
576 **47**, 209–216 (2015).

21. Williams, M. J. *et al.* Quantification of subclonal selection in cancer from bulk
578 sequencing data. *Nat. Genet.* **50**, 895–903 (2018).

22. Lengauer, C., Kinzler, K. W. & Vogelstein, B. Genetic instability in colorectal cancers.
580 *Nature* **386**, 623–627 (1997).

23. Siegmund, K. D., Marjoram, P., Woo, Y.-J., Tavaré, S. & Shibata, D. Inferring clonal
582 expansion and cancer stem cell dynamics from DNA methylation patterns in
583 colorectal cancers. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 4828–4833 (2009).

24. Bollen, Y. *et al.* Reconstructing single-cell karyotype alterations in colorectal cancer
585 identifies punctuated and gradual diversification patterns. *Nat. Genet.* **53**, 1187–
586 1195 (2021).

25. Alexandrov, L. B. *et al.* Clock-like mutational processes in human somatic cells. *Nat.*
588 *Genet.* **47**, 1402–1407 (2015).

26. Eldredge, N. & Gould, S. Punctuated Equilibria: An Alternative to Phyletic Gradualism.
in *Models in Paleobiology* **82**, 82–115 (1971).

27.   Vlachogiannis, G. *et al.* Patient-derived organoids model treatment response of
metastatic gastrointestinal cancers. *Science* **359**, 920–926 (2018).

28.   Priestley, P. *et al.* Pan-cancer whole-genome analyses of metastatic solid tumours.
Nature **575**, 210–216 (2019).

29.   Kaufmann, T. L. *et al.* MEDICC2: whole-genome doubling aware copy-number
phylogenies for cancer evolution. *Genome Biol.* **23**, 241 (2022).

30.   Cornish, A. J. *et al.* Whole genome sequencing of 2,023 colorectal cancers reveals
mutational landscapes, new driver genes and immune interactions. *bioRxiv*
2022.11.16.515599 (2022). doi:10.1101/2022.11.16.515599

31.   Gao, R. *et al.* Punctuated copy number evolution and clonal stasis in triple-negative
breast cancer. *Nat Genet* **48**, 1119–1130 (2016).

32.   Hu, Z. *et al.* Quantitative evidence for early metastatic seeding in colorectal cancer.
Nat. Genet. **51**, 1113–1122 (2019).

33.   Salehi, S. *et al.* Clonal fitness inferred from time-series modelling of single-cell cancer
genomes. *Nature* **595**, 585–590 (2021).

34.   Kreso, A. *et al.* Variable clonal repopulation dynamics influence chemotherapy
response in colorectal cancer. *Science* **339**, 543–548 (2013).

35.   Boumahdi, S. & de Sauvage, F. J. The great escape: tumour cell plasticity in resistance
to targeted therapy. *Nat. Rev. Drug Discov.* **19**, 39–56 (2020).

610