1   Title page

4   **Attentional Focus Modulates Automatic Finger-tapping**

5   **Movements**

Xilei. Zhang [a, b, *], Xingxun. Jiang [a, b], Xiangyong. Yuan [c], Wenming. Zheng [a, b, *]

[a] School of Biological Sciences & Medical Engineering, Southeast University, China

[b] Key Laboratory of Child Development and Learning Science, Ministry of Education, China

[c] Institute of Psychology, Chinese Academy of Sciences, China

[*] To whom correspondence should be addressed. Email: xilei.zhang@seu.edu.cn or

wenming_zheng@seu.edu.cn.

The authors declare no competing interests.

Abstract

The majority of human behaviors are composed of automatic movements (e.g., walking or finger-tapping) which are learned during nurturing and can be performed simultaneously without interfering with other tasks. One critical and yet to be examined assumption is that the attention system has the innate capacity to modulate automatic movements. The present study tests this assumption. Setting no deliberate goals for movement, we required sixteen participants to perform personalized and well-practiced finger-tapping movements in three experiments while focusing their attention on either different component fingers or away from movements. Using cutting-edge pose estimation techniques to quantify tapping trajectory, we showed that attention to movement can disrupt movement automaticity, as indicated by decreased inter-finger and inter-trial temporal coherence; facilitate the attended and inhibit the unattended movements in terms of tapping amplitude; and re-organize the action sequence into distinctive patterns according to the focus of attention. These findings demonstrate compelling evidence that attention can modulate automatic movements and provide an empirical foundation for theories based on such modulation in controlling human behavior.

Keywords: attention, automatic movements, finger tapping, DeepLabCut, constrained action hypothesis, Norman-Shallice theory

Introduction

The majority of human behaviors are composed of automatic actions [1-3]. Other than reflexive actions (e.g., the eye-blink reflex evoked by a startling sound), which are natural and can severely interrupt the ongoing task, automatic movements in this study are operationally defined as movements learned during nurturing and can simultaneously be performed without interfering with other tasks [4-7]. Requiring little attentional control, these automatic actions offer the advantage of allowing us to devote our limited attentional resources to other important matters. For example, we can drink a cup of coffee while immersed in deep thinking without worrying about the muscle contractions and relaxations used to execute these movements. However, this does not mean that automatic movements are disconnected from our attention. On the contrary, attention has long been conceptualized as a system of great significance that modulates automatic movements [5,8,9].

On the one hand, it has been proposed that focusing attention on body movements can disrupt movement automaticity and thus deteriorate the performance of well-practiced skills (i.e., constrained action hypothesis [10,11]). Taking a vertical jump-and-reach task as an example, focusing attention at the fingertips versus the rungs to be touched leads to decreased maximum vertical jump height [12]. However, it should be noted that these studies involved a hidden common confounding factor. Although these actions are generally deemed skillful and automatic, most are guided by deliberate goals. For example, participants were required to jump as high as possible [12], swim as fast as possible [13], or throw a dart as accurately as possible [14]. With specific goals to fulfill, it is natural to assume that participants would deliberately control certain aspects of their movements (e.g., force, timing, speed, coordination). As a result,

3

such studies are incapable of elucidating the complex relationship between attention and

automatic movement, given it is impossible to conclude that such actions are not entirely

voluntary.

On the other hand, attention has long been conceptualized as the mediator in translating

deliberate goals (e.g., drinking coffee) into specific body (muscle) movements (i.e., Normal-

Shallice theory [5]). As one of the core proposals, attention is given an innate potential to

modulate automatic movements by facilitating attended actions while inhibiting unattended

actions [5]. To our surprise, this proposal has never been empirically tested (see Discussion).

Recently, one neuroimaging study [15] found that evidence indicating the interaction between

attention and automatic movements is absent for healthy participants (but present for a group

of Parkinson's disease patients). For this study, the authors trained a visually cued keypress

movement to automaticity at first (see [16,17] for debates on this paradigm in training movement

automaticity) and examined neural dynamic changes when attention was directed back to

movement. In the healthy control group (as compared to Parkinson's disease patients), however,

data revealed that attention modulates activity and connectivity in those regions responsible for

goal-directed, controlled movements (the dorsolateral prefrontal cortex, anterior cingulate

cortex, and rostral supplementary motor area) [15,18], but not in the region responsible for well-

learned, automatic movements (the striatum) [7]. Though worthy of further scrutinization using

more sensitive analyses (e.g., pattern analysis), the non-significant effect of attention on neural

activities in the striatum poses a challenge for determining whether attention truly modulates

automatic movements of healthy people free of movement disorders.

The present study re-investigates the interaction between attention and automatic

movements. Crucially, we did not ask participants to perform automatic movements under any

specific goals, given the aforementioned issue of previous studies. Moreover, the automatic

movement chosen for this study was not one newly learned in the lab. During both the entrance

exam and the main task (see Methods section), participants were required to repetitively and

sequentially tap their fingers at a usual, comfortable pace. Even when attention was directed

toward these movements, the lack of a specific goal denied the need for participants to try and

control their tapping movements. To quantify the trajectories of tapping fingers, we used a new

marker-less video-based pose estimation technique [19]. In the present study, we manipulated

attention by asking participants to focus their attention on the movements of either all fingers

or a single finger (the index or middle finger), and contrasted these movement-focused

conditions with a reference condition in which attention was directed away from the tapping

fingers. Given everyday movements are hard to quantify in the laboratory due to techniques

used [20], evaluation of the direct relationship between attention and automatic movements is a

blank to be filled. In the present study, we aimed to fill this blank for the first time, based on

methodological improvements introduced above.

To fully elucidate how attention modulates automatic finger-tapping movements, we

examined the following three predictions. First, we tested the core proposition of the

constrained action hypothesis [10,11] that attention disrupts movements' automaticity. For a

sequence rhythmically tapped in perfect automaticity, we assumed that trajectory of finger

movements, as well as phase lag between fingers, would be duplicated cycle by cycle, leading

to ceiling inter-finger/inter-trial coherence. However, when attention is directed to movement,

we predicted a decrease in temporal coherence relative to the reference condition when

attention was directed away from movements. Secondly, using tapping amplitude as a measure,

we tested a core proposition from the Norman-Shallice theory. This theory, focusing on how

attention translates goals into behavior, proposes that attention modulates automatic

movements by facilitating attended actions while inhibiting unattended actions. Our primary

interest was whether the tapping amplitude would increase for attended fingers while

decreasing for unattended fingers. Finally, we performed a pattern analysis on the tapping

trajectories of all fingers to examine the prediction that attention focused at different fingers

would create distinct patterns of finger-tapping movement. The above predictions were tested

in three experiments. Experiments 1 and 2 directed participants' attention to the respective

fingers using two different paradigms, and we expected similar findings in both. For the control,

experiment 3 directed attention toward the conceptual label, rather than movement, of

respective fingers, and we expected no modulating effect of attention.

Methods

Participants

Sixteen college students (7 females) with a mean age of 24.44 ± 0.48 years [hereafter

the value following the symbol '±' represents one standard error of the mean (SEM)] took part

in experiments 1, 2, and 3 successively over three days spanning eight weeks. The sample size

was determined according to previous publications [21-23]. Procedures and protocols for this study

adhered to tenets of the Declaration of Helsinki and were approved by the review board of the

School of Biomedical Sciences and Engineering, Southeast University, China. All participants

were naive to the purpose of the experiments, gave prior informed written consent, and were

debriefed after each experiment.

Materials

In each experiment, participants were asked to tap their right-hand fingers on a pad

fixed to the table and to rest their thumbs at an anchor position marked by a small rubber patch

on the pad. Participants' finger-tapping behaviors were recorded by a full HD-camera (TC-

UV8000, 20X-zoom, resolution = 1920×1080, sampling rate = 60 Hz) facing the right hand

and placed on the table at a distance of 30 cm from the anchor. When participants were

informed to tap their fingers using a pure tone, a visual cue was simultaneously presented on

the screen in an isolated area. This area was shielded from the sight of participants but

continuously monitored by another HD-camera (TC-980S, 12X-zoom, 1920×1080, 60 Hz). We

used a video broadcasting station (model: iRBS-V8, with two video engines) to synchronize

videos from these two cameras so we could determine from which frame participants were

informed to tap their fingers.

Procedure

Stimuli were presented using Psychophysics Toolbox extension [24] with MATLAB (The

MathWorks, Natick, MA). Upon arrival at the laboratory, participants received an entrance

exam in which they had to concurrently perform two tasks. In one task of the exam, we required

them to tap the fingers (excepting thumbs) of their right hand at their own tempo, ensuring

familiarity and comfort with the movements as much as possible. In a simultaneous secondary

task, we asked participants to read a text paragraph clearly and loudly. Following prior literature

[6], we surmised that if one could tap his/her fingers fluently without being interrupted by the

secondary task, he/she was regarded as qualified for the subsequent experiment. In fact,

7

144   participants had no difficulty in passing the exam after some practice.

Figure 1. Schematic illustration of the experimental paradigm of (a) experiment 1, (b)

experiment 2, and (c) experiment 3. In experiments (a) 1 and (c) 3, a secondary letter-counting

or label-counting task was performed simultaneously with the finger-tapping task. The focus

of attention was manipulated in each experiment so that either different tapping fingers

(indicated by the pink shade on hand) were attended (experiments (a) 1 and (b) 2), or different

labels of particular fingers were attended (experiment (c) 3).

In experiment 1 (Fig. 1a), we examined whether and how these automatic movements

are influenced by the focus of directed attention. In a dual-task paradigm used to check for

movement automaticity [25,26], participants were required to perform the finger-tapping task

simultaneously with a secondary, more exhausting task (i.e., a letter-counting task). This

paradigm allows us to evaluate the validity of our manipulation of attentional focus [27,28] by

measuring the accuracy of the secondary task. Specifically, if participants focused their

attention on finger-tapping, the accuracy of the letter-counting task would decrease. The four

conditions in experiment 1 included three movement-focused conditions and a reference

condition. In the movement-focused condition, attention was directed at the movement of the

tapping sequence (sequence-focused condition), the index finger only (index-focused

condition), or the middle finger only (middle-focused condition). Under these conditions,

participants had to focus their best on movement of the required finger(s), while paying less

attention to the letter-counting task. In the reference condition, attention was focused on the

sequence of letters (letter-focused condition), and participants were required to count for the

appearance of a pre-defined target letter as accurately as possible while neglecting their finger-

tapping movements.

At the beginning of each trial, participants were informed of the experimental

conditions and target letter (O, G, L, or A, determined randomly). Once their attention was

given the required focus, participants pressed the enter key. An initiation beep (500 Hz, 500

ms) indicated participants were to start tapping their fingers until hearing the termination beep

(250 Hz, 500 ms). They were advised in advance that any deliberate control of movement was

forbidden. During this period, a random sequence of the letters O, G, L, and A were presented

at a frequency of 2.5 Hz (stimulus onset asynchrony: 400 ms). Each trial lasted either 6.71 sec

(short-duration trial, in which the target letter was presented 3–6 times) or 11.14 sec (long-

duration trial, in which the target letter was presented 5–9 times), in order to prevent

participants from being able to predict when the trial would terminate. At the end of each trial,

participants reported (by pressing keys using their left hands): (1) how many times they saw

the target letter, and (2) the amount of attention (ranging from 0 to 100) they had directed to

their finger-tapping movements. The next trial began after these responses. A total of 72 trials

were segmented into 12 blocks, each corresponding to one condition. In each block, there were

three short-duration trials and three long-duration trials. A rest followed every three blocks.

Experiment 2 (Fig. 1b) aimed to replicate the findings of experiment 1 in a single-task

paradigm, where participants were instructed to concentrate their efforts on finger-tapping with

no secondary task. In line with experiment 1, there were three movement-focused conditions

and one reference condition. However, differently and crucially, in the movement-focused

conditions, participants were not only told to attend towards, but to forcibly stare at, the

particular finger(s), continuously tracking their movements and neglecting the movements of

other fingers. In the reference condition, they were told to stare at their static thumbs. The target

finger was determined randomly for each trial, with no more than two repetitions. At the end

of each trial, participants reported the amount of attention paid to the target finger(s). To save

time and alleviate fatigue, the total trial number was reduced to 56, with 7 trials for each

duration (short and long) and attention condition. All other aspects were the same as in

experiment 1.

Experiment 3 (Fig. 1c) served as a control experiment to examine whether the possible

influence of attentional focus on automatic finger-tapping is derived from movement being

attended or from its conceptual label embedded in the mind. To this end, we used the dual-task

paradigm, as in experiment 1, but replaced the letter-counting task with a label-counting task,

in which participants were asked to neglect finger tapping but view a random sequence of finger

labels displayed on the screen and count for the target labels. The target label was the "thumb

finger" in the reference condition, and the "whole sequence", "index finger'," and "middle

finger" in three movement-related conditions. To direct attention away from their finger-

tapping movements, participants were instructed to look straight ahead at a screen and devote

all attention to the counting task. All labels were written in the participants' native language

(i.e., Chinese), and the target label was determined randomly for each trial. At the end of each

trial, participants reported: (1) the number of target labels, and (2) the amount of attention they

had directed to the movement of related finger(s). Other procedures were the same as those

used in experiment 1. There were 56 trials in total, with 7 trials for each duration and attention

condition.

Data Analysis

Accuracy of secondary counting task. For experiments 1 and 3, we calculated the

accuracy (ACC) of the secondary letter-counting task and label-counting task, respectively,

taking ACC as 1 if and only if the reported number of targets equalized with that of the actual

number of targets. Otherwise, we took ACC as 0. In the algorithm, missing and false alarms

had equal influence on accuracy.

Video-based estimation of tapping trajectories. Participants' finger-tapping

movements were extracted using a cutting-edge video-based pose estimation algorithm.

Specifically, by using DeepLabCut [19] running on a Linux (Ubuntu 16.04 LTS) operating system

with a graphics processing unit (GPU, NVIDIA TITAN XP, 12 GB of memory), we trained a

deep neural network with 50 layers (i.e., ResNet-50) to automatically recognize the coordinates

of seventeen key points of interest (Fig. 2a) for each frame during finger-tapping movements

(Fig. 2b). In the first step, we selected 25 representative frames for each participant from videos

recorded during experiment 1, manually marked coordinates of key points of interest in these

frames, and generated a total of 400 marked frames. We then randomly assigned 380 frames to

the training dataset and the remaining 20 frames as the test dataset. By using default parameter

settings, we trained a ResNet-50 model through 800,000 iterations. To evaluate its recognition

performance, we calculated the mean difference (in pixels) between the manually marked

coordinates and predicted coordinates. It was 3.19 ± 0.07 pixels for the training dataset and

4.65 ± 0.90 pixels for the test dataset, a negligible error considering there were $1500 \times 1080 =$

$1,620,000$ pixels within each frame. Thus, we applied this model to extract coordinates for each

frame, participant, and experiment, and only imported the vertical (y-axis) trajectories of

fingers into subsequent analyses, since fingers were tapped mainly in the vertical direction. For

preprocessing, vertical trajectories were segmented (from 30 frames before onset of the

initiation beep to 90 frames after onset of the termination beep), duration-normalized (the

duration of each trial was normalized to the mean duration of short- or long-duration trials

using function *interpft*.m), and temporally smoothed (sliding window = 6 frames, equaling 100

ms). Finally, outlier values that deviated the previous frame by a minimum distance of 100

pixels were replaced using the interpolation method (*inpaint_nans.m*). Relative to other

approaches quantifying finger movements, for example, those using retro-reflective markers [29],

force transductors [30,31] and cyber glove kinematic sensors [32], the marker-less video-based pose

estimation technique used in the present study provides an advantage of quantifying

movements in high precision while allowing participants to tap their fingers freely.

Defining the temporal and frequency ranges of tapping trajectory. The temporal

and frequency ranges for our interests were defined as those during which finger-tapping

behaviors were automatically performed without deliberate control. First, as the initiation and

termination of tapping behaviors inevitably require deliberate control [5], we selected the

temporal range from 1,000 ms post-onset of the initiation beep to the moment before the onset

of the termination beep for further analysis. There were 344 and 609 time points in total for

short-duration and long-duration trials, respectively. Second, as automatic finger-tapping

behaviors can be deconstructed as repetitive movements of each finger at a constant speed, the

frequency range of interest for each participant was defined as the frequency band, which

showed the strongest inter-trial temporal coherence. We therefore performed a wavelet

transform coherence analysis (WTC, http://www.glaciology.net/wavelet-coherence) for every

two trials (Fig. 2c) to produce a time-by-frequency 2-D coherence map (Fig. 2d). As a measure

of correlation between two time series [33], each coherence intensity out of the coherence map

ranges from 0 to 1, with 1 reflecting complete coherence (absolute phase synchrony) and 0

reflecting no coherence (no phase synchrony) at a given time-frequency point. The coherence

intensity was then averaged across the selected time range of interest, trial-durations, fingers,

and attention conditions. The mean coherence intensity can now be seen as a function of

frequency, among which the frequency with the strongest coherence intensity can be located

(*findpeaks.m*) and named as the tapping rate. The frequency band corresponding to full width

at half maximum (FWHM) centered at the tapping rate was determined as the frequency range

of interest.

Inter-finger and inter-trial coherence of tapping trajectory. We calculated the inter-

finger and inter-trial coherence of tapping trajectory to examine whether and how attention

modulates the temporal coupling of finger-tapping movements. For each experiment and

condition, mean inter-finger (or inter-trial) coherence was calculated by averaging the

coherence intensity within the temporal-frequency range of interest and across trials (or

fingertips).

Amplitude of tapping trajectory. We calculated the amplitude of tapping trajectories

for each fingertip to further test the hypothesis that attention increases tapping amplitude for

target fingers and decreases tapping amplitude for non-target fingers. The trajectory of each

fingertip was filtered into the frequency range of interest (*filter.m*) and analyzed by Hilbert

transform (*hilbert.m*). The absolute value of the resulting signal was averaged within the

temporal range of interest, yielding an estimation of the raw tapping amplitude for each

condition. For each experiment and fingertip, modulation index I_A was computed as $I_A =$

$(A_{attention}/A_{reference})-1$, where $A_{reference}$ is the amplitude of the reference condition and $A_{attention}$ is

the amplitude of each attention condition. In this way, the I_A of the reference condition for each

experiment was normalized to 0. Thus, an I_A above 0 indicates an excitation effect (i.e., increase

in amplitude), while an I_A below 0 indicates an inhibition effect (i.e., decrease in amplitude)

relative to the reference condition.

Pattern analysis of tapping trajectories. Supposing attentional focus indeed interferes

with automatic movements, different patterns of finger-tapping behaviors would be observed

when the focus of attention is changed. To test this hypothesis, a pattern analysis was performed

on the trajectory of all key points from the tapping fingers (a total of 14 key points). First, WTC

14

analysis was performed for each trial between every two key points, resulting in a trial × time

(time points in the temporal range of interest) × frequency (40 frequency points) × paired key

points ($C_{14}^2 = 91$) matrix for each participant. The resultant 4-D matrix was then concatenated

into a pooled time (trial × time points) × pooled frequency (frequency points × paired key

points) 2-D matrix. Second, to reduce dimensional space, we performed a principle component

analysis on the pooled frequency dimension, and only retained the top 80 components with the

highest variance, which together accounted for over 80% of the total variance. Consequently,

the pooled frequency dimension of the 2-D matrix was reduced from 40 × 91 to 80. Third, time

points from trials with the same condition were extracted in order and categorized into 6

independent and equivalent datasets. A sliding window method (length = 60 time points, step

size = 4 time points, with each step producing a sample containing 60 time points × 80

frequency components) was applied on the pooled time dimension, generating 1188 (or 928)

samples for each dataset in experiment 1 (or 2 and 3). Finally, we trained Support Vector

Machine (SVM) classifiers (radial basis function kernel, γ = 0.014, as default) for each

participant following a six-fold cross-validation approach. In each fold, five datasets were used

to train a classifier while the remaining was used for testing. After six folds, all datasets were

tested once. To evaluate the performance of the models, we calculated two indices. The

decoding accuracy was quantified as the ratio of correctly classified samples to the total number

of samples for each condition, while the mean decoding accuracy was calculated as the

decoding accuracy averaged across four conditions for each participant and experiment.

Beyond the above analyses on temporal coherence and tapping amplitude, analysis on the

decoding accuracy provided us the additional advantage to quantitatively evaluate whether and

to what extent the two conditions formed the same movement pattern, even if they had

comparable mean temporal coherence and/or tapping amplitude.

Statistics. Parametric tests used included the one-way and two-way repeated measures

analysis of variance (ANOVA), paired-sample t-test, one-sample t-test, and Pearson correlation

analysis. Unless specifically denoted, all statistical tests were conducted in a two-tailed manner

with a threshold of p = 0.05. All data analyses and statistics were performed using standard and

custom scripts adapted to MATLAB and Python.

Figure 2. Recording and quantification of automatic finger tapping movements. *(a)* One

exemplar frame illustrating the location of 17 key points of interest of the right-hand fingers.

For each frame, the coordinates of these points were automatically recognized using

DeepLabCut [19] . *(b)* Concrete frames illustrating changes in the position of key points during

repetitive finger tapping movements. For visibility, only key points on the fingertips are shown.

(c) Y-axis trajectories of the index finger from two exemplar trials. Dotted lines represent the

onset of pure tones informing participants to initiate and terminate finger tapping, respectively.

The solid line indicates the beginning of the temporal range of interest. *(d)* The 2-D map

produced by wavelet transform coherence (WTC) analysis between the above two trials. The

color bar indicates the coherence intensity, and the orientations of arrows indicate the circular

phase lag at each time-frequency point. The yellow rectangle indicates the temporal-frequency

range of interest determined for this representative participant. To exclude the distorting effect

of edge artifacts, values in the area masked by translucent white were not included in further

analyses. *(e)* Tapping rate is individually different but stable across experiments. Though

experiments were performed under different task demands and in separate days, the rate of

tapping movements between experiments was highly correlated (Pearson $r > 0.91$).

332   **Results**

333   **Constant and Fast Tapping Rates between Experiments**

334   As acknowledged, automatic movement operationally refers to those movements that

can be performed without interfering with other tasks [4,5]. From this point of view, if finger-

tapping movements were automatic, they would not interfere with the secondary task and

would remain stable across experiments, even if conducted on separate days or under different

task requirements. Given that each participant's finger-tapping behavior consisted of a

sequence of repetitive movements with which he/she was familiar, the tapping rate of each

participant was hypothesized to be stable across experiments. As expected, the tapping rates

from three experiments were highly correlated ($0.91 < rs < 0.94$, $ps < 0.001$, Fig. 2e). Moreover,

participants tapped their fingers at a speed as fast as $5.66\ HZ \pm 0.35\ HZ$ (ranging from $3.07\ HZ$

to $8.43\ HZ$). It seems impossible for participants to make deliberate plans and intentionally

control their movements at such a fast rate [34]. No participants reported attempting to control

their finger movements when tapping. This converging evidence supports that participants'

finger-tapping movements were automatic and under no intentional control.

Results of Experiment 1

In experiment 1, a one-way repeated measures ANOVA found a significant effect of

attentional focus on both letter-counting accuracy ($F_{3,45} = 35.30$, $p < 0.001$, $\eta_p^2 = 0.702$, Fig.

3a) and self-rated amount of attention to movement ($F_{3,45} = 204.11$, $p < 0.001$, $\eta_p^2 = 0.932$, Fig.

3b). Relative to the reference condition, post hoc tests revealed that directing attention to

movements led to an increase in self-rated amount of attention to movement (all $t_{15} > 14.95$, ps

< 0.001, Cohen's $d > 6.33$) and decrease in letter-counting accuracy (all $t_{15} < -6.06$, $ps < 0.001$,

Cohen's $d > 1.71$). Crucially, these two measures correlated significantly for all three

movement-focused conditions ($-0.73 < rs < -0.61$, $0.001 < ps < 0.006$, one-tailed, Fig. 3c).

These results provide compelling evidence that attention had been effectively directed toward

movement in experiment 1.

Figure 3. Objective and subjective measures of attention directed to finger-tapping movement

in experiment 1. *(a)* Objective measure of attention: accuracy (ACC) of the secondary letter-

counting task as a function of attentional focus. *(b)* Subjective measure of attention: self-rated

amount of attention directed to movement as a function of attentional focus. *(c)* More attention

directed to movement relative to the reference condition (letter-focused condition) led to less

attention available for the secondary task, thus possibly contributing to more decrease in ACC

of the secondary task. The error bar indicates one SEM.

One of the primary goals of experiment 1 was to examine the prediction derived from

the constrained action hypothesis that attention to movement may lead to impaired temporal

coherence between tapping trajectory across fingers and across trials. We thus performed two

one-way repeated measures ANOVAs on inter-finger coherence and inter-trial coherence,

respectively. As shown in Fig. 4a, results revealed a significant effect of attentional focus on

both types of coherence measures ($F_{3,45} = 7.56$, $p < 0.001$, $\eta_p^2 > 0.335$). Post hoc tests revealed

that temporal coherence was strongest in the reference condition (i.e., letter-focused condition;

inter-finger coherence: all $t_{15} > 1.96$, ps < 0.069; inter-trial coherence: all $t_{15} > 2.21$, ps < 0.043)

and weakest in the middle-focused condition (inter-finger coherence: all $t_{15} < -3.14$, ps < 0.007;

inter-trial coherence: all $t_{15} < -2.08$, ps < 0.055). There were no significant differences between

sequence- and index-focused conditions (inter-finger coherence: $t_{15} = 0.45$, p $= 0.661$; inter-

trial coherence: $t_{15} = 0.77$, p $= 0.453$). These results clearly indicate that focusing attention at

movement reduces the temporal coherence of automatic finger-tapping movements.

Figure 4. Temporal coherence of finger trajectories varying as a function of attentional focus.

Each panel indicates results with respect to inter-finger (shown in blue) and inter-trial (shown

in orange) temporal coherence, with x-axis labels indicating four attention conditions with

different attentional foci. The error bar indicates one SEM.

Another primary goal of experiment 1 was to examine the prediction derived from the

Norman-Shallice theory that attention would facilitate (inhibit) movement of attended

(unattended) actions. Following prior literature characterizing the function of the motor system

according to the amplitude of tapping movements [35], we tested this prediction taking tapping

amplitude as the dependent variable. We first examined for each tapping finger to determine

whether the I_A values in movement-focused conditions were significantly larger than zero using

a one-sample t-test, with zero indicating the normalized I_A value of the reference condition (i.e.,

letter-focused condition, see Methods section). For target fingers, expectedly, we revealed I_A

values significantly above 0 (increased tapping amplitude) for all target fingers in the three

movement-focused conditions (all $t_{15} > 2.34$, ps < 0.033, Fig. 5a). For non-target fingers,

however, only the I_A value of the little finger was found to have a < 0 tendency (attenuated

tapping amplitude, in the index-focused condition: $t_{15} = -1.89$, $p = 0.078$, in the middle-focused

condition: $t_{15} = -2.28$, $p = 0.038$). The I_A values of all other non-target fingers in the index-

focused and middle-focused conditions did not significantly differ from 0 (all $|t_{15}| < 1.44$, ps $>$

0.171). These results suggest that attention can facilitate the movement of target fingers and

possibly inhibit movement of (some) non-target fingers.

Figure 5. Tapping amplitude of finger trajectories varying as a function of attentional focus.

Each panel indicates results with respect to the amplitude for each tapping finger normalized

to the reference condition. X-axis labels indicate four attentional focus conditions, with the first

label indicating the reference condition whose amplitude had been normalized to 0. The error

bar indicates one SEM.

Next, we examined whether the I_A values in movement-focused conditions were

modulated by tapping fingers. A two-way repeated measures ANOVA on the I_A values revealed

that the main effects of condition ($F_{2,30} = 8.95$, $p < 0.001$, $\eta_p^2 = 0.374$) and tapping finger ($F_{3,45}$

$= 8.87$, $p < 0.001$, $\eta_p^2 = 0.372$) as well as the interaction ($F_{6,90} = 24.10$, $p < 0.001$, $\eta_p^2 = 0.616$)

were significant. To interpret this significant interaction, two follow-up statistical analyses

were performed. Firstly, we performed a one-way repeated measures ANOVA on I_A values

across the three movement-focused conditions for each tapping finger and found significant

differences (all $F_{2,30} > 6.08$, ps < 0.006, $\eta_p^2 > 0.289$). Post hoc tests found a larger tapping

amplitude when the index finger was attended than unattended (index-focused condition vs.

middle-focused condition: $t_{15} = 4.33$, p < 0.001; although not significant for sequence-focused

condition vs. middle-focused condition: $t_{15} = 1.41$, p = 0.180). This attention-induced

facilitation effect was replicated for the middle finger (sequence-focused and middle-focused

conditions vs. index-focused condition: both $t_{15} > 2.33$, ps < 0.034), and the ring and little

fingers (sequence-focused condition vs. index-focused and middle-focused conditions: all $t_{15} >$

3.26, ps < 0.005). These comparisons further support the effect of attentional focus on tapping

amplitude. Secondly, we performed one-way repeated measures ANOVA on the I_A values for

each movement-focused condition to assess differences in tapping amplitude between fingers,

revealing significant results (all $F_{3,45} > 4.43$, ps < 0.008, $\eta_p^2 > 0.228$). Post-hoc tests showed

the tapping amplitude of the little finger increased the greatest among target fingers when

attended in the sequence-focused condition (all $t_{15} > 2.81$, ps < 0.013) and decreased the

greatest among non-target fingers when unattended in the index-focused and middle-focused

conditions (all $t_{15} < -3.25$, ps < 0.005). These results suggest that tapping amplitude of the little

finger is most sensitive to the effect of attentional focus.

Figure 6. Distinct patterns of automatic movements varying as a function of attentional focus.

For each panel and cell, the value corresponds to the ratio of samples of the true condition,

which were classified as the predicted condition. Statistics were conducted against the chance

level (0.25) by using a one-sample t-test. Darker cells indicate values above chance while

whiter cells indicate values below chance. *, p < 0.05, ***, p < 0.001.

Finally, we performed a pattern analysis to examine whether attentional focus

determined the pattern of tapping movements, or, in other words, whether correctly recognizing

a particular mode of movement through pattern decoding significantly out-performed guessing

(25%). As expected, a one-sample t-test revealed that both the decoding accuracy for each

condition (i.e., correct classifications displayed in the diagonal direction of Fig. 6a, all $t_{15} >$

10.49, ps < 0.001) and the mean decoding accuracy across conditions (64.05% ± 1.87%, $t_{15} =$

20.91, p < 0.001, Cohen's *d* = 5.23, 95% confidence interval, or CI = [3.31, 17.14]) were

significantly above chance level. Furthermore, the percentage values in non-diagonal cells (i.e.,

incorrect classifications) were all significantly below chance level. These results indicate that

directing attention to different fingers creates distinct patterns of finger-tapping movements.

Results of Experiment 2

In experiment 2, we examined whether the effects found in experiment 1 could be

observed when participants' attention was directed toward the target fingers by looking directly

at them with no exhausting secondary task. Participants reported that most of their attention

(88.27% ± 2.16%) was directed to the target fingers, although they also stated their attention

was sometimes distracted by non-target fingers that moved simultaneously with the target

finger.

Following the same analyses as experiment 1, the results of experiment 2 first replicated

the effect that focusing attention at one's movement can desynchronize the temporal coherence

of automatic finger-tapping movements (Fig. 4b). Specifically, a one-way repeated measures

ANOVA found a significant effect of attentional focus on both inter-finger coherence ($F_{3,45}$ =

5.31, p < 0.003, η_p^2 = 0.261) and inter-trial coherence ($F_{3,45}$ = 3.72, p < 0.018, η_p^2 = 0.199).

Post hoc tests revealed that both inter-finger and inter-trial temporal coherence of the new

reference condition (i.e., thumb-focused condition) were larger than those of the index-focused

(t_{15} > 1.92, ps < 0.075) and middle-focused conditions (both t_{15} > 2.20, ps < 0.044), but were

comparable with those of the sequence-focused condition (both t_{15} < 0.44, ps > 0.668).

Consistent with experiment 1, the two indicators of temporal coherence were weakest in the

middle-focused condition (inter-finger coherence: all t_{15} < -2.46, ps < 0.027; inter-trial

coherence: all t_{15} < -2.05, ps < 0.059), and comparable between the sequence-focused and

index-focused conditions (both t_{15} < 1.61, p > 0.129).

With respect to the role of attentional focus on tapping amplitude, we once again

compared I_A values in the movement-focused conditions with zero, which represented the

normalized reference condition (i.e., thumb-focused condition). For the target fingers, as shown

in Fig. 5b, one-sample t-tests revealed increased tapping amplitude for the little and ring fingers

in the sequence-focused condition (t_{15} > 2.80, ps < 0.014; but for the index and middle fingers:

both t_{15} < 1.69, ps > 0.111), and the index finger in the index-focused condition (t_{15} = 2.34, p =

0.034), but not for the middle finger in the middle-focused condition (t_{15} = 1.17, p = 0.260).

For non-target fingers, tapping amplitude was attenuated for all non-target fingers in the index-

focused condition ($t_{15} < -3.30$, ps < 0.005) and the index and little fingers in the middle-focused

condition (both $t_{15} < -1.83$, ps < 0.088; but for the ring finger: $t_{15} = -1.31$, $p = 0.209$).

Collectively, these results suggest that attention can facilitate the movement of (some) target

fingers and simultaneously inhibit movement in (some) non-target fingers.

Next, we conducted a two-way repeated measures ANOVA on the I_A values, with the

condition (three movement-focused conditions) and tapping finger as independent variables,

revealing a significant main effect of condition ($F_{2,30} = 8.10$, $p = 0.002$, $\eta_p^2 = 0.351$) and a

marginally significant main effect of tapping finger ($F_{3,45} = 2.27$, $p = 0.093$, $\eta_p^2 = 0.131$); the

latter also showed a significant interaction ($F_{6,90} = 11.29$, $p < 0.001$, $\eta_p^2 = 0.429$). On one hand,

a one-way repeated measures ANOVA indicated significant differences of I_A values between

the three movement-focused conditions for each finger (all $F_{2,30} > 8.03$, ps < 0.002, $\eta_p^2 > 0.349$).

Post hoc tests replicated the attention-induced facilitation effect (i.e., larger tapping amplitude

for the attended fingers versus unattended fingers) for the index finger (sequence-focused and

index-focused conditions vs. middle-focused condition: both $t_{15} > 3.24$, ps < 0.006), the middle

finger (sequence-focused and middle-focused conditions vs. index-focused condition: both $t_{15} >$

3.54, ps < 0.003), and the ring and little fingers (sequence-focused condition vs. index-focused

and middle-focused conditions: all $t_{15} > 2.14$, ps < 0.049). On the other hand, a one-way

repeated measures ANOVA showed significant differences between tapping fingers for each

condition (all $F_{3,45} > 4.54$, ps < 0.007, $\eta_p^2 > 0.232$). Consistent with experiment 1, post-hoc

tests revealed that amplitude of the little finger increased most when attended in the sequence-

focused condition (all $t_{15} > 2.81$, ps < 0.013) and decreased most when unattended in the index-

focused and middle-focused conditions (all $t_{15} < -3.25$, ps < 0.005). In line with experiment 1,

the above results indicate that the effect of attentional focus on tapping amplitude can be

observed when participants look directly at the fingers they are instructed to notice.

For pattern analysis, a one-sample t-test yielded an above-chance decoding accuracy

for each condition (cells in the diagonal direction: all $t_{15} > 5.66$, ps < 0.001), and mean decoding

accuracy across conditions ($48.26 \pm 2.61\%$, $t_{15} = 8.91$, p < 0.001, Cohen's $d = 2.23$, 95% CI =

[1.29, 3.14]). In addition, as shown in Fig. 6b, such above-chance decoding performance was

observed only for cells in the diagonal direction (i.e., correct predictions), while percentage

values in most other cells (i.e., incorrect predictions) were significantly below chance level.

These results confirm the findings of experiment 1, indicating that attention directed to different

fingers can create distinct patterns of finger-tapping movements.

Results of Experiment 3

In experiment 3, we directed participants' attention to the conceptual label instead of

the movements of their tapping finger(s). Accuracy of the label-counting task was at a level

($72.77\% \pm 3.82\%$) comparable with that of the letter-focused reference condition of experiment

1. After the experiment, participants reported that they paid little attention ($8.66\% \pm 1.43\%$) to

finger movements. Thus, participants might direct their main attention toward the conceptual

label of target finger(s) and hardly notice their finger-tapping behaviors.

We hypothesized that directing attention towards label and away from movement would

have little, if any, impact on automatic finger-tapping movements. To test our hypothesis, we

analyzed the data of experiment 3 following the same analyses as experiments 1 and 2. First, a

one-way repeated measures ANOVA, taking inter-finger coherence and inter-trial coherence as

dependent variables and the focused condition as the independent variable, failed to find a

significant effect (both $F_{3,45} < 0.69$, ps > 0.564, $\eta_p^2 < 0.045$, Fig. 4c). Second, one-sample t-

tests showed no significant changes in I_A values of the three movement-related conditions

relative to zero ($|t_{15}| < 1.65$, ps > 0.120). Third, two-way repeated measures ANOVA on I_A

revealed that the main effect of condition ($F_{2,30} = 0.23$, p $= 0.795$, $\eta_p^2 = 0.015$), main effect of

the tapping finger ($F_{3,45} = 1.36$, p $= 0.267$, $\eta_p^2 = 0.083$), and their interaction ($F_{6,90} = 1.51$, p $=$

0.183, $\eta_p^2 = 0.092$) were all non-significant (Fig. 5c). Finally, we performed a pattern analysis.

As shown in Fig. 6c, above-chance decoding accuracy was present only in the three movement-

related conditions ($t_{15} > 2.45$, ps < 0.027), but not in the reference condition ($t_{15} = 1.26$, p $=$

0.229). Although a one-sample t-test revealed that mean decoding accuracy was still

significantly above-chance ($30.12\% \pm 0.81\%$, $t_{15} = 6.29$, p < 0.001, Cohen's $d = 1.57$, 95% CI

$= [0.82, 2.30]$), a paired sample t-test revealed a significant decrease relative to the above two

experiments (both $t_{15} < -6.78$, ps < 0.001, Cohen's $d > 2.265$). As the decoding accuracy for the

reference condition did not exceed chance level, it is likely that the partially remaining

capability of decoding in movement-related conditions was due to occasionally spreading

attention to the movement of respective fingers during the label-counting task, rather than

methodological artifacts [36] [37]. Therefore, the above findings confirmed our hypothesis,

indicating that it is attention to movement, rather than to conceptual label of respective fingers,

that modulates automatic movement.

535   **Discussion**

The present study was, to our best knowledge, the first to demonstrate that attention can modulate automatic movements without deliberate goals. By directing attention to different fingers of an automatic multi-finger repetitive tapping sequence, our main findings were that attention, if focused on movement rather than the conceptual label of different fingers, could lower inter-finger and inter-trial temporal coherence, increase (or decrease) the tapping amplitude of attended (or non-attended) fingers, and create distinct patterns of finger-tapping movements.

A clear distinction should be made between automatic movements we targeted in the present study and reflexive actions (e.g., the eye-blink reflex elicited by a startling sound) that are instinct and prone to severely interrupting the ongoing tasks. Though reflexive actions can also be modulated by attention [38,39], they only account for a small proportion of actions in our daily life since they usually emerge in threatening situations. On the contrary, most human behaviors are learned during nurturing and have been well-practiced to automaticity (e.g., walking, talking, or finger-tapping) [4-7]. As is known, attention is critical for scheduling a variety of our daily movements. Even for movements guided by particular goals, it is the attention, rather than the goals, that directly modulates automatic movements [5]. From this point of view, we evaluate for the first time in the present study, how attention alone modulates automatic movements when no specific goals are set. Our findings could serve as an indispensable and empirical foundation for theories taking attentional modulation on automatic movements as a prerequisite condition when accounting for the role of attention in complex behaviors such as skill performance and learning [11,40] and goal-directed action control [5].

First, our findings provide empirical foundation for the core idea of the well-known constrained action hypothesis that attention to movement disrupts movement automaticity [10,11]. In line with this proposal, we found reduced temporal coherence, indicating disrupted automaticity, in almost all movement-focused conditions compared with the reference condition. The impaired coherence may be accounted for by mismatch on the coordination speed between systems supporting deliberate attention and automatic movements, respectively [34,41]. As revealed via a task requiring maintaining balance on an unstable surface, attention to foot movement (internal focus condition) leads to slower movement adjustments, while attention directed away from foot (external focus condition) leads to faster movement adjustment [42]. These results indicate a slower pace when attention is manipulated than in automatic mode. Applying the same logic to the present study, the attention may have actively intervened in the fast-paced automatic movement, leading to timing mismatch (as reflected by decreased temporal coherence) between the attended and unattended fingers.

Our findings also provide an empirical foundation for the renowned Norman-Shallice theory [5]. This theory focuses on how deliberate goals control automatic movements by directing attention. A critical and yet to be examined assumption is the bi-directional modulation effect of attention upon automatic movements. In other words, attention by itself would facilitate attended actions and inhibit unattended actions. Since daily action is hard to study in the laboratory [20], researches testing the Norman-Shallice theory put more effort in addressing the proposed role of the frontal lobe in executive control (inhibitory control and/or conflict monitoring) [43], for example using tasks like the Stroop and Wisconsin Card Sort [44-46]. However, they relatively neglected how attention by itself modulates automatic movements. In

the present study, our finding that attention could increase (or decrease) the tapping amplitude

of attended (or non-attended) fingers supplies direct and compelling evidence supporting this

view. Although a similar effect of attention has been widely proven and accepted in the field

of perception [47], to our best knowledge, this study is the first to extend this attentional effect to

automatic movements.

Results of the pattern analysis provide more insight about the interaction between

attention and automatic movements. As revealed, the above-chance decoding accuracy

indicates that attention can dramatically reorganize automatic movements into distinctive

patterns, even when those movements cannot be distinguished from each other according to

their temporal coherence or tapping amplitude. Highlighting the intimate interaction between

attention and automatic movements, the pattern analysis findings support both the constrained

action hypothesis [10,11] and the Norman-Shallice theory [5]. The dependence of automatic

movement patterns varying as a function of attentional focus may be double-edged. On one

hand, newly created movement patterns may fail to satisfy previous requirements and

consequently lead to deteriorated task performance [40]. On the other hand, disruption of the

previously established movement automaticity is necessary to acquire new motor skills,

especially in the early learning stage [8,9].

All findings above conclude with the fundamental view that attention can modulate

automatic movements, even without specific goals. However, the neural modulations, when

attending to automatic movements, have barely been investigated. Prior studies have revealed

that the neural dynamics underlying the development of movement automaticity (task expertise)

are mainly two-folds. On the one hand, neural activation decreases in a wide variety of motor-

related areas (e.g., cerebellum, cingulate motor area, primary motor cortex and supplementary

motor area)[15,48,49] and the attention system (especially the dorsolateral prefrontal cortex, i.e.,

DLPFC, and anterior cingulate cortex, i.e., ACC [18,50]). These results suggest that attention

becomes less important when there evolves a more efficient neural code for controlling

movements. On the other hand, neural activation increases in a critical subcortical region, i.e.,

the sensorimotor territory of the striatum [7], as well as its effective connectivity with these

motor-related regions [15,48]. These results suggest that motor skills are likely encoded across

multiple representations within a subcortico-cortical loop [51] centered on the striatum.

609   When attention is directed back to automatic movements, we surmise that the

subcortical striatum would be of different neural activity and connectivity with motor-related

regions and the attention system. Though prior studies examining this possibility [15,18] provided

no support for this hypothesis, two critical issues should be considered, pointing out future

studies' direction. At first, the task they used (i.e., initiation of visually cued keypress

movements) might have mainly measured dynamics related to deliberate control processes,

because initiation of a movement sequence, which usually requires deliberate control [5], can

hardly be trained to automatic despite the level of training [16,17]. For this reason, the striatum

may not be effectively engaged in related tasks. Alternatively, the paradigm used in the present

study may provide a promising solution. On the other hand, the fact that they did not find

differences in the striatum does not necessarily mean differences do not exist. For this reason,

future studies are encouraged to employ more sensitive analyses (e.g., pattern analysis) to

delineate neural dynamic changes, if any, in the striatum and other related regions.

Future studies are also encouraged to compare the neural dynamic changes between the

healthy and motor deficit (e.g., Parkinson's disease) patients, to obtain a better understanding

of motor-deficit pathophysiology. Inspired by the Norman-Shallice theory [5], continued

investigations should also bring goals into consideration [52], to further examine whether and

how attention translates deliberate goals into automatic movement sequences. Limitations of

this study include that we did not track eye movements, which prevented us from evaluating to

what extent participants had locked their gaze on the required focus. The fact that we did not

counterbalance the order of experiments constitutes another limitation, although the order

effects, if any, would decay during the interval (spanning weeks) between experiments.

In sum, we found for the first time that attention can modulate automatic movements

without assistance from deliberate goals. Once being directed to perform a sequence of

automatic movements, focused attention disrupts movement automaticity, facilitates attended

and inhibits unattended movements, and re-organizes the movement sequence into distinct

patterns according to the focus of attention.

636
637

Reference

1   Strack, F. & Deutsch, R. Reflective and impulsive determinants of social behavior. *Personality and*
social psychology review **8**, 220-247 (2004).

2   Fourneret, P. & Jeannerod, M. Limited conscious monitoring of motor performance in normal subjects.
Neuropsychologia **36**, 1133-1140 (1998).

3   Wegner, D. M. & Bargh, J. A. Control and automaticity in social life.   (1998).

4   Shiffrin, R. M. & Schneider, W. Controlled and automatic human information processing: II. Perceptual
learning, automatic attending and a general theory. *Psychological review* **84**, 127 (1977).

5   Norman, D. A. & Shallice, T. Attention to action: Willed and automatic control of behavior. in
Consciousness and Self-Regulation: Advances in Research and Theory Volume 4   (eds Richard J.
Davidson, Gary E. Schwartz, & David Shapiro)   1-18 (Springer US, 1986).

6   Wu, T., Kansaku, K. & Hallett, M. How Self-Initiated Memorized Movements Become Automatic: A
Functional MRI Study. *Journal of Neurophysiology* **91**, 1690-1698, doi:10.1152/jn.01052.2003 (2004).

7   Lehéricy, S. *et al.* Distinct basal ganglia territories are engaged in early and advanced motor sequence
learning. *Proceedings of the National Academy of Sciences* **102**, 12566-12571 (2005).

8   Doyon, J., Penhune, V. & Ungerleider, L. G. Distinct contribution of the cortico-striatal and cortico-
cerebellar systems to motor skill learning. *Neuropsychologia* **41**, 252-262 (2003).

9   Willingham, D. B. A neuropsychological theory of motor skill learning. *Psychological review* **105**, 558
(1998).

10   Wulf, G., McNevin, N. & Shea, C. H. The automaticity of complex motor skill learning as a function of
attentional focus. *The Quarterly Journal of Experimental Psychology Section A* **54**, 1143-1154 (2001).

11   Wulf, G., Shea, C. & Park, J.-H. Attention and motor performance: preferences for and advantages of
an external focus. *Research quarterly for exercise and sport* **72**, 335-344 (2001).

12   Wulf, G., Dufek, J. S., Lozano, L. & Pettigrew, C. Increased jump height and reduced EMG activity
with an external focus. *Human movement science* **29**, 440-448 (2010).

13   Freudenheim, A. M., Wulf, G., Madureira, F., Pasetto, S. C. & Corrěa, U. C. An external focus of
attention results in greater swimming speed. *International Journal of Sports Science & Coaching* **5**,
533-542 (2010).

14   Lohse, K. R., Sherwood, D. E. & Healy, A. F. How changing the focus of attention affects performance,
kinematics, and electromyography in dart throwing. *Human Movement Science* **29**, 542-555 (2010).

15   Wu, T. *et al.* Attention to automatic movements in Parkinson's disease: Modified automatic mode in the
striatum. *Cerebral Cortex* **25**, 3330-3342 (2015).

16   de Wit, S. *et al.* Shifting the balance between goals and habits: Five failures in experimental habit
induction. *Journal of Experimental Psychology: General* **147**, 1043 (2018).

17   Watson, P. & de Wit, S. Current limits of experimental research into habits and future directions.
Current Opinion in Behavioral Sciences **20**, 33-39 (2018).

18   Jueptner, M. *et al.* Anatomy of motor learning. I. Frontal cortex and attention to action. *Journal of*
neurophysiology **77**, 1313-1324 (1997).

19   Mathis, A. *et al.* DeepLabCut: markerless pose estimation of user-defined body parts with deep
learning. *Nature Neuroscience*, doi:10.1038/s41593-018-0209-y (2018).

20   Schwartz, M. F. The cognitive neuropsychology of everyday action and planning. *Cognitive*
Neuropsychology **23**, 202-221, doi:10.1080/02643290500202623 (2006).

680	21	Wu, T., Wang, L., Hallett, M., Li, K. & Chan, P. Neural correlates of bimanual anti-phase and in-phase
681		movements in Parkinson's disease. *Brain* **133**, 2394-2409, doi:10.1093/brain/awq151 (2010).
682	22	Thunell, E. & Thorpe, S. J. Memory for Repeated Images in Rapid-Serial-Visual-Presentation Streams
683		of Thousands of Images. *Psychological Science* **30**, 989-1000, doi:10.1177/0956797619842251 (2019).
684	23	Zhou, L., Deng, C., Ooi, T. L. & He, Z. J. Attention modulates perception of visual space. *Nature*
685		*Human Behaviour* **1**, 0004, doi:10.1038/s41562-016-0004 (2016).
686	24	Brainard, D. The Psychophysics Toolbox. *Spat Vis* **10**, 433-436 (1997).
687	25	Beilock, S. L., Carr, T. H., MacMahon, C. & Starkes, J. L. When paying attention becomes
688		counterproductive: impact of divided versus skill-focused attention on novice and experienced
689		performance of sensorimotor skills. *Journal of Experimental Psychology: Applied* **8**, 6 (2002).
690	26	Beilock, S. L. & Carr, T. H. On the fragility of skilled performance: What governs choking under
691		pressure? *Journal of experimental psychology: General* **130**, 701 (2001).
692	27	Castaneda, B. & Gray, R. Effects of focus of attention on baseball batting performance in players of
693		differing skill levels. *Journal of Sport and Exercise Psychology* **29**, 60-77 (2007).
694	28	Gray, R. Attending to the execution of a complex sensorimotor skill: Expertise differences, choking,
695		and slumps. *Journal of Experimental Psychology: Applied* **10**, 42 (2004).
696	29	Jobbágy, Á., Harcos, P., Karoly, R. & Fazekas, G. Analysis of finger-tapping movement. *Journal of*
697		*Neuroscience Methods* **141**, 29-39, doi:https://doi.org/10.1016/j.jneumeth.2004.05.009 (2005).
698	30	Aoki, T., Francis, P. R. & Kinoshita, H. Differences in the abilities of individual fingers during the
699		performance of fast, repetitive tapping movements. *Experimental Brain Research* **152**, 270-280 (2003).
700	31	Kilbreath, S. L., Gorman, R. B., Raymond, J. & Gandevia, S. C. Distribution of the forces produced by
701		motor unit activity in the human flexor digitorum profundus. *Journal of Physiology* **543**, 289-296
702		(2010).
703	32	Ingram, J. N., Körding, K. P., Howard, I. S. & Wolpert, D. M. The statistics of natural hand
704		movements. *Experimental brain research* **188**, 223-236 (2008).
705	33	Grinsted, A., Moore, J. C. & Jevrejeva, S. Application of the cross wavelet transform and wavelet
706		coherence to geophysical time series. *Nonlinear processes in geophysics* **11**, 561-566 (2004).
707	34	Ballard, D. H., Hayhoe, M. M., Pook, P. K. & Rao, R. P. Deictic codes for the embodiment of
708		cognition. *Behavioral and Brain Sciences* **20**, 723-742 (1997).
709	35	Espay, A. J. *et al.* Impairments of speed and amplitude of movement in Parkinson's disease: A pilot
710		study. *Movement Disorders* **24**, 1001-1008, doi:10.1002/mds.22480 (2009).
711	36	Martin, J. K. & Hirschberg, D. S. *Small sample statistics for classification error rates I: Error rate*
712		*measurements*. (Information and Computer Science, University of California, Irvine, 1996).
713	37	Chen, Y., Garcia, E. K., Gupta, M. R., Rahimi, A. & Cazzanti, L. Similarity-based classification:
714		Concepts and algorithms. *Journal of Machine Learning Research* **10**, 747-776 (2009).
715	38	De la Casa, L., Mena, A. & Ruiz Salas, J. Effect of Stress and Attention on Startle Response and
716		Prepulse Inhibition. *Physiology & Behavior* **165**, doi:10.1016/j.physbeh.2016.07.022 (2016).
717	39	Lang, P., Bradley, M. & Cuthbert, B. Emotion, Attention, and the Startle Reflex. *Psychological review*
718		**97**, 377-395, doi:10.1037/0033-295X.97.3.377 (1990).
719	40	Wulf, G. Attentional focus and motor learning: a review of 15 years. *International Review of sport and*
720		*Exercise psychology* **6**, 77-104 (2013).
721	41	Passingham, R. E. Attention to action. *Philosophical transactions of the Royal Society of London.*
722		*Series B, Biological sciences* **351**, 1473-1479, doi:10.1098/rstb.1996.0132 (1996).
723	42	McNevin, N. & Shea, C. Increasing the distance of an external focus of attention enhances learning.

Psychological research **67**, 22-29, doi:10.1007/s00426-002-0093-6 (2003).

43   Badgaiyan, R. D. Executive control, willed actions, and nonconscious processing. *Human brain*
mapping **9**, 38-41, doi:doi:10.1002/(SICI)1097-0193(2000)9:1<38::AID-HBM4>3.0.CO;2-T (2000).

44   Y, D., Kimberg, J, M. & Farah. A unified account of cognitive impairments following frontal lobe
damage: the role of working memory in complex, organized behavior. *Journal of experimental*
psychology. General (1993).

45   Stanislas, D. & Jean-Pierre, C. The Wisconsin Card Sorting Test: Theoretical Analysis and Modeling in
a Neuronal Network. *Cerebral Cortex*, 62-79 (1991).

46   Cohen, J., Dunbar, K. & Mcclelland, J. On the control of automatic processes: A parallel distributed
processing model of the Stroop effect. *Psychological Review* **97**, 332-361 (1988).

47   Herrmann, K., Montaser-Kouhsari, L., Carrasco, M. & Heeger, D. J. When size matters: attention
affects performance by contrast or response gain. *Nature neuroscience* **13**, 1554 (2010).

48   Wu, T., Hallett, M. & Chan, P. Motor automaticity in Parkinson's disease. *Neurobiology of Disease* **82**,
226-234, doi:https://doi.org/10.1016/j.nbd.2015.06.014 (2015).

49   Poldrack, R. A. *et al.* The Neural Correlates of Motor Skill Automaticity. *The Journal of Neuroscience*
25, 5356-5364, doi:10.1523/jneurosci.3880-04.2005 (2005).

50   Isomura, Y. Neural coding of "attention for action" and "response selection" in primate anterior
cingulate cortex. *J. Neurosci* **23** (2003).

51   Berlot, E., Popp, N. J. & Diedrichsen, J. R. In search of the engram, 2017. *Current Opinion in*
Behavioral Sciences **20**, 56-60 (2018).

52   Custers, R. & Aarts, H. The Unconscious Will: How the Pursuit of Goals Operates Outside of
Conscious Awareness. *Science* **329**, 47-50, doi:10.1126/science.1188595 (2010).

Acknowledgements. This research was supported by grants from the National Key Research

& Development Program of China (2018YFB1305200) and the National Natural Science

Foundation of China (61921004) for W.Z., and Postdoctoral Science Foundation of China

(2019M661703) and the Fundamental Research Funds for the Central Universities

(2242020R20021) for X.Z.

Author Contributions Statement. X.Z. developed the study concept and designed the study.

Testing and data collection were performed by X.Z. and X.J; X.Z., X.J., and X.Y. performed

the data analysis and interpretation under the supervision of W.Z; X.Z. drafted the manuscript,

and X.Y. and W.Z. provided critical revisions. All authors approved the final version of the

manuscript for submission.

Additional Information. The authors declare no competing interests.

757   **Figure Legends**

Figure 1. Schematic illustration of the experimental paradigm of (a) experiment 1, (b)

experiment 2, and (c) experiment 3. In experiments (a) 1 and (c) 3, a secondary letter-counting

or label-counting task was performed simultaneously with the finger-tapping task. The focus

of attention was manipulated in each experiment so that either different tapping fingers

(indicated by the pink shade on hand) were attended (experiments (a) 1 and (b) 2), or different

labels of particular fingers were attended (experiment (c) 3).

Figure 2. Recording and quantification of automatic finger tapping movements. *(a)* One

exemplar frame illustrating the location of 17 key points of interest of the right-hand fingers.

For each frame, the coordinates of these points were automatically recognized using

DeepLabCut [19]. *(b)* Concrete frames illustrating changes in the position of key points during

repetitive finger tapping movements. For visibility, only key points on the fingertips are shown.

(c) Y-axis trajectories of the index finger from two exemplar trials. Dotted lines represent the

onset of pure tones informing participants to initiate and terminate finger tapping, respectively.

The solid line indicates the beginning of the temporal range of interest. *(d)* The 2-D map

produced by wavelet transform coherence (WTC) analysis between the above two trials. The

color bar indicates the coherence intensity, and the orientations of arrows indicate the circular

phase lag at each time-frequency point. The yellow rectangle indicates the temporal-frequency

range of interest determined for this representative participant. To exclude the distorting effect

of edge artifacts, values in the area masked by translucent white were not included in further

analyses. *(e)* Tapping rate is individually different but stable across experiments. Though

experiments were performed under different task demands and in separate days, the rate of

tapping movements between experiments was highly correlated (Pearson $r > 0.91$).

Figure 3. Objective and subjective measures of attention directed to finger-tapping movement

in experiment 1. *(a)* Objective measure of attention: accuracy (ACC) of the secondary letter-

counting task as a function of attentional focus. *(b)* Subjective measure of attention: self-rated

amount of attention directed to movement as a function of attentional focus. *(c)* More attention

directed to movement relative to the reference condition (letter-focused condition) led to less

attention available for the secondary task, thus possibly contributing to more decrease in ACC

of the secondary task. The error bar indicates one SEM.

Figure 4. Temporal coherence of finger trajectories varying as a function of attentional focus.

Each panel indicates results with respect to inter-finger (shown in blue) and inter-trial (shown

in orange) temporal coherence, with x-axis labels indicating four attention conditions with

different attentional foci. The error bar indicates one SEM.

Figure 5. Tapping amplitude of finger trajectories varying as a function of attentional focus**.**

Each panel indicates results with respect to the amplitude for each tapping finger normalized

to the reference condition. X-axis labels indicate four attentional focus conditions, with the first

label indicating the reference condition whose amplitude had been normalized to 0. The error

bar indicates one SEM.

Figure 6. Distinct patterns of automatic movements varying as a function of attentional focus.

For each panel and cell, the value corresponds to the ratio of samples of the true condition,

which were classified as the predicted condition. Statistics were conducted against the chance

level (0.25) by using a one-sample t-test. Darker cells indicate values above chance while

whiter cells indicate values below chance. *, $p < 0.05$, ***, $p < 0.001$.