1   **Evaluating Human Autosomal Loci for Sexually Antagonistic Viability Selection in Two**
2   **Large Biobanks**
3

4   Authors: Katja R. Kasimatis*[,†,**], Abin Abraham[‡,**], Peter L. Ralph*, Andrew D. Kern*, John A.
5   Capra[‡,§], Patrick C. Phillips*
6
7   *Institute of Ecology and Evolution, University of Oregon, Eugene, USA
8   [†]Department of Ecology and Evolutionary Biology, University of Toronto, Canada
9   [‡]Vanderbilt Genetics Institute, Vanderbilt University Medical Center, Nashville, USA
10 [§]Department of Biological Sciences, Vanderbilt University, Nashville, USA
11
12 **These authors contributed equally.

13   **Running Title:** Evaluating Sexually Antagonistic Variants in Humans
14
Keywords: sexually antagonistic selection, male-female F_{ST}, biobank, association study, probe
mapping

Corresponding Authors:
John A. Capra
U5221 BSB/MRB III
VU Station B, Box 35-1634
Nashville, TN 37235
tony.capra@vanderbilt.edu
615-343-3671
25
Patrick C. Phillips
272 Onyx Bridge
5289University of Oregon
Eugene, OR 97403
pphil@uoregon.edu
541-346-0916

ABSTRACT

Sex and sexual differentiation are ubiquitous across the tree of life. Because females and males often have substantially different functional requirements, we expect selection to differ between the sexes. Recent studies in diverse species, including humans, suggest sexually antagonistic viability selection creates allele frequency differences between the sexes at many different loci. However, theory and population-level simulations indicate that sex-specific differences in viability would need to be very extreme in order to produce and maintain reported levels of between-sex allelic differentiation. We address this paradox between theoretical predictions and empirical observations by evaluating evidence for sexually antagonistic viability selection on autosomal loci in humans using the largest cohort to date (UK Biobank, n=438,427) along with a second large, independent cohort (BioVU, n=93,864). We performed association tests between genetically ascertained sex and genotypes. Although we found dozens of genome-wide significant associations, none replicated across samples. Moreover, closer inspection revealed that all associations are likely due to cross-hybridization with sex chromosome regions during genotyping. We report loci with potential for mis-hybridization found on commonly used genotyping platforms that should be carefully considered in future genetic studies of sex-specific differences. Despite being well-powered to detect allele frequency differences of up to 0.8% between the sexes, we do not detect evidence for this signature of sexually antagonistic viability selection on autosomal variation. These findings suggest a lack of strong ongoing sexually antagonistic viability selection acting on single locus autosomal variation in humans.

INTRODUCTION

Understanding the relationship between genotype and sexually dimorphic phenotypes, and how

selection shapes this relationship, is fundamental to understanding sex-specific responses in

aging (Archer *et al.* 2018), fertility (Farquhar *et al.* 2019), disease susceptibility (Morrow 2015;

Ferretti *et al.* 2018; Dumitrescu *et al.* 2019), and treatment (Khramtsova *et al.* 2018). Sexual

dimorphism is common across a range of plant and animal taxa (Rowe *et al.* 2018; Deegan and

Engel 2019). Differences in optimal fitness values between the sexes may result in sexually

antagonistic selection (Arnqvist and Rowe 2005) – i.e., selection on autosomal variants that

affect fitness in different directions for each sex. Surveys of natural selection suggest that the

repeated evolution of sexual dimorphism is commonly associated with sexually antagonistic

selection (Cox and Calsbeek 2009). Yet, we still lack an understanding of how this process

shapes genomic variation within and between species. A major obstacle in assessing the

genomic consequences of sexually antagonistic selection is that most of the hypothesized

genomic signatures are not unique to this mode of selection. However, when the alleles at a

single locus have opposite effects on viability between the sexes, namely intralocus sexual

conflict (Rice and Chippindale 2001), then sexually antagonistic viability selection is

hypothesized to favor different alleles in each sex. This process is predicted to generate allele

frequency differences between the sexes among adults (Mank 2017; Kasimatis *et al.* 2017).

Recent research has looked for this signature of selection by identifying alleles with high

male-female F_{ST} (Cheng and Kirkpatrick 2016; Kasimatis *et al.* 2017; 2019), a normalized

measure of allele frequency difference. Such studies across a range of taxa have suggested

that potentially hundreds of autosomal loci are subject to ongoing sexually antagonistic selection

with many differentiated loci having male-female divergence values of at least 10% (Lucotte *et*

al. 2016; Flanagan and Jones 2017; Wright *et al.* 2018; Dutoit *et al.* 2018; Bissegger *et al.*

2019), and some reaching even as high as 45% (Vaux *et al.* 2019). These results are surprising

because the production and maintenance of such large male-female differences on autosomes

requires strong, ongoing selection to overcome the homogenization of genotypes during meiotic

segregation every generation (Cheng and Kirkpatrick 2016; Kasimatis *et al.* 2019). Theory

suggests that a male-female F_{ST} value of 1% requires at least a 33% viability cost per sex per

generation (Kasimatis *et al.* 2019). Given the high sex-specific viability cost, factors such as

population structure, sampling variance due to small sample sizes, or bioinformatic artifacts may

contribute to the high divergence values observed (Kasimatis *et al.* 2019). Of particular concern

are the small sample sizes (15 to 100 individuals) used by many previous studies. Detecting the

level of allelic differentiation expected at sexually antagonistic loci with moderate sex-specific

mortality (≤10% per sex) requires substantially larger sample sizes and accounting for other

possible confounding effects, such as population structure (Kasimatis *et al.* 2019). Indeed, a

meta-analysis of 51 studies that included more than 100,000 European-ancestry individuals did

not find any common variants associated with sex ratio (Boraska *et al.* 2012).

We aim to reconcile empirical observations with theoretical predictions using a robust

statistical framework to identify intralocus sexually antagonistic viability selection in the largest

human cohort to date. We use two large-scale biobanks, the UK Biobank and the Vanderbilt

Biobank (BioVU) to analyze >500,000 human genomes for signals of male-female divergence

driven by sexually antagonistic selection. We rigorously control for population stratification and

potential molecular and informatic artifacts. Compared to previous studies examining sexual

antagonism, these datasets significantly improve our statistical power to detect allele frequency

differences among the sexes by providing the largest available sample sizes to date – several

orders greater than previous studies in humans (Lucotte *et al.* 2016; Cheng and Kirkpatrick

2016) and non-model taxa (Lucotte *et al.* 2016; Flanagan and Jones 2017; Wright *et al.* 2018;

Dutoit *et al.* 2018; Bissegger *et al.* 2019; Vaux *et al.* 2019). Our association framework differs

from traditional association studies as genetic sex is the phenotype of interest and the

mechanism generating a true effect would be sex-specific viability. After controlling for multiple

confounders, we are unable to detect evidence for ongoing sexually antagonistic viability

selection at individual autosomal loci.

MATERIALS AND METHODS

Genotyping and quality control in BioVU

The DNA biobank at Vanderbilt University, consists of DNA extracted from blood collected

during routine clinical testing. For 93,864 individuals, GWAS-level genotyping was performed

using the Illumina MEGA-Ex chip which includes >2 million common and rare variants before

imputation. We obtained genotyped data in PLINK format from the Vanderbilt sequencing core

after the following quality control steps: excluding either samples or variants with ≥5%

missingness, mismatched identifiers as detected by identity by descent checks, and non-

concordance between reported gender and genetically determined sex. Overlapping variants

with 1000 Genomes demonstrated ≥99.98% variant call concordance using HapMap sample

aliquots. Using PLINKv1.90b3s (Chang *et al.* 2015), we additionally performed the following

quality control steps. We first confirm that duplicate samples and those with high missing rate

(≥5%) are not present and exclude samples with high heterozygosity on autosomes (>3 S.D.

from observed data), or high relatedness (%IBD ≥ 0.2). Next, we removed duplicated variants

and variants with high missing rate (≥5%) or significantly different missing rate between cases

and controls (p < 0.00001, Fisher's Exact test). We then included only samples with a self or

third party reported race as 'white' and variants with minor allele frequency >0.01. This

additional quality control resulted in a final European-ancestry dataset of 61,760 samples

(34,269 females and 27,491 males) and 1,763,607 variants. We calculated the top 12 principal

components on this cohort. We imputed variants that reached nominal or genome-wide

statistical significance (P < 5E-8) in the UK Biobank data but were not genotyped in the BioVU

cohort. These variants were imputed using the Michigan Imputation Server (v1.2.4) (Das *et al.*

2016) using the HRC (Version r1.1 2016) reference panel and retaining variants with $R^2 > 0.3$.

Imputed allele dosages were converted to hard calls using PLINK/2.00-alpha2 (Chang *et al.*

2015) and filtered to exclude variants with minor allele frequency <1% and genotyping rate

<95%. All PLINK code is available on the GitHub repository https://github.com/abraham-

abin13/sexually_antagonistic_sel.git.

Genotyping and quality control in the UK Biobank

The UK Biobank is an international health resource with data from approximately 500,000

participants. Genotyping and quality control procedures have previously been described in detail

by Bycroft et al. (2018). Briefly, two arrays – the UK Biobank Axiom Array (n = 438,427

participants) and the UK BiLEVE Axiom Array (n = 49,950 participants) – were used to genotype

participants (71 bp oligos). Quality control procedures carried out before the data were released,

included: removal of participants with excess heterozygosity or missingness, removal of

markers with batch, plate, array, or sex effects, and removal of markers with discordance across

control replications (Bycroft *et al.* 2018). The removal of sex effects, namely allele frequency

differences between females and males at a given marker, does not preclude our analysis as

the conservative threshold (P < 10E-12) removed only eight markers and the sex differences at

these markers were due to technical artifacts, such as the probe sequence mapping to the Y

chromosome (C. Bycroft pers. comm.). The released genotype data contains 805,462 markers

from 488,377 participants (Field IDs 22100-22124). Additionally, the genetic sex (Field ID

22001), year of birth (Field ID 34), date of assessment (Field ID 53), and assessment center

(Field ID 54) were requested for each participant. The top 40 genetic principal components

(Field ID 22009) were previously calculated using fastPCA (Bycroft *et al.* 2018).

150   Using PLINKv1.90b3s (Chang *et al.* 2015), we additionally performed the following

quality control steps. We excluded samples with high missing rate (≥5%) and high

heterozygosity on autosomes (>3 S.D. from observed data). Next, we pruned markers in

linkage-disequilibrium (window size = 50kb, step rate = 5, r^2 threshold = 0.2). Finally, we

removed variants with significantly different missing rate between cases and controls (P <

0.00001, Fisher's Exact test). We included only variants with minor allele frequency > 0.01 to

exclude inaccurate calls made for low frequency alleles (Wright *et al.* 2019; Weedon *et al.*

2019). This additional quality control resulted in a final dataset of 488,291 samples (264,813

females and 223,478 males) and 653,632 variants. A binomial test was used to test for a lack of

minor allele homozygotes relative to that expected under HWE (this is conservative, because

most human population dynamics is expected to lead to an excess of homozygosity). All PLINK

code is available on the GitHub repository https://github.com/abraham-

abin13/sexually_antagonistic_sel.git

Imputed genotype and phased haplotype values were used to compare significant loci in

the BioVU cohort, which were not directly genotyped in the UK BIOBANK arrays. Again,

imputation was completed prior to the data release using the Haplotype Reference Consortium

and UK10K haplotype resource. The imputation methods are described in detail in Bycroft et al.

(2018). Imputed allele dosages were converted to hard calls using PLINK/2.00-alpha2 (Chang

et al. 2015).

Genome-wide association for an individual's sex

We performed a GWAS in UK Biobank and BioVU separately using a logistic regression testing

the association between an individual's sex (binary variable, concordant with their genetic sex)

and the effect allele, defined as the minor allele by PLINKv1.90b3s (Chang *et al.* 2015), using

an additive model. For BioVU analysis, we controlled for genetic ancestry using 12 genetic

principal components and included year of birth as a covariate. For the UK Biobank analysis, we

again controlled for genetic ancestry using 12 genetic principal components, along with age at

assessment and UK Biobank sampling center as covariates. All genome wide association tests

were done using PLINKv1.90b3s (Chang *et al.* 2015). We focused our analyses on the

autosomes, where genomic divergence between the sexes is not confounded by sex

chromosome processes. During our quality control steps before association testing, we did not

remove variants based on deviations from Hardy Weinberg Equilibrium (HWE) since theory

indicates that sex-specific selection can violate the assumptions of HWE (Kasimatis *et al.* 2019).

Resampling of sex and generating a null distribution

To determine if p-values were well calibrated (i.e., uniformly distributed on [0,1]) at non-

associated variants, we performed a permutation analysis to calculate the distribution of p-

values within the UK Biobank cohort. We resampled genetic sex 100 times per chromosome to

generate a set of random associations between genotype and this phenotype. We then reran

the logistic regression, again including 12 genetic principal components, age, and sampling

center as covariates, for only those variants that had a p-value < 0.01 in the original association

analysis (n = 8,868 SNPS). These analyses generated a distribution of 100 p-values at each

variant. Permuted p-values were uniformly distributed (Supplementary Figure 1A), even when

the values were small (Supplementary Figure 1B), indicating the p-values for this association

analysis were well-calibrated and therefore a genome-wide Bonferroni significance threshold of

P < 5E-8 was appropriate. All R and PLINK code are available on the GitHub repository

https://github.com/abraham-abin13/sexually_antagonistic_sel.git.

Identifying SNPs with sequence similarity to sex chromosomes

Incorrectly mapped sex-chromosome variants to an autosomal region can result in statistically

significant GWAS hits for an individual's sex due to the different effects on allele counts

between females and males. We used BLAT (Kent 2002) with default parameters (stepSize=5,

repMatch=2253, minScore=20, minIdentity=0) to identify sequence similarity between the probe

sequences used on the genotyping array and sex chromosome regions. The MEGA-Ex array

probe sequences used to genotyped the BioVU cohort were obtained directly from Illumina.

Probes sequence for the UK Axiom Biobank array (Resource 149601) and UK BiLEVE array

(Resource 149600) were download from https://biobank.ctsu.ox.ac.uk/crystal/label.cgi?id=263.

MEGA-Ex probes are 50 base pair sequences adjacent to the variant being tested; MEGA-Ex

uses single base extension to detect the variant allele. UK Biobank array probes are 71 base

pairs long with the variant being genotyped located in the middle. BLAT hits to the X or Y

chromosome were further filtered to identify regions likely to cross-hybridize by requiring at least

40 base pair overlap, sequence similarity ≥90%, and that the matching sequence overlaps (UK

Biobank arrays) or flanks (MEGA-Ex array) the variant being tested. Similar criteria were used in

a previous a study that reported cross-hybridization on the Illumina Infinium

HumanMethylation27K microarray platform (Chen *et al.* 2012). Next, we identified the best

BLAT hit to a sex chromosome for each probe sequence by selecting the hit with the highest

BLAT score, which accounts for match length and sequence similarity. For this step, we

considered the UK Axiom and UK BiLEVE array together thus selecting the probe sequence

with the highest BLAT score from one of the two arrays per variant tested in the GWAS. In the

BioVU (MEGA-Ex array) and UK Biobank arrays, 83,083 out of 798,051 and 128,090 out of

620,040 autosomal probes had at least one BLAT match (BLAT score ≥ 20) to a sex

chromosome region. To further focus on sequence similarity with potential to cause genotyping

error, we identified sex chromosome matches with the following criteria (Chen *et al.* 2012): 1)

≥40 base pairs in length, 2) ≥90% sequence similarity, and 3) overlap between the match and

the variant being genotyped.

Power Analysis

We conducted a power analysis to determine how the minimum allelic divergence between the

sexes that could be detected within the BioVU and UK Biobank cohorts (Supplemental File 9).

Specifically, we determined the probability that we would reject the null observation that the

population frequency of each allele is equal at a p-value threshold of P = 1E-8. Suppose we

have N males and M females, and the allele frequencies in the two groups are P and Q. Since

the cohort sample sizes are large, if the population frequencies are p and q, then P ~ Normal(p,

p (1-p) / 2N) and Q ~ Normal(q, q (1-q) / 2M). The difference in population allele frequencies is

then given by P - Q ~ Normal(p - q, p (1-p) / 2N + q (1-q) / 2M). The variance is maximized

when p = q = 1/2, so is at most the variance in the population is: V = (1/N + 1/M)/8. The two-

sided p-value for P-Q being nonzero will be below 1E-8 if |P-Q| is larger than z(0.5e-8) *sqrt(V),

where z(p) is the p-th quantile for the standard Normal distribution. Even, then, an allele with |P-

Q| = z(0.5E-8) * sqrt(V) will only have a two-sided p-value half the time; alleles must by slightly

farther apart (by z(0.025) * sqrt(V)) to have a 95% probability that statistical noise does not put

them above the p=1e-8 threshold. Therefore, we will have 95% power to detect any SNP with

true |p-q| > (z(0.5E-8) + z(0.025)) * sqrt(V).

Data Accessibility

All the data generated from this study (Supplemental Files 1-9) were deposited in the figshare

repository https://figshare.com/s/e863ea11cc9dab30c1b9 to be made public upon publication.

All the code generated for this study were deposited in a GitHub repository to be made public

upon publication.

246   **RESULTS**

247   Throughout this paper when we refer to an individual's sex, we are referencing that individual's

248   sex chromosome composition as estimated in each biobank dataset and binarized (i.e.,

249   metadata reports each individual as XY or XX, although the datasets almost certainly include

250   individuals not falling into these two categories (Lanfranco *et al.* 2004)). We make no

251   statements in relation to gender, which is determined by many factors beyond genetics.

Seventy-seven variants show genome-wide significance as candidates for sexually antagonistic selection

To identify autosomal variants that could be under sexually antagonistic selection, we performed a genome-wide association study (GWAS) between females and males in two large, independent cohorts (BioVU: 34,269 females and 27,491 males; UK Biobank: 264,813 females and 223,478 males). We first applied standard quality control steps to remove samples with high relatedness, discordant sex, or high heterozygosity and excluded genotyped variants with high overall missing rate (Methods). We account for potential confounders by including age and 12 principal components for population stratification as covariates. The resulting p-values are well-calibrated, as verified by permuting the sex labels in the UK Biobank cohort (Supplementary Figure 1A), and so the standard genome-wide significance threshold of P < 5E-8 is appropriate for the association analysis (Methods, Supplementary Figure 1B). Applying this threshold resulted in five and 72 genome-wide significant variants in BioVU and UK Biobank, respectively.

Since different amount of missing data for each variant between cases and controls can lead to spurious associations (Moskvina *et al.* 2006), we tested variants for a statistically significant difference in the missing rate between females and males (Methods). This control excluded what would have been 64 genome-wide significant variants in the UK Biobank and none in the BioVU cohort (Supplemental Figure 2, Supplementary File 1), leaving us with eight and five variants in the two datasets, respectively (Figure 1, Table 1). One intriguing genome-wide significant variant in the UK Biobank cohort, (rs11032483; OR = 1.25, P < 1.3E-53), which lies in a known regulatory region on chromosome 11 and has evidence from association studies for increasing risk in males and being protective in females for a number of sex-specific reproductive pathologies (Cortes *et al.* 2018).

No candidate loci replicate across BioVU and the UK Biobank

Comparing the five autosomal significant hits from BioVU to the eight from the UK Biobank, none of the associations are genome-wide significant in both cohorts (Table 1). Furthermore, none of the significant hits in one cohort even meet a nominal significance threshold (P < 0.05) in the other cohort. For example, the variant with the strongest association in the UK Biobank cohort (rs11032483) had no evidence for association with sex in the BioVU cohort (P = 0.99).

The regions surrounding each of the significant variants do not exhibit the expected association signal clusters arising from variants in strong linkage disequilibrium (LD) with the causal variant. For example, the most strongly associated variant overall (rs9870157) has 33

variants with R^2 of at least 0.8 in the 1000 Genomes Phase 3 European-ancestry (EUR)

populations. However, there are no other strong associations among these variants. The lack of

replication across the two cohorts and the missing association peaks among variants in strong

LD suggest that these signals could be false positives driven by technical or biological artifacts.

Significant associations are likely due to mis-hybridization with sex chromosome regions

Genotyping error can occur due to probe cross-reactivity between different regions of the

genome. Sex-biased error has been observed in array-based studies of DNA methylation (Chen

et al. 2013) and has been reported in the canid genome (Tsai *et al.* 2019), the stickleback

genome (Bissegger *et al.* 2019), and on the Y chromosome in humans (Boraska *et al.* 2012;

Lucotte *et al.* 2016). For instance, if an autosomal variant is assayed with a probe sequence that

has sufficient sequence similarity to a Y chromosome region carrying the reference allele, then

males homozygous for the alternate allele at the autosomal locus may instead be genotyped as

heterozygous for the alternate allele. Females would not be subject to this bias, and thus there

would appear to be an allele frequency difference between the sexes. Similarly, an autosomal

variant with a probe sequence with high similarity to the X chromosome could result in a lack of

homozygotes for the allele not on the X chromosome in both sexes, but the strength of this

effect would differ between females and males. Furthermore, such cross-reactivity can lead the

normalized intensities produced by genotyping arrays to lie outside of the regions corresponding

to each genotype, and thus a missing genotype (Zhao *et al.* 2018). Cross reactivity to a sex

chromosome could therefore cause a differential missingness rate between the sexes. Indeed,

we observe an almost complete lack of minor allele homozygotes in males across all thirteen

genome-wide significant SNPs, as well as for females in all but four genome-wide significant

SNPs (Supplemental Table 1). The same explanation is likely behind the 64 SNPs discarded for

association between missingness and sex, as 26 of these SNPs have almost no male minor

allele homozygotes and 47 have a p-value for lack of minor allele homozygotes of less than 1E-

6.

To quantify the potential for mis-hybridization of sex chromosome regions to autosomal

probes, we used BLAT (Kent 2002) to find all regions across the genome with high sequence

similarity to autosomal probe sequences on the MEGAEx (BioVU) and UK Axiom/BilEVE (UK

Biobank) genotyping arrays (Methods). We assign each probe sequence to the sex

chromosome region with the highest BLAT score.

The probes for each significantly associated variant have high sequence similarity to a

sex chromosome region (Figure 2, Table 2). In contrast, the majority of probes (79% in UK

Biobank, 89% in BioVU) do not have any detectable similarity (BLAT score < 20) to a sex

chromosome sequence. Compared to the distribution of BLAT scores for probes with a match to

a sex chromosome region, all genome-wide significant variants had BLAT scores greater than

the 99th and 95th percentile for BioVU and UK Biobank respectively (inset Figure 2A, 2B).

Using a stricter criteria to define potential sex chromosome sequence similarity (Methods), we

find that all genome-wide significant variants in BioVU (Supplemental Figure 3A) and six out of

eight genome-wide significant variants in UK Biobank (Supplemental Figure 3B) still have strong

sequence similarity to a sex chromosome region (Table 2). Only 0.57% (4,587 probes) and

3.3% (20,528) of all probes in BioVU and UK Biobank respectively have such a sex

chromosome match (Supplemental Figure 3). The difference in percentage is likely due to the

UK Biobank arrays having longer probe sequences. Probes of genome-wide significant variants

have similar BLAT matching properties as non-significant probes (Supplementary Figure 4) in

UK Biobank and BioVU. Many of the 64 SNPs discarded for between-sex differences in

missingness also demonstrated high sequence similarity to sex chromosome regions

(Supplementary Table 8). Overall, the lack of homozygotes and the high sequence similarity

between significant probes and sex chromosomes strongly suggests that sex-specific

genotyping error is the source of the significant associations rather than sexually antagonistic

selection.

The lack of sex-specific allele frequency differences is not due to being statistically

underpowered

To determine if the lack of significant associations might be a result of being underpowered to

detect plausible effect sizes, we conducted a power analysis (Methods). Based on the large

cohort sizes, we have 95% power to detect a variant with a true allele frequency difference

greater than 2% between the sexes in the BioVU cohort and greater than 0.8% in the UK

Biobank (Figure 3A). A frequency difference of f% caused by sex-specific antagonistic selection

at a locus requires a mortality of roughly $f/2$%, so we should be able to detect segregating

variants with sex-specific mortality effects of at least 0.4% (Supplemental File 9). For

comparison, a cohort of 100 individuals, as used in a previous HapMap study (Lucotte *et al.*

2016), only has 95% power to detect allele frequency differences between the sexes of 38% or

greater (Figure 3B).

DISCUSSION

352
Understanding how sex-specific effects are transmitted by autosomal variation is critical for

understanding how sexual dimorphisms arise and fix in populations. Sexually antagonistic

selection maintains sexual dimorphisms and is predicted to be a pervasive driver of genome

evolution (Rowe *et al.* 2018). Yet, empirically, the genomic signature of this process is not well

characterized. In this study, we sought to identify the extent of one genomic signature of

sexually antagonistic viability selection acting on autosomal variation in human populations.

Capitalizing on two of the largest available biobanks, we performed genome-wide association

tests for genetic sex that failed to identify and replicate any genome-wide significant variants.

On closer inspection, a number of promising genome-wide significant variants were driven by

technical artifacts, most likely due to high sequence similarity to a sex chromosome. We

conclude there is no conclusive signal in these data of sexually antagonistic viability selection

acting on genetic variants at individual loci based on a male-female divergence statistic.

These results stand in contrast to recent male-female F_{ST} studies, that have reported

tens to hundreds of significantly differentiated variants (Lucotte *et al.* 2016; Flanagan and Jones

2017; Wright *et al.* 2018; Dutoit *et al.* 2018; Bissegger *et al.* 2019; Vaux *et al.* 2019). These

studies suggest strong, pervasive sexually antagonistic viability selection acting across the

genomes of various species, which would be puzzling in light of theoretical observations and

simulations indicating that strong allelic divergence between the sexes requires high sex-

specific mortality rates to overcome the homogenizing effect of meiotic segregation occurring

every generation (Kasimatis *et al.* 2019). In contrast to these studies, the sample size of our

study provided statistical power to distinguish true signal of plausible magnitude from stochastic

noise. Additionally, our use of larger sample sizes provided power to detect smaller allelic

divergence between the sexes – within the range predicted to be generated by weak sexually

antagonistic selection. Our results are in line with a previous meta-analysis of sex-specific

common variant differences in humans (Boraska *et al.* 2012), though our direct, replicated

approach with larger sample sizes mitigates against potential confounders across different

studies.

We found strict quality control measures for population structure and multiple testing

essential. In particular, rigorous testing for sequence similarity to the sex chromosomes showed

that all significant SNPs had strong sequence matches. The potential for high sequence

similarity between autosomes and sex chromosomes to generate sex-biased genotyping errors

has been reported previously (Chen *et al.* 2012; 2013). However, the potential for these sex

chromosome artifacts to affect population genetic statistics has not been fully appreciated until

recently (Bissegger *et al.* 2019; Tsai *et al.* 2019) or has only been examined for the Y

chromosome (Lucotte *et al.* 2016). In particular, probe sequences with high sequence similarity

to one the sex chromosomes can lead to skewed allele frequency estimates in a sex-specific

manner due to sequence mis-hybridization and the different sex chromosome content between

females and males. This problem extends beyond SNP-based genotyping to read-based

sequencing data, where inaccurate mapping of reads to an autosome instead of the sex

chromosome could generate a similar skew in allele frequencies. This sex chromosome effect is

potentially very common, and therefore, must be explicitly considered in any sex-specific or sex-

stratified analyses to prevent technical and bioinformatic artifacts from generating false signals.

Participation bias rather than differential mortality can also generate a signal of male-female

divergence (Pirastu *et al.* 2020), though this source of error is not relevant in this study since we

did not find candidate SNPs for sexually antagonistic selection that passed our quality controls.

Such artifacts will be especially problematic in species with new sex chromosomes, poorly

assembled genomes, or rapidly evolving sex chromosome systems. In our case, filtering out

SNPs with large differences in missingness between sexes and/or lack of homozygotes was

sufficient to remove problematic SNPs.

Comparison of sequence similarity and match length for all probes indicates that

thousands of other probes have similarly strong sex chromosome matches as the candidate

variants analyzed here (Supplementary Figure 2). While previous studies have detected similar

hybridization effects (Chen *et al.* 2012; 2013), the extent to which they can skew association

results has not yet been reported on the UKBiobank and BioVU arrays. This high sequence

similarity could suggest that more variants should show false positive signatures of sex-specific

allele frequency differences. However, multiple factors contribute to the potential for mis-

hybridization and inaccurate genotyping. For example, hybridization strength and kinetics are

determined by sequence attributes beyond simple sequence identity, including local GC content

and the potential for DNA secondary structures to form (Zhang *et al.* 2018). Furthermore, the

sequence region matched on the sex chromosome (i.e., pseudo-autosomal versus non-

recombining) also matters. It is also likely that different quality control strategies used on

different genotyping array platforms filter different problematic sites.

Although sexually antagonistic selection is certainly an important selective pressure, we

see no evidence of it generating substantial autosomal allelic divergence between the sexes in

human populations. This strong negative result is unusual, as genome-wide association studies

for most traits on a biobank-scale find significantly associated SNPs. We know that humans

have the opportunity for sexually antagonistic effects, as seen through sex-specific mortality and

disease susceptibility (Morrow 2015; Khramtsova *et al.* 2018). However, randomization of

alleles every generation by meiotic segregation means that a large selective pressure is

required to create a large difference in allele frequencies, and thus, this genetic process makes

it harder to detect the results of sexually antagonistic selection. Furthermore, some sexually

antagonistic variants are not stably polymorphic; we would not detect these because they move

rapidly to fixation (Rowe *et al.* 2018; Kasimatis *et al.* 2019).

Given the confounding factors, technical artifacts, and high sampling variance,

identifying variants with small sex-specific effect sizes is a formidable challenge. We strongly

recommend that future studies avoid simple metrics, like the male-female F_{ST}, and instead

incorporate strict quality filters and control for known confounders into their association tests.

Sexually antagonistic viability selection is not the only action of sex-specific selection nor is

male-female allelic divergence at a single locus the only possible signature of sexual

antagonism. Given the extent of sexual dimorphisms in nature, there are almost surely

autosomal loci subject to sexually antagonistic selection, which may be detectable through other

genomic signatures. However, our work illustrates that the field must reconsider our

assumptions and develop new metrics for identifying signatures of sexual antagonism in the

light of theoretical expectations to understand how this process affects the genome. Such

studies will help us understand the translation of sex across the genotype-phenotype map and

apply this to human health.

Acknowledgements

We thank Locke Rowe and the Phillips and Capra lab groups for their comments on this work,

as well as Clare Bycroft, Adrián Cortés, and Gil McVean for information about UK Biobank

genotyping. A.K. was supported by NIH award R01GM117241. J.A.C. was supported by NIH

award R35GM127087. P.C.P. was supported by NIH award R01GM102511 and

R35GM131838. A.A. is supported by NIGMS of the National Institutes of Health under award

number T32GM007347. This research has been conducted using the UK Biobank Resource

under Application Number 43626. This work was conducted in part using the resources of the

Advanced Computing Center for Research and Education at Vanderbilt University and the

resources of the Research Advanced Computing Services at the University of Oregon. The

content is solely the responsibility of the authors and does not necessarily represent the official

views of the National Institutes of Health.

Contributions

K.R.K. and P.C.P. devised the project. K.R.K., A.A., P.L.R., A.D.K., J.A.C., and P.C.P. designed

the analyses. K.R.K. and A.A. performed analyses. K.R.K., A.A., P.L.R, and J.A.C wrote the

manuscript with the support of the other authors.

The authors declare no competing interests.

REFERENCES

Archer C. R., Recker M., Duffy E., Hosken D. J., 2018 Intralocus sexual conflict can resolve the male-female health-survival paradox. Nat. Commun. **9**: 1–7.

Arnqvist G., Rowe L., 2005 *Sexual conflict*. Princeton University Press, Princeton, NY.

Bissegger M., Laurentino T. G., Roesti M., Berner D., 2019 Widespread intersex differentiation across the stickleback genome - The signature of sexually antagonistic selection? Mol. Ecol. **77**: 1–10.

Boraska V., Jeroncic A., Colonna V., Southam L., Nyholt D. R., *et. al.*, 2012 Genome-wide meta-analysis of common variant differences between men and women. Hum. Mol. Genet. **21**: 4805–4815.

Bycroft C., Freeman C., Petkova D., Band G., Elliott L. T., *et. al.*, 2018 The UK Biobank resource with deep phenotyping and genomic data. Nature **562**: 203–209.

Chang C. C., Chow C. C., Tellier L. C., Vattikuti S., Purcell S. M., *et. al.*, 2015 Second-generation PLINK: rising to the challenge of larger and richer datasets. Gigascience **4**: 2–16.

Chen Y.-A., Choufani S., Grafodatskaya D., Butcher D. T., Ferreira J. C., *et. al.*, 2012 Cross-Reactive DNA Microarray Probes Lead to False Discovery of Autosomal Sex-Associated DNA Methylation. Am. J. Hum. Genet. **91**: 762–764.

Chen Y.-A., Lemire M., Choufani S., Butcher D. T., Grafodatskaya D., *et. al.*, 2013 Discovery of cross-reactive probes and polymorphic CpGs in the Illumina Infinium HumanMethylation450 microarray. Epigenetics **8**: 203–209.

Cheng C., Kirkpatrick M., 2016 Sex-specific selection and sex-biased gene expression in humans and flies. PLoS Genetics **12**: e1006170.

Cortes A., Dendrou C. A., Fugger L., McVean G., 2018 Systematic classification of shared components of genetic risk for common human diseases. bioRxiv: 1–22.

Cox R. M., Calsbeek R., 2009 Sexually antagonistic selection, sexual dimorphism, and the resolution of intralocus sexual conflict. Am. Na.t **173**: 176–187.

Das S., Forer L., Schönherr S., Sidore C., Locke A. E., *et. al.*, 2016 Next-generation genotype imputation service and methods. Nat. Genet. **48**: 1284–1287.

Deegan D. F., Engel N., 2019 Sexual dimorphism in the age of genomics: how, when, where. Front. Cell. Dev. Biol. **7**: 1–7.

Dumitrescu L., Barnes L. L., Thambisetty M., Beecham G., Kunkle B., *et. al.*, 2019 Sex differences in the genetic predictors of Alzheimer's pathology. Brain **142**: 2581–2589.

Dutoit L., Mugal C. F., Bolívar P., Wang M., Nadachowska-Brzyska K., *et. al.*, 2018 Sex-biased gene expression, sexual antagonism and levels of genetic diversity in the collared flycatcher (*Ficedula albicollis*) genome. Mol. Ecol. **27**: 3572–3581.

Farquhar C. M., Bhattacharya S., Repping S., Mastenbroek S., Kamath M. S., *et. al.*, 2019
Female subfertility. Nat. Rev. Dis. Primers. **5**: 1–22.

Ferretti M. T., Iulita M. F., Cavedo E., Chiesa P. A., Schumacher Dimech A., *et. al.*, 2018 Sex
differences in Alzheimer disease - the gateway to precision medicine. Nat. Rev. Neurol. **14**:
457–469.

Flanagan S. P., Jones A. G., 2017 Genome-wide selection components analysis in a fish with
male pregnancy. Evol. **71**: 1096–1105.

Kasimatis K. R., Nelson T. C., Phillips P. C., 2017 Genomic signatures of sexual conflict. J.
Hered. **108**: 780–790.

Kasimatis K. R., Ralph P. L., Phillips P. C., 2019 Limits to genomic divergence under sexually
antagonistic selection. G3 **9**: 3813–3824.

Kent W. J., 2002 BLAT - The BLAST-like alignment tool. Genome Res. **12**: 656–664.

Khramtsova E. A., Davis L. K., Stranger B. E., 2018 The role of sex in the genomics of human
complex traits. Nat. Rev. Genet **62**: 1–190.

Lanfranco F., Kamischke A., Zitzmann M., Nieschlag E., 2004 Klinefelter's syndrome. Lancet
364: 273–283.

Lucotte E. A., Laurent R., Heyer E., Ségurel L., Toupance B., 2016 Detection of allelic
frequency differences between the sexes in humans: a signature of sexually antagonistic
selection. Genome Biol. Evo.l **8**: 1489–1500.

Mank J. E., 2017 Population genetics of sexual conflict in the genomic era. Nat. Rev. Genet. **7**:
1–10.

Morrow E. H., 2015 The evolution of sex differences in disease. Biol. Sex Differ. **6**: 1–7.

Moskvina V., Craddock N., Holmans P., Owen M. J., O'Donovan M. C., 2006 Effects of
differential genotyping error rate on the type I error probability of case-control studies. Hum.
Hered. **61**: 55–64.

Pirastu N., Cordioli M., Nandakumar P., Mignogna G., Abdellaoui A., *et. al.*, 2020 Genetic
analyses identify widespread sex-differential participation bias. bioRxiv: 1–54.

Rice W. R., Chippindale A. K., 2001 Intersexual ontogenetic conflict. J. Evol. Biol. **14**: 685–693.

Rowe L., Chenoweth S. F., Agrawal A. F., 2018 The genomics of sexual conflict. Am. Nat. **192**:
274–286.

Tsai K. L., Evans J. M., Noorai R. E., Starr-Moss A. N., Clark L. A., 2019 Novel Y chromosome
retrocopies in canids revealed through a genome-wide association study for sex. Genes **10**:
320–11.

Vaux F., Rasmuson L. K., Kautzi L. A., Rankin P. S., Blume M. T. O., *et. al.*, 2019 Sex matters:
otolith shape and genomic variation in deacon rockfish (*Sebastes diaconus*). Ecol Evol **27**:
477–21.

Weedon M. N., Jackson L., Harrison J. W., Ruth K. S., Tyrrell J., *et. al.*, 2019 Assessing the
analytical validity of SNP-chips for detecting very rare pathogenic variants: implications for
direct-to-consumer genetic testing. bioRxiv: 696799.

Wright A. E., Fumagalli M., Cooney C. R., Bloch N. I., Vieira F. G., *et. al.*, 2018 Male-biased
gene expression resolves sexual conflict through the evolution of sex-specific genetic
architecture. Evol. Letters **215**: 403–10.

Wright C. F., West B., Tuke M., Jones S. E., Patel K., *et. al.*, 2019 Assessing the pathogenicity,
penetrance, and expressivity of putative disease-causing variants in a population setting.
Am. J. Hum. Genet. **104**: 275–286.

Zhang J. X., Fang J. Z., Duan W., Wu L. R., Zhang A. W., *et. al.*, 2018 Predicting DNA
hybridization kinetics from sequence. Nat. Chem. **10**: 91–98.

Zhao S., Jing W., Samuels D. C., Sheng Q., Shyr Y., *et. al.*, 2018 Strategies for processing and
quality control of Illumina genotyping arrays. Brief Bioinform. **19**: 765–775.
546

547 **Table 1. Genome-wide significant variants in BioVU and UK Biobank cohorts.** Variants

548 passing genome-wide significance (P < 5E-8) in the BioVU or UK Biobank cohorts are reported.

549 Genome-wide significant variants did not replicate across the cohorts. Location is reported in

550 GRch37/hg19 coordinates. Allele refers to the effect allele with which odds ratio (OR) is

551 calculated. Individuals refers to the total number of individuals tested for the variant.

Location (chr:position)	SNP ID	Allele	Max. OR	Individuals	BioVU p-value	UK Biobank p-Value
BioVU Significant SNPs						
3:16652240	rs9870157	T	1.31	61,709	2.82e-83	0.42
7:100351596	rs145369881	T	0.78	60,499	3.25e-08	0.06
7:121147858	rs77638744	A	1.12	61,361	1.52e-10	0.13
13:20119336	rs9508454	C	1.19	61,694	1.26e-31	0.87
14:35761675	rs1048990	G	1.16	61,712	1.94e-20	0.75
UK Biobank Significant SNPs						
1:162075684	rs75745570	T	0.92	471,060	0.66	7.60e-14
4:88457099	rs114928327	T	0.89	413,257	0.30	4.09e-22
10:39006198	rs11598874	T	1.07	478,329	0.44	1.94e-12
11:4515024	rs11032483	T	1.25	482,581	0.99	1.33e-53
11:34104213	rs75212444	T	0.88	482,788	0.20	2.35e-13
12:118926685	rs7298104	T	0.91	475,771	0.93	5.97e-10
19:53535248	rs116890400	A	0.88	485,047	0.51	1.02e-11
21:18068575	rs73196350	A	0.94	479,137	0.76	5.19e-10

553
554

555 **Table 2. Best sex chromosome sequence match for genome-wide significant variant**
556 **probes.** Variants with genome-wide significant associations with genetic sex are reported with
557 GWAS p-value (P-value) and the matched sex chromosome region (Matched Sex
558 Chromosome) with the highest BLAT score (BLAT score). The sequence similarity and length of
559 the matching region (Match Length) are also reported.
560

Dataset	SNP ID	Location (chr:position)	P-value	Matched Sex Chromosome	BLAT Score	Sequence Similarity (%)	Match Length (bp)
BioVU	rs9870157	3: 16652240	2.82E-83	Y: 26964471-26964521	46	96.0	50
BioVU	rs145369881	7: 100351596	3.25E-8	X: 26864979-2685116	39	90.0	50
BioVU	rs77638744	7: 121147858	1.52E-10	Y: 23315613-23316169	45	98.0	50
BioVU	rs9508454	13: 20119336	1.26E-31	Y: 28612640-28612689	46	91.9	50
BioVU	rs1048990	14: 35761675	1.94E-20	Y: 15398460-15398510	46	96.0	50
UKBB	rs75745570	1: 162075684	7.60E-14	X: 121952043-121952114	57	90.2	71
UKBB	rs114928327	4: 88457099	4.09E-22	X: 79084149-79084223	60	93.0	71
UKBB	rs11598874	10: 39006198	1.94E-12	Y: 13568059-13568130	67	97.2	71
UKBB	rs11032483	11: 4515024	1.33E-53	Y: 19070733-19070803	48	84.3	70
UKBB	rs75212444	11: 34104213	2.35E-13	X: 36967486-36967536	46	96.0	50
UKBB	rs7298104	12: 118926685	5.97E-10	Y: 1513524-1513899	43	93.9	59
UKBB	rs116890400	19: 53535248	1.02E-11	X: 38605892-38605963	57	90.2	71
UKBB	rs73196350	21: 18068575	5.19E-10	X: 80780038-80780101	49	88.9	63

561
562

Figure 1

564
565
Figure 1. Genome-wide association tests for genetic sex reveals candidate variants for sexually antagonistic selection. To identify candidate variants for sexually antagonistic selection, we performed genome-wide association tests between females (cases) and males (controls) in two large biobank cohorts: **(A)** BioVU (females = 34,269, males = 27,491) and **(B)** UK Biobank (females = 264,813, males = 223,478). After standard quality control and sex-specific missingness filters (Methods), we identified five variants with genome-wide statistically significant associations (P < 5E-8, solid red line) in BioVU and eight in the UK Biobank. None of the significant variants in BioVU and UK Biobank replicated at genome-wide or nominal significance (P < 0.05) across the two cohorts (Table 1). The probe sequence for each associated variant (except rs11032483) had >90% sequence identity to at least one sequence on a sex chromosome (Table 2). Each point represents one variant. Each variant is colored by whether the best match of its probe sequence to a sex chromosome (according to BLAT score) is on X (pink) or Y (green). If it has no strong match to either sex chromosome it is colored black. The size of each point indicates the degree of sequence similarity.

581   **Figure 2**

582
583

Figure 2. Probes for autosomal variants associated with genetic sex show high sequence

similarity to sex chromosomes. We searched probe sequences used to genotype autosomal

variants in the BioVU (798,051 autosomal probes) and UK Biobank (620,040 autosomal probes)

cohorts for high sequence similarity to sex chromosome regions using BLAT (Methods). (**A**)

More than 80% of BioVU autosomal probes do not have any sequence similarity (BLAT score ≤

20) to a sex chromosome region; these are plotted at 0. Among the 83,083 BioVU probes with

similarity to a sex chromosome sequence (inset), the probes for the variants with genome-wide

significant associations with sex (blue triangles) are all in the tail of the distribution beyond the

99[th] percentile of the BLAT match score. (**B**) Patterns are similar for the UK Biobank probes;

however, a higher fraction (20%, 128,090) have detectable similarity to a sex chromosome,

likely due to their greater length than the BioVU probes.

595
596

597   **Figure 3**

598
599
Figure 3. **Statistical power was sufficient to detect small allelic divergence between the**
sexes. (A) The power to detect different levels of allelic divergence between the sexes was
calculated for the BioVU (blue) and UKBB (green) cohorts. The dashed line shows the 95%
power threshold. **(B)** Statistical power for the analyzed cohorts compared to previous analysis of
human sequences (Lucotte *et al.* 2016) based on approximately 100 individuals per HapMap
population (gray).
606

SUPPLEMENTARY FILES

Supplementary File 1: "TableS1_gwas_significant_hits.xlsx" - Summary statistics for genome-wide significant variants (including those removed for uneven missing rate between the sexes) associated with genetic sex in BioVU and UK Biobank. MALE_HOM1, MALE_HET, and MALE_HOM2 are the counts of genotypes of the minor allele homozygote, heterozygote, and major allele homozygote genotype calls in males, respectively; MALE_MISSING is the number of missing genotypes in males reported by plink. FEM_ prefixes similar columns for females. MISSING_PVAL gives the p-value from Fisher's exact test comparing proportions of missing genotypes between males and females as reported by plink. HOM1_PVAL gives the p-value for a binomial test for the proportion of minor allele homozygotes (of either sex) being equal to the marginal allele frequency squared. OR, STAT and ASSOC_PVAL gives the maximum odds ratio, t-statistic and p-value from logistic regression as described in the text.

Supplementary File 2: "TableS2_HWE_genotype_counts.xlsx" - Genome-wide significant variants associated with genetic sex in BioVU and UK Biobank with Hardy-Weinberg Equilibrium statistics.

Supplementary File 3: "TableS3_raw_blat_xy_hits_bv.tsv" - Raw BLAT results for hits to chromosome X or Y for probes in the MEGAex genotyping array.

Supplementary File 4: "TableS4_raw_blat_xy_hits_ukaxiom.tsv" - Raw BLAT results for hits to chromosome X or Y for probes in the UK Biobank Axiom genotyping array.

Supplementary File 5: "TableS5_raw_blat_xy_hits_ukbil.tsv" - Raw BLAT results for hits to chromosome X or Y for probes in the UK BilEVE genotyping array.

Supplementary File 6: "TableS6_best_blatscore_xy_hit_length_filtered_bv_gwas.tsv" - Best BLAT matches to chromosome X or Y based on highest BLAT score for each probe in the MEGAex genotyping array.

Supplementary File 7: "TableS7_best_blatscore_xy_hit_length_filtered_uk_gwas.tsv" - Best BLAT matches to chromosome X or Y based on highest BLAT score for each probe in the UK Biobank. BLAT results for hits to chromosome X or Y for probes are chosen after pooling across UK Biobank Axiom and UK BilEVE genotyping arrays.

Supplementary File 8: "TableS8_uk_var_w_missingness_best_blatscore_xy.tsv" – Best sex chromosome match based on highest BLAT score for UK Biobank genome-wide significant variants with statistically significant difference in missing rate between females and males.

Supplementary File 9: "FileS9_mortality_selection_derivation.pdf" – Mathematical derivation for estimating the sex-specific mortality cost.

649
650
Supplementary Figure 1. Permutation of genetic sex to generate a null distribution
demonstrates that p-values are well calibrated. We randomly permuted genetic sex and ran
a genome-wide association test between the permuted females and males in the UKBB cohort
100 times. Only those variants with a p-value < 0.01 under the association with the true genetic
sex are considered (n = 8,868 SNPs). **(A)** Q-Q plot of all the permuted variants shows they are
uniformly distributed. **(B)** Permuted variants are uniformly distributed even at very small p-
values.
658

659
660
Supplementary Figure 2. Significantly different variant missingness between females and
males contribute many spurious association in the UK Biobank GWAS for genetic sex.
After running a GWAS for genetic sex in **(A)** BioVU and **(B)** UK Biobank cohorts, we identify five
and 72 variants with genome-wide significant associations (solid red line, P < 5E-8; dashed red
line P < 5E-6) respectively. Variants with a statistically significant difference (p < 0.00001,
Fisher's Exact test) in the missing rate between females and males are colored in red. In the UK
Biobank cohort, 64 genome-wide significant variants also have a statistically significant
difference in the missingness between cases and control, suggesting that these associations
are spurious.
670
671

672
673
674

Supplementary Figure 3. Sequence similarity distribution of probes after applying strict matching criteria to a sex chromosome. To identify probes most likely to mis-hybridize between autosomal and sex chromosome sequences, we filtered those whose best BLAT match met the following criteria in (A) BioVU and (B) UK Biobank: sex chromosome match ≥40 base pairs in length, ≥90% sequence similarity, and overlap of the matching region with the genotyped variant. Out of all autosomal probes, 0.57% and 3.3% met the aforementioned criteria in BioVU and UK Biobank, respectively.

Supplementary Figure 4. Probes of genome-wide significant variants with a match to sex chromosome regions have similar matching properties as non-significant variant when comparing BLAT score and match length to sequence similarity. Using BLAT, we identify array probe sequences with high sequence similarity (≥90%) to a sex chromosome region, have a match length ≥40 base pairs, and overlap or is adjacent on the probe sequence to the variant being genotyped. We plot bivariate kernel density estimates comparing **(A)** BLAT score and **(B)** match length against sequence similarity (y-axis) for BioVU and UK Biobank probe sequences. Darker blue represents areas of higher density. The position of probe sequences for genome-wide significant variants are overlaid as red markers on each plot. Comparing against the densities of the non-significant variants, probes of genome-wide significant variants occur in areas of high density suggesting they have similar matching properties as non-significant probes.