A novel setup for simultaneous two-photon functional imaging and precise spectral and spatial visual stimulation in *Drosophila*

Authors: R. C. Feord[1] & T. J. Wardill[1,2*]

Affiliations:

[1]Physiology, Development & Neuroscience, University of Cambridge, CB2 3EG, UK.

[2]Ecology, Evolution and Behavior, University of Minnesota, St Paul, MN, 55108, USA.

*Correspondence to: twardill@umn.edu

ABSTRACT

Motion vision has been extensively characterised in *Drosophila melanogaster*, but substantially less is known about how flies process colour, or how spectral information affects other visual modalities. To accurately dissect the components of the early visual system responsible for processing colour, we developed a versatile visual stimulation setup to probe combined spatial, temporal and spectral response properties. Using flies expressing neural activity indicators, we tracked visual responses in the medulla to a projected colour stimulus. The introduction of custom bandpass optical filters enables simultaneous two-photon imaging and visual stimulation over a large range of wavelengths without compromising the temporal stimulation rate. With monochromator-produced light, any spectral bandwidth and centre wavelength from 390 to 730 nm can be selected to produce a narrow spectral hue. A specialised screen material scatters each band of light across the visible spectrum equally at all locations of the screen, thus enabling presentation of spatially structured stimuli. We show layer-specific shifts of spectral response properties in the medulla correlating with projection regions of photoreceptor terminals.

INTRODUCTION

The fruit fly, *Drosophila melanogaster*, is a key model for invertebrate vision research[1]. The small diameters of cells in the early visual neuropil have long limited electrophysiological approaches to recording neural activity in the optic lobes, a problem recently overcome with advances in technology and genetic tools that make functional imaging methods possible [2–4]. Though these imaging methods have facilitated extensive characterisation of motion vision in flies [5], how flies process colour information, or how the spectral content of light affects other visual modalities, remains largely unknown.

Photoreceptors in the *Drosophila* eye express different classes of photoreceptive cells, referred to as R1-R8. Within an ommatidium, the six outer photoreceptors (R1-R6) express the broadband rhodopsin (Rh) 1. The more centrally located cells (inner photoreceptors), R7 and R8, express Rh3-6 photopigments, which are narrowly tuned to UV, blue and green bands of the spectrum [6–9]. Additional parameters, notably screening pigments, exert influence on the spectral response properties of the photoreceptor cells [10]. Light-absorbing screening pigment encircling each ommatidium restricts off-axis light from reaching the photoreceptors [10] and yellow-coloured pigment granules in R1-R6 cells contribute to spectral tuning via a dynamic and light-dependent migratory pattern [11,12].

The overall picture of colour vison in *Drosophila* is being pieced together with improved knowledge of photoreceptors, candidate cells involved in circuitry, and colour-guided behaviour [13–19], but much still remains to be characterised. The long-standing dogma advocates that the motion and colour vision processing streams are neatly separated, with R1-R6 providing input to the motion detection pathway and R7-R8 providing input to the colour vision pathway [13,20–22]. More recent evidence, however, demonstrates that signals arising from R7 and R8 improve motion processing in *Drosophila* [23]. This paradigm of combining modalities at an early stage of visual processing has previously been demonstrated as a strategic mechanism for improved perceptual discrimination and retention of salient visual features [24–26].

Commercially available visual displays are tailored to the primate retina. Their three-colour channel configuration provides an excellent experience to the trichromatic user but fall short for visual systems with different photoreceptor sensitivities. Furthermore, the high flicker-fusion frequencies in invertebrates, beyond 120 Hz in *Drosophila melanogaster* [27], outpace the refresh rates of commercial displays. An array of experimental systems for visual stimulation have been developed, but many constraints still limit the investigation of combined colour and motion processing. The production of bands of monochromatic light is commonly achieved via a broadband light source coupled either to a monochromator [8], individual colour filters [28,29], LEDs [30], or more recently LED-based monochromator systems [31]. Such monochromatic light provides full-field stimulation but lacks any spatial structure. Paradigms for spatially patterned stimuli include LCD displays, LED panels or projectors, none of which offer the option of many colours [17,32–34]. In order to probe visual response properties to combined modalities, the integration of spectral and spatial resolution within a stimulation paradigm is essential. In addition, the spectrally broad and high detection sensitivity of two-photon imaging systems restricts the visual stimuli's spectral range and intensity: light applied within the detection range of the microscope will result in an artefact on the acquired image consequently restricting the range of wavelengths available for visual stimulation.

In order to determine the precise contribution of spectral information to visual computations, whether general or colour-specific, we designed a system that offers fine-wavelength resolution across a large portion of the spectrum, that can be calibrated to produce isoluminant stimuli over a biologically-relevant range of intensities and that allows the presentation of spatially- and temporally-structured stimuli. This system presents several advantages not offered in any previous imaging-based setups for characterising physiological responses to colour. These include (1) the delivery of coloured stimuli simultaneous to the acquisition of two-photon images (previous methods rely on the scanner fly-back periods for stimulation [17,34], (2) the ability to arbitrarily chose the centre wavelength and bandwidth), and (3) a rear projection screen which maintains isoluminant hues across the visible spectrum. Using this setup, we characterised intensity-response relationships and spectral response profiles of the pan-neuronally labelled medulla in several genetically modified strains of *Drosophila*, varying in opsin functionality and screening pigment density.

RESULTS

A modified monochromator-projector-microscope system

A visual stimulus was presented to a fly while imaging its neural responses via a two-photon microscope. To produce a range of colours (selectable narrow bands of the spectrum), we modified a projector to use a monochromator as its light source (**Figure 1A**). We introduced custom Semrock bandpass optical filters to the imaging and visual stimulation pathways (**Figure 1B**). These filters allowed for the presentation of a visual stimulus with minimal contamination to the calcium signal read. One filter set, added to the monochromator, blocks the wavelengths of light within the microscope's detection range. The other set of optical filter combination is integrated into the microscope to limit the wavelengths of light entering the detectors and works to reject bleed-through of detectable light wavelengths from the visual stimulus by means of an arrangement of high optical density bandpass filters (further details of the filters in **Supplementary Figure S1**).

Combined spectral and spatial precision

The light produced by the monochromator can be modified for two parameters, wavelength and intensity, providing ample versatility to produce stimuli across the visible spectrum and across several log units of intensity. To test for spectral response independent of brightness, all monochromatic bands of light were calibrated to produce the same radiance with minimal variation over time (**Supplementary Figure S2**). Bands spanned from 385 to 725 nm at approximately 5 nm centre wavelength intervals. Gaps in the transmitted light between 500-540 nm and 610-650 nm (**Supplementary Figure S2**) exist because of the dichroic filters integrated to the monochromator. Furthermore, we sought to ensure minimal variation of projected light across the two-dimensional plane. As the distance between the fly's eye and the screen increases towards the outer edges of the screen, the brightness diminishes accordingly (**Figure 2A**). A more problematic source of variation, however, stems from light scattering. Scattering is an inherent and necessary property of the screen to ensure light diffuses over the array of angles required to reach the fly's eye (**Figure 2A**). Scattering properties of a material, however, are coupled to the wavelength of the light. This creates the challenge of identifying a screen material that scatters light in an equivalent manner across the UV and visible spectrum. We identified a screen fitting these requirements (Da-Lite, Polacoat® Flex Plex Video Vision), ensuring that calibrated isoluminant bands of light at the screen centre retain their flat isoluminant calibration across the screen from 385 to 720 nm (**Figure 2B** and **2C**). A small variation between the UV and red light occurs at the outer edges of the screen. However, by delimiting a circular ellipse (diameter: 400 pixels/38.8 degrees of visual field), with its centre realigned to correspond to maximal brightness (blue dot, **Figure 2B**), the projected stimulus retains spectral constancy (**Figure 2D**).

Proof of concept: spectral response properties in the *Drosophila* medulla

General medulla responses

To probe spectral response properties in the medulla of *Drosophila*, we established intensity-response relationships and spectral response profiles by means of calcium activity indicator imaging in pan-neuronally labelled flies. Transgenic fly strains differed from each other for one or more of the following parameters: screening pigment density (red: high, orange: low), photoreceptor function (intact or Rh1-only) and calcium activity indicator (GCaMP6f: green, RGECO: red, **Figure 3A**).

A set of full-field light pulses of increasing intensity were applied to the fly's eye (**Supplementary Figure S3**) and sigmoid curves fitted to establish intensity-response relationships for UV, blue and

green light (390, 460 and 565 nm centre wavelength respectively, **Figure 3B**). We observed a left
shift of the green intensity-response curve and half-maximum values in orange-eye flies by
comparison with their red counterparts (**Figure 3B** and **3C**), indicative of increased sensitivity to
longer wavelengths of light. Next, we applied a series of light pulses ranging from UV light through to
red light set at approximately 5 nm centre wavelength intervals (**Supplementary Figure S3**) and
plotted normalised spectral response curves (**Figure 3D**). Our results revealed strain-specific
sensitivity to colours: orange-eye flies exhibited a decreased sensitivity in the blue range and an
increased sensitivity in the green range by comparison with their red eye counterparts (**Figures 3D**
and **Supplementary Figure S5**). Coefficient values extracted from fitted intensity response curves
(slope and half maximum, **Supplementary Figure S4**) suggest minor response property discrepancies
between flies expressing GCaMP6f and RGECO (**Supplementary Figure S4**), attributable to
differences in amplitude and decay times [35]. Consequently, we did not pool data from RGECO-
expressing flies with GCaMP6f data for statistical analyses. Nonetheless, this red-emitting indicator
serves the valuable purpose of completing the spectral profile as the red GaAsP is used to record
RGECO signals, thus allowing the GCaMP-restricted green wavelengths of the spectral sweep to be
filled in (**Figure 3E**).
160
161 *Layer-specific responses*
162
Pan neuronal labelling of the medulla reveals clearly discernible and identifiable layer structures
(**Figure 4A**) allowing us to delve into layer-specific responses for the intensity-response relationships
and spectral response curves as above (**Figures 4C** and **Supplementary Figure S6**). Our results
demonstrate that half maximum values for UV, blue and green light exhibit layer-specific variability
between fly strains (**Figure 4C**). Although most layer groupings retain the higher sensitivity of
orange-eye flies to green light, this difference disappears in layers M2 and layers M6-M7. The
decreased sensitivity of orange-eye/Rh1-rescue flies relative to the red eye/wildtype photoreceptor
flies in the UV-blue range of the spectral response curve (**Figure 3D**) is only apparent in layers M6-
M7 and M9-M10. Strikingly, the red eye/Rh1-rescue flies exhibit increased sensitivity to UV, blue and
green light in M2, but this increased sensitivity does not occur in the orange eye/Rh1 rescue flies.
We also see this trend in layer M3, and we attribute the absence of statistical significance to the
lower number of ROIs with successfully fitted intensity-response curves as response size is smaller
than other layers.
176
177
178 **DISCUSSION**
179
We developed a novel experimental setup that enables simultaneous two-photon functional brain
imaging and visual stimulation across the spectrum in *Drosophila*. With this system, a fly can be
presented with over 50 different wavelength bands of the visual spectrum, causing minimal
disruption to the image quality produced by the microscope. Furthermore, our visual stimulation
paradigm provides combined high spatial resolution and high spectral resolution, overcoming the
customary trade-off between the two parameters.
186
The modifications to the optical pathway in both the monochromator and the microscope overcome
the major limitation for spectrally-varied stimuli with two-photon imaging. The study of motion
vision in *Drosophila* is usually carried out using blue light for visual stimulation, which offers an
adequate wavelength range for Rh1-derived visual responses that sits outside of the PMT's
detection range. For the use of spectrally diverse stimuli, a workaround method was developed
whereby the visual stimulus delivery occurs during the two-photon scanner fly-back period [17,18].
This approach, however, is limited by the discontinuous nature of the visual stimulus, which could
introduce aliasing problems with moving patterns, but also reduces the microscope photon

collection time thereby reducing the imaging sensitivity (this particularly affects calcium indicators
with slow temporal responses). The introduction of custom Semrock bandpass filters in our system
results in an extensive range of possible wavelengths for visual stimulation simultaneous to the
acquisition of imaging data by the microscope. A gap in the spectrum corresponding to each PMT's
detection range still exists, but is considerably narrowed, and can be overcome by switching
between neural activity indicators with different emission spectra detected by alternate PMT
detectors.
202
In addition to the combination of simultaneous spectral visual stimulation and two-photon imaging,
we also present a screen material that maintains near-constancy of the brightness of a wavelength
band at all points of the screen from UV through to red light. To our knowledge this, surpasses all
currently published screen materials in this regard, and offers a host of possibilities for
characterisation of combined visual modalities such as motion and colour. The inevitable drop in
radiance and spectral constancy towards the outer edges of the screen can be eliminated by
restricting the stimulus to the central portion of the screen. Alternatively, if some variation of the
brightness and spectral properties of the stimulus is not deemed to impact response properties, use
of the extended screen area provides a greater coverage of the visual field.
212
Our understanding of the spectral quality of visual information transmitted by the different
photoreceptive cells to downstream visual neurons is primarily based on the generation of spectral
sensitivity curves for opsins or electroretinogram measurements of the surface of the eye [8,36]. A
recent study, however, demonstrated that R7 and R8 terminals interact to transform the inner
photoreceptor output to a biphasic sensitivity curve [17], exemplifying the modification of spectral
tuning at the early stages of visual processing. Very little is known about how outputs of different
photoreceptors might be combined by colour-encoding circuits, highlighting the need for high
resolution spectral stimulus capabilities. Our system uses narrow bands of monochromatic light well
suited to our investigation into spectral response properties, but it is entirely possible to modify the
light source to produce multispectral light using a system such as the LED-based monochromator [31].
Methods commonly employed in mammalian vision research, such as the sophisticated combination
of multispectral light to effectively target only one class of photoreceptors [37], are limited in
invertebrates. Further characterisation of the effect of pre-receptoral filtering, such as eye pigment
screening, as well as instability and adaptation properties of visual pigments, is required. With this
information, a smaller, but carefully selected, array of colour channels can be coupled to a display
technology (screen or panel) such as the modified projector system with a five-primary light engine
used in mouse vision research [33] or the arbitrary-spectrum spatial visual stimulator presenting up to
six chromatic channels [34].
231
An ideal spectral stimulation range would extend further into the UV portion of the spectrum to
match the known spectral sensitivities of Rh1, Rh3 and Rh5. Monochromators providing this range of
UV light exist. However, the high-energy photons of the shorter wavelengths are lost by
transmission throughout the optical pathway, with the DLP chip in the projector, in particular,
drastically reducing the UV content of light. The ongoing progress in the development of UV
transmitting projectors (e.g. Texas Instruments) and optics might soon provide an adequate and
affordable solution and extend the capabilities of our system to include UV visual stimulation.
239
The experimental setup we describe here enabled us to record the responses to a range of narrow
bands of light across the spectrum, over a range of intensities spanning several log units of light in
the medulla. We find several modifications of spectral response properties in the summed activity of
the pan-neuronally labelled neuropil between transgenic fly strains, demonstrating the precision,
reliability and sensitivity of our setup. A reduction of screening pigment density results in increased

sensitivity of medulla neurons to longer wavelength of light, consistent with the Goldsmith's
recruitment hypothesis and prior ERG recordings in flies [38–40].
247
Assessment of inter-layer variability revealed a diversity of responses. Most intriguing is the
increased sensitivity of layers M2 and M3 in red eye flies that have lost R7 and R8 function,
suggestive of inhibitory control of postsynaptic neurons in wild type flies by the R8 cells that
terminate in M3 [41]. Several reasons may explain why orange eye flies lacking functional
photoreceptors do not also exhibit increased sensitivity in M2 and M3. Flies homozygous for the
white null allele 1118 are known to have retina degeneration [42] and this may not be rescued with
the mini-white vector expression. This effect could also be attributed to prolonged depolarisation
afterpotential (PDA) [43], more common in white-eye flies [44], where the temporary light insensitivity
of the photoreceptor requires red to shift metarhodopsin back to rhodopsin. This could also explain
the decreased responsiveness to short wavelengths observed in the spectral response profile of this
strain. The decreased sensitivity in the orange-eye flies to UV light in layers M6-M7 could be
attributable to the lack of input from UV-sensitive R7 photoreceptors that project to this layer [41,45].
It is less clear, however, why these flies lose their increased sensitivity to green light uniquely in this
layer structure. If the shift in response properties is caused by the photoreceptor terminals, green
light should inhibit R7 activity [17,18], resulting in decreased sensitivity in the red-eye flies expressing
all functional opsins while maintaining unaffected high sensitivity in the orange-eye strain. Our data
appears to suggest layer-specific shifts in spectral response properties in the medulla corresponding
to projection regions of photoreceptor terminals. We did not detect summation of R7-R8 with R1-R6
signals proposed to drive lobular plate responses reported previously [23] as we found no difference
between wild type and Rh1-only red-eye flies in most layers, however further investigation is
required to establish this with certainty.
269
270 Our setup overcomes many limitations that have hindered the study of colour vision and the
271 integration of visual modalities in *Drosophila*. Going forward, it will serve as a highly versatile tool to
272 further recent breakthroughs in the field [17,18] and unravel the neural circuitry underlying spectral
273 processing.
274
275 **MATERIALS AND METHODS**
276
Fly stocks
Expression of the fluorescent calcium indicators GCaMP6f [3] and RGECO [35] was achieved using the
GAL4/UAS expression systems [46]. We used the pan-neuronal promoter nSyb (Bloomington Stock
Center, 39171) and the following stocks for the red-eye norpA[36] X chromosome mutant [23], white-
eye norpA[36] mutant (Bloomington Stock Center, 52276) and rhodopsin 1 rescue construct for norpA
(Bloomington Stock Center, 9048). Four different fly stocks were used in this study:
283
284 (1) w[+]; P{yw [20XUAS-IVS-GCaMP6f}attP40/+; P{yw [GMR57C10-GAL4}attP2/+ (n=5)
285 (2) w[+]; +; P{yw [GMR57C10-GAL4}attP2, PBac{w, 20xUAS-jRGECO1a-p10}VK5 (n=5)
286 (3) w[+] norpA[36]; UAS-GCaMP6f / P{w[+mC]=ninaE-norpA.W}2; 39171-Gal4/+ (n=4)
287 (4) w[-] norpA[36]; UAS-GCaMP6f / P{w[+mC]=ninaE-norpA.W}2; 39171-Gal4/+ (n=5)
288
Fly stocks were made using standard fly crossing techniques using balancer chromosomes and in the
case of stock 2, recombination was used to bring two insertions onto the same chromosome. Note
that P{yw [20XUAS-IVS-GCaMP6f}attP40 is abbreviated as UAS-GCaMP6f, P{yw [GMR57C10-
GAL4}attP2 is abbreviated as 39171- Gal4. Stock 1 is referred to throughout as red eye/wild type
photoreceptor function/GCaMP. Stock 2 is referred to throughout as red eye/wild type
photoreceptor function/RGECO. Stock 3 is referred to throughout as red eye/Rh1 photoreceptors

only/GCaMP. Stock 4 is referred to throughout as orange eye/Rh1 photoreceptors only/GCaMP.
Females were used for stocks 1 and 2 and males for stocks 3 and 4.
297
Fly Care
All flies were reared at 25-27°C with approximately 60% humidity under a 12:12 light/dark cycle and
fed a standard cornmeal and molasses diet. Flies were collected one day after eclosion using CO2 for
sedation and were transferred into fresh food vials. A prior study [42] cautions against using flies aged
beyond 4-5 days if they carry the w1118 mutation for reasons of retinal degeneration, recordings
were carried out in 4 to 7-day old flies. Slightly older flies were used due to ease of making the
imaging preparation and did not show different results across this age range.
305
Fly preparation
Following cold-anaesthesia (~15 min in a vial placed on ice), the fly was positioned on a sheet of
aluminium foil such that the back of the head capsule protruded through a small cut-out and was
tilted to form an approximate 10° angle with the foil sheet. This configuration allowed the back of
the head to be exposed for imaging, while leaving the majority of the compound eye below the foil
for visual stimulation. Using UV-curing adhesive, the fly was then secured to the sheet with special
care to reduce brain movement by attaching the proboscis and legs to the upper thorax and
immobilising the abdomen. A saline solution (103 mM NaCl, 3 mM KCl, 20 mM BES, 10 mM
trehalose, 20 mM sodium bicarbonate, 1 mM sodium phosphate monobasic, 2 mM CaCl2 and 4 mM
MgCl2, balanced to pH 7.4) was added to create a bath over the back surface of the head. The
cuticle and underlying trachea were then gently removed to expose the optic neuropil. The trachea
extending from the rim of the eye outward over the medulla was removed using forceps to gently
pull it away. A gravity/suction pump was used to circulate saline over the course of the experiment
to prevent build of up of metabolites or other resulting from the dissection. Following the removal
of the cuticle, an interval of 45 minutes, or more, was allowed to ensure stabilisation of neural
activity and dark adaptation of the fly.
322
Visual stimuli
A projector (DepthQ 360 DLP, WXGA resolution) coupled to a screen was modified to use an
independent light source, a monochromator (Cairn Research Optoscan Monochromator). This
allowed us to project selectable narrow bands of the spectrum onto a screen material (Da-Lite,
Polacoat® Flex Plex Video Vision) that minimised wavelength-dependent scattering properties. Only
the central portion of the screen was used, segmented by a circular ellipse of 400 pixels in diameter
corresponding to 38.8 degrees of visual field. By removing the corners and edges, spectral variation
resulting from light scattering is reduced (**Figure 2D**). The projector was used for patterned stimuli
up to 360 Hz frame rate as some fly species, such as *Coenosia attenuata* can see up to 300 Hz, while
Drosophila melanogaster can see just beyond 120 Hz [27]. The light guide, the projector, the screen
and the stage/holder were adjusted to ensure an ideal alignment of the optical pathway and visual
field of the fly. A first adjustment was made to the light guide and projector mirror position to place
the maximal brightness at the screen projection centre with an even drop in optical power towards
the edges of the screen in all directions. Next, the holder was positioned in relation to the screen
such that the holder centre is perpendicular to the projection centre and equidistant to all four
corners. Furthermore, the holder and screen/projector are adjusted such that when the holder is
positioned to be parallel to the screen, the spectrometer samples light directly at the centre of the
screen. A motorised tip/tilt system allowed us to precisely tilt the holder in any direction such that
the central point of the holder where the spectrometer detector (or fly's eye) is positioned remains
in the same point of space but the angle of the detector varies such that it samples light from a
different location of the screen. Radiance spectra were measured using a NIST calibrated Avantes
AvaSpec 2048 Single Channel spectrometer coupled to an Avantes UV-VIS 600 µm fibre (numerical
aperture NA = 0.22, acceptance angle = AA = 25.4° and solid angle SA = 0.1521°).

346
Intensity-Response Relationship
The first set of visual stimuli applied to the fly were used to establish intensity-response
relationships of medulla neurons. The stimulation protocol consisted of three repeated 1-second
light flashes intercalated with 3 seconds of darkness. This motif was repeated for a range of twenty
incremental intensities spanning four log units of light (from 10^8 to 10^{11} photons/s/m^2) to establish
the dynamic range of the visual response. Intensity-response relationships were probed for three
different bands of the spectrum defined by the following centre wavelengths: 385 nm (UV), 440 nm
(blue) and 565 nm (green). These colours were chosen with the intent to match the known spectral
sensitivity peaks of the fruit fly opsins within the limitations of the system, such as the lack of UV
light transmission through the projector and the restriction of most green wavelengths by the filters
to prevent sampling by the GaAsP detectors.
358
Spectral Response
Next, in order to probe spectral response properties, the fly was presented with a spectral sweep of
randomised light flashes of varying centre wavelengths ranging from 390 to 720 nm, all calibrated to
produce the same radiance of 2.63 x 10^9 photons/s/m^2. This intensity was chosen as it sits in the
middle of the intensity-response range across all tested wavelengths and all tested fly genotypes. A
gap in the spectral sweep was introduced spanning 490 to 656 nm and 595 to 700 for the green-
emitting GCaMP and the red-emitting RGECO, respectively, to accommodate the red and green
GaAsP detector sensitivities. Flashes of light were applied for 1 second with a 3-second interval
between each pulse, and the entire spectral sweep was repeated five times with a different colour
randomisation for each repetition. Spacing between centre wavelengths was originally set to be 5
nm increments, as specified by the voltages supplied by the monochromator software. However, an
assessment of the Gaussian curve describing the band of light revealed that the peak of each curve
is, in fact, not aligned with the expected centre wavelength, but shifted in either direction by a
negligible amount. Nonetheless, in the interest of precision, centre wavelengths have been adjusted
in graphs throughout.
374
Further visual stimulus considerations
Daily calibrations were carried out for all stimuli, and a post-calibration check was conducted to
ensure precision of the stimuli. Each experiment consisted of all stimuli types and these were always
presented in the following order: spectral sweep, green-, UV- and blue intensity-response stimuli. A
ten-minute dark adaptation period preceded the five repetitions of the spectral sweep and preceded
each presentation of the intensity-response stimulus.
381
Two-photon imaging
Calcium signals in neurons of the medulla were imaged using a two-photon Bruker (Prairie
Technologies) In Vivo Microscope using GFP and RFP detection channels, with a 20X water
immersion objective (Zeiss, W Plan-Apochromat 20x/1.0 DIC M27; Cat # 421452-9600-000), modified
with the addition of specialised optical bandpass filters (**Figure 1B** and **Supplementary Figure 1S**). An
insight DS+ laser (Newport Spectra-Physics), with 920 nm infrared excitation applied to the sample.
For functional imaging of calcium responses, data was acquired at a 512 x 512-pixel resolution at a
rate of approximately 30 frames per second. Visual stimuli were generated in StimGL (Howard
Hughes Medical Institute) and controlled via Matlab (Mathworks, MA, USA). The start of a visual
stimulus sequence was indicated to the imaging software (PrairieView) by a trigger, which
subsequently initiated the acquisition of images, but also controlled the monochromator and
projector. Images were acquired using the green channel (GaAsP detector 1) for flies expressing
GCaMP and the red channel (GaAsP detector 2) for flies expressing RGECO. A 200X zoom (20X on the
objective and 10X on the software) was used to visualise the fly medulla. This allowed imaging of all
layers of the neuropil from proximal to distal across a number of cartridges (ranging from a half a

dozen to a dozen in different experiments, 0.104 to 0.115 µm/pixel with a field of view of 53 to 59
µm).
399
400 **Analysis**
401
Fluorescent signal extraction
Analyses were conducted using custom-made scripts in Matlab (Mathworks, MA, USA). For region of
interest (ROI) selection, a reference image was used corresponding to averaged frames across the
response period for a blue light pulse. Circular ROIs 12 pixels in diameter were manually positioned
to tile the sections of the 512 x 512-pixel image containing the medulla that are visually
distinguishable from their fluorescence (both baseline and light response). The aim of this selection
process was to assess responses in a given discernible layer without assuming response uniformity
across the layer. Each ROI covers a sufficient number of pixels to ensure a good response could be
extracted whilst limiting noise issues that may arise in larger ROIs. Each imaging sequence from a fly
medulla averaged approximately 150 ROIs that were extracted for analysis. A cross-correlation
analysis [47] was implemented to compensate for motion in the x-y plane resulting from
physiologically driven movement of the visual neuropil (e.g. muscular displacement), vibrations
resulting from equipment, or other. With this method, images across the experiment were realigned
to an initial template image in order to ensure continuity of all features, and consequently continuity
of ROIs from one image to another throughout the stack. A voltage recording paired to the two-
photon image stack was used to segment the experiment into individual stimulus presentations. For
each segment, baseline fluorescence was averaged over a pre-stimulus darkness period (5 frames
prior to stimulus onset). Baseline fluorescence was averaged for each ROI over the pre-stimulus
darkness period. Calcium activity was assessed by changes in fluorescence. These were calculated as
the difference between average fluorescence for a given frame and the baseline fluorescence and
then divided by the baseline for each ROI:
423
424 $$\Delta F = \frac{F - F_0}{F_0}$$
425
426 where ΔF is the change in fluorescence, F is the fluorescent signal and F_0 is the baseline fluorescence.
427 Extracted responses were then averaged across stimulus repeats.
428
Curve fitting parameters and data selection
ΔF traces extracted for individual ROIs were averaged across repetitions of identical stimuli. In the
case of data resulting from the spectral sweep, only three of the five repetitions were retained via a
selection method whereby the two responses furthest from the mean of all five responses for a
stimulus were removed. This methodology was introduced in an effort to reduce noise in the data.
Next, a stimulus response value was calculated by integrating the response over time. The time
segment for response integration was matched to the excitatory phase of the response profile
observed to ensure maximal signal. In practical terms, integral values of the trace were calculated
either over the one-second duration of the stimulus, over the one-second duration post stimulus or
over the time course of both. Sigmoid curves were fitted to the intensity-response data using the
Naka-Rushton equation classically used for electroretinogram data [48]:
440
441 $$\frac{V}{Vmax} = \frac{I^n}{I^n + K^n}$$
442
V is the response and $Vmax$ the maximum amplitude of the response, I is the stimulus intensity, K is
the intensity at the half-maximum response and n is the exponential slope measured at the half-
maximum response. The curves were fitted in a two-step process. First, the intensity-response data

was transformed for each stimulus intensity: $\log\left[\left(\frac{Vmax}{V}\right)-1\right]$. Then a linear regression was fitted to
the transformed data to determine an initial approximation of K and n [49,50]. Next, the estimated K
and n values were used to fit the curve with the above equation. The goodness-of-fit of the curve
was evaluated by means of the mean square error value (MSE) and curves presenting MSE values
superior to 0.05 were discarded. Furthermore, data was retained only if a curve could be fitted to all
three intensity-response stimuli colours (UV, blue and green) within the set MSE parameter
threshold.

A selection process was also applied to spectral sweep data. Spectral response curves were only
retained for a given ROI if the following conditions were fulfilled: (1) all three intensity-response
curves were fit as per the above criteria and (2) the light intensity used for the spectral sweep sits at
least 1/5th below the maximal value of the intensity-response curve for all three colours (to ensure
saturation was not reached at any point of the spectral sweep).

Statistics
All stages of data analysis (image analysis, curve fitting, data representation and statistical analyses)
were conducted using custom scripts which used statistical packages available in Matlab
(Mathworks, MA, USA). Individual statistical tests are reported in figure legends and the results of
post hoc tests, original data points, mean, standard deviation, and standard error of the mean are
reported throughout using notBoxPlot Matlab function (version 1.31.0.0, Rob Cambell).

REFERENCES

1.   Behnia, R. & Desplan, C. Visual circuits in flies: Beginning to see the whole picture. *Current*
Opinion in Neurobiology (2015) doi:10.1016/j.conb.2015.03.010.
2.   Nakai, J., Ohkura, M. & Imoto, K. A high signal-to-noise ca2+ probe composed of a single
green fluorescent protein. *Nat. Biotechnol.* (2001) doi:10.1038/84397.
3.   Chen, T. W. *et al.* Ultrasensitive fluorescent proteins for imaging neuronal activity. *Nature*
(2013) doi:10.1038/nature12354.
4.   Simpson, J. H. & Looger, L. L. Functional imaging and optogenetics in drosophila. *Genetics*
(2018) doi:10.1534/genetics.117.300228.
5.   Borst, A., Haag, J. & Reiff, D. F. Fly Motion Vision. *Annu. Rev. Neurosci.* (2010)
doi:10.1146/annurev-neuro-060909-153155.
6.   Franceschini, N., Kirschfeld, K. & Minke, B. Fluorescence of photoreceptor cells observed in
vivo. *Science (80-.).* (1981) doi:10.1126/science.7268434.
7.   Chou, W. H. *et al.* Patterning of the R7 and R8 photoreceptor cells of Drosophila: Evidence for
induced and default cell-fate specification. *Development* (1999).
8.   Salcedo, E. *et al.* Blue- and green-absorbing visual pigments of Drosophila: Ectopic expression
and physiological characterization of the R8 photoreceptor cell- specific Rh5 and Rh6
rhodopsins. *J. Neurosci.* (1999) doi:10.1523/jneurosci.19-24-10716.1999.
9.   Mazzoni, E. O. *et al.* Iroquois complex genes induce co-expression of rhodopsins in
Drosophila. *PLoS Biol.* (2008) doi:10.1371/journal.pbio.0060097.
10.   Stavenga, D. G., Wehling, M. F. & Belušič, G. Functional interplay of visual, sensitizing and
screening pigments in the eyes of Drosophila and other red-eyed dipteran flies. in *Journal of*
Physiology (2017). doi:10.1113/JP273674.
11.   Kirschfeld, K. & Franceschini, N. Ein Mechanismus zur Steuerung des Lichtflusses in den
Rhabdomeren des Komplexauges von Musca. *Kybernetik* (1969) doi:10.1007/BF00288624.
12.   Franceschini, N. Chromatic Organization and Sexual Dimorphism of the Fly Retinal Mosaic. in
Photoreceptors (1984). doi:10.1007/978-1-4615-9382-9_19.
13.   Gao, S. *et al.* The Neural Substrate of Spectral Preference in Drosophila. *Neuron* (2008)

doi:10.1016/j.neuron.2008.08.010.

14. Schnaitmann, C., Garbers, C., Wachtler, T. & Tanimoto, H. Color discrimination with
broadband photoreceptors. *Curr. Biol.* (2013) doi:10.1016/j.cub.2013.10.037.

15. Karuppudurai, T. *et al.* A Hard-Wired Glutamatergic Circuit Pools and Relays UV Signals to
Mediate Spectral Preference in Drosophila. *Neuron* (2014) doi:10.1016/j.neuron.2013.12.010.

16. Melnattur, K. V. *et al.* Multiple redundant medulla projection neurons mediate color vision in
drosophila. *J. Neurogenet.* (2014) doi:10.3109/01677063.2014.891590.

17. Schnaitmann, C. *et al.* Color Processing in the Early Visual System of Drosophila. *Cell* (2018)
doi:10.1016/j.cell.2017.12.018.

18. Heath, S. L. *et al.* Circuit Mechanisms Underlying Chromatic Encoding in Drosophila
Photoreceptors. *Curr. Biol.* (2020) doi:10.1016/j.cub.2019.11.075.

19. Schnaitmann, C., Pagni, M. & Reiff, D. F. Color vision in insects: insights from Drosophila.
Journal of Comparative Physiology A: Neuroethology, Sensory, Neural, and Behavioral
Physiology (2020) doi:10.1007/s00359-019-01397-3.

20. Rister, J. *et al.* Dissection of the Peripheral Motion Channel in the Visual System of Drosophila
melanogaster. *Neuron* (2007) doi:10.1016/j.neuron.2007.09.014.

21. Yamaguchi, S., Wolf, R., Desplan, C. & Heisenberg, M. Motion vision is independent of color in
Drosophila. *Proc. Natl. Acad. Sci. U. S. A.* (2008) doi:10.1073/pnas.0711484105.

22. Joesch, M., Schnell, B., Raghu, S. V., Reiff, D. F. & Borst, A. ON and off pathways in Drosophila
motion vision. *Nature* (2010) doi:10.1038/nature09545.

23. Wardill, T. J. *et al.* Multiple spectral inputs improve motion discrimination in the drosophila
visual system. *Science (80-.).* (2012) doi:10.1126/science.1215317.

24. Takeuchi, T., De Valois, K. K. & Hardy, J. L. The influence of color on the perception of
luminance motion. *Vision Res.* (2003) doi:10.1016/S0042-6989(03)00086-5.

25. Nishida, S., Watanabe, J., Kuriki, I. & Tokimoto, T. Human Visual System Integrates Color
Signals along a Motion Trajectory. *Curr. Biol.* (2007) doi:10.1016/j.cub.2006.12.041.

26. Gollisch, T. & Meister, M. Eye Smarter than Scientists Believed: Neural Computations in
Circuits of the Retina. *Neuron* (2010) doi:10.1016/j.neuron.2009.12.009.

27. Gonzalez-Bellido, P. T., Wardill, T. J. & Juusola, M. Compound eyes and retinal information
processing in miniature dipteran species match their specific ecological demands. *Proc. Natl.*
Acad. Sci. U. S. A. (2011) doi:10.1073/pnas.1014438108.

28. Kien, J. & Menzel, R. Chromatic properties of interneurons in the optic lobes of the bee - II.
Narrow band and colour opponent neurons. *J. Comp. Physiol. ■ A* (1977)
doi:10.1007/BF00610452.

29. Meinertzhagen, I. A., Menzel, R. & Kahle, G. The identification of spectral receptor types in
the retina and lamina of the dragonfly Sympetrum rubicundulum. *J. Comp. Physiol. □ A* (1983)
doi:10.1007/BF00623906.

30. Little, C. M., Rizzato, A. R., Charbonneau, L., Chapman, T. & Hillier, N. K. Color preference of
the spotted wing Drosophila, Drosophila suzukii. *Sci. Rep.* (2019) doi:10.1038/s41598-019-
52425-w.

31. Belušič, G., Ilić, M., Meglič, A. & Pirih, P. A fast multispectral light synthesiser based on LEDs
and a diffraction grating. *Sci. Rep.* (2016) doi:10.1038/srep32012.

32. Paulk, A. C. *et al.* Selective attention in the honeybee optic lobes precedes behavioral
choices. *Proc. Natl. Acad. Sci. U. S. A.* (2014) doi:10.1073/pnas.1323297111.

33. Allen, A. E., Storchi, R., Martial, F. P., Bedford, R. A. & Lucas, R. J. Melanopsin Contributions to
the Representation of Images in the Early Visual System. *Curr. Biol.* (2017)
doi:10.1016/j.cub.2017.04.046.

34. Franke, K. *et al.* An arbitrary-spectrum spatial visual stimulator for vision research. *Elife*
(2019) doi:10.7554/eLife.48779.

35. Dana, H. *et al.* Sensitive red protein calcium indicators for imaging neural activity. *Elife* (2016)
doi:10.7554/eLife.12727.

36. Stavenga, D. G. On visual pigment templates and the spectral shape of invertebrate
rhodopsins and metarhodopsins. *J. Comp. Physiol. A Neuroethol. Sensory, Neural, Behav.*
Physiol. (2010) doi:10.1007/s00359-010-0568-7.
37. Estévez, O. & Spekreijse, H. The 'silent substitution' method in visual research. *Vision Res.*
(1982) doi:10.1016/0042-6989(82)90104-3.
38. Goldsmith, T. H. Do flies have a red receptor? *J. Gen. Physiol.* (1965)
doi:10.1085/jgp.49.2.265.
39. Stark, W. S. & Wasserman, G. S. Transient and receptor potentials in the electroretinogram of
Drosophila. *Vision Research* (1972) doi:10.1016/0042-6989(72)90049-1.
40. Stark, W. S. & Wasserman, G. S. Wavelength-specific ERG characteristics of pigmented- and
white-eyed strains of Drosophila. *J. Comp. Physiol.* (1974) doi:10.1007/BF00694472.
41. Takemura, S. Y., Lu, Z. & Meinertzhagen, I. A. Synaptic circuits of the Drosophila optic lobe:
The input terminals to the medulla. *J. Comp. Neurol.* (2008) doi:10.1002/cne.21757.
42. Ferreiro, M. J. *et al.* Drosophila melanogaster white mutant w1118 undergo retinal
degeneration. *Front. Neurosci.* (2018) doi:10.3389/fnins.2017.00732.
43. Wright, R. & Cosens, D. Blue-adaptation and Orange-adaptation in white-eyed Drosophila:
Evidence that the prolonged afterpotential is correlated with the amount of M580 in R1-6. *J.*
Comp. Physiol. ■ *A* (1977) doi:10.1007/BF00610456.
44. Belusic, G. ERG in Drosophila. in *Electroretinograms* (2011). doi:10.5772/21747.
45. Takemura, S. Y. *et al.* A visual motion detection circuit suggested by Drosophila
connectomics. *Nature* (2013) doi:10.1038/nature12450.
46. Brand, A. H. & Perrimon, N. Targeted gene expression as a means of altering cell fates and
generating dominant phenotypes. *Development* (1993).
47. Guizar-Sicairos, M., Thurman, S. T. & Fienup, J. R. Efficient subpixel image registration
algorithms. *Opt. Lett.* (2008) doi:10.1364/ol.33.000156.
48. Naka, K. I. & Rushton, W. A. H. An attempt to analyse colour reception by electrophysiology.
J. Physiol. (1966) doi:10.1113/jphysiol.1966.sp008002.
49. Evans, L. S., Peachey, N. S. & Marchese, A. L. Comparison of three methods of estimating the
parameters of the Naka-Rushton equation. *Doc. Ophthalmol.* (1993)
doi:10.1007/BF01203279.
50. McCulloch, K. J., Osorio, D. & Briscoe, A. D. Determination of photoreceptor cell spectral
sensitivity in an insect model from in vivo intracellular recordings. *J. Vis. Exp.* (2016)
doi:10.3791/53829.

581
582
583

Acknowledgements

TJW held a Biotechnology and Biological Sciences Research Council (BBSRC) David Phillips Fellowship
(BB/L024667/1) and RCF was on the BBSRC Doctoral Training Partnership. We thank the University
of Cambridge for supporting the partial funding of our system and space to conduct our research.
We also thank Bruker Fluorescence Microscopy (particularly John Rafter) for their support in
integrating optics to their system, Semrock for assisting with custom fabrication of high quality
optics, Cairn Research (particular Jeremy Graham) for assistance with customisation of the optoscan
monochromator, HHMI Janelia farm for specifications to modify the DepthQ projector to accept a
light guide source, the University of Cambridge fabrication workshops and Milly Sharkey for
assistance with monochromator calibration. We also thank Mathias Wernet for anatomical insights
and the many people who helped with other resources and comments both along the way and for
final manuscripts.

Author information

599
600 T.J.W. developed the initial study idea, designed and built optics system and generated transgenic
601 *Drosophila* lines. R.C.F. conceptualised the monochromator light stimuli and projector screen
602 system. T.J.W. and R.C.F. designed the experiments. R.C.F. conducted the experiments, analysed the
603 data and wrote the manuscript. T.J.W. and R.C.F. finalised the manuscript.
604
605 **Additional information**
606
607 *Competing interests*
608
The authors declare no competing interests

***Figure 1.** A novel setup enabling simultaneous two-photon functional imaging and precise colour*

visual stimulation.

(A) Placement of the fly in the holder with its cuticle removed to expose the optic lobes allows

simultaneous neural activity imaging and visual stimulation. (Ai) A modified projector with a

monochromator light source projects over 50 different bands of the visual spectrum onto a screen.

These colour bands are precisely calibrated using a spectrometer to measure the radiance value of

the visual stimulus. Both the monochromator and the spectrometer are jointly controlled: automated

calibration of the stimulus produces the required light intensity and spectral content via a feedback

loop. A tilted holder allows (Aii) the spectrometer position to be adjusted for measurements of

specific points of the screen, as well as (Aiii) to maximise the coverage of the fly's eye by the screen.

(B) Modifications of the optical pathway in the monochromator and the microscope are necessary to

increase the bandwidths of stimulus wavelengths without detection by the microscope. Light from a

broadband tungsten 150W bulb (lamp) is selectively transmitted through the monochromator via an

input slit (IS), several mirrors (M), a grating (Gr) and an exit slit (ES). Three custom Semrock filters are

*added along this pathway. Filters **1** and **2** are long-pass filters that prevent the transmission of*

*harmonics in the UV range. Filter **1** is moveable and is only used for light above 460 nm. Filter **3***

prevents the transmission of the bands of red and green light that are detected by the microscope. A

*custom Semrock filter (**4**) replaces the dichroic mirror at the start of the imaging pathway, and*

*combined with **5** and **6**, these three filters serve to block the transmission of any light beyond the*
narrow bands of red and green detected by the microscope. A further three filters are combined for
*each GaAsP detector. Filters **7** and **10** block light that falls outside the red and green band ranges.*
*Filters **8** and **11** are long-pass filters. Finally, filters **9** and **12** selectively transmit only specific red and*
green band of light for GaAsP detectors 1 and 2 respectively. Unlike classic quad housing designs,
there are only two detectors instead of four and 200-500 nm and 550-600 nm light along the GaAsp1
and GaAsp2 detector pathways respectively is discarded. Filters denoted with asterisks (*) are a
custom addition to the commercially available version of the system. Filter spectra are reported in
Supplementary Figure S1.
638

Figure 2.** **Optical properties of the screen allow for combined high spatial and high spectral
precision.
*(**A**) Rear-projection of the stimulus from a point light source onto a screen causes variation in light*
intensity across the surface (brightest at the centre and decreased towards the edges). Light
scattering is a necessary property of the screen material to ensure light reaches the fly's eye despite
its angle variance to the screen. However, scattering properties are wavelength-dependent
*perturbing spectral constancy across the screen. (**B**) The percentage variation between minimum and*
maximum radiance curve integrals from an isoluminant calibration of wavelength bands between
385 nm and 730 nm (385 nm light at screen centre approx. 2.35x10^{10}). This variation is measured for
24 different screen locations along the orthogonal and diagonal axes of the screen. A 400-pixel
diameter circle delineated by the dotted line corresponding to 40 degrees of visual field encompasses
a minimally-varying portion of the screen. The blue dot represents a small shift of the circle centre
*relative to the screen centre as maximal brightness is not perfectly centred. (**C**) The percentage*
decrease in intensity from the screen centre plotted as a function of distance for a brighter (385 nm
light at screen centre approx. 2.35x10^{10} and a dimmer (385 nm light at screen centre approx.
1.42x10^{10} calibration. A linear regression fitted to the data shows the two intensities exhibit similar
*trends for variance across the spectrum. (**D**) Example traces of a flat calibration at the screen centre,*
measured along one diagonal of the screen. Minimal variation occurs within the circle specified in
*(**B**).*
658

*Figure 3. **Reduced screening pigment expression biases the fly's spectral sensitivity towards longer wavelengths of light.***

*(**A**) Experiments were performed in four different fly strains expressing pan-neuronal calcium activity indicators – red eye, wildtype photoreceptors and GCaMP6f (RE-WT-GCaMP, bordeaux); red eye, Rh1 only and GCaMP6f (RE-Rh1-GCaMP, red); orange eye, Rh1 only and GCaMP6f (OE-Rh1-GCaMP, orange) and red eye, wildtype photoreceptors and RGECO (RE-WT-RGECO, grey). Intensity-response relationship curves (**B**) were established for three different bands of monochromatic light (UV, blue and green; centre wavelengths 390, 460 and 565 nm respectively) and the half maximum values (**C**) were extracted from fitted curves. Individual data points correspond to the average of ROI responses across an individually discernible layer structure (see methods) for a given fly preparation. Black line = mean, inner grey box = SEM, outer grey box = SD. Significant differences are noted with star values, with the P values from left to right: *P = 0.0105 and ***P = 0.0002 (one-way ANOVA) (**D**) Normalised spectral response profiles, determined by a sweep of equal intensity light pulses across the spectrum. Superimposed means-only traces are represented in the upper panel and the same mean traces are depicted staggered with the addition of standard deviations in the lower panel. Stars indicate statistically significant differences between fly strains for a given centre wavelength, further statistical information is reported in Figure S4 and supplementary table S1. For GCaMP-expressing flies, a gap in the spectral sweep is necessary in the green wavelengths of light corresponding to the microscope's green detection range. (**E**) Mean trace ± SD of the spectral response profile for RGECO-expressing flies. The spectral sweep gap for RGECO-expressing flies sits at the red end of the spectrum.*

Figure 4. Spectral response properties vary across medulla layers.
(A) Pan-neuronal GCaMP labelling of the medulla reveals six distinguishable layer structures (Ai,
schematic; Aii, 2-photon image of baseline fluorescence), each exhibiting discernible cartridge-width
substructures (Aiii). Response profiles of ROIs (segmentation examples shown in Ai) to a blue (440
nm) light pulse are stereotyped within a layer structure. These vary in temporal profile and size of
response across layers M1, M2, M3, M5, M6-M7 and M9-M10 (Aiv). (B) Fly strains expressing pan-
neuronal GCaMP6f– red eye/wild type photoreceptors (RE-WT, bordeaux); red eye/Rh1 only (RE-Rh1,
red) and orange eye/Rh1 only (OE-Rh1, orange). (C) Half maximum values extracted from fitted
intensity-response curves for three different bands of monochromatic light (UV, blue and green;
centre wavelengths 390, 460 and 565 nm respectively) across six layer-groupings of the medulla as
determined in (A). Black line = mean, inner grey box = SEM, outer grey box = SD. Significant
differences are noted with star values, with one-way ANOVA P values reported in supplementary
table S2.

A novel setup for simultaneous two-photon functional imaging and precise spectral and spatial visual stimulation in *Drosophila*

Authors: R. C. Feord[1] & T. J. Wardill[1,2*]

Affiliations:

[1]Physiology, Development & Neuroscience, University of Cambridge, CB2 3EG, UK.

[2]Ecology, Evolution and Behavior, University of Minnesota, St Paul, MN, 55108, USA.

*Correspondence to: twardill@umn.edu

SUPPLEMENTARY INFORMATION

Figure S1. Filter spectra for the monochromator and the microscope.
*Filter spectra for the modified optical pathways in the monochromator and the microscope depicted in **Figure 1B**.*

Figure S2. Detailed optical specifications of the monochromator-projector setup.
(A) Radiance curves for each centre wavelength (385 to 720 nm in approximate 5 nm increments) for monochromator input and exit slits calibrated to produce equal brightness corresponding to a set reference value (here the radiance of the dimmest band of light, 385 nm centre wavelength, for a 30 nm slit width). (B) Calibrated radiance values determined from the area under the curve for each radiance spectra, measured at half hour intervals across the experimental day. (C) Brightness fluctuation of calibrated light across the spectrum reported as the maximum and minimum percentage variance from the mean at any given time (see B), and over the course of the entire day. (D) Power fluctuations for individual centre wavelengths at maximum slit width and calibrated slit widths over the course of 5 minutes. Percentage change corresponds to maximum and minimum change from the mean of measurements taken every second over the course of five minutes.

Figure S3. Intensity-response and spectral sweep stimuli: example responses.
(A) Example fluorescence responses of an ROI to the intensity-response stimulus protocol consisting of light pulses (L.P.) of increasing intensity. Traces for a given intensity represent the mean of the response from all three stimulus repeats. (B) Example fluorescence responses of an ROI to the spectral sweep stimulus protocol consisting of light pulses of varying centre wavelengths. Traces for a given intensity represent the mean of the response from all three stimulus repeats. Response traces are not represented beyond 590 nm for the sake of clarity as no discernible change in fluorescence is detected beyond this point. Note that pulses were presented randomly but have been reordered here in ascending wavelength.

Figure S4. Intensity-response relationship coefficient comparison across fly strains.
(A) Experiments were performed in four different fly strains expressing pan-neuronal calcium activity indicators– red eye, wild type photoreceptors and GCaMP6f (RE-WT-GCaMP, bordeaux); red eye, Rh1 only and GCaMP6f (RE-Rh1-GCaMP, red); orange eye, Rh1 only and GCaMP6f (OE-Rh1-GCaMP, orange) and red eye, wild type photoreceptors and RGECO (RE-WT-RGECO, grey). (B) Slope values extracted from intensity-response relationship curves for three different bands of monochromatic light (UV, blue and green; centre wavelengths 390, 460 and 565 nm respectively) for GCaMP-expressing fly strains. (C) Slope (i) and half maximum (ii) values extracted as in (B) comparing responses in red eye/wild type photoreceptors flies expressing either GCaMP6f or RGECO. Individual data points correspond to the average of ROI responses across a given layer structure (see methods) for a given fly preparation. The curves in (A) are a mean of the fitted curves of individual data points. Black line = mean, inner grey box = SEM, outer grey box = SD. Significant differences are noted with star values, with the P values from top to bottom: ***P = 0.0005 and *P = 0.0415 (one-way ANOVA).

Figure S5. Spectral response profiling – breakdown for individual wavelengths
(A) *Fly strains expressing pan-neuronal GCaMP6f– red eye/wild type photoreceptors (RE-WT, bordeaux); red eye/Rh1 only (RE-Rh1, red) and orange eye/Rh1 only (OE-Rh1, orange).* ***(B)*** *Responses to the individual wavelengths of the spectral sweep in Figure 3D that exhibit significant differences. Black line = mean, inner grey box = SEM, outer grey box = SD. Significant differences are noted with star values, with one-way ANOVA P values reported in table S1.*

Figure S6. Spectral response profiles of layer groupings in the medulla
(*A*) *Schematic of layer structures in the medulla as distinguished from pan-neuronal GCaMP labelling (see Figure 4). (B) Fly strains expressing pan-neuronal GCaMP6f– red eye/wild type photoreceptors (RE-WT, bordeaux); red eye/Rh1 only (RE-Rh1, red) and orange eye/Rh1 only (OE-Rh1, orange). (C) Mean spectral response profiles for layers M1, M2, M3, M5, M6-M7 and M9-M10.*

Table S1. 1-way ANOVA p-values from Figure 3D and Figure S5B

	RE-WT/RE-Rh1	RE-WT/OE-Rh1	RE-Rh1/OE-Rh1
396	1	3.2094e-05	3.2094e-05
398	1	3.2094e-05	3.2094e-05
401	0.8996	3.2094e-05	3.2094e-05
406	0.9999	3.2094e-05	3.2094e-05
411	1	3.2094e-05	3.2094e-05
416	1	3.2094e-05	3.2094e-05
421	1	3.2094e-05	3.2094e-05
425	1	3.2094e-05	3.2094e-05
430	1	3.2094e-05	3.2094e-05
435	1	3.2094e-05	3.2094e-05
440	1	3.2094e-05	3.2094e-05
445	0.0294	0.6676	3.2094e-05
451	1.9732e-04	4.3811e-05	1
569	0.0283	3.2094e-05	3.2094e-05
574	0.0098	3.2094e-05	3.2094e-05
578	0.1550	3.2094e-05	3.2094e-05
583	0.9891	3.2094e-05	3.2094e-05
589	1	3.2094e-05	3.2094e-05
593	1	3.2094e-05	3.2309e-05
598	1	2.5220e-04	0.2610

Table S2. 1-way ANOVA p-values from Figure 4C

		RE-WT/RE-Rh1	RE-WT/OE-Rh1	RE-Rh1/OE-Rh1
M1	UV	1	1	1
	Blue	1	0.0039	0.6421
	Green	1	7.2533e-06	7.2533e-06
M2	UV	7.2533e-06	1	7.2533e-06
	Blue	7.2533e-06	0.7172	7.2533e-06
	Green	7.2533e-06	7.2533e-06	7.2533e-06
M3	UV	0.9962	1	0.9960
	Blue	0.4919	1	0.9999
	Green	0.9957	0.3603	1
M5	UV	1	0.1753	0.9403
	Blue	1	1	1
	Green	1	7.2534e-06	7.2534e-06
M6-M7	UV	1	0.0335	0.9671
	Blue	1	0.2098	0.9167
	Green	1	1	1
M9-M10	UV	1	0.0022	0.8434
	Blue	0.9999	1	0.9981
	Green	1	7.2534e-06	7.2534e-06