(Title): Impact of a candidate vaccine on the dynamics of salmon lice (*Lepeophtheirus salmonis*)

infestation and immune response in Atlantic salmon (*Salmo salar* L.)

(Authors): Jaya Kumari Swain[1,4,†,*], Yamila Carpio[2,†,*], Lill-Heidi Johansen[1], Janet Velazquez[2], Liz

Hernandez[2], Yeny Leal[2], Ajey Kumar[3], Mario Pablo Estrada[2,*]

(Affiliations): [1]Nofima, Tromsø, NO-9291, Norway

[2]Animal Biotechnology Division, Center for Genetic Engineering and Biotechnology, Havana, 10600,

Cuba

[3]Symbiosis Centre for Information Technology, Symbiosis International (Deemed University), Pune,

411057, Maharashtra, India

[4]Current address: Fish Immunology and Vaccinology Research Group, Norwegian College of Fishery

Science, UiT- The Arctic University of Norway, NO-9037 Tromsø, Norway

*Corresponding authors:

E-mail: jaya.k.swain@uit.no

E-mail: yamila.carpio@cigb.edu.cu

E-mail: mario.pablo@cigb.edu.cu

[†] Both authors contributed equally to this work

Abstract

Infection with parasitic copepod salmon louse *Lepeophtheirus salmonis*, represents one of the most important limitations to sustainable Atlantic salmon (*Salmo salar* L.) farming today in the North Atlantic region. The parasite exerts negative impact on health, growth and welfare of farmed fish as well as impact on wild salmonid populations. It is therefore central to ensure continuous low level of salmon lice with the least possible handling of the salmon and drug use. To address this, vaccination is a cost-effective and environmentally friendly control approach. In this study, efficacy of a vaccine candidate, containing a peptide derived from ribosomal protein P0, was validated post infestation with *L. salmonis*, at the lab-scale. The sampling results showed good potential of the vaccine candidate when administered intraperitoneally in the host, in reducing the ectoparasite load, through reduction of adult female lice counts and fecundity and with greater presumptive effect in F1 lice generation. The sampling results correlated well with the differential modulation of pro-inflammatory, Th1, Th2 and T regulatory mediators at the transcript level at different lice stages. Overall, the results supports the effectiveness of the vaccine candidate in controlling salmon lice infestation load. However, further validation is necessary under field conditions.

Keywords: salmon louse; Atlantic salmon; Lepeophtheirus salmonis; vaccine; gene expression; salmon aquaculture; copepod

Introduction

Atlantic salmon (*Salmo salar* L.) is the most important economical species in aquaculture with a production value of 14.7 billion US dollars in 2014 [1] with Norway, Chile and Scotland being the top three salmon producers. However, with increased production the alarm about the impact and number of diseases has also augmented, with parasitic salmon lice emerging as one of the most important in recent years in all the three major salmon-producing countries.

Two lice species represents the primary concern for salmon farming: *Lepeophtheirus salmonis* in the Northern Hemisphere and *Caligus rogercresseyi* in the Southern Hemisphere [2]. In this study, we focused on, a single caligid copepod species *L. salmonis*, which predominates in the North Atlantic, causing year-round infestations of Atlantic salmon housed in marine cages, with concomitant ramifications for fish health in both farmed and wild salmonids as well as for aquaculture economics and sustainability [3]. However, the introduction of more and more salmon farms has significantly increased both the number and density of available susceptible hosts as well as parasite abundance in the coastal waters round the year [4].

Sea lice parasitize salmon during the marine phase of the life cycle, in both wild and farmed salmon, by attaching to their skin often close to gills and fins, feeding on the mucus, epithelial tissues and blood; reproducing on the host and releasing the eggs into the seawater. In seawater, the eggs hatch and develop into planktonic infective stages to parasitize the available host repeatedly [5], thus causing increased parasitic burden on the hosts. If left untreated, this might lead to impaired growth, osmoregulatory stress and open wounds, which can facilitate the entry of other pathogens [5, 6]. The impaired growth and secondary infections cause significant negative animal welfare and economic impact [7]. Moreover, relative to other salmonids, Atlantic salmon have limited ability to resist infection by *L. salmonis* and is therefore highly susceptible to the parasite [8]. The transfer of

sea lice infestation from farmed to wild salmonids is of great concern [9]. Therefore, control of lice

is the first basic priority for the industry, for further sustainable development. As a result, regulatory

affairs departments in salmon producing countries have enforced strict limitations to the allowed

sea lice levels in a farm. These regulations in turn impose treatments through different chemical,

physical and biological methods at frequent intervals and thus directly increase the control-related

costs.

However, pesticide use is significantly reduced now-a-days due to widespread resistance to these

drugs and environmental pollution [10-12]. Increased frequency of treatment methods and salmon

handling by drug-free treatments in the salmon farms has led to challenges with production cost,

handling stress, injury, risk of secondary infection, mortality and thus impaired fish welfare. This has

increased the necessity to develop new and alternative preventive measures [13, 14], which can

document effect on lice and guarantee the fish welfare [15, 16]. To address this, vaccination against

salmon lice could be an important alternative, since it is well-known that fish vaccines have greatly

contributed to reducing the use of drugs (especially antibiotics) against several fish diseases.

Although *L. salmonis* has been an area of research for several decades [3, 5-6, 17, 18], understanding

the mechanisms behind the protection and development of prototype vaccines has been relatively

slow and is still in its infancy. Approaches so far used have met with little or no success due to

challenges in identification of protective antigens and mechanisms. Most strategies for sea lice

vaccines have adopted similar approaches used for vaccines against other ectoparasites in

mammals, for example vaccines against ticks [19].

The present study utilized a previously obtained vaccine candidate based on ribosomal protein P0

for its validation at the laboratory scale [20]. The P0 protein, having a molecular mass between 34-

38 kDa, is highly conserved among eukaryotes [21]. The P0 peptide selected as antigen is located in

a region of low sequence similarity between the lice P0 protein and those of its salmon host, in order

to avoid the induction of tolerance in the parasite or production of auto-antibodies in the salmon

host. In addition, to increase its immunogenicity, promiscuous T-cell epitopes (TCEs) from tetanus

toxin and measles virus were fused to the N-terminus of a 35 amino acids peptide from the

ribosomal P0 protein of *L. salmonis* [20]. These TCEs are universally immunogenic in mammalian

immune systems [22] and reported to improve vaccine efficacy in salmonids [23]. In our previous

study, the candidate vaccine has shown to induce specific IgM response against pP0 compared to

only synthetic pP0, in different teleost species including Atlantic salmon [20].

The purpose of this study was to investigate if the candidate vaccine is able to provide protection,

either in terms of reduced lice count or reduced fecundity or both. Therefore, we targeted to study

the impact of the candidate vaccine at different stages of parasite infestation, post immunization,

under controlled laboratory conditions. Moreover, to highlight the vaccine's further impact on F1

generation hatching efficiency, egg strings collected from the parasitized adult female lice were

hatched and compared. Simultaneously, host-lice interaction studies at the gene level were

performed to explore the immune modulation in response to vaccination for the first time at

different life stages of lice infestation.

Materials and Methods

Antigen purification

The antigen is P0 based protein, which was granted patent for vaccine antigen (Vaccine composition

for controlling ectoparasite infestations PCT/CU2011/000005). P0 based antigen protein was

purified as described previously by Leal et al. 2019 [20]. Briefly, inclusion bodies were obtained by

harvesting induced bacteria cells and centrifugation at 10,000 x g for 10 min at 4°C. The cell pellets

were resuspended in 300 mM NaCl, 10 mM Tris, pH 6 and were disrupted in French Press (Ohtake,

Japan) at 1 200 kgf/cm^2. The disrupted cell suspension was centrifuged at 10,000 x g for 10 min at

4°C and the cell pellet containing the protein was suspended in 1M NaCl, 1% Triton X-100 using

politron Ultra-Turrax T25, IKA WERKE and centrifuge again at 10,000 x g for 10 min at 4°C. This step

was repeated once again and purified inclusion bodies were suspended in PBS (16 mM Na_2HPO_4, 4

mM NaH_2PO_4, 120 mM NaCl, pH 7.4). Protein concentration was determined (Pierce, USA) according

to the manufacturer's instructions and by densitometry scanning of protein gels. Protein samples

were checked by SDS-PAGE on 15% polyacrylamide gels and *western blotting*..

Fish immunization and lice challenge

The experiment was approved by the Norwegian Food Safety Authority, application ID 14617

(https://www.mattilsynet.no/sok/?search=ID+14617) and performed at the Aquaculture Research

Station (Tromsø, Norway). Atlantic salmon (AquaGen standard, average weight 40 ± 6), at a density

of 10 kg/m^3 were kept in circular 500 L tanks supplied with recirculating fresh water for 2 weeks at

an ambient temperature of approximately 10 °C with 24 h light (summer stimuli) for acclimation.

Fish were fed with a commercial pellet diet (Nutra Olympic, Skretting). One hundred and twenty fish

were placed in each tank and three experimental groups were settled. For vaccine formulation,

recombinant antigen protein or PBS control were adjuvanted in Montanide ISA50 V2 (Seppic,

France). at a ratio of 50/50. Montanide ISA50 V2 has been successfully used in salmon as an adjuvant

(24). Our previous observation has also shown that ISA50 works better compared to fish specific

adjuvant ISA763 (unpublished observation).

Immunization and challenge schedule are outlined in Fig 1. The fish were starved for one day before

vaccination. Prior to vaccination, fish were anaesthetized in 0.005% benzocaine. First immunization

was performed as follows: each fish in the control group (Group 1) received 0.05 mL of PBS

emulsified in adjuvant by intraperitoneal (ip) injection; second group (Group 2) received ip injection

at a dose of 1 µg/gram body weight (gbw) of the recombinant antigen protein emulsified in

adjuvant; and third group (Group 3), received ip injection at 1 µg/gbw of the recombinant antigen

protein emulsified in adjuvant plus bath immunized with recombinant antigen protein as inclusion

bodies (200 µg/L) for 1 hour (120 fish in 200 L aerated static bath), immediately after ip injection.

Fifteen days post immunization, fish were transferred to seawater. After 37 days post first

immunization, a booster dose was given to each fish (average weight 60 ± 10 g) at a similar dose per

gram of body weight as first vaccination. Each experimental group were kept in duplicate.

Throughout the experiment, the following experimental conditions were maintained: Temperature:

10°C; Light: 24 h; Oxygen level at outlet: ~80-90 %; Salinity: 34-35 ppt.

***In vivo* lice challenge:** After one month of booster dose, 90 fish (average weight 94 ± 16 g) from

each group were bath challenged with infective copepodids of *L. salmonis* (Oslo/Gulen strain from

Norwegian Institute of Marine Research, IMR). The groups were bath challenged in separate tanks

for one hour with stopped water supply. Each tank received approx. 3150 copepodids to have an

average distribution of about 35 copepodids/fish. Post challenge each group was distributed into

triplicate tanks with 30 fish per tank. The parasitized fish were kept in seawater with a salinity of

34.5‰, oxygen level: 80/90 % and at a temperature of approximately 10 °C, until the salmon lice

reached desired developmental stage i.e at matured adult stage when females have developed egg

strings.

Sampling and lice counting

To evaluate at which developmental life stage of lice the vaccine was effective, counting of lice on

10 parasitized fish per tank i.e. 30 fish per group, were performed at 17 days post infestation (dpi)

(chalimus), 28 dpi (pre-adult) and 50 dpi at mature adult stage with first reproductive egg strings

(Fig 1). At 50 dpi i.e at the adult stage, both egg string number and egg string length per female were

noted, for all the immunized groups and compared. Fish were given an overdose of anesthesia

(0.01% benzocaine) before counting. To avoid counting error of detached lice due to anesthesia and

handling, counting of chalimus at 17 dpi, pre-adults at 28 dpi and adults at 50 dpi on individual

parasitized fish were performed under water in a white tray. All empty trays were checked for lice.

At pre-challenge sampling points, fish were killed using 0.01% benzocaine prior to measurement of

length and weight as well as collection of different tissues (skin, spleen and head kidney) for gene

expression study. Further, spleen, head kidney and skin tissues were sampled aseptically from 18

fish per group (6 fish/tank). Tissue samples were immediately transferred to RNA later (Ambion)

and kept at 4 °C overnight and then stored at -20°C. Overall sampling time-points as outlined in Fig

1 were at 0 (prior to 1st vaccination), 69 (31 days post booster) days post vaccination or 0 day

challenge, and 17 (chalimus), 28 (pre-adult) and 50 (adult) dpi.

Incubation of collected egg strings for F1 generation hatching

To analyze the effect of vaccine on F1 generation copepodids production, the first reproductive egg

strings, obtained from gravid females at 50 dpi were incubated in well-aerated filtered seawater.

This was to determine the effect, vaccine candidate had on hatching efficiency of the F1 generation

copepodids. Fifty egg strings (sampled from the first reproductive event at 50 dpi) from each

experimental group were incubated in 5 parallel aerated flow-through incubators (containing 500

mL filtered seawater/incubator at ~10 °C) for 8 days, to study the hatching success to F1 generation

copepodids. First visual observation was done on day eight post incubation and final counting was

performed at day ten. Copepodid density was estimated by taking 10 mL water samples from each

replicate and counting of copepodid was performed using dissecting microscope. This observation

was repeated four times for each replicate.

Vaccine efficacy

The overall efficacy (in percentage) of the candidate vaccine was calculated using lice count data collected from different dpi, including female lice fecundity parameters and F1 generation copepodid count compared to control group, using a similar approach as used to assess vaccine efficacy for candidate vaccines against ticks [25]:

Vaccine efficacy (%) = 100 x [1-(NCh x NPA x NF x NM x FE x NE x LE x CC)]

NCh: number of chalimus in vaccinated group/ number of chalimus in control group

NPA: number of pre-adults in vaccinated group/number of pre-adults in control group

NF: number of adult females in vaccinated group/number of adult females in control group

NM: number of males in vaccinated group/number of males in control group

FE: number of females with eggs in vaccinated group/number of females with eggs in control group

NE: number of egg strings in vaccinated group/number of egg strings in control group

LE: length of egg strings in vaccinated group/length of egg strings in control group

CC: F1 generation copepodids count from vaccinated group/F1 generation copepodids count in control group

Observation of side effects

Usually, adjuvanted vaccines may cause inflammation, granuloma and pigmentation at the site of injection. To check the side effects of the candidate vaccine having the Montanide ISA50 as an adjuvant, visual scoring and analysis of the vaccine side effects resulting in adhesion was performed using the Speilberg scoring method according to the criteria detailed by Midtlyng et al. 1996 [26]. A separate score for pigmentation for each fish was assigned according to the table in Fig 4B. Fish weight and length were registered and the condition factor (K) was calculated according to Barnham and Baxter, 1998 [27].

Gene expression studies

All organs from the sampled fish, kept in RNA-later (Ambion, Austin, TX, USA) were subsequently

processed for RNA isolation. Total RNA was extracted by MagMAX™-96 Total RNA Isolation Kit

(Invitrogen), including turbo DNase treatment (Invitrogen) according to manufacturer's instruction.

Analysis of gene expression by Real-time PCR (QPCR) was performed in duplicates with a

QuantStudio 5 Real-Time PCR System (Applied Biosystems) using SYBR Green (Applied Biosystems)

in 384 well plates. For each mRNA, gene expression was normalized to the geometric mean of the

3 house-keeping genes (EF-1a, 18S and β-actin) in each sample and fold change was calculated

according to Pfaffl method [28] using the primer efficiency (E). Primer sequences used for gene

expression studies are listed in Table 1.

Table 1. Primer sequences used for the real-time PCR analysis. (*) indicates reference genes used in

this study for normalization.

GENE TARGET	NAME	ACCESSION No.	FORWARD (5-3')	REVERSE (5-3')	AMPLI-CON
Immuno-globulin M (secretory)	IgMs	BT060420	CTACAAGAGGGAGACCGGAG	AGGGTCACCGTATTATCACTAGTTT	90
Immuno-globulin T	IgT	GQ907004	CAACACTGACTGGAACAACAAGGT	CGTCAGCGGTTCTGTTTTGGA	97
Tumor necrosis factor alpha1	TNFα1	AY929385	ACTGGCAACGATGCAGGACAA	GCGGTAAGATTAGGATTGTATTCACCCTCT	144
Interleukin 1 beta	IL-1β	AY617117	GCTGGAGAGTGCTGTGGAAGAAC	CGTAGACAGGTTCAAATGCACTTTGTG	220
Interferon gamma	IFN-γ	AY795563	GATGGGCTGGATGACTTTAGGATG	CCTCCGCTCACTGTCCTCAAA	166
Interleukin-4/13A	IL-4/13A	EG837625	CCACCACAAAATGCAAGGAGTTCT	CCTGGTTGTCTTGGCTCTTCAC	147
Cluster of Differentiation 4	CD4	EU585750	CGGAAGCGAGGGATATAAATGGTG	GGCATCATCACCCGCTGTCT	215
Cluster of Differentiation 8 alpha	CD8α	AY693393	GACAACAACAACCACCACGACTACAC	GCATCGTTTCGTTCTTATCCGGTT	211
Matrix metallo-proteinase-9	MMP-9	AGKD01108865	TGGAGAGAACTACTGGAGGCTGGA	CCGACAGAAGTAGATGTGGCCCTT	142
Interleukin 8	IL-8	HM162835	TCCTGACCATTACTGAGGGGATGA	AGCGCTGACATCCAGACAAATCTC	200
Interleukin 10	IL-10	EF165028	CTGTTGGACGAAGGCATTCTAC	GTGGTTGTTCTGCGTTCTGTTG	129
Interleukin 22	IL-22	DW572073	GGCCCGAGTCAGCAGAGACCT	CTCCTCCATCCCGGCCAACTTC	106
Beta actin*	β-actin	BT059604	CAGCCCTCCTTCCTCGGTAT	CGTCACACTTCATGATGGAGTTG	72

Elongation factor 1-α*	EF1α	AF498320	CAAGGATATCCGTCGTGGCA	ACAGCGAAACGACCAAGAGG	327
18 S ribosomal RNA*	18 S rRNA	AJ427629	TGTGCCGCTAGAGGTGAAATT	CGAACCTCCGACTTTCGTTCT	101

Statistical analysis

The results were analyzed and expressed as mean ± standard deviation (SD) unless otherwise stated.

Statistical analysis was performed and graphs were made using the Prism 6.01 software for

Windows (GraphPad software, San Diego, CA, USA). Experimental groups were conducted in

triplicates. Prior to data analysis, outliers were identified and removed from subsequent analyses.

Normal distribution was assessed using D'Agostino & Pearson omnibus normality test. Multiple

comparison were performed using analysis of variance (ANOVA) or Kruskal Wallis test depending on

the normal distribution and equal variance of the data followed by Tukey or Dunn's Multiple

Comparison *post hoc* tests. P-values < 0.05 were considered statistically significant. Two-way

hierarchical clustering analysis heat map and dendrogram of relative gene expression data and

experimental groups were generated in R language using ComplexHeatmap package by Gu, Z et al.

2016 [29]. In the Principal component analyses, "FactoMineR" package of the R statistical software

(v3.6.2) was used to calculate the principal components and visualizations were constructed using

"factoextra" package. Ellipses in the PCA graph are confidence ellipses with a confidence level of

0.95 and the centroids represent the center of the mass of the points per group.

Results

Impact of the vaccine candidate post lice infestation

Immunization and challenge schedule were performed as outlined in Fig 1. Post infestation, lice

from infected salmon were counted at 17, 28 and 50 dpi, corresponding to different developmental

stages: chalimus, pre-adult (PA) and adult stage (Fig 2A). At 17 dpi, mean number of chalimus (± SD)

attached per fish was 20.00 (± 8.08), 25.17 (± 10.02) and 23.70 (± 12.41) for group 1, group 2 and

group 3, respectively. No significant differences among groups were detected at 17 dpi. At 28 dpi,

mean PA count per fish was 12.83 (± 6.29), 12.73 (± 5.13) and 17.77 (± 7.28) for control, group 2 and

3, respectively, where group 3 showed more PA per fish as compared to groups 1 and 2 (P<0.01).

Finally, at 50 dpi, mean infestation rate of adult lice per fish was reduced to 5.13 (± 2.94), 4.06 (±

2.53) and 5.50 (± 2.63) for group 1, 2 and 3, respectively, and compared to control, group 2 showed

an overall reduction tendency of 21 % (Table 2), although not significant. Moreover, development

rates of *L. salmonis* throughout the experiment was nearly identical between the immunized and

the control group.

Table 2: Effect of vaccination on salmon lice infestation following adult stage of lice.

Experimental groups	Number of Fish (n)	Reduction of adult lice number	Reduction of adult females	Reduction of gravid females with eggs	Reduction of egg string length (mm)	Reduction in F1 copepodids
Group 2	30	21%	40 % * (*P< 0.02*)	42% *(*P< 0.03*)	5%	23%
Group 3	30	-7%	5%	12%	6%* (*P<0.02*)	4%

244
shows significant difference with respect to control group. Group2 is ip vaccinated group; Group 3

is ip vaccinated + bath immunization group. The percentage values were derived compared to control

from the data shown in Fig 2 and 3.

Although there were no great differences between the total lice counts per fish on the immunized

and control fish regardless of louse life stages, statistically fewer adult female lice (40 % reduction),

fewer number of female lice with eggs (42 % reduction) and total number of egg strings per fish

were present on group 2 immunized fish compared to control. However, no differences were

observed in group 3 compared to control (Figs 2 B, D, and E; Table 2). During the 50 dpi sampling,

all egg strings were collected from the gravid females. Most of the gravid females had two egg

strings and less numbers were found in group 2 compared to group 1 (P<0.05) (Fig 2F). No

differences among the groups were observed in the female numbers with one egg string per fish (Fig 2G). Furthermore, gravid female lice removed from the immunized fish showed shorter egg string length compared to the control group, of which group 3 had significant reduction of 6 % (P<0.05) (Fig 2H; Table 2). The results mentioned above clearly showed reduced number of eggs produced by females in group 2 (42 % reduction, P<0.03) and thus supports significantly reduced fecundity in terms of reduced egg string data and less gravid females in group 2 (Table 2). Overall, lice-induced damage on the parasitized fish was low and no wounds were visually observed on any of the experimental fish. Furthermore, there was no evidence of any secondary infections either on the surface or in internal organs of the infected fish.

Impact on hatching efficiency of F1 generation copepodids

Post egg string measurement, a total of 50 egg strings from each experimental group were divided into 5 replicates (10 egg strings per replicate), for the F1 incubation experiment. Fig 2I shows the leftover egg strings (from the total egg strings collected from gravid females at 50 dpi), after the removal of 50 egg strings for the hatching experiment. During incubation, hatching of the egg strings were followed in each group to check if the reduced female fecundity of group 2 in F0 generation had any consequences in the early F1 generation. Subsequently, the F1 copepodids were observed on day 8 and counted on day 10 post-incubation, and data were analyzed. At day 8, the hatching success of egg strings removed from lice on the immunized group were delayed and reduced, especially in group 2, compared to the control group (Fig 3A). This correlates well with the reduced fecundity in the F0 generation of group 2 gravid females. However, the counting at day 10 showed a reduction of 23 and 4 % infective copepodids in the vaccinated group 2 and 3, respectively (Fig 3B; Table 2). The percentage reduction of copepodids on day 10 was not high, as expected based on observation made on day 8 (Fig 3A). This was due to some unseen or technical problem occurring

during the weekend, resulting in some unexpected mortality of the copepodids before counting on

day 10. The experiment was not possible to repeat due to limited time and resources available.

Overall, the results from lice counting and analysis of different parameters at different lice stages

post infestation, showed that the vaccine efficacy of group 2 was the best among the groups with

an efficacy of 86 %. However, group 3 efficacy was negative compared to control since some of the

parameters were lower than the control group. The terminology "vaccine efficacy" used here should

not be interpreted as protection obtained. This is the overall vaccine effects based on different

parameters studied, as described in the materials and methods section.

Side effects analysis

Fish weight, length and condition factor (K) were analyzed at all sampling points. Immunized fish

had less weight and length post lice challenge as compared to the control group (Fig 4A). The

condition factor was acceptable (1.2) and it was the same for all the groups at different sampling

times (Fig 4A). For salmonids, K values usually fall in the range 0.8 to 2.0 [27].

Moreover, side effects of the candidate vaccine having the Montanide ISA50 V2 adjuvant were

analyzed using Speilberg and pigmentation scoring at 50 dpi. Speilberg scoring at 50 dpi showed

that the control group had an average score of 2.0 compared to group 2 and 3, which showed an

average score of 2.8, i.e below 3, which is in an acceptable range for adhesion (Fig 4B).

Pigmentation score was significantly less in the immunized groups compared to the control group,

as shown in Fig 4B. Pigmentation was observed only on the epithelial lining and not in muscle or

tissue within the peritoneum.

Simultaneously, individual fish checked for vaccine depots had vaccine residues, which were

encapsulated by connective tissue as small pockets. The injection site was checked for redness and

lesions and looked normal in all the fish.

Effect of vaccination combined with *L. salmonis* infestation on tissue specific gene expression

Gene expression of pro-inflammatory mediators (TNF-α, IL-1β, IL-8); Th17 and regulatory mediators (IL-22, IL-10); Th1 and Th2 mediators (IFN-γ, IL-4/13A); immunoglobulin genes and cellular markers (IgM, IgT, CD4, CD8α); and tissue remodeling gene matrix metalloprotease 9 (MMP9), were studied to evaluate the immune response of vaccinated fish to salmon lice infestation at different stages of their life-cycle, compared to control fish. Both anterior kidney and spleen, the main immune organs in teleost fish, were used to evaluate systemic responses and skin was used to evaluate the local immune response to salmon lice infestation.

Global assessment: Heat map and hierarchical clustering

To obtain an overview of the expression profiles of the different groups tested at different sampling points corresponding to different lice stages, heat map were constructed with hierarchical clustering. Hierarchical clustering of all the genes studied, identified three clusters representing a differential clustered expression pattern with respect to spleen tissue (Fig 5A). Hierarchical clustering of the experimental groups at different sampling time points pre and post infestation (Fig 5B), also identified 3 clusters for all the tissues studied, showing differences in gene expression under different lice infection stages and treatment groups. Gene expression cluster comparison showed that the pro-inflammatory cytokines, T-regulatory mediators, Th1 and Th2 mediators and T cell surface markers were strongly clustered. A clear pattern of different upregulated gene clusters were visible in different tissues, showing highly upregulated cluster of pro-inflammatory cytokines genes in spleen, highly upregulated regulatory cytokine genes in head kidney and mixed upregulated gene expression of Th1, Th2, T reg, IgM and IL-8 in skin. These results showed that, apart from lymphoid organs, local response played a major role during the host-parasite interaction in later

stages post infestation i.e 28 dpi in the vaccinated groups (group 2 and 3). On the other hand,

column-wise comparison based on different sampling time-points, within respective groups,

showed strong clusters with respect to substantial gene upregulation at 28 dpi in vaccinated groups

(group 2 and 3) in skin, at 17 and 50 dpi (group 3), and 28 dpi (group 2) in spleen and at 17 dpi (group

2 and 3) as well as 28 dpi (group 2) in head kidney. Consequently, evaluating the two-way

hierarchical clustering analysis for all the tissues, vaccinated group 2 at 28 dpi showed the highest

number of upregulated genes compared to the control group. However, vaccinated group 3 showed

higher number of upregulated genes at 17 dpi in spleen and head kidney and at 28 dpi in skin. Heat

map with two-way clustering of genes studied in the individual tissue is given in S1 Fig.

Principal component analysis (PCA)

We performed exploratory data analyses using principal component analysis (PCA) in all the tissues

studied. The PCA analysis of the expression profile of the 12 selected genes in skin samples at

different time points post lice infestation (Fig 6) showed that samples taken at the early stages of

infection [0 day challenge (69 d) and 17 dpi] in vaccinated and control groups were very similar and

with low variability. Consecutive samples (28 and 50 dpi) displayed an increasing deviation along

the principal component 1 (PC1) that contributed to most (78.5 %) of the observed variation.

Samples taken at 28 and 50 dpi formed clearly distinct clusters, and variability among individual

sampling points within groups increased with infection time. Moreover, 28 dpi in vaccinated group

2 contributes to maximum percentage variation (~43%) in PC1 (Fig 6A and C). All the 12 genes

studied showed significant (P< 0.05) contribution in PC1 (Fig 6C) and in addition, IgT expression

showed significant contribution in PC2 where 50 dpi in group 3 had maximum contribution. For head

kidney and spleen, PC1 component contributed to 67.9 and 64.3 % variation, respectively at 17 and

28 dpi in both the vaccinated groups (S2 and S3 Figs). Similar to skin, in head kidney also 28 dpi in

vaccinated group 2 contributes to maximum percentage variation (~37%) in PC1. On the contrary,

in spleen 17 dpi vaccinated group 2 showed maximum contribution of ~34 % followed by 28 dpi

from vaccinated group 2 (~19 %) (S2C and S3C Figs). All genes showed significant contribution in PC1

expect for IL-10 and MMP9 in head kidney and CD8α in spleen (S2D and S3D Figs). This shows that

vaccination together with lice infestation has significant effect on the overall gene expression profile

with more significant contribution at 28 dpi in group 2 than group 3.

Detailed assessment by individual gene expression analysis

The results from the overview of gene expression profiles and the exploratory data analyses clearly

showed changes related to different sampling points post lice infestation and vaccination groups.

Therefore, we proceeded to study these changes in detail, to further characterize gene expression

levels. The overview of the relative gene expression of all the genes analyzed in this study is

graphically represented in S4 Fig. The gene expression results in spleen (S4A Fig), showed that the

pro-inflammatory cytokines TNF-α, IL-1β and chemokine IL-8 were significantly up-regulated

starting from pre-challenge (69 d) and this trend was maintained until 50 dpi in the immunized

groups compared to the control. The same results were obtained for metalloproteinase 9 (MMP-9)

except at 28 dpi in spleen (S4A Fig). For IL-1β, there was also significant expression in head kidney

across all sampling time-points except 17 dpi in both vaccinated groups. On the other hand, in skin

IL-8 expression was up-regulated only at 28 dpi in group 2 and at both 17 dpi and 28 dpi in group 3

(S4A Fig). However, IL-1β and TNF-α in skin was downregulated in infected salmon at 50 dpi in both

the vaccinated group, respectively compared to control except at day 69 (pre-challenge), where

TNF-α was significantly upregulated in group 3.

The gene expression results also showed that both IgM and IgT transcript levels were significantly

upregulated in the vaccinated groups compared to control in all the tissues and sampling time-

points studied (S4B Fig) with exception of IgT in head kidney at 69 day post vaccination and 17 dpi.

Both genes followed almost similar pattern of expression in different groups and sampling time-

points, suggesting its important role in host-parasite interaction. IgT transcription in skin was up-regulated earlier, at 69d (0 day infestation) in group 3.

On the other hand, the activation of T-cell related genes: CD4, CD8α, IL-4/13A and IFN-γ showed significantly higher expression levels in the spleen at 50 dpi (S4B Fig). This trend was also seen at 28 dpi in the head kidney showing the activation of T-cell mediated immunity and the involvement of Th1/Th2 response. Significant decreasing trends of expression levels were also found in these genes at other sampling points showing different patterns of regulation depending on sampling time or experimental groups. For example, cytotoxic T cell marker, CD8α transcript was downregulated compared to the control group in most of the sampling points other than the time-points mentioned above.

In addition, T-cell mediated Th17 and regulatory cytokines, IL-22 and IL-10 shared a common trend of gene expression (S4B Fig) without any specific significant up-regulation except group 3 at 50 dpi in spleen. They were significantly downregulated in group 2 fish at 17 dpi in spleen and in head kidney at 28 dpi, whereas in group 3 fish at 28 dpi only in spleen.

Discussion

The importance of Atlantic salmon in aquaculture and its susceptibility to infection with *L. salmonis* has led researchers to investigate efficient non-medicinal, cost effective and eco-friendly measures to control the sea-lice load through the possibility of vaccine development. While development of a vaccine against *L. salmonis* is still on its way, developing a better understanding of host-pathogen interaction and its modulation in relation to vaccine candidate will provide a lead, to further understand the vaccine efficacy. In the current study, we used a vaccine based on the peptide of 35 amino acids from the ribosomal P0 protein of *L. salmonis* fused to the C-terminal of TCE's from tetanus toxin and measles virus positioned in tandem and previously tested for better antibody

response [20]. Normally, housekeeping proteins are highly conserved among species and the

development of a vaccine candidate based on housekeeping proteins such as P0 ribosomal protein

is very challenging due to its high degree of identity between the P0 sequence of the vertebrate

host and the ectoparasite. Consequently, the peptide P0 used as a vaccine candidate in this study

was selected from the less conserved region between the *L. salmonis* and salmon.

According to the sampling results in this study, initially an overall average of about 23 attached lice

at the chalimus stage were recorded from each fish sampled at 17 dpi and by the end of the

experiment, these numbers reduced to about 5 adult lice per fish. The total number of lice attached

at a particular developmental stage post infestation did not vary statistically between the

immunized or control groups of fish, although there was a tendency of reduction at the adult lice

stage in ip vaccinated group 2. However, significant impact on gravid female lice count and its

reproductive efficacy with delayed hatching and reduced trend of copepodids count in F1

generation was documented also in group 2 compared to control group . This showed that the

major effect of vaccine was apparent in the adult female lice and its fecundity. A similar impact on

female's *R. B. microplus* population was seen after challenge when a 20 aa P0 peptide derived from

Rhipicephalus ticks conjugated to KLH was used to immunize cattle [25]. They reported decrease in

female's yield and weight as well as decrease in egg mass and eggs hatched compared to only KLH

injected group. Similar results have also been reported using sea lice whole extract or lice protein

involved in midgut function and blood digestion, as a vaccine in Atlantic salmon, resulting in fewer

oviparous female lice and lower fecundity [24, 30]. Based on our results, it was expected that a

reduction in parasite fecundity due to vaccination will have an exponential reduction effect on the

overall lice population and thus lice load on the host at later generations, and consequently will

warrant a reduction in chemical or drug free treatments to control lice.

Analogous to the overall efficacy calculated for the pP0 antigen against *R. B. microplus* ticks as well

as other authors [31-33], we extrapolated a similar formula to our experiment for estimating vaccine

efficacy by using the results obtained from lice count post infestation and other sampling results.

Vaccinated group 2 obtained an overall vaccine efficacy of 86% whereas group 3 showed negative,

thus exploring the potential effectiveness of the vaccine candidate through ip method only

However, further in-depth work has to be done. Thus, this method showed that a careful analysis

of formulas, to evaluate vaccine efficacy and its application against salmon lice, could improve the

understanding of these vaccines in teleost. Moreover, lice count as a proxy for resistance had been

questioned, since individual lice counts vary between trials and certain immune genes are affected

negatively by increasing number of lice [34]. Therefore, large number of experimental animals must

be used in these types of immunization and infection trials and treatment efficacy parameters other

than lice count should be considered.

In Atlantic salmon, normally IgM transcripts are most abundant followed by IgT, especially in spleen

and head kidney [35]. In the present study, the increase in relative expression level of IgM and IgT

in spleen, head kidney and skin in vaccinated groups, indicated their important role in systemic and

mucosal immune response in the context of copepodid infestation. In agreement with these results,

Tadiso et al. observed 10-fold increases in IgT expression in the skin from infested Atlantic salmon

and up-regulation of IgT and IgM in spleen and skin two weeks post lice infection [35], but until now,

IgM and IgT responses observed in Atlantic salmon have not been associated with protection against

copepodid infestation. The role of antibodies in protection against copepodid infestation in teleost

has not been fully explored and needs further understanding. For future studies, it will be of greater

importance to measure antigen-specific IgM in serum and in in mucus by ELISA, to understand their

role in protection and crosstalk during salmon lice infestation, post vaccination.

To understand the underlying immune mechanism, we assessed transcriptomic responses at systemic and local level in immunized salmon focusing on mid and late response post infestation. The results showed substantial increase in relative expression of pro-inflammatory mediators (IL-1β, TNF-α, IL-8) at the systemic level (spleen and to some extent in head kidney). This is in line with the sustained response of systemic pro-inflammatory cytokines seen in the more resistant species such as the pink salmon throughout the infection and even after rejection in these fish [36]. Barker et al. (2019) also obtained similar results with significantly higher levels of IL-1β expression at 17 dpi with sea lice [37]. The same pattern of expression held true when investigating tissue repair enzyme MMP 9 gene expression that was used as an indicator to evaluate the wound healing response of the fish to sea lice infestation. The increase in MMP 9, had been suggested by several groups as a possible mechanism for sea lice resistance in Atlantic salmon [34, 37, 38]. In addition, induced high IL-8 transcript levels in skin and spleen post vaccination (69 d) has been implicated as an inducer of neutrophil migration and antibody secreting cells locally. Furthermore, it can also be speculated that elevated systemic expression of inflammatory and T regulatory mediators, pre and post lice challenge in the vaccinated fish compared to only adjuvant control, might have been involved in local expression of IgM and IgT transcript. Moreover, early upregulation of immunoglobulin like genes in spleen, head kidney and skin, in addition to panels of immune genes, indicates a rapid activation of the systemic as well as local anti-parasitic response to some extent, which is in accordance with the results obtained by Skugor et al. (2008) [39]. This demonstrates a facilitated cross talk between immune genes in vaccinated group pre and post infection.

On the other hand, the pro-inflammatory response in skin post infestation appeared to be at the basal level compared to adjuvant control, except for IL-8, which was significantly upregulated at 17 dpi and 28 dpi of sampling for both vaccination types. It is possible that by the time systemic inflammatory response was mounted, the cytokine expression had already returned to its basal level

in skin. Microarray experiment looking at the effects of early stage *L. salmonis* attachment showed

that the local expression in skin decreased at early time points from 5 dpi, although the systemic

response in the spleen remained throughout the study period [35]. As the earliest samples for gene

expression in our study was taken at 17 dpi, it is possible that early transient increase of

inflammatory cytokines in the skin was missed. Another possible explanation can be the sampling

of skin from the standard area of the fish (near the dorsal fin and above the lateral line), regardless

of louse attachment. Therefore, if the cutaneous inflammatory response is directed exclusively at

the site of attachment, it would not have been targeted by the standardized skin sampling,

especially if infection intensity was not as dramatic as those reported previously [35]. Matrix

metalloprotease plays a role in the reconstruction process of the extracellular matrix during wound

healing. In sea lice infested salmon, the slow repair of extracellular matrix is in parallel with stable

up-regulation of MMP9 and MMP13 at the damaged sites, and whose excessive activity may

contribute to the development of chronic wounds [39]. Here, absence of MMP-9 stimulation in skin

could suggest less damage to the host with no chronic wound and subsequently less tissue repair

required. This was confirmed by no visible damages to the skin during the experiment.

Despite that, immersion bath stimulates immune response, mainly in mucosal tissues such as skin

[40]. The intraperitoneal injection of the candidate vaccine plus immersion bath with inclusion

bodies received by group 3 was ineffective in terms of vaccine efficacy, although some immune

parameters were improved. This could be the effect of an exacerbated immune priming. The

response to other parasites have often been described in terms of Th1/Th2 dichotomy, but recent

studies have shown that host-pathogen interactions are more complex. A T cell effector subset

Th17, characterized by the production of IL-17 and IL-22, were identified along with signature

cytokines for regulatory T cell subset (T reg), being inhibitory IL-10 and/or TGF-β. Th1, Th2 and Th17

reciprocally regulates the development and function of each other, while Treg cells suppress all

three subsets [35, 39, 41]. The regulatory cytokines control inflammation and thus protect against

immunopathology, but in doing so they reduce the effectiveness of immune mechanisms

responsible for the expulsion of the parasites. Here, pro-inflammatory response in skin might be

regulated by IL-10, IL4 and IL-22 at 28 dpi of the immunized salmon but requires further study to

match with the earlier time points than 17dpi. This is somewhat in accordance with the results

obtained in resistant coho salmon (*Oncorhynchus kisutch*), although at an earlier time-point up to

72 hours [42]. We observed down regulation of IL-22 and IL-10 in spleen of salmon at chalimus

stage of infection and an increase in IL-1β, TNF-α and IL-8 at subsequent pre-adult stage in group 2.

In group 3, down regulation of IL-22 and IL-10 was seen in spleen at pre-adult stage of infection (28

dpi), which in turn is related to the increase observed in pro-inflammatory cytokines at adult stage

(50 dpi). These differences in the regulation of inflammation could explain the differences found in

the results between different vaccination methods i.e group 2 and group 3. This can be possibly a

consequence of the *E. coli* proteins present in inclusion bodies of the bath vaccine given in group 3.

Further studies targeting more immunological markers could clarify the mechanisms responsible for

the differences between the two groups.

Previous studies have shown that the pathological effects of sea lice become especially profound

for the host fish when they reach free-ranging stage (colloquially 'mobile') on the host compared to

attached chalimus stage [43]. This explores the important strategy the host should develop to avoid

damage on the skin through early free-ranging pre-adult lice interaction and develop resistance

against it. The use of hierarchical clustering heat map and PCA analysis in this study showed a clear

overview of the gene expression in different tissues across the groups at different time-points post

infection and the way the genes were regulated in the vaccinated and the non-vaccinated group.

Most of the genes were highly to moderately upregulated at 28 dpi in only ip vaccinated, group 2,

when the infestation was at the mobile stage (pre-adult), while they were upregulated to some

extent in the ip plus bath vaccinated group 3 at both chalimus and pre-adult phase. In addition, in

only ip vaccinated group 2, differential gene expression, cluster analysis and principal component

analysis also showed the dynamics of T-cell response as mixed Th1/Th2/T17/Treg at the pre-adult

lice stage of infestation. This reflects the importance and potential of the gene modulation strategy

of only ip vaccinated group compared to ip plus bath vaccinated group , against the early mobile lice

stage, for effective vaccine efficacy at the later adult stage and that correlated well with the adult

female lice count and fecundity data documented in the ip injected vaccine group.

Taken together, our result provided new insight into the potential of the candidate vaccine in

reducing salmon lice load and its effect on host-parasite interaction with minimal side-effects. The

cumulative sampling results showed an overall vaccine efficacy of 86 % in the ip injected vaccine

group with an expected larger impact on F1 parasite generation by reduced re-infection loads via

fewer females and decreased fecundity. In addition, the results revealed the priming of immune

response post vaccination and pre-challenge, leading to simultaneous involvement of both systemic

and local immunity during the salmon lice interaction for vaccinated fish, at the mobile lice stages.

These findings provided valuable leads for the effectiveness of the P0 antigen as a vaccine candidate

against salmon lice (*L. salmonis*). However, long-term challenge trials and studies of re-infection

post vaccination is necessary to fully understand and explore the protection potential of candidate

vaccine and underlying molecular mechanism of protection at the gene level. Another aspect to

have in mind is that in experimental challenge conditions, the infestation load is usually very high

(i.e in this validation study: 35 copepodids per fish) and is far higher compared to the natural

conditions in the field. Therefore, performing a challenge experiment under field conditions will be

the next step for further evaluation of the vaccine efficacy in controlling salmon lice infestation.

Acknowledgements

We would like to thank staffs at Aquaculture Research Station in Tromsø for assistance in fish
maintenance, copepodid production, performing lice challenge and lice counting. We also thank Dr.
Trilochan Swain for the valuable suggestions during the development of project concept and
manuscript preparation.

Author Contributions

541 Conceptualization, J.K.S., Y.C., and M.P.E.; Investigation, J.K.S., Y.C., L.H.J., L.H., J.K., Y.L.; validation
542 and visualization, J.K.S., and Y.C.; resources, J.K.S., Y.C., and M.P.E.; data curation and formal
543 analysis, J.K.S., Y.C. and, A.K; writing—original draft preparation, J.K.S., and Y.C.; writing—review
544 and editing, J.K.S., Y.C., L.H.J., A.K., and M.P.E.; project administration, funding acquisition and
545 supervision, J.K.S. All authors have read and agreed to the published version of the manuscript.

Funding

547 This work was funded by Norwegian Seafood Research Fund (FHF), grant number 901461
548 (https://www.fhf.no/prosjekter/prosjektbasen/901461/).

Conflicts of Interest

550 The authors have declared that no competing interests exist. The funders had no role in study
551 design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

1. Food and Agriculture Organization of the United Nations (2016). The State of World Fisheries
and Aquaculture 2016. Rome: Contributing to food security and nutrition for all.
2. Johnson SC, Treasurer JW, Bravo S, Nagasawa K, Kabata Z. A review of the impact of parasitic
copepods on marine aquaculture. Zool. Stud. 2004; 43: 229–243.

3. Pike AW, Wadsworth SL. "Sealice on Salmonids: Their Biology and Control." Advances in

Parasitology. 1999; 44:233–337.

4. Ugelvik MS, Skorping A, Moberg O, Mennerat A. Evolution of virulence under intensive farming:

salmon lice increase skin lesions and reduce host growth in salmon farms. J. Evol. Biol. 2017;

30(6):1136-42.

5. Costello MJ. "Ecology of Sea Lice Parasitic on Farmed and Wild Fish." Trends in Parasitology.

2006; 22(10):475–83.

6. Tully O, Nolan DT. "A Review of the Population Biology and Host-parasite Interactions of the Sea

Louse *Lepeophtheirus salmonis* (Copepoda : Caligidae)." Parasitology. 2002; 124:S165–S182.

7. Frazer LN, Morton A, Krkosek M. Critical thresholds in sea lice epidemics: evidence, sensitivity

and subcritical estimation. Proc. Biol. Sci. 2012; 279: 1950–1958.

8. Fast MD. Fish immune responses to parasitic copepod (namely sea lice) infection. Dev. Comp.

Immunol. 2014; 43:300–312.

9. Provan F, Jensen LB, Uleberg KE, Larssen E, Rajalahti T, Mullins J, et al. Proteomic analysis of

epidermal mucus from sea lice–infected Atlantic salmon, Salmo salar L. J. Fish Dis. 2013;

36(3):311-21.

10. Lees F, Baillie M, Gettinby G, Revie CW. The Efficacy of Emamectin Benzoate against Infestations

of *Lepeophtheirus salmonis* on Farmed Atlantic Salmon (*Salmo salar* L) in Scotland, 2002–2006.

PLoS ONE 2008; 3, e1549. (doi:10.1371/journal.pone.0001549).

11. Jones PG, Hammell KL, Gettinby G, Revie CW. Detection of emamectin benzoate tolerance

emergence in different life stages of sea lice, *Lepeophtheirus salmonis*, on farmed Atlantic

salmon, Salmo salar L. J. Fish Dis. 2013; 36: 209–220.

12. Aaen SM, Helgesen KO, Bakke MJ, Kaur K, Horsberg TE. Drug resistance in sea lice: a threat to

salmonid aquaculture. Trends Parasitol. 2015; 31: 72–81.

13. Igboeli OO, Burka JF, Fast MD. *Lepeophtheirus salmonis*: a persisting challenge for salmon

aquaculture. Anim. Front. 2013; 4:22–32.

14. McNair CM. Ectoparasites of medical and veterinary importance: drug resistance and the need

for alternative control methods. J. Pharm. Pharmacol. 2015; 67:351–363.

15. Liu Y, Bjelland HV. Estimating costs of sea lice control strategy in Norway. Prev. Vet. Med. 2014;

117: 469–477. (doi:10.1016/j.prevetmed.2014.08.018).

16. Brooks KM. Considerations in developing an integrated pest management programme for

control of sea lice on farmed salmon in Pacific Canada. J. Fish Dis. 2009; 32:59–73.

(doi:10.1111/j.1365-2761. 2008.01013.x).

17. Johnson SC, Treasurer JW, Bravo S, Nagasawa K, Kabata Z. A Review of the Impact of Parasitic

Copepods on Marine Aquaculture. Zool. Studies 2004; 43(2):229–43.

18. Boxaspen K. "A Review of the Biology and Genetics of Sea Lice." ICES Journal of Marine Science

2006; 63(7):1304–16.

19. Raynard RS, Bricknell IR, Billingsley PF, Nisbet AJ, Vigneau A, Sommerville C. Development of

vaccines against sea lice. Pest Management Science. 2002;58(6):569-75.

20. Leal Y, Velazquez J, Hernandez L, Swain JK, Rodríguez AR, Martínez R, et al. Promiscuous T cell

epitopes boosts specific IgM immune response against a P0 peptide antigen from sea lice in

different teleost species. Fish. Shellfish Immunol. 2019; 92:322-30.

21. Goswami A, Chatterjee S, Sharma S. Cloning of a ribosomal phosphoprotein P0 gene homologue

from *Plasmodium falciparum*, Mol. Biochem. Parasitol., 1996;82:117–120.

22. Panina-Bordignon P, Tan A, Termijtelen A, Demotz S, Corradin G, Lanzavecchia A. Universally

immunogenic T cell epitopes: promiscuous binding to human MHC class II and promiscuous

recognition by T cells. Eur J Immunol 1989; 19:2237–42.

23. Kuzyk MA, Burian J, Machander D, Dolhaine D, Cameron S, Thornton JC, Kay WW. An efficacious

recombinant subunit vaccine against the salmonid rickettsial pathogen *Piscirickettsia salmonis*.

Vaccine 2001; 19:2337-2344.

24. Contreras M, Karlsen M, Villar M, Olsen RH, Leknes, LM et al. Vaccination with Ectoparasite

Proteins Involved in Midgut Function and Blood Digestion Reduces Salmon Louse Infestations.

Vaccines, 2020; 8: 32. Doi:10.3390/vaccines8010032.

25. Rodríguez-Mallon A, Encinosa PE, Méndez-Pérez L, Bello Y, Rodríguez-Fernández R, Garay H,

Cabrales A et al. High efficacy of a 20 amino acid peptide of the acidic ribosomal protein P0

against the cattle tick, *Rhipicephalus microplus*. Ticks and Tick-borne Dis. 2015; 6: 530-537.

26. Midtlyng PJ, Reitan LJ, Speilberg L. Experimental studies on the efficacy and side-effects of

intraperitoneal vaccination of Atlantic salmon (*Salmo salar* L.) against furunculosis. Fish Shellfish

Immunol. 1996; 6:335-350.

27. Barnham Ch, Baxter A. Condition Factor, K, for Salmonid Fish. Fisheries notes. March 1998

FN0005, ISSN 1440-2254.

28. Pfaffl MVA. New mathematical model for relative quantification in real-time RT-PCR. Nucleic

Acids Res. 2001; 29, e45.

29. Gu Z, Eils R, Schlesner M. Complex heatmaps reveal patterns and correlations in

multidimensional genomic data. Bioinformatics 2016; 32(18):2847-2849.

30. Grayson TH, John RJ, Wadsworth S, Greaves K, Cox D, Roper J, et al. Immunization of Atlantic

salmon against the salmon louse: identification of antigens and effects on louse fecundity.

1995;47:85-94.

31. de la Fuente J, Rodriguez M, Montero C, Redondo M, Garcia-Garcia JC, Mendez L et al.

Vaccination against ticks (*Boophilus spp.*): the experiencewith the Bm86-based vaccine Gavac.

Genet. Anal. 1999; 15:143–148.

32. Hajdusek O, Almazan C, Loosova G, Villar M, Canales M, Grubhoffer L et al. Characterization of

ferritin 2 for the control of tick infestations. Vaccine 2010; 28:2993–2998.

33. Almazan C, Kocan KM, Bergman DK, Garcia-Garcia JC, Blouin EF, de la Fuente J. Identification of

protective antigens for the control of Ixodes scapularis infestations using cDNA expression

library immunization. Vaccine. 2003; 21:1492–1501.

34. Holm H, Santi N, Kjoglum S, Perisic N, Skugor S, Evensen O. Difference in skin immune responses

to infection with salmon louse (*Lepeophtheirus salmonis*) in Atlantic salmon (*Salmo salar L.*) of

families selected for resistance and susceptibility. Fish Shellfish Immunol. 2015;42(2):384–94.

35. Tadiso TM, Krasnov A, Skugor S, Afanasyev S, Hordvik I, Nilsen F. Gene expression analyses of

immune responses in Atlantic salmon during early stages of infection by salmon louse

(*Lepeophtheirus salmonis*) revealed biphasic responses coinciding with the copepod-chalimus

transition. BMC Genomics. 2001; 12: 141.

36. Jones SRM, Fast MD, Johnson SC, Groman DB. Differential rejection of salmon lice by pink and

chum salmon: disease consequences and expression of proinflammatory genes. Dis. Aquatic

Org. 2007; 75:229–238.

37. Barker SE, Bricknell IR, Covello J, Purcell S, Fast MD, Wolters W, et al. Sea lice, *Lepeophtheirus*

salmonis (Krøyer 1837), infected Atlantic salmon (*Salmo salar* L.) are more susceptible to

infectious salmon anemia virus. PLOS ONE. 2019;14(1): e0209178.

38. Skugor S, Holm HJ, Bjelland AK, Pino J, Evensen O, Krasnov A, et al. Nutrigenomic effects of

glucosinolates on liver, muscle and distal kidney in parasitefree and salmon louse infected

Atlantic salmon. Parasites & Vectors. 2016; 9.

39. Skugor S, Glover KA, Nilsen F, Krasnov A. Local and systemic gene expression responses of

Atlantic salmon (*Salmo salar* L.) to infection with the salmon louse (*Lepeophtheirus salmonis*).

BMC Genomics. 2008; 9:18.

40. Mweemba Munang'andu H, Mutoloki S, Evensen O. A review of the immunological mechanisms

following mucosal vaccination of finfish. Front. Immunol. 2015; 6:427. doi:

10.3389/fimmu.2015.00427.

41. Belkaid Y, Blank RB, Suffia I. Natural regulatory T cells and parasites: a common quest for host

homeostasis. Immunol. Rev. 2006;212:287-300.

42. Braden LM, Koop BF, Jones SR. Signatures of resistance to *Lepeophtheirus salmonis* include a

TH2-type response at the louse-salmon interface. Dev. Comp. Immunol. 2015; 48(1):178-91.

https://doi.org/10.1016/j.dci.2014.09.015

43. Revie C, Dill L, Finstad B, Todd CD. Sea Lice Working Group Report. NINA Special Report 2009;

39: 1– 17.

Figure Captions

Fig 1. Experimental Outline. Experimental design depicting immunization, challenge, post challenge

schedule along with sampling time-points. Booster timeline of vaccination is shown by darker arrow.

Fig 2. Efficacy of TT-P0 vaccine on lice counts and fecundity of adult female lice post infestation.

(A) Bar graph showing average lice count / fish for different immunized groups at different lice

stages post infestation (dpi): chalimus (17 dpi), pre-adult (28 dpi) and adults (50 dpi).

Adult lice on the experimental fish were sampled at 50 days post infestation (dpi). The sampled lice

were counted for total number of males, females and female's fecundity parameters per fish. Data

showing, (B) Female numbers, (C) Male numbers, (D) Females with eggs, (E) Number of egg strings

per fish, (F) Females with two egg strings, (G) Females with one egg string, (H) Egg string length, for

different groups per fish (n=30) at 50 dpi. Data shown as mean + SD. A Mann-Whitney test was

performed due to unequal variances to compare vaccinated groups (Group 2 or 3) with control

(Group 1). Asterisk indicates statistically significant differences compared to control group (*P<0.05,

** *P*<0.01). (I) Photograph of leftover egg strings (after removal of 50 egg strings for F1 generation

hatching experiment) to represent the visual number of total egg strings in different groups. Group

details: Group 1 is control group; Group2 received ip injection of adjuvant emulsified vaccine

antigen; Group 3 received ip injection of adjuvant emulsified vaccine antigen + bath immunization.

Fig 3. TT-P0 vaccine's effects on F1 generation hatching and copepodid number.

(A) Observation noted on day 8 for hatching efficiency and visual health status of the hatched

copepodids and (B) Total number of copepodids counted on day 10 post incubation of egg strings.

The bar shows the mean value + SD in 5 replicate tanks for each experimental group. Group details:

Group 1 is control group; Group2 received ip injection of adjuvant emulsified vaccine antigen; Group

3 received ip injection of adjuvant emulsified vaccine antigen + bath immunization.

Fig 4. Side effects of TT-P0 vaccine post immunization.

(A) Growth (weight and length) and condition factor of the fish post immunization and challenge at

different sampling points: Pre-immunization (0 d), 69d post immunization (69 d) and at different

days post infestation (dpi) based on the different lice stages during infection: 17 dpi (chalimus), 28

dpi (pre-adult) and 50 dpi (adult). (B) Visual scoring and analysis of the vaccine side effects resulting

in adhesion (left panel) and pigmentation (right panel) near the vaccination site. Data are shown as

the mean + SD of the parameters under analysis (n = 30). Based on normal distribution test, one-

way ANOVA or Kruskal Wallis test was done followed by Tukey or Dunn's Multiple Comparison.

Asterisk (*) indicates statistical difference *(*P* <0.05), **(*P* <0.01), ***(*P*<0.001) between the groups

with respect to control (Group1). Group details: Group 1 is control group; Group2 received ip

injection of adjuvant emulsified vaccine antigen; Group 3 received ip injection of adjuvant emulsified

vaccine antigen + bath immunization.

Fig 5. Hierarchical clustering analysis heat map and dendrogram of relative gene expression data

over different sampling time points pre and post infection within the vaccinated groups, and for

three different tissues. (A) indicates the pattern of gene expression across different groups and

tissues. It also shows one-way clustering of differentially expressed genes on the right with respect

to spleen whereas (B) shows two way hierarchical clustering of genes on the right and group wise

sampling time-points on the top. Differential gene expression is represented for all genes as a colour

gradient across all sampling points within different groups from brick red (lowest) to black (highest)

for spleen, green (lowest) to dark orange (highest) for head kidney, blue (lowest) to red (highest)

for skin. Group details: Group 1 is control group; Group2 received ip injection of adjuvant emulsified

vaccine antigen; Group 3 received ip injection of adjuvant emulsified vaccine antigen + bath

immunization.

Fig 6. PCA analysis of skin samples pre and post lice challenge. PCA analysis for the *in vivo* challenge

samples representing the distribution of lice infested host skin samples in vaccinated (group 2 and

3) and only adjuvant vaccinated control (group 1) groups at 0, 17, 28 and 50 dpi (A and B). Analysis

was based on mean fold-changes of all genes for each individual sample at each sampling point

(smaller symbols) relative to the unvaccinated control. The ellipses indicate the group

dispersion/variability from the centroid (larger symbols) calculated using all individual fold-changes

values/group (A). (C) shows the contribution of sampling points to different components. (D) shows

the contribution of genes on different components and the significant genes contributing in

principal component 1 and 2.

Supporting information

S1 Fig. Two-way hierarchical clustering heat map for each tissue. The rows represent gene

expression and the column represents different sampling points within respective groups. Group

details: Group 1 is control group; Group2 received ip injection of adjuvant emulsified vaccine

antigen; Group 3 received ip injection of adjuvant emulsified vaccine antigen + bath immunization.

S2 Fig. PCA analysis of head kidney samples post immunization and lice infestation. PCA analysis

of head kidney samples from vaccinated (group 2 and 3) and only adjuvant vaccinated (group 1)

groups at 0, 17, 28 and 50 dpi (A and B). Analysis was based on mean fold-changes of all genes for

each individual sample at each sampling point (smaller symbols) relative to the unvaccinated

control. The ellipses indicate the group dispersion/variability from the centroid (larger symbols)

calculated using all individual fold-changes values/group (A). (C) shows the contribution of sampling

points to different components. (D) shows the contribution of genes on different components and

the significant genes contributing in principal component 1 and 2.

S3 Fig. PCA analysis of spleen samples post immunization and lice infestation. PCA analysis of

spleen samples from vaccinated (group 2 and 3) and only adjuvant vaccinated (group 1) groups at

0, 17, 28 and 50 dpi (A and B). Analysis was based on mean fold-changes of all genes for each

individual sample at each sampling point (smaller symbols) relative to the unvaccinated control. The

ellipses indicate the group dispersion/variability from the centroid (larger symbols) calculated using

all individual fold-changes values/group (A). (C) shows the contribution of sampling points to

different components. (D) shows the contribution of genes on different components and the

significant genes contributing in principal component 1 and 2.

S4 Fig. Transcriptional analysis of immune genes post immunization and lice infection.

Transcript levels of the pro-inflammatory cytokines (A) and immune genes (B) in spleen, head kidney

and skin at different sampling points: 69 days from first vaccination (69d) or zero day challenge and

after challenge (dpi: days post infestation), were analysed by real-time QPCR. The QPCR data were

normalized to the geometric mean of the 3 house-keeping genes (EF-1a, 18S and β-actin) and

expression is relative to the pre-immunized level. Fold change was calculated using the primer

efficiency. Data shown represent the mean ± SD of experiments performed in triplicate, n=18

fish/group (6 fish/replicate). Statistical analysis was carried out using one-way ANOVA or Kruskal

Wallis test followed by Tukey or Dunn's Multiple Comparison compared to control group (*P < 0.05,

**P < 0.01, ***P < 0.001). Group details: Group 1 is control group; Group2 received ip injection of

the adjuvant emulsified vaccine antigen; Group 3 received ip injection of adjuvant emulsified

vaccine antigen + bath immunization.

A

Incubator number/ Group number	Observations
I1-G1	Less copepodite number in the water column and some at the tank bottom
I3-G1	Good numbers in the water column and at bottom
I4-G1	Good numbers in the water column and at bottom
I5-G1	Good numbers in the water column and at bottom
I6-G1	Good numbers in the water column and bottom
I8-G2	Less number of copepodids at the bottom
I7-G2	Less copepodids at the bottom
I9-G2	Less in the water column. Half of an egg was still to hatch
I6-G2	Less at the bottom and in the water column. One egg string still to hatch
I3-G2	Less number at the bottom. Nothing visually observed in the water column
I10-G2	Less number at the bottom
I11-G3	Many in the water column. Less number at the bottom
I12-G3	Many in the water column. Less number at the bottom
I13-G3	More number in the water column and at bottom
I14-G3	Good numbers in the water column and bottom
I15-G3	Less numbers in the water column and bottom

B

A

B

C

D

Significant gene variables in PC1

	correlation	p.value
TNF-α	0.9796554	2.652822e-08
IFN-γ	0.9687562	2.224960e-07
IL-1ß	0.9643262	4.285462e-07
IL-8	0.9546093	1.405862e-06
MMP-9	0.9288714	1.271394e-05
IL-10	0.9160907	2.841690e-05
IL-4/13A	0.9105080	3.883910e-05
IL-22	0.8747809	1.957688e-04
CD4	0.8531028	4.187136e-04
IgM	0.8314264	8.018853e-04
CD8α	0.7592495	4.182072e-03
IgT	0.6207972	3.122735e-02

Significant gene variables in PC2

	correlation	p.value
IgT	0.6843255	0.01409954