**Title:** Anticipatory responses along motion trajectories in awake monkey area V1

**Authors**

Giacomo Benvenuti[1‡], Sandrine Chemla[1], Arjan Boonman[1†], Laurent Perrinet[1], Guillaume S Masson[1], Frédéric Chavane[1]*

**Affiliations**

[1]Institut de Neurosciences de la Timone (INT), CNRS and Aix-Marseille Université, UMR 7289, Campus Santé Timone, 27 boulevard Jean Moulin, Marseille 13005, France.

‡ Current address: Center for Perceptual Systems, University of Texas, Austin, Texas 78712, USA

†Current address: Department of Zoology, Faculty of Life Sciences, Tel Aviv University, Tel Aviv 6997801, Israel

**\*Correspondence: frederic.chavane@univ-amu.fr**

**Highlights**

- Our hypothesis is that lateral propagation of activity in V1 contributes to the integration of translating stimuli

- Consistent with this hypothesis, we find that a translating bar induces anticipatory spiking activity in V1 neurons.

- A V1 model describes how this anticipation can arise from inter and intra-cortical lateral propagation of activity.

- The dynamic of VSDi and LFP signals in V1 is consistent with the predictions made by the model.

- The intra-cortical origin is further confirmed by the fact that a bar moving from the ipsilateral hemifield does not evoke anticipation.

- Horizontal and feedback input are not only modulatory but can also drive spiking responses in specific contexts.

**ABSTRACT**

What are the neural mechanisms underlying motion integration of translating objects? Visual motion integration is generally conceived of as a feedforward, hierarchical, information processing. However, feedforward models fail to account for many contextual effects revealed using natural moving stimuli. In particular, a translating object evokes a sequence of transient feedforward responses in the primary visual cortex but also propagations of activity through horizontal and feedback pathways. We investigated how these pathways shape the representation of a translating bar in monkey V1. We show that, for long trajectories, spiking activity builds-up hundreds of milliseconds before the bar enters the neurons' receptive fields. Using VSDI and LFP recordings guided by a phenomenological model of propagation dynamics, we demonstrate that this anticipatory response arises from the interplay between horizontal and feedback networks driving V1 neurons well ahead of their feedforward inputs. This mechanism could subtend several perceptual contextual effects observed with translating objects.

**INTRODUCTION**

In standard information processing models of the visual system, low-level visual features are extracted locally within selective receptive fields and are rapidly relayed to downstream areas in order to encode increasingly complex features (Fukushima, 1980; Riesenhuber and Poggio, 1999). When probed with spatially stationary stimuli centered on their receptive fields, properties of visual neurons from V1 to extra-striate areas are indeed well captured by a simple cascade of *linear-nonlinear models* encoding low-level visual properties such as local orientation or motion direction (Carandini et al., 2005; Rust et al., 2006) as well as more complex visual pattern (David et al., 2006; DiCarlo et al., 2012). A well-known example is given by *visual motion integration* where three decades of studies have been conducted almost exclusively with motion signals presented within a stationary aperture, such as drifting periodic stimuli or random dot patterns, centered on neuronal receptive fields (**RF**). The reported neuronal properties have provided strong evidences for the *linear-nonlinear cascade* theoretical framework, where V1 neurons encode local ambiguous motion signals that are integrated by MT neurons to compute direction and speed of pattern motion (Movshon et al., 1985; Rust et al., 2006; Simoncelli and Heeger, 1998; Wilson et al., 1992).

However, natural objects often move along extended and smooth spatio-temporal trajectories within the visual field. A handful of psychophysical, computational and physiological studies (Watamaniuk & Mc Kee 1995, Watamaniuk et al 1995, Burgi et al., 2000; Perrinet and Masson, 2012; Tlapale et al., 2010) have proposed that the visual system can use the spatio-temporal regularity of smooth trajectories to solve many of the uncertainties inherent to visual motion such as aperture (Marr, 1982; Wallach, 1935) and correspondence problems (Ullman, 1979; Wertheimer, 1912) and to improve the *signal to noise ratio* of visual input evoked by moving stimuli (e.g. Burgi et al., 2000). This idea is consistent with the view that visual systems can

benefit from contextual information and internal models of the statistics of the visual world
(Berkes et al., 2011; Geisler, 2008).
For example, the property of smoothness associated to motion trajectories can be used by the
visual system to generate internal models of moving objects (i.e. priors). In accordance with this
hypothesis, there is a rich psychophysical literature demonstrating how spatio-temporal
coherence affects the way we perceive moving stimuli attributes such as (1) position (MacKay,
1958; Nijhawan, 1994; W. Metzger, 1932), (2) direction (Anstis and Ramachandran, 1987;
Lorenceau et al., 1993; Ramachandran and Anstis, 1983; Welch et al., 1997), (3) motion (Van
Doorn and Koenderink, 1984; Nakayama and Silverman, 1984; Neri, 2014; Snowden and
Braddick, 1989b; Verghese and Watamaniuk, 1999; Verghese et al., 2000; Watamaniuk and
McKee, 1995; Watamaniuk et al., 1995), (4) speed (Castet et al., 1993; Georges et al., 2002;
McKee and Welch, 1985; Scott-Brown and Heeley, 2001) (5) orientation (Alais and Lorenceau,
2002; Fredericksen et al., 1994; Guo et al., 2004; Pavan et al., 2011; Werkhoven et al., 1990) and
(6) color (Watanabe and Nishida, 2007). For example, a target stimulus presented at the leading
edge of a smooth trajectory has been shown to be easier to detect than when presented in
isolation or at the trailing edge of the trajectory (Arnold et al., 2014; Lenkic and Enns, 2013; Liu
et al., 2006; Roach et al., 2011; Schwiedrzik et al., 2007). This phenomenon cannot be simply
explained by a shift in spatial attention, lateral facilitation or retinal motion and it is more likely
implemented at an early stage of visual processing (Roach et al., 2011). Furthermore, stimuli
presented at the leading edge of long smooth trajectories are perceived having  higher resolution
(i.e. lower blurring due to motion) than stationary moving stimuli (Burr and Ross, 1986;
Watanabe and Nishida, 2007) and are more easily trackable behind occlusions (Watamaniuk and
McKee, 1995).

Several computational studies suggest that a simple hierarchical feedforward model with
contiguous local filters sequentially activated by the translating stimulus is insufficient to
account for these contextual effects. In contrast, they proposed that a *mechanism involving*
*lateral propagation across the sequence of filters activated along the trajectory* could explain the
emergence of motion trajectory effects (Burgi et al., 2000; Grzywacz et al., 1995; Kaplan et al.,
2013; Khoei et al., 2017; Perrinet and Masson, 2012; Tlapale et al., 2010; Yuille and Grzywacz,
1989). These models postulate that lateral interactions in early visual areas could participate in
processing the responses evoked by smooth trajectories of translating stimuli.  Early visual areas
are "retinotopically organized" (Van Essen and Newsome, 1984; Hubel and Wiesel, 1974), such
that translating stimuli generate a sequential activation of contiguous locations on the retinotopic
map.  Since neurons in these areas are tightly connected to their neighbors through horizontal
connections and feedback loops (Angelucci et al., 2002; Benucci et al., 2007; Bringuier et al.,
1999; Bullier, 2001; Chavane et al., 2011; Girard et al., 2001; Grinvald et al., 1994; Jancke et al.,
2004a; Muller et al., 2014; Reynaud et al., 2012; Slovin et al., 2002) see review (Muller et al.,
2018) (**Figure 1A**), feedforward responses at each location can rapidly spread to contiguous
areas of the map before the translating stimulus enters their RF (Jancke et al 2004, Chemla et al

2019). Consequently, the neural population mapping the leading edge of the translating stimulus
will receive convergent lateral and FF inputs. These two signals may be recursively integrated
along the motion trajectory and generate a gradual build-up of neural activity that would carry
trajectory information. The advantage of such integration mechanism is that it would combine
only the spatio-temporally coherent responses evoked by translating stimuli.

The aim of this study is to investigate this intriguing hypothesis. The primary visual cortex (**V1**)
is a good candidate circuit to support such trajectory-based computation as it has the largest and
most precise retinotopic representation of the outside world (Hubel and Wiesel, 1974; Tootell et
al., 1982). To understand how V1 networks map motion trajectories, it is thus necessary to
elucidate the interactions between feedforward, feedback and lateral networks at the proper
spatio-temporal recording scales.

To achieve this goal, we recorded both single-unit and neuronal population activities in the
primary visual cortex of four awake, fixating, monkeys to map the dynamics of the cortical
responses to a translating bar. By using different trajectory lengths, we manipulated the history
of cortical activation elicited by the moving bar before reaching the classical RF of the recorded
neuron. Using single-unit recordings, we found that for long trajectories, about half of V1
neurons demonstrate a significant build-up of anticipatory spiking responses long before the bar
enters their classical receptive field. Using a computational model, we show that the specific
dynamics of spiking anticipatory responses could be explained by an intra-cortical horizontal and
feedback network propagating information faster than the feedforward input sequence. Using
voltage-sensitive dye imaging (**VSDi**) to measure the aggregate membrane potential of a large
V1 area, we showed indeed that the neural population response to a translating bar exhibits a
spatio-temporal structure consistent with the predictions of the model. To test whether feedback
signals could also contribute to this anticipatory activity, we used a multi-electrode array to
measure local field potential (**LFP**) responses across the V1 area and showed that a bar moving
over a long trajectory elicits a strong and very early decrease of low frequency power, a signal
recently attributed to feedback modulation (Bastos et al., 2014; Engel and Fries, 2010; van
Kerkoerle et al., 2014). The intra-cortical origin of these anticipatory responses was further
demonstrated by the observation that anticipation was abolished when the bar translation was
initiated from the ipsi-lateral visual hemi-field. Altogether, these results dissect out the multiple
scales at which an intricated network of connectivities within and between visual areas leads to
the emergence of an anticipatory spread of activity that shall contribute to processing of
translating objects.

**RESULTS**

The goals of this study were to test whether trajectory-dependent responses to a translating bar
can be observed in V1 and, if so, understand the underlying network mechanisms.

**Anticipatory spiking responses**

We first recorded spiking responses from 80 V1 single-units in two alert, fixating macaques presented with three visual stimulation paradigms. Neurons were selected regardless of their tuning properties. For each neuron, we first mapped the spatial profile of the classical RF using sparse noise sequences and the orientation and direction tunings, using a translating bar (12 directions). Then, a vertical bright bar (0.5x4 degrees of visual angle, **dva**) was moved horizontally at 6.6 dva/s starting at either 1.5, 3 or 6dva (short, medium and long trajectories respectively) from the RF centre and disappearing 1.5-2dva beyond it (**Figure 1A**). Critically, these three trajectories stimulated the RF in exactly the same way. However, in the medium and long trajectory conditions, the translating bar had a different history, activating a large area outside of the RF through a smooth path before reaching it.

For each cell, the responses for the three trajectory conditions were aligned relative to the absolute position of the bar (**Figure 1B-F**). On the other hand, for comparing the evoked responses across cells (**Figure 1E-F**), we aligned all time-courses relative to the latency of the response evoked by the short path condition. This operation led to the automatic exclusion of 31 cells for which the detection of this important reference point was not possible due to low signal-to-noise ratio (see Methods) and the fact that stimulus direction and orientation were not optimized for the cells' preference.

In the remaining 49 cells, the time-course of the responses (PSTH) to the shortest (1.5dva) motion trajectory presented a typical fast transient rise and a slower decrease (**Figure 1B-F, black trace**). However, when the bar trajectory started 3dva or 6dva before the RF centre (dark and light blue curves, respectively), a ramp-like increase of the discharge-rate was observed before the response-onset for the shortest motion trajectory, in half (53%) of the neurons (**Figure 2A**). Such anticipatory build-up scaled up with the trajectory length. In the other neurons, responses were identical for the three trajectory conditions (**Figure 1D & figure supplement 1**), excluding that anticipatory responses were due to a systematic bias in our specific experimental approach (e.g. stimulation protocol, general bias due to eye movements, generalized allocation of spatial attention, generalized effect due to the input from LGN). On average, latencies for the medium (3dva) and long (6dva) trajectories started respectively 116ms (std =93ms) and 220ms (std = 189ms) before the latency to the shortest path (1.5dva), a decrease which was found significant over the whole population (two-sample t-test P<0.05, see **Figure 2D**). This timing corresponded to an average distance of 2.3 and 3dva between the bar position and the RF centre. In contrast, both time-to-peak and response offset remained similar across all trajectories (two-sample t-test P>0.05), excluding a spatial shift of the RF or an unspecific gain effect.

**Figure 1. Single units responses to a translating bar as a function of its trajectory length outside the RF.** (**A**) Experimental approach. Schematic of the retino-thalamo-cortical neural circuit. A translating bar (bottom) is going to cross consecutive V1 RFs (grey disks) eliciting a sequence of feedforward activations, propagating laterally across the retinotopic map through intra (i.e. horizontal) and inter-cortical (feedback-loops) connections. Neurons with the RF at the end of the trajectory are recorded extracellularly using an electrode (top-right). The inset on the right depicts the paths of the three trajectories (dotted lines, 1.5, 3 and 6dva, respectively) with respect to the RF (grey circle). The color-code is consistent throughout figures. (**B-D**) Examples of single unit responses for the three trajectory lengths presented with raster plots (top) and PSTHs (bottom) (B and D**:** Monkey PD, C**:** Monkey NO). B

and C are examples of neurons presenting anticipation, D of a neuron that does not. Colored lines
(bottom) indicate the stimulus presentation time, for each condition. Latency is indicated by triangles.
Shades represent the standard error of the mean (SEM) across trials. (**E-F**) Average PSTHs across
anticipatory neurons for the three trajectories, for monkeys PD (n=13) and NO (n=13), respectively.
Responses are aligned to the latency of the short trajectory condition. Colored lines (top) indicate
significant difference with respect to the short trajectory condition (two-sample t-test P<0.05).

This anticipatory activity extended well beyond the size of the measured RF, similar to classical
results (Angelucci et al., 2002), and was neither correlated with the RF sizes nor could be
explained by a systematic bias in monkeys' fixation (see **figure supplement 2**). The gradual rise
of this anticipatory response, furthermore, contrasts with the transient profile of the neural
response for a bar swept over the RF and suggests a slower cumulative process. These
anticipatory responses merged ~100ms before the short-trajectory response onset and from then
followed an identical dynamic. This asymmetry eliminates the possibility that the response build-
up would be related to a systematic misestimation of the classical RF position or size, that would
have resulted in a change in response peak time for instance. Next, we aimed at better
characterizing the origin of the cell-to-cell variability in the observed anticipatory responses, and
its relationship to the tuning of the cells.
223
**Understanding the diversity of degrees of anticipation**
To estimate the proportion of cells showing a significant modulation of firing rate over time, we
compared spike counts across trials in short (i.e. 1.5 dva) *vs* long (3 & 6 dva) trajectories
conditions. The proportion of cells significantly activated (Two-sample t-test P<0.05) at different
time bins along the motion trajectory gradually increased (**Figure 2A**), such that 53% of them
showed a significant anticipatory response 100ms before the bar entered the classical RF, for
long (3dva and/or 6dva, green) but not short (1.5dva, black) trajectories. By comparison, when
the bar entered the classical RF (i.e. vertical dotted line), all three conditions drove the same
number of cells.

**Figure 2. Amplitudes, latencies and tunings of trajectory dependent responses.** (**A**) Proportion of
neurons  (out of a total of n=49) showing a significant response over time for all 3 trajectory conditions
(see cartoon in insert). Green color represents proportion of neurons showing a significant response only
for the medium and/or long trajectory conditions. Colored lines at the bottom indicate stimulus
presentation for the three trajectories. (Top-left) same inset than **Fig1A**. (**B**) Difference of firing rate
between responses to long (3-6dva) and short (1.5dva) trajectory lengths before entering the RF (thick
when significant) vs within the RF (red filled when significant). Before and within the RF was estimated
using the latency of the short trajectory response. Table on the top: number of cells presenting a
significant increase (1), decrease (-1) or no change (0) in firing-rate rate in long trajectories compared to
short trajectories. Table on the bottom: same than table on the top but for medium trajectories (**C**) Mean
onset latencies (square), peak response times (diamond) and offset response time (stars) for the three
trajectories conditions as a function of trajectory length averaged across neurons and monkeys (n=49).
Time 0 is when the bar crosses the RF center. Dots indicate latencies of individual medium and long
trajectories responses. An second abscissae is drawn at the top to indicate the equivalent visual distance
(dva) from the receptive field center. (**D**) Polar histogram representing the normalized averaged
anticipatory response strength (computed in a time window 100ms before the latency of the short
trajectory) as a function of neurons direction preference (n=32, only cells with DI>0.1). Inset: number of
cells preferring each direction.

In a few neurons, the two longest trajectories resulted in a significantly different spiking
response in the RF center (e.g. the example in **Figure 1C**). This result could provide an
important evidence in support of the predictive coding theory, as prediction would result in
smaller error signal and hence a smaller neuronal response (Alink et al., 2010; Mumford, 1992;

Srinivasan et al., 1982). However, this effect was not evidenced in the average response profiles
(**Figure 1E-F**) although it could have been smoothed out because of the variability in response
latencies. Hence, we compared individual discharge rates between short and long trajectories in
two different time-windows of 100ms: one placed just before (ordinate) and one just after
(abscissae) response latency (as estimated by detecting the first significant change with respect to
the baseline) for the short trajectory condition (i.e. the RF border) (**Figure 2B**). A small fraction
of the cells exhibiting an anticipatory build-up of activity responded slightly less when the bar
crossed their RF center (upper left quadrant in Figure 2B). However, such decrease was
significant in only 4% of the cells while, on the contrary, 16% displayed a significant increase of
their response in the RF (insets). These results rule out a systematic relationship between the
occurrence of an anticipatory build-up and a modulation of the stimulus-driven responses within
the RF center.
269
Next, we compared, for every neuron, the latencies for medium and long trajectories vs. the
latency for the short one (**Figure 2C**). This analysis revealed a continuum in the timing of
anticipation as shown by the latency scatter for the medium and long trajectories that fully
covered a wide range of values from 0 to -400 and -800ms, respectively. While the distributions
of response onset latencies were significantly different for the three trajectories conditions (two-
sample t-test p<0.05) both time-to-peak and response offset latencies distributions were not
significantly different across all trajectories (two-sample t-test p>0.05). We checked if the
variability in anticipatory responses across all neurons can be due to differences in their tuning
properties. Selecting neurons with a direction index > 0.1 we probed the relationship between the
relative firing rate in the 100ms time-window before the bar enters the RF and direction
preferences (**Figure 2D**). Indeed, neurons with a preferred direction aligned to the stimulus
motion direction exhibited a stronger anticipatory response. Therefore, the diversity of the
anticipation timing and strength of the response across neurons may simply reflect a direction-
selective anticipatory mechanism.
284
Overall, we demonstrate that a large population of V1 neurons exhibits a slow build-up of
activity prior a moving input hits their receptive field. Such anticipatory activity occurs sooner
for longer trajectories (**Figure 2C,** squares). In comparison, both the response peak and offset
response time did not change with trajectory length (Figure 2C, diamond and stars respectively).
Our next goal was to understand what neural mechanisms could generate this anticipatory
response.
**Anticipation is an expected emergent property of intra-cortical propagation**
These characteristics of observed anticipatory responses can be used to theoretically constrain
the putative underlying neural mechanisms. First, these responses emerge well before the
stimulus enters the classical RF of the recorded neurons. Second, the decrease in onset latency is
not followed by a similar shift in time-to-peak and offset latency. Eventually, anticipatory
responses present a slow dynamic consisting of a gradual rise of activity. A feedforward network

alone is unlikely to account for the slow dynamics and the asymmetry of these responses.
Alternatively, these features are consistent with surround modulation (Bair et al., 2003; Klink et
al., 2017; Nassi et al., 2013; Reynaud et al., 2012). Visual information from the RF surround can
be provided to V1 neurons from intra-cortical horizontal (**H**) and inter-cortical feedback (**FB**)
networks (Angelucci and Bressloff, 2006; Roelfsema, 2006). We therefore reasoned that the
anticipation we observed in V1 shall arise from the interactions between feedforward and intra-
cortical inputs carried by horizontal and feedback connections. To test this hypothesis, we
disentangled the contribution of these signals to anticipatory responses by using the well know
intrinsic differences in the spatial and temporal scales of these inputs. To do so, we developed a
simple phenomenological, but biologically-plausible, model simulating feed-forward and
cortico-cortical responses within a cortical retinotopic map (**Figure 3A,** see Methods and **figure**
**supplement 3**).
309
310

**Figure 3. A physiologically plausible model of lateral cortical integration.** (**A**) Cartoon describing the model developed to predict the time-course of the neural population response to a translating bar in the space-time domain. The translating bar evokes a sequence of spatio-temporal feedforward activation of the cortex at a speed of $v_{ff}$ (thick blue line, blue hue codes for time). The feedforward activation is then relayed by horizontal propagation at speed $v_h$ and over a distance of $d_h$ represented by thinner obliques lines. (**B**). Activation of the model when $v_{ff}$, $v_h$ and $d_h$ are set to values compatible with our visual stimulation paradigm and the know horizontal network properties of the cortex (Angelucci et al., 2002; Girard et al., 2001; Grinvald et al., 1994; Muller et al., 2014). Color-code represents cortical activation strength (arbitrary units). (**C**) Time-course of the neural population response at different positions along the trajectory. Traces are aligned to the time of the feedforward arrival in the receptive field centre (the

stimulus onset is displayed as a slant dark line). For each response time-course, latency is indicated by a
disk. The profile of latency changes with the increase of the trajectory length is captured by a piecewise
linear regression (red and orange lines) showing a first decrease with a slope S1 up to a position P,
followed by a second slope S2. (**D**) Effect of changing the speed $v_{ic}$ on the slope S1 (left). The middle and
right panel shows, for a speed example (black circle in the left panel), an example of time-courses and the
plot of latencies as a function of distance along the trajectory (with extraction of S1, P and S2). (**E**) Same
as in D but for changes of $d_{ic}$. (**F**) Same as in D but for changes of feedback strength.

The response to a bar translating along an extended, straight, trajectory was modelled as a
sequence of feedforward activations at different spatiotemporal positions, changing with speed
$v_{ff}$ (**Figure 3A**). At each spatiotemporal position, this feedforward input also gave rise to an
isotropic horizontal propagation of activity with speed $v_h$ over a cortical distance $d_h$ (**Figure 3A**).
When $v_h > v_{ff}$, the model qualitatively predicts the emergence of a gradual anticipatory build-up
of activity whose onset latency depends on trajectory length (**Figure 3C**). This effect is a
consequence of the speed difference between the sequence of feedforward activations (i.e.
stimulus speed) and the horizontal propagation: after an initiation phase, a given cortical position
will receive information about the approaching bar first from the intra-cortical propagation (i.e.
anticipatory response) and second from the feedforward input. For each position, after realigning
the time-course of activation patterns to the end of all trajectories, we observed a gradual
emergence of an anticipatory activation, whose latency (blue dots) decreases with trajectory
length (color-coded as hue of the blue) and reaches saturation for distances larger than $d_h$ (**Figure
3D**). Beyond this distance, the latencies of anticipatory responses do not change anymore simply
because of the absence of direct anatomical horizontal connections. In other words, to generate a
latency decrease, the bar needs to reach a region that is directly connected to the recording site.

To capture this behavior, we used a piecewise linear regression of the latency as a function of
trajectory length. From this regression, we extracted the slope of the change in the response
latency at the beginning and at the end of the trajectory (**S1** and **S2** respectively) and the position
of intersection between these regression lines (**P**). These 3 parameters correspond to the speed at
which latency of anticipatory activity decrease along the trajectory (S1), the spatial limit of this
decrease (P) and the rate of latency changes for distances beyond this position (S2). To
understand how does the spatio-temporal properties of the intra-cortical propagations affect these
three parameters, we varied $v_h$ and $d_h$ over a large range of values. As predicted, we found that
**S1** and **P** are under control of $v_h$ and $d_h$ respectively, which linearly modulate these parameters
over a large range of values (**Figure 3D and E**). In short, the properties of the intra-cortical
propagation directly influence the speed and the extent of the latency decrease.

This observation has two theoretical consequences. First, if anticipation would result from a
large divergence of the feedforward inputs, we should observe a latency decrease at a fixed delay
after stimulus onset since all thalamic input arrive with the same latency.  As a result,
anticipation should not be trajectory-dependent and should generate a very high speed S1
(**Figure 3D**). Second, the spatial extent of the observed latency decrease should be equivalent to

anatomical extent of the anatomical divergence (**Figure 3E**). In the following section, we tested
the validity of this model with further neural recordings at different scales.
366

**VSD imaging unveils an intra-cortical propagation of anticipatory activity**

Our model produces testable predictions for the spatiotemporal profile of V1 population
responses to a translating stimulus. To test these predictions, we need to record cortical activity
at both high resolution and large field of view in order to measure the fine change of response
latency as a function of trajectory length. To do so, we used voltage-sensitive dye imaging
(**VSDi**), a mesoscopic imaging technique with high spatial and temporal resolution. Since this
technique reveals the aggregate membrane potential oscillations of the cortex (Grinvald and
Hildesheim, 2004), we can dissect out the effect of cortical propagation with a high sensitivity.
In a third, awake and fixating, monkey (WA), we measured population synaptic activity using
VSDi (**Figure 4A**) over a large cortical region representing retinotopically -2dva to 0dva of
azimuthal eccentricity.
378

The translating bar elicited a wave of VSD activity propagating across the retinotopic map at the
expected speed and direction of the bar motion trajectory. More important, a strong anticipatory
activity was also detectable at the early frames for the medium and long trajectories (**Figure 4B**).
Notice that, since the VSDi signal reflects the combination of both the sub and supra-threshold
response components (Chemla and Chavane, 2010; Chen et al., 2012), this observation at
population level indirectly confirms the spiking results in a third monkey but with a different
technique. For similar retinotopic positions, the anticipatory response evoked by the moving bar
was stronger for the long trajectory than the one for the medium trajectory (compare first 4
panels in Figure 4B) confirming the slow build-up of activity reported above. Our model
demonstrates how such slow dynamics can be a signature of lateral propagation of activity and
not a divergent feedforward network.
390

To facilitate the qualitative comparison with the model simulations shown in **Figure 3C**, we
displayed the time course of the VSDi responses as being aligned to the end of the trajectories
and sorted accordingly to their distance from the initial bar position (blue hue color code, **Figure**
**4D**, top panel). For each cortical position, the anticipatory VSD activity started growing slowly a
few hundreds of milliseconds before the bar reached the corresponding retinotopic positions.
Beyond this point, responses were identical across all conditions, similarly to what was observed
at the spiking level, and in close agreement with model predictions. We measured both onset
(vertical ticks at the bottom of Figure 4D top panel) and peak latencies for each condition and
plotted them as a function of the distance (in visual angle) between the initial position of the bar
and the position encoded by each cortical image pixel (**Figure 4D** bottom panel). Responses
peaked (triangles) at a fixed delay along the bar trajectory, dynamically mapping the bar motion
with the accurate speed (6.8dva/s), corresponding to a cortical speed of 0.016m/s (the equivalent

of $v_{ff}$ in our model). Notice that, in **Figure 4D** we aligned the responses to the end of the
trajectories, hereby correcting for bar motion. As a consequence, the vertical alignment of peaks
demonstrates that it always occurs when the bar was at the retinotopic position encoded by the
retinotopic map (in other words, when it reaches the population RF). In contrast, response onset
latencies for pixels close (<2dva) to the starting position of the bar motion decreased rapidly. By
fitting this spatio-temporal profile as in the model, we estimated an apparent cortical speed of
0.06m/s, consistent with the speed of intra-cortical horizontal propagation (Bringuier et al., 1999;
Girard et al., 2001; Muller et al., 2014; Reynaud et al., 2012). This result suggests that intra-
cortical inputs, propagating faster on the cortex than the feedforward activation sequence elicited
by the moving bar (0.06 vs 0.016 m/s), is most probably at the origin of the anticipatory
activation observed in macaque area V1.
414
However, for pixels located more than 2dva away from the stimulus starting position (i.e. the
position is estimated from the piecewise linear regression fit procedure), responses latencies
decreased much less, corresponding to a cortical speed of ~0.02m/s. Such temporal dynamics
corresponds to a visual speed of 8.8dva/s, only slightly higher than the bar translation speed
(6.6dva/s). Accordingly to our model, such values suggest that the large anticipatory responses
seen for long distances, result from the sequential activation of a nested horizontal network
driving responses with a spatial range of 2dva. It is compatible with the spatial extent of the
horizontal network in macaque V1 (Angelucci et al., 2002). These measured values for both
speed and spatial extent are compatible with the intra-cortical connectivity but not with a
feedforward origin. Indeed, if anticipation would result from divergence of the feedforward
network, our model predicts a faster latency decrease and the spatial extent of this decrease
would be smaller than the one we measured (Blasdel and Lund, 1983; Florence et al., 1988;
Perkel et al., 1986). Overall, both model prediction and VSDi measurements are in strong
agreement to support an intra-cortical origin of anticipatory responses in macaque V1.
429

**Figure 4. Neural population responses to a translating bar as a function of trajectory length**
**(A)** Experimental rationale. Similar to Figure1A but VSD responses are captured using a CCD camera.
**(B)** Time sequence of the VSD response map (2 sessions, n=37 trials) for the three trajectory conditions.
**(C)** Retinotopic map of the horizontal dimension (grayscale code for the horizontal eccentricity, contours delimit the progression for every 0.5dva). Top-right: Cartoon representing corresponding bar positions in the visual field in the blue color-code. **(D)** Top: Time course of the VSD responses for all conditions broken down by the distance of the initial bar position to the pixel's retinotopic position. Vertical lines: responses latencies. Color-code same as Figure3C. Bottom: Latency (circle) and time-to-peak (triangle) of VSD responses as a function of the distance of the initial position of the bar to the retinotopic representation (similar to Figure 3C). Red and orange lines are linear regressions.

What could be the origin of the difference observed between cortical (8.8dva/s) and stimulus (6.6dva/s) speeds? One hypothesis is that it result from gain modulation exerted by long-distance feedback inputs, since only those can mediate such long-range lateral interactions (Angelucci et al., 2002). Therefore, we added a feedback component to our model that acts by varying the relative gain between horizontal and feedforward inputs, as already suggested by others (Liang et al., 2017; Piech et al., 2013; Ramalingam et al., 2013). Increasing the gain of lateral inputs changed the shape of the response time-course (**Figure 3F**) resulting in an increased detectability of the response and, thereby, slightly increasing the slope of the second component in our spatiotemporal profile. These effects are consistent with the results obtained in VSD. To verify this hypothesis, our next step was to test whether the time-scale of such gain modulation was coherent with the dynamics of feedback inputs.

**Low-frequency LFP signals support the existence of an early and diffuse anticipation signal**

Convergent studies have recently suggested that the power of low-frequency (i.e. alpha and beta bands) local-field potentials (LFP) signals in V1 can be used as a proxy to quantify the amount of feedback inputs onto V1 (Bastos et al., 2014; van Kerkoerle et al., 2014). Therefore, in order to test our model prediction, that the contribution of feedback signals could participate to the dynamic of anticipatory responses, we studied the time-course of low-frequency LFP signals in V1 in response to our 3-trajectories paradigm (**Figure 5A**). To do so, we measured LFP signals from arrays of 10x10 electrodes (Utah arrays, **MEA**) chronically implanted in two monkeys (BA and PD). The arrays covered a V1 cortical territory of 4x4mm, representing 2-3dva of the motion path. Similar to VSDi, the LFP response (0.3 - 250Hz) to the translating bar across the array followed the projection of the bar to the retinotopic map, swiping across the cortex at the matching cortical speed of 0.015m/s (**Figure 5B**).

For longer trajectories (lower rows), clear anticipatory LFP responses were observed all over the array. As for above, we aligned the responses from all electrodes based on their retinotopic position and calculated the time-course of the LFP power-frequency spectrum for the responses to the three trajectories conditions (**Figure 5C**). The anticipatory response was evidenced as a power increase or decrease at high or low frequencies, respectively. Across all frequencies, the response peak always occurred when the bar crossed the centre of the retinotopic area covered by the electrodes array. To detail the dynamics of low and high frequency LFP signals as a function of the bar trajectory length, we filtered the LFP responses using two bandwidths (low: bandwidth 7-25Hz and high: 40-100Hz) and aligned these signals based on the distance between the RF position for each electrode and the initial position of the trajectory (**Figure 5D-E,** top panels). We found a very early and strong anticipatory decrease in the low frequency band. At higher frequencies, however, the anticipatory build-up was more similar to the spiking activities described above, both in terms of amplitude and latency. For both frequency bands, changes were graded according to the length of the motion path. To interpret these results in the framework of our model, we extracted the latency of low- and high-frequency bands and plotted them as a function of trajectory length (**Figure 5D-E**, bottom panels). At high frequencies, in the gamma-band regime, we observed a dynamic similar to single-units and VSDi with an initial phase of latency decrease well captured by a propagation of 0.14m/s. For distances larger than 2dva, the decrease in latency is stable. However, at low-frequencies, in the alpha to beta-band regimes, we observed a constant delay after stimulus onset for all trajectories.

This temporal dynamic could be due to divergent inputs from either thalamo-cortical or feedback pathways, since both can generate synchronous inputs over a large extent. However, given the limits on the spatial extent of the thalamic axonal arbors in primates (less than 0.5mm, (Blasdel and Lund, 1983; Florence et al., 1988; Perkel et al., 1986)) and the divergence of thalamo-cortical inputs (less than 0.5mm in diameter, (Perkel et al., 1986)), this effect is more probably due to feedback from downstream areas that, in contrast, have the appropriate spatial and

493 temporal scales (Angelucci et al., 2002; Girard et al., 2001; Stettler et al., 2002). Overall, these
494 results are consistent with the predictions of our model. In the last part of our study, we further
495 support the hypothesis that the generation of anticipatory responses necessitates lateral
496 propagation of activity using a complementary approach.

497
498

**Figure 5. Multi-unit LFP responses as a function of stimulus' trajectory length** (**A**) Experimental
approach. Similar to Fig. 1A but the LFP of the neural population response is recorded using a square
multi-electrode-array (MEA) (**B**) Time sequence of the map of LFP responses in MEA space for the three
trajectory lengths, (3 sessions, 52 trials per condition). Colored lines at the bottom indicate stimulus
presentation and condition type (color-code) (**C**) Time-frequency representation of the LFP power
averaged across 96 electrodes. (**D-E**) <u>Top:</u> Time-course of the averaged LFP power for low (D) and high
(E) frequency bands, for all conditions broken down by the distance of the initial bar position to each
electrode retinotopic position (t=0ms). Black-to-cyan color codes for the increasing distance of the
electrode retinotopic position to the initial position of the bar. <u>Bottom:</u> Latency (circle) of LFP responses
as a function of the distance of the initial position of the bar to the retinotopic representation. The black
line corresponds to the bar trajectory time-course. The orange and red dotted lines are the result of a
piecewise linear regression and the black dotted line of a single linear regression.

511
512

## Anticipation is cancelled for inter-hemifield stimulation

In all conditions tested so far, bar trajectories were all presented in the contra-lateral visual hemi-
field, activating a continuous cortical path confined within the recorded hemisphere. If the intra-
cortical origin hypothesis holds true, anticipatory responses should not be evoked when the
trajectory has its origin within the ipsilateral visual hemifield. This is due to the fact that, since
our recordings are retinotopically located very close to the vertical meridian, such stimuli would

activate first the opposite hemisphere, generating activation to the recorded location through the
myelinated callosal V1-V1 connections, that are much faster than the amyelinated lateral
connections and are mostly restricted to a region close to the vertical meridian (Kennedy and
Dehay, 1988; Kennedy et al., 1986).
523
In three monkeys, we tested this prediction using both LFP (**Figure 6A**) and VSDI recordings
(**Figure 6B**). As expected, for the opposite stimulus direction, the peaks of the LFP and VSDi
responses move towards the opposite cortical direction (compared to **Figures 4** and **5**). However,
in stark contrast to a stimulus restricted to the contra-lateral hemi-field (**Figures 4,5**), moving a
visual stimulus from the ipsi-lateral hemifield did not generated anticipatory responses, for both
types of recordings and for both low and high frequency power values (see below). This result
provides another strong evidence that the observed anticipatory responses are mainly driven by
propagations within the intra- an inter-cortical, horizontal and feedback networks.
532

**Figure 6. Neural population responses to trajectories with different lengths presented in the ipsi-**
**lateral visual hemi-field.** (**A,C**) The experimental rational for LFP (A) and VSDi (C) recordings. Neural
responses are measured in the retinotopic area close to the vertical meridian for a bar starting to move
from the ipsi-lateral visual hemifield (i.e. activating the opposite hemisphere) and finishing its trajectory
just after crossing the vertical meridian (i.e. activating the recorded hemisphere). Red discs indicate the
position of the fixation spot. (**B**) and (**D**) same legend as in Figure 4B and 5B respectively.
540

Lastly, we compared the average responses latencies across all monkeys, for the three
complementary recordings, the three different trajectory length and the two trajectory directions
(**Figure 7**). For a stimulation within the contra-lateral hemifield, anticipation gradually develops
along the trajectory length in the various measures of neuronal activity. The strongest
anticipation was seen for low-frequency band LFP responses (squares), followed by VSDi
(triangle) and high-frequency LFP (diamond) responses, leading to a final anticipation expressed
at suprathreshold level in single unit activity ( circle, **Figure 7A**). For the opposite direction, in
both LFP and VSDi recordings, no sign of anticipation is evident for both LFP and VSD signals.
Moreover, delayed responses are observed in the VSDi responses for the long trajectories.
(**Figure 7B**).

552
553 **Figure 7. Comparison of results across all recording techniques and 4 monkeys.** (**A**) Averaged
554 latencies of SUA (2 monkeys), LFP power and VSDi responses as a function of trajectory length for
555 stimulation within the contra-hemifield. (**B**) Same than A but for a bar coming from the ipsi-lateral
556 hemifield. SEM is calculated across all neurons (SUA) electrodes (LFP) or pixels (VSD).
557

**DISCUSSION**

In primary visual cortex of awake, fixating monkeys, we show that a bar translating along a
smooth and extended trajectory significantly activates half of the recorded neurons several
hundreds of milliseconds and a few degrees before actually entering their classical RF. This
trajectory-dependent anticipatory activity has a specific slow temporal profile composed of a
ramping response that is intrinsically different from the classical transient and fast responses
driven by pure feed-forward cortical inputs. Such anticipatory activity cannot be explained by an
underestimation of the RF sizes or by the scatter in the eye positions. These surprising
anticipatory spiking responses present a wide range of timings across neurons, longer trajectory
lengths leading to earlier response build-up onsets. This variability in anticipatory response onset
is partially accounted for by the neurons' direction preference: cells preferring the motion
direction of the bar display the strongest and earliest anticipatory build-up of spiking activity. In
contrast, after entering the classical RF of the neurons, the translating bar produces the same
spiking activity regardless of its trajectory length and therefore its cortical mapping history.

Anticipatory responses in the visual system have already been reported in a few physiological
studies, first in vitro in the retina of rabbit and salamander (Berry et al., 1999), in area 17 of
anesthetized cat (Jancke et al., 2004b; Orban et al., 1985), but also monkeys areas V1 (Guo et al.,
2007; Subramaniyan et al., 2015) and V4 (Sundberg et al., 2006). However, these previous
observations all differ on several key features when compared to our present study. First, these
previous studies (with the exception of Guo et al., 2007, see below) were inspired by the flash-
lag effect, in which a translating stimulus is judged ahead of a stimulus flashed at the same
position at the same time. In this context, these studies have compared the latency of responses
evoked by bars in translation to bars flashed alone in the RF centre. The RF was therefore
stimulated differently by the flashed and the moving stimuli, in the sense that the moving
stimulus activated a sequence of positions in the RF before reaching the position in which the
flashed stimuli was presented for comparison. In contrast, in our approach, the classical RF is
stimulated exactly in the same way for all trajectory conditions. Consistent with their initial
hypothesis, these previous studies showed a global shift in the time-course of their responses,
while maintaining the same shape (i.e. time-to-peak and response offset present similar
anticipation than response onset). In our study, anticipatory responses consist in the appearance
of an early ramping activity for long trajectory. Afterwards, the response peaks and decrease at
the same time than shorter trajectories.

Our results are in accordance with an earlier study carried out in the anesthetized macaque on a
small number of neurons, where a bar flashed just outside the classical RF of a V1 neuron can
evoke a response when presented after a sequence of 3 other collinear bars in apparent motion,
outside the RF (Guo et al., 2007). However, the phenomenon reported in this study is again
different from the one we report, since the apparent motion stimulus used will stimulate V1 in a
sequence of discrete and transient feedforward activations, creating an intrinsically different

activity pattern than that caused by a smooth translating stimulus. Indeed, it has been recently
demonstrated, using VSDi in the awake monkey, that a two-strokes apparent motion stimulus
generates two waves of propagation interacting non-linearly, instead of many cumulative
propagations produced by each position visited by a translating object (Chemla et al., 2019).
602
Crossing different physiological signatures at single cell and population levels, and guided by a
computational model, we provide convergent evidence that the observed spiking anticipation is
not a mere consequence of similar events occurring at earlier processing stages but rather results
from a specific propagation of activity within V1 intra- and inter-cortical networks. Our
biologically realistic-model shows that the cumulative recruitment of converging horizontal input
can produce a gradual build-up of anticipatory activity preceding the feedforward activation.
Such build-up is evident at both sub-threshold and spiking levels. The trajectory-dependent shift
in response onset timing was also observed with high resolution and large field-of-view using
VSDi. From the time-course of these neural population responses across the retinotopic map, we
have been able to estimate the speed and spatial extent of the propagation underlying the
observed anticipation. The spatio-temporal properties of such propagation are fully compatible
with the well-known characteristics of the cortical horizontal network in primate area V1 for
trajectories up to 2dva (Angelucci et al., 2002; Girard et al., 2001; Muller et al., 2014).
616
On the other hand, the further decrease in the response onset we observed using VSDi for
trajectories extending beyond 2dva (i.e. covering more than 6mm of cortical space) cannot be
explained by this horizontal network alone because of the limitation in horizontal connections
length. Therefore, we hypothesized that a divergent feedback input from higher visual areas
could generate such fast and long-distance effect (Angelucci and Bressloff, 2006; Bair et al.,
2003; Bullier, 2001). Using our model, we found that a feedback mechanism can indeed induce a
further, though smaller, decrease of response onset for these larger trajectory distances, as
observed with VSDi. To support this hypothesis, we show that the bar appearance, even for
trajectories longer than 2dva,  triggers a fast and strong decrease of low-frequency LFP power,
that is likely to indicate a feedback modulation (Bastos et al., 2014; van Kerkoerle et al., 2014).
Furthermore,  the spatial and temporal properties of this signal are compatible with the known
properties of the feedback connectivity (Angelucci and Bressloff, 2006; Bair et al., 2003; Bullier,
2001; Girard et al., 2001). As suggested for sensory-motor networks, such decrease could be
related to reporting an expected change in the input, a deviation from the "status quo"  (Engel
and Fries, 2010).
632
A last and strong piece of evidence in support of the intra-cortical propagation mechanism comes
from our control experiment in which we moved the bar from the ipsilateral visual hemifield,
first activating the cortical hemisphere opposite to our recordings. In fact, while slow lateral
interactions, mediated by amyelinated horizontal connections, are restricted to the same
hemisphere, inter-hemispheric interactions, mediated by fast myelinated callosal connections, are

restricted to a region of the visual field close to the vertical meridian (Kennedy and Dehay, 1988;
Kennedy et al., 1986). Therefore, a bar moving within the ipsi-lateral visual hemifield would not
activate the horizontal network of the recorded cortex. Fast, inter-hemispheric callosal
interactions will signal the arrival of the bar in the contra-lateral visual hemifield but only when
it approaches the vertical meridian and without any precise retinotopic projection. This is
consistent with fact that we did not observe any anticipatory responses in our population activity
measurements (VSDi, LFP) for ipsi-lateral approaching bar. While we did not record single unit
activities under this condition, VSDi and gamma power LFP signals indicate also the presence of
spiking activities. We shall therefore not expect to find any significant slow ramping of spiking
activity when a moving object approaches from the other side of the vertical meridian. This does
not preclude, however, that inter-hemispheric trajectories could be linked at other spatio-
temporal scales in extra-striate cortex that can exhibit larger receptive fields with ipsilateral
representations  (Desimone et al., 1993; Gross et al., 1977; Pigarev et al., 2001).

All these converging evidences thus suggest that trajectory-dependent, anticipatory, spiking
activity results from the interplay of intra- and inter-cortical networks, both propagating activity
faster than the feedforward sequence of inputs. Incidentally, our results also show that, through
the accumulation of convergent sub-threshold activations, intra-cortical network can drive
spiking responses several hundreds of millisecond before the arrival of feedforward inputs,
instead of being only modulatory (see also (Jancke et al., 2004a). A possible functional
consequence could be that V1 neurons can signal the direction of a moving object before it
reaches their receptive fields. Indeed, the fact that anticipatory responses are stronger for neurons
whose direction preference matches the trajectory motion direction suggests that direction-
selective anticipatory signals can reach downstream populations of neurons (for example MT
cells). Thus, motion integration might be facilitated by propagation of anisotropic and
anticipatory signals, alleviating motion uncertainties (Burgi et al., 2000; Perrinet and Masson,
2012; Tlapale et al., 2010).

Our study documents for the first time that the neural activity evoked along the trajectory travels
across the retinotopic map at the cadence imposed by the FF response, representing at each
moment in time information about the stimulus recent history but also its likely future location
and velocity. The cortical representation of the leading edge of a motion trajectory is gradually
spreading out over a larger retinotopic area than stimuli presented along short trajectories.
Downstream extra-striate visual neurons with large RFs could therefore integrate both the
feedforward evoked population response with this trajectory-dependent population response.
This could be relevant in explaining why stimuli presented at the leading edge of motion
trajectories are more detectable (Arnold et al., 2014; Lenkic and Enns, 2013; Liu et al., 2006;
Roach et al., 2011; Schwiedrzik et al., 2007) and their location are perceived ahead of a stimulus
flashed in the same position (Jancke et al., 2004b; Nijhawan, 1994).

678 Additionally, this phenomenon could be at the origin of the diffusion of motion information, a
679 mechanism postulated to be necessary for enhancing motion discriminability (Burgi et al., 2000;
680 Grzywacz et al., 1995; Yuille and Grzywacz, 1989)' and inferring non-ambiguous motion
681 direction (Anstis and Ramachandran, 1987; Perrinet and Masson, 2012; Ramachandran and
682 Anstis, 1983). Consistently with our observations indeed, such psychophysical effects are
683 modulated by motion trajectory length (McKee and Welch, 1985; Nakayama and Silverman,
684 1984; Snowden and Braddick, 1989a, 1989b; Watamaniuk et al., 1995). Hence, this circuitry
685 may serve as a plausible mechanism for a predictive coding of future locations and to resolve
686 motion integration computations (Kaplan et al., 2013; Khoei et al., 2017; Perrinet and Masson,
687 2012). Indeed, such diffusion could subserve two key computational elements of predictive
688 coding: taking advantage of temporal coherency to predict future events and removing highly
689 correlated events within the sensory inflow (Alink et al., 2010; Mumford, 1992; Rao and Ballard,
690 1999).

692 In conclusion, our study suggests that propagation of activity within or between cortical areas
693 may play an important role in the dynamical processing of non-stationary stimuli (see also
694 (Chemla et al., 2019; Muller et al., 2018)). We document an effect occurring at the specific
695 spatial and temporal scales of macaque area V1. The anticipatory effect propagates only over a
696 few degrees ahead of the translating bar. We shall envision that the lateral interactions will
697 propagate information across the overlaid functional maps not only in V1, but also in all the
698 other visual areas and maps. In other words, we may imagine a generalization of the effect across
699 the various scales of the encoded retinotopic maps along the visual system, or across different
700 feature maps such as the orientation map in V1 (Chavane et al 2011, Rankin & Chavane 2016,
701 Tlapale et al, 2010), the direction map in MT (DeAngelis and Newsome, 1999) or more complex
702 maps in other extra-striate visual areas. Analyzing systematically the anticipation across these
703 feature maps may help to understand to what extent this anticipatory signal is a simple predictive
704 warning signal or whether it carries more information about the nature of the incoming stimulus.
705 We propose that parallel propagation of information within all these maps, combined with rapid
706 exchange of information between these maps, may be a canonical computational operation by
707 which the visual system process complex, natural, non-stationary inputs.

708
709
710

711 **METHODS**

712 **1. Animals and surgery procedures**

714 Experiments were conducted on a total of five hemispheres, in four adult male rhesus monkeys
715 (*Macaca mulatta):* monkeys NO (right hemisphere), PD (left and right), BM (right), BR (right).
716 Experimental protocols have been approved by the Marseille Ethical Committee for Animal
717 Research (approval #A10/01/13, official national registration #71-French Ministry of Research).

All procedures complied with the French and European regulations for animal research, as well
as the guidelines from the Society for Neuroscience. All the monkeys were chronically implanted
with a head-holder. Monkey NO, PD and BR were also implanted with a recording chamber
located above areas V1/V2. Monkey PD and monkey BM were implanted with a Utah array.
Monkey BR had a third surgery for durotomy and transparent artificial dura mater insertion, in
order to perform voltage-sensitive dye imaging. Monkeys NO and PD were implanted with a
scleral search coil for right eye position monitoring.

**Multi-electrodes array:** Ninety days after head holder implantation, once monkeys PD and BM
were trained to achieve a good fixation, a second surgery was performed. Area V1 of the left
(monkey PD) and the right hemisphere (monkey BM) were surgically implanted with a 100-
electrode Utah array (Blackrock Microsystems, Salt Lake City, UT, USA). The array had an
arrangement of 10×10 (PD: Platinum ~400kΩ 1kHz, BM: Iridium oxide ~50kΩ, at 1kHz)
electrodes, each of them 1mm long, with an inter-electrode distance of 400μm. The surgery was
performed under deep general anesthesia and full aseptic procedures. Anesthesia was induced
with 10mg/kg *i.m*. ketamine and maintained with 2–2.5% isoflurane in 40:60 O2–air. To prevent
cortical swelling, 2ml/kg of mannitol *i.v*. was slowly injected over a period of 10min. A 2×2 cm
craniotomy was performed over the visual cortex and the dura was incised and reflected. The
Utah array was inserted in the cortex using a pneumatic inserter (Array Inserter, Blackrock
Microsystems) ~5mm under the lunate sulcus, in perifoveal position and covered with a sheet of
an artificial non-absorbable dura (Gore-tex). The real dura was sutured back and covered with a
piece of an artificial absorbable dura (Seamdura, Codman). The bone flap was put back at its
original position and attached to the skull by means of a 4×40 mm strip of titanium (Bioplate,
Codman). The array connector was fixed to the skull on the opposite side with titanium bone
screws (Bioplate, Codman). The skin was sutured back over the bone flap and around the
connector. The monkey received a full course of antibiotics and analgesic before returning to the
home cage.

**Voltage sensitive dye imaging (VSDi):** The surgical preparation and VSD imaging protocol
have been previously described in Reynaud et al., 2012 (Reynaud et al., 2011). Briefly, after
head holder and chamber implantation on the right hemisphere, a second surgery was performed,
once the monkey BR was trained to achieve a good fixation. The dura was removed surgically
over a surface corresponding to the recording aperture (18mm diameter), and a silicon-made
artificial dura was inserted under aseptic conditions (Arieli et al., 2002). Such silicon-dura is
necessary to have a good optical access to the cortex.

**2. Recordings**

**Eye Movements:** For monkeys NO and PD, horizontal and vertical positions of the contra-
lateral eye to the recorded hemisphere, were recorded using the scleral search coil technique

(Judge et al., 1980; Robinson, 1963). For those monkeys, a second surgery was performed to
insert a search coil below the ocular sclera, to record eye movements with the electromagnetic
technique (Robinson, 1963).  For monkey PD and BM we used the Eyelink 1000 (SR Research
sampling frequency of 1,000Hz). For monkey BR, we used an ISCAN eye tracking system (512
x 256 x 60Hz, ETL-200).

**Single electrode (SE):** A computer-controlled microdrive (MT-EPS, AlphaOmega, Israel) was
used to insert through the dura a tungsten micro-electrode (FHC, 0.5-1.2MΩ at 1,000Hz) on the
right hemisphere of monkey PD and NO. Spikes were sorted on-line using a template-matching
algorithm (MSD, AlphaOmega).

**Multi-electrodes array (MEA):** Recordings were carried out through a 128-channel Cerebus
acquisition system. The signal from each active electrode (96 out of the 100 electrodes were
connected) was pre-processed by a head stage with unity gain and then amplified with a gain of
5000. The signal was filtered in two different frequency bands to split it into LFPs (0.3–250Hz)
and spiking activity (0.5–7.5kHz). The LFPs were sampled at 1kHz and saved on disk. To
improve the electrode independence among LFPs, we subtracted from all electrodes the common
LFP averaged across all electrodes.

**Voltage sensitive dye imaging (VSDi):** After removing of the artificial dura-mater, the cortex
was stained with the voltage-sensitive dye RH-1691 (Optical Imaging), prepared in artificial
cerebrospinal fluid (aCSF) at a concentration of 0.2mg/ml, and filtered through a 0.2µm filter.
After a 3 hours staining period, the chamber was rinsed thoroughly with filtered aCSF, to wash
off any supernatant dye. The artificial dura was inserted back and the chamber closed with
transparent agar and cover glass. Optical signals were recorded from a focal place ~300µm
below the surface using a Dalstar camera (512x512pixels resolution, frame rate of 110Hz) driven
by the Imager 3001 system (Optical Imaging). Excitation light was provided by a 150W halogen
lamp filtered at 630nm and fluorescent signals were high-pass filtered at 665nm during 1363ms
(trial duration).

**3. Experimental protocols**

**Behavioral task and Training:** Monkeys were trained to fixate a red target presented on the
center of the screen for all the duration of the trial (~1.5sec) within a window of 1-2dva of
diameter. We monitored the eye positions and controlled online the behaviour (sampling rate: 1
KHz) using the REX package under the QNX operating system (Hays et al., 1982). For VSDI
experiments, both online behavioral control and image acquisition were heartbeat triggered;
heartbeat was detected with a pulse oximeter (Nonin 8600V). After a successful trial, i.e. the

monkey has maintained his fixation during the whole acquisition period (~2s), monkeys received
a drop of water (NO, PD, BM) or mashed apple (BR) as a reward. For VSDi recordings, an inter-
trial interval of 8 sec was set for dye bleaching prevention.
801
**Visual stimulation:** Visual stimulation protocols were displayed using different devices
depending on the technique. For both SE and MEA recordings, they have been produced using
in-house software (Gérard Sadoc, Acquis1-Elphy, Biologic CNRS-UNIC/ANVAR) but with
different displays. In each system, the display was gamma calibrated by means of a lookup table.
For the SE experiments, stimuli were back-projected on a translucent screen covering of visual
field of the monkey, at a distance of 100 cm, using a video projector (resolution:
$1280 \times 1024$ pixels at 60 Hz, operating range 0-24 cd/m$^2$). The luminance of the motion stimulus
was 22.2 cd/m$^2$ (background 2.2cd/m$^2$). For the MEA experiments, stimuli were presented on a
LCD screen 27' (resolution: 1920x1080 pixels at 60Hz, operating range 0-294cd/m$^2$) at a
distance of 57cm. Mean luminance of the motion stimulus was 80cd/m2 (background 5cd/m$^2$).
For VSDI, the visual stimuli were computed on-line using VSG2/5 (VSL v8) libraries on Matlab
(The MatWorks Inc., Natick, MA, USA) and were displayed on a 22' CRT monitor at a
resolution of 1024x768 pixels and a refresh rate of 100Hz, operating range 0-75cd/m$^2$. The
viewing distance was set to 57cm, the mean luminance of the motion stimulus was set to
71cd/m2 and the background was kept constant to 12cd/m$^2$
817
**SE stimulation protocol:**   Each neuron was characterized using three different stimulation
paradigms. First a sparse noise (SN) paradigm, to quantitatively map out the receptive field (RF)
of the neuron. The SN consisted in the sequential presentation of small dark and light squares of
0.6x0.6dva, flashed for 50ms randomly positioned within a 10x10 grid (11.2+/-11 cd/m2,
~15trials, see (Bringuier et al., 1999). The stimulation for the following paradigms was centered
on the RF centre (RFc). To estimate the RFc, we extracted the center from the ellipse that was
best adjusted to the activity contour at a 3.8 z-score (equivalent to a p-value of 0.1% corrected
for 100 positions). Then, a direction tuning (DT) paradigm, using a bright bar (0.5 x 4 dva)
moving at 6.6 dva/s. The bar was crossing the stimulation centre in the middle of its trajectory (6
dva altogether) and randomly presented at 12 different directions (spaced by 30° of circular
angle). The bar orientation was always perpendicular to the motion direction. Finally, a trajectory
length paradigm (3T) in which the same bar, oriented perpendicularly to the motion direction
was moved horizontally at 6.6 dva/s, starting at 1.5, 3 or 6 dva from the RF centre and
disappearing 1.5 (NO) or 2 dva (PD) beyond the RF centre. The bar trajectory started from a
lateral position in the contra-lateral visual hemi-field towards the vertical meridian. In those three
protocols, control trials with no stimulation were presented within each block (blank condition).
All stimuli were presented in random order.
835
836    **MEA stimulation protocol:** Electrodes LFP RF were mapped by the same sparse-noise
837    paradigm used for SE and stimulation was centered on the centre of mass of the population RFs.

838    Then, we used the same protocols (DT and 3T) than for single units recordings. However, we
839    also extended the protocol 3T using a symmetrical trajectory moving towards the RFs from the
840    ipsilateral visual hemi-field.
841
**VSDi stimulation protocol:** The VSDi stimulation protocol for the 3T paradigm was established
using the exact same stimuli (see above). The trajectory vertical position was centered on 2dva
below the horizontal, the imaging area being roughly centered on [-2; -1dva] (see retinotopic
map using intrinsic optical imaging Figure 4C), while horizontally moving at 6.6dva/s from 7, 4
or 2.5dva on the left of the vertical meridian to 0dva (vertical meridian position). Here again, we
extended the protocol using a symmetrical trajectory moving towards the RFs from the ipsilateral
visual hemi-field. In addition to these three evoked conditions, two blank conditions, i.e. a grey
screen is presented during the acquisition period, were used and all stimuli were randomly
interleaved.
851
852    **4. Data Analysis**
853    All data have been analyzed using custom-written routines in Matlab.
854
**SUA PSTH plots:** For every neuron, we calculated the peristimulus time histograms (PSTH) for
the three trajectory lengths conditions, averaging across the trials PSTHs (total number of trials
between 15 and 35, bin size of 10ms), smoothed by a low pass filter (window of 120 ms) and
expressed in spikes per second. Standard error of the mean (SEM) was calculated across trials.
First, the histogram of every trial (bin size 10ms) was smoothed with a low pass filter (window
of 120ms). Then we calculated the SEM across these smoothed histograms. Response latency
was calculated on PSTHs that had a peak response higher than 1.96 times the standard deviation
over the mean of the amplitude of the blank condition PSTH (amplitude threshold). The latency
was computed on the PSTH derivative (estimated on each time frame "t" by calculating the
linear regression of the PSTH response over 20 points, corresponding to 200ms, centred on t).
The latency was defined on each condition as the moment the PSTH derivative was crossing a
threshold, (calculated as the mean plus 1.96 times the standard deviation of the derivative of the
blank condition PSTH) for 5 consecutive points.
868
**Averaged PSTHs:** For each monkey, we averaged the PSTHs of cells presenting a statistically
significant anticipatory response (two-sample t-test between the distribution of spikes numbers
for trial of the longer trajectory length conditions and the short one in a 100ms time window
placed just before the response latency for the short trajectory, i.e. just outside the classical RF).
Before averaging, single cells PSTHs were aligned on the latency of the short trajectory
condition. Second, to remove the unselective component of the response for each cell, the
average across time of the PSTH baseline (the first 400ms before stimulus presentation) was
subtracted from the full PSTH. To align the PSTHs of each cell on the latency for the short
trajectory, for each condition, we first calculated the error between the expected (40ms after

stimulus presentation) and measured latency and then we adjusted the PSTHs offset in time to
remove it. The statistical significance of the averaged responses was calculated over time (within
a sliding time window of 100ms, shifted of steps of 10ms). For each neuron and each condition,
the distribution of spikes counts across trials, in each time window, was compared by a two-
sample t-test (P<0.05) to the distribution of spikes counts across trials, over all the time windows
in the blank condition.

**Time course of the neurons proportion with statistically significant response:** For each
neuron, we calculated at which moment in time (within contiguous time windows of 100ms) the
distribution of spikes counts across trial, for the evoked conditions, was significantly different
from the distribution of spikes counts across trials and all the time windows of the blank
condition (one-tailed two-sample t-test, P<0.05). For each time window, we calculated the
percentage of neurons with a significant response, with respect to the number of neurons
presenting a significant response in the time window corresponding to the RF center (49
neurons).

**Relative firing rate within a window of 100ms just before and just after the RF border:** For
each neuron, we calculated the difference in spike rate between the short and the longer
trajectories conditions (aligned on the RF centre) within a 100ms window, placed just before the
response latency if the response to the short trajectory or just after. The statistically significant
response differences between the distribution of spikes numbers for trial of the longer trajectory
length conditions and the short one, were calculated by one-tailed two-sample t-test (P<0.05).

**Neurons preferred direction and direction index:** Neurons preferred direction was calculated
fitting the neurons tuning curves (built from the PSTH responses within a time window of 200ms
centred on the RF centre for all the 12 conditions of the DT paradigm) with a double
multiplicative Von Mises function, see (Swindale, 1998). Neuron direction index was calculated
as 1, minus the ratio of the response in the opposite direction to the response in the preferred
direction.

**Probability of anticipation:** Probability of anticipation was calculated for each cell by one-
tailed two-sample t-test (P<0.05) between the short and the longer trajectory conditions (aligned
on the latency for the short trajectory condition) within a 100ms window place just before the
response latency for the short trajectory.

**Spikes representation in visual space corrected by eye position deviation from the fixation**
**point (control):** For 22 cells in monkey PD, we checked whether the response anticipation could
be explained by eye movements towards the approaching bar. For that purpose, for each trials the
time occurrence of each spikes was converted into a visual space coordinate corresponding to the
stimulus position in visual space for that particular time assuming a latency of 50ms ((skp

timing-50) *stimulus speed). This position was then converted by a position in retinal space
taking into account the distance of eye position (low pass filtered across 100 ms) from the
fixation centre (the average across all the eye positions), at -50ms from the spike time. Then we
calculated the PSTHs for the evoked conditions, in space coordinates, for the rasters before and
after correction of spike positions.

**LFP power spectral analysis:** The LFP spectrum (sampling rate 1000Hz) was computed using
Chronux open source MATLAB toolbox (http://www.chronux.org/, multi-taper method,
*mtspecgramc.m* function). Taper parameters were bandwidth (W): 2.5, taper duration (T): 1,
integer (p) such as 2TW-p tapers are used: 2. In order to optimize the frequency/time resolution,
we used two different sliding time windows for low (7-25Hz) and high (40-100Hz) frequencies
(respectively 400ms and 150ms). The LFP spectrum of the evoked conditions was divided for
the mean spectrum of the blank condition (across time) and the response offset at zero frame was
subtracted. LFP was recorded through a Utah array in 3 sessions for a total of 52 trials for
condition. The spectrogram was obtained by averaging across 96 electrodes, re-aligned on the
LFP RF centre of every electrode, determined through the sparse-noise paradigm (as described
above).

**VSD data analysis:** Stacks of images were stored on hard-drives for off-line analysis. The
analysis was carried out using the Optimization, Statistics, and Signal Processing Toolboxes. The
evoked response to each stimulus was computed in three successive steps. First, the recorded
value at each pixel was divided by the average value before stimulus onset (frames 0 division), to
remove slow stimulus-independent fluctuations in illumination and background fluorescence
levels. Second, this value was subsequently subtracted by the value obtained for the blank
condition (blank subtraction) to eliminate most of the noise due to heartbeat and respiration
(Shoham et al., 1999). Finally, a linear detrending of the time series was applied to remove
residual slow drifts induced by dye bleaching (Chen et al., 2008; Meirovithz et al., 2010). For
this study, VSD data from two sessions were accumulated by registering them onto a reference
session (using a quadratic registration algorithm described by Takerkart et al., 2008(Takerkart et
al., 2008)). Then responses amplitudes were normalized to a reference condition and data were
*Z*-scored(Jancke et al., 2004a) for further analysis.

**Retinotopic maps:** To obtain a retinotopic representation, we presented the bar stimulus
(0.5x4dva) at 6 different static positions (from -2.5 to 0.5dva compared to the vertical meridian)
for 600ms (after a delay period of 200ms) and we recorded intrinsic optical imaging signals from
the cortical surface(Grinvald et al., 1994). Each trial consisted of 4181ms (92frames) and was
followed by an inter-trials interval of 5s. For intrinsic-signal imaging, the cortex was illuminated
with a 605nm light. Off-line analysis of intrinsic maps started with the same normalization
procedures than for VSD images, i.e. frames 0 division and blank subtraction) providing the
cortical activation profile for each static positions of the bar. Then for each pixel, we extracted

the mean values in each of the six cortical profiles and computed the centre of mass. The
retinotopic map was finally constructed after conversion into visual field positions.

**VSDI Latency estimations:** Response latency was defined as the point in time at which the
signal derivative crossed a threshold set at 2.57 times (99% confidence) the SD of its baseline
computed during a 100-ms-long window right before stimulus onset.

**Spatio-temporal phenomenological cortical Model:** The model was built to represent V1
activity over a 2D matrix representing retinotopic cortical space in one dimension (full extent of
21mm) and time in the other dimension (full extent of 1sec). In this spatio-temporal
representation, we modelled the feedforward input as an activation moving across the matrix at a
speed of $v_{ff}$ set to 0.016 m/s (equivalent to 6.6dva/s in cortical coordinate at the eccentricity of
our neural recordings). At each moment, the dot-like feedforward input generated an activation
profile implemented as a gaussian function in the spatial domain, with a standard deviation of
0.42mm.

For each of these feedforward activations, we computed V1 activity as the interaction of this
activation with that of similarly defined horizontal and feedback maps. First, horizontal activity
is the result of a spatio-temporal convolution with a kernel ($k_h$) which is then scaled with a non-
linearity. The kernel is modelled as a spreading gaussian (see eq 1) for which the standard
deviation increases linearly with time with a celerity $c$ and the amplitude decreases exponentially
with the square of time relative to $\tau$. The weight $w_h$ of this kernel is adjusted to account for the
strength of the horizontal input. The speed of the horizontal spread $v_h$ was therefore controlled by
$c$ (set by default to 0.1m/s) and the spatial extent is controlled by a trade-off between $\tau$ and $c$, i.e.
the maximum extent being reached for the latest point in time for which the horizontal activity
amplitude is still above baseline. As a consequence, we used the value of $\sigma$ at time $t=1.5\tau$ (the
amplitude will have reach 10% of its maximum) as a proxy for the horizontal extent $d_h$ (set by
default to 6mm)

$$k_h = \frac{\alpha(t)}{2\pi} e^{-\frac{t^2}{2\sigma(t)}} \quad \text{for } t>=0, \text{ where } \quad \sigma(t)=ct \quad \text{and} \quad \alpha(t)=we^{-\left(\frac{t}{\tau}\right)^2} \quad (1)$$

To account for the observation that the feedforward response amplitude does not change along
the trajectory, the horizontal activity contribution was scaled through a sigmoid non-linearity (**S**)
controlled by the difference between feedforward and horizontal activities. Horizontal and
feedforward signals were then linearly combined with a factor of $\rho$.

$$V_1 = \rho ff + (1-\rho)S\left(\frac{ff-h}{g}\right)h \quad \text{where} \quad S(y) = \frac{1}{1+e^{-y}}$$

Feedback effect was thereafter modelled as a spatio-temporal multiplicative gain to the horizontal activity. The kernel of the feedback was similarly designed as the one for the horizontal, albeit with much faster speed and larger spatial extent. The resulting activity of the feedback amplified multiplicatively the activity of the horizontal activity (Liang et al., 2017; Piech et al., 2013; Ramalingam et al., 2013). The feedback strength was controlled by a scalar weight ($w_{fb}$) that we modulated between 0 (no feedback) and 1.2 (feedback stronger than horizontal).

The evolution of the expected response time-course for different positions along the trajectory can be simply read-out from the matrix representing the spatio-temporal activity. Similarly to what was done in experimental data, latency was calculated as threshold crossing of derivative above a given value (set to 0.01).

**AUTHOR CONTRIBUTIONS**

GB and AB performed single unit recordings. GB performed MEA recording. SC performed VSDI recording. FC and LP did the computational model. FC supervised all the experiments and designed the study with the help of GB. GB, GSM & FC wrote the manuscript with the help of AB and SC.

**ACKNOWLEDGMENTS**

This work was supported by the Marie Curie Program (FACETS-ITN) the FET Program (FACETS, BrainScales) of the European Union and the ANR Trajectory and Horizontal-V1. We thank Ivan Balansard, Xavier DeGiovanni, Marc Martin and Fréderic Barthélemy for their excellent technical support. We are grateful to Cyril Monier who provided an important help in setting up the MEA experiments and to Lyle Muller for careful reading of the manuscript.

**REFERENCES**

Alais, D., and Lorenceau, J. (2002). Perceptual grouping in the Ternus display: evidence for an "association field" in apparent motion. Vision Res. *42*, 1005–1016.

Alink, A., Schwiedrzik, C.M., Kohler, A., Singer, W., and Muckli, L. (2010). Stimulus predictability reduces responses in primary visual cortex. J. Neurosci. *30*, 2960–2966.

Angelucci, A., and Bressloff, P.C. (2006). Contribution of feedforward, lateral and feedback connections to the classical receptive field center and extra-classical receptive field surround of primate V1 neurons. Prog. Brain Res. *154*, 93–120.

Angelucci, A., Levitt, J.B., Walton, E.J.S., Hupe, J.-M., Bullier, J., and Lund, J.S. (2002). Circuits for Local and Global Signal Integration in Primary Visual Cortex. J. Neurosci. *22*, 8633–8646.

Anstis, S., and Ramachandran, V. (1987). Visual inertia in apparent motion. Vision Res. *27*, 755–764.

Arieli, A., Grinvald, A., and Slovin, H. (2002). Dural substitute for long-term imaging of cortical activity in behaving monkeys and its clinical implications. J. Neurosci. Methods *114*, 119–133.

Arnold, D.H., Marinovic, W., and Whitney, D. (2014). Visual motion modulates pattern sensitivity ahead, behind, and beside motion. Vision Res. *98*, 99–106.

Bair, W., Cavanaugh, J.R., and Movshon, J.A. (2003). Time course and time-distance relationships for surround suppression in macaque V1 neurons. J. Neurosci. *23*, 7690–7701.

Bastos, A.M., Vezoli, J., Bosman, C.A., Schoffelen, J.-M., Oostenveld, R., Dowdall, J.R., De Weerd, P., Kennedy, H., and Fries, P. (2014). Visual Areas Exert Feedforward and Feedback Influences through Distinct Frequency Channels. Neuron *85*, 390–401.

Benucci, A., Frazor, R.A., and Carandini, M. (2007). Standing waves and traveling waves distinguish two circuits in visual cortex. Neuron *55*, 103–117.

Berkes, P., Orban, G., Lengyel, M., and Fiser, J. (2011). Spontaneous Cortical Activity Reveals Hallmarks of an Optimal Internal Model of the Environment. Science (80-. ). *331*, 83–87.

Berry, M.J.I., Brivanlou, I.H., Jordan, T.A., and Meister, M. (1999). Anticipation of moving stimuli by the retina. Nature *398*.

Blasdel, G.G., and Lund, J.S. (1983). Termination of afferent axons in macaque striate cortex. J. Neurosci. *3*, 1389–1413.

Bringuier, V., Chavane, F., Glaeser, L., and Frégnac, Y. (1999). Horizontal propagation of visual activity in the synaptic integration field of area 17 neurons. Science *283*, 695–699.

Bullier, J. (2001). Integrated model of visual processing. Brain Res. Rev. *36*, 96–107.

Burgi, P.Y., Yuille, A.L., and Grzywacz, N.M. (2000). Probabilistic motion estimation based on temporal coherence. Neural Comput. *12*, 1839–1867.

Burr, D., and Ross, J. (1986). Visual processing of motion. Trends Neurosci. *9*, 304–307.

Carandini, M., Demb, J.B., Mante, V., Tolhurst, D.J., Dan, Y., Olshausen, B.A., Gallant, J.L., and Rust, N.C. (2005). Do we know what the early visual system does? J. Neurosci. *25*, 10577–10597.

Castet, E., Lorenceau, J., Shiffrar, M., and Bonnet, C. (1993). Perceived speed of moving lines
depends on orientation, length, speed and luminance. Vision Res. *33*, 1921–1936.

Chavane, F., Sharon, D., Jancke, D., Marre, O., Frégnac, Y., and Grinvald, A. (2011). Lateral
Spread of Orientation Selectivity in V1 is Controlled by Intracortical Cooperativity. Front. Syst.
Neurosci. *5*, 4.

Chemla, S., and Chavane, F. (2010). Voltage-sensitive dye imaging: Technique review and
models. J. Physiol. *104*, 40–50.

Chemla, S., Reynaud, A., di Volo, M., Zerlaut, Y., Perrinet, L., Destexhe, A., and Chavane, F.
(2019). Suppressive traveling waves shape representations of illusory motion in primary visual
cortex of awake primate. J. Neurosci. 2718–2792.

Chen, Y., Geisler, W.S., and Seidemann, E. (2008). Optimal temporal decoding of neural
population responses in a reaction-time visual detection task. J. Neurophysiol. *99*, 1366–1379.

Chen, Y., Palmer, C.R., and Seidemann, E. (2012). The relationship between voltage-sensitive
dye imaging signals and spiking activity of neural populations in primate V1. J. Neurophysiol.
*107*, 3281–3295.

David, S. V., Hayden, B.Y., and Gallant, J.L. (2006). Spectral Receptive Field Properties
Explain Shape Selectivity in Area V4. J. Neurophysiol. *96*, 3492–3505.

DeAngelis, G.C., and Newsome, W.T. (1999). Organization of disparity-selective neurons in
macaque area MT. J Neurosci *19*, 1398–1415.

Desimone, R., Moran, J., Mishkin, M., and Schein, S.J. (1993). A role for the corpus callosum in
visual area V4 of the macaque. Vis. Neurosci.

DiCarlo, J.J., Zoccolan, D., and Rust, N.C. (2012). How does the brain solve visual object
recognition? Neuron *73*, 415–434.

Van Doorn, A.J., and Koenderink, J.J. (1984). Spatiotemporal integration in the detection of
coherent motion. Vision Res. *24*, 47–53.

Engel, A.K., and Fries, P. (2010). Beta-band oscillations--signalling the status quo? Curr. Opin.
Neurobiol. *20*, 156–165.

Van Essen, D.C., and Newsome, W.T. (1984). The visual field representation in the striate cortex
of the macaque monkey: Asymmetries, anisptropies, and indiviual variability. Vision Res. *24*,
429–448.

Florence, S.L., Casagrande, V.A., and Biology, C. (1988). Organization of Individual Cortex in a
Primate Afferent Axons in Layer IV of Striate. *7*.

Fredericksen, R.E., Verstraten, F.A., and van de Grind, W.A. (1994). Spatial summation and its
interaction with the temporal integration mechanism in human motion perception. Vision Res.
*34*, 3171–3188.

Fukushima, K. (1980). Neocognitron: A Self-organizing Neural Network Model for a
Mechanism of Pattern Recognition Unaffected by Shift in Position. Biol. Cybern. *202*, 193–202.

Geisler, W.S. (2008). Visual Perception and the Statistical Properties of Natural Scenes. Annu.
Rev. Psychol. *59*, 167–192.

Georges, S., Seriès, P., Frégnac, Y., and Lorenceau, J. (2002). Orientation dependent modulation

of apparent speed: psychophysical evidence. Vision Res. *42*, 2757–2772.

Girard, P., Hupé, J.M., and Bullier, J. (2001). Feedforward and feedback connections between
areas V1 and V2 of the monkey have similar rapid conduction velocities. J. Neurophysiol. *85*,
1328–1331.

Grinvald, A., and Hildesheim, R. (2004). VSDI: a new era in functional imaging of cortical
dynamics. Nat. Rev. Neurosci. *5*, 874–885.

Grinvald, A., Lieke, E.E., Frostig, R.D., and Hildesheim, R. (1994). Cortical point-spread
function and long-range lateral interactions revealed by real-time optical imaging of macaque
monkey primary visual cortex. J. Neurosci. *14*, 2545–2568.

Gross, C.G., Bender, D.B., and Mishkin, M. (1977). Contributions of the corpus callosum and
the anterior commissure to visual activation of inferior temporal neurons. Brain Res.

Grzywacz, N.M., Watamaniuk, S.N., and McKee, S.P. (1995). Temporal coherence theory for
the detection and measurement of visual motion. Vision Res. *35*.

Guo, K., Nevado, A., Robertson, R.G., Pulgarin, M., Thiele, A., and Young, M.P. (2004). Effects
on orientation perception of manipulating the spatio-temporal prior probability of stimuli. Vision
Res. *44*, 2349–2358.

Guo, K., Robertson, R.G., Pulgarin, M., Nevado, A., Panzeri, S., Thiele, A., and Young, M.P.
(2007). Spatio-temporal prediction and inference by V1 neurons. Eur. J. Neurosci. *26*, 1045–
1054.

Hays, A.V.J., Richmond, B.J., and Optican, L.M. (1982). Unix-based multiple-process system,
for real-time data acquisition and control.

Hubel, D.H., and Wiesel, T.N. (1974). Uniformity of monkey striate cortex: a parallel
relationship between field size, scatter, and magnification factor. J. Comp. Neurol. *158*, 295–305.

Jancke, D., Chavane, F., Naaman, S., and Grinvald, A. (2004a). Imaging cortical correlates of
illusion in early visual cortex. Nature *428*, 423–426.

Jancke, D., Erlhagen, W., Schöner, G., and Dinse, H.R. (2004b). Shorter latencies for motion
trajectories than for flashes in population responses of cat primary visual cortex. J. Physiol. *556*,
971–982.

Judge, S.J., Richmond, B.J., and Chu, F.C. (1980). Implantation of magnetic search coils for
measurement of eye position: An improved method. Vision Res. *20*, 535–538.

Kaplan, B. a., Lansner, A., Masson, G.S., and Perrinet, L.U. (2013). Anisotropic connectivity
implements motion-based prediction in a spiking neural network. Front. Comput. Neurosci. *7*, 1–
14.

Kennedy, H., and Dehay, C. (1988). Functional implications of the anatomical organization of
the callosal projections of visual areas V1 and V2 in the macaque monkey. Behav Brain Res *29*,
225–236.

Kennedy, H., Dehay, C., and Bullier, J. (1986). Organization of the callosal connections of visual
areas V1 and V2 in the macaque monkey. J. Comp. Neurol. *247*, 398–415.

van Kerkoerle, T., Self, M.W., Dagnino, B., Gariel-Mathis, M. -a., Poort, J., van der Togt, C.,
and Roelfsema, P.R. (2014). Alpha and gamma oscillations characterize feedback and
feedforward processing in monkey visual cortex. Proc. Natl. Acad. Sci.

Khoei, M.A., Masson, G.S., and Perrinet, L.U. (2017). The Flash-Lag Effect as a Motion-Based
Predictive Shift. PLoS Comput. Biol. *13*, 1–31.

Klink, P.C., Dagnino, B., Gariel-Mathis, M.A., and Roelfsema, P.R. (2017). Distinct
Feedforward and Feedback Effects of Microstimulation in Visual Cortex Reveal Neural
Mechanisms of Texture Segregation. Neuron *95*, 209-220.e3.

Lenkic, P.J., and Enns, J.T. (2013). Apparent Motion Can Impair and Enhance Target Visibility:
The Role of Shape in Predicting and Postdicting Object Continuity. Front. Psychol.

Liang, H., Gong, X., Chen, M., Yan, Y., Li, W., and Gilbert, C.D. (2017). Interactions between
feedback and lateral connections in the primary visual cortex. Proc. Natl. Acad. Sci. *114*, 8637–
8642.

Liu, J. V., Ashida, H., Smith, A.T., and Wandell, B.A. (2006). Assessment of Stimulus-Induced
Changes in Human V1 Visual Field Maps. J. Neurophysiol.

Lorenceau, J., Shiffrar, M., Wells, N., and Castet, E. (1993). Different motion sensitive units are
involved in recovering the direction of moving lines. Vision Res. *33*, 1207–1217.

MacKay, D.. (1958). Perceptual stability of a stroboscopically lit visual field containing self-
luminous objects. Nature *181*, 507–508.

Marr, D. (1982). Vision: A computational investigation into the human representation and
processing of visual information. (San Francisco).

McKee, S.P., and Welch, L. (1985). Sequential recruitment in the discrimination of velocity. J.
Opt. Soc. Am. A. *2*, 243–251.

Meirovithz, E., Ayzenshtat, I., Bonneh, Y.S., Itzhack, R., Werner-Reiss, U., and Slovin, H.
(2010). Population response to contextual influences in the primary visual cortex. Cereb. Cortex
*20*, 1293–1304.

Movshon, J. a, Adelson, E.H., Gizzi, M.S., and Newsome, W.T. (1985). The analysis of moving
visual patterns. Pattern Recognit. Mech. *54*, 117–151.

Muller, L., Reynaud, A., Chavane, F., and Destexhe, A. (2014). The stimulus-evoked population
response in visual cortex of awake monkey is a propagating wave. Nat. Commun. *5*, 3675.

Muller, L., Chavane, F., Reynolds, J., and Sejnowski, T.J. (2018). Cortical travelling waves:
Mechanisms and computational principles. Nat. Rev. Neurosci.

Mumford, D. (1992). On the computational architecture of the neocortex. Biol. Cybern. *66*, 241–
251.

Nakayama, K., and Silverman, G. (1984). Temporal and spatial characteristics of the upper
displacement limit for motion in random dots. Vision Res.

Nassi, J.J., Lomber, S.G., and Born, R.T. (2013). Corticocortical feedback contributes to
surround suppression in V1 of the alert primate. J. Neurosci. *33*, 8504–8517.

Neri, P. (2014). Dynamic Engagement of Human Motion Detectors across Space-Time
Coordinates. J. Neurosci. *34*, 8449–8461.

Nijhawan, R. (1994). Motion extrapolation in catching. Nature *370*, 256–257.

Orban, G.A., Hoffmann, K.P., and Duysens, J. (1985). Velocity selectivity in the cat visual
system. I. Responses of LGN cells to moving bar stimuli: a comparison with cortical areas 17

and 18. J. Neurophysiol. *54*, 1026–1049.

Pavan, A., Casco, C., Mather, G., Bellacosa, R.M., Cuturi, L.F., and Campana, G. (2011). The
effect of spatial orientation on detecting motion trajectories in noise. Vision Res. *51*, 2077–2084.

Perkel, D.J., Bullier, J., and Kennedy, H. (1986). Topography of the afferent connectivity of area
17 in the macaque monkey: a double-labelling study. J. Comp. Neurol. *253*, 374–402.

Perrinet, L.U., and Masson, G.S. (2012). Motion-based prediction is sufficient to solve the
aperture problem. Neural Comput. *10*, 24.

Piech, V., Li, W., Reeke, G.N., and Gilbert, C.D. (2013). Network model of top-down influences
on local gain and contextual interactions in visual cortex. Proc. Natl. Acad. Sci. *110*, E4108–
E4117.

Pigarev, I.N., Nothdurft, H.C., and Kastner, S. (2001). Neurons with large bilateral receptive
fields in monkey prelunate gyrus. Exp. Brain Res.

Ramachandran, V.S., and Anstis, S.M. (1983). Extrapolation of motion path in human visual
perception. Vision Res. *23*, 83–85.

Ramalingam, N., McManus, J.N.J., Li, W., and Gilbert, C. (2013). Top-down modulation of
lateral interactions in visual cortex. J. Neurosci. *33*, 1773–1789.

Rao, R.P., and Ballard, D.H. (1999). Predictive coding in the visual cortex: a functional
interpretation of some extra-classical receptive-field effects. Nat. Neurosci. *2*, 79–87.

Reynaud, A., Takerkart, S., Masson, G.S., and Chavane, F. (2011). Linear model decomposition
for voltage-sensitive dye imaging signals: application in awake behaving monkey. Neuroimage
*54*, 1196–1210.

Reynaud, A., Masson, G.S., and Chavane, F. (2012). Dynamics of local input normalization
result from balanced short- and long-range intracortical interactions in area V1. J. Neurosci. *32*,
12558–12569.

Riesenhuber, M., and Poggio, T. (1999). Hierarchical models of object recognition in cortex.
Nat. Neurosci. *2*, 1019–1025.

Roach, N.W., McGraw, P. V., and Johnston, A. (2011). Visual motion induces a forward
prediction of spatial pattern. Curr. Biol. *21*, 740–745.

Robinson, D.A. (1963). A Method of Measuring Eye Movemnent Using a Scieral Search Coil in
a Magnetic Field. Ire Trans. Biomed. Electron. *10*, 137–145.

Roelfsema, P.R. (2006). Cortical Algorithms for Perceptual Grouping. Annu. Rev. Neurosci. *29*,
203–227.

Rust, N.C., Mante, V., Simoncelli, E.P., and Movshon, J.A. (2006). How MT cells analyze the
motion of visual patterns. Nat. Neurosci. *9*, 1421–1431.

Schwiedrzik, C.M., Alink, A., Kohler, A., Singer, W., and Muckli, L. (2007). A spatio-temporal
interaction on the apparent motion trace. Vision Res. *47*, 3424–3433.

Scott-Brown, K.C., and Heeley, D.W. (2001). The effect of the spatial arrangement of target
lines on perceived speed. Vision Res. *41*, 1669–1682.

Shoham, D., Glaser, D.E., Arieli, A., Kenet, T., Wijnbergen, C., Toledo, Y., Hildesheim, R., and
Grinvald, A. (1999). Imaging cortical dynamics at high spatial and temporal resolution with

novel blue voltage-sensitive dyes. Neuron *24*, 791–802.

Simoncelli, E.P., and Heeger, D.J. (1998). A model of neuronal responses in visual area MT.
Vis. {R}esearch *38*, 743–761.

Slovin, H., Arieli, A., Hildesheim, R., and Grinvald, A. (2002). Long-term voltage-sensitive dye
imaging reveals cortical dynamics in behaving monkeys. J. Neurophysiol. *88*, 3421–3438.

Snowden, R., and Braddick, O. (1989a). Extension of displacement limits in multiple-exposure
sequences of apparent motion. Vision Res. *29*, 1777–1787.

Snowden, R.J., and Braddick, O.J. (1989b). The combination of motion signals over time. Vision
Res. *29*, 1621–1630.

Srinivasan, M. V, Laughlin, S.B., and Dubs, A. (1982). Predictive coding: a fresh view of
inhibition in the retina. Proc. R. Soc. Lond. B. Biol. Sci. *216*, 427–459.

Stettler, D.D., Das, A., Bennett, J., and Gilbert, C. (2002). Lateral connectivity and contextual
interactions in macaque primary visual cortex. Neuron *36*, 739–750.

Subramaniyan, M., Ecker, A.S., Patel, S.S., Cotton, R.J., Bethge, M., Berens, P., and Tolias, S.A.
(2015). Faster processing of moving compared to flashed bars in awake macaque V1 provides a
neural correlate of the flash lag illusion. BioRxiv Prepr. 1–26.

Sundberg, K.A., Fallah, M., and Reynolds, J.H. (2006). A motion-dependent distortion of
retinotopy in area V4. Neuron *49*, 447–457.

Swindale, N. V (1998). Biological Cybernetics Orientation tuning curves : empirical description
and estimation of parameters. *56*, 45–56.

Takerkart, S., Fenouil, R., Piovano, J., Reynaud, A., Hoffart, L., Chavane, F., Papadopoulo, T.,
Conrath, J., and Masson, G.S. (2008). A quantification framework for post-lesion neo-
vascularization in retinal angiography. In 2008 5th IEEE International Symposium on
Biomedical Imaging: From Nano to Macro, Proceedings, ISBI, pp. 1457–1460.

Tlapale, E., Masson, G.S., and Kornprobst, P. (2010). Modelling the dynamics of motion
integration with a new luminance-gated diffusion mechanism. Vision Res. *50*, 1676–1692.

Tootell, R.B.H., Silverman, M.S., Switkes, E., and De Valois, R.L. (1982). Deoxyglucose
analysis of retinotopic organization in primate striate cortex. Science (80-. ).

Ullman, S. (1979). The interpretation of structure from motion. Proc. R. Soc. Lond. B. Biol. Sci.
*203*, 405–426.

Verghese, P., and Watamaniuk, S. (1999). Local motion detectors cannot account for the
detectability of an extended trajectory in noise. Vision Res. *39*, 19–30.

Verghese, P., McKee, S.P., and Grzywacz, N.M. (2000). Stimulus configuration determines the
detectability of motion signals in noise. J. Opt. Soc. Am. A. Opt. Image Sci. Vis. *17*, 1525–1534.

W. Metzger (1932). No TitleVersuch einer gemeinsamen Theorie der Phänomene Fröhlichs und
Hazelhoffs und Kritik ihrer Verfahren zur Messung der Empfindungszeit. Psychol. Forsch. *16*,
176–200.

Wallach, H. (1935). Über visuell wahrgenommene Bewegungsrichtung. Psychol. Forsch. *20*,
325–380.

Watamaniuk, S.N., and McKee, S.P. (1995). Seeing motion behind occluders. Nature *377*, 729–

730.

Watamaniuk, S.N., McKee, S.P., and Grzywacz, N.M. (1995). Detecting a trajectory embedded
in random-direction motion noise. Vision Res. *35*, 65–77.

Watanabe, J., and Nishida, S. (2007). Veridical perception of moving colors by trajectory
integration of input signals. J. Vis. *7*, 3.1-16.

Welch, L., Macleod, D.I., and Mckee, S.P. (1997). Motion interference: perturbing perceived
direction. Vision Res. *37*, 2725–2736.

Werkhoven, P., Snippe, H.P., and Koenderink, J.J. (1990). Effects of element orientation. *47*,
509–525.

Wertheimer, M. (1912). Experimentelle studien über das sehen von bewegung. Z. Psychol. *61*.

Wilson, H.R., Ferrera, V.P., and Yo, C. (1992). A psychophysically motivated model for two-
dimensional motion perception. Vis. Neurosci. *9*, 79–97.

Yuille, A.L., and Grzywacz, N.M. (1989). A mathematical analysis of the motion coherence
theory. Int. J. Comput. Vis. *3*, 155–175.

1277

1278    **SUPPLEMENTARY MATERIAL**

1279

1280    **Figure supplement 1.**

1281
**Suppl. Figure 1: Average PSTHs across non-anticipatory neurons for the three trajectories**.
(**A**) Average PSTHs across non-anticipatory neurons for the three trajectories for monkey PD (n=20).
Responses are aligned to the latency of the short trajectory condition. Color lines (bottom) indicate the
stimulus presentation time, for each condition. No significant difference with respect to the short
trajectory condition were detected (Two-sample t-test P>0.05). (**B**) Same than A but for monkey NO
(n=3).

1288

1289

1290

**Figure supplement 2: Anticipation is not caused by eye movements or excessively large RF**

In order to completely exclude potential effects of eye movements or changes in receptive field
size to the observed anticipation, we conducted several control experiments. First, we computed
the distribution of eye positions across all successful trials of the 3-trajectories paradigm for 22
cells (a smaller number of cells is used here because the eye-movements recorded in some
experimental sessions were deteriorated due to an error in the data acquisition software). Our
results show that 99% of eye positions were distributed within 1dva around the fixation spot and
with identical distribution for the 3 conditions (**Supp. Figure 2A**). Thus, any changes in RF
location due to eye movements would have been much smaller than the 2.3 (PD) and 3 dva (NO)
equivalent distance reported for the average anticipatory responses in the two monkeys.
1301
1302   Second, comparing the skewness of the distributions in Figure 1A for long (y-axis) and short
1303   trajectory conditions (x-axis), we showed that there is not difference in the slant towards of
1304   (negative values) or away from (positive values) the direction of the approaching bar (**Supp.**
1305   **Figure 2B**).
1306
Third, we recomputed mean response profiles as a function of the actual distance of the bar from
the RF center when considering the real-time position of eye fixation (acquisition rate 2Hz)
(**Supp. Figure 2C**). This analysis (A-C) was applied to the 22 cells for which we had high
quality eye-movements recordings. The anticipatory responses for medium (3dva) and long
(6dva) trajectories were still significant in all the anticipatory cells tested (n=7), rejecting the
possibility that they could be a mere consequence of a systematic bias or transient events in
fixation behavior. On the contrary, the overall response across the 22 cells tested presents an
increase in the anticipatory response after correcting for the real-time position of the eye.
1315
Fourth, we confirmed that anticipatory responses are not simply explained by a sampling bias for
large receptive fields (**Supp. Figure 2D, E, F**). The anticipation is plotted in spatial coordinate
as a function of the receptive field radius for medium (1D) and long (1E) trajectories. The
anticipation started systematically outside of the classical RF (above the main diagonal) and
most of the points are also further away than twice the size of the classical RF (above y=2x). We
also compared the p-value of the statistical test for the significance of the anticipatory response
(relative to baseline) and showed that there is not trends, actually the smallest p-values are
observed for the smallest RF.
1324

**Suppl. Figure 2: Test existence of biases from eye movements and RF sizes in the**
**emergence of anticipatory responses.** (**A**) Gaze positions distribution across all successful trials
for all 3-trajectories conditions (black), thin traces are single cells, thick trace the average, which
are plotted for the 3 conditions in the same graph in the bottom (n = 22 cells). (**B**) for all cells we
measured the skewness of the distribution in A to test whether a slant towards (negative) or away
(positive values) of the direction of the approaching bar was existing for long trajectories (y-axis)
compared to the short trajectory condition (x-axis). The average and std across cells is presented
as error bars. (**C**) Eye movement correction control. Average PSTHs among cells (n=22 cells)
for the three conditions before and after eye movement correction of the spikes timing relative to
the exact distance between the receptive field and the bar. Neural response is therefore expressed
in number of spikes per degree of visual angle of the bar displacement (spk/dva), assuming a
constant latency of 50ms. Upper left inset sketch the RF position correction (grey before, dark
after correction) due to eye position (red dot) relative to the central fixation ('+'), allow to
compute the exact distance (d in dva) between the bar and the RF. Colored lines (top) indicate
significant difference with respect to the short trajectory condition (two-sample t-test P<0.05)
before the bar enters the RF (i.e. 0 dva). (**D-E**) Position of the bar at response onset (y-axis) as a
function of RF size for the medium and long-trajectory conditions. The main diagonal (y=x) and
twice the main diagonal (y=2x) are shown to compare the position of the bar that evoked
anticipatory activity with extreme limits of RF when mapped using classical procedures. (**F**) p-
value of the anticipatory activity t-test (relative to spontaneous activity) as a function of its RF
size (n = 36).

1347

1348

1349

1350

1351

1352

1353

1354

1355

1356

1357

1358

1359

1360

1361     **Figure supplement 3.**

1362

**Suppl. Fig 3: Block diagram sketch of the computational model.** From left to right is depicted
the logical flow of our model. The matrices of the diagram all represents space in y and time in
x. On the left is represented the feedforward predicted activation of V1 to a bar moving at
6.6dva/s other an extended trajectory. At that level, random noise is added to the simulated
activation. To generate horizontal or feedback activation maps, the feedforward matrix is
convolved with kernels with low extent low speed (horizontal) and large extent fast speed
(feedback), and then weighted by $w_h$ and $w_{fb}$. The feedback activation map is then added a scalar
of 1 and multiplied to the horizontal activation map to amplify of the horizontal activation
(Liang et al., 2017; Ramalingam et al., 2013, Piech et al 2013). The resulting activation map is
then passing through a sigmoid non-linearity controlled by the difference between the
feedforward and the horizontal-feedback activation maps. The relative contribution of the
feedforward and horizontal-feedback maps are then summed with a ratio of $\rho$, to generate the
predicted V1 activity.