Evaluating and integrating spatial capture-recapture models with data of

variable individual identifiability

JOEL S. RUPRECHT[1,*], CHARLOTTE E. ERIKSSON[1], TAVIS D. FORRESTER[2], DARREN A. CLARK[2],

MICHAEL J. WISDOM[3], MARY M. ROWLAND[3], BRUCE K. JOHNSON[2,†], AND TAAL LEVI[1]

[1]*Department of Fish and Wildlife, 104 Nash Hall, Oregon State University, Corvallis, OR 97331*

[2]*Oregon Department of Fish and Wildlife, 1401 Gekeler Lane, La Grande, OR 97850*

[3]*Pacific Northwest Research Station, USDA Forest Service, 1401 Gekeler Lane, La Grande, OR*

97850

†Retired

*Email: joel.ruprecht@oregonstate.edu

Abstract. Spatial capture-recapture (SCR) models have become the preferred tool for estimating

densities of carnivores. Within this family of models are variants requiring identification of all

individuals in each encounter (SCR), a subset of individuals only (generalized spatial mark-

resight, gSMR), or no individual identification (spatial count or spatial presence-absence).

Although each technique has been shown through simulation to yield unbiased results, the

consistency and relative precision of estimates across methods in real-world settings are seldom

considered. We tested a suite of models ranging from those only requiring detections of

unmarked individuals to others that integrate remote camera, physical capture, genetic, and

global positioning system (GPS) data into a 'hybrid' model, to estimate population densities of

black bears, bobcats, cougars, and coyotes. For each species we genotyped fecal DNA collected

with detection dogs during a 20-day period. A subset of individuals from each species was

affixed with GPS collars bearing unique markings and resighted by remote cameras over 140

days contemporaneous with scat collection. Camera-based gSMR models produced density

estimates that differed by less than 10% from genetic SCR for bears, cougars, and coyotes once

important sources of variation (sex or behavioral status) were controlled for. For bobcats, SCR

estimates were 33% higher than gSMR. The cause of the discrepancies in estimates was likely

attributable to challenges designing a study compatible for species with disparate home range

sizes and the difficulty of collecting sufficient data in a timeframe in which demographic closure

could be assumed. Unmarked models estimated densities that varied greatly from SCR, but

estimates became more consistent in models wherein more individuals were identifiable. Hybrid

models containing all data sources exhibited the most precise estimates for all species. For

studies in which only sparse data can be obtained and the strictest model assumptions are

unlikely to be met, we suggest researchers use caution making inference from models lacking

individual identity. For best results, we further recommend the use of methods requiring at least

a subset of the population is marked and that multiple datasets are incorporated when possible.

Key words: abundance, black bear, bobcat, camera trapping, carnivore, count, cougar, coyote,

density estimation, mark-resight, noninvasive genetic sampling, spatial capture-recapture

INTRODUCTION

Population abundance is a state variable of paramount importance in both applied and

basic ecology but remains a challenge to estimate for most free-ranging animal populations. This

is particularly the case for terrestrial carnivore species that are cryptic and occur at low densities

(Wilson and Delahay 2001). Many statistical models accounting for imperfect detection have

been used to estimate abundance, but the data required for each method vary along a continuum

of identifiability of individual animals, ranging from no individual recognition to full individual

recognition. For situations in which no animals can be uniquely identified, models to estimate

abundance have been developed based on presence-absence of unmarked individuals in a

sampling session (Royle and Nichols 2003, Ramsey et al. 2015) or counts of unmarked

individuals (Buckland et al. 1993, Royle 2004, Rowcliffe et al. 2008, Chandler and Royle 2013,

Moeller et al. 2018). Other models rely on partially-observable encounter histories when a subset

of the population is identifiable (e.g. White 1996, McClintock and White 2009, Sollmann et al.

2013a, Augustine et al. 2018), and the most data-intensive models utilize fully-observable

encounter histories such that all individuals can be uniquely identified (e.g., Otis et al. 1978,

Nichols 1992, Efford 2004, Royle et al. 2013). Collecting higher resolution data where

individuals are identifiable is substantially more challenging and costly, but models based on

these data may produce more robust estimates of abundance particularly in situations with

suboptimal study designs or when model assumptions cannot strictly be upheld (Chandler and

Royle 2013, Augustine et al. 2019). Knowing what type and how much data to collect to achieve

satisfactory results is therefore a central challenge to estimate population abundance for

conservation, management, or research objectives.

Spatial capture-recapture (SCR; Efford 2004, Borchers and Efford 2008, Royle and

Young 2008, Royle et al. 2013) has emerged as the most prominent method of abundance

estimation for carnivores over the past five years (Appendix S1: Fig. S1). SCR models exploit

the spatial locations of animal detections and assume the rate or probability an individual is

detected is highest at its home range center and declines as distance from that location increases

(Royle and Young 2008). The spatially-referenced detections of an individual allow the

estimation of the animal's activity center, and the number of estimated activity centers yields an

estimate of abundance. Population density, often preferable to abundance, is calculated by

dividing the estimated number of activity centers by the area of the state space which eliminates

the arbitrarily-defined effective sampling area required in traditional capture-mark-recapture

approaches (Royle et al. 2013). Furthermore, SCR can accommodate individual and trap-level

covariates to explain spatial and individual variation in detection rates.

SCR models are well-suited for noninvasive genetic sampling in which individuals are

uniquely identified, typically using DNA from scat or hair samples, to produce fully-observable

encounter histories (Gardner et al. 2009, Kery et al. 2011, Morin et al. 2016, 2018). Scat-

detection dogs in particular have become a common tool to efficiently collect scat on large

landscapes within a narrow time window to ensure demographic closure even for species that

occur at low densities (Wasser et al. 2004, 2011). While effective for SCR density estimation,

scat detection requires highly-trained dogs and specialization in genetic techniques or the

resources to contract out those tasks. As a more inexpensive and accessible alternative, wildlife

researchers increasingly use remote cameras to study and monitor wildlife. However, cameras

can only be used for conventional SCR analyses when all individuals of a species possess natural

markings (e.g. unique pelage patterns) that permit individual identification (an exception is the

spatial partial identity model; Augustine et al. 2018).

Spatial mark-resight (SMR) and generalized spatial mark-resight (gSMR) models are

variants of SCR that allow spatially-explicit density estimation when only a subset of the

population can be uniquely identified (Chandler and Royle 2013, Sollmann et al. 2013a,

Whittington et al. 2018). A common example is identification of individual study animals that

are marked with GPS (global positioning system) collars or other telemetry devices. For many

species it would be inefficient to capture and mark animals solely for the purpose of density

estimation; however, many researchers and wildlife management agencies routinely monitor

populations with telemetry devices. In these cases, resighting marked animals on an array of

remote cameras may offer a relatively straightforward option for estimating density at little extra

cost (Whittington et al. 2018).

Due to the challenge of individually marking animals, spatial presence-absence (SPA;

Ramsey et al. 2015) and spatial count (SC) or 'unmarked SCR' models (Chandler and Royle

2013) have been developed to exploit spatial autocorrelation in counts of unmarked animals to

make inference on their activity centers, and by extension, population density. The attractiveness

of these approaches is that they require no individual identification so the population need not be

artificially nor naturally marked. Theoretically, the only data required for SPA or SC models are

spatially-referenced binary detections or counts of animal detections, respectively, which allows

use of standard camera trapping datasets that can be collected noninvasively without the cost and

specialized equipment needed for genotyping. Although Chandler and Royle (2013) caution

researchers to mark a subset of the population whenever possible, others have successfully

estimated population densities using datasets lacking any individual recognition (e.g., Evans and

Rittenhouse 2018). If this were universally true, SC methods would represent a major advance in

our ability to rapidly and affordably count carnivores.

These three model types (i.e., SC, SMR/gSMR, and SCR) require data of vastly different

levels of individual identifiability which raises concerns over the consistency of results one may

observe if using multiple methods. Understanding the relative performance of these commonly-

used density estimators is critical to guide future population monitoring for conservation and

management. We hypothesized, *a priori*, that genetic SCR methods would yield the most robust

density estimates because they require fully-observable encounter histories. As the use of gSMR,

SMR, and SC methods proliferates in the literature, however, it is important to evaluate their

credibility with respect to the 'gold standard' SCR approach. A variety of additional data sources

can be incorporated into these models, which could potentially increase the concordance of

results from unmarked models to fully-marked SCR models while also increasing precision and

accuracy. For example, all of these models can be informed by telemetry data in various

capacities (Sollmann et al. 2013a) to provide independent information on detection parameters,

and the SMR model should additionally be informed by the capture process used to mark

animals to reduce bias (i.e. generalized SMR; Whittington et al. 2018). Since research projects

and monitoring programs have limited budgets, it is also critical to know how much and what

type of data must be collected to achieve satisfactory results.

Here we use a unique dataset consisting of 1) genotyped scats located by detection dogs,

2) images of both marked and unmarked animals from an array of remote cameras, 3) encounter

histories from live-trapping, and 4) GPS collar locations to estimate the abundances of four

carnivore species: black bears (*Ursus americanus*), bobcats (*Lynx rufus*), cougars (*Puma*

concolor), and coyotes (*Canis latrans)*. Our objective is to evaluate a suite of existing spatial

density models across a range of individual identifiability to evaluate agreement in results and

precision under realistic data collection scenarios. In addition, we present a new hybrid model

that can incorporate SCR and gSMR data sources (i.e. including genotypes, the physical marking

process, camera data, and GPS data) into a single hierarchical model, and assess the resulting

gains in precision. Notably our model comparisons occur in the context of a real system rather

than simulation. Each species varies in its life history, from highly territorial but group living

(coyotes), to non-territorial solitary individuals (black bears), to solitary territorial individuals

with territory size varying by sex (bobcat females are territorial while males are not; cougar

males are territorial while females are not). By evaluating a suite of methods with different data

requirements, our findings directly address how population densities of terrestrial carnivores can

most efficiently be estimated, and whether the investment in genotyping or marking and

resighting animals is warranted given the availability of models that do not rely on individual

identification.

METHODS

Study Area

Our study was conducted in and adjacent to the Starkey Experimental Forest and Range

(hereafter Starkey) in the Blue Mountains of northeastern Oregon (Fig. 1). Starkey is bounded by

a 2.4-meter fence, enclosing 100.7 km^2, which prevents the passage of large herbivores

(Rowland et al. 1997); however, carnivores are undeterred by the fence and regularly cross in

and out of the enclosure (this study, Oregon Department of Fish and Wildlife, unpublished data).

Starkey is part of the Wallowa-Whitman National Forest and is administered by the US Forest

Service. Land immediately adjacent to Starkey is predominantly public and managed by the US

Forest Service (Wallowa-Whitman and Umatilla National Forests) but also includes private

inholdings. The study area contains a mosaic of Ponderosa pine (*Pinus ponderosa*) and mixed

pine-fir forests (*Pinus*, *Abies* and *Pseudotsuga* spp.), interspersed with grasslands dominated by

native bunchgrasses (*Poa*, *Danthonia*, and *Pseudoroegneria* spp.) and invasive annual grasses

(*Bromus* and *Ventenata* spp.) (Rowland et al. 1997). Elevation ranges between 1,122 and 1,500

m within Starkey (Rowland et al. 1997). See Wisdom et al. (2005) for additional details about

Starkey.

Data collection

Scat dataset

Three teams of scat detection dogs from the University of Washington Conservation

Canine program surveyed a 224 km^2 study area encompassing all of Starkey and roughly an

equal area immediately to the north and west of the enclosure between 6 and 26 June 2017 (Fig.

1). The area surveyed was composed of 56 grid cells each with an area of 4 km^2. Detection dogs

surveyed 6–8 km linear distance within each cell to distribute effort across the study area. Dog

handlers were not given specific survey routes but were encouraged to follow natural travel

corridors such as ridgelines, saddles, drainage bottoms, game trails, and fencelines. No more than

50% of the distance traveled per grid cell was permitted to be on linear features such as trails or

roads. When scats were located, the handler recorded the GPS position and placed the entire scat

in triplicate paper bags (Fig. 2a). Within 72 hours of collection, scats were desiccated in a drying

oven for 24 hours at 40°C (Murphy et al. 2000). Detection dogs in our study were trained to

locate black bear, coyote, cougar, and bobcat scats.

We used genotyping by amplicon sequencing (Eriksson et al. 2020) to identify unique

individuals with single-nucleotide polymorphisms (SNPs), a process that uses the power of high-

throughput sequencers to yield high genotype success rates and low error rates. Coyote scats

were genotyped following methods in Eriksson et al. (2020). We genotyped bobcat, cougar, and

black bears scat using the same approach but with novel primers designed and tested specifically

for this study (Appendix S2). Prior to genotyping, the identity of the defecator for each scat was

identified using DNA metabarcoding of ~100 base pairs of the mitochondrial 12S gene region as

part of a separate diet study (primers used in Eriksson et al. 2020, Massey et al. 2020, and

modified from Riaz et al. 2011). See Appendix S2 for SNP discovery methods and Eriksson et

al. (2020) for detailed primer design and genotyping protocols.

Camera dataset

A 1- x 1-km grid of 94 unbaited remote cameras (Bushnell TrophyCam Aggressor,

Overland, KS) was placed inside the Starkey fence perimeter and was operational between April

and September 2017 (Fig. 1). All cameras were pointed north. We visited each camera station

every 6-8 weeks to change batteries and download data. Each camera was set to take a burst of 3

photos when triggered.

Photos were tagged to species or to individual when animals were uniquely identifiable

due to GPS collars and ear tags. We used the open source photo management software DigiKam

(www.digikam.org) for tagging and database management and the R library camtrapR (Niedballa

et al. 2016) to extract metadata from photos. We censored photos in which we could not

determine whether an animal was wearing a GPS collar or if we could not verify the identity of

an animal photographed with a GPS collar. However, these records constituted < 5% of all

detections. Because animals were typically photographed more than once during a visit, we

recorded information for each independent photo sequence and considered photo sequences > 30

minutes from the next detection of the same species to be independent.

Telemetry dataset

We captured and GPS-collared a sample of individuals from each of the four carnivore

species as part of concurrent research on predator-prey relationships. Coyotes, bobcats, and bears

were captured inside Starkey only, and cougars were captured in Starkey plus a buffer of

approximately 20 km which was encompassed by the state space. Coyotes were captured using

padded foothold traps (Oneida Victor SoftCatch No. 3, Euclid, OH) with a tranquilizer tab

(Balser 1965) containing 50 mg propriopromazine hydrochloride attached to each trap to subdue

captured animals until researchers arrived. Captured animals were immobilized with tiletamine-

zolazepam (Telazol®) at a concentration of 10 mg/kg. A GPS collar (Lotek MiniTrack, Lotek

Wireless Inc., Newmarket, ON, Canada or Vectronic Vertex, Vectronic Aerospace GmbH,

Berlin, Germany) was placed on each adult coyote and was programmed to record locations

every 2 or 3 hours.

We captured bobcats using cage traps baited with visual and olfactory attractants.

Captured bobcats were administered ketamine (10 mg/kg) and xylazine (1.5 mg/kg) for

immobilization, and upon release, yohimbine (0.125 mg/kg; Yobine®) was given as an

antagonist for xylazine. Each bobcat was fit with a GPS collar (Lotek MiniTrack, Lotek Wireless

Inc., Newmarket, ON, Canada) scheduled to take fixes every 2 hours.

Black bears were captured using culvert traps or padded foot snares (Lemieux and

Czetwertynski 2006). Captured bears were immobilized with Telazol® at a concentration of 7

mg/kg. Bears were fit with GPS collars (Lotek GPS 7000 or LiteTrack Iridium 420, Lotek

Wireless Inc., Newmarket, ON, Canada) that recorded positions every 15 min to 2 hours.

We captured cougars using trained pursuit hounds. We systematically searched roads for

fresh cougar tracks in snow and when located, released hounds to pursue tracks until the cougar

was treed. Dogs were allowed to pursue tracks of any individual cougar. When treed, cougars

were immobilized with ketamine (10 mg/kg) and xylazine (2 mg/kg) via dart gun, and before

release were administered yohimbine (0.125 mg/kg; Yobine®) as an antagonist for xylazine. A

GPS collar (Lotek GPS 4400S, IridiumTrack M, Lotek Wireless Inc., Newmarket, ON, Canada

or Vectronic Vertex Lite, Vectronic Aerospace GmbH, Berlin, Germany) was placed on each

cougar and was programmed to record locations every 3 hours.

We permanently marked the belting of GPS collars with unique numbers used to identify

individuals on camera (Fig. 2b). When possible (i.e., for the larger collars), we also affixed

sections of colored heat-shrink plastic tubing over the collar belting to assist with identification

(Fig. 2c). We ear-tagged each captured animal such that females received ear tags on the left ear

and males on the right ear which provided an additional cue when identifying GPS-collared

animals on camera.

We collected biological samples from each captured individual for SNP primer

development and to match the genotypes of GPS-collared animals with genotypes detected by

scat detection dogs. For each animal we sampled some combination of tissue (ear-tag punch),

hair, and scat (see Eriksson et al. 2020 for additional details). All animal handling was performed

in accordance with protocols approved by the USDA Forest Service, Starkey Experimental

Forest Institutional Animal Care and Use Committee (IACUC No. 92-F-0004; protocol #STKY-

16-01) and followed the guidelines of the American Society of Mammalogists for the use of wild

mammals in research (Sikes 2016).

Modelling Approach

Our approach was to fit a suite of spatially-explicit models across a spectrum of data

resolutions (Table 1) for each species to evaluate the value of each data source and to test

whether incorporating multiple data sets improved precision. Each model considered here is a

variation of a conventional SCR model (Royle and Young 2008) in which the detection

probability $p_{i,j}$ (or rate $\lambda_{i,j}$) decays as a function of the Euclidean distance between the detector j

and the activity center s for individual i, $d_{i,j}$, to yield

$p_{i,j} = p0 \times exp(-d_{i,j}^2/2\sigma^2)\,,$ (1)

where $p0$ is the baseline detection probability when the detector is located exactly at the activity

center of the animal, and σ is a spatial scale parameter related to home range size that determines

the rate at which detection probability declines with distance from the activity center. The

location of each activity center (s_i) is latent but estimated from the locations in which detections

and non-detections occurred.

The observed encounter frequencies $y_{i,j}$ can be modeled as either a binomial or Poisson

random variable, depending on whether the detection process produces binary or count

outcomes,

$y_{i,j} \sim \text{Binomial}(p_{i,j} \times z_i, K)$ or $y_{i,j} \sim \text{Poisson}(\lambda_{i,j} \times z_i \times K),$ (2)

where K is the number of sampling occasions, and z_i is the data augmentation parameter (Royle

et al. 2007) that takes the values of 0 or 1 for $i = 1, 2…M$, where M is the number of known

individuals plus some number of all zero encounter frequencies representing individuals in the

population not detected during sampling. The value of M must be chosen to be larger than the

maximum possible population size and is sufficiently large when it does not affect the estimation

of N. Data augmentation is used to estimate the number of animals in the state space, S (an area

prescribed to be sufficiently large such that animals with activity centers at the outermost regions

of this area have a negligible probability of being detected). z_i determines whether each

hypothetical individual in the augmented population is real or not and is governed by a Bernoulli

distribution,

$z_i \sim \text{Bernoulli}(\psi)$ for $i = 1, 2 \ldots M$ (3)

in which ψ is the expected number of activity centers divided by M. The estimated population

size N (i.e. the number of activity centers in the state space) is then determined by summing over

all z_i,

$N = \sum_{i=1}^{M} z_i,$ (4)

and density D is obtained by dividing the population size by the area of the state space, A,

$D = N/A.$ (5)

To ensure that the specification of the state space and number of augmented individuals is

sufficiently large, M must be substantially higher than the estimated N, the upper bound of ψ

must not be near 1, and the buffer around the survey area forming the state space must be > 2.5

times σ (Royle et al. 2013).

We assumed a homogenous point process for the distribution of activity centers (s_i) such

that the x and y dimensions of each activity center were uniformly distributed within the defined

limits of the state space:

$s_i \sim \text{Uniform}(S)$ (6)

Each model subsequently described is based on the same framework as the conventional SCR

model described above but incorporates variations due to different data sources or resolutions

which are described next.

Spatial Count or 'unmarked SCR'

The spatial count model (Chandler and Royle 2013) does not require individual recognition, so

the encounter frequency $y_{i,j}$ is replaced by a vector of counts of unmarked individuals at each

detector (n_j) and encounter rate λ replaces p. The counts are now modeled as a Poisson random

variable,

$n_j \sim \text{Poisson}(\Lambda_j \times K)$ (7)

where

$\Lambda_j = \lambda 0 \sum_{i=1}^{M} exp(-d_{i,j}^2/2\sigma^2) \times z_i.$ (8)

Spatial Presence Absence

The Poisson distribution assumes that the mean is equal to the variance, but when animal

detections are aggregated in time and/or space the variance may greatly exceed the mean. This

overdispersion could bias SC results due to the violation of the model assumption. The spatial

presence-absence model (SPA; Ramsey et al. 2015) is similar to the spatial count model, but

only detection-non detection data are utilized so a binomial distribution replaces the Poisson. By

reducing counts to binary detections and using a binomial distribution, the SPA model may be

more appropriate in situations with overdispersion.

$n_j \sim \text{Binomial}(P_j, K)$ (9)

where $n_j \in \{0,1\}$ and

$P_j = 1 - \prod_{i=1}^{M}(1 - p_{i,j} \times z_i)$ (10)

Above, P_j is the probability that at least one individual is detected in detector J, and $p_{i,j}$ is the

detection probability for individual i at detector J which like the other models decreases as a

function of the distance between the activity center and the trap according to a half-normal

function (as in equation 1).

Spatial Mark-Resight

Spatial mark-resight models (Chandler and Royle 2013, Sollmann et al. 2013a) combine

SCR and SC. Here, a subset of the population is individually recognizable due to either natural or

artificial markings but the remaining individuals are observed without individual identity. The

Poisson-distributed encounter history:

$y.marked_{i,j} \sim \text{Poisson}(\lambda.resight_{i,j} \times z_i \times K)$ (11)

is obtained for marked individuals and the resulting expected resighting rate:

$\lambda.resight_{i,j} = \lambda0.resight \times exp(-d_{i,j}^2/2\sigma^2)$ (12)

is applied to the counts of unmarked individuals n_j by replacing $\lambda0$ in Eq. 8 with $\lambda0.resight$.

Assumptions of this model are that marked and unmarked individuals have the same rate of

detection and are always accurately identified as marked or not.

Generalized Spatial Mark-Resight

Generalized spatial mark-resight (Whittington et al. 2018) is identical to the conventional

SMR described above but incorporates a submodel for the marking (i.e. physical capture and

tagging) process and is therefore suitable only when a portion of the population is captured and

marked. Generalized SMR alleviates bias introduced due to heterogeneity of individual

encounter rate arising from situations when the marked portion of the population resides nearer

to the resighting detectors (and are therefore detected more frequently) than unmarked

individuals at the periphery of the state space (Whittington et al. 2018). That is, it accounts for

the fact that the marked segment of the population is seldom distributed randomly throughout the

state space (Whittington et al. 2018). The marking process submodel is itself a SCR model for

the encounter histories of individuals captured and marked:

$y.marking_{i,j} \sim \text{Binomial}(p.marking_{i,j} \times z_i, K)$ (13)

where the probability of capturing individual i in trap j similarly decays with distance according

to

$p.marking_{i,j} = p0.marking \times exp(-d_{i,j}^2/2\sigma^2)$ (14)

The animal capture and marking process and the resighting of marked animals on cameras shares

information on z_i, s_i, and possibly σ, which has the potential to increase model performance and

precision. Generalized spatial mark-resight should be used whenever possible because

conventional SMR models have been shown to be biased (Whittington et al. 2018).

Hybrid Generalized Spatial Mark-Resight + Spatial Capture-Recapture

When genetic SCR data are obtained concurrently with a capture and marking effort, and

the marked individuals are resighted on an array of detectors (e.g. trail cameras), we propose data

from all three processes can be modeled together in a 'hybrid' gSMR+SCR model (Fig. 3). This

model incorporates equations 1-6, and 11-14. This data-intensive model thus utilizes multiple

encounter methods (physical capture, genetics, and cameras) and requires that the identities of

captured and marked individuals can always be reconciled when encountered in the genetic

dataset, so captured animals must always be genotyped. Further, the encounter histories from

each of the three data sources must be aligned such that each individual known from either the

marking/resighting process or scat detection process corresponds to the same individual, i in the

combined model. However, not all known individuals are likely to be detected in all data

sources, so all zero detection histories must be given to known individuals when not encountered

in a certain encounter method. For marked animals which are known to exist because they were

physically captured, the possible outcomes of the detection process are 1) detected by neither

camera nor genetics, 2) detected by camera only, 3) detected by genetics only, or 4) detected by

both camera and genetics. For unmarked animals (i.e. those not physically captured), the possible

outcomes of the detection process are 1) detected by neither camera nor genetics, 2) detected by

genetics only (and without a genotype matching any of the marked individuals), 3) detected by

camera only without individual identity, and 4) detected by camera and genetics (and without the

possibility of deterministically linking the genotype to the photographed individual). Information

on z_i, s_i, and σ can be shared, and baseline detection rates can be informed by the different data

sources when an individual is detected by one but not all detection methods. For example,

knowing a captured animal exists but was not detected by genetics can help inform the

probability of detection in the genetic SCR component. Similarly, knowing a captured animal

exists but was not photographed informs the detection probability for the SMR model

component.

The hybrid gSMR+SCR model, below with the same distributional assumptions and

priors as in our empirical study, has the following joint posterior distribution:

$$[z_i, \psi, \sigma, s_i, p0, \lambda 0.\,marking, \lambda 0.\,resight \,|$$

$$y.\,genetics_{i,j,k},\, y.\,mark_{i,j,k},\, y.\,resight_{i,j,k},\, n_{j,k},\, x.\,genetics_j,\, x.\,marking_j,\, x.\,camera_j,\, telemetry_{i,l} \,]$$

$$\propto Genetic\ SCR: \prod_{i=1}^{M}\prod_{j=1}^{J}\prod_{k=1}^{K} \text{Bernoulli}\left(y.genetics_{i,j,k}\mid f(p0, z_i, \psi, \sigma, s_i, x.genetics_j)\right)$$

$$\times Marking\ Process\ SCR: \prod_{i=1}^{M}\prod_{j=1}^{J}\prod_{k=1}^{K} \text{Poisson}\left(y.marking_{i,j,k}\mid f(\lambda0.marking, z_i, \psi, \sigma, s_i, x.marking_j)\right)$$

$$\times Camera\ SMR: \prod_{i=1}^{n.marked}\prod_{j=1}^{J}\prod_{k=1}^{K} \text{Poisson}\left(y.resight_{i,j,k}\mid f(\lambda0.resight, z_i, \psi, \sigma, s_i, x.camera_j)\right)$$

$$\times \prod_{i=n.marked+1}^{M}\prod_{j=1}^{J}\prod_{k=1}^{K} \text{Poisson}\left(n_{j,k}\mid f(\lambda0.resight, z_i, \psi, \sigma, s_i, x.camera_j)\right)$$

$$\times Telemetry\ Data: \prod_{i=1}^{n.marked}\prod_{l=1}^{n.telem.locs} \text{Multivariate Normal}\left(Telemetry_{i,l}\mid s_i, \sigma^2\mathbf{I}\right)$$

$$\times Parameters: \text{Bernoulli}(z_i\mid\psi) \times \text{Beta}(\psi\mid1,1) \times \text{Uniform}(\sigma\mid0, 20{,}000) \times \text{Beta}(p0\mid1,1)$$

$$\times \text{Uniform}(\lambda0.marking\mid0,5) \times \text{Uniform}(\lambda0.resight\mid0,5) \times \text{Uniform}(s_i\mid S_{min}, S_{max})$$

where *x* corresponds to the geographic coordinates of the detectors of a given data source. A

directed acyclic graph (DAG) of the hybrid gSMR+SCR model is presented in Fig. 3.

Incorporating telemetry data

If some animals within the population are fit with GPS collars (or other telemetry

methods) during the time sampling is conducted, the telemetry positions can be incorporated into

any of the models previously described (Royle et al. 2013, Sollmann et al. 2013a). This is

achieved by modelling the *l* GPS positions as a two-dimensional multivariate normal distribution

(i.e. bivariate normal) centered on the activity center coordinates of marked individual *i* with no

covariance (Garton et al. 2001):

$Telemetry_{i,l} \sim \text{Multivariate Normal}(s_i, \sigma^2\mathbf{I})$ (14)

Thus, the incorporation of telemetry data informs the location of the activity center s for

telemetered individuals, the scale parameter in the detection function σ, and baseline detection

rate or probability. However, in the spatial count models, individual recognition is not known,

which prevents the GPS data from being deterministically linked to specific individuals, so in

those models the telemetry data typically only inform σ.

Application

For each of the four study species (black bear, bobcat, cougar, coyote), we first fit "base"

models that did not consider differences in sex or behavioral status in detection parameters. SC

models were fit using only camera data. Individual identity was ignored in SC models even when

it was known so that we could evaluate an approach that required no individual recognition.

Although this was an inefficient use of the information available, our objective was to determine

whether investing in the marking and resighting of individuals was necessary. SMR and gSMR

models were constructed from the same camera data as the SC models but GPS-collared animals

were identified to individual, with the latter method also including the encounter histories from

the marking process. SCR models were constructed only from genetic samples (scats) because

we could not identify individuals from cameras unless they were artificially marked. The gSMR

+ SCR hybrid model that we developed combined data from cameras, physical captures to mark

animals, and genetic samples. We assumed that camera and scat data were independent because

1) both data collection methods passively sampled the population in different ways and at

different sites within the superpopulation, 2) the detection of a scat did not influence the

probability of that individual being detected by camera (and vice-versa), and 3) neither camera

placements nor scat dog survey routes were influenced by the knowledge gained from the other

method.

For every modeling method implemented (Table 1), we fit models both with and without

GPS data. When telemetry data were included, we randomly selected 100 GPS locations (Fig.

2d) from each individual to alleviate lack of independence arising from temporal autocorrelation.

Because GPS data were collected contemporaneously with camera and genetic sampling, they

could be deterministically linked to the detection data of that individual which helped us estimate

the activity center for that individual, as well as σ and $p0$ or $\lambda0$. The easting and northing

locations were modeled as a bivariate normal distribution with no covariance, with a mean of

that individual's activity center (s_i) and standard deviation equal to the detection scale parameter

σ.

For the genetic SCR analyses, we used a single occasion because detection dogs only

surveyed each grid cell once. We overlaid a 1 x 1 km grid over the survey area and attributed the

location of any scat inside that cell to the grid cell center. To control for variation in survey

effort, we included the distance walked by scat dog handlers within each grid cell as a covariate

influencing $p0$ using a linear model with a logit link. Samples belonging to the same individual

that were found in the same grid cell were recorded as a single detection of individual i. Because

some carnivore species defecate nonrandomly as a means of scent marking, we believe removing

recaptures within the same grid cell helped ensure independence of samples.

For camera analyses, we parsed the encounter data into ten 14-day occasions beginning

15 April. We used a trap operation matrix in our detection function to account for varying

camera operability so that the probability of detection was zero if a camera was not functional.

Similarly, if an animal marked at some point during the sampling period was not available to be

detected during one or more occasions due to death or because it was not yet collared, we did not

allow that individual to contribute to the likelihood for that occasion.

To assess and compare dispersion across model types, we calculated the coefficient of

variation as the posterior standard deviation divided by the posterior mean. We refer to precision

as the inverse of the coefficient of variation rather than the inverse of the variance.

Model implementation

For models lacking individual identity of all individuals (i.e. all except SCR) we used the

marginal model implementation described by Chandler and Royle (2013). We used Markov

Chain Monte Carlo (MCMC) to draw samples from the posterior distributions for the parameters

of interest. We used JAGS (Plummer 2003) in program R version 3.6.1 (R Development Core

Team 2019) and the jagsUI package (Kellner 2015) and NIMBLE (de Valpine et al. 2017,

NIMBLE Development Team 2019). For each model we ran 3 chains consisting of 50,000

iterations per chain and discarded the first 15,000 as burn in. We assessed model convergence by

visually inspecting traceplots and ensuring the $\hat{R}$ values were less than 1.1 (Gelman 1996). If

models failed to converge after the first run, we updated them until convergence was satisfactory.

For all parameters, we calculated the 95% highest posterior density intervals (HPDI) and 95%

Bayesian credible intervals (BCI) (Chen and Shao 1999). For estimates of population density,

which may be sensitive to the amount of data augmentation prescribed, we report the posterior

mode in the text but also present the posterior means and medians in tables.

RESULTS

Our capture efforts yielded GPS collars placed on 6 black bears, 4 bobcats, 6 cougars,

and 9 coyotes that could be resighted. During the time of camera sampling, 1 cougar died but the

remainder of marked individuals remained alive and in the sampling area. However, 3 coyotes

were transient, meaning they roamed widely across an area much larger than the study area. We

obtained 159 independent photo sequences for bears (28 marked, 131 unmarked), 34 photo

sequences of bobcats (9 marked and 25 unmarked), 48 photo sequences of cougars (24 marked,

24 unmarked), and 479 photo sequences of coyotes (55 marked, 424 unmarked; Table 1). Scat

detection dogs located 86 bear scats, 95 bobcat scats, 17 cougar scats, and 772 coyote scats. Of

these, we successfully genotyped 43 bear scats out of 72 attempts (60% success rate), 86 bobcat

scats out of 95 attempts (91% success rate), 15 cougar scats out of 17 attempts (88% success

rate), and 201 coyote scats out of 216 attempts (93% success rate). After removing duplicate

scats from the same individual in the same grid cell, we had 40 bear scats from 31 individuals, 68

bobcat scats from 32 individuals, 13 cougar scats from 7 individuals, and 165 coyote scats from

83 individuals (Table 1). Maps of spatially-explicit detection data for camera and genetic

analyses are provided as supplementary information (Appendix S3: Figures S1–S4).

Telemetry Data

Across all species and methods, the inclusion of GPS data resulted in a mean decrease in

the posterior modes for animal densities of 25.4% (Fig. 4, Appendix S4: Tables S1–S8). For all

methods except spatial count models, including GPS data led to a mean decrease in the

coefficients of variation of 18.6% (Fig. 5, Appendix S4: Tables S1–S8). But in the unmarked

models, including GPS data increased the coefficients of variation by an average of 154.5% (Fig.

5, Appendix S4: Tables S1–S8). However, very low SC+GPS density estimates were responsible

for the increases in the coefficients of variation in those models—standard deviations were

actually smaller in SC models including GPS data than SC models lacking GPS data. Because

models containing GPS data were more precise, likely because GPS data provided valuable

information aiding in the identifiability of detection parameters, we consider them more reliable

than models that do not include telemetry data.

Spatial Capture-Recapture

We report SCR+GPS estimates as the benchmark to which the other methods are

compared because they have the strictest requirements of individual identifiability and thus

should yield the most robust estimates. The base SCR+GPS estimate of the posterior mode (and

95% BCI) for black bears was 9.9/100 km^2 (6.1–14.9 /100 km^2); for bobcats was 12.1/100 km^2

(8.4–16.2/100 km^2), for cougars was 1.7/100 km^2 (0.9–3.6/km^2); and for coyotes was 25.0/100

km^2 (21.0–30.9/100 km^2) (Fig. 4, Appendix S4: Tables S1–S8).

Spatial Count

Spatial count models produced highly different results depending on whether telemetry

data were included (Fig. 4; Appendix S4: Tables S1–S8). When GPS data were not included,

densities from SC models were on average 773.9% higher than SCR+GPS estimates (Fig. 4, Fig.

6, Appendix S4: Tables S1–S8). However, when GPS data were included, densities from SC

models were on average 84.4% lower than SCR+GPS estimates (Fig. 4, Fig. 6, Appendix S4:

Tables S1–S8). Coefficients of variation for SC and SC+GPS models were 89.9% or 290.8%

greater than base SCR+GPS, respectively (Fig. 5, Appendix S4: Tables S1–S8).

Spatial Presence Absence

Spatial presence-absence models were even more inconsistent with base SCR+GPS than were SC models. Animal densities were on average 1,267% higher or 83% lower than SCR+GPS depending on whether telemetry data were included or not (Fig. 4, Fig. 6, Appendix S4: Tables S1–S8). Coefficients of variation were 125.5% higher than the SCR+GPS model when telemetry was not included, and 470.1% higher with telemetry included (Fig. 5, Appendix S4: Tables S1–S8).

Base Spatial Mark-Resight vs Base Spatial Capture-Recapture

Averaged over all species, base SMR+GPS models estimated animal densities that were only 0.67% lower than base SCR+GPS models (Fig. 4, Fig. 6, Appendix S4: Tables S1–S8). However, coefficients of variation were 51.8% higher in base SMR+GPS models than base SCR+GPS models (Fig. 5, Appendix S4: Tables S1–S8).

Base Spatial Mark-Resight vs Base Generalized Spatial Mark-Resight

Across all models and species, adding the marking process to base SMR models (i.e. SMR vs gSMR) decreased estimates of animal density by an average of only 2.2% (Fig. 4, Fig. 6, Appendix S4: Tables S1–S8). However, the coefficients of variation were on average 9.8% lower in base gSMR models than base SMR models (Fig. 5, Appendix S4: Tables S1–S8). Averaged across all species, base gSMR+GPS models were 3.6% closer to the base SCR+GPS estimates than were base SMR+GPS estimates.

Base Spatial Capture-Recapture vs Base Hybrid

The base Hybrid+GPS models estimated, on average, densities that were only 2.1% lower

than base SCR+GPS models (Fig. 4, Fig. 6, Appendix S4: Tables S1–S8). But the coefficients of

variation were on average 16.0% lower in the base Hybrid+GPS models than the base SCR+GPS

models (Fig. 5, Appendix S4: Tables S1–S8).

Post-hoc Analyses

The notable differences in estimates across certain base model types compelled us to

conduct post-hoc analyses to attempt to explain unmodeled sources of individual heterogeneity

and more strictly adhere to important model assumptions. Thus, for every species we fit an

additional series of models with 1) sex-specific detection and spatial scale parameters (for all

except unmarked models), 2) a reduced sampling period for all methods using camera data (70

days vs 140 days), and 3) for coyotes only, models with behavioral status-specific (i.e. resident

vs transient, Appendix S5) detection and spatial scale parameters (transient coyotes in our study

used home ranges that were 10–20 times the area of resident coyotes home ranges). For each of

the new model types, we calculated the percent differences in posterior modes of animal

densities between SCR+GPS models and SMR+GPS, gSMR+GPS, or hybrid+GPS models to

assess whether the consistency of results improved. We also assessed how the coefficients of

variation changed between the base and post-hoc models.

Sex-specific Models

Including sex-specific detection and spatial scale parameters increased the SCR+GPS

estimates of animal density by 4.4% averaged across species. The change in agreement between

sex-specific gSMR+GPS and sex-specific SCR+GPS posterior modes of animal density relative

to the same comparisons for the base models varied by species. By including sex, the difference

between gSMR+GPS and SCR+GPS estimates was reduced by 5.5 percentage points (from

11.7% to 6.2%) and 10.8 percentage points (from 44.4% to 33.6%) for black bears and bobcats,

respectively, and was only increased by 2.3 percentage points (from 6.9% to 9.1%) and 0.1

percentage points (25.8% to 25.9%) for cougars and coyotes (Fig. 7–8, Appendix S6: Tables S1–

S8). The sex-specific SCR+GPS models had coefficients of variation that were 9.4% higher than

the base SCR+GPS models when averaged across species. For gSMR+GPS, coefficients of

variation were 15.3% higher in the sex-specific models than the base gSMR+GPS estimates (Fig.

9).

Reduced Sampling Period Models

When the duration of camera sampling was reduced in half for gSMR+GPS models, the

estimates of animal densities were closer to the base SCR+GPS estimates for coyotes only. For

that species, the percent difference in the posterior mode for animal density was reduced by 25.7

percentage points (from 25.8% to 0.1%; Fig. 7, Appendix S7: Tables S7–S8) when the reduced

sampling period gSMR+GPS estimates were compared to the base SCR+GPS estimates.

However, the percent difference in animal density was increased by 6.7 percentage points (from

11.7% to 18.5%), 4.9 percentage points (from 44.4% to 49.3%), and 8.1 percentage points (from

6.9% to 15.0%) for bears, bobcats, and cougars, respectively (Fig. 7, Appendix S7: Tables S1–

S6). Averaged across species, the coefficients of variation for gSMR+GPS models were 13.7%

higher in the reduced sampling period models than the base gSMR+GPS models (Fig. 9).

Behavioral Status Models for Coyotes

Allowing detection and spatial scale parameters to vary by the behavioral status (i.e.

resident vs transient as determined from GPS data; Appendix S5) of coyotes increased the

estimates of density for models based on both camera and genetic data (Appendix S8: Tables

S1–S2). However, incorporating behavioral status into models reduced the percent difference of

coyote density between gSMR+GPS and SCR+GPS by 22.6 percentage points (from 25.8% to

3.16%; Fig. 7–8, Appendix S8: Tables S1–S2) relative to the same comparison for base models.

DISCUSSION

Many applications in ecology depend on accurate estimates of animal density. Despite

tremendous advancements in wildlife monitoring technology (e.g. remote cameras), molecular

methods (e.g. genotyping), and statistics (e.g. models accounting for imperfect detection),

estimating population density can still be a formidable task. Using base models (i.e. those not

considering variation in detection parameters induced by sex or behavioral status), we found that

gSMR+GPS models based on resighting marked animals on cameras produced density estimates

consistent with genetic SCR+GPS models for bears (posterior mode was 11.7% higher for

gSMR+GPS than SCR+GPS) and cougars (gSMR+GPS was 6.9% lower than SCR+GPS).

However, there were greater discrepancies in base models for bobcats for which the posterior

mode of gSMR+GPS estimates were 44.4% lower than the SCR+GPS results and for coyotes for

which gSMR+GPS estimates were 25.8% higher than the SCR+GPS results. For both species,

the posterior mode of the SCR models was contained in both the 95% HPDI and BCI of SMR

models, but the converse was not true (Fig. 4, Appendix S4: Tables S3 and S7).

Although base gSMR models and base SCR models produced qualitatively similar

density estimates, the moderate differences in results for bobcats and coyotes may be due to

challenges of simultaneously camera trapping for multiple species with distinct habitat

preferences, home range sizes, and life history strategies. We attempted to refine estimates by

fitting additional post-hoc models to explain more variation in heterogeneity of detection rate

using individual covariates on detection parameters for sex, and, for coyotes only, behavioral

status to consider resident and transient states. After controlling for these important sources of

variation, we found much greater concordance in estimates between camera-based gSMR and

genetic SCR (Fig. 8).

We also fit models using half the sampling occasions for camera analyses to assess

whether a lack of demographic closure could have driven the discrepancies between analyses

based on camera versus genetic data. While the available data indicated that demographic closure

was not grossly violated in this study (one marked cougar died during the time of camera

sampling), it is nonetheless possible that other deaths or dispersal occurred which would cause

our estimates to be biased high. If, however, recruitment occurred and juveniles were not

detected then our estimates could be biased low. But because the species considered here are

generally long-lived and our sampling was designed to miss the birth pulse, we believe the

population was reasonably closed during our study.

Base SCR+GPS estimates were 44% higher than base gSMR+GPS estimates for bobcats,

but this was reduced to 33% by including sex-specific detection parameters. This modest

improvement in consistency was likely driven by sex-specific differences in home range size,

which induced heterogeneity in detection rate that was previously unaccounted for. However, a

moderate discrepancy in estimates remained. Based on GPS data, bobcats in this system used

dense forest cover and our camera placements were biased away from this habitat type because

they had extremely limited viewsheds that would limit photos of other species. In contrast,

detection dogs readily located bobcat scats and the higher detection probability of scats may have

driven the difference in results between cameras and genetics. Alternatively, the assumption of

demographic closure may have been violated since bobcats can give birth nearly year-round

(Winegarner and Winegarner 1982). It is possible that juvenile bobcats were detected via genetic

sampling but that they were missed on cameras. Genetic techniques are currently unable to

accurately determine the age of the individual from its scat, so unless scats of juveniles can be

censored due to size or other morphological characteristics, recruitment has the potential to be

problematic for all genetic mark-recapture studies.

For coyotes, the inconsistency between base gSMR and base SCR models likely involved

unmodeled sources of individual heterogeneity in detection rates across methods as was observed

by Murphy et al. (2018a) who estimated lower densities when models were constructed solely

from scat data than from both scat and hair. The authors speculated that the behavioral status of

coyotes (i.e. resident vs transient) led to cohorts that were disproportionately detected using one

detection method over another. Specifically, they reasoned that territorial resident coyotes could

be more likely to be detected from scats because of smaller home ranges and increased scent-

marking with scat for territorial defense than wide-ranging transients. Our study may exhibit an

analogous issue if cameras detected both resident and transient coyotes at similar rates but the

detected scats disproportionately belonged to residents. Similarly, Morin et al. (2016) suggested

that SCR density estimation in coyotes could be improved by modeling heterogeneity in σ and

baseline detection rates arising as a result of behavioral status. Without modelling such

heterogeneity, the bimodality in home range size can lead to estimates of detection and scale

parameters that are averaged between the two behavioral states (Murphy et al. 2018). Transient

coyotes in our study used home ranges that were 10–20 times greater than those of resident

coyotes.

To explore whether failing to acknowledge these sources of heterogeneity were

responsible for the inconsistencies between base gSMR and base SCR models, we conducted a

post-hoc analysis where σ and baseline detection parameters were allowed to vary as a function

of transient or resident behavioral status (see Appendix S5 for details). Including behavioral

status increased the estimates of density from base models in both gSMR+GPS and SCR+GPS

models but reduced the difference between them substantially from 26% to 3% (Fig. 7, Appendix

S8: Tables S1–S2). We thus speculate that the unmodeled heterogeneity arising from the distinct

behavioral states explained much of the disagreement between model types. We recommend

others control for behavioral states in coyotes if GPS data are available to classify individuals

into residents or transients.

Spatial count and spatial presence-absence models reliant on photos of unmarked animals

produced density estimates that were very inconsistent with methods in which individuals were

identifiable. For all four species, we observed the same pattern in which estimates from SC and

SPA models without GPS data were higher than SCR estimates and the same models with GPS

data were lower than SCR estimates. Informing σ with telemetry data did not lead to accurate

enumeration of activity centers and instead falsely attributed unmarked photo detections to too

few individuals. Overdispersion in the count data was probably not responsible for imprecise

results in the SC models because after collapsing counts to independent binary detections and

using the SPA model, the estimates became even more different from SCR results.

SC and SPA models require sufficiently close trap spacing so that spatial autocorrelation

can be detected and used to identify activity centers (Chandler and Royle 2013, Ramsey et al.

2015). Poor density estimation from SC models can result from insufficient spatial structure of

animal detections that prevent individual activity centers from being clearly demarcated.

However, our 1 km camera grid spacing allowed for many cameras per animal home range,

suggesting that these models are not robust for carnivores under sampling scenarios that are

feasible to accomplish in most studies. Further, SC and SPA results can be unreliable, unstable,

or suffer from convergence failure when data are sparse (Chandler and Royle 2013, Ramsey et

al. 2015, Kane et al. 2015, Burgar et al. 2018a, Burgar et al. 2018b, Murphy et al. 2018b),

suggesting that this will be a consistent challenge in carnivore studies using unbaited cameras

within a time frame over which demographic closure can be reasonably assumed. Populations

with large values of σ and/or high densities can produce biased results in SC models (Chandler

and Royle 2013, Augustine et al. 2019) limiting their potential for use in many species. Including

habitat covariates that influence spatial density in SC models may lead to results that are

consistent with SCR (e.g. Evans and Rittenhouse 2018), but our results suggest that fully

unmarked SCR models (i.e. SC and SPA) may unsuitable for density estimation of the carnivores

we studied, particularly when detectors are widely spaced, count data are sparse, and auxiliary

data are not available.

Previous studies have successfully estimated population densities of multiple carnivore

species using camera trapping (Jiménez et al. 2017, Burgar et al. 2018a, Rich et al. 2019). Our

study also suggests a multi-species monitoring approach is possible yet highlights the challenges

of simultaneous density estimation of multiple species. For example, the substantial variation in

home range size among carnivores requires fine-scale sampling for smaller ranges (e.g. coyotes

and bobcats) while allowing for the larger extents needed for greater home range sizes (e.g. bears

and cougars). This can be accomplished with either a finely-spaced grid (this study) or through

cameras that are clustered with variable spacing intervals (Rich et al. 2019, Murphy et al. 2019).

Similarly, assumptions of demographic closure may be challenging to meet for species that roam

widely unless the extent of the study area is large enough to contain many home ranges. For

example, our small survey area relative to the species with the largest home range and lowest

density (e.g. only 7 distinct cougars were sampled from genetics) may have made our estimates

sensitive to the assumption of demographic closure given that the death, dispersal, or recruitment

of even one individual could have a substantial influence on estimates.

For all species, the inclusion of telemetry data to inform density estimation generally

produced consistent results with increased precision and smaller coefficients of variation for all

methods except unmarked models (Fig. 5, Appendix S4: Tables S1–S8). The gains in precision

were greatest for the models/species with the sparsest data (e.g. cougar SCR model). This

highlights the challenge of estimating σ when telemetry data are excluded and spatial recaptures

are sparse. For example, transient coyotes with large home ranges may rarely or never be

recaptured, which would falsely suggest low detectability rather than the large value for σ that is

evident from telemetry data. The absence of telemetry data can thus lead to both imprecise and

biased estimates of σ with consequent effects on other model parameters (i.e. $\lambda 0$ or $p0$, s_i, and

ultimately, N) (Whittington et al. 2018). Improving the estimate of σ by incorporating telemetry

data should aid in identifiability of other parameters and lead to greater accuracy in the models

incorporating GPS data. SC and SPA models were particularly problematic because they

produced biologically unrealistic density estimates that changed from very high to very low with

the inclusion of telemetry data. Although telemetry data decreased precision in unmarked

models, the large reduction in the density estimate when including telemetry data led to an

increase in the coefficient of variation because the denominator became smaller. An inaccurate

result with a low CV invites the possibility of overconfidence in an incorrect result when using

unmarked models.

Models that included more information on individual identity exhibited more precise

density estimates (Fig. 5), except when the number of genotyped scats was very low. In

particular, the small sample of cougar scats led to SCR models that were substantially less

precise than SMR and gSMR models (Fig. 5, Appendix S4: Table S5). For all species, the hybrid

model containing camera, physical capture (gSMR), and genetic (SCR) data had the lowest

coefficients of variation (Fig. 5) highlighting the benefit of leveraging multiple datasets even if

some are sparse.

A recent movement in ecological modeling of demographic rates has sought to strengthen

inference by utilizing multiple independent data sources (e.g. Besbeas et al. 2002, Schaub et al.

2007). SMR and SCR models require a substantial investment of time and money yet may still

only yield sparse data. Integrating multiple sparse data sources is thus an efficient use of

resources in addition to the benefits of improving estimation. Others have integrated camera and

genetic data for spatially-explicit density estimation (Sollmann et al. 2013b, Clare et al. 2017,

Gopalaswamy et al. 2012), but to our knowledge this has only occurred when animals on camera

could always be identified to individual, so the result was a SCR model composed of two

different data sources. And recently, both identified and unidentified genetic samples were

combined in a SCR model using multiple observation processes (Tourani et al. 2020). These

studies, however, contrast with our approach of combining SMR for a partially-marked

population (i.e. those in which not all individuals can be uniquely identified) with genetic SCR

data in which every sample was identifiable to individual, and sharing information via

deterministic linkages between individuals that were genotyped, GPS collared, and resighted on

cameras.

Ancillary data associated with genetic or mark-resight data may motivate the decision to

collect one data type over another. For example, telemetry data obtained in conjunction with a

mark-resight study may be useful not only to inform σ, $p0/\lambda0$, and the activity centers of the

telemetered individuals, but also to classify an animal into a behavioral state such as resident vs

transient or breeder vs nonbreeder, and these distinctions may explain important sources of

individual heterogeneity in detection rate. The photographs in a mark-resight analysis may also

permit identification of juveniles which can easily be censored if the researcher desires to

estimate the density of the adult population segment only. While genetic sampling is unlikely to

yield these sources of information, it offers other benefits such as the ability to study diet (e.g.

Kartzinel et al. 2015), hormone levels (e.g. Gobush et al. 2014), and population structure or

genetic diversity (e.g. Goosens et al. 2005), which if useful for other objectives may steer

researchers toward that method of data collection. Genetic samples also readily yield the sex of

the individual which may not be possible from camera trapping for species that are not sexually

dimorphic. The cost of each data collection method may also play a factor in the decision. While

both genetics and mark-resight studies may be expensive, when a segment of the population is

already being monitored via telemetry devices, the extra cost to resight marked individuals using

cameras may be minimal compared to the cost of a separate genetics SCR study (Whittington et

al. 2018). However, in the absence of an existing effort to obtain telemetry data, genetic SCR

methods are substantially more cost effective given the expense associated with capturing,

collaring, and resighting animals.

Another extension to our efforts might implement spatial partial-identity models to

extract even more information from the existing datasets (Augustine et al. 2018). This would

allow partial genotypes (Augustine et al. 2019) or photos in which the marked status or

individual identity (Augustine et al. 2018, Jiménez et al. 2019, Tourani et al. 2020) cannot be

ascertained to be included in SCR estimation. However, this model extension would minimally

influence our results because of the high genotyping success from scats using genotyping by

amplicon sequencing (Eriksson et al. 2020) and our high success in identifying marked status or

individual identity in our camera trap photos. Further, if the discrepancies between model types

were driven by violations of model assumptions (e.g. demographic closure), partial identity

models would likely not overcome the inherent bias and may instead only increase precision.

Knowing how much and what type of data to collect is a challenge confronted by most

researchers aiming to efficiently estimate population densities. Our results from a suite of

spatially-explicit models for bears, bobcats, cougars and coyotes in a natural system highlight the

relative performances of models within the spatial capture-recapture family constructed from

data of varying sources and resolutions. We acknowledge that in some instances, such as setting

management objectives for common species, absolute population densities may not be essential

and simply being able to accurately characterize relative changes in population trends may be

adequate. In other cases, such as the calculation of extinction risk for imperiled species,

obtaining the most accurate estimates of population size is crucial. We found that the most

common method for estimating population size of terrestrial carnivores in the past five years was

(spatial) capture-recapture and that relatively few studies used methods that do not require

individual recognition (Appendix S1: Fig. 1). This suggests that in most cases, researchers are

willing to spend the resources necessary to achieve robust results. Regardless, the method chosen

must be commensurate with the objective of the study.

We conclude by suggesting that future research employing SCR or its variants to estimate

the density of carnivores (1) ensures the assumptions and study design requirements of unmarked

methods can strictly be met before using them, (2) controls for important sources of variation in

detection probability such as those arising from sex or behavioral status, (3) incorporates GPS

data when possible, and (4) considers combining multiple data sources, e.g. using the hybrid

gSMR+SCR model presented here. We note that SCR models with few scats produced density

estimates consistent with, but less precise than, a years-long effort to capture carnivores and

resight them on camera traps. Given the inadequacy of the unmarked models and expense

associated with capturing carnivores to mark them for resighting, we recommend that new

studies to estimate population density of carnivores (that are not distinguishable with natural

markings) first consider using genetic SCR. The efficacy of this approach will continue to

increase as noninvasive genetic sampling transitions toward low-cost and high-success

genotyping of SNPs by high-throughput sequencing (Eriksson et al. 2020). Finally, newer

models to estimate density using unmarked animals, such as the time-to-event and space-to-event

family of models (Moeller et al. 2018), provide hope for a cost-effective method to estimate

population densities for multiple species. These models, however, have their own set of

assumptions such as random camera placement with respect to animal movement, accurate

measurement of the camera viewshed area and perfect detection of animals within that area, and

in the case of the time-to-event model, accurate and independent estimates of animal movement

speed. Whether those assumptions are tenable for typical field applications will require a similar

test to that presented here prior to their integration into large-scale carnivore monitoring

programs.

ACKNOWLEDGMENTS

We thank M. Bianco, K. Bowman, C. Brown, L. Carr, L. Chodelski, T. Craddock, E. Crain, C.

Eckrich, M. Esquibel, S. Gillman, M. Goldman, A. Hilger, R. Jensen, C. Kidd, M. Olsen, J.

Smith, A. Yancey, and J. Yancey for field support. We thank K. Cronin, S. Gillman, M.

Goldman, J. Hart, V. Heilman, R. Jensen, J. Neeley, N. Owen, S. Muñoz for photo tagging

support. We thank J. Allen for lab work. We thank R. Chandler and B. Augustine for coding

assistance. We thank R. Chandler and an anonymous reviewer for helpful comments on an

earlier draft. We thank B. Dick, R. Kennedy, B. Naylor, and D. Rea and the USDA Forest

Service Pacific Northwest Research Station for logistical support. Funding and/or field

equipment was provided from Oregon Hunter's Association, Oregon Trapper's Association,

Wildlife Restoration Act, Oregon Department of Fish and Wildlife, USDA Forest Service, and

Oregon State University.

LITERATURE CITED

Augustine, B. C., J. A. Royle, M. J. Kelly, C. B. Satter, R. S. Alonso, E. E. Boydston, and K. R.

Crooks. 2018. Spatial capture–recapture with partial identity: an application to camera traps. The

Annals of Applied Statistics 12:67–95.

Augustine, B. C., J. A. Royle, S. M. Murphy, R. B. Chandler, J. J. Cox, and M. J. Kelly. 2019.

Spatial capture–recapture for categorically marked populations with an application to genetic

capture–recapture. Ecosphere 10:e02627.

Balser, D. S. 1965. Tranquilizer tabs for capturing wild carnivores. The Journal of Wildlife

Management 29:438–442.

Besbeas, P., S. N. Freeman, B. J. T. Morgan, and E. A. Catchpole. 2002. Integrating

mark-recapture-recovery and census data to estimate animal abundance and demographic

parameters. Biometrics 58:540–547.

Borchers, D. L., and M. G. Efford. 2008. Spatially explicit maximum likelihood methods for

capture–recapture studies. Biometrics 64:377–385.

Buckland, S. T., D. R. Anderson, K. P. Burnham, and J. L. Laake. 1993. Distance sampling:

estimating abundance of biological populations. Oxford University Press, Oxford, UK.

Burgar, J. M., A. C. Burton, and J. T. Fisher. 2019. The importance of considering multiple

interacting species for conservation of species at risk. Conservation Biology 33:709–715.

Burgar, J. M., F. E. Stewart, J. P. Volpe, J. T. Fisher, A. C. Burton. 2018b. Estimating density for

species conservation: Comparing camera trap spatial count models to genetic spatial capture-

recapture models. Global Ecology and Conservation 15:e00411.

Chandler, R. B., and J. A. Royle. 2013. Spatially explicit models for inference about density in

unmarked or partially marked populations. The Annals of Applied Statistics 7:936–954.

Chen, M. H. and Q. M. Shao. 1999. Monte Carlo estimation of Bayesian credible and HPD

intervals. Journal of Computational and Graphical Statistics 8:69–92.

Clare, J., S. T. McKinney, J. E. DePue, and C. S. Loftin. 2017. Pairing field methods to improve

inference in wildlife surveys while accommodating detection covariance. Ecological

Applications 27:2031–2047.

de Valpine, P., D. Turek, C. J. Paciorek, C. Anderson☐Bergman, D. T. Lang, and R. Bodik.

2017. Programming with models: writing statistical algorithms for general model structures with

NIMBLE. Journal of Computational and Graphical Statistics 26:403–413.

Efford, M. 2004. Density estimation in live☐trapping studies. Oikos 106: 598–610.

Eriksson, C. E., J. Ruprecht, and T. Levi. 2020. More affordable and effective noninvasive SNP

genotyping using high-throughput amplicon sequencing. Molecular Ecology Resources

https://doi.org/10.1111/1755-0998.13208.

Evans, M. J., and T. A. Rittenhouse. 2018. Evaluating spatially explicit density estimates of

unmarked wildlife detected by remote cameras. Journal of Applied Ecology 55:2565–2574.

Gardner, B., J. A. Royle, and M. T. Wegan. 2009. Hierarchical models for estimating density

from DNA mark–recapture studies. Ecology 90:1106–1115.

Garton, E. O., M. J. Wisdom, F. A. Leban, and B. K. Johnson. 2001. Experimental design for

radiotelemetry studies. Pages 15–42 in J. Millspaugh and J. Marzluff, editors. Radiotelemetry

and animal populations. Academic Press, San Diego.

Gelman, A. 1996. Inference and monitoring convergence. Pages 131–143 in W. R. Gilks, S.

Richardson and D. J. Spiegelhalter, editors. Markov chain Monte Carlo in practice. Chapman and

Hall/CRC, Boca Raton, Florida, USA.

Gobush, K. S., R. K. Booth and S. K. Wasser. 2014. Validation and application of noninvasive

glucocorticoid and thyroid hormone measures in free-ranging Hawaiian monk seals. General and

Comparative Endocrinology 195:174– 182.

Goossens, B., L. Chikhi, M. F. Jalil, M. Ancrenaz, I. Lackman-Ancrenaz, M. Mohamed, P.

Andau, and M. W. Bruford. 2005. Patterns of genetic diversity and migration in increasingly

fragmented and declining Orangutan (*Pongo pygmaeus*) populations from Sabah, Malaysia.

Molecular Ecology 14:441– 456.

Gopalaswamy, A. M., J. A. Royle, M. Delampady, J. D. Nichols, K. U. Karanth, and D. W.

Macdonald. 2012. Density estimation in tiger populations: combining information for strong

inference. Ecology 93:1741–1751.

Jiménez, J., J. C. Nuñez-Arjona, C. Rueda, L. M. González, F. García-Domínguez, J. Muñoz-

Igualada, and J. V. López-Bao. 2017. Estimating carnivore community structures. Scientific

Reports 7:1–10.

Kane, M. D., D. J. Morin, and M. J. Kelly. 2015. Potential for camera-traps and spatial mark-

resight models to improve monitoring of the critically endangered West African lion (*Panthera*

leo). Biodiversity and Conservation 24:3527–3541.

Kartzinel, T. R., P. A. Chen, T. C. Coverdale, D. L. Erickson, W. J. Kress, M. L. Kuzmina, D. L.

Rubenstein, W. Wang, and R. M. Pringle. 2015. DNA metabarcoding illuminates dietary niche

partitioning by African large herbivores. Proceedings of the National Academy of Sciences of

the United States of America 112:8019– 8024.

Kellner, K. F. jagsUI: A wrapper around rjags to streamline JAGS analyses. R package version

1.3.1. 2015.

Kery, M., B. Gardner, T. Stoeckle, D. Weber and J. A. Royle. 2011. Use of spatial

capture-recapture modeling and DNA data to estimate densities of elusive animals. Conservation

Biology 25:356–364.

Lemieux, R., and S. Czetwertynski. 2006. Tube traps and rubber padded snares for capturing

American black bears. Ursus, 17:81–92.

Massey, A. L., G. Roffler, T. Vermeul, J. M. Allen, and T. Levi. 2019. Comparison of

mechanical sorting and DNA metabarcoding for diet analysis with degraded wolf scats. bioRxiv.

https://doi.org/10.1101/2019.12.13.875898.

McClintock, B. T., and G. C. White. 2009. A less field-intensive robust design for estimating

demographic parameters with mark–resight data. Ecology 90:313–320.

Moeller, A. K., P. M. Lukacs, and J. S. Horne. 2018. Three novel methods to estimate abundance

of unmarked animals using remote cameras. Ecosphere 9:e02331.

Morin, D. J., M. J. Kelly, and L. P. Waits. 2016. Monitoring coyote population dynamics with

fecal DNA and spatial capture–recapture. The Journal of Wildlife Management 80:824–836.

Morin, D. J., L. P. Waits, D. C. McNitt, and M. J. Kelly. 2018. Efficient single-survey estimation

of carnivore density using fecal DNA and spatial capture-recapture: a bobcat case

study. Population Ecology 60:197–209.

Murphy, M. A., L. P. Waits, and K. C. Kendall. 2000. Quantitative evaluation of fecal drying

methods for brown bear DNA analysis. Wildlife Society Bulletin 28:951–957.

Murphy, S. M., B. C. Augustine, J. R. Adams, L. P. Waits, and J. J. Cox. 2018a. Integrating

multiple genetic detection methods to estimate population density of social and territorial

carnivores. Ecosphere 9:e02479.

Murphy, A., B. D. Gerber, Z. J. Farris, S. Karpanty, F. Ratelolahy, and M. J. Kelly. 2018b.

Making the most of sparse data to estimate density of a rare and threatened species: a case study

with the fosa, a little-studied Malagasy carnivore. Animal Conservation 21:496–504.

Murphy, S. M., D. T. Wilckens, B. C. Augustine, M. A. Peyton, and G. C. Harper. 2019.

Improving estimation of puma (*Puma concolor*) population density: clustered camera-trapping,

telemetry data, and generalized spatial mark-resight models. Scientific Reports 9:1–13.

Nichols, J.D. 1992. Capture-recapture models. BioScience 42:94–102.

Niedballa, J., R. Sollmann, A. Courtiol, and A. Wilting. 2016. camtrapR: an R package for

efficient camera trap data management. Methods in Ecology and Evolution 7:1457–1462.

NIMBLE Development Team. 2019. NIMBLE: MCMC, particle filtering, and programmable

hierarchical modeling. R package version 0.9.0.

Otis, D. L., K. P. Burnham, G. C. White, and D. R. Anderson. 1978. Statistical inference from

capture data on closed animal populations. Wildlife Monographs 62:3–135.

Plummer, M. 2003. JAGS: a program for analysis of Bayesian graphical models using Gibbs

sampling. Proceedings of the 3rd International Workshop on Distributed Statistical Computing

(DSC 2003), March 20–22, Vienna, Austria.

R Development Core Team. 2019. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria.

Ramsey, D. S., P. A. Caley, A. Robley. 2015. Estimating population density from presence–

absence data using a spatially explicit model. The Journal of Wildlife Management 79:491–499.

Riaz, T., W. Shehzad, A. Viari, F. Pompanon, P. Taberlet, and E. Coissac. 2011. ecoPrimers:

inference of new DNA barcode markers from whole genome sequence analysis. Nucleic Acids

Research 39:e145.

Rich, L. N., D. A. Miller, D. J. Muñoz, H. S. Robinson, J. W. McNutt, and M. J. Kelly. 2019.

Sampling design and analytical advances allow for simultaneous density estimation of seven

sympatric carnivore species from camera trap data. Biological Conservation 233:12–20.

Rowcliffe, J. M., J. Field, S. T. Turvey, and C. Carbone. 2008. Estimating animal density using

camera traps without the need for individual recognition. Journal of Applied Ecology 45:1228–

1236.

Rowland, M. M., L. D. Bryant, B. K. Johnson, J. H. Noyes, M. J. Wisdom, and J. W. Thomas.

1997. The Starkey project: history, facilities, and data collection methods for ungulate research.

PNW-GTR-396. USDA Forest Service Pacific Northwest Research Station, Portland, Oregon,

USA.

Royle, J. A. 2004. N-mixture models for estimating population size from spatially replicated

counts. Biometrics 60:108–115.

Royle, J. A., and J. D. Nichols. 2003. Estimating abundance from repeated presence–absence

data or point counts. Ecology 84:777–790.

Royle, J. A., R. M. Dorazio, and W. A. Link. 2007. Analysis of multinomial models with

unknown index using data augmentation. Journal of Computational and Graphical Statistics 16:

67–85.

Royle, J. A., and K. V. Young. 2008. A hierarchical model for spatial capture–recapture data.

Ecology 89:2281–2289.

Royle, J. A., R. B. Chandler, R. Sollmann, and B. Gardner. 2013. Spatial capture-recapture.

Academic Press.

Schaub, M., O. Gimenez, A. Sierro, and R. Arlettaz. 2007. Use of integrated modeling to

enhance estimates of population dynamics obtained from limited data. Conservation Biology

21:945– 955.

1003

1004   Sikes, R. S., & Animal Care and Use Committee of the American Society of Mammalogists.

1005   2016. 2016 Guidelines of the American Society of Mammalogists for the use of wild mammals

1006   in research and education. Journal of Mammalogy 97:663–688.

1007

1008   Sollmann, R., M. M. Furtado, B. Gardner, H. Hofer, A. T. Jácomo, N. M. Tôrres, and L. Silveira.

1009   2011. Improving density estimates for elusive carnivores: accounting for sex-specific detection

1010   and movements using spatial capture–recapture models for jaguars in central Brazil. Biological

1011   Conservation 144:1017–1024.

1012

1013   Sollmann, R., B. Gardner, A. W. Parsons, J. J. Stocking, B. T. McClintock, T. R. Simons, K. H.

1014   Pollock, and A. F. O'Connell. 2013a. A spatial mark–resight model augmented with telemetry

1015   data. Ecology 94:553–559.

1016

1017   Sollmann, R., N. M. Tôrres, M. M. Furtado, A. T. de Almeida Jácomo, F. Palomares, S. Roques,

1018   and L. Silveira. 2013b. Combining camera-trapping and noninvasive genetic data in a spatial

1019   capture–recapture framework improves density estimates for the jaguar. Biological Conservation

1020   167:242–247.

1021

1022 Tourani, M., P. Dupont, M. A. Nawaz, and R. Bischof. 2020. Multiple observation processes in

1023 spatial capture–recapture models: How much do we gain? Ecology 101:e03030.

1024

1025 Wasser, S. K., B. Davenport, E. R. Ramage, K. E. Hunt, M. Parker, C. Clarke, and G. Stenhouse.

1026 2004. Scat detection dogs in wildlife research and management: application to grizzly and black

1027 bears in the Yellowhead Ecosystem, Alberta, Canada. Canadian Journal of Zoology 82:475–492.

1028

1029 Wasser, S. K., J. L. Keim, M. L. Taper, and S. R. Lele. 2011. The influences of wolf predation,

1030 habitat loss, and human activity on caribou and moose in the Alberta oil sands. Frontiers in

1031 Ecology and the Environment 9:546–551.

1032

1033 White, G. C. 1996. NOREMARK: population estimation from mark-resighting surveys. Wildlife

1034 Society Bulletin 24:50–52.

1035

1036 Whittington, J., M. Hebblewhite, and R. B. Chandler. 2018. Generalized spatial mark–resight

1037 models with an application to grizzly bears. Journal of Applied Ecology 55:157–168.

1038

1039 Wilson, G. J., and R. J. Delahay. 2001. A review of methods to estimate the abundance of

1040 terrestrial carnivores using field signs and observation. Wildlife Research 28:151–164.

1041

1042 Winegarner, C. E., and M. S. Winegarner. 1982. Reproductive history of a bobcat. Journal of

1043 Mammalogy 63:680–682.

1044

1045   Wisdom, M. J., M. M. Rowland, B. K. Johnson, and B. L. Dick. 2005. Overview of the Starkey

1046   Project: mule deer and elk research for management benefits. Pages 17–28 in M. J. Wisdom,

1047   editor. The Starkey project: a synthesis of long-term studies of elk and mule deer. Alliance

1048   Communications Group, Lawrence, Kansas, USA.

1049

1050

1051

TABLE 1: Description of data requirements for each model evaluated to estimate densities of four

carnivore species in northeast Oregon. The data collection sources used in this study are given in

parentheses. Models are ordered in terms of increasing data requirements. Sample sizes for the

number of independent detections for each data collection method are given in the last column,

separated by commas, for black bears, bobcats, cougars, and coyotes, respectively. For sample

sizes of genetic samples, we provide both the number of unique individuals and number of total

samples, respectively, separated by a slash (e.g. 83/165 indicates 83 unique coyotes were

detected from 165 genotyped scats). For brevity we only present sample sizes the first time a

certain data type is introduced.

1061

Model	Data Inputs	Sample size (for bear, bobcat, cougar, coyote)
Spatial Presence-Absence	1. Binary detections of unmarked individuals (cameras)	103, 30, 41, 283
Spatial Count	1. Counts of unmarked individuals (cameras)	159, 34, 48, 479
Spatial Mark-Resight	1. Encounter history of marked individuals (cameras) 2. Counts of unmarked individuals (cameras)	28, 9, 24, 55 131, 25, 24, 424
Generalized Spatial Mark-Resight	1. Encounter history of marked individuals (cameras) 2. Counts of unmarked individuals (cameras) 3. Encounter history of marking process (live traps)	 6, 4, 6, 9
Spatial Capture-Recapture	1. Encounter history of genotypes (scat)	31/40, 33/68, 7/13, 83/165
Spatial Capture-Recapture + Marking Process	1. Encounter history of genotypes (scat) 2. Encounter history of marking process (live traps)	
Hybrid Generalized Spatial Mark-Resight +	1. Encounter history of marked individuals (cameras) 2. Counts of unmarked individuals (cameras) 3. Encounter history of marking process (live traps) 4. Encounter history of genotypes (scats)	

Spatial Capture- Recapture

1062

1063

1065   FIG. 1: The study area in northeast Oregon in which the densities of black bears, bobcats,

1066   cougars, and coyotes were estimated using a suite of spatially-explicit models fit to data

1067   collected by remote cameras and noninvasive genetic sampling. The black polygon represents

1068   the Starkey Experimental Forest and Range boundary, green points are camera locations, and

1069   blue lines are scat detection dog survey tracks.

1070

FIG. 2: a) Scat located by detection dogs subsequently analyzed for species ID and genotyped to

determine individual identity. b) A GPS-collared bear detected on remote camera. The number

on the collar belting and location of the ear tag permitted individual identification of this animal.

c) Immobilized coyote fit with GPS collar. Note the colored shrinkwrap around collar banding

and unique number aiding in individual identification when photographed on remote cameras. d)

GPS collar positions for coyotes (light green), cougars (blue), bobcats (purple), and bears (red)

during the time sampling was conducted. The red outline is the Starkey Experimental Forest and

Range boundary.

1079

FIG. 3: A directed acyclic graph (DAG) displaying the inputs (labeled outside curly brackets) and

relationships between data and parameters for the hybrid gSMR+SCR (generalized spatial mark-

resight and spatial capture-recapture) model. Dependent variables (i.e. detection data) are

represented by ovals. Solid lines connect parameters to data and independent variables are

connected to dependent variables by dotted lines. Parameters appearing inside overlapping boxes

are jointly estimated across different data sources and submodels. Telemetry data can optionally

be incorporated into every model. Note that gSMR is composed of 1) conventional spatial mark-

resight (SMR) and 2) the marking process, a SCR model describing the process by which a

subset of animals are physically captured and marked allowing them to be resighted.

1089

FIG. 4: Split violin plots displaying posterior distributions for the densities of black bears,

bobcats, cougars, and coyotes across a suite of methods for base models (i.e. those without sex or

behavioral-specific detection parameters). Within each panel, models are ordered in terms of

increasing data requirements with the most data-hungry models on the right. For each model

type, the left (blue) half of the violin excludes GPS telemetry data and the right (green) half

includes GPS data from a telemetered subset of individuals of each species. The white boxes

display the 95% highest posterior density interval, and the middle black bar indicates the

posterior mode. Densities are presented as number of animals per 100 km^2. Note that the spatial

count and spatial presence-absence models for each species required a different y-axis scale than

the other model types. SC = spatial count, SPA = spatial presence absence, SMR = conventional

spatial mark-resight, gSMR = generalized spatial mark-resight, SCR = spatial capture-recapture,

and hybrid = combined gSMR+SCR.

1102

FIG. 5: Coefficients of variation for density estimates for bears, bobcats, cougars, and coyotes

(top left, clockwise). For each model, the left (blue) bar excludes GPS data and the right (green)

bar includes GPS data from a telemetered subset of individuals of each species. The coefficient

of variation was calculated as the posterior standard deviation divided by the posterior mean of

animal density. Note that the sample size for the cougar SCR models was small ($N = 13$ scats)

and likely caused the relative imprecision for those models.

1109

FIG. 6: Percent difference in the posterior mode of animal density from a suite of base models

compared to the base SCR+GPS (spatial capture-recapture including telemetry data) model for

black bears, bobcats, cougars, and coyotes. Note that the spatial count (SC) and spatial presence-

absence (SPA) models for each species required a different y-axis scale than the other model

types. The dotted red line refers to no difference from the base SCR+GPS model; bars above the

line indicate higher estimates and bars below the line indicate lower estimates than the base

SCR+GPS model. SMR = conventional spatial mark-resight, gSMR = generalized spatial mark-

resight, and Hybrid = the combined SCR+gSMR model.

1118

FIG. 7: Comparison of SMR+GPS, gSMR+GPS, and Hybrid+GPS density estimates from base

models and a set of post-hoc models that were fit to attempt to more fully adhere to model

assumptions and/or explain more heterogeneity in detection rate in relation to SCR+GPS density

estimates. Bars indicate the percent difference in the posterior mode of animal density compared

to genetic SCR+GPS estimates for base models, sex-specific models (i.e. those in which

detection and spatial scale parameters were allowed to vary by sex), and reduced sampling

period models (reduced from 140 days to 70 days of camera sampling) for black bears, bobcats,

cougars, and coyotes. For coyotes, an additional model type with detection and spatial scale

parameters varying by behavioral status (i.e. transient vs resident) is compared. The horizontal,

dotted red line indicates no difference from the SCR+GPS model; bars above the line indicate

higher estimates and bars below the line indicate lower estimates than the SCR+GPS model.

Comparisons are made with respect to sex-specific SCR+GPS estimates for 'sex' models,

behavioral status-specific SCR+GPS estimates for 'behavioral status' models, and base

SCR+GPS estimates otherwise.

Fɪɢ. 8: Split violin plots displaying posterior distributions for animal density estimates for sex-

specific models (bobcat, bear, cougar) and behavioral specific models (resident vs transient

status in coyotes). For each model type, the left (blue) half of the violin excludes GPS telemetry

data and the right (green) half includes GPS data from a telemetered subset of individuals of each

species. The white boxes display the 95% highest posterior density interval, and the middle black

bar indicates the posterior mode. Densities are presented as number of animals per 100 km^2.

SMR = conventional spatial mark-resight, gSMR = generalized spatial mark-resight, SCR =

spatial capture-recapture, and hybrid = combined gSMR+SCR.

1141

1142   Fɪɢ. 9: Coefficients of variation (defined as the standard deviation divided by the posterior mean)

1143   displaying the relative precision of density estimates using SMR+GPS, gSMR+GPS, and

1144   Hybrid+GPS for black bears, bobcats, cougars, and coyotes for base models, sex-specific

1145   models, and reduced sampling period models. For coyotes, an additional model type with

1146   detection and spatial scale parameters varying by behavioral status is presented.

1147

1148 **Figure 1**

1149

1150

Figure 2

1154 **Figure 3**

1155

1156

1157 **Figure 4**

1158

1159

1160

1161 **Figure 5**

1162

1163

1164

1165　**Figure 6**

1166

1167

1168 **Figure 7**

1169

1170

1171 **Figure 8**

1172

1173 **Figure 9**

1174