1

The transcription factor TRF2 has a unique function in regulating

cell cycle and apoptosis

Adi Kedmi[*], Anna Sloutskin[*], Natalie Epstein[*], Lital Gasri-Plotnitsky[*], Debby

Ickowicz[*], Irit Shoval[*], Tirza Doniger[*], Eliezer Darmon[*], Diana Ideses[*], Ziv Porat[†],

Orly Yaron[*], and Tamar Juven-Gershon[*1]

[*] The Mina and Everard Goodman Faculty of Life Sciences, Bar-Ilan University,

Ramat Gan 5290002, Israel; and

[†] The Flow Cytometry Unit, Life Sciences Core Facilities, Weizmann Institute of

Science, Rehovot 7610001, Israel

[1] Correspondence: The Mina and Everard Goodman Faculty of Life Sciences, Bar-

Ilan University, Ramat Gan 5290002, Israel. E-mail: tamar.gershon@biu.ac.il

ABSTRACT

Background: Diverse biological processes and transcriptional programs are

regulated by RNA polymerase II (Pol II), which is recruited by the general transcription

machinery to the core promoter to initiate transcription. TRF2 (TATA-box-binding

protein-related factor 2) is an evolutionarily conserved general transcription factor that

is essential for embryonic development of *Drosophila melanogaster*, *C. elegans*,

zebrafish and *Xenopus*. Nevertheless, the cellular processes that are regulated by

TRF2 are largely underexplored.

Results: Here, using *Drosophila* Schneider cells as a model, we discovered that TRF2

regulates apoptosis and cell cycle progression. We show that TRF2 knockdown

results in increased expression of distinct pro-apoptotic genes and induces apoptosis.

Using flow cytometry, high-throughput microscopy and advanced imaging-flow

cytometry, we demonstrate that TRF2 regulates cell cycle progression and exerts

distinct effects on G1 and specific mitotic phases. RNA-seq analysis revealed that

TRF2 controls the expression of *Cyclin E* and the mitotic cyclins, *Cyclin A*, *Cyclin B*

and *Cyclin B3*, but not *Cyclin D* or *Cyclin C*. To identify proteins that could account for

the observed regulation of these cyclin genes, we searched for TRF2-interacting

proteins. Interestingly, mass spectrometry analysis of TRF2-containing complexes

identified GFZF, a nuclear glutathione S-transferase implicated in cell cycle regulation,

and Motif 1 binding protein (M1BP). TRF2 has previously been shown to interact with

M1BP and M1BP has been shown to interact with GFZF. Furthermore, available ChIP-

exo data revealed that TRF2, GFZF and M1BP co-occupy the promoters of TRF2-

regulated genes. Using RNAi to knockdown the expression of either M1BP, GFZF,

TRF2 or their combinations, we demonstrate that although GFZF and M1BP interact

with TRF2, it is TRF2, rather than GFZF or M1BP, that is the main factor regulating

the expression of *Cyclin E* and the mitotic cyclins.

Conclusions: Our findings uncover a critical and unanticipated role of a general

transcription factor as a key regulator of cell cycle and apoptosis.

Keywords

Basal transcription machinery, RNA polymerase II, gene expression, TATA box-

binding protein (TBP), TBP-related factor 2 (TRF2), cyclin genes.

BACKGROUND

Multiple biological processes and transcriptional programs are regulated by RNA

polymerase II (Pol II). The initiation of transcription of protein-coding genes and

distinct non-coding RNAs occurs following the recruitment of Pol II to the core

promoter region by the general/basal transcription machinery (1-4). The core

promoter, which directs accurate initiation of transcription and encompasses the

transcription start site (TSS), may contain short DNA sequence elements/motifs,

which confer specific properties to the core promoter (1, 4-10). The first step in the

recruitment of Pol II to initiate transcription is the binding of TFIID, which is

composed of TATA-box-binding protein (TBP) and TBP-associated factors.

Remarkably, although TBP is considered a universal general transcription factor,

robust Pol II transcription is observed in mouse TBP-/- blastocysts, indicating the

existence of TBP-independent Pol II transcription *in vivo* (11). The complexity of

transcription is also manifested by the existence of diverse transcriptional regulators,

among which are the TBP family members. There are three TBP family members in

Drosophila melanogaster: TBP, TRF1 and TRF2 (reviewed in (12-16)). TRF1, the

first *Drosophila* TBP family member identified, is insect specific (17). An evolutionary

conservation analysis indicated that TRF2 (also known as TLP (TATA-like protein),

TLF (TBP-like factor), TRP (TBP-related protein) and TBPL1 (TBP-like 1)), is highly

conserved in evolution (12, 18-22) and is present in all bilaterian organisms, but not

in any of the non-bilaterian genomes available (19). It was further discovered that

TRF2, which is involved in Pol II transcription, evolved by duplication of the TBP

gene (19). Yet, unlike TBP and TRF1, TRF2 does not bind TATA-box containing

promoters (19, 20, 22). There are two *Drosophila* TRF2 protein isoforms that result

from an internal translation initiation: the evolutionarily conserved short isoform (632

aa; typically referred to as "TRF2") and a long *Drosophila*-only isoform (1715 aa), in

which the same short amino acid sequence is preceded by an N-terminal domain

(23). TRF2 affects early embryonic development of *Drosophila*, *C. elegans*, zebrafish

and *Xenopus*, differentiation and morphogenesis (23-32). Mouse TRF2 is essential

for spermiogenesis (33-35).

One of the open questions in the transcriptional regulation field is what are the

cellular functions of TRF2. Despite its importance in development, the cellular

processes that are regulated by TRF2 remain largely underexplored. To identify and

characterize the cellular processes that are regulated by TRF2, we used *Drosophila*

S2R+ cells as a model and knocked-down the expression of TRF2. We discovered

that reduced expression of TRF2 (but not TBP or TRF1) results in apoptosis and

increased expression of key, yet not all, pro-apoptotic genes (including *rpr, hid, p53*),

suggesting this is not a general stress response. Surprisingly, not only that TRF2

regulates apoptotic cell death, reduced expression of TRF2 (but not its family

members, TBP or TRF1), exerts distinct effects on G1, G2/M and specific mitotic

phases, as demonstrated by quantitative high-throughput imaging flow cytometry.

We further discovered that TRF2 controls the expression of *Cyc E* and the mitotic

Cyc A, *Cyc B* and *Cyc B3* genes. Using mass spectrometry analyses of TRF2-

interacting proteins and available ChIP-exo data, we demonstrate the co-occupancy

of TRF2, GFZF (GST-containing FLYWCH zinc-finger protein) and M1BP (motif 1

binding protein) in the majority of promoters bound by each of the three factors.

Remarkably, the promoters of the TRF2-regulated mitotic cyclins and *Cyclin E* are

bound by the three factors, whereas the promoters of *Cyclin C* and *Cyclin D*, which

are not regulated by TRF2, are not bound. Furthermore, the Motif 1 sequence

element is enriched in the promoters of genes bound by all three proteins,

suggesting the involvement of GFZF and M1BP as co-factors in TRF2-regulated cell

cycle progression. Moreover, we demonstrate that TRF2, rather than M1BP or

GFZF, is the main factor that regulates the expression of the mitotic cyclins and

Cyclin E. Importantly, while general/basal transcription factors might be viewed as

having a somewhat "generic" role, our findings emphasize the unique, unanticipated

functions of *Drosophila* TRF2 as an essential factor for cell cycle progression and

apoptotic cell death.

RESULTS

Knockdown of TRF2 expression results in apoptotic cell death and induced

expression of key pro-apoptotic genes

The TBP-related transcription factor TRF2 is a key general/basal transcription factor

(reviewed in (12-16)), yet the cellular processes that are regulated by TRF2 remain

largely underexplored. To investigate the cellular functions of TRF2, we used

Drosophila S2R+ cells as a model, and knocked down its expression by RNAi using

non-overlapping dsRNA probes (Additional file 1: Figure S1a). TBP knockdown was

used as a control throughout this study. The resulting reduction in protein expression

was verified by western blot analysis (Additional file 1: Figure S1b, c). We

consistently observed significant cell death following TRF2 knockdown, as evident by

microscopic examination and by the reduced amounts of total RNA purified from

TRF2-RNAi treated cells, as compared to mock-treated cells. To specifically

investigate whether TRF2 plays an important role in apoptosis, we performed

Annexin V/PI analysis to detect early and late apoptotic cell death by flow cytometry

analysis. PI is excluded from cells with intact membranes, while dead and damaged

cells have membranes that are permeable to PI. Annexin V binds phospholipids that

are exposed in cells undergoing apoptosis. Hence, cells that are both Annexin V and

PI negative are considered viable, while cells that are in early apoptosis are Annexin

V positive and PI negative, and cells that are in late apoptosis or already dead are

both Annexin V and PI positive. S2R+ cells were incubated for three days with either

one of the four non-overlapping dsRNA probes directed against *Trf2*, a dsRNA probe

against *Tbp*, a dsRNA probe against *Trf1* or a dsRNA probe against the

homeodomain transcription factor *exd*, as a negative control. Cells were harvested

and stained with Annexin-V FITC/PI. Approximately 30% of cells stain positive for

Annexin V following TRF2 knockdown by the four different dsRNA probes, ~2 fold

higher than the mock and the exd-RNAi treated cells (Fig. 1). Notably, cell death

induced by TBP or TRF1 dsRNA probes (an average of 20%) was not as

pronounced as the cell death induced by either of the four TRF2 dsRNA probes.

These findings emphasize the unique effects of TRF2, as compared to TBP, on

apoptosis.

To identify the targets that are unique to TRF2, we used RNAi to knockdown

either TRF2 or TBP in *Drosophila* S2R+ cells and performed RNA-seq analysis at

two time points, 48h and 72h. In order to further understand how TRF2 knockdown

results in cell death, we searched our RNA-seq data for pro-apoptotic genes that are

regulated by TRF2. The inhibitor of apoptosis protein (IAP) family has already been

shown to play an important role in cell survival (reviewed in (36-38)). *Drosophila*

Death-associated IAP-1 (DIAP1), a key member of the IAP family, inhibits apoptosis

by binding to the active Caspase 9-like Dronc, and functioning as an E3-ubiquitin

ligase to promote its degradation. The *reaper* (*rpr*), *head involution defective* (*hid*),

and *grim* genes encode IAP antagonists that bind DIAP1, disrupt its interactions with

caspases and target it for degradation, resulting in caspase activation and cell death.

Scylla (*scyl*) has been implicated in developmental cell death (39). Indeed, the RNA-

seq analysis revealed a significant upregulation of *rpr* and *scyl* (Additional file 2:

Table S1). We thus decided to knockdown either TRF2 (probes #1 and #2), exd,

TBP or TRF1 and examine by reverse transcription-qPCR the expression of multiple

genes implicated in the apoptotic machinery: the IAP antagonists *rpr*, *hid* and *grim*

(reviewed in (37)), *scyl* (39), the BCL-2 family members *buffy* (40) and *Death*

executioner Bcl-2 (*Debcl*) (41-44), and the Caspase 9-like *Dronc* (reviewed in (37)).

Remarkably, reducing the expression levels of TRF2 (but not exd, TBP or TRF1)

resulted in increased expression levels of *rpr*, *hid* and *scyl* (Fig. 2a). The detected

increase in *Buffy* expression levels was not statistically significant. The expression of

grim, *Debcl* and *Dronc* was not significantly altered by TRF2 knockdown (Fig. 2a),

suggesting that this is not a general stress response, and that TRF2 specifically

regulates the expression of distinct, but not all, pro-apoptotic genes.

The p53 tumor suppressor gene is a key regulator of both cell cycle

progression and programmed cell death (reviewed, for example, in (45, 46)). p53 is

expressed at low levels in S2R+ cells (www.flybase.org). Surprisingly, the RNA-seq

analysis revealed its upregulation upon TRF2 knockdown (Additional file 2: Table

S1). Examination of p53 expression by reverse transcription-qPCR following

knockdown of either TRF2 (probes #1 and #2) or TBP reproduced the upregulation

trend of p53 expression by TRF2. Notably, TBP knockdown did not alter the

expression of p53, manifesting the unique characteristics of gene regulation via

TRF2. Interestingly, it was previously demonstrated that p53 regulates the

expression of *rpr* and *hid*, which in turn, induce apoptosis (47, 48). Hence, it is

possible that the observed upregulation of *rpr* and *hid* following the knockdown of

TRF2 (Fig. 2a, b) is mediated via p53. Unfortunately, multiple attempts to reduce the

levels of endogenous p53 by two different dsRNA probes, were unsuccessful. Thus,

one cannot exclude the possibility that the increased expression of p53 following

TRF2 knockdown, may partially contribute to the upregulation of *rpr* and *hid*

expression.

TRF2 exerts distinct effects that are independent of TBP

TRF2 may repress transcription involving TBP or TFIID, probably by recruiting TFIIA

(20, 49). One can suggest that TRF2 is, in some respects, antagonistic to TBP, and

the observed upregulation of pro-apoptotic genes following TRF2 knockdown (Fig. 2)

results from TBP activity, which is no longer obstructed by TRF2. To examine

whether the observed upregulation of *rpr* and *hid* results from TBP activity, we

overexpressed TBP in S2R+ cells that were either incubated with TRF2 dsRNA

probes or mock treated. Although overexpression of TBP was clearly evident (>100

fold; Additional file 3: Figure S2), no differences in the levels of *rpr* and *hid* were

observed upon its overexpression (Additional file 3: Figure S2). Furthermore,

overexpression of TBP following TRF2 knockdown did not alter *rpr* or *hid* expression.

Thus, we conclude that the observed upregulation of *rpr* and *hid* following TRF2

knockdown results from the activity of TRF2 as a unique transcription factor, rather

than a TBP antagonist.

Knockdown of endogenous TRF2 expression results in altered cell cycle

distribution and G1 arrest

To identify specific cellular processes that are regulated by TRF2, we performed

Gene Ontology (GO) terms analysis of genes that were either downregulated or

upregulated following TRF2 knockdown (using string.db.org). Surprisingly, we

discovered that genes that were downregulated following TRF2 knockdown are

enriched for cell cycle and mitotic cell cycle processes, while genes that were

upregulated following TRF2 knockdown are enriched for response to stimulus and

stress (Additional file 2: Table S1). Interestingly, while the number of genes that were

downregulated was only half of the number of the upregulated genes following TRF2

knockdown (337 vs. 684 genes, respectively), the enrichment scores ($-\log_{10}$(P

values)) of the downregulated genes were 5-fold higher. We thus decided to

examine the effects of TRF2 knockdown on cell cycle distribution. To knockdown the

expression of the endogenous genes, S2R+ cells were incubated for three days with

dsRNA probes against *Trf2*, *Tbp*, *Trf1* or *exd*, as a negative control. Cells were

harvested, fixed and analyzed by flow cytometry. Control S2R+ cells display a

normal profile with an average of ~31% of cells in G1 phase and with an average of

~34% of cells in G2/M phase, similarly to mock treated cells (which were processed

similarly, but were not incubated with any dsRNA; Fig. 3). Following TRF2

knockdown by either one of the four dsRNA probes, we observed a distinct decrease

in the fraction of cells in G2/M phase (an average of ~20%), as well as the fraction of

cells in S phase (an average of ~20%) with a concomitant increase in the fraction of

cells in G1 phase (an average of ~55%). Remarkably, these effects are unique to

TRF2 knockdown, as the knockdown of its family members TBP or TRF1 resembles

the cell cycle distribution of control and mock treated cells (Fig. 3).

Our RNA-seq analysis reveals subsets of genes that are involved in S and

G2/M phases. In addition, the cell cycle analysis indicates that TRF2 plays a role in

S and/or G2/M phases. To determine if TRF2 affects cell cycle progression to S

phase, S2R+ cells were arrested in G1 with 1mM hydroxyurea (HU) for 18h following

knockdown of either TRF2 (dsRNA probes #1 and #2) or TBP, and then released to

cycle by replacing the medium with fresh medium. As can be seen in Figure 4, HU

treatment (0h) resulted in accumulation of cells in G1 (~55%). Two hours following

the release, mock treated cells returned to cycle (indicated by the decrease in the

number of cells in G1), and the fraction of cells in S and G2/M increased. Similarly,

cells in which TBP was depleted by RNAi, returned to cycle. Surprisingly, unlike

mock or TBP RNAi-treated cells, cells in which TRF2 was depleted by either one of

the two probes, remained in G1 (~55%) and did not return to cycle (Fig. 4, Additional

file 4: Figure S3). Moreover, even 8h following the removal of HU, cells in which

TRF2 was knocked-down remained G1-arrested. Our findings imply that

endogenous TRF2 is involved in progression into S phase.

TRF2 regulates the expression of specific cyclin genes

To further explore the connection between TRF2 and cell cycle progression, we turned to our RNA-seq data for cell cycle-related genes that may be influenced by knockdown of TRF2. We discovered that following TRF2 knockdown, the expression of the *Cyc A*, *Cyc B*, *Cyc B3* and *Cyc E* cell cycle regulators was significantly reduced (over 2.3-fold). Notably, the expression levels of *Cyc C* and *Cyc D* were unchanged (Additional file 2: Table S1). In order to verify the effect of TRF2 knockdown on the expression of these genes, reverse transcription-qPCR analysis of endogenous cyclin genes was performed on mock, TRF2 (probes #1 and #2), exd, TBP or TRF1 RNAi-treated cells. Knockdown of TRF2 by either probe #1 or #2 significantly reduces the expression of *Cyc A*, *Cyc B* and Cyc *B3*, as compared to mock treated cells, much more than knockdown of TBP or TRF1 (Fig. 5a). As there was a difference between the effects of probe #1 and #2 on *Cyc E* expression, 4 non-overlapping TRF2 dsRNA probes were used to assess the effect of TRF2 knockdown on *Cyc E* expression (Fig. 5b). Notably, each of the 4 non-overlapping TRF2 dsRNA probes reduces *Cyc E* expression. Unlike *Cyc D* (which is required for G1 progression (50, 51)), the expression of *Cyc E* (which promotes G1-S transition (52, 53)) was reduced following TRF2 knockdown (Fig. 5a, b). We next tested whether TRF2 knockdown affects the protein levels of Cyc A, Cyc B and Cyc E. Unfortunately, we were unable to detect endogenous Cyc A and Cyc B protein expression using publicly available anti- *Drosophila* Cyc A and Cyc B antibodies (data not shown). Remarkably, using anti-*Drosophila* Cyc E antibodies, we observed a distinct reduction in Cyc E protein levels following knockdown of TRF2 (but not TBP), using both TRF2 probes (Fig. 5c), further suggesting that TRF2 regulates G1-S transition by modulating the expression of *Cyc E*.

TRF2 regulates distinct mitotic phases

We were intrigued by the downregulation of the mitotic cyclin gene expression (*Cyc A*, *Cyc B* and *Cyc B3*) following TRF2 knockdown (Fig. 5a) and decided to explore the effect of TRF2 downregulation on mitotic phases. To this end, we knocked-down the expression of TRF2 or TBP and stained cells for mitotic chromatin (anti-phospho-Histone H3 (Ser10)), DNA (Hoechst) and filamentous Actin (Phalloidin). These allowed us to analyze the fraction of cells in mitosis following TRF2 knockdown (Fig. 6a). To determine the number of mitotic cells, we developed a pipeline to automatically detect the Hoechst and phospho-Histone H3 signals. Notably, there was a reduction in the number of cells undergoing mitosis following TRF2 knockdown, as compared to mock or TBP RNAi treated cells (Fig. 6b).

To explore the effect of TRF2 on G2/M, we sought to synchronize cells in G2/M. Unfortunately, we were unable to synchronize cells in G2/M in a reversible manner (see methods), and thus we could not perform G2/M block-release experiments. Nevertheless, we succeeded in discerning the effects of TRF2 on specific mitotic phases by employing advanced imaging-flow cytometry analysis (ImageStreamX mark II imaging flow-cytometer, Amnis Corp, Seattle, WA, Part of EMD Millipore). Imaging-flow cytometry analysis combines the high-quality imaging and functional insights of microscopy with the speed, sensitivity, and phenotyping abilities of flow cytometry. We knocked down the expression of TRF2 or TBP and stained cells with anti-phospho-Histone H3 (Ser10) antibodies and Hoechst. A total of 40,000 cells of each treatment were analyzed by an ImageStream flow cytometer to determine the number of cells in each mitotic phase, according to their nuclear morphology (Fig. 6c-f, Additional file 5: Figure S4, Additional file 6: Table S2). Remarkably, although 40,000 cells were analyzed in each experiment, only a few

hundred cells were mitotic, and following TRF2 knockdown there was an even bigger

reduction in the total number of mitotic cells (Fig. 6b, d). Notably, despite the overall

reduction in the mitotic cell population, knockdown of TRF2 (but not TBP) resulted in

a significant accumulation of cells in anaphase and telophase (Fig. 6e, f).

To validate the accurate identification of mitotic cells, we used Colchicine as a

control. Colchicine treatment resulted in accumulation of cells in mitosis, specifically

in anaphase (Fig. 6e). Multiple studies have shown that Colchicine disrupts the

metaphase to anaphase transition (see for example, (54)), yet, the increase in

anaphase has also been documented (55). Furthermore, it is established that

different cell types behave differently during mitosis in the presence of drugs that

disrupt microtubules function (56). Interestingly, morphological examination of the

Colchicine-treated S2R+ cells indicated aberrant DNA staining patterns and mal-

oriented clumped chromosomes in prophase, metaphase and anaphase cells

(Additional file 7: Figure S5), in line with the absence of a spindle. Notably, mal-

oriented un-centered chromosomes, such as those observed in Colchicine-treated

cells, were not observed in TRF2-RNAi treated cells.

The effects of TRF2 knockdown on cyclin gene expression correlates with the

promoter occupancies of TRF2 and its co-factors GFZF and M1BP

To better understand how TRF2 regulates the expression of the cyclin genes, we

turned our attention to TRF2-interacting proteins. TRF2 was recently shown to

interact with M1BP (motif 1 binding protein) (57). Interestingly, M1BP was recently

demonstrated to interact with GFZF, a nuclear glutathione S-transferase protein that

has been implicated in cell cycle regulation (58). We suspected that GFZF may

interact with TRF2. Indeed, using FLAG immuno-affinity purification from FLAG-HA

TRF2-inducible S2R+ cells followed by mass spectrometry analysis, we discovered

that both M1BP and GFZF are in complex with the evolutionarily conserved TRF2

(also known as short TRF2), but not with the long *Drosophila*-only TRF2 isoform or

with TBP (Table 3 and Additional file 8: Table S3). This prompted us to examine the

occupancy of TRF2, M1BP and GFZF in the vicinity of the TSSs (-100 to +100

relative to the TSS) of TRF2-regulated cyclin genes, using publicly available TRF2,

M1BP and GFZF ChIP-exo analyses in *Drosophila* S2R+ cells (GSE97841,

GSE105009) (57, 58). We examined the number of bound sites, the average peak

scores and the maximum peak scores of cyclin genes and several ribosomal protein

genes for comparison (Table 4). As expected, M1BP, TRF2 and GFZF co-occupy

the promoters of the ribosomal protein genes. Interestingly, while M1BP occupies the

-100 to +100 regions of all the examined cyclin genes, both TRF2 and GFZF occupy

the -100 to +100 regions of *Cyc A*, *Cyc B*, *Cyc B3* and *Cyc E*, and to a lesser extent

the promoters of *Cyc C* and *Cyc D*, which are not regulated by TRF2. The

occupancies of the three proteins is especially striking in the vicinities of the *Cyc B*

and *Cyc B3* TSSs. Thus, the effects of TRF2 knockdown on cyclin gene expression

generally correlate with the occupancies of both TRF2 and GFZF in the -100 to +100

regions of *Cyc A*, *Cyc B*, *Cyc B3* and *Cyc E.*

To characterize the co-occupancies of TRF2, GFZF and M1BP in a genome-

wide manner, we examined the binding of each factor to *Drosophila* promoter

regions (± 50 bp relative to FlyBase annotated TSSs). Remarkably, a major fraction

of promoters is bound by all three transcription factors (Fig. 7a). Reassuringly, the

co-bound promoters include the TRF2-regulated *Cyc A*, *Cyc B*, *Cyc B3* and *Cyc E*,

but not *Cyc C* and *Cyc D* promoters, which are not regulated by TRF2.

To better decipher the characteristics of the co-bound promoters, we used MEME

(59) to detect enriched sequence motifs. Interestingly, the top enriched motif in

promoters that are bound by all three proteins (Fig. 7b) closely resembles Ohler

Motif 1 (60), also detected in M1BP ChIP-exo analysis (57). We next analyzed the

core promoter composition of the co-bound promoters, using the ElemeNT algorithm

(61). Strikingly, the co-bound promoters are depleted for the TATA-box motif and

enriched for the TCT and Motif 1 core promoter elements, as compared to the

genomic distribution of core promoter elements (Fig. 7c).

To examine the contribution of M1BP and GFZF to the effect of TRF2

knockdown on cyclin gene expression, we used RNAi to knockdown the expression

of either M1BP, GFZF, TRF2 or their combinations. The use of each of the dsRNA

probes resulted in a significantly reduced expression of the targeted gene (Fig. 7d).

Surprisingly, M1BP knockdown resulted in increased expression of *Trf2*, *gfzf* and

Cyc E. Since both TRF2 and M1BP were previously shown to affect the expression

of ribosomal protein genes (32, 62), we tested whether their knockdown affects the

expression of several ribosomal target genes, namely, *RpL30*, *RpLP1* and *RpLP2*.

While TRF2 and GFZF knockdown did not affect their expression, M1BP knockdown

resulted in significantly increased expression of *RpLP2* (Additional file 9: Figure

S6a). Notably, this effect was not general, but rather specific to distinct cyclin and

ribosomal protein genes (Fig. 7d, e and Additional file 9: Figure S6), as the

expression of *CG12493* and *SgII*, two previously identified M1BP targets (62), was

reduced following M1BP knockdown (Additional file 9: Figure S6b).

The expression levels of *Cyc A* and *Cyc B* were specifically reduced following TRF2

knockdown, but not following GFZF or M1BP knockdown (Fig. 5a and Fig. 7d, e).

Cyc D expression was not affected by either of these single factor knockdowns (Fig.

7d), as in Figure 5. As can be observed by TRF2 knockdown, as well as the

combined knockdown of TRF2, GFZF and M1BP, the expression pattern of *Cyc A*

and *Cyc B* are mostly influenced by TRF2 knockdown. *Cyc E* exhibits a composite

pattern: it is reduced following TRF2 knockdown, but the combined knockdowns of

TRF2 and M1BP or GFZF seem to restore its expression as compared to mock

treatment. Notably, *Cyc D* expression pattern is the least affected by the different

knockdown combinations.

Taken together, these data suggest that the observed effects of TRF2

knockdown on cell cycle progression (Figs. 3, 4 and 6) are, at least partially,

mediated by the reduced expression of *Cyc E*, *Cyc A* and *Cyc B* following TRF2

knockdown (Figs. 5 and 7d, e). Importantly, the co-occupancy of TRF2, GFZF and

M1BP in the promoters of these cyclin genes, the enrichment of Motif 1 in their

promoters and the expression patterns of the cyclin genes following the knockdown

of either TRF2 alone, or in combination with GFZF and/or M1BP, imply that GFZF

and M1BP may serve as co-factors in TRF2-regulated cell cycle progression. Yet, it

is TRF2 that is the major transcription factor regulating the expression pattern of the

abovementioned cyclin genes.

389   DISCUSSION

In this study, we discovered that knockdown of the general/basal transcription factor

TRF2 results in increased expression of the *rpr*, *hid* and *p53* pro-apoptotic genes,

which is in line with the observed increased apoptotic cell death following TRF2

knockdown and with the extensively characterized involvement of p53 in apoptosis,

G1 arrest and G2 arrest (reviewed, for example, in (45)). A recent study has

demonstrated that human TRF2 interferes with MDM2 binding and ubiquitination of

p53, leading to p53 protein stabilization (63). Our results provide evidence for

another level of p53 regulation, *i.e.*, transcriptional regulation. Furthermore, in a

similar manner to the p53-MDM2 negative feedback loop in vertebrates (reviewed in

(45)), activation of *Drosophila* p53 transcriptionally activates the *companion of reaper*

(*corp*) gene (64-66), which in turn, negatively regulates its activity (67). The fact that

corp expression in *Drosophila* S2R+ is rather low and not altered by TRF2

knockdown (Additional file 2: Table S1), may provide support for p53-independent

regulation of *rpr* and *hid* by TRF2.

Furthermore, we examined the expression of multiple additional genes

implicated in apoptosis following TRF2 knockdown. *Scylla* is pro-apoptotic (39) and

its upregulation is in line with the observed cell death following TRF2 knockdown.

Buffy, a *Drosophila* Bcl2 family member, however, has been shown to act in an anti-

apoptotic manner, but has also been shown to cause a G1-S arrest (40). Thus, its

upregulation following TRF2 knockdown may contribute to the observed G1-S

accumulation of cells. These effects are in line with studies suggesting the existence

of TRF2-regulated transcriptional systems (10, 16, 32, 68, 69) and are unlikely to

represent a general stress response, as the expression of *Debcl* and *Dronc* is not

affected by TRF2 knockdown.

Interestingly, we discovered that TRF2 knockdown results in accumulation of

cells in G1 and in reduction in the number of cells in S and G2/M phases. G1/S

transition is regulated by Cyclin E activity, while S phase and G2/M transition are

regulated by Cyclin A activity, and transition into and within mitosis is regulated by

the activities of Cyclin B and Cyclin B3. Remarkably, TRF2 knockdown in S2R+ cells

resulted in reduced expression of *Cyc E*, *Cyc A*, *Cyc B* and *Cyc B3* (but not *Cyc D*),

suggesting that TRF2 regulates cell cycle progression by modifying the expression of

specific cyclins. The reduced expression of *Cyc A*, *Cyc B* and *Cyc B3* and the

reduction of the number of cells undergoing mitosis following TRF2 knockdown, are

in line with previous studies, which demonstrated inhibition of nuclear mitotic entry in

Drosophila embryos following simultaneous knockdown of *Cyc A*, *Cyc B* and *Cyc B3*

(70). Interestingly, Cyclin A and Cyclin B have previously been shown to inhibit

metaphase-anaphase transition, whereas Cyclin B3 promotes it (71). Thus, the

specific accumulation of cells in anaphase and telophase observed by imaging flow

cytometry (Fig. 6e, f), could result from the reduced expression of *Cyc A* and *Cyc B*

following TRF2 knockdown (Fig. 5). Notably, to the best of our knowledge, this study

is the first to employ imaging-flow cytometry in the analysis of *Drosophila* cells.

To examine whether TRF2-interacting proteins could account for the observed

regulation of *Cyc E*, *Cyc A*, *Cyc B* and *Cyc B3* (but not *Cyc D* or *Cyc C*), we

searched for TRF2-interacting proteins. TRF2 has been shown to interact with M1BP

(57) and M1BP has been shown to interact with GFZF, a nuclear glutathione S-

transferase implicated in cell cycle regulation (57). Our proteomic analyses revealed

that both M1BP and GFZF preferentially interact with TRF2, but not with TBP (Table

3 and Additional file 8: Table S3). Remarkably, examination of publicly available

TRF2, M1BP and GFZF ChIP-exo data from *Drosophila* S2R+ cells (57, 58),

indicated that the effects of TRF2 knockdown on the expression of cyclin genes

correlate with TRF2, GFZF and M1BP co-occupancies of the promoters of the TRF2-

regulated cyclin genes (Table 4), in line with the reported regulatory effects of

GFZF. Genome-wide examination of promoters bound by all three transcription

factors revealed the enrichment of the TCT and Motif 1 core promoter elements.

Whereas one would expect the TCT to be enriched as TRF2 and M1BP have

previously been implicated in the regulation of ribosomal protein genes (32), the

enrichment of Motif 1 within promoters bound by all three factors indicates a shared

function for TRF2, GFZF and M1BP. Notably, in our experimental system, TRF2

knockdown by RNAi resulted in a two-fold reduction in *Trf2* levels (Fig. 2a, Additional

file 2: Table S1). Under these conditions, we did not detect any change in ribosomal

protein gene expression (Additional file 2: Table S1). Hence, the reduced expression

of *Cyc A*, *Cyc B* and *Cyc E* following a two-fold reduction in TRF2 expression, does

not result from a general inhibition of protein synthesis. As direct binding of TRF2 to

DNA could not be demonstrated (32), it is likely that TRF2 indirectly regulates the

expression of these cyclin genes and that there are TRF2-associated factors that

enable DNA binding. Our analysis suggested that GFZF and M1BP could serve as

such factors for specific TRF2-regulated processes. Interestingly, the expression

patterns of the cyclin genes following the knockdown of either TRF2 alone, or in

combination with GFZF and/or M1BP, indicated that among these three factors,

TRF2 is the major contributor to the expression pattern of *Cyc A*, *Cyc B* and *Cyc E*.

Moreover, while *Cyclin A*, *Cyclin B* and *Cyclin E* expression levels are reduced

following TRF2 (but not GFZF) knockdown, the expression levels of both *Cyclin A*

and *Cyclin B* , but not *Cyclin E*, are reduced following the combined knockdown of

TRF2 and GFZF (Fig. 7d, e), suggesting that the reduced expression levels of *Cyclin*

A and *Cyclin B* are not mediated via *Cyclin E*.

Notably, mouse TBP was recently shown to remain bound to mitotic

chromosomes during mitosis of mouse embryonic stem cells (mESCs), and to recruit

a small population of Pol II molecules to mitotic chromosomes (72). Nevertheless,

active Pol II transcription occurs in the absence of mouse TBP, whereas Pol I and

Pol III, are significantly reduced (11, 72). It remains to be determined whether

Drosophila TBP is bound to mitotic chromosomes during mitosis. As we did not

observe significant effects on cell cycle progression or mitosis following *Drosophila*

TBP knockdown (Figs. 3, 4 and 6), it is likely that *mouse* TBP may exert different

functions as compared to *Drosophila* TBP, perhaps via its associated proteins.

The effects of TRF2 knockdown on cell cycle progression and apoptosis of

Drosophila cultured S2R+ cells are in line with the early embryonic lethality of TRF2

knockout flies. TRF2 has also been shown to be essential for embryonic

development of *C. elegans*, zebrafish and *Xenopus*. It remains to be determined

whether knockdown of TRF2 in cellular systems from these species results in similar

effects.

CONCLUSIONS

Taken together, using *Drosophila* cells as a model system, we discovered that the

knockdown of TRF2, rather than TBP or TRF1, regulates apoptosis and cell cycle

progression via distinct target genes. Importantly, we discovered that TRF2 is

associated with the GFZF and M1BP proteins, and that TRF2, GFZF and M1BP co-

occupy the promoters of the TRF2-regulated cyclins. Furthermore, we show that

TRF2, rather than GFZF or M1BP, is the major contributor to the expression pattern

of *Cyc E*, *Cyc A* and *Cyc B*. Importantly, while a general transcription factor may be

regarded as having a "generic function", our findings emphasize the unique,

unanticipated functions of *Drosophila* TRF2 as an essential factor for specific major

cellular processes.

MATERIALS AND METHODS

Drosophila melanogaster Schneider S2R+ Cells

Drosophila melanogaster Schneider S2R+ adherent cells were cultured in

Schneider's *Drosophila* Media (Biological Industries) that was supplemented with

10% heat-inactivated FBS and Penicillin 100 units/ml Streptomycin 0.1mg/ml

(Biological Industries).

Generation of dsRNA probes

All dsRNA probes were chosen based on http://www.dkfz.de/signaling/e-rnai3// and

http://www.flyrnai.org/snapdragon as described in (69). Primer sequences used for

the generation of dsRNA probes are provided in Table 1. DNA fragments

corresponding to each dsRNA were subcloned into both pBlueScript SK+ and KS+.

The dsRNA probes were generated by PCR amplification of the DNA using T7 and

T3 primers, followed by *in vitro* transcription of templates in both pBlueScript

orientations using T7 RNA polymerase. Resulting RNA products were annealed to

generate the dsRNA probes.

RNA interference (RNAi)

 For 6 well plate, 1.25×10^6 cells/well were resuspended and seeded in empty

Schneider's *Drosophila* Media (Biological Industries) with 30μg/ml dsRNA directed

against different genes for 1 hour. Next, two volumes of complete medium were

added to the wells and cells were incubated for 3 additional days.

Western blot analysis

Knockdown of TRF2 and TBP was verified by western blot analysis using anti-TRF2

and anti-TBP polyclonal antibodies (generous gift from Jim Kadonaga). Cyclin E

levels were analyzed by the 8B10 antibodies (generous gift from Helena Richardson)

[73]. The levels of Actin or γ-Tubulin, as a loading control, were detected using either

mouse monoclonal anti-Actin (Abcam, 8224) or anti- γ-Tubulin (Sigma, GTU-88)

antibodies. Anti-Cyclin A and -Cyclin B concentrated monoclonal antibodies

(Developmental Studies Hybridoma Bank, A12 and F2F4, respectively) were tested

as well, however no endogenous proteins were detected, possibly due to technical

limitations.

TBP expression vector

The coding sequence of *Drosophila* TBP was amplified by PCR and cloned with an

N-terminal Flag-HA tag into the pAc5.1 vector (Life Technologies) using cDNA from

S2R+ cells as template and the following primers: Forward (containing an XhoI site,

underlined)

5' CCG<u>CTCGAG</u>GACCAAATGCTAAGCCCCA 3' and reverse (containing an AgeI

site, underlined) 5' AGC<u>ACCGGT</u>TTATGACTGCTTCTTGAACTTCTTTAA 3'

Plasmid sequence was verified by sequencing.

RNAi-coupled overexpression

For 6 well plate, 1.25×10^6 *Drosophila* S2R+ cells/well were resuspended and seeded

in empty medium with 30µg/ml dsRNA directed against TRF2 for 1 hour. Next, two

volumes of complete medium were added to the wells and cells were incubated for 3

days. Three days post dsRNA treatment, cells were transfected with the TBP-pAc

expression vector (930 ng) or an empty vector control using the Escort IV reagent

(Sigma). Media was replaced 18-24 hrs post transfection. Cells were harvested 36-

48 hrs post transfection and RNA was purified and analyzed by RT-qPCR. Each

qPCR experiment was performed in triplicates. The graphs represent an average of

3 independent experiments. Error bars represent SEM.

RNA-seq analysis

S2R+ cells were treated with dsRNA probes against *Trf2* (probe #1) and *Tbp*, and

harvested at two time points - 48h and 72h post RNAi treatment. For each time point,

a matched mock control was collected separately.

RNA was extracted using Quick-RNA™ MiniPrep (Zymo Research), and 800ng of

each sample was purified using NEBNext Poly(A) mRNA Magnetic Isolation Module

(NEB #E7490). Libraries were prepared using NEBNext Ultra II RNA Library Prep Kit

for Illumina (NEB #E7770), following the manufacturer's instructions. NEBNext

Multiplex Oligos for Illumina (NEB #E7335, NEB #E7500, NEB #E7710, NEB

#E7730) were used. Libraries were pooled and a 1% PhiX library control was added.

Single-end sequencing was performed on an Illumina NextSeq 500 machine.

Reads were aligned to dm6 genome build using STAR (version 2.6.0a), and htseq-

count (version 0.5.1p3, (74)) was used to count the reads mapped to each gene.

Differential expression analysis of conditions was performed using the DESeq2 R

package (75). Only genes with adjusted p-value < 0.1 were considered for

subsequent analysis. GO terms analysis was carried out using STRING v11 (76).

For all experiments, three independent biological replicates were compared and

merged for subsequent analysis. RNA-seq Data is available at GSE133685.

RT-PCR

Total RNA was isolated using the PerfectPure RNA Cultured Cell kit (5 PRIME) or

Quick-RNA™ MiniPrep (Zymo Research). One microgram of the total RNA was

reverse-transcribed into cDNA with M-MLV (Promega) or qScript Flex cDNA Kit

(Quanta). Control reactions lacking reverse transcriptase were also performed to

ensure that the levels of contaminating genomic DNA were negligible. Quantitation

was performed by real-time PCR to determine the transcription levels of the

endogenous genes. The expression levels were compared to *Gapdh2*. Primer

sequences for real-time PCR are provided in Table 2. For all quantifications, the

error bars represent ±S.E.M of at least 3 independent experiments; NS, not

significant; **$P < 0.01$; ***$P < 0.001$. Statistical analyses were performed on log-

transformed relative quantification (RQ) values using one-way ANOVA followed by

Tukey's post hoc test, unless otherwise stated in the figure legend.

Flow cytometry analysis

For cell cycle distribution by Propidium Iodide (PI) staining, cells were harvested

following 72h incubation with dsRNA, centrifuged for 5 min at 300g and fixed with

80% ethanol at 4ºC overnight. Before subjecting the cells to flow cytometry, the cells

were centrifuged for 5 min at 300g, washed in 1 ml of Phosphate-buffered saline

(PBS) and incubated for 40 min at 4°C. The cells were then stained in PBS

containing 50 µg/ml PI (Sigma) and 50 µg/ml RNase A (Roche). After incubation for

15 min at room temperature, fluorescence was measured using a FACSCalibur

Becton Dickinson flow cytometer.

For analyzing apoptosis by Annexin V and PI staining, cells were harvested 72h

following incubation with dsRNA probes and stained with Annexin V and PI

(MEBCYTO Apoptosis Kit; MBL). Fluorescence was measured using a FACSCalibur

Becton Dickinson flow cytometer.

For G1 phase cell arrest by hydroxyurea (HU) and BrdU (5-Bromo-2´-Deoxyuridine)-

PI staining, 72h following incubation with dsRNA, the medium was replaced with

medium containing a final concentration of 1mM HU for 18h (77) (for control cells,

the medium was replaced with a fresh medium). BrdU (40µM final concentration)

was added to the medium for 2h. Cells were released from HU by medium

replacement and, at 0, 2, 4, 6 or 8 hours following the release, cells were harvested,

centrifuged and fixed with 80% ethanol at 4ºC overnight. Following fixation, cells

were stained with FITC *(Fluorescein isothiocyanate)*-conjugated anti-Brdu antibodies

(BD) and PI according to the provided protocol, and fluorescence was measured

using BD FACSARIA III. All flow cytometry data was analyzed using the FlowJo

software. Statistical analyses of flow cytometry and imaging flow cytometry data

were performed in SPSS using two-tailed Students t-test. The number of times each

experiment was repeated, is detailed in the figure legends.

It is of note that unfortunately, we were unable to synchronize cells in G2/M in

a reversible manner using either Nocodazole or Colchicine, which cause

microtubules depolymerization. Specifically, Nocadazole did not arrest the S2R+

cells in G2/M, while Colchicine, which has been used since the 1950s to inhibit

mitotic progression, did cause enrichment of mitotic cells (Fig. 6b, d). However, this

effect was irreversible (data not shown). Thus, Colchicine could not be used for

G2/M block-release synchronization experiments.

Immunostaining for fluorescence microscopy and Imaging flow cytometry

analysis

Cells were RNAi-treated as described above. On day 4, 2ml of fresh medium was

added to the wells. To enrich for G2/M, as a control, Colchicine (Sigma) was added

to a final concentration of 350ng/ml. On the following day, the cells were harvested

and fixed with 4% Paraformaldehyde (PFA)/PBS (30 min), washed in PBST (PBS

containing 0.5% Triton x), blocked with PBS containing 1% BSA and 1% serum (1h),

and incubated with phospho-Histone H3 (Ser10) antibody (1:200, Cell Signaling

Technology #9701) for 1h at RT, followed by overnight at 4°C. Cells were washed,

stained with the secondary antibody (1:500, DyLight 488, ab96883), and then

counter-stained with 10µg/ml Hoechst 33342 (Sigma). For microscopy analysis,

samples were also stained for filamentous Actin with 3.5µM Acti-stain 670 Phalloidin

(Cytoskeleton, Inc. Cat. # PHDN1). Following staining, samples were subjected to

imaging flow cytometry, confocal microscopy or wide-field fluorescence microscopy

analysis.

Microscope image analyses

High resolution images were acquired using a Leica SP8 confocal microscope, and

high-throughput images for quantitative analysis were acquired using a Leica DMi8

microscope. Three separate experiments were performed and captured at 20x

magnification. For each treatment, approximately 275 frames were acquired and

analysed. The total number of Alexa 488 anti-phosphor-Histone H3 (Ser10) (PH3)

labeled cells, and the total number of Hoechst stained cells, were calculated using

the Fiji distribution of ImageJ.

Analysis workflow:

1. The raw PH3 channel images were enhanced using brightness and contrast, and

then the background was subtracted by reducing the Gaussian blurred filtered

image of the enhanced image. Next, a median filter was applied to smoothen the

image and an Otsu threshold was applied to get the binary image of all mitotic

nuclei. Finally, watershed was implemented to separate touching nuclei. Mitotic

cells were counted, eliminating small debris and noise.

2. To analyze the Hoechst channel, the background was subtracted by reducing the

Gaussian blurred filtered image of the original image. Next, a Moments threshold

was applied to get the binary image of total nuclei. Finally, watershed was

implemented to separate touching nuclei. Nuclei were counted while eliminating

small debris and noise.

The ratio between the number of mitotic cells and the total number of cells yields the

mitotic index for each treatment.

All manipulations in the images were made evenly across the entire field.

The Fiji macros will be shared upon request.

Multispectral imaging flow-cytometry (IFC) analysis

Cells were imaged using multispectral imaging flow cytometry (ImageStreamX mark

II imaging flow-cytometer; Amnis Corp, Seattle, WA, Part of EMD Millipore). Each

experiment was performed 3 times. In each experiment, at least 40,000 cells were

collected from each sample, and data were analyzed using the image analysis

software (IDEAS 6.2; Amnis Corp). Images were compensated for fluorescent dye

overlap by using single-stain controls. Imaging flow cytometry results were analyzed

by calculation of a set of parameters, termed "features", performed on a defined area

of interest, termed "mask". The serial gating strategy to identify the mitotic cell

population was as follows: Single cells were first gated using the area and aspect

ratio features on the bright-field (BF) image (the aspect ratio, which indicates how

round or oblong an object is, is calculated by division of the minor axis by the major

axis). Uncropped cells were gated using the centroid X (the number of pixels in the

horizontal axis from the upper left corner of the image to the center of the mask) and

area features. Focused cells were gated using the Gradient RMS feature, as

previously described (78) (Additional file 5: Figure S4a-c). Following this standard

gating series, the mitotic cell fraction of the entire cell population was identified using

the staining intensity for PH3 AF488 (channel 2), and mitotic cells were gated as the

high intensity population of PH3 staining within all focused cells (Additional file 5:

Figure S4d,e).

For a more complex analysis, we performed a second gating series. Focused cells

were first gated for G2/M based on DNA (Hoechst) intensity (Additional file 5: Figure

S4) and then gated for mitotic cells, as previously, by high PH3 intensity. To further

subdivide into the specific cell division phases, we gated according to nuclear

morphology based on the spot count and aspect ratio intensity features (see

Additional file 5: Figure S4f, g for detailed masking and gating). As it was previously

shown that serine 10 of histone H3 becomes dephosphorylated during telophase

(79), the telophase population was derived from the negative PH3-stained cells,

based on the BF circularity feature and DNA aspect ratio intensity (see Additional file

5: Figure S4h for detailed masking and gating). Full details of all masking, features

and analysis strategies are included in the legend of Additional file 5: Figure S4.

Identification of unique TRF2-interacting proteins

To identify the proteins that are in complex with TRF2 (the evolutionarily conserved

short TRF2), we used inducible FLAG-HA-tagged TRF2 S2R+ cells (69). As

controls, we used inducible S2R+ for FLAG-HA-long TRF2 (69) or FLAG-HA-TBP

(generated as in (69)). Cells were either induced by copper sulfate or left untreated.

Protein extracts were prepared and TRF2-containing complexes were immuno-

precipitated using anti-FLAG M2 affinity gel (Sigma). Following the IP, TRF2-

containing complexes were released with a FLAG peptide (Sigma). Samples were

resolved by SDS-PAGE. Proteins that were purified from TRF2-induced cells were

separated to two samples: proteins larger or smaller than 40 kDa. Samples were

subjected to Mass spectrometry analyses (The Smoler Protein Research Center,

Technion). Briefly, samples were digested by trypsin, analyzed by LC-MS/MS on Q-

Exactive Plus (ThermoFisher) and identified by the Discoverer software (with two

search algorithms: Sequest (ThermoFisher) and Mascot (Matrix science) against the

Drosophila melanogaster section of the NCBI non-redundant and Uniprot databases,

and a decoy database (in order to determine the false discovery rate). All the

identified peptides were filtered with high confidence, top rank, mass accuracy, and a

minimum of 2 peptides. High confidence peptides have passed the 1% FDR

threshold. Semi-quantitation was done by calculating the peak area of each peptide.

The area of the protein is the average of the three most intense peptides from each

protein. The results are provided in Additional file 8: Table S3.

Visualization of publicly available ChIP-exo data

A genome browser session

(https://genome.ucsc.edu/s/Anna%20Sloutskin/dm3_ChIP_Exo) based on available

TRF2, GFZF and M1BP ChIP-exo bedgraph files (GSE97841, GSE105009) (57, 58)

was created. The session contains an "Overlap" track (the ChIP-exo peaks that were

identified as overlapping in the ±50bp window relative to FlyBase TSS) and the

"trustedTSS" track that is based on 5' GRO-seq (GSE68677) and PRO-Cap

(GSM1032759) data. The relevant interval (±50bp or 100bp) is indicated.

List of abbreviations

ChIP-exo - chromatin immunoprecipitation combined with exonuclease digestion

followed by high-throughput sequencing

DPE - downstream core promoter element

GFZF - GST-containing FLYWCH zinc-finger protein

M1BP - Motif 1 binding protein

Pol II - RNA polymerase II

TBP - TATA-box-binding protein

TRF2 - TBP-related factor 2

TSS - Transcription start site

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

RNAseq data generated during the current study is available in the GEO repository

(GSE133685).

ChIP-exo data analyzed during the current study was downloaded from the GEO

(GSE97841, GSE105009).

Competing interests

The authors declare that they have no competing interests

Funding

This work was supported by grants from the Israel Science Foundation to T.J.-G.

(no. 798/10 and no. 1234/17).

Authors' contributions

A. Kedmi prepared RNA samples and O. Yaron and A. Sloutskin performed RNA-

seq experiments. T. Doniger and A. Sloutskin analyzed the RNA-seq experiments

and performed bioinformatics analysis. A. Kedmi, N. Epstein, L. Gasri-Plotnitsky and

D. Ickowicz performed and/or analyzed flow cytometry experiments. A. Kedmi, A.

Sloutskin and D. Ideses performed reverse transcription-qPCR analysis. I. Shoval

and Z. Porat advised and performed the analysis of ImageStream® experiments, A.

Kedmi and I. Shoval performed and analyzed fluorescence microscopy experiments.

E. Darmon and D. Ideses performed western blot analyses. A. Sloutskin performed

the statistical analysis. A. Kedmi, A. Sloutskin and T. Juven-Gershon designed the

study, planned experiments, analyzed results, and wrote the manuscript with input

from all authors.

Acknowledgements

We thank Jim Kadonaga for the generous gift of anti-TRF2 and anti-TBP antibodies,

for invaluable advice and assistance and for critical reading of the manuscript. We

thank Helena Richardson for the generous gift of anti-Cyclin E antibodies. We thank

Moshe Oren, Doron Ginsberg, Ze'ev Paroush, Sivan Henis-Korenblit, Shaked Cohen

and Itay Lazar for invaluable advice and assistance. We thank Dr. Jennifer Israel-

Cohen for assistance with statistical analysis. We thank Doron Ginsberg, Yaron

Shav-Tal, Galit Shohat-Ophir and Orit Adato for critical reading of the manuscript.

We thank Orit Adato and Sarit Lampert for technical assistance.

REFERENCES

1. Danino YM, Even D, Ideses D, Juven-Gershon T. The core promoter: At the heart of gene expression. Biochimica et biophysica acta. 2015;1849(8):1116-31.
2. Goodrich JA, Tjian R. Unexpected roles for core promoter recognition factors in cell-type-specific transcription and gene regulation. Nature reviews Genetics. 2010;11(8):549-58.
3. Muller F, Tora L. Chromatin and DNA sequences in defining promoters for transcription initiation. Biochimica et biophysica acta. 2014;1839(3):118-28.
4. Thomas MC, Chiang CM. The general transcription machinery and general cofactors. Crit Rev Biochem Mol Biol. 2006;41(3):105-78.
5. Dikstein R. The unexpected traits associated with core promoter elements. Transcription. 2011;2(5):201-6.
6. Haberle V, Stark A. Eukaryotic core promoters and the functional basis of transcription initiation. Nature reviews Molecular cell biology. 2018;19(10):621-37.
7. Kadonaga JT. Perspectives on the RNA polymerase II core promoter. Wiley Interdiscip Rev Dev Biol. 2012;1(1):40-51.
8. Lenhard B, Sandelin A, Carninci P. Metazoan promoters: emerging characteristics and insights into transcriptional regulation. Nature reviews Genetics. 2012;13(4):233-45.
9. Ohler U, Wassarman DA. Promoting developmental transcription. Development. 2010;137(1):15-26.
10. Vo Ngoc L, Wang YL, Kassavetis GA, Kadonaga JT. The punctilious RNA polymerase II core promoter. Genes & development. 2017;31(13):1289-301.
11. Martianov I, Viville S, Davidson I. RNA polymerase II transcription in murine cells lacking the TATA binding protein. Science. 2002;298(5595):1036-9.
12. Akhtar W, Veenstra GJ. TBP-related factors: a paradigm of diversity in transcription initiation. Cell & bioscience. 2011;1(1):23.
13. Muller F, Zaucker A, Tora L. Developmental regulation of transcription initiation: more than just changing the actors. Current opinion in genetics & development. 2010;20(5):533-40.
14. Reina JH, Hernandez N. On a roll for new TRF targets. Genes & development. 2007;21(22):2855-60.
15. Torres-Padilla ME, Tora L. TBP homologues in embryo transcription: who does what? EMBO reports. 2007;8(11):1016-8.
16. Zehavi Y, Kedmi A, Ideses D, Juven-Gershon T. TRF2: TRansForming the view of general transcription factors. Transcription. 2015;0.
17. Hansen SK, Takada S, Jacobson RH, Lis JT, Tjian R. Transcription properties of a cell type-specific TATA-binding protein, TRF. Cell. 1997;91(1):71-83.
18. Dantonel JC, Wurtz JM, Poch O, Moras D, Tora L. The TBP-like factor: an alternative transcription factor in metazoa? Trends in biochemical sciences. 1999;24(9):335-9.
19. Duttke SH, Doolittle RF, Wang YL, Kadonaga JT. TRF2 and the evolution of the bilateria. Genes & development. 2014;28(19):2071-6.
20. Moore PA, Ozer J, Salunek M, Jan G, Zerby D, Campbell S, et al. A human TATA binding protein-related protein with altered DNA binding specificity inhibits transcription from multiple promoters and activators. Molecular and cellular biology. 1999;19(11):7610-20.

21. Persengiev SP, Zhu X, Dixit BL, Maston GA, Kittler EL, Green MR. TRF3, a
TATA-box-binding protein-related factor, is vertebrate-specific and widely
expressed. Proceedings of the National Academy of Sciences of the United
States of America. 2003;100(25):14887-91.
22. Rabenstein MD, Zhou S, Lis JT, Tjian R. TATA box-binding protein (TBP)-related
factor 2 (TRF2), a third member of the TBP family. Proceedings of the National
Academy of Sciences of the United States of America. 1999;96(9):4791-6.
23. Kopytova DV, Krasnov AN, Kopantceva MR, Nabirochkina EN, Nikolenko JV,
Maksimenko O, et al. Two isoforms of Drosophila TRF2 are involved in
embryonic development, premeiotic chromatin condensation, and proper
differentiation of germ cells of both sexes. Molecular and cellular biology.
2006;26(20):7492-505.
24. Bashirullah A, Lam G, Yin VP, Thummel CS. dTrf2 is required for transcriptional
and developmental responses to ecdysone during Drosophila metamorphosis.
Developmental dynamics : an official publication of the American Association of
Anatomists. 2007;236(11):3173-9.
25. Dantonel JC, Quintin S, Lakatos L, Labouesse M, Tora L. TBP-like factor is
required for embryonic RNA polymerase II transcription in C. elegans. Molecular
cell. 2000;6(3):715-22.
26. Jacobi UG, Akkers RC, Pierson ES, Weeks DL, Dagle JM, Veenstra GJ. TBP
paralogs accommodate metazoan- and vertebrate-specific developmental gene
regulation. The EMBO journal. 2007;26(17):3900-9.
27. Kaltenbach L, Horner MA, Rothman JH, Mango SE. The TBP-like factor CeTLF
is required to activate RNA polymerase II transcription during C. elegans
embryogenesis. Molecular cell. 2000;6(3):705-13.
28. Levine M, Cattoglio C, Tjian R. Looping back to leap forward: transcription enters
a new era. Cell. 2014;157(1):13-25.
29. Muller F, Lakatos L, Dantonel J, Strahle U, Tora L. TBP is not universally
required for zygotic RNA polymerase II transcription in zebrafish. Current biology
: CB. 2001;11(4):282-7.
30. Neves A, Eisenman RN. Distinct gene-selective roles for a network of core
promoter factors in Drosophila neural stem cell identity. Biology open. 2019;8(4).
31. Veenstra GJ, Weeks DL, Wolffe AP. Distinct roles for TBP and TBP-like factor in
early embryonic gene transcription in Xenopus. Science. 2000;290(5500):2312-
5.
32. Wang YL, Duttke SH, Chen K, Johnston J, Kassavetis GA, Zeitlinger J, et al.
TRF2, but not TBP, mediates the transcription of ribosomal protein genes.
Genes & development. 2014;28(14):1550-5.
33. Martianov I, Fimia GM, Dierich A, Parvinen M, Sassone-Corsi P, Davidson I.
Late arrest of spermiogenesis and germ cell apoptosis in mice lacking the TBP-
like TLF/TRF2 gene. Molecular cell. 2001;7(3):509-15.
34. Zhang D, Penttila TL, Morris PL, Roeder RG. Cell- and stage-specific high-level
expression of TBP-related factor 2 (TRF2) during mouse spermatogenesis.
Mechanisms of development. 2001;106(1-2):203-5.
35. Zhang D, Penttila TL, Morris PL, Teichmann M, Roeder RG. Spermiogenesis
deficiency in mice lacking the Trf2 gene. Science. 2001;292(5519):1153-5.
36. Deveraux QL, Reed JC. IAP family proteins--suppressors of apoptosis. Genes &
development. 1999;13(3):239-52.

37. Fuchs Y, Steller H. Live to die another way: modes of programmed cell death
and the signals emanating from dying cells. Nature reviews Molecular cell
biology. 2015;16(6):329-44.
38. Martin SJ. Destabilizing influences in apoptosis: sowing the seeds of IAP
destruction. Cell. 2002;109(7):793-6.
39. Scuderi A, Simin K, Kazuko SG, Metherall JE, Letsou A. scylla and charybde,
homologues of the human apoptotic gene RTP801, are required for head
involution in Drosophila. Dev Biol. 2006;291(1):110-22.
40. Quinn L, Coombe M, Mills K, Daish T, Colussi P, Kumar S, et al. Buffy, a
Drosophila Bcl-2 protein, has anti-apoptotic and cell cycle inhibitory functions.
The EMBO journal. 2003;22(14):3568-79.
41. Brachmann CB, Jassim OW, Wachsmuth BD, Cagan RL. The Drosophila bcl-2
family member dBorg-1 functions in the apoptotic response to UV-irradiation.
Current biology : CB. 2000;10(9):547-50.
42. Colussi PA, Quinn LM, Huang DC, Coombe M, Read SH, Richardson H, et al.
Debcl, a proapoptotic Bcl-2 homologue, is a component of the Drosophila
melanogaster cell death machinery. The Journal of cell biology.
2000;148(4):703-14.
43. Igaki T, Kanuka H, Inohara N, Sawamoto K, Nunez G, Okano H, et al. Drob-1, a
Drosophila member of the Bcl-2/CED-9 family that promotes cell death.
Proceedings of the National Academy of Sciences of the United States of
America. 2000;97(2):662-7.
44. Zhang H, Huang Q, Ke N, Matsuyama S, Hammock B, Godzik A, et al.
Drosophila pro-apoptotic Bcl-2/Bax homologue reveals evolutionary conservation
of cell death mechanisms. The Journal of biological chemistry.
2000;275(35):27303-6.
45. Levine AJ, Oren M. The first 30 years of p53: growing ever more complex. Nat
Rev Cancer. 2009;9(10):749-58.
46. Vousden KH, Lane DP. p53 in health and disease. Nature reviews Molecular cell
biology. 2007;8(4):275-83.
47. Brodsky MH, Nordstrom W, Tsang G, Kwan E, Rubin GM, Abrams JM.
Drosophila p53 binds a damage response element at the reaper locus. Cell.
2000;101(1):103-13.
48. Fan Y, Lee TV, Xu D, Chen Z, Lamblin AF, Steller H, et al. Dual roles of
Drosophila p53 in cell death and cell differentiation. Cell Death Differ.
2010;17(6):912-21.
49. Teichmann M, Wang Z, Martinez E, Tjernberg A, Zhang D, Vollmer F, et al.
Human TATA-binding protein-related factor-2 (hTRF2) stably associates with
hTFIIA in HeLa cells. Proceedings of the National Academy of Sciences of the
United States of America. 1999;96(24):13720-5.
50. Baldin V, Lukas J, Marcote MJ, Pagano M, Draetta G. Cyclin D1 is a nuclear
protein required for cell cycle progression in G1. Genes & development.
1993;7(5):812-21.
51. Matsushime H, Roussel MF, Ashmun RA, Sherr CJ. Colony-stimulating factor 1
regulates novel cyclins during the G1 phase of the cell cycle. Cell.
1991;65(4):701-13.
52. Dulic V, Lees E, Reed SI. Association of human cyclin E with a periodic G1-S
phase protein kinase. Science. 1992;257(5078):1958-61.

53. Koff A, Giordano A, Desai D, Yamashita K, Harper JW, Elledge S, et al.
Formation and activation of a cyclin E-cdk2 complex during the G1 phase of the
human cell cycle. Science. 1992;257(5077):1689-94.

54. Taylor EW. The Mechanism of Colchicine Inhibition of Mitosis. I. Kinetics of
Inhibition and the Binding of H3-Colchicine. The Journal of cell biology.
1965;25:SUPPL:145-60.

55. Hervás JP, Fernández-Gómez ME, Giménez-Martín G. Colchicine Effect on the
Mitotic Spindle: Estimate of Multipolar Anaphase Production. Caryologia.
1974;27(3):359-68.

56. Rieder CL, Palazzo RE. Colcemid and the mitotic cycle. Journal of cell science.
1992;102 (Pt 3):387-92.

57. Baumann DG, Gilmour DS. A sequence-specific core promoter-binding
transcription factor recruits TRF2 to coordinately transcribe ribosomal protein
genes. Nucleic acids research. 2017;45(18):10481-91.

58. Baumann DG, Dai MS, Lu H, Gilmour DS. GFZF, a glutathione S-transferase
protein implicated in cell cycle regulation and hybrid inviability, is a transcriptional
co-activator. Molecular and cellular biology. 2017.

59. Bailey TL, Boden M, Buske FA, Frith M, Grant CE, Clementi L, et al. MEME
SUITE: tools for motif discovery and searching. Nucleic acids research.
2009;37(Web Server issue):W202-8.

60. Ohler U, Liao GC, Niemann H, Rubin GM. Computational analysis of core
promoters in the Drosophila genome. Genome Biol.
2002;3(12):RESEARCH0087.

61. Sloutskin A, Danino YM, Orenstein Y, Zehavi Y, Doniger T, Shamir R, et al.
ElemeNT: a computational tool for detecting core promoter elements.
Transcription. 2015;6(3):41-50.

62. Li J, Gilmour DS. Distinct mechanisms of transcriptional pausing orchestrated by
GAGA factor and M1BP, a novel transcription factor. The EMBO journal.
2013;32(13):1829-41.

63. Maeda R, Tamashiro H, Takano K, Takahashi H, Suzuki H, Saito S, et al. TBP-
like Protein (TLP) Disrupts the p53-MDM2 Interaction and Induces Long-lasting
p53 Activation. The Journal of biological chemistry. 2017;292(8):3201-12.

64. Akdemir F, Christich A, Sogame N, Chapo J, Abrams JM. p53 directs focused
genomic responses in Drosophila. Oncogene. 2007;26(36):5184-93.

65. Brodsky MH, Weinert BT, Tsang G, Rong YS, McGinnis NM, Golic KG, et al.
Drosophila melanogaster MNK/Chk2 and p53 regulate multiple DNA repair and
apoptotic pathways following DNA damage. Molecular and cellular biology.
2004;24(3):1219-31.

66. Zhang Y, Lin N, Carroll PM, Chan G, Guan B, Xiao H, et al. Epigenetic blocking
of an enhancer region controls irradiation-induced proapoptotic gene expression
in Drosophila embryos. Developmental cell. 2008;14(4):481-93.

67. Chakraborty R, Li Y, Zhou L, Golic KG. Corp Regulates P53 in Drosophila
melanogaster via a Negative Feedback Loop. PLoS Genet.
2015;11(7):e1005400.

68. Isogai Y, Keles S, Prestel M, Hochheimer A, Tjian R. Transcription of histone
gene cluster by differential core-promoter factors. Genes & development.
2007;21(22):2936-49.

69. Kedmi A, Zehavi Y, Glick Y, Orenstein Y, Ideses D, Wachtel C, et al. Drosophila
TRF2 is a preferential core promoter regulator. Genes & development.
2014;28(19):2163-74.

70. McCleland ML, O'Farrell PH. RNAi of mitotic cyclins in Drosophila uncouples the
nuclear and centrosome cycle. Current biology : CB. 2008;18(4):245-54.
71. Yuan K, O'Farrell PH. Cyclin B3 is a mitotic cyclin that promotes the metaphase-
anaphase transition. Current biology : CB. 2015;25(6):811-6.
72. Teves SS, An L, Bhargava-Shah A, Xie L, Darzacq X, Tjian R. A stable mode of
bookmarking by TBP recruits RNA polymerase II to mitotic chromosomes. eLife.
2018;7:e35621.
73. Richardson H, O'Keefe LV, Marty T, Saint R. Ectopic cyclin E expression
induces premature entry into S phase and disrupts pattern formation in the
Drosophila eye imaginal disc. Development. 1995;121(10):3371-9.
74. Anders S, Pyl PT, Huber W. HTSeq--a Python framework to work with high-
throughput sequencing data. Bioinformatics. 2015;31(2):166-9.
75. Love MI, Huber W, Anders S. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550.
76. Szklarczyk D, Gable AL, Lyon D, Junge A, Wyder S, Huerta-Cepas J, et al.
STRING v11: protein-protein association networks with increased coverage,
supporting functional discovery in genome-wide experimental datasets. Nucleic
acids research. 2019;47(D1):D607-D13.
77. Peng Q, Wang Y, Li M, Yuan D, Xu M, Li C, et al. cGMP-Dependent Protein
Kinase Encoded by foraging Regulates Motor Axon Guidance in Drosophila by
Suppressing Lola Function. J Neurosci. 2016;36(16):4635-46.
78. George TC, Fanning SL, Fitzgerald-Bocarsly P, Medeiros RB, Highfill S, Shimizu
Y, et al. Quantitative measurement of nuclear translocation events using
similarity analysis of multispectral cellular images obtained in flow. J Immunol
Methods. 2006;311(1-2):117-29.
79. Hendzel MJ, Wei Y, Mancini MA, Van Hooser A, Ranalli T, Brinkley BR, et al.
Mitosis-specific phosphorylation of histone H3 initiates primarily within
pericentromeric heterochromatin during G2 and spreads in an ordered fashion
coincident with mitotic chromosome condensation. Chromosoma.
1997;106(6):348-60.
80. Filby A, Perucha E, Summers H, Rees P, Chana P, Heck S, et al. An imaging
flow cytometric method for measuring cell division history and molecular
symmetry during mitosis. Cytometry A. 2011;79(7):496-506.
1003

Figure Legends

Fig. 1.

Trf2 Knockdown enhances early and late apoptosis in S2R+ cells. *Drosophila* S2R+

cells were incubated for three days with dsRNA directed against *Trf2*, *exd*, *Tbp* and

Trf1. To examine whether *Trf2* knockdown triggers apoptotic cell death, cells were

harvested following 72 h and stained with Annexin-V FITC and PI. **a** FACS analysis

of a representative experiment. **b** Average percentages of cells undergoing early and

late apoptosis. (n=4, *$0.01 < p \leq 0.05$, **$0.005 \leq p \leq 0.01$, ***$p < 0.005$, two-tailed

Students t-test; comparison to mock).

1013

Fig. 2.

The expression levels of distinct pro-apoptotic genes increase following knockdown

of TRF2. *Drosophila* S2R+ cells were incubated for three days with dsRNA directed

against TRF2, exd, TBP and TRF1. RNA was isolated from the cells and reverse

transcribed to cDNA. qPCR experiments were used to analyze the RNA levels of the

endogenous genes: **a** *Trf2*, *TBP, TRF1*, *rpr*, *hid*, *grim*, *scyl*, *Buffy*, *Debcl* and *Dronc*

and **b** *Trf2*, *p53* and *TBP.* qPCR experiments were performed in triplicates, and the

graph represents the average of three to eight experiments. Error bars represent the

SEM. *$p < 0.05$, one-way ANOVA followed by Tukey's post hoc test as compared to

the mock treatment of the relevant gene.

1024

Fig. 3.

Trf2, but not *Tbp* or *Trf1* knockdown, affects cell cycle distribution. *Drosophila* S2R+

cells were incubated for three days with dsRNA directed against *Trf2*, *Tbp*, *Trf1* and

exd. Cells were fixed with 80% ethanol and stained with Propidium-Iodide (PI) for

flow cytometry (FACS) analysis. **a** Cell cycle distribution histograms of a

representative experiment. **b** Average cell cycle distribution determined by FACS

analyses of eight independent experiments (*0.01 < p ≤ 0.05, **0.005 ≤ p ≤ 0.01, ***p

< 0.005, two-tailed Students t-test; comparison to mock).

1033

Fig. 4.

TRF2 is involved in S phase progression. *Drosophila* S2R+ cells were incubated for

three days with dsRNA probes directed against *Trf2* or *Tbp*. Next, cells were either

left untreated or treated with 1mM Hydroxyurea for 18h. The cells were allowed to

resume cell cycle for 2h, 4h, 6h or 8h in fresh medium containing 40µM 5-Bromo-2´-

deoxyuridine (BrdU), and were then fixed with 80% ethanol overnight and analyzed

by FACS using BrdU-PI staining. Each histogram plots the PI fluorescence intensity

(representing DNA content) on the X-axis, and cell count on the Y-axis.

1042

1043

Fig. 5.

Knock down of *Trf2* expression by RNAi reduces the expression of cyclin genes.

Drosophila S2R+ cells were incubated for three days with dsRNA probes directed

against *Trf2*, *exd*, *Tbp* and *Trf1*. RNA was isolated from the cells and reverse

transcribed (RT) to cDNA. Real-time PCR (qPCR) experiments were used to analyze

the RNA levels of the endogenous genes: **a** *Trf2*, *CycA*, *CycB*, *CycB3*, *CycC*, *CycD*

and **b** *CycE*. As there were differences in *CycE* expression following RNAi with

probe #1 compared to probe #2, four non-overlapping probes were used to

knockdown *Trf2* expression towards the analysis of *CycE* expression. qPCR

experiments were performed in triplicates, and the graph represents the average of

three to eight experiments. Error bars represent the SEM. *p< 0.05, one-way

ANOVA followed by Tukey's post hoc test as compared to the mock treatment of the

relevant gene. **c** Western blot analysis following TRF2 and TBP knockdown in S2R+

cells, using anti-TRF2 polyclonal antibodies and anti-Cyc E monoclonal antibodies.

Actin was used as a loading control.

1059

Fig. 6.

TRF2 regulates cell cycle progression through mitosis. *Drosophila* S2R+ cells were

incubated for three days with dsRNA directed against TRF2 or TBP. Cells were then

fixed with 4% PFA and stained with a phospho-Histone H3 (Ser10) antibody (PH3,

mitotic marker; green), Hoechst (DNA visualization; Blue) and Phalloidin (filamentous

Actin visualization; red). **a** Representative confocal microscopy images of *Drosophila*

S2R+ cells in different mitotic phases. **b** Comparison of mitotic indices following each

treatment, calculated based on microscopic analysis. Shown are the averages of

three independent experiments, in which a total of 100,000-800,000 cells were

analyzed for each treatment (*$0.01 < p \le 0.05$, **$0.005 \le p \le 0.01$, ***$p < 0.005$, two-

tailed Students t-test; comparison to mock). **c** Representative images obtained by

imaging flow cytometry analysis. **d** Comparison of mitotic indices following each

treatment, calculated based on imaging flow cytometry. Shown are the averages of

three independent experiments, in which a total of 40,000 cells were analyzed for

each treatment (*$0.01 < p \le 0.05$, **$0.005 \le p \le 0.01$, ***$p < 0.005$, two-tailed

Students t-test; comparison to mock). **e-f** Distribution of mitotic phases among all

mitotic cells, based on imaging flow cytometry. Shown are the averages of three

independent experiments, in which a total of 40,000 cells were analyzed for each

treatment (*$0.01 < p \le 0.05$, **$0.005 \le p \le 0.01$, ***$p < 0.005$, two-tailed Students t-

test; comparison to mock). Filled triangles indicate aberrant chromosomal

morphology in Colchicine-treated cells. Phospho-Histone H3 Ser10-positive cells (**e**),

were analyzed separately from the Phospho-Histone H3 Ser10-negative cells

undergoing mitosis (**f**). Phospho-Histone H3 Ser10-negative cells undergoing mitosis

(as identified by the imaging flow cytometer) were defined as cells in telophase (f).

Notably, cell counts of telophase cells likely include Phospho-Histone H3 Ser10-

negative doublet cells, which, even using the high-resolution Imagestream, could not

be distinguished from telophase cells.

1087
Fig. 7.

CycA and *CycB* expression is affected by TRF2 knockdown, and less by GFZF and

M1BP. **a** Schematic representation of genes containing at least one binding site of

the specified transcription factor at ±50bp relative to its TSS, as determined by

FlyBase. ChIP-exo data was retrieved from GSE97841 and GSE105009. **b** Top

enriched motif among the 4331 commonly bound promoters, as detected by MEME

analysis. Its resemblance to Ohler Motif 1 is depicted by the motif logo derived by

M1BP ChIP-exo (57). **c** The 4331 commonly bound regions were analyzed for core

promoter composition. This promoter group was found to be depleted for the TATA-

box motif and enriched for dTCT and the Motif 1 core promoter elements. p-values

were adjusted using Bonferroni correction. ***$p < 10^{-5}$. **d-e** *Drosophila* S2R+ cells

were incubated for three days with dsRNA probes directed against *Trf2* (probe #1),

gfzf, *M1BP* or their combinations. RNA was isolated from the cells and reverse

transcribed (RT) to cDNA. Real-time PCR (qPCR) experiments were used to analyze

the RNA levels of the endogenous of *Trf2*, *gfzf* and *M1BP*, as well as *CycA*, *CycB*,

CycD and *CycE* genes, as indicated. **d** Single knockdowns of *Trf2*, *gfzf* or *M1BP*. **e**

Knockdown of multiple genes, as indicated. qPCR experiments were performed in

triplicates, and the graph represents the average of 4 independent experiments.

Error bars represent the SEM. *p< 0.05, one-way ANOVA followed by Tukey's post

hoc test as compared to the mock treatment of the relevant gene.

1108

1109

Additional files

Additional file 1: Figure S1. .pdf

A) Schematic representation of dsRNA TRF2 probes used in this study. *B*) Western

blot analysis following TRF2 and *C)* TBP knockdown in S2R+ cells, using anti-TRF2

and anti-TBP polyclonal antibodies, respectively. Actin and γ-Tubulin were used as

loading controls.

Additional file 2: Table S1. .xlsx

GO terms analysis of the RNA-seq data. Summary, as well as the exact GO terms

are provided, according to the datasheet name. Analysis was performed sing

STRING. Only genes with pAdj <0.1 were considered. DEseq2 output is presented

for either *Trf2* or *Tbp* knockdown, as compared to mock at 72h post silencing.

1121

Additional file 3: Figure S2. .pdf

TBP overexpression does not result in induction of pro-apoptotic gene expression.

Drosophila S2R+ cells were depleted of *Trf2* by RNAi and then transfected with

either pAc-empty (mock) or TBP expression vector. Cells were harvested 36–48h

following transfection, RNA was purified and reverse transcribed to cDNA for RT-

qPCR analysis. qPCR experiments were performed in triplicates, and the graph

represents the average of 3 experiments. Error bars represent the SEM. *p< 0.05,

two-tailed Students t-test; comparison to the same treatment without TBP

overexpression.

1131

Additional file 4: Figure S3. .pdf

TRF2 is involved in S phase progression. *Drosophila* S2R+ cells were incubated for

three days with dsRNA probes directed against *Trf2* or *Tbp*. Next, cells were either

left untreated or treated with 1mM Hydroxyurea for 18h. The cells were allowed to

resume cell cycle for 2h, 4h, 6h or 8h in fresh medium containing 40µM BrdU, and

were then fixed with 80% ethanol overnight and analyzed by FACS using BrdU-PI

staining. The PI fluorescence intensity (representing DNA content) is plotted on the

X-axis (linear scale), and the BrdU-FITC fluorescence intensity (representing BrdU

incorporation into the DNA) is plotted on the Y-axis (log scale).

1141

Additional file 5: Figure S4. .pdf

Gating and masking strategy for imaging-flow cytometry data analysis. *A*) Cells were

gated for single cells, using the area and aspect ratio features on the BF image. *B*)

Centered and uncropped cells were gated based on the centroid X (the number of

pixels in the horizontal axis from the upper left corner of the image to the center of

the mask) and area features. *C*) Focused cells were gated, using the Gradient RMS

feature, as previously described (78). *D*) The G2/M population was gated out of the

focused cells, based on DNA (Hoechst) intensity. *E*) Mitotic cells were gated from the

G2M population, as the high intensity population of pH3 staining, based on the

intensity feature of pH3 and Max pixel feature of pH3 (the largest value of the

background-subtracted pixels contained in the input mask). *F*) To include only single

positive pH3 stained cells, doublet cells were eliminated by gating the single cells

from the mitotic population, using the area and aspect ratio features of the BF. *G*) To

discriminate between the different mitotic phases subpopulations, several masks

were created: 1. A morphology mask that includes all pixels within the outermost

image contour. 2. A threshold mask that includes the highest intensity pixels

(indicated as percentages). 3. A range mask that selects components in an image

within a selected size (μm), was used to eliminate small components.

Furthermore, the following features were used on the combined masks:

1. Spot count - the number of connected components in an image.

2. Aspect ratio intensity - the aspect ratio weighted for fluorescence intensity.

These masks were combined and the features were calculated and plotted as

follows:

Spot count of the combined mask: range (threshold 75%, M07) 20-5000, was plotted

against the aspect ratio intensity of the combined mask: range

(Threshold(Morphology(M07), 82%) 15-5000.

The prophase population was defined as having a more circular nuclear staining

(aspect ratio intensity should be high), and was hence gated as one nuclear spot

count with aspect ratio intensity bigger than 0.6. On the other hand, the metaphase

population was defined as having a more elongated DNA distribution and was gated

as one nuclear spot count with aspect ratio intensity smaller than 0.6. Finally, the

anaphase population was gated as cells with two nuclear spots having aspect ratio

intensity less than 0.6. (80). *H*) As serine 10 of histone H3 becomes

dephosphorylated during telophase, the telophase population was derived from the

negative pH3-stained cells. To identify telophase pairs, an object mask was created

(which segments images to closely identify the area corresponding to the cell). The

circularity feature (which measures the degree of the mask's deviation from a circle)

of the object mask (BF), was plotted against the aspect ratio intensity of the M07

DNA mask. Telophase cells were gated as having the lowest BF circularity and as

the most elongated, based on DNA stain (lowest aspect ratio intensity (M07)).

1182

1183   **Additional file 6: Table S2. .doc**

1184   Number of cells in each mitotic phase within phospho-Histone H3 Ser10-positive

1185   mitotic cells or telophase.

1186

Additional file 7: Figure S5. .pdf

Drosophila S2R+ cells treated with Colchicine display aberrant chromosomal

morphology. Colchicine-treated cells were fixed with 4% PFA and stained with a

phospho-Histone H3 (Ser10) antibody (green), Hoechst (DNA visualization; blue)

and Acti-stain 670 Phalloidin (filamentous Actin visualization; red). *A*) Shown are

representative images of cells in different mitotic phases obtained by imaging flow

cytometry analysis. *B*) Quantitation of cells with aberrant DNA morphology within

mock and colchicine-treated cells using imaging flow cytometry analysis. Prophase-

and metaphase-gated cells within either mock or colchicine-treated S2R+ cells, were

further gated using calculations of the following features:

1. The Delta centroid XY feature (which measures the distance in microns between

the centroid feature of two images using the user provided masks) was calculated

using the BF default mask and Hoechst channel mask of the 60% most highly

intense pixels. Cells with centered nucleus will get a lower value while polar located

nucleus will get a higher value. 2. The Max contour position feature (the location of

the contour in the cell that has the highest intensity concentration; the score is

between 0 to 1, with 0 being the object center and 1 the object perimeter). To

distinguish between central vs. polar location of the dividing nucleus, the Delta-

centroid XY was plotted against the Max contour position, and polar DNA was gated

as having the highest values of both features. Depicted are the analyses of cells in

prophase and metaphase. Cells in anaphase are not shown due to the low number

of cells observed in anaphase (Additional file 6).

1209

1210 **Additional file 8: Table S3. .xlsx**

1211 Mass-spectrometry data for short TRF2, long TRF2 and TBP proteins.

1212

Additional file 9: Figure S6. .pdf

Knockdown of *M1BP* specifically elevates the expression of *CycA* and ribosomal

protein genes. *Drosophila* S2R+ cells were incubated for three days with dsRNA

probes as indicated in the legend. RNA was isolated from the cells and reverse

transcribed (RT) to cDNA. Real-time PCR (qPCR) experiments were used to analyze

the RNA levels of the endogenous genes. qPCR experiments were performed in

triplicates, and the graph represents the average of 3 experiments ± SEM. *A) CycA*

expression levels are reduced following *Trf2* knockdown, but not *gfzf* or *M1BP*

knockdown. $*p < 0.05$, one-way ANOVA followed by Tukey's post hoc test as

compared to the mock treatment of the relevant gene. *B)* Additional ribosomal genes

are influenced by *M1BP* knockdown, however not all genes are upregulated in

response to *M1BP* knockdown. *CG12493* and *sgll* genes were previously shown to

be downregulated upon *M1BP* knockdown (62).$**p < 0.01$, two-tailed Students t-test;

compared to mock treatment.

1227

Table 1.

Primers for generation of dsRNA probes

dsRNA probe	Forward primer	Reverse primer
Trf2 #1	ATAGGTACCGGCAACCGGCAGTAAAAATA	ATAACTAGTACTCCACATTTGATCCCTGC
Trf2 #2	ATACTCGAGAACAGAAGGAGCAGCATCGT	ATAACTAGTTATTTTTACTGCCGGTTGCC
Trf2 #3	ATAGGTACCAAGGAGAACCAATCGCCGAAT	ATAACTAGTATTAGAAGAACTTAAGCGATC
Trf2 #4	ATACTCGAGCAATCTGACTTGAATCCCGG	ATAACTAGTTCATCTGAAGCTTGTCGCG
exd	ATAACTAGTTCGATGGTGCTGACAATGCC	ATAGGTACCGGGGGCTTAGATCCTGATGGAG
Trf1	GGGGTACCGGACAGGGATAATGTGGCTG	AAACTAGTGGCTTGACCATGCGATAGAT
Tbp	GGGGTACCACATGATGCCCATGAGTGA	AAACTAGTAATGGGGAATATCTTGTCGAAG
rpr	AAGGTACCACGAAAGAAAGTGTGTGCG	AAACTAGTTGCAATTTTTAGCCAACTTCG
scyl	AAGGTACCTACTACGCTGCTGACGAGGA	AAACTAGTATCACCATTAGTTGGTGGGCG
p53	AAGGTACCGATGCTGCAGGACATTCAGA	AAACTAGTCTCGGCTATCATTGCTCTCC
M1BP	GGGGTACCATATTAACACGAAACACCGGG	AAACTAGTACCTTGGTGTCGTCGATCTC
gfzf	GGGGTACCTCAGCATCTGTTCCACTTCG	AAACTAGTGTGTGTGAATGTGGGTCGAG

Table 2.

Real-time PCR (qPCR) primers

Gene	Forward primer	Reverse primer
Gapdh2	TTCCTCAGCGACACCCACTC	ATGACGCGGTTGGAGTAGCC
Trf2	GGAATCGTCTTCTGGGGACT	GACGACTCCTGTTGGCTTTG
CycA	TGGGCACGGCAGCTATGTAT	CCTGCGCCTTGGTGTAACTG
CycB	CGAGCACCATACGATGTCCA	TTGAGCAAGTGCAGCGACAG
CycB3	TCCCAGAGACTGCTCCAAGC	CATGGCGTAGTGGGACACCT
CycC	CACCGATGTCTGCCTGCTC	GCACGATCTCCTGGACCTTG
CycD	AGGTCGAGGAGAAGCACCAC	CCTCGGCACACACTTCCAT
CycE	CTCGGTTTTGAGCCTCCATC	AGACAACGGGCGAGGTGTAG
Tbp	TCAGCTCCGGCAAGATGGTG	GCAGGGAAACCGAGCTTTTGG
Trf1	AGAAGCTGGGATTCCCCGTA	GCACGTGGTTGAGGTTCTCC
exd	GCGAAATCAAGGAGAAGACCGTCC	CCTCGGCAATCAGCATGTTGTCC
rpr	CATACCCGATCAGGCGACTC	GTGTACTGGCGCAGGGTTTC
hid	CGACCTCCACGCCGTTATC	GCTCTGGTACTCGCGCTCAT
grim	TTTGGCCCAGATCTTCTGCT	GCATCAGTCACGTCGTCCTC
Debcl	ACAGCATGGGCGAGGAACT	ATGTCGCTGTCCTCCAGCTC
scyl	ATAATCCGCGTGTCGGAGAA	CCGTATCCGAATCGACCTTG
buffy	TTCTCAGGGTCGTTGCCTGT	TGGAGGTGGAGCCCAGTATG
p53	TTAGCGTTGAGCCTTTGACG	CAGGGGGACTACAACGGAAA
M1BP	AATTTGGCTGCGAACTCTGT	CAGCGGCCACAGTACTTACA
gfzf	GAACCCACCGGATATGTCAC	TGCTGGCAGGGTCTTAAGTT
RpLP2	GACATGGGCTTCGCTCTCTT	GTGAACGGATGGGTGCTACA
RpS12	CAAGCGTCAGGCTGTTCTGT	CAGCTTCTTGTGCGAGTCCA
CG12493	GACAACCAATTGGATCAGGAAAG	AGATTCACCATCGGGCATATT
sgll	ATTCGAAGGATGAGCCAAGG	CCAGTCTCGGAATACACAGAAG
RpL30	CAAATACTGCCTGGGCTACA	TACTCGATCTCGGACTTCCTC
RpLP1	CACTTCGACATGTCCACCAA	CCTTCAGGATGGTGTTGATCTT

Table 3.

Enrichment of TRF2, TBP, M1BP and GFZF in FLAG-immuno-affinity purified

complexes from inducible S2R+ cells

Name of purified (/co-purified) protein	Enrichment of purified (/co-purified) proteins within		
	Short TRF2-associated proteins	Long TRF2-associated proteins	TBP-associated proteins
TRF2	78.695	6.128	-
TBP	-	-	18.3
M1BP	1.795E7	-	-
GFZF	1.683E7	-	-

The values in the table represent the enrichment of each purified (/co-purified)

protein in the indicated sample. The enrichment was calculated as the ratio between

the mass spectrometry area (the average of the three most intense peptides from

each protein) of the induced sample and the un-induced sample.

Table 4.

TRF2, M1BP and GFZF occupancy within -100 to +100 relative to the TSSs

Gene name	TSS Location	TRF2			M1BP			GFZF		
		# of bound sites	Max. peak score	Avg. peak score	# of bound sites	Max. peak score	Avg. peak score	# of bound sites	Max. peak score	Avg. peak score
CycA_1	chr3L:11826617	1	54.16	54.16	59	126.93	30	2	112.85	87.78
CycA_2	chr3L:11826310	1	36.11	36.11	1	20.04	20.04	0	0	0
CycB_1	chr2R:18694432	15	150.45	38.11	47	106.89	26.72	3	114.64	62.1
CycB_2	chr2R:18694437	15	150.45	38.11	51	106.89	25.28	3	114.64	62.1
CycB_3	chr2R:18694449	15	150.45	38.11	60	106.89	27.28	3	114.64	62.1
CycB3	chr3R:20696533	4	90.27	40.62	128	140.29	36.69	9	157.63	72.05
CycE_1	chr2L:15746609	2	66.2	42.13	1	6.68	6.68	0	0	0
CycE_2	chr2L:15748123	4	60.18	31.6	8	53.44	30.9	1	136.14	136.14
CycC_1	chr3R:10715915	4	36.11	24.07	4	66.81	23.38	1	42.99	42.99
CycC_2	chr3R:10715922	4	36.11	24.07	3	66.81	26.72	1	42.99	42.99
CycD_1	chrX:15803691	2	42.13	24.08	7	46.76	20.04	0	0	0
CycD_2	chrX:15803682	2	42.13	24.08	7	46.76	20.04	0	0	0
RpL5	chr2L:22429377	74	210.63	51.56	142	140.29	33.07	29	148.68	48.36
RpL7	chr2L:10201108	23	210.63	50.50	121	374.11	62.44	5	118.23	65.92
RpL23	chr2R:18741912	38	132.4	46.25	200	327.35	51.27	68	449.61	99.00
RpL30	chr2L:19009229	21	102.31	30.38	141	233.82	47.81	16	231.08	76.58
RpLP1	chr2L:419957	14	156.47	44.28	148	180.38	37.78	19	195.25	74.10
RpLP2	chr2R:12473638	48	234.71	67.08	175	173.70	37.37	8	175.55	65.16

TRF2, M1BP and GFZF occupancy data (number of bound sites, the maximum peak scores and the average peak scores) was retrieved from ChIP-exo experiments performed in *Drosophila* S2R+ cells (GSE97841 and GSE105009) (64, 65). Due to variations in TSSs obtained by different methods, we relate to -100 to +100 relative to the TSSs peaks from available 5'GRO-seq and focused TSS analysis in S2 cells (GSE68677 and http://labs.biology.ucsd.edu/Kadonaga/drosophila.tss.data/) and PRO-cap analysis in S2 cells (GSM1032759).