Brain structures associated with reading and their abnormalities in dyslexia: a whole-brain analysis

Running title: Neuroanatomy of dyslexia

Kujala, T.[1], Thiede, A.[1], Palo-oja, P.[1], Virtala, P.[1], Laasonen, M.[2,3,4], Numminen, J.[5] & Sihvonen, A. J.[1,6]

[1]Cognitive Brain Research Unit, Department of Psychology and Logopedics, University of Helsinki, Finland.

[2]Department of Psychology and speech-language pathology, Speech and Language Pathology & Special psychologist education in child and adolescent psychology, University of Turku, Finland.

[3]Department of Psychology and Lopopedics, University of Helsinki, Finland.

[4]Department of Phoniatrics, Helsinki University Hospital, Finland.

[5]Department of Radiology, Töölö Hospital, Helsinki University Central Hospital, Finland.

[6]Department of Neurosciences, University of Helsinki, Finland.

Address correspondence to: Teija Kujala, Cognitive Brain Research Unit, Department of Psychology and Logopedics, P.O.Box 21, Haartmaninkatu 3 B, University of Helsinki, Finland.
E-mail: teija.m.kujala@helsinki.fi

Abstract

Developmental dyslexia (DD) is a highly prevalent neurodevelopmental disorder, which is devastating for individuals in modern societies in which fluent reading skill is mandatory for leading a normal life. Research on the neural origins of DD has continued for half a century, yielding, however, inconsistent results. It has also lacked a thorough characterization of the association between abnormal neuroanatomy and skills vital for reading. The current study was set out to determine abnormalities of grey and white matter volumes in adults with DD and associations between brain structures and reading and related skills. To this end, we conducted a whole-brain voxel based morphometry following current guidelines on state-of-the-art analysis approaches and rigorous neuropsychological testing. We found decreased volumes of grey matter in DD, comprising a left-hemispheric network including superior temporal and inferior frontal gyri, insula, the limbic system, and basal ganglia, and white matter, including the right middle temporal gyrus and hippocampus. These results are both consistent with the most robust previous findings on structural abnormalities in the left frontotemporal network in DD and yield novel insight to the role of subcortical structures in DD, scarcely studied so far. Moreover, areas with decreased grey matter volumes in DD were positively associated with technical reading scores (both groups included). This provides particularly strong support for the conclusion that the grey matter regions that we identified to have a low volume in DD comprise the core areas vital for reading.

Keywords: Developmental Dyslexia, Neuroanatomy, Voxel-based Morphometry, Grey Matter, White Matter

1. Introduction

Developmental dyslexia (DD) is a reading-skill impairment, which may emerge irrespective of adequate intelligence and reading instruction (Manual of Mental Disorders, 2000). The prevalence of dyslexia ranges between 5-17,5% (Shaywitz, 1998), which makes it the most common neurodevelopmental disorder. Due to its high prevalence and devastating influences on the individual's academic achievements, career, self-esteem, and coping in the modern society, it is pertinent to understand the neural basis of DD. Yet, this task is very challenging due to the heterogeneity of its phenotype (Zoubrinetzky et al., 2014; McArthur et al., 2013) and the complexity of the neural network underlying reading (Kujala et al., 2007).

According to functional imaging studies, the key neural network for reading comprises frontotemporoparietal circuits predominantly in the left hemisphere (e.g., Levy et al., 2009; Welcome and Joanisse, 2012). During word reading, the early visual processing occurs in the left inferior occipitotemporal (OT) cortex (Stoodley and Stein, 2013). The OT along with cerebellum are the starting points for the two major forward-driving nodes in the network of reading (Kujala J. et al., 2007). After the left OT areas, reading involves left parietal cortex and left inferior frontal gyrus (IFG; Levy et al., 2009). Pseudo-word reading relies more on the route using left OT than left parietal cortex, presumably due to the higher demand on phonological analysis when reading pseudo-words than familiar words (Levy et al., 2009).

The endeavor to find anomalies in the reading circuitry in DD to reveal its neural basis has continued for over 50 years (e.g., Drake, 1968; Ramus et al., 2018). However, the previous morphological studies, including those utilizing modern neuroimaging methods, have offered relatively few strong and replicated results on the neural basis of DD. Moreover, only few of

them have properly assessed and discussed the association between neuroanatomy and skills pertinent for reading.

The current study employed a rigorous neuropsychological testing of cognitive functions vital for reading and voxel-based morphometry (VBM), an automated MRI method for assessing focal brain changes (Ashburner and Friston, 2000). Since its introduction, VBM has become a standard method to analyze neuroanatomical abnormalities in various disorders, including DD with over 20 published studies (Brambati et al., 2004, Brown et al., 2001, Dole et al., 2013, Eckert et al., 2016, Eckert et al., 2005, Evans et al., 2014, Hoeft et al., 2007, Jednoróg et al., 2015, Jednoróg et al., 2014, Krafnick et al., 2014, Kronbichler et al., 2008, Menghini et al., 2008, Pernet et al., 2009a, Pernet et al., 2009b, Silani et al., 2005, Siok et al., 2008, Steinbrink et al., 2008, Tamboer et al., 2015, Vinckenbosch et al., 2005, Xia et al., 2016, Yang et al., 2016). Most frequently reported brain regions in VBM findings in DD include left posterior temporal and temporoparietal areas, but with both increased and decreased grey matter (GM) volume reported in DD (e.g., Brambati et al., 2004; Silani et al., 2005; Hoeft et al., 2007). In addition, DD has been associated with reduced GM (both modulated and non-modulated data) bilaterally in the frontal lobe (Brown et al., 2001) as well as in the left superior and inferior frontal gyrus (Brown et al., 2001; Brambati et al., 2004), bilateral OT regions (Brambati et al., 2004; Eckert et al., 2005; Kronbichler et al., 2008), subcortical structures (caudate and thalamus, Brown et al., 2001), and cerebellum (Brown et al., 2001; Brambati et al., 2004; Eckert et al., 2005; Kronbichler et al., 2008). In contrast, a number of studies with no significant morphological findings in DD have also been published (e.g., Pernet et al., 2009a, b; Casper et al., 2018).

Three recent meta-analyses have probed the heterogeneous morphometric findings in DD. In 2013, Richlan and colleagues evaluated nine VBM studies on DD and found reduced GM in

the superior temporal areas bilaterally in DD (Richlan et al., 2013), but with less stringent statistical thresholding. A few years later, Eckert and colleagues evaluated 11 VBM studies on DD, and reported lower GM volume (i.e. modulated data; adjusted for total GM volume) in left superior temporal sulcus, left orbitofrontal cortex, and the right cerebellum in participants with DD (Eckert et al., 2016). Interestingly, one of the most robust findings is reduced total brain volume in subjects with DD compared with typical readers, a finding confirmed by a meta-analysis including 1164 participants across 18 studies (Ramus et al., 2018). However, the direction of association between the reduced brain volume and DD, let alone its possible aetiological factor for DD remain currently unknown.

Overall, there is enormous variation between the results obtained on the structural brain anomalies in DD. As pointed out by Ramus and colleagues in their cogent review (Ramus et al., 2018), a few key problems render the majority of the results unreliable. First, a mixed selection of statistical thresholding and corrections for multiple tests has been applied, if any. When voxel-based neuroimaging methods (e.g. VBM) are used to reliably identify the neuroanatomical changes in DD, a combination of reasonable cluster-size threshold and voxel extent is needed to produce a desirable balance between Types I and II error rates (Lieberman and Cunningham, 2009). Second, most of the studies consist of small sample sizes, with only three studies including more than 16 dyslexics (see Eckert et al., 2016, for a meta-analysis). Another core problem is the bias for publishing studies reporting group differences. Due to these reasons, it is very hard to judge which findings are robust and reliable (see Ramus et al., 2018, for a review).

To attain reliable VBM results, it is critical how the preprocessing of MRI data is carried out, in other words, how the GM and white matter (WM) probabilistic maps are formed. As in any

voxel-based imaging analysis, following published guidelines, ensuring efficient registration, as well as controlling for factors that affect brain size (e.g. age) aid to achieve dependable results and extra sensitivity (Barnes et al., 2010; Ashburner and Friston, 2001; Pell et al., 2008; Li X et al., 2013; Da Ma et al., 2018). One crucial preprocessing step is modulation, which allows testing for regional differences in the absolute amounts (volume) of GM or WM (Ashburner and Friston, 2000; Good et al., 2001). Unfortunately, modulation has not been used consistently in morphological studies on DD (Brown et al., 2001; Vinckenbosch et al., 2005), but yet the results have been interpreted and discussed similarly whether the probabilistic maps were representing volume or not. Ramus et al. (2018) published a well-founded list of methodological recommendations for more reliable research on the neuroanatomy of DD and encouraged to systematically use relevant covariates to diminish their confounding effects on between-subject comparisons. In general, it is strongly recommended to include an adjustment for head size and other nuisance variables like gender and age in the morphological analyses (Barnes et al., 2010; Da Ma et al., 2018; Li X et al., 2013; Pell et al., 2008). Moreover, other seldom controlled variables in DD analyses are verbal and full-scale intelligence quotient (IQ), which were suggested to be correlated with brain size (McDaniel, 2005).

Based on this information, and following the previously published recommendations (Ramus et al., 2018), we set out to evaluate the GM and WM volume abnormalities in adult dyslexic individuals, whose reading-skill and neuropsychological profiles were rigorously assessed with a carefully composed test battery. We investigated brain properties while systematically using relevant covariates in the analyses. Furthermore, the connection between regional GM and WM volume and reading-related skills were analyzed as well as the association between these skills and group findings.

2. Materials and methods

2.1 Procedure

The data were collected in three separate sessions. In the first session neuropsychological testing was carried out. The second session included functional imaging (combined magneto- and electroencephalography, the data from which will be reported elsewhere) and responding to questionnaires. In the third session, anatomical MRI and diffusion tensor imaging data were acquired.

2.2 Neuropsychological tests and questionnaires

Participants filled out questionnaires concerning background information, DD in family, Adult Reading History Questionnaire (ARHQ; Lefly & Pennington, 2000), and Attention-Deficit/Hyperactivity Disorder (ADHD) Self-Report Scale Symptom Checklist ASRS-v1.1 (Kessler et al., 2005). The questions on the background included the participants' language skills and usage; language difficulties, neurological, psychiatric, and hearing disorders of him/her and the relatives (parents and siblings); the participant's other health issues and vision; music education and listening; and education and employment status. One questionnaire assessed the family history of DD.

The neuropsychological test battery was designed to assess phonological processing, reading, IQ, and working memory functions (Table 1). The domain of phonological processing included neuropsychological tests as follows: 'Pig Latin' test for assessing phonological awareness (Nevala et al., 2006), non-word span length for phonological short-term memory (Laasonen et al., 2002), and rapid alternating stimulus naming for rapid serial naming (Wolf, 1986). Technical reading skills (accuracy and speed) were tested with word and pseudoword list reading, and reading comprehension with text reading (Nevala et al., 2006). The verbal IQ was evaluated with the subtests Similarities and Vocabulary, and the performance IQ was assessed

with the subtests Block Design and Matrix Reasoning from the Wechsler Adult Intelligence Scale (Wechsler, 2005). Working memory functions were evaluated with Letter-number Series and Visual series subtests from the Wechsler Memory Scale, WMS-III (Wechsler, 2008). Composite scores of the test results (bolded in Table 1) were formed for phonological processing and technical reading by converting the raw scores (of subtests listed in Table 1 below the respective composite) to *z*-scores and averaging them, and for working memory the composite was formed according to WMS-III (Wechsler, 2008).

Please, insert Table 1 around here

2.3 Participants

Participants were recruited via social media, website of a related project, and through a local learning disabilities association. The exclusion criteria were as follows: attention deficit, developmental language disorder, some other language impairment, neurological disorders, psychiatric disorders, early bilingualism, left-handedness, uncorrected hearing or visual deficit, medication affecting central nervous system, alcohol or drug addiction, performance IQ below 80, and non-detachable metal in the body, large tattoos in the upper body, individualized school curriculum, and pregnancy. Furthermore, from the dyslexic group we excluded participants with a compensated DD (i.e. those who had experienced DD spectrum symptoms, but did not meet the criteria in our reading-skill tests).

The participant was classified as dyslexic if 1) a recent statement on dyslexia diagnosis was available from a psychologist, special education teacher, or similar, or the participant had both 2) reading-related problems in childhood as evaluated with the adult ARHQ and confirmed in an interview, and 3) a performance of at least one standard deviation (SD) below the average

of age-matched standardized control data (Laasonen et al., 2010) in at least two different reading subtests (word list reading, pseudoword list reading, text reading) in either speed or accuracy (Table 1). Participants classified to the Control group 1) had no language-related problems (and neither did their parents nor siblings), 2) reported no childhood problems in reading or writing as indicated by the ARHQ, and 3) performed within norm in at least two reading subtests in both speed and accuracy.

Forty-five participants completed the MR-imaging, the final sample consisting of 22 typically reading and 23 dyslexic participants. The groups were balanced in age, years of education, years of music education (since it may improve language skills; see, e.g., Kraus and Chandrasekaran, 2010), and sex (see Table 2 for descriptive statistics). The groups showed no significant differences in the demographic variables, whereas they significantly differed in the composite scores of phonological processing, technical reading, and working memory (Table 2). However, they differed in full-scale IQ (FIQ), which was, therefore, used as a covariate in the analyses.

The experiments were undertaken with the understanding and written consent of each subject, and following the Code of Ethics of the World Medical Association (Declaration of Helsinki). The Coordinating Ethics Committee of The Hospital District of Helsinki and Uusimaa approved the study protocol.

Please, insert Table 2 around here

2.4 MRI data acquisition

Participants were scanned with a 3T MRI Siemens Skyra scanner (Siemens Healthcare, Erlangen, Germany) using a 32-channel head coil, at AMI center in Aalto University, Finland.

High-resolution magnetization prepared rapid acquisition gradient-recalled (MPRAGE) T1 images were obtained (flip angle = 7°, TR = 2530 ms, TE = 3.3 ms, voxel size = 1.0 x 1.0 x 1.0 mm^3).

2.5 Data analysis

2.5.1 Voxel-based morphometry

Morphometric analysis was carried out using VBM (Ashburner and Friston, 2000) and the Statistical Parametric Mapping software (SPM8, Wellcome Department of Cognitive Neurology, UCL) under MATLAB 8.0.0 (The MathWorks Inc., Natick, MA, USA, version R2014b). VBM is an MRI analysis technique that allows comparison of GM and WM differences in focal brain regions between groups (Ashburner and Friston, 2000). After reorienting the individual T1 images using the anterior commissure as a landmark for the origin, Unified Segmentation (Ashburner and Friston, 2005) with medium regularization was applied to the T1 images, segmenting them precisely into GM, WM, and cerebrospinal fluid probability maps before normalizing them into the Montreal Neurological Institute (MNI) space. To preserve the original signal strength during the normalization, GM and WM segmented images were modulated. After this, to reduce residual inter-individual variability, GM and WM probability maps were smoothed using an isotropic spatial filter (FWHM = 6 mm). During each step, the images were visually checked for potential registration errors.

2.5.2 Statistical analyses

Preprocessed GM and WM images were then entered into a second-level analysis. First, three one-sample t-tests including all subjects (N=45) were calculated to evaluate the focal GM and WM structures associated with better performance in technical reading, phonological, and working memory tests. Then, using independent-sample t-tests, two different contrasts

(Controls>Dyslexics, Dyslexics>Controls) were calculated to evaluate the GM and WM volumetric differences between the dyslexics and controls. All results were thresholded at a whole-brain uncorrected P<.005 threshold and a familywise error rate (FWE) corrected P<.05 at the cluster level with a cluster extent of more than 100 contiguous voxels (Lieberman and Cunningham, 2009). Exact neuroanatomical regions were identified using the Automated Anatomical Labeling Atlas (Tzourio-Mazoyer et al., 2002) included in the xjView toolbox (http://www.alivelearn.net/xjview/).

All VBM analyses were adjusted for age, gender, and total intracranial volume (TIV) (Barnes et al., 2010; Da Ma et al., 2018; Li X et al., 2013; Pell et al., 2008), and to follow the previously published recommendations (Ramus et al., 2018) and to take into account the group difference, FIQ, was also added as a covariate in the VBM analyses. Partial correlations (two-tailed) with previously mentioned covariates were calculated between each individual significant cluster and the three composite scores (technical reading, phonological processing, working memory; see Table 1) using SPSS (IBM Corp. Released 2012. IBM SPSS Statistics for Windows, Version 24.0. Armonk, NY: IBM Corp.). To control for multiple comparisons, false discovery rate (FDR) approach was used and only significant results are reported.

3. Results

3.1 Anatomical correlates of neuropsychological tests

We first determined how reading, phonological processing, and working memory were associated with brain structures including both groups in the analysis (Fig. 1; Table 3). Better performance in technical reading (accuracy and speed combined) was associated with greater GM volume in two clusters comprising the left temporal (superior and middle temporal gyrus, fusiform gyrus), limbic (amygdala, hippocampus, parahippocampal gyrus) and insular areas

(R=.641, P<.001) as well as the left cerebellum (R=.499, P=.001). In addition, better technical reading performance was associated with greater WM volume in two clusters comprising right frontal areas (inferior frontal and precentral gyrus), basal ganglia (putamen, globus pallidus), insula, pons, and left (R=.625, P<.001) and right cerebellum (R=.490, P=.001). Higher working memory scores were associated with greater WM volume in the brainstem and bilaterally in the cerebellum (R=.518, P=.001).

Please, insert Fig. 1 around here

Please, insert Table 3 around here

3.2 Neuroanatomical group differences

In whole-brain analyses comparing the groups, we found both greater GM and WM volumes in the control than dyslexic participants in several brain areas (Fig. 2; see Table 4 for descriptive statistics). Greater GM volumes (Controls>Dyslexics) were found in a cluster comprising the left superior temporal gyrus (STG), IFG, insula, as well as in the left limbic (amygdala, hippocampus) and basal ganglia areas (claustrum, putamen). Greater GM volume in the observed areas correlated significantly with higher technical reading score (R=.431, P=.015). In addition, the control participants had greater WM volume in the right middle temporal gyrus (MTG) and hippocampus compared with the dyslexic participants. Again, greater WM volume in these areas correlated significantly with a higher technical reading score (R=.426, P=.015). Importantly, these results did not correlate with FIQ and there were no significant correlations between FIQ and the volumes of GM and WM.

Crucially, there was an overlap between brain areas associated with technical reading skills and brain areas in which participants with DD had a reduced GM volume. This overlapping area

comprised the left STG, insula, and limbic areas (amygdala, hippocampus). Greater GM volume in these areas correlated with higher technical reading score (R=.577, P<.001). Interestingly, the groups showed no significant differences in total GM volume, total WM volume, total cerebrospinal fluid volume, total brain volume, or TIV.

Please, insert Fig. 2 around here

Please, insert Table 4 around here

4. **Discussion**

There is an obvious need to understand neural underpinnings of DD, yet, no clear consensus has so far been reached (e.g., Ramus et al., 2018, for a review). Our study was set out at illuminating the neuroanatomical basis of DD using a whole-brain volumetric analysis and a thorough reading and neuropsychological test battery. We used multiple VBM analyses which were refined so that the commonly present limitations in VBM studies on DD would be curtailed. These analyses included hypothesis-free whole-brain analysis and correlational analyses of neuropsychological tests and neuroanatomy. We found decreased volumes of GM in participants with DD, comprising a left-hemispheric network including superior temporal and inferior frontal gyri, insula, the limbic system, and basal ganglia, and of WM, including the right middle temporal gyrus and hippocampus. However, unlike many previous studies (Ramus et al., 2018, for a review), we found no group differences in total, GM, or WM volumes. Thus, overall volumetric differences do not explain the deficits in reading and related skills in our dyslexic sample. Crucially, a significant overlap was found between the brain areas with decreased volumes in participants with DD and the brain areas that correlated with technical reading abilities. These regions also largely overlap with the neural network that has been

associated with reading (e.g., Kujala et al., 2007). These converging results highlight the robustness of our current findings on DD.

4.1 Associations between anatomy and reading-related skills

As could be expected, better performance in technical reading was significantly correlated with a larger GM volume of an extensive neural network, since a wide range of functions subserved by multiple brain areas are needed for fluent and accurate reading (e.g., Paulesu et al., 2000; Kujala et al., 2007; Levy et al., 2009; Welcome and Joanisse, 2012; Oberhuber et al., 2013). This network comprised the left temporal (STG, MTG, fusiform gyrus), limbic (amygdala, hippocampus, parahippocampal gyrus), and insular areas, as well as the left cerebellum. In addition, higher technical reading scores were associated with a larger WM volume in right frontal areas (inferior frontal and precentral gyrus), basal ganglia (putamen, globus pallidus), insula, pons, as well as left and right cerebellum. These results are compatible with a large body of previous research associating reading skills and subskills with brain structures in frontal, temporal, cerebellar, and subcortical areas (Paulesu et al., 2000; Kujala et al., 2007; Levy et al., 2009; Welcome and Joanisse, 2012; Oberhuber et al., 2013; Stoodley and Stein, 2013).

Higher working memory scores were associated with increased WM volume in the brainstem and bilaterally in the cerebellum. This is consistent with a previously found association between working memory load and the amplitude of brainstem responses (Sörqvist et al., 2012), indicating the involvement of brainstem in working memory functions. Brainstem also regulates vigilance (Iovino et al., 2019). Therefore, it could be speculated that these associations are related to attention, which is tightly involved in working memory functions requiring "attentional spotlight" to items that are actively consciously processed (e.g., Rhodes and Cowan, 2018).

The association between the cerebellar WM volume and working memory is in accordance with currently known cerebellar functions. Cerebellum is involved in regulating the direction of attention, detecting errors, timing and sequencing, as well as in associative learning, all vitally involving working memory (Stoodley and Stein, 2013, for a review). This is also compatible with the rich cerebellar interconnections with the prefrontal cortex and other association cortices (Stoodley and Stein, 2013), which belong to the neural network responsible for working memory functions (Knight et al., 1999).

We found no significant correlations between phonological scores and neuroanatomical findings. A potential reason for this might be that only relatively small brain areas, suggested to be restricted to the left temporal and angular gyri (Glezer et al., 2016) process phonology and that the present analyses controlling for various nuisance variables impede small effects to become significant. Moreover, while both groups had more than 20 subjects, our overall sample size remained moderate, which in turn decreases power to observe small effects in the data. Furthermore, using our phonological composite score in the correlation analysis might not be optimal, since the tests included tap also into other functions besides pure phonological processing. Another possible explanation is that phonological deficits might primarily arise from subtle anatomical and functional changes in pathways interconnecting posterior and anterior superior temporal regions (Richardson et al., 2011) rendering the overall methodology (i.e. VBM) disadvantageous here.

4.2 Neuroanatomical group differences

We found reduced GM volume in dyslexic participants in left-hemispheric regions of STG, IFG, insula, limbic system, and basal ganglia, and WM volume in right temporal and hippocampal areas. However, even though the most robust finding in previous studies has been a reduced total brain volume in DD (see Ramus et al., 2018, for a meta-analysis), we found no

group differences in the brain, GM, or WM volumes. Thus, overall volumetric differences do not explain the deficits in reading and related skills in the current sample of dyslexics. Importantly, greater GM and WM volumes in left STG, insula, and limbic areas correlated with better technical reading skills, that is, reading accuracy and speed. This finding, directly linking the same brain areas to the core reading skills and reduced neural volumes in DD underlines the robustness and reliability of our results. Our study included 23 dyslexic and 22 control participants, which just about exceeds the minimum N (>20) suggested for neuroanatomical studies on DD (Ramus et al., 2018). However, since the group differences and reading-skill associations pinpoint the same areas, which also belong to the reading network (Kujala et al., 2007), we are confident that our findings are not "false positive" results, a concern which has been raised often in this field of research due to the varying results reported (Ramus et al., 2018).

Unlike in many other studies (see Ramus et al., 2018, for a meta-analysis) the dyslexic sample in our study did not differ from controls in total brain, GM, WM, or cerebrospinal fluid volume or TIV. The lack of group differences in the total brain volume is particularly interesting since it has been speculated that the reduced total brain volume in DD could either be associated with the etiology of DD or a consequence of this disorder (Ramus et al., 2018). However, contradicting results have been published (Frye et al., 2010) and the finding could depend on the population studied (adults vs. children). Our data showing no group differences in a range of brain measures (total brain, GM, WM, cerebrospinal fluid volume, and TIV) suggest that the occurrence of DD may not (only) rely on brain volume reduction as a predisposing factor or as a de rigueur developmental consequence.

Our group and correlational results on the GM volume in STG, IFG, and insula are consistent with earlier neuroanatomical findings. For example, according to the meta-analysis of Richlan et al. (2013) the most consistent GM reductions in DD have been found in superior temporal

areas and according to the meta-analysis of Ramus et al. (2018) in left perisylvian and OT regions. Furthermore, a large number of studies proposed that these areas belong to the network, which is vital for reading (Kujala et al., 2007; Paulesu et al., 2000; Kujala et al., 2007; Levy et al., 2009; Welcome and Joanisse, 2012; Oberhuber et al., 2013). Our results are also compatible with functional imaging studies on DD, which have revealed hypoactivations in superior, middle, and inferior temporal areas as well as inferior frontal gyrus in the left hemisphere (Richlan et al., 2009, for a meta-analysis). Next, we will inspect our findings in the light of the currently known functions of these areas.

Superior temporal areas were consistently found to be active during reading (Sandak et al., 2004; Richlan, 2014), and proposed to reflect semantic (Helenius et al., 1998; Halgren et al., 2002) and phonological analysis (Jobard et al., 2003). A restricted lesion to left STG was reported to result in pure word deafness, with the patient being able to read, write, and perceive nonspeech auditory stimuli but not to correctly process speech input (Maffei et al., 2017). The reduced GM in the STG of our dyslexic participants might, therefore, reflect a speech-specific deficit, consistent with one of the leading theories on DD according to which reading impairments in DD are largely based on poor speech sound representations (Peterson and Pennington, 2012; see, however, Ramus, 2014). This result is highly compatible with a meta-analysis based on which it was proposed that the phonological processing deficit is an endophenotype of DD (Snowling and Melby-Lervåg, 2016).

Activations of **left IFG** have been reported for words and legal pseudowords (pseudowords not violating the rules of the language) but not for consonant strings (Wilson et al., 2005; Cornelissen et al., 2009) or faces (Cornelissen et al., 2009). Its subregion BA 44 was found to be activated by both words and pseudowords, whereas subregion 45 was only activated by words (during a lexical decision task; Heim et al., 2005). These areas were also found to be involved in rapid learning of novel word-forms (Kimppa et al., 2015; 2018), an effect not found

in dyslexic children (Kimppa et al., 2018). Furthermore, it was proposed that left IFG participates in nonlexical phonological grapheme-phoneme conversion during reading (Heim et al., 2005). Overall, these results, as well as a meta-analysis (Bookheimer 2002), suggest that left IFG regions are involved in word and phonological speech processing. The reduced left IFG GM in our dyslexic participants might, therefore, reflect their observed problems of word-form processing and learning (Kimppa et al., 2018), as well as deficits in phonological processing (e.g., Snowling and Melby-Lervåg, 2016) and grapheme-phoneme conversion (Snowling, 1980).

Whereas the involvement of cortical areas in reading and reading impairment has been extensively studied, less attention has been paid to **subcortical areas** so far (see Krishnan et al., 2016, for a review). Our study revealed a reduced GM volume in the dyslexic sample in limbic (hippocampus and amygdala) and basal ganglia (claustrum and putamen) areas. Similar findings were obtained in the left hemisphere of dyslexic males by Casanova et al. (2005), reporting anomalies in amygdala, hippocampus, putamen, and globus pallidus in their volumetric analysis. Furthermore, in dyslexic children, a successful reading intervention was found to result in increased GM volume in several brain areas including hippocampus (Krafnick et al., 2011). Hippocampus is known to have a central role in memory functions, particularly in memory consolidation (see Buzsáki and Moser, 2013, for a review). A reduced GM in the hippocampus of the dyslexic participants could be associated with their poor ability to "tune in" or form memory representations of novel repetitive stimuli (see Ahissar, 2007, for a review; see also Kimppa et al., 2018). This was suggested to influence the efficacy of short-term memory and underlie a wide range of difficulties associated with DD (Ahissar, 2007).

Amygdala, besides having a central role in processing fear-eliciting stimuli (Markowitsch, 1998) was also suggested to be involved in memory and reward networks and to process valence, salience, and stimulus relevance (Sander et al., 2003; Janak and Tye, 2015, for

reviews), the left amygdala having a higher affinity to language than the right one (Markowitsch, 1998). Processing of relevance and salience, in turn, are strongly linked with attention functions, which are known to be dysfunctional in at least some subgroups of dyslexic individuals (Hari et al., 2001; see Krause, 2017, for a review). Despite our participants having been screened for ADHD with a questionnaire, it is possible that they had some attentional difficulties that do not lead to a suspicion of ADHD.

The putamen has been associated with the initiation of movements (e.g., Tricomi et al., 2009). However, it was also found to be activated more by reading words and pseudowords than by naming pictures and colours (Oberhuber et al., 2013), the effect being stronger in the left than right putamen, suggesting that it belongs to the reading network. The left putamen was found to predominantly coactivate with left-hemispheric regions that have a direct association with language processes (see Vinas-Guasch and Wu, 2017, for a meta-analysis). Our finding of reduced GM in the left putamen in participants with DD is therefore highly compatible with these results.

Globus pallidus is a major component of basal ganglia, and it is associated with voluntary movements (Turner and Anderson, 2005). Since globus pallidus receives its major input from striatum, which includes putamen in addition to caudate nucleus, it could be speculated that its decreased GM volume in DD results from the decreased GM volume of putamen.

The claustrum is a narrow structure located between insula and putamen, the functions of which are not well-known. It is thought to be the most densely connected brain structure, involved in integrating a range of cortical inputs and in segregating attention (Goll et al., 2015). Perhaps this diminished GM volume in claustrum in our dyslexic sample might be related to the observed problems in integrating sensory information (e.g., Widmann et al., 2012) and in regulating attention in DD (see Krause, 2015, for a review).Yet, at this stage these suggestions are highly speculative and need confirmation from further studies.

In the current study, group differences in WM volume were limited to right MTG and hippocampus, with dyslexic participants having a smaller volume than controls. Some previous studies have shown reduced right-hemispheric WM volumes in DD (e.g., in inferior longitudinal fasciculus, Lu et al., 2016, Banfi et al., 2019) as well as absent rightward asymmetry of the inferior fronto-occipital fasciculus (IFOF) in a subpopulation of DD (Banfi et al., 2019). IFOF is a ventral associative bundle that interconnects occipital, temporal and orbitofrontal areas (Sarubbo et al., 2013). It encompasses MTG and runs next to hippocampus, so it is possible that our results showing decreased WM volume in DD in right MTG and hippocampus reflect an anomalous rightward asymmetry. Close to hippocampus, the IFOF is intertwined with another ventral tract, uncinate fasciculus, which interconnects anterior temporal and orbitofrontal regions (Catani et al., 2002; Bakhtiari et al., 2014). Recently, the right uncinate fasciculus integrity has been associated with better reaction time in a reading task (Bakhtiari et al., 2014). Overall, it has been proposed that the right hemisphere might have a compensatory function in reading in DD (Temple et al., 2003; Eden et al., 2004; Hoeft et al., 2011). In light of these suggestions, it could be speculated that the diminished right-hemispheric WM volume of our adults with DD might reflect inefficient compensatory mechanisms, since these participants still had a persistent dyslexia in the adult age.

4.3 Conclusions

The current study evaluated the neuroanatomical basis of DD using rigorous neuropsychological testing and multiple VBM analyses refined to curtail the common limitations of previous VBM studies on DD. We found reduced GM volumes in adults with DD in left STG and IFG, which is compatible with proposed core nodes of reading (Kujala et al., 2007) and findings in meta-analyses on GM volumes in DD (e.g., Richlan et al., 2013; Ramus et al., 2018). Furthermore, GM volume reductions were found in our DD sample in subcortical structures, which have so far scarcely been studied, including left limbic and basal ganglia

areas. WM reductions, in turn, included right temporal and hippocampal structures. Importantly, these brain areas were also correlated with reading accuracy and speed in the whole sample, suggesting that we were reliably able to illuminate the core dysfunctional neural reading network in DD. Next, it would be important to determine which of these volumetric changes in DD underlie the reading deficit and which ones have resulted from, for example, reduced exposure to print. Moreover, as multiple genetic and environmental risks may lead to reading deficits (e.g., Peterson and Pennington, 2015, for a review), a delineation of subgroups of distinct reading deficit profiles from a larger sample would increase our understanding of DD. This, in turn, is vital for developing interventions that are applicable to individuals with various reading deficit profiles.

Funding

This study was financially supported by the Academy of Finland (grants 276414 and 316970), Jane and Aatos Erkko Foundation, and Kela.

Competing interests

The authors report no competing interests.

References

Ahissar, M., 2007. Dyslexia and the anchoring- deficit hypothesis. TICS 11, 458–465.

American Psychiatric Association, 2000. Diagnostic and statistical manual of mental disorders (4th ed., text rev.). Washington, DC: American Psychiatric Association.

Ashburner, J., Friston, K.J., 2005. Unified segmentation. Neuroimage 26, 839–851.

Ashburner, J., Friston, K. J., 2001. Why voxel-based morphometry should be used. Neuroimage 14, 1238–1243.

Ashburner J., Friston K.J., 2000. Voxel-based morphometry–the methods. Neuroimage 11, 805–821.

Bakhtiari, R., Boliek, C., Cummine, J., 2014. Investigating the contribution of ventral-lexical and dorsal-sublexical pathways during reading in bilinguals. Front Hum Neurosci 8, 507.

Banfi C., Koschutnig K., Moll K., Schulte-Korne G., Fink A., Landerl K., 2019. White matter alterations and tract lateralization in children with dyslexia and isolated spelling deficits. Hum Brain Mapp 40, 765–776.

Barnes J., Ridgway G.R., Bartlett J., Henley S.M., Lehmann M., Hobbs N., Clarkson M.J., MacManus D.G., Ourselin S., Fox N.C., 2010. Head size, age and gender adjustment in MRI studies: a necessary nuisance? Neuroimage 53, 1244–1255.

Bookheimer S., 2002. Functional MRI of language: new approaches to understanding the cortical organization of semantic processing. Annu Rev Neurosci 25, 151–188.

Brambati, S. M., Termine, C., Ruffino, M., Stella, G., Fazio, F., Cappa, S. F., & Perani, D., 2004. Regional reductions of gray matter volume in familial dyslexia. Neurology 63, 742–745.

Brown, W.E., Eliez, S., Menon, V., Rumsey, J.M., White, C.D., Reiss, A.L., 2001. Preliminary evidence of widespread morphological variations of the brain in dyslexia. Neurology 56, 781–783.

Buzsáki, G., Moser, E., 2013. Memory, navigation and theta rhythm in the hippocampal-entorhinal system. Nat Neurosci 16, 130–138.

Casanova, M.F., El-Baz, A.S., Giedd, J., Rumsey, J.M., Switala, A.E., 2005. Increased White Matter Gyral Depth in Dyslexia: Implications for Corticocortical Connectivity. J Autism Dev Disord 40, 21–29.

Catani M., Howard R.J., Pajevic S., Jones D.K., 2002. Virtual in vivo interactive dissection of white matter fasciculi in the human brain. Neuroimage 17, 77–94.

Cornelissen, P.L., Kringelbach, M.L., Ellis, A.W., Whitney, C., Holliday, I.E., Hansen, P.C., 2009. Activation of the Left Inferior Frontal Gyrus in the First 200 ms of Reading: Evidence from Magnetoencephalography (MEG). Plos One 4, e5359.

Dole, M., Meunier, F., Hoen, M., 2013. Gray and white matter distribution in dyslexia: a VBM study of superior temporal gyrus asymmetry. Plos One 8, e76823.

Drake, W.E., 1968. Clinical and Pathological Findings in a Child with a Developmental Learning Disability. J Learn Disabil 1, 486–502.

Eckert, M.A., Berninger, V.W., Vaden, K.I. Jr., Gebregziabher, M., Tsu L., 2016. Gray Matter Features of Reading Disability: A Combined Meta-Analytic and Direct Analysis Approach. eNeuro 3, ENEURO.0103-15.2015.

Eckert, M. A., Leonard, C. M., Wilke, M., Eckert, M., Richards, T., Richards, A., & Berninger, V., 2005. Anatomical signatures of dyslexia in children: unique information from manual and voxel-based morphometry brain measures. Cortex 41, 304–315.

Eden, G.F., Jones, K.M., Cappell, K., Gareau, L., Wood, F.B., Zeffiro, T.A., Dietz N.A.E., Agnew J.A., Flowers D.L., 2004. Neural changes following remediation in adult developmental dyslexia. Neuron 44, 411–422.

Evans, T. M., Flowers, D. L., Napoliello, E. M., & Eden, G. F., 2014. Sex-specific gray matter volume differences in females with developmental dyslexia. Brain Structure and Function 219, 1041–1054.

Frye, R. E., Liederman, J., Malmberg, B., McLean, J., Strickland, D., & Beauchamp, M. S., 2010. Surface area accounts for the relation of gray matter volume to reading-related skills and history of dyslexia. Cerebral Cortex 20, 2625–2635.

Glezer, L.S., Eden, G., Jiang, X., Luetje, M., Napoliello, E., Kim, J., Riesenhuber, M., 2016. Uncovering phonological and orthographic selectivity across the reading network using fMRI-RA. Neuroimage 138, 248–256.

Goll, Y., Atlan, G., Citri, A., 2015. Attention: the claustrum. TINS 38, 486–495.

Good, C.D., Johnsrude, I.S., Ashburner, J., Hensona, R.N.A., Friston, K.J., Frackowiaka, R.S.J., 2001. A voxel-based morphometric study of ageing in 465 normal adult human brains. Neuroimage 14, 21–36.

Halgren, E., Dhondac, R.P., Christensen, N., Van Pettene, C., Marinkovica, K., Lewine, J.D., Dale, A.M., 2002. N400-like Magnetoencephalography Responses Modulated by Semantic Context, Word Frequency, and Lexical Class in Sentences. Neuroimage 17, 110–116.

Hari, R., Renvall, H., Tanskanen, T,. 2001. Left minineglect in dyslexic adults. Brain 124, 1373–1380.

Heim, S., Alter, K., Ischebeck, A.K., Amunts, K., Eickhoff, S.B., Mohlberget, H., Zilles, K., von Cramon, D.Y., Friederici, A.D., 2005. The role of left Brodmann's areas 44 and 45 in reading words and pseudowords. Cogn Brain Res 25, 982–993.

Helenius, P., Salmelin, R., Service, E., Connolly, J.F., 1998. Distinct time courses of word and context comprehension in the left temporal cortex. Brain 121, 1133–1142.

Hoeft, F., McCandliss, B. D., Black, J. M., Gantman, A., Zakerani, N., Hulme, C., Lyytinen H., Whitfield-Gabrieli, S., Glover, G.H., Reiss, A.L. & Gabrieli, J. D., 2011. Neural systems predicting long-term outcome in dyslexia. Proceedings of the National Academy of Sciences 108, 361–366.

Hoeft, F., Meyler, A., Hernandez, A., Juel, C., Taylor-Hill, H., Martindale, J. L., McMillon, G., Kolchugina G., Black, J.M., Faizi, A., Deutsch, G. K., Siok, W.T., Reiss, A.L., Whitfield-Gabrieli, S., Gabrieli, J.D.E., 2007. Functional and morphometric brain dissociation between dyslexia and reading ability. Proceedings of the National Academy of Sciences 104, 4234–4239.

Iovino, M., Messana, T., De Pergola, G., Iovino, E., Guastamacchia, E., Giagulli, V.A., Triggiani, V., 2019. Vigilance States: Central Neural Pathways, Neurotransmitters and Neurohormones. Endocr Metab Immune 19, 26–37.

Janak, P.H., Tye, K.M., 2015. From circuits to behaviour in the amygdala. Nature 517, 284–292.

Jednoróg, K., Marchewka, A., Altarelli, I., Monzalvo Lopez, A. K., van Ermingen- Marbach, M., Grande, M., Grabovska, A., Heim, S. & Ramus, F. 2015. How reliable are gray matter disruptions in specific reading disability across multiple countries and languages? Insights from a large- scale voxel- based morphometry study. Human brain mapping 36, 1741–1754.

Jednoróg, K., Gawron, N., Marchewka, A., Heim, S., Grabowska, A. 2014. Cognitive subtypes of dyslexia are characterized by distinct patterns of grey matter volume. Brain Structure and Function 219, 1697–1707.

Jobard, G., Crivello, F., Tzourio-Mazoyera, N., 2003. Evaluation of the dual route theory of reading: a metanalysis of 35 neuroimaging studies. Neuroimage 20, 693–712.

Kessler, R.C., Adler L., Ames, M., Demler, O., Faraone, S., Hiripi, E., Howes, M.J., Jin, R., Secnik, K., Spencer, T., Ustun, T.B., Walters, E.E., 2005. World Health Organization Adult ADHD Self-report Scale (ASRS): A Short Screening Scale for Use in the General Population. Psychological Medicine 35, 245–256.

Kimppa, L., Shtyrov, Y., Partanen, E., Kujala T., 2018. Impaired neural mechanism for online novel word acquisition in dyslexic children. Scientific reports 8, 1-12.

Kimppa, L., Kujala, T., Leminen, A., Vainio, M., Shtyrov, Y., 2015. Rapid and automatic speech-specific learning mechanism in human neocortex. Neuroimage 118, 282–291.

Knight, R.T., Staines, W.R., Swick, D., Chao, L.L., 1999. Prefrontal cortex regulates inhibition and excitation in distributed neural networks. Acta Psychologica 101, 159–178.

Krafnick, A. J., Flowers, D. L., Luetje, M. M., Napoliello, E. M., & Eden, G. F., 2014. An investigation into the origin of anatomical differences in dyslexia. Journal of Neuroscience 34, 901–908.

Krafnick, A. J., Flowers, D. L., Napoliello, E. M., & Eden, G. F., 2011. Gray matter volume changes following reading intervention in dyslexic children. Neuroimage 57, 733–741.

Kraus, N., Chandrasekaran, B., 2010. Music training for the development of auditory skills. Nature Reviews Neuroscience 11, 599–605.

Krause, M.B., 2015. Pay Attention!: Sluggish Multisensory Attentional Shifting as a Core Deficit in Developmental Dyslexia. Dyslexia 21, 285–303.

Krishnan, S., Watkins, K.E., Bishop, D.V.M., 2016. Neurobiological Basis of Language Learning Difficulties. TICS 20, 701–714.

Kronbichler, M., Wimmer, H., Staffen, W., Hutzler, F., Mair, A., Ladurner, G., 2008. Developmental dyslexia: gray matter abnormalities in the occipitotemporal cortex. Human brain mapping 29, 613–625.

Kujala, J., Pammer, K., Cornelissen, P., Roebroeck, A., Formisano, E., Salmelin, R., 2007. Phase coupling in a cerebro-cerebellar network at 8-13 Hz during reading. Cerebral Cortex 17, 1476–1485.

Laasonen, M., Service, E., Virsu, V., 2002. Crossmodal Temporal Order and Processing Acuity in Developmentally Dyslexic Young Adults. Brain Lang 80, 340–354.

Lefly, D.L., Pennington, B.F., 2000. Reliability and Validity of the Adult Reading History Questionnaire. J Learn Disabil 33, 286–296.

Levy, J., Pernet, C., Treserras, S., Boulanouar, K., Aubry, F., Démonet, J.-F., Celsis, P., 2009. Testing for the Dual-Route Cascade Reading Model in the Brain: An fMRI Effective Connectivity Account of an Efficient Reading Style. Plos One 4, e6675.

Li, X., Pu, F., Fan, Y., Niu, H., Li, S., Li, D., 2013. Age-related changes in brain structural covariance networks. Frontiers in human neuroscience 7, 98.

Lieberman, M.D., Cunningham, W.A., 2009. Type I and Type II error concerns in fMRI research: re-balancing the scale. Soc Cogn Affect Neurosci 4, 423–428.

Lu, C., Qi, Z., Harris, A., Weil, L.W., Han, M., Halverson, K., Perrachione, T.K., Kjelgaard, M., Wexler, K., Tager-Flusberg, H., Gabrieli, J.D., 2016. Shared neuroanatomical substrates

of impaired phonological working memory across reading disability and autism. Biol Psychiatry Cogn Neurosci Neuroimaging 1, 169–177.

Ma, D., Popuri, K., Bhalla, M., Sangha, O., Lu, D., Cao, J., Jacova, C., Wang, L., Beg, M.F., Alzheimer's Disease Neuroimaging Initiative, 2019. Quantitative assessment of field strength, total intracranial volume, sex, and age effects on the goodness of harmonization for volumetric analysis on the ADNI database. Human brain mapping 40, 1507–1527.

Maffei, C., Capasso, R., Cazzollia, G., Colosimo, C., Dell'Acqua, F., Piludu, F., Catani, M., Miceli, G., 2017. Pure word deafness following left temporal damage: Behavioral and neuroanatomical evidence from a new case. Cortex 97, 240–254.

Maldjian, J.A., Laurienti, P.J., Kraft, R.A., Burdette, J.H., 2003. An automated method for neuroanatomic and cytoarchitectonic atlas-based interrogation of fmri data sets. Neuroimage 19, 1233–1239.

Markowitsch, H.J., 1998. Differential contribution of right and left amygdala to affective information processing. Behav Neurol 11, 233–244.

McArthur, G., Kohnen, S., Larsen, L., Jones, K., Anandakumar, T., Banales, E., 2013. Getting to grips with the heterogeneity of developmental dyslexia. Cogn Neuropsychol 30, 1–24.

McDaniel, M.A., 2005. Big-brained people are smarter: a meta-analysis of the relationship between in vivo brain volume and intelligence. Intelligence 33, 337–346.

Menghini, D., Hagberg, G. E., Petrosini, L., Bozzali, M., Macaluso, E., Caltagirone, C., Vicari, S., 2008. Structural correlates of implicit learning deficits in subjects with developmental dyslexia. Annals of the New York Academy of Sciences 1145, 212–221.

Nevala, J., Kairaluoma, L., Ahonen, T., Aro, M., Holopainen, L., 2006. Lukemis- ja kirjoittamistaitojen yksilötestistö nuorille ja aikuisille [reading and writing test for adolescents and adults]. Jyväskylä (FI): Niilo Mäki Institute.

Oberhuber M., Jones, Ō.P., Hope, T.M.H., Prejawa, S., Seghier, M.L., Green, D.W., Price, C.J., 2013. Functionally distinct contributions of the anterior and posterior putamen during sublexical and lexical reading. Front Hum Neurosci 19, 787.

Paulesu, E., McCrory, E., Fazio, F., Menoncello, L., Brunswick, N., Cappa, S.F., Cotelli, M., Cossu, G., Corte, F., Lorusso, M., Pesenti, S., Gallagher, A., Perani, D., Price, C., Firth, C.D., Frith, U., 2000. A cultural effect on brain function. Nat Neurosci 3, 91–96.

Pell, G. S., Briellmann, R. S., Chan, C. H. P., Pardoe, H., Abbott, D. F., Jackson, G. D., 2008. Selection of the control group for VBM analysis: influence of covariates, matching and sample size. Neuroimage 41, 1324–1335.

Pernet, C., Andersson, J., Paulesu, E., Demonet, J. F., 2009a. When all hypotheses are right: a multifocal account of dyslexia. Human brain mapping 30, 2278–2292.

Pernet, C. R., Poline, J. B., Demonet, J. F., Rousselet, G. A., 2009b. Brain classification reveals the right cerebellum as the best biomarker of dyslexia. BMC neuroscience 10, 67.

Peterson, R.L, Pennington, B.F., 2015. Developmental dyslexia. Annu Rev Clin Psychol 11, 238–307.

Peterson, R.L., Pennington, B.F., 2012. Developmental dyslexia. Lancet 379, 1997–2007.

Ramus, F., Altarelli, I., Jednorógc, K., Zhao, J., Covella, L.S., 2018. Neuroanatomy of developmental dyslexia: Pitfalls and promise. Neurosci Biobehav Rev 84, 434–452

Ramus, F., 2014. Neuroimaging sheds new light on the phonological deficit in dyslexia. TiCS 18, 274–275.

Rhodes, S., Cowan, N., 2018. Attention in working memory: attention is needed but it yearns to be free. Ann N Y Acad Sci 1424, 52–63.

Richardson, F.M., Seghier, M.L., Leff, A.P., Thomas, M.S., Price, C.J., 2011. Multiple routes from occipital to temporal cortices during reading. J Neurosci 31, 8239–8247.

Richlan, F., 2014. Functional neuroanatomy of developmental dyslexia: the role of orthographic depth. Front Hum Neurosci 8, 347.

Richlan, F., Kronbichler, M., Wimmer, H., 2013. Structural abnormalities in the dyslexic brain: a meta-analysis of voxel-based morphometry studies. Hum Brain Mapp 34, 3055–3065.

Richlan, F., Kronbichler, M., Wimmer, H., 2009. Functional abnormalities in the dyslexic brain: A quantitative meta- analysis of neuroimaging studies. Human brain mapping 30, 3299–3308.

Sandak, R., Mencl, W.E., Frost, S.J., Rueckl, J.G., Katz, L., Moore, D.L., Mason, S.A., Fulbright, R.K., Constable, R.T., Pugh, K.R., 2004. The neurobiology of adaptive learning in reading: A contrast of different training conditions. Cogn Affect Behav Neurosci 4, 67–88.

Sander, D., Grafman, J., Zalla, T., 2003. The Human Amygdala: An Evolved System for Relevance Detection. Rev Neurosci 14, 303–316.

Sarubbo, S., De Benedictis, A., Maldonado, L.L., Basso, G., Duffau, H., 2013. Frontal terminations for the inferior fronto-occipital fascicle: Anatomical dissection, DTI study and functional considerations on a multi-component bundle. Brain Struct Funct 218, 21-37.

Shaywitz, S.E., 1998. Dyslexia. N Engl J Med 338, 307–312.

Silani, G., Frith, U., Demonet, J. F., Fazio, F., Perani, D., Price, C., Frith, C.D., Paulesu, E., 2005. Brain abnormalities underlying altered activation in dyslexia: a voxel-based morphometry study. Brain 128, 2453–2461.

Siok, W. T., Niu, Z., Jin, Z., Perfetti, C. A., Tan, L. H., 2008. A structural–functional basis for dyslexia in the cortex of Chinese readers. Proceedings of the National Academy of Sciences 105, 5561–5566.

Snowling, M.J., Melby-Lervåg, M., 2016. Oral language deficits in familial dyslexia: A meta-analysis and review. Psychol Bull 142, 498–545.

Snowling, M.J., 1980. The development of grapheme-phoneme correspondence in normal and dyslexic readers. J Exp Child Psychol 29, 294–305.

Steinbrink, C., Vogt, K., Kastrup, A., Müller, H. P., Juengling, F. D., Kassubek, J., Riecker, A., 2008. The contribution of white and gray matter differences to developmental dyslexia: insights from DTI and VBM at 3.0 T. Neuropsychologia 46, 3170–3178.

Stoodley, C.J., Stein, J.F., 2013. Cerebellar Function in Developmental Dyslexia. Cerebellum 12, 267–276.

Sörqvist P, Stenfelt S, Rönnberg J. 2012. Working Memory Capacity and Visual–Verbal Cognitive Load Modulate Auditory–Sensory Gating in the Brainstem: Toward a Unified View of Attention. J Cognitive Neurosci. 24:2147–54.

Tamboer, P., Scholte, H. S., Vorst, H. C., 2015. Dyslexia and voxel-based morphometry: correlations between five behavioural measures of dyslexia and gray and white matter volumes. Annals of dyslexia 65, 121–141.

Temple, E., Deutsch, G.K., Poldrack, R.A., Miller, S.L., Tallal, P., Merzenich, M.M., Gabrieli J.D.E., 2003. Neural deficits in children with dyslexia ameliorated by behavioral remediation: Evidence from functional MRI. PNAS 100, 2860–2865.

Tricomi, E., Balleine, B.W., O'Doherty, J.P., 2009. A specific role for posterior dorsolateral striatum in human habit learning. Eur J Neurosci 29, 2225–2232.

Turner, R.S., Anderson, M.E., 2005. Context-dependent modulation of movement-related discharge in the primate globus pallidus. J Neurosci 25, 2965–2976.

Tzourio-Mazoyer, N., Landeau, B., Papathanassiou, D., Crivello, F., Etard, O., Delcroix, N., Mazoyer, B., Joliot, M., 2002. Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. Neuroimage 15, 273–289.

van Oers, C. A., Goldberg, N., Fiorin, G., van den Heuvel, M. P., Kappelle, L. J., Wijnen, F. N., 2018. No evidence for cerebellar abnormality in adults with developmental dyslexia. Experimental brain research 236, 2991–3001.

Viñas-Guasch, N., & Wu, Y. J. 2017. The role of the putamen in language: a meta-analytic connectivity modeling study. Brain Structure and Function. 222:3991–4004.

Vinckenbosch, E., Robichon, F., Eliez, S., 2005. Gray matter alteration in dyslexia: converging evidence from volumetric and voxel-by-voxel MRI analyses. Neuropsychologia 43, 324–331.

Wechsler, D., 2008. WMS-III manual. Helsinki (FI): Psykologien Kustannus Oy.

Wechsler, D., 2005. Wechsler Adult Intelligence Scale-Third edition: Manual. Helsinki (FI): Psykologien Kustannus Oy.

Welcome, S.E., Joanisse, M.F., 2012. Individual differences in skilled adult readers reveal dissociable patterns of neural activity associated with component processes of reading. Brain Lang 120, 360–371.

Widmann, A., Schröger, E., Tervaniemi, M., Pakarinen, S., Kujala, T., 2012. Mapping symbols to sounds: electrophysiological correlates of the impaired reading process in dyslexia. Front Psychol 3, 1–12.

Wilson, T.W., Leuthold, A.C., Lewis, S.M., Georgopoulos, A.P., Pardo, P.J., 2005. The time and space of lexicality: the neuromagnetic view. Exp Brain Res 167, 1–13.

Wolf, M., 1986. Rapid alternating stimulus naming in the developmental dyslexias. Brain Lang 27, 230–279.

Xia, Z., Hoeft, F., Zhang, L., Shu, H., 2016. Neuroanatomical anomalies of dyslexia: disambiguating the effects of disorder, performance, and maturation. Neuropsychologia 81, 68–78.

Yang, Y. H., Yang, Y., Chen, B. G., Zhang, Y. W., Bi, H. Y., 2016. Anomalous cerebellar anatomy in Chinese children with dyslexia. Frontiers in psychology 7, 324.

Zoubrinetzky, R., Bielle, F., Valdois, S., 2014. New Insights on Developmental Dyslexia Subtypes: Heterogeneity of Mixed Reading Profiles. Plos One 9, e99337.

Table 1. Neuropsychological tests and composites (bold).

Neuropsychological composites (bold) and individual tests	Reference
Phonological processing	
Pig Latin (accuracy)	Nevala *et al.*, 2006
Nonword span length (accuracy)	Laasonen, Service & Virsu, 2002
Rapid Alternate Stimulus naming (RAS) (speed of second trial)	Wolf, 1986
Reading, technical	
word list reading (speed and accuracy)	Nevala *et al.*, 2006
pseudoword list reading (speed and accuracy)	Nevala *et al.*, 2006
Reading comprehension (speed and accuracy)	Nevala *et al.*, 2006 *)
Intelligence Quotient	
Verbal IQ [Wechsler Adult Intelligent Scale (WAIS)-III Similarities and Vocabulary]	Wechsler, 2005
Performance IQ (WAIS-III Block design and Matrix reasoning)	Wechsler, 2005
Working memory functions	
WMS-III Letter-number series	Wechsler, 2008
WMS-III Visual Series	Wechsler, 2008

*) Not included in the technical reading composite score.

Table 2. Demographic, morphological, and neuropsychological data

Group sizes (n) and mean (median*) values of background variables in the Dyslexic and Control groups with standard deviation (interquartile range*) in parentheses. On the bottom, neuropsychological test results (full IQ and the composite scores) are presented. P-values show Chi Squared (χ^2), independent-samples t-test (t), and Mann-Whitney U-test (U) statistics for group comparisons. Non-normally distributed variables in one or both groups are marked with an asterisk (*).

	Dyslexic (*n* = 23)	Control (*n* = 22)	*P* value
Demographic			
Gender (male/female)	11/12	10/12	1.000 (χ^2)
Age (years)	31.3 (8.6)	29.8 (5.9)	.530 (*t*)
Education (years)	15.7 (5.2)	16.1 (4.4)	.817 (*t*)
Musical education (years)	3.0 (7.8)	3.7 (5.5)	.730 (*t*)
Morphological			
Grey matter volume (total)	0.8 (0.1)	0.8 (0.1)	.680 (*t*)
White matter volume (total)	0.5 (0.1)	0.5 (0.1)	.655 (*t*)
Cerebrospinal fluid volume (total)	0.6 (0.2)	0.5 (0.2)	.736 (*t*)
Total intracranial volume	1.8 (0.3)	1.8 (0.2)	.966 (*t*)
Total brain volume	1.3 (0.1)	1.3 (0.1)	.638 (*t*)
Neuropsychological			
Full IQ	104.6 (9.4)	117.3 (7.0)	.000 (*t*)
Phonological processing*	-0.2 (1.2)	0.5 (0.5)	.000 (*U*)
Technical reading*	-0.3 (1.0)	0.6 (0.2)	.000 (*U*)
Working memory	19.8 (4.8)	24.3 (4.8)	.003 (*t*)

Table 3. Focal grey and white matter volume associations with the neuropsychological composite scores.

Contrast	GMV/WMV	Area name	Coordinates	Cluster size	t-alue	Correlation
Technical reading	**GMV**	Left Superior Temporal Gyrus (BA 38)	-32 5 -21	2846	4.94*	R(39)=.641, P<.001
		Left Middle Temporal Gyrus (BA 21)	-48 7 -18			
		Left Fusiform Gyrus (BA 36)	-27 -5 -41			
		Left Insula (BA 13)	-36 10 -14			
		Left Amygdala	-25 -2 -21			
		Left Parahippocampal Gyrus (BA 34)	-16 -3 -20			
		Left Hippocampus	-16 -7 -14			
		Left Cerebellum	-36 -54 -34	4215	3.68*	R(39)=.499, P=.001
Technical reading	**WMV**	Right Putamen	29 -8 10	15095	5.24*	R(39)=.625, P<.001
		Right Insula	33 -14 9			
		Right Globus Pallidus	26 -15 -8			
		Right Inferior Frontal Gyrus	46 9 11			
		Right Precentral Gyrus	50 -9 24			
		Left Parahippocampal Gyrus	-18 -19 -14			
		Right Parahippocampal Gyrus	25 -19 -9			
		Pons	-14 -34 -29			
		Left Cerebellum	-17 -45 -33			
		Right Cerebellum	23 -52 -33	3142	3.88*	R(39)=.490, P=.001
Working memory	**WMV**	Brainstem	2 -31 -31	9857	4.69*	R(39)=.518, P=.001
		Left Cerebellum	-16 -42 -31			
		Right Cerebellum	26 -43 -31			

*p < 0.005 FWE-corrected at the cluster level

All results are thresholded at a whole-brain uncorrected p < 0.005 threshold at the voxel level with a minimal cluster size set to 100 voxels.

Correlations are partial correlations with 2-tailed p-value controlling for age, sex, TIV and FIQ.

BA, Brodmann area, GMV = grey matter volume, WMV = white matter volume

Table 4. Grey and white matter volume differences between the dyslexic and control groups.

Contrast	GMV/WMV	Area name	Coordinates	Cluster size	t-value	Correlation
Controls > Dyslexics	**GMV**	Left Hippocampus	-20 -8 -12	4866	6.09*	TR: R(39)=.431, P=.015
		Left Amygdala	-29 -3 -22			
		Left Claustrum	-33 1 -1			
		Left Insula (BA 13)	-36 13 -10			
		Left Putamen	-32 0 -4			
		Left Globus Pallidus	-17 -8 -5			
		Left Superior Temporal Gyrus (BA 34)	-40 9 -25			
		Left Inferior Frontal Gyrus (BA 44)	-51 12 4			
Controls > Dyslexics	**WMV**	Right Hippocampus	38 -10 -16	2245	4.53*	TR: R(39)=.426, P=.015
		Right Middle Temporal Gyrus	61 -33 -7			
Overlap of technical reading (one sample and t-test)	**GMV**	Left Insula (BA 13)	-33 0 -13	769		TR: R(39)=.577, P<.001
		Left Superior Temporal Gyrus (BA 38)	-32 4 -18			
		Left Amygdala	-20 -6 -13			
		Left Hippocampus	-19 -9 -13			

*p < 0.05 FWE-corrected at the cluster level

All results are thresholded at a whole-brain uncorrected p < 0.005 threshold at the voxel level with a minimal cluster size set to 100 voxels.

Correlations are partial correlations with 2-tailed p-value controlling for age, sex, TIV and FIQ.

BA, Brodmann area, GMV = grey matter volume, TR = Technical reading score, WMV = white matter volume

Figure legends

Figure 1.

Grey and white matter volume associations with reading-related skills (see also Table 3). Top: Grey matter volume and technical reading score; Middle: White matter volume and technical reading score; Bottom: White matter volume and working memory score. N = 45. Neurological convention is used with MNI coordinates at the bottom left of each slice. All statistical maps are thresholded at a cluster-level FWE-corrected p < 0.05 threshold. Grey and white matter volume correlations to reading-related skills are shown with scatter plots. AMY = amygdala, BS = brainstem, CER = cerebellum, FG = fusiform gyrus, GB = globus pallidus, GMV = grey matter volume, IFG = inferior frontal gyrus, L = left, PCG = precentral gyrus, PHG = parahippocampal gyrus, R = right, STG = superior temporal gyrus, WMV = white matter volume.

Figure 2.

Grey and white matter volume group differences (see also Table 4). Top: Controls > Dyslexics, grey matter volume; Middle: Controls > Dyslexics, white matter volume; Bottom: Overlap image (purple) of Controls > Dyslexics, grey matter volume (red) and grey matter volume and technical reading score correlation analysis (blue). N = 45. Neurological convention is used with MNI coordinates at the bottom left of each slice. All statistical maps are thresholded at a cluster-level FWE-corrected p < 0.05 threshold. Grey and white matter volume correlations to reading-related skills are shown with scatter plots. Bar plots for mean GMV in significant clusters (Table 4) are shown: bar = mean (centered), error-bar = standard error of mean. CLAU = claustrum, GB = globus pallidus, GMV = grey matter volume, HIP = hippocampus, IFG = inferior frontal gyrus, INS = insula, L = left, MTG = middle temporal gyrus, PUT = putamen, R = right, STG = superior temporal gyrus, WMV = white matter volume.

Controls > Dyslexics GMV

Controls > Dyslexics WMV

Cluster 1: Technical Reading score association with GMV
Cluster 2: Controls > Dyslexics GMV