Identification of a Novel PP2A Regulator, WNK1, as Critical For Uterine Function

Ru-pin Alicia Chi[1], Tianyuan Wang[2], Chou-Long Huang[3], San-pin Wu[1], Steven L. Young[4], John Lydon[5], and Francesco DeMayo[1*]

[1] Reproductive and Developmental Biology Laboratory, National Institute of Environmental Health Sciences, Durham, NC, 27709, USA

[2] Integrative Bioinformatics Support Group, National Institute of Environmental Health Sciences, Durham, NC, 27709, USA

[3] Department of Internal Medicine, University of Iowa Carver College of Medicine, Iowa, IA, 52242, USA

[4] Department of Obstetrics and Gynecology, University of North Carolina at Chapel Hill, Chapel Hill, NC, 27599, USA

[5] Department of Molecular and Cellular Biology, Baylor College of Medicine, Houston, TX, 77030, USA

[*] Correspondence: demayofj@niehs.nih.gov

Keywords:

WNK1, reproduction, implantation, AKT, PP2A, FOXO1, adenomyosis, hyperplasia

Abstract

WNK1 is critical for uterine function as a mediator of stromal cell decidualization *in vitro*.

Here, we employed a mouse model with conditional WNK1 ablation from the female

reproductive tract to define its *in vivo* role in uterine biology. Loss of WNK1 altered uterine

morphology, causing endometrial epithelial hyperplasia, adenomyosis and a delay in

embryo implantation, ultimately resulting in compromised fertility. Mechanistic

investigations through transcriptomic and proteomic approaches uncovered the

regulatory role of WNK1 in controlling the PP2A-AKT-FOXO1 signaling axis. We show

that WNK1 interacts directly with PPP2R1A, which is crucial for PP2A phosphatase

activity. PP2A phosphatase in turn dephosphorylates AKT, thereby reducing its inhibitory

effect on FOXO1. This permits the nuclear entry of FOXO1 to transcriptionally regulate

implantation-associated genes. Our findings revealed a novel function of WNK1 in

regulating AKT-FOXO1 post-translational modification, and demonstrated that this

signaling pathway is critical in normal uterine physiology and pregnancy.

35
36
37
38

Introduction

The ability of the uterus to support and maintain the development of an embryo is critically dependent upon the process of implantation, the aberrant occurrence of which causes a ripple effect leading to pregnancy complications and miscarriage[1]. Embryo implantation occurs during the "window of receptivity" and requires a fully prepared and receptive uterus. Investigation of uterine function mediators during receptivity identified with no lysine kinase (K) 1 (WNK1) as potentially functional in the uterus acting downstream of EGFR, whose inhibition impaired decidualization[2]. Functional analysis of human endometrial stromal cells *in vitro* demonstrated that WNK1 is needed for proliferation, migration and differentiation[3]. Collectively, these findings indicate a previously unrecognized function of WNK1 in the female reproductive tract and led us to hypothesize that WNK1 is a mediator of uterine function.

WNK1 belongs to a family of serine/threonine protein kinases[4, 5], with its name derived from the unusual placement of the catalytic lysine in subdomain I[6]. To date, Wnk1's function is the most extensively explored in the kidney and the nervous system due to the link between its mutation and familial hypertension and autonomic neuropathy[7-9]. In the renal system, WNK1 controls ion homeostasis through diverse mechanisms including activation of the SGK1/epithelial sodium channel pathway[10], regulating the potassium channel Kir1.1 cell surface localization[11], as well as controlling the activity of Na-K-Cl cotransporter through phosphorylating OSR1(OXSR1) and SPAK[12, 13]. Interestingly, WNK1's regulatory function on OSR1 is critical for cardiovascular development, thereby contributing to embryonic lethality when WNK1 is ablated from the endothelium[14, 15]. These findings suggest that although WNK1 exhibits organ-specific

physiological functions, the underlying cellular components regulated by WNK1 may share

similarity between the different tissues.

Despite its ubiquitous expression pattern, WNK1's role in organs other than those described

above remain unexplored. Given the role of WNK1 in regulating uterine stromal cell biology *in*

vitro, we hypothesized that WNK1 is essential in regulating uterine functions. To test this idea, we

established a conditional uterine WNK1 ablated mouse model, which resulted in severely

compromised fertility. We demonstrated that WNK1 is critical in maintaining uterine morphology,

regulating epithelial proliferation and permitting appropriate embryo implantation. Additionally, we

defined the molecular mechanism through both transcriptomic and proteomic approaches to show

that the WNK1 signaling cascade begins by its direct interaction with the scaffold subunit of the

phosphatase complex PP2A, PPP2R1A. This interaction stabilized PP2A which prevented

unregulated AKT phosphorylation, ultimately alleviating AKT's inhibitory effect on FOXO1 nuclear

localization, an indispensable mediator of implantation[16].

RESULTS

WNK1 is expressed in the uterus during early pregnancy in both humans and mice

WNK1 expression was examined by immunohistochemistry in human endometrium during the

proliferative and mid-secretory phases as well as in the peri-implantation uterus of mice. In

humans, WNK1 is expressed in both the epithelial and stromal cells during the proliferative and

mid-secretory phases (Fig. 1.A). Similarly in mice, WNK1 is expressed during and after

implantation on gestation days (GDs) 4.5 and 5.5 (Fig. 1.B). These findings support the *in vivo*

involvement of WNK1 in regulating functions of the female reproductive tract.

WNK1 ablation altered uterine morphology and microenvironment

To examine WNK1's function in the female reproductive tract, we established a mouse model with

WNK1 ablation in progesterone receptor (PGR) expressing cells, which comprised all

compartments of the uterus. The conditional *Wnk1* allele mice (Wnk1$^{f/f}$) were crossed to the

PGR^{Cre} mice[14, 17], and confirmed that in the PGR$^{Cre/+}$;Wnk1$^{f/f}$ offspring (Wnk1$^{d/d}$), Cre mediated

excision led to the removal of exon 2 (Fig. S1. A and S1.B). Analysis of protein expression in

uterine tissue via western blot confirmed reduced WNK1 protein levels in the Wnk1d/d mice (Fig.

S1.C), indicating that exon 2 excision led to reduced total protein levels.

We employed a tissue clearing technique to visualize uterine morphology in 3D during the window

of receptivity (Fig. 2.A and Videos 1 and 2)[18]. WNK1 deficiency substantially impacted uterine

morphology, exhibiting increased number of, as well as altered structure of the endometrial glands

(Fig. 2.A, Videos 1 and 2). Among the abnormalities observed in the Wnk1$^{d/d}$ uteri was the failure

of gland reorientation surrounding the embryo[19, 20]. Examination of uterine cross sections from

older mice (26 and 50 weeks) further demonstrated invasion of glands into the myometrium,

suggesting that WNK1 ablation caused adenomyosis (Fig. 2.B). This was supported by the

elevated expression of *Moesin* (*Msn*) in the Wnk1$^{d/d}$ uteri, a biomarker for adenomyosis in humans

(Fig. 2.C)[21]. Quantification of gland number and *Foxa2* gene expression showed significant

elevation in the Wnk1$^{d/d}$ uteri (Fig. 2.D and E), confirming the substantial increase in glandular

tissues. To examine whether the increased glands were a result of increased proliferation in the

uterus, we examined the expression of two mitotic markers - cyclin D1 (CCND1) and

phosphorylated histone H3 (H3S10p). Elevated levels of both proteins in the glandular epithelial

cells of the Wnk1$^{d/d}$ uteri was observed (Fig. 2.F). In addition to the increased CCND1 and H3S10p

in the glandular epithelium, higher expression of both proteins was also observed in the luminal

epithelium of Wnk1$^{d/d}$ uteri demonstrating that WNK1 ablation induced epithelial hyperplasia was

not restricted to the glands. Moreover, we observed increased extracellular matrix deposition

especially surrounding the glands in the Wnk1$^{d/d}$ uteri, as shown by Masson's trichrome staining

(Fig. 2.G). These results suggest that the adenomyotic phenotype could be associated with

increased epithelial proliferation as well as excessive extracellular matrix deposition[22, 23].

Uterine WNK1 ablation compromised fertility in mice through impaired implantation

A six month breeding trial was conducted to determine the impact of WNK1 ablation on female

fertility. Of the 8 control mice, 7 were able to complete the breeding trial, with 1 found dead

midtrial. Necropsy showed neither pregnancy nor abnormality associated with the reproductive

tract in this one mouse, indicating the cause of death was not related to abnormal uterine function.

The 7 mice produced 31 litters totaling 245 pups, which was equivalent to 4.4 litters and 35 pups

per mouse during the 6 months (Fig. 3.A and B). In contrast, only 4 of the 8 Wnk1$^{d/d}$ mice initiated

in the trial were able to complete the trial. This was due to 4 females succumbing to complications

during pregnancy or delivery. Of the 4 females that completed the trial, 10 litters and 18 pups were produced, which averaged to 2.5 litters and 4.5 pups per mouse (Fig. 3.A and B). The average litter size was also significantly smaller, averaging 3.3 pups per litter in the Wnk1$^{d/d}$ mice, compared to 7.7 in the control Wnk1$^{f/f}$ mice (Fig. 3.C). While the Wnk1$^{f/f}$ mice bred consistently, producing the last litters in the twentieth week of the trial, most of the Wnk1$^{d/d}$ mice stopped breeding after 3 litters less than 15 weeks into the trial, with only one mouse producing after the twentieth week (Fig. 3.D). Taken together, these results illustrated compromised ability to support pregnancy and progressive sterility with uterine loss of WNK1.

We next examined whether the subfertile phenotype was associated with an implantation defect in the Wnk1$^{d/d}$ mice. Dams were euthanized on GD 4.5, and embryo implantation was visualized by Evan's blue dye staining. As expected, 84.2% of the control mice successfully permitted embryo implantion on GD 4.5 while the remaining 15.8% had no embryos present in the uterus as indicated by uterine flushing (Fig. 3.E, top panel and Fig. 3.F). On the other hand, only 29.4% of the mated Wnk1$^{d/d}$ mice were able to form implantation sites on GD 4.5; however, 52.9% of the mice harboured fertilized embryos inside the uterus as identified by uterine flushing (Fig. 3.E, top panel and Fig. 3.F). Examination of the uterus on GD 5.5 showed 76.9% of the Wnk1$^{d/d}$ mice with implantation sites which is, at this time point, comparable to their Wnk1$^{f/f}$ control littermates (Fig. 3.E, bottom panel). Histological examination showed that the control Wnk1$^{f/f}$ mice had already degraded the epithelium, enabling the embryo to invade the underlying stroma (Fig. 3.G, left), while Wnk1$^{d/d}$ uteri had intact maternal epithelium at this time point with the embryo trapped inside the luminal space (Fig. 3.G). These findings demonstrate that the majority of the implantation events in the Wnk1$^{d/d}$ mice were delayed.

In addition to the uterine expression, PGR is also expressed in the ovaries and pituitary, hence

WNK1 was also ablated from those tissues. We next questioned whether the implantation

phenotype could be attributed to a dysfunction in the hapothalamic-pituitary-ovarian axis. Ovarian

function was evaluated by assaying ovulation and ovarian steroid hormone levels. The mice were

subjected to superovulatory regimen of donadotropins, followed to mating to wildtype male mice,

and euthanized on GD 1.5 to simultaneously monitor fertilization of the oocytes. We found no

significant decrease in the number of 2-cell embryos produced by the Wnk1$^{d/d}$ dams indicating

that ovulation and fertilization was not affected (Fig. S2.A). Additionally, serum estradiol (E$_2$) and

progesterone (P$_4$) levels were similar between the Wnk1$^{f/f}$ and WNK1$^{d/d}$ mice on GD 4.5 (Fig. S2.B

and C), demonstrating that the ovaries were able to produce and maintain hormone levels. These

results indicate that the main contributing factor for the delayed implantation was not a malfunction

of the hypothalamic-pituitary-ovarian axis.

Prerequisites for uterine receptivity are the production of leukemia inhibitory factor (LIF) from the

uterine glands and the cessation of epithelial proliferation prior to implantation (GD 3.5)[24, 25]; as

well as suppression of epithelial PGR expression and the nuclear localization of FOXO1 during

implantation (GD 4.5)[16, 26]. Hence, we examined the uterus to see whether the delayed

implantation was associated with impairment of the those parameters. *Lif* gene expression on GD

3.5 was similarly induced in the Wnk1$^{f/f}$ and Wnk1$^{d/d}$ mice, which decreased to basal level on GD

4.5 in the Wnk1$^{f/f}$ mice (Fig. 3.H). The Wnk1$^{d/d}$ mice showed significantly higher *Lif* levels on GD

4.5 (Fig. 3.H), possibly resulting from the increased glandular tissue. The proliferative markers Ki-

67, PGR and FOXO1 were all deregulated in the Wnk1$^{d/d}$ mice on GD 4.5. High levels of Ki-67

were observed in the luminal epithelium of Wnk1$^{d/d}$ uteri (Fig. 3.I), indicating failure to impede

epithelial proliferation. Increased PGR coupled to decreased nuclear FOXO1 and increased

cytoplasmic FOXO1 were simultaneously observed in the luminal epithelium, indicating aberrant

regulation of this signaling pathway. Moreover, the underlying stromal cells also exhibited reduced

nuclear FOXO1, suggesting that the cellular impact of WNK1 on FOXO1 localization may be

similar in both the epithelial and stromal compartments. Our findings illustrated that several crucial

implantation-associated molecular events were deregulated in the Wnk1$^{d/d}$ mice.

Delayed implantation negatively impacted pregnancy and embryo development in the

Wnk1$^{d/d}$ mice

Interestingly, of the 29.4% mated Wnk1$^{d/d}$ mice that were able to permit embryo implantation on

time (GD 4.5), the number of implantation sites were similar to their Wnk1$^{f/f}$ control littermates

(Fig. 4.A). However, the number of implantation sites present on GD 5.5 was significantly lower

in the Wnk1$^{d/d}$ mice (Fig. 4.B). This finding indicated that the delay in implantation is associated

with reduced number of implantation sites. Additionally, spacing between the implantation sites in

the Wnk1$^{d/d}$ mice were irregular whereas the implantation sites observed in the Wnk1$^{f/f}$ mice were

evenly distributed (Fig. 4.C). Interestingly, for the Wnk1$^{d/d}$ mice that were able to implant promptly,

implantation spacing was more evenly distributed (Fig. 3.F), suggesting that the delay impacted

both implantation numbers and spacing. Examination of the uterus and embryo during mid-

pregnancy (GD 8.5) further demonstrated that the Wnk1$^{d/d}$ mice carried either resorbed embryos

(Fig. 4.D, middle panel, blue arrows) or abnormally formed decidual balls (Fig. 4.D.right panel),

compared to the normally sized decidual balls observed in the control mice (Fig. 4.D, left panel).

Moreover, we also observed multiple embryos within one decidual zone (Fig. 4.D, red arrows),

possibly from the cluttered/delayed implantation. Morphology was evaluated by examining cross

sections through the center of the decidual ball, which showed that the Wnk1$^{f/f}$ mice have

vascularized and initiated placentation (Fig. 4.E, black arrows and dashed lines, respectively),

both of which were lacking in the Wnk1$^{d/d}$ uteri. Ultrasound scans demonstrated decreased

gestation sac size (Fig. 4.F and G) and decreased embryo size at both GD 8.5 and GD 10.5 (Fig.

4.F and H). By GD 12.5, embryo resorption was frequently observed in the Wnk1$^{d/d}$ mice (Fig.

4.F, bottom panel). Collectively, these findings demonstrate that uterine WNK1 ablation led to

abnormal implantation and negatively impacted embryo development, resulting in the

compromised pregnancy outcome and subfertility..

Loss of uterine WNK1 altered the transcriptome and kinase phosphorylation network

during the receptive phase

To fully characterize the molecular mechanisms underlying the loss of WNK1 induced-

implantation defect, we next examined global gene expression profile by RNA sequencing (RNA-

seq) in the uterus during receptivity. To ensure that the analysis was conducted only on the

maternal uterine tissues and not the embryos, we used vasectomized wild-type males to induce

pseudopregnancy in the Wnk1$^{f/f}$ and Wnk1$^{d/d}$ mice, which was confirmed by serum progesterone

levels on pseudopregnancy day (PPD) 4.5 (Table S1). In total, there were 14,423 and 14,337

genes expressed in the Wnk1$^{f/f}$ and Wnk1$^{d/d}$ uterus, respectively; of which 14,024 were expressed

in both. The transcriptomes were subjected to principle component analysis (PCA) as a measure

of quality control, which segregated according to genotype indicating that the samples were well-

characterized by genotype (Fig. S3). Using a defining threshold of *q*-value under 0.05 for

significance and fold change (FC) over 1.5 as differential expression, we identified 1,727

significantly and differentially expressed genes (DEGs) in the Wnk1$^{d/d}$ uterus during receptivity

(Table S2). We then conducted detailed analyses to characterize the molecular alterations

associated with uterine WNK1 ablation using the DAVID Bioinformatic Database and Ingenuity

Pathway Analysis (IPA). The top biological processes associated with the DEGs were adhesion,

cell movement and locomotion, inflammation and blood vessel development (Table S3). Many

important molecular functions associated with implantation were also deregulated in the Wnk1$^{d/d}$

uteri, such as cell proliferation and apoptosis, Notch signaling, cell differentiation, epithelial to

mesenchymal transition (EMT), cytokine production, and response to estrogen. Prediction of

upstream regulator activity further showed altered activity for many important receptivity

mediators, including the suppression of JAG, HEY2, PTEN and SERPINE1 (Fig. 5.A). On the

other hand, TGFB1, ERBB2, AKT, estrogen, ERK, MUC1 and KLF5 were predicted to show

increased activity (Fig. 5.A, for the complete list see Table S4).

We have earlier shown that there was a decrease in nuclear FOXO1 during the window of

implantation in the Wnk1$^{d/d}$ mice (Fig. 3.I). Since FOXO1 is a transcription factor that is critical for

implantation, its nuclear exclusion would likely results in its reduced activity in transcribing genes.

Therefore, we compared the DEGs to known FOXO1-controlled endometrial genes[16], and found

that of the 631 FOXO1-regulated endometrial genes, 313 showed altered expression in Wnk1$^{d/d}$

uterus (Fig. 5.B). Detailed comparison further illustrated that 90% of these common genes were

deregulated in the same direction under WNK1 and FOXO1 deficient conditions (Fig. 5.C). These

findings suggest that the impaired implantation observed in the Wnk1$^{d/d}$ mice was mediated

partially through insufficient FOXO1 signaling during receptivity.

As WNK1 is a kinase, we next examined the alterations in the kinase phosphorylation network to

understand the impact of WNK1 ablation. To this end, we employed an image-based

phosphokinase array to simultaneously evaluate the phosphorylation status of multiple kinases

in the uterus during receptivity (Fig. 5.D). Loss of WNK1 altered the phosphorylation of various

kinases, including TOR (mTOR), SRC, PRAS40 (AKT1S1), JNK, AMPKα1 (PRKAA1), GSK-3α/β

(GSK3A and GSK3B) and AKT (Fig. 5.E, all kinases with > 2 foldchange in phosphorylation are

shown in Fig. S4.A). The phosphorylation of AKT, GSK-3α/β and PRAS40 were independently

validated via western blotting (Fig. S4.B) and all showed elevated phosphorylation in Wnk1$^{d/d}$ uteri

during receptivity. Interestingly, AKT was identified as an activated upstream regulator by IPA,

and the phosphokinase array demonstrated its elevated phosphorylation in the Wnk1[d/d] uteri

during receptivity (Fig. 5.D, E and Fig. S4.B). Furthermore, elevated AKT phosphorylation on GD

4.5 was confirmed in both the epithelium and the stroma (Fig. 5.F). Indeed, we found that in the

control mice, phosphorylation of AKT was actively suppressed as the mice transitioned into the

receptive phase from GD 3.5 to PPD 4.5, however, the Wnk1[d/d] mice maintained high AKT

phosphorylation both prior to and during receptivity (Fig. 5.G). As AKT directly phosphorylates

FOXO1 leading to its nuclear exclusion[27, 28], and decreased nuclear FOXO1 was observed in the

Wnk1[d/d] uteri, we next examined FOXO1 phosphorylation. Western blot analysis showed a robust

induction in FOXO1 phosphorylation during receptivity in the Wnk1[d/d] mice, while total FOXO1

level remained constant (Fig. 5.H). These findings suggest that the elevated FOXO1

phosphorylation, and hence nuclear exclusion may be attributed to increased AKT

phosphorylation.

WNK1 ablation led to FOXO1 nuclear exclusion via AKT phosphorylation, which is

associated with decreased PP2A expression and activity

Having demonstrated that the loss of WNK1 led to increased phosphorylation of AKT and FOXO1

in mouse uteri, we next examined whether this regulatory axis was similarly maintained in human

endometrial HEC1A (epithelial) and THESC (stromal) cells. Using small interfering RNA against

WNK1 (siWNK1), WNK1 protein expression was inhibited, which robustly induced AKT and

FOXO1 phosphorylation in both cell lines (Fig. 6.A). In order to test whether AKT facilitated

FOXO1 nuclear exclusion downstream of WNK1, we next treated these cells with an AKT

inhibitor, GDC0941, and examined whether it could rescue WNK1 ablation-induced

phosphorylation and nuclear exclusion of FOXO1. FOXO1 localization clearly decreased in the

nucleus of both cells after transfection with siWNK1 (Fig. 6.B, panels 1 VS 2, and 4 VS 5).

However, when the siWNK1 transfected cells were treated with GDC0941, nuclear FOXO1 was

readily restored (Fig. 6.B, panels 3 and 6). This suggested that WNK1 inhibition-induced nuclear

exclusion of FOXO1 is mediated through AKT. This is further supported by the findings that AKT

inhibition rescued WNK1 knock-down induced FOXO1 phosphorylation (Fig. 6.C). Interestingly,

GDC0941 treatment reduced the phosphorylation of AKT and FOXO1 to a level that is lower than

seen in the siCTRL transfected, untreated cells (considered basal level). As GDC0941 inhibits

AKT through its upstream regulator PI3K[29], it is likely that PI3K lies upstream of WNK1 in

regulating AKT. Indeed, none of the PI3K family members were impacted by WNK1 inhibition,

including p110-α, p110-β, p110-γ, Tyr458 phosphorylated p85 and Tyr199 phosphorylated p55

(Fig. 6.C). Similarly in mice, the expression of these proteins in the uterus were comparable

between the Wnk1$^{f/f}$ and Wnk1$^{d/d}$ mice during receptivity (Fig. 6.D).

A search of the upstream regulators predicted by IPA identified candidate AKT regulators with

altered activities in the Wnk1$^{d/d}$ uteri, including PTEN, PPP2CA, and sirolimus (rapamycin, Table

S4). PTEN and PPP2CA are both phosphatases that regulate AKT phosphorylation, and both

displayed repressed activities in the Wnk1$^{d/d}$ mice during receptivity (Z-scores of -2.079 and -

1.195, respectively, Table S4). Sirolimus, on the other hand, is a drug targeting the kinase mTOR,

which was strongly inhibited (Z-score of -2.95, Table S4). We found that mTOR phosphorylation

and PP2A subunits A and C were altered in the Wnk1$^{d/d}$ mice, while PTEN level was not

significantly different (Fig. 6.E). This finding suggested that increased AKT phosphorylation in the

Wnk1$^{d/d}$ mice may be mediated through elevated mTOR or repressed PP2A activity. As mTOR is

both a regulator and a substrate of AKT[30, 31], we examined whether WNK1 ablation-induced AKT

phosphorylation is mediated through mTOR. We inhibited mTOR activity using rapamycin and

examined AKT/FOXO1 phosphorylation as well as FOXO1 localization as a readout of AKT

activity. As shown in Fig. S5.A, rapamycin treatment did not reverse the nuclear exclusion of

FOXO1 induced by WNK1 inhibition. Additionally, AKT and FOXO1 phosphorylation was not

rescued by rapamycin treatment (Fig. S5.B). Similar results were observed in HEC1A cells where

WNK1 and mTOR double knock-down failed to rescue AKT and FOXO1 phosphorylation (Fig.

S5.C). Thus, mTOR is likely not the WNK1 mediator controlling AKT activity, and its elevated

phosphorylation is a result of elevated AKT activity, rather than its cause.

WNK1 regulates AKT phosphorylation through direct interaction with PPP2R1A

WNK1 ablation led to impaired implantation arising from dysregulated AKT-FOXO1 signaling with

a concomitant repression of PP2A activity and reduced PP2A subunits A and C expression. We

explored the possible regulatory link between WNK1 and PP2A/AKT using a non-biased WNK1

immunoprecipitation-mass spectrometry (IP-MS) approach to identify WNK1 binding partners.

Successful WNK1 IP was confirmed by examining the lysate for WNK1 expression after

immunoprecipition using a a rabbit IgG (negative control) or WNK1 targeting antibody from

HEC1A cells (Fig. S6), and the peptides identified by mass-spectrometry are listed in Table S5.

Amongst those were peptides belonging to WNK1 itself, as well as a known WNK1 substrate,

oxidative stress responsive kinase 1 (OXSR1/OSR1)[12], confirming the validity of the pull-down

results (Table S5).

Putative WNK1 binding proteins identified in this experiment include Wnt regulators (OFD1 and

CCDC88C), chromosome modulating and DNA repair proteins (SMCA1, KIF11, FANCI, RAD50

and SLC25A5), proteins associated with the endoplasmic reticulum and ribosomal functions

(UGGT1, SEC23A, HYOU1, EMC1, AIFM1, HM13, SCFD1) as well as the mitochondria (AIFM1,

SLC25A5). Of particular interest were the components of protein phosphatase complexes PP2A

(PPP2R1A) and PP6 (PPP6R3), as both regulate AKT phosphorylation. Since the enzymatic

activity of PP2A was predicted by IPA as repressed in the Wnk1$^{d/d}$ mice during receptivity

(PPP2CA, Table S4), we postulated that these observations were associated with the direct

interaction of WNK1 and PPP2R1A, the alpha isoform of the scaffold subunit A of PP2A. In order

to confirm the interaction of WNK1 and PPP2R1A, a YFP-tagged WNK1 (c4161, Fig. S7) was

expressed in HEC1A cells, then immunoprecipitated using a YFP nanobody, followed by detection

for PPP2R1A in the pulldown. We first confirmed that c4161 transfection induced exogenous

WNK1 expression when compared to the control cells transfected with YFP only expressing

construct (cYFP, Fig. 7.A). WNK1 was subsequently detected in the lysate immunoprecipitated

for YFP (Fig. 7.B, upper panel), which co-immunoprecipitated with PPP2R1A (Fig. 7.B, middle

panel).

Having confirmed the interaction of WNK1 and PPP2R1A, we next explored the biological

implications of this interaction. The PP2A phosphatase complex is comprised of the scaffold

subunit A with 2 isoforms, the regulatory subunit B with 13 isoforms and the enzymatic subunit C

with 2 isoforms. As shown earlier, uterine WNK1 ablation led to decreased protein levels of

subunits A and C (Fig. 6.E), yet RNA-seq showed no alteration in transcription of the 4 genes

encoding these 2 subunits in the Wnk1$^{d/d}$ mice. It has been reported that the stability of the PP2A

subunits depends on their association with each other[32]. Hence, reduced subunit levels could be

an indication that the complexing capacity of the subunits were reduced in the absence of WNK1,

leading to their degradation. We therefore postulated that the WNK1-PPP2R1A interaction is

necessary for the association of the PP2A subunits. To test this idea, we examined the levels of

PPP2R1A, total PP2A subunit A and total PP2A subunit C in WNK1 knock-down HEC1A cells,

and accordingly found their reduced levels upon WNK1 inhibition (Fig. 7.C). Lastly, to validate

that PP2A mediates AKT/FOXO1 signaling, we inhibited PPP2R1A expression in HEC1A cells

using siRNA, and examined the components of the PP2A-AKT-FOXO1 signaling axis. As

expected, PPP2R1A knock-down caused a reduction in total subunits A and C of PP2A (Fig. 7.D).

Interestingly, AKT phosphorylation was selectively induced on threonine 308, but not serine 473

after PPP2R1A knock-down (Fig. 7.D). This nonetheless, translated to elevated FOXO1

phosphorylation, indicating that loss of PP2A activity-induced AKT phosphorylation on this residue

alone is sufficient to alter FOXO1 phosphorylation (Fig. 7.D). These findings confirmed that in

endometrial cells, WNK1 controls FOXO1 phosphorylation through directly interacting with

PPP2R1A, which leads to stabilization of the PP2A complex and induces AKT dephosphorylation

(Fig. 7.E). In the Wnk1$^{d/d}$ uterus, the loss of WNK1 destabilizes and reduces PP2A activity, and

this in turn, causes an elevation of AKT and subsequently FOXO1 phosphorylation, rendering

FOXO1 cytoplasmic. As such, implantation-mediating genes reliant on FOXO1 transcriptional

regulation becomes deregulated, ultimately resulting in impaired implantation in these mice.

Discussion

Reproductive biology has relied profoundly on transcriptomic analyses to identify novel players that may serve crucial functions in the regulation of fertility. While this approach has uncovered many key components in the reproductive tract, it is unable to detect alterations at the proteomic level, such as post translational modifications (PTMs). In many cases, the PTMs control protein activity and stability, and hence are the actual determinants of functional output. Using a proteomic approach, we identified WNK1 as a potential regulator of uterine biology with previously unreported reproductive functions. In this study, we examined the *in vivo* function of WNK1 using a whole-uterus WNK1 knock-out mouse model. We demonstrate that loss of WNK1 led to hyperplasia, adenomyosis and impaired implantation, which could all contribute to compromised fertility. Importantly, we demonstrate for the first time that WNK1 is a direct regulator of the PP2A-AKT-FOXO1 signaling pathway. As FOXO1 is critical for embryo implantation in mice[16], and the subfertile phenptype of Wnk1$^{d/d}$ mice is associated with implantation impairments, it is likely that aberrant FOXO1 localization is at least partially responsible for the subfertility. It is worth noting that the Wnk1$^{d/d}$ mice did not fully recapitulate the uterine FOXO1 knock-out phenotype, possibly due to decreased FOXO1 signaling, rather than complete inhibition.

Using a combination of *in vivo* and *in vitro* studies, we demonstrate that FOXO1 deregulation resulted from elevated AKT phosphorylation. This in turn was attributed to the reduced levels of PP2A subunits A and C, and hence reduced PP2A phosphatase activity. Interestingly, the RNA-seq results did not show altered transcription of *Ppp2ca*, *Ppp2cb*, *Ppp2r1a* and *Ppp2r1b,* which encode the different isoforms of subunits A and C. This suggests that the altered protein level was mediated through post-transcriptional or post-translational measures. As the half-life of the PP2A subunits is dependent on their ability to complex with each other[32], a possibility is that WNK1 facilitates the binding of the subunis, resulting in PP2A complex stabilization. Therefore,

in the absence of WNK1, the decreased coupling of the subunits to each other led to shorter half-

lives and decreased protein levels. It is worth noting that the antibody against PP2A subunit B

used in this study targets only one of the thirteen isoforms (PPP2R2A). While this particular

isoform was unaffected by loss of WNK1, it is possible that other subunit B isoforms may be

affected. Mechanistically, WNK1 directly interacts with PPP2R1A and hence this interaction may

be crucial for PP2A complex formation. Future studies will aim to understand the biochemical

nature of the PPP2R1A-WNK1 interaction, for example, whether WNK1 phosphorylates

PPP2R1A, and if so, whether the phosphorylation interferes with PP2A subunits coupling. As

WNK1, PP2A, AKT and FOXO1 are all ubiquitously expressed with crucial functions, this signaling

cascade is likely to have broad-spectrum functions in other tissues.

Whilst we were able to demonstrate the dysregulation of PP2A-AKT-FOXO1 signaling pathway

in the Wnk1[d/d] mice and the associated implantation defect, it is likely that the effects of WNK1

ablation on uterine biology is not limited to implantation. Given that 30% of the Wnk1[d/d] mice were

able to implant promptly with normal numbers of embryos, we rarely observed normal sized litters

from these mice. This suggests that even after a seemingly normal implantation, there are likely

other impairments in subsequent pregnancy development accounting for the compromised

fertility. Indeed, gene expression analysis indicated altered signaling in many receptivity

regulators including suppressed Notch signaling (HEY2 and JAG2)[33], elevated ERK[34] and MUC

signaling[35], which are all important decidualization mediators. Additionally, WNK1 is a major ion

channel regulator in renal cells[36], and ion channels are also significant players in endometrial

functions[37]. For example, the cAMP-activated anion channel cystic fibrosis transmembrane

conductance regulator (CFTR) and epithelial sodium channel (ENaC) are two ion channels

responsible for regulating uterine fluid flow[38, 39], and both are WNK1-regulated in other systems[10,

40]. Interestingly, there was a 2.7-fold increase in *Cftr* expression in the Wnk1[d/d] uteri during

implantation (Table S2), suggesting that WNK1 may regulate the expression of this gene's RNA

in addition to its reported regulatory function at the post-translational level. Moreover, WNK1 is

expressed in the placenta, indicating that WNK1 may also have placental functions and hence a

role in supporting embryo development during mid-pregnancy[41]. These findings offer additional

mechanisms through which WNK1 ablation could potentially impair fertility in mice, and will be

addressed in future studies to explore WNK1's alternative uterine functions.

As aforementioned, uterine WNK1 ablation exhibited pleiotropic effects, one which was

particularly apparent was the epithelial abnormality in the Wnk1$^{d/d}$ uteri. We observed epithelial

hyperplasia which manifested through abnormal gland growth and associated with increased

numbers of CCND1 and H3S10P positive cells. This was accompanied by adenomyosis which is

characterized by extension of the glands into the myometrium. Adenomyosis is known to reduce

implantation[42], which is associated with dysregulated adhesion molecules expression, including

ITGB3 and SPP1[43]. Interestingly, we found elevated *Spp1* expression in the Wnk1$^{d/d}$ uteri during

implantation, revealing another possible mechanism of impaired implantation in the Wnk1$^{d/d}$ mice.

In addition to implantation, adenomyosis negatively impact pregnancy in several other ways

including abnormal muscle contractility, inflammation and metabolism[42, 44, 45]. Interestingly, the

ERM protein family member moesin (encoded by the *MSN* gene) has been correlated with

adenomyosis in humans[21], and we accordingly found higher *Msn* expression in the Wnk1$^{d/d}$ mice.

This finding suggests that WNK1 ablation-induced adenomyosis may share molecular similarities

to this disease in humans. We propose that the uterine WNK1 ablation mouse model could be

employed to study this disease.

Although neither epithelial hyperplasia nor adenomyosis are cancerous, both are progressive

conditions which may lead to malignant transformation[46, 47]. Functional interpretation of the

transcriptome reiterated this, where many cancer development and progression associated

signaling pathways were altered in the Wnk1$^{d/d}$ uteri, including elevated TGFB, AKT and

estrogen[47-50]. Interestingly, a recurrent mutation of *Ppp2r1a* is associated with serous endometrial

carcinoma[51, 52], and this mutation has been found to impact oncogenic signaling through a

dominant negative effect[53]. It is possible that WNK1 could act to protect cells against cancer

progression through stabilizing PP2A subunits and hence activity. We speculate that WNK1

signaling in the uterus also acts to maintain tissue homeostasis, such as preventing unregulated

epithelial proliferation. As such, uterine loss of WNK1 displayed epithelial hyperplasia and may

promote cancer development with aging.

In summary, this study is a first to explore the reproductive function of WNK1 *in vivo*. We

demonstrate that WNK1 is critical in maintaining normal uterine morphology, mediating epitheial

homeostasis and implantation. Future studies will further explore the molecular actions of this

complex protein in female reproductive tract, as well as the significance in human fertility and

pathological outcomes when WNK1 is deregulated.

Online Methods

Ethics statement

This study was conducted according to the federal regulations regarding the use of human

subjects. Procedures were approved by the following ethics committee: Institutional Review

Board/Committee-A (IRB) of Greenville Health System under IRB file #Pro0000093 and

Pro00013885 and the University of Chapel Hill at North Carolina IRB under file #: 05-1757.

Written, informed consents were obtained from all patients prior to participation.

All animal studies were conducted in accordance with the Guide for the Care and Use of

Laboratory Animals, as published by the National Institute of Health. Animal protocols were

approved by the Animal Care and Use Committee (ACUC) of National Institute of Environmental

Health Sciences (protocol numbers 2015-0012 and 2015-0023). The mice were housed with a

maximum of 5 per cage with a 12-hour light and dark cycle, and fed irradiated Teklad global soy

protein-free extruded rodent diet (Harlem Laboratories, Inc., Indianapolis, IN) and fresh water ad

libitum. Euthanization was carried out by carbon dioxide inhalation followed by cervical

dislocation.

Generation of transgenic mice

The Wnk1$^{f/f}$ mice were a kind gift from Dr CL Huang (University of Iowa Healthcare), the

generation of which was described in a previous publication[14]. Briefly, the *Wnk1* floxed allele was

established by insertion of *loxP* sites into the 5' and 3' region of exon 2 of the mouse *Wnk1* gene.

The Wnk1$^{f/f}$ mice were crossed to mice carrying Cre under the control of the progesterone

receptor (PGR^{Cre}) to generate conditional uterine Wnk1 ablated mice (Wnk1$^{d/d}$, Fig. S1) [17].

Fertility trial

Seven-week old Wnk1$^{f/f}$ (control) and Wnk1$^{d/d}$ (experimental) mice were housed with wild-type

CD1 males for a period of 6 months, during which the mice were monitored daily for pregnancy

and delivery. Upon the first observation of delivery, the total number of pups including both live

and dead were recorded.

Implantation determination and pseudopregnancy

Mice of age 6-10 weeks were housed with wild-type C57BL/6J males, and monitored each

morning until vaginal plug is observed (indicating that mating has occurred). The first noon

following which the vaginal plug was seen was defined as gestation day (GD) 0.5. Mice were

anesthetized by isoflurane inhalation on GD 4.5 and 5.5, followed by retro-orbital administration

of 200 µL of 1% Evans Blue Dye prepared in phosphate buffered saline (PBS). Mice were then

euthanized, and the uterine horns harvested for implantation determination and imaging. For mice

sacrificed on GD 4.5, the uterine horns were flushed using a p200 pipette if no implantation sites

were observed, and the eluant was examined under brightfield microscope to determine presence

of blastocysts. The uterine horns were then fixed 48 hours in 4% paraformaldehyde (PFA)

prepared in PBS for histology and immunohistochemstry, or frozen on dry ice for RNA and protein

extraction. For RNA-seq, pseudopregnancy was induced by mating the females to vasectomized

wild-type male mice, and all procedures were conducted as described above except for the Evans

Blue Dye injection, as no implantations were expected.

Superovulation assay

Three-week old Wnk1$^{f/f}$ and Wnk1$^{d/d}$ mice were subjected to a superovulation regimen, which began with intraperitoneal administration of 5 IU of pregnant mare's serum gonadotropin (catalog no. 493-10-2.5, Lee Biosolutions), followed by 5 IU of human chorionic gonadotropin (catalog no. 869031, EMD Millipore) 48 hours later. Superovulated mice were placed with wild-type CD1 males overnight. Mating was confirmed by presence of vaginal plug the next morning (GD 0.5), and mice were euthanized on GD 1.5 followed by oviduct flushing. The number of embryos was determined by counting under a brightfield microscope.

Serum collection

On GD or pseudopregnancy day (PPD) 4.5, mice were anesthetized by intraperitoneal administration of Fetal Plus (1mg/10g body mass) and whole blood was collected via retro-orbital puncture. Blood was allowed to clot at room temperature for approximately 30 minutes, then centrifuged at 1000 X G for 10 minutes at 4^{O}C. The supernatant (serum) was moved into a fresh tube and stored at -80^{O}C until hormone testing. Hormone testing was conducted by the Ligand Core Laboratory of University of Virginia, Center for Research in Reproduction.

High frequency ultrasound imaging

On GDs 8.5, 10.5 and 12.5, high frequency ultrasound imaging was used to evaluate the uterus and embryo development. Dams were anesthetized by isoflurane inhalation and placed onto an electric heating pad to maintain body temperature. Abdominal hair was removed using depilatory cream (Nair$_{TM}$ Church & Dwight Co. Trenton, NJ), and eye lubricant was applied to prevent desiccation. Dams were manipulated into a supine position for the scan while heart rate and body

temperature were continuously monitored. Images were visualized and captured using the

VisualSonics VevoR 2100 Imaging System with a 550s scan head (Fujifilm VisualSonics Inc.,

Toronto, ON) at 55 megahertz. Each scanning session was limited to maximum 15 minutes, after

which the dams were monitored until full recovery.

Tissue processing, histology, immunohistochemical and immunofluorescence staining

For histology, immunohistochemistry and immunofluorescence, the tissues were similarly

processed as described below. After 48 hour fixation, tissues were placed into 70% ethanol for

a minimum of 48 hours. Tissues were then dehydrated and embedded in paraffin blocks and

sectioned to 5 μm thickness onto glass slides. Slides were heated at 60^OC for 10 minutes, followed

by 5 minutes cooling. Sections were deparaffinized by 3 serial incubations in Citrisolv clearing

agent (catalog no. 22-143-975, Thermo Fisher), followed by rehydration through decreasing % of

ethanol. For histology, sections were subjected to hematoxylin and eosin (H&E) and Masson's

trichrome staining, followed by dehydration through increasing % of ethanol, incubation in Citrisolv

and coverslipping. For immunohistochemistry, sections were subjected to antigen retrieval after

rehydration by boiling in the Vector Labs Antigen Unmasking Solution as per manufacturer's

instructions (H-3300, Vector Laboratories, Burlingame, CA, USA). Blocking of endogenous

peroxidase was performed by treating the sections with 3% hydrogen peroxide diluted in distilled

water for 10 minutes at room temperature. Tissues were blocked in 5% normal donkey serum

(NDS) for 60 minutes at room temperature, prior to overnight incubation with the primary antibody

at 4^OC. The slides were washed twice in PBS for a total of 10 minutes at room temperature and

secondary antibody diluted in 1% w/v bovine serum albumin (BSA) prepared in PBS was applied.

The ABC reagent was applied to tissue according to the manufacturer's instructions (Vector Labs

ABC PK-6100, Vector Laboratories). Signals were developed using the Vector Labs DAB

ImmPACT Staining Kit (Vector Labs SK-4105, Vector Laboratories). Finally, the tissues sections

were counterstained with hematoxylin and dehydrated through increasing ethanol concentration,

followed by Citrisolv incubation and coverslipping. For immunofluorescence, tissue sections were

subjected to antigen retrieval as described above. Tissues were blocked in 0.4% v/v Triton X-100,

1% BSA and 5% NDS for 30 minutes at room temperature followed by overnight incubation in

primary antibody prepared in 0.4% Triton X-100/PBS at 4^{O}C. Sections were washed 3 times 5

minutes in PBS and incubated with secondary antibodies diluted in 0.4% Triton X-100/PBS for 90

minutes at room temperature. Finally, the slides were washed 3 times 5 minutes in PBS, and

coverslipped in DAPI containing mounting medium (Vectorshield Hardset™ Antifade Mounting

Medium, catalog no. H-1400, Vector Laboratories). Details of antibodies used in this study are

provided in table S6.

RNA extraction and cDNA conversion

The frozen tissues were disrupted in TRIzol reagent (Thermo Fisher) by bead milling, followed by

2 aqueous phase separations using 1-Bromo-3-chloropane and chloroform. Pure ethanol was

added to the aqueous layer, and the RNA was extracted using the Qiagen RNEasy RNA mini

prep kit columns as per manufacturer's instructions (Qiagen, Valencia, CA). Resulting RNA

concentration and quality was determined using the NanoDrop ND-1000. cDNA was generated

by reverse transcription using the M-MLV Reverse Transcriptase (catalog number 28025013

Thermo Fisher) following the manufacturer's instructions.

qRT-PCR

qRT-PCR was performed using the SsoAdvancedTM Universal SYBR Green Supermix

(1725274, Bio-Rad) with the following primers (from 5' to 3', F = forward and R = reverse): *Wnk1*

– AGGCAGAGATTCAAAGAAGAGG (F) and CCCAGGAATCATAGAATCGAACA (R); *Msn* –

CCATGCCGAAGACGATCA (F) and CCAAACTTCCCTCAAACCAATAG (R); and *Foxa2* –

GAGACTTTGGGAGAGCTTTGAG (F) and GATCACTGTGGCCCATCTATTT (R). *Lif* expression

was determined using the Taqman Master Mix (Life Technologies) and Taqman probes (Applied

Biosystems). The Delta delta Ct values were calculated using 18S RNA control amplification

results to acquire the relative mRNA expression for each sample.

RNA-sequencing

For each mouse, 1 µg of total uterine RNA was sent to the NIH Intramural Sequencing Center to

create a libarary using the TruSeq RNA Kit (Illumina, San Diego, CA, USA) following the

manufacturer's instructions. The RNA libraries were sequenced with a HiSeq 2000 System

(Illumina). The raw RNA reads (75 nt, paired-end) were processed by filtering with average quality

score greater than 20. Reads that passed the initial processing were aligned to the mouse

reference genome (mm10; Genome Reference Consortium Mouse Build 38 from December

2011) using TopHat version 2.0.4 [54]. Expression values of RNA-seq were expressed as fragments

per kilobase of exon per million fragments (FPKM). Differential expression was calculated using

Cuffdiff function from Cufflinks version 2.2[55]. Transcripts with the average FPKM > 1 in at least

one group, *q*-value < 0.05 and at least 1.5-fold difference in FPKM were defined as differentially

expressed genes (DEGs). The data discussed in this publication have been deposited in NCBI's

Gene Expression Omnibus and are accessible through GEO Series accession number

GSE144802.

Human phospho-kinase antibody array

Site specific phosphorylation levels of 43 kinases were measured using the Human Phospho-

Kinase Array Kit (catalog no. ARY 003B, R&D Systems) according to the manufacturer's

instructions with the experimental design as described below. Pseudopregnancy was induced in

the Wnk1$^{f/f}$ and Wnk1$^{d/d}$ mice as previously described, and mice were euthanized on PPD 4.5.

Uterine tissues were frozen at -80^{O}C until ready to proceed. Lysate were extracted independently

from 6 mice per group by bead milling in the lysis buffer provided within the kit, and protein

concentrations were determined using the BCA Protein Assay Kit (catalog no. 23225, Pierce).

Equal amounts from each mouse were pooled in each group (to a total of 900 μg), and the

remaining steps followed the standard protocol of the kit. Signal intensity was quantified by

ImageJ[56].

Protein extraction from uterine tissues and protein expression analysis

Tissues were homogenized in RIPA Lysis and Extraction Buffer (Thermo Fisher) supplemented

with protease inhibitor cocktail (cOmplete Mini, EDTA-free, catalog no. 11836170001, Roche

Diagnostics) and phosphatase inhibitor cocktail (phosSTOP, catalog no. 4906837001, Roche

Diagnostics), followed by centrifugation at 10,000 X G for 10 minutes at 4^{O}C, and the supernatant

was moved into fresh eppendorf. Protein concentrations were measured using the BCA Protein

Assay Kit (Pierce). Heat denatured protein samples were resolved using 7.5%, 10% or gradient

4-20% Criterion Tris-HCl precast gels (Bio-Rad), followed by transferring using the Trans-Blot

Turbo Transfer System (Bio-Rad), as according to the manufacturer's instructions. PVDF and

nitrocellulose membranes were used for target proteins > 200 KDa and < 200 KDa, respectively.

After transfer, the membranes were blocked in 5% w/v non-fat milk or BSA prepared in Tris

buffered saline with 0.1% Tween-20 (TBST). Membranes were incubated with primary antibody

at 4^{O}C with shaking overnight, followed by three 10 minute washes in TBST the next morning.

Membranes were proceeded to secondary antibody incubation at room temperature for at least

one hour with shaking, and washed another 3 times in TBST. Depending on the expected signal

strength, different peroxidase chemiluminescent substrates were used: KPL LumiGLO^R (catalog

no. 546101, Seracare), Clarity Western ECL Substrate (catalog no. 1705060, Bio-Rad), and

Amersham ECL Prime Western Blotting Detection Reagent (catalog no. RPN2232, GE

Healthcare Life Sciences). Antibody sources and dilutions are summarized in supplemental table

6. For each western blot, GAPDH or β-tubulin were detected as the loading control, and in cases

where the target protein is in the same region as the loading control proteins, a duplicate gel was

ran and transferred in parallel. For each set of samples, a representing GAPDH or β-tubulin blot

is shown. Details of antibodies used in this study are provided in table S6.

Tissue clearing and three-dimensional reconstruction

Uterine tissues were fixed in 4% PFA for 16 hours, followed by 3 rinses in PBS. Tissues were

then incubated in hydrogel monomer solution AP40 (4% v/v acrylamide and 0.25% w/v VA-044 in

PBS) for 72 hours at 4$^\text{O}$C, protected from light. Oxygen was then removed from the samples using

a chamber connected to vacuum and nitrogen, followed by incubation at 37$^\text{O}$C for 3 hours to

initiate tissue-hydrogel hybridization. Hydrogel was removed from the tissues via 3 PBS washes,

and tissues were subsequently incubated in 8% SDS prepared in PBS for 7 days at 37$^\text{O}$C with

shaking, and the SDS solution replaced twice during incubation. The tissues were then washed

5 times one hour in PBS and blocked in 5% NDS prepared in PBS/triton X-100 with 0.01% of

sodium azide. The samples were then incubated in primary antibody prepared in antibody diluent

(2% v/v NDS, 0.01% w/v sodium azide in PBST) for 6 days at room temperature with constant

rotation, followed by 5 one hour washes in 0.1% v/v Triton in PBS (PBS-T). Secondary antibody

was similarly prepared in antibody diluent and incubated for another 6 days at room temperature

with constant rotation and protected from light, replacing antibody half way through incubation.

Finally, the samples were washed an additional 5 times one hour in PBS-T and incubated in

Refractive Index Matching Solution (80% w/v Histodenz (catalog no. D2158, Sigma-Aldrich)

prepared in 0.02M phosphate buffer, pH7.5 with 0.1% Tween-20 and 0.01% sodium azide,

refractive index = 1.46) for 1-3 days, and samples were mounted in fresh Reflective Index

Mounting Solution using a 1 mm deep iSpacer (www.sunjinlabs.com). Details of antibodies used

in this study are provided in table S6.

Cell culture

Human endometrial epithelial cell line HEC1A and telomerase-transformed human endometrial

stromal cells (THESC) were obtained from American Type Culture Collection (ATCC, Rockville,

MD, USA). HEC1A cells were cultured in McCoy's 5A modified medium (catalog no. 16600082,

Gibco) and the THSEC cells were maintained in DMEM/F12 (1:1) (catalog no. 11330-032, Gibco),

both supplemented with 10% fetal bovine serum (FBS, catalog no. 10437-028, Gibco) and 100

U/mL penicillin and 100 µg/mL streptomycin, unless otherwise stated.

siRNA transfection and drug treatments

Cells were transfected with siRNAs using the Lipofectamine RNAiMax transfection reagent

(catalog no. 13778150, Thermo Fisher) following the manufacturer's protocol. Cells were

transfected with 24 – 72 nM siRNA in transfection medium supplemented with 2% charcoal-

stripped FBS (catalog no. 12676-029, Gibco) for 24-48 hours before replacing with fresh growth

medium. Proteins were harvested from cells 72 hours after transfection unless otherwise stated.

The siRNAs used in this study were: nontargeting siRNA (siCTRL, catalog no. D-001810-10-20,

Dharmacon), *Wnk1* targeting siRNA (siWNK1, catalog no. L-005362-02-0005, Dharmacon), *Mtor*

targeting siRNA (simTOR, catalog no. L-003008-00-0005, Dharmacon), and *Ppp2r1a* targeting

siRNA (siPPP2R1A, catalog no. L-060647-00-0005, Dharmacon). AKT and mTOR inhibitors

GDC0941 and rapamycin (catalog no. S1065 and S1039, respectively, Selleckchem) were

dissolved in DMSO, and cells were treated with 5 µM GDC 0941 and 10 – 40 µM rapamycin for

24 hours, while the control cells received equivalent volumes of DMSO.

Immunofluorescence of cultured cells

Cells were cultured in 4-chambered coverglass (catalog no. 155382, Thermo Fisher) as described

above. Following transfection and/or drug treatment for the appropriate time period, cells were

rinsed in cold PBS, fixed in 4% PFA and permeabilized in 0.5% Triton X-100/PBS for 10 and 5

minutes, respectively, at room temperature. Cells were then incubated in blocking buffer (5% v/v

NDS, 0.2% v/v fish gelatin (catalog no. G7765, Sigma-Aldrich), 0.2% v/v Tween-20 in PBS) for

30 minutes at 37^{O}C. Primary antibody was diluted in blocking buffer and added to the cells for 60

minutes, followed by secondary antibody for another 60 minutes; both incubation steps were

performed at 37^{O}C in a humidified chamber. Finally, cells were rinsed 3 times with 0.2% Tween-

20/PBS and coverslipped using a DAPI containing mounting medium (Vectorshield Hardset^{TM}

Antifade Mounting Medium, catalog no. H-1400, Vector Laboratories). Details of antibodies used

in this study are provided in table S6.

WNK1 Immunoprecipitation Mass-spectrometry

HEC1A cells were grown to 70% confluency, followed by collection using trypsin. Cells were

washed 2 X in cold PBS, followed by resuspension in cell lysis buffer (50 mM Tris-HCl pH 7.5,

150 mM NaCl, 1 mM EDTA, 1% NP-40, 1% sodium deoxycholate, 0.1% SDS, with protease and

phosphatase inhibitors added fresh to 1 X). Cells were incubated on ice for 10 minutes, followed

by sonication on medium power (3 X 5 seconds). Lysate was centrifuged at 13,000 X G for 10

minutes at 4^{O}C. WNK1 targeting antibody was added at 1:100 to the supernatant, and incubated

with rotation at 4^{O}C overnight. Prewashed beads (50% protein A and 50% protein G, catalog

numbers 10002D and 10004D, respectively, Thermo Fisher) were added to the immunocomplex

and incubated for 30 minutes at room temperature with rotation. Beads were pelleted using a

magnetic separation rack, followed by 3 washes in lysis buffer. Beads were heated to $100^{O}C$ with

3 X SDS buffer (150 mM Tris-HCl pH 6.8, 6% SDS, 0.3% BPB, 30% glycerol, 3% B-

mercaptoethanol) for 5 minutes, before electrophoresis through a 7.5% Criterion Tris-HCl precast

gel (Bio-Rad). Gel regions were excised from the SDS-PAGE gel and minced, and digests were

performed with a ProGest robotic digester (Genomic Solutions) where the gel pieces were

destained by incubation in 25 mM ammonium bicarbonate with 50% acetonitrile (v/v) twice for a

total of 30 minutes. The gel pieces were dehydrated in acetonitrile, followed by drying under a

nitrogen stream, and further incubated with 250 ng trypsin (Promega) for 8 hours at $37^{O}C$. The

digests were collected, and peptides were re-extracted three times. The extractions were pooled

for each sample, lyophilized and resuspended in 20 µL 0.1% formic acid. The protein digests were

analyzed by LC/MS on a Q Exactive Plus mass spectrometer (Thermo Fisher) interfaced with a

nanoAcquity UPLC system (Waters Corporation), and equipped with a 75 µm x 150 mm BEH

dC18 column (1.8 µm particle, Waters Corporation) and a C18 trapping column (18 µm x 20 mm)

with a 5 µm particle size at a flow rate of 400 nL/min. The trapping column was positioned in-line

of the analytical column and upstream of a micro-tee union which was used both as a vent for

trapping and as a liquid junction. Trapping was performed using the initial solvent composition. A

volumn of 5 µL of digested sample was injected into the column, and peptides were eluted by

using a linear gradient from 99% solvent A (0.1% formic acid in water (v/v)) and 1% solvent B

(0.1%formic acid in acetonitrile (v/v)), to 40% solvent B over 60 minutes. For the mass

spectrometry, a data dependent acquisition method was employed with an exclusion time of 15

seconds and an exclusion of +1 charge states. The mass spectrometer was equipped with a

NanoFlex source and was used in the positive ion mode. Instrument parameters were as follows:

sheath gas, 0; auxiliary gas, 0; sweep gas, 0; spray voltage, 2.7 kV; capillary temperature, $275^{O}C$;

S-lens, 60; scan range (m/z) of 200 to 2000; 2 m/z isolation window; resolution: 70,000; automated

gain control (AGC), 2 X 10^5 ions; and a maximum IT of 200 ms. Mass calibration was performed

before data acquisition using the Pierce LTQ Velos Positive Ion Calibration mixture (Thermo

Fisher). Peak lists were generated from the LC/MS data using Mascot Distiller (Matrix Science)

and the resulting peak lists were searched using the Spectrum Mill software package (Agilent)

against the SwissProt database. Searches were performed using trypsin specificity and allowed

for one missed cleavage and variable methionine oxidation. Mass tolerance were 20 ppm for MS

scans and 50 ppm for MSMS scans.

Generation of mammalian YFP-WNK1 expression constructs

The coding region of the WNK1 sequence (NM_014823.3) with attL sites and N-terminal TEV

cleavage site was synthesized by GeneWiz Inc. and cloned into pUC57 (Kanamycin) plasmid.

Gateway Cloning using LR Clonase II mix (Thermo Fisher) was used to transfer the WNK1

sequence into the Vivid Colors pcDNA6.2/N-YFP vectors (Thermo Fisher), which created the

mammalian expression vectors with YFP fused to the N-terminal end of WNK1 (Fig. S7, c4161).

Co-Immunoprecipitation

HEC1A cells were transfected with cYFP or c4161 for 48 hours, followed by trypsinization, 3

washes and resuspension in lysis buffer (50 mM Tris pH8.0, 400 mM NaCl, 0.1% NP-40 and 0.5

mM DTT, with protease and phosphatase inhibitors freshly added to 1 X). The lysate was

incubated at 4^OC with rotation for 30 minutes. Lysates were centrifuged at 21,100 X G for 10

minutes at 4^OC, and the supernatant was added to 1.5 volumes of 25% glycerol, followed by

centrifugation at 21,100 X G for 10 minutes at 4^OC. Anti-GFP resin slurry was added to the

supernatant and nutated for 1 hour at 4^OC. Beads were centrifuged at 1,000 X G for 5 minutes,

4^OC, followed by 6 washes in 100 µL of PBST in Bio-Spin columns (catalog number 7326204,

Bio-Rad). The bound immunocomplexes were eluted via 0.1 M glycine, pH 2.0, and eluent was

neutralized using 2M Tris-HCl, pH 8.0.

Confocal Microscopy

All fluorescent images presented in this study were captured using the Zeiss LSM 780 UV

confocal microscope.

Data analysis

GraphPad Prism versions 7 and 8 were used for data analysis. Each set of data points were first

subjected for normality tests. Student's t tests and Mann-Whiteney tests were performed for

normally distributed data and non-normally distributed data, respectively. In each case, a p-value

less than 0.05 was considered as significant. Functional annotation for the differentially expressed

genes derived from RNA-seq were analyzed by Ingenuity Pathway Analysis (IPA) and Database

for Annotation, Visualization, and Integrated Discovery (DAVID)[57].

Acknowledgements

This work was supported in part by Intramural Research Program of the National Institute of Health (Z1AES103311-01 (F.J.D.)); the Eunice Kennedy Shriver National Institute of Child Health & Human Development (RO1 HD042311 (J.P.L)); and National Institute of Diabetes and Digestive and Kidney Diseases (RO1 DK111542 (C.-L.H)). The authors thank Dr Sheng Song for guidance on the CLARITY technique; Dr Nyssa Adams for conducting the initial breeding trial, Dr Carmen Williams and Dr Sophia Tsai for reviewing the manuscript. We appreciate support from the NIEHS animal facility, Knockout Mouse Core, Digital Imaging Core, the Epigenomics and DNA Sequencing Core, the Fluorescent Microscopy and Imaging Core, the Mass Spectrometry Research and Support Group, the Structural Biology Core of NIEHS for their support and guidance with specialized techniques, as well as the Ligand Assay and Analysis Core at the University of Virginia.

Author Contributions

Conceptualization, R.A.C., S.P.W. and F.J.D.; Methodology, R.A.C. and F.J.D.; Validation, R.A.C.; Formal Analysis, R.A.C. and T.W.; Investigation, R.A.C.; Resources, S.L.Y., J.L., C.L.H. and F.J.D.; Data Curation, R.A.C. and T.W.; Writing – Original Draft, R.A.C.; Writing – Review & Editing, R.A.C. and F.J.D.;Visualization, R.A.C.; Supervision, F.J.D.; Project Administration, R.A.C. and F.J.D.; Funding Acquisition, F.J.D.

773 **Declaration of Interests**

774 The authors declare no competing interests.

References

1. Wang, H. & Dey, S.K. Roadmap to embryo implantation: clues from mouse models. *Nature reviews. Genetics* **7**, 185-199 (2006).

2. Large, M.J. *et al.* The epidermal growth factor receptor critically regulates endometrial function during early pregnancy. *PLoS genetics* **10**, e1004451 (2014).

3. Adams, N.R. *et al.* WNK lysine deficient protein kinase 1 regulates human endometrial stromal cell decidualization, proliferation, and migration in part through mitogen-activated protein kinase 7. *Biology of reproduction* **97**, 400-412 (2017).

4. Moore, T.M. *et al.* PSK, a novel STE20-like kinase derived from prostatic carcinoma that activates the c-Jun N-terminal kinase mitogen-activated protein kinase pathway and regulates actin cytoskeletal organization. *The Journal of biological chemistry* **275**, 4311-4322 (2000).

5. Nagase, T. *et al.* Prediction of the coding sequences of unidentified human genes. VII. The complete sequences of 100 new cDNA clones from brain which can code for large proteins in vitro. *DNA research : an international journal for rapid publication of reports on genes and genomes* **4**, 141-150 (1997).

6. Xu, B. *et al.* WNK1, a novel mammalian serine/threonine protein kinase lacking the catalytic lysine in subdomain II. *The Journal of biological chemistry* **275**, 16795-16801 (2000).

7. Wilson, F.H. *et al.* Human hypertension caused by mutations in WNK kinases. *Science (New York, N.Y.)* **293**, 1107-1112 (2001).

8. Disse-Nicodème, S. *et al.* A new locus on chromosome 12p13.3 for pseudohypoaldosteronism type II, an autosomal dominant form of hypertension. *American journal of human genetics* **67**, 302-310 (2000).

9. Lafreniere, R.G. *et al.* Identification of a novel gene (HSN2) causing hereditary sensory and autonomic neuropathy type II through the Study of Canadian Genetic Isolates. *American journal of human genetics* **74**, 1064-1073 (2004).

10. Xu, B.E. *et al.* WNK1 activates SGK1 to regulate the epithelial sodium channel. *Proceedings of the National Academy of Sciences of the United States of America* **102**, 10315-10320 (2005).

11. Wade, J.B. *et al.* WNK1 kinase isoform switch regulates renal potassium excretion. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 8558-8563 (2006).

12. Anselmo, A.N. *et al.* WNK1 and OSR1 regulate the Na+, K+, 2Cl- cotransporter in HeLa cells. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 10883-10888 (2006).

13. Vitari, A.C., Deak, M., Morrice, N.A. & Alessi, D.R. The WNK1 and WNK4 protein kinases that are mutated in Gordon's hypertension syndrome phosphorylate and activate SPAK and OSR1 protein kinases. *The Biochemical journal* **391**, 17-24 (2005).

14. Xie, J. *et al.* Endothelial-specific expression of WNK1 kinase is essential for angiogenesis and heart development in mice. *The American journal of pathology* **175**, 1315-1327 (2009).

15. Xie, J., Yoon, J., Yang, S.S., Lin, S.H. & Huang, C.L. WNK1 protein kinase regulates embryonic cardiovascular development through the OSR1 signaling cascade. *The Journal of biological chemistry* **288**, 8566-8574 (2013).

16. Vasquez, Y.M. *et al.* FOXO1 regulates uterine epithelial integrity and progesterone receptor expression critical for embryo implantation. *PLoS genetics* **14**, e1007787 (2018).

17. Soyal, S.M. *et al.* Cre-mediated recombination in cell lineages that express the
progesterone receptor. *Genesis (New York, N.Y. : 2000)* **41**, 58-66 (2005).
18. Yang, B. *et al.* Single-cell phenotyping within transparent intact tissue through whole-
body clearing. *Cell* **158**, 945-958 (2014).
19. Yuan, J. *et al.* Tridimensional visualization reveals direct communication between the
embryo and glands critical for implantation. *Nature communications* **9**, 603 (2018).
20. Yuan, J. *et al.* Planar cell polarity signaling in the uterus directs appropriate positioning
of the crypt for embryo implantation. *Proceedings of the National Academy of Sciences*
of the United States of America **113**, E8079-e8088 (2016).
21. Ohara, R. *et al.* Moesin overexpression is a unique biomarker of adenomyosis.
Pathology international **64**, 115-122 (2014).
22. Frantz, C., Stewart, K.M. & Weaver, V.M. The extracellular matrix at a glance. *Journal of*
cell science **123**, 4195-4200 (2010).
23. Kishi, Y. *et al.* Phenotypic characterization of adenomyosis occurring at the inner and
outer myometrium. *PloS one* **12**, e0189522 (2017).
24. Stewart, C.L. *et al.* Blastocyst implantation depends on maternal expression of
leukaemia inhibitory factor. *Nature* **359**, 76-79 (1992).
25. Dey, S.K. *et al.* Molecular cues to implantation. *Endocrine reviews* **25**, 341-373 (2004).
26. Wetendorf, M. *et al.* Decreased epithelial progesterone receptor A at the window of
receptivity is required for preparation of the endometrium for embryo attachment. *Biology*
of reproduction **96**, 313-326 (2017).
27. Brunet, A. *et al.* Akt promotes cell survival by phosphorylating and inhibiting a Forkhead
transcription factor. *Cell* **96**, 857-868 (1999).
28. Tang, E.D., Nunez, G., Barr, F.G. & Guan, K.L. Negative regulation of the forkhead
transcription factor FKHR by Akt. *The Journal of biological chemistry* **274**, 16741-16746
(1999).
29. Folkes, A.J. *et al.* The identification of 2-(1H-indazol-4-yl)-6-(4-methanesulfonyl-
piperazin-1-ylmethyl)-4-morpholin-4-yl-t hieno[3,2-d]pyrimidine (GDC-0941) as a potent,
selective, orally bioavailable inhibitor of class I PI3 kinase for the treatment of cancer.
Journal of medicinal chemistry **51**, 5522-5532 (2008).
30. Malley, C.O. & Pidgeon, G.P. The mTOR pathway in obesity driven gastrointestinal
cancers: Potential targets and clinical trials. *BBA clinical* **5**, 29-40 (2016).
31. Dan, H.C. *et al.* Akt-dependent activation of mTORC1 complex involves phosphorylation
of mTOR (mammalian target of rapamycin) by IkappaB kinase alpha (IKKalpha). *The*
Journal of biological chemistry **289**, 25227-25240 (2014).
32. Seshacharyulu, P., Pandey, P., Datta, K. & Batra, S.K. Phosphatase: PP2A structural
importance, regulation and its aberrant expression in cancer. *Cancer letters* **335**, 9-18
(2013).
33. Afshar, Y. *et al.* Notch1 mediates uterine stromal differentiation and is critical for
complete decidualization in the mouse. *FASEB journal : official publication of the*
Federation of American Societies for Experimental Biology **26**, 282-294 (2012).
34. Lee, C.H. *et al.* Extracellular signal-regulated kinase 1/2 signaling pathway is required
for endometrial decidualization in mice and human. *PloS one* **8**, e75282 (2013).
35. Zhang, L., Patterson, A.L., Zhang, L., Teixeira, J.M. & Pru, J.K. Endometrial stromal
beta-catenin is required for steroid-dependent mesenchymal-epithelial cross talk and
decidualization. *Reproductive biology and endocrinology : RB&E* **10**, 75 (2012).
36. Shekarabi, M. *et al.* WNK Kinase Signaling in Ion Homeostasis and Human Disease.
Cell metabolism **25**, 285-299 (2017).
37. Ruan, Y.C., Chen, H. & Chan, H.C. Ion channels in the endometrium: regulation of
endometrial receptivity and embryo implantation. *Human reproduction update* **20**, 517-
529 (2014).

38.   Chan, L.N. *et al.* Inhibition of amiloride-sensitive Na(+) absorption by activation of CFTR
in mouse endometrial epithelium. *Pflugers Archiv : European journal of physiology* **443**
Suppl 1, S132-136 (2001).
39.   Yang, J.Z. *et al.* Differential expression and localization of CFTR and ENaC in mouse
endometrium during pre-implantation. *Cell biology international* **28**, 433-439 (2004).
40.   Yang, C.L. *et al.* WNK1 and WNK4 modulate CFTR activity. *Biochemical and biophysical*
research communications **353**, 535-540 (2007).
41.   O'Reilly, M., Marshall, E., Speirs, H.J. & Brown, R.W. WNK1, a gene within a novel
blood pressure control pathway, tissue-specifically generates radically different isoforms
with and without a kinase domain. *Journal of the American Society of Nephrology :*
JASN **14**, 2447-2456 (2003).
42.   Harada, T. *et al.* The Impact of Adenomyosis on Women's Fertility. *Obstetrical &*
gynecological survey **71**, 557-568 (2016).
43.   Xiao, Y. *et al.* Expression of integrin beta3 and osteopontin in the eutopic endometrium
of adenomyosis during the implantation window. *European journal of obstetrics,*
gynecology, and reproductive biology **170**, 419-422 (2013).
44.   Vannuccini, S. *et al.* Pathogenesis of adenomyosis: an update on molecular
mechanisms. *Reproductive biomedicine online* **35**, 592-601 (2017).
45.   Garavaglia, E. *et al.* Adenomyosis and its impact on women fertility. *Iranian journal of*
reproductive medicine **13**, 327-336 (2015).
46.   Yeh, C.C., Su, F.H., Tzeng, C.R., Muo, C.H. & Wang, W.C. Women with adenomyosis
are at higher risks of endometrial and thyroid cancers: A population-based historical
cohort study. *PloS one* **13**, e0194011 (2018).
47.   Sobczuk, K. & Sobczuk, A. New classification system of endometrial hyperplasia WHO
2014 and its clinical implications. *Przeglad menopauzalny = Menopause review* **16**, 107-
111 (2017).
48.   Daikoku, T. *et al.* Conditional loss of uterine Pten unfailingly and rapidly induces
endometrial cancer in mice. *Cancer research* **68**, 5619-5627 (2008).
49.   Kriseman, M. *et al.* Uterine double-conditional inactivation of Smad2 and Smad3 in mice
causes endometrial dysregulation, infertility, and uterine cancer. *Proceedings of the*
National Academy of Sciences of the United States of America **116**, 3873-3882 (2019).
50.   Pavlidou, A. & Vlahos, N.F. Molecular alterations of PI3K/Akt/mTOR pathway: a
therapeutic target in endometrial cancer. *TheScientificWorldJournal* **2014**, 709736
(2014).
51.   Kuhn, E. *et al.* Identification of molecular pathway aberrations in uterine serous
carcinoma by genome-wide analyses. *Journal of the National Cancer Institute* **104**,
1503-1513 (2012).
52.   Zhao, S. *et al.* Landscape of somatic single-nucleotide and copy-number mutations in
uterine serous carcinoma. *Proceedings of the National Academy of Sciences of the*
United States of America **110**, 2916-2921 (2013).
53.   Haesen, D. *et al.* Recurrent PPP2R1A Mutations in Uterine Cancer Act through a
Dominant-Negative Mechanism to Promote Malignant Cell Growth. *Cancer research* **76**,
5719-5731 (2016).
54.   Trapnell, C., Pachter, L. & Salzberg, S.L. TopHat: discovering splice junctions with RNA-
Seq. *Bioinformatics (Oxford, England)* **25**, 1105-1111 (2009).
55.   Trapnell, C. *et al.* Transcript assembly and quantification by RNA-Seq reveals
unannotated transcripts and isoform switching during cell differentiation. *Nature*
biotechnology **28**, 511-515 (2010).
56.   Schneider, C.A., Rasband, W.S. & Eliceiri, K.W. NIH Image to ImageJ: 25 years of
image analysis. *Nature methods* **9**, 671-675 (2012).

57.   Huang da, W., Sherman, B.T. & Lempicki, R.A. Systematic and integrative analysis of
large gene lists using DAVID bioinformatics resources. *Nature protocols* **4**, 44-57 (2009).

FIGURES & FIGURE LEGENDS

933 **Figure 1. WNK1 is expressed in the uterus during the window of implantation in both**

934 **humans and mice.** (A and B) Immunohistochemical staining of WNK1 in proliferative and mid-

935 secretory phased endometrial tissues from fertile women (A), and during receptive gestation day

936 (GD) 4.5 and post implantation GD 5.5 in the uterus of wild-type mice (B). IgG served as negative

937 controls, scale bars = 50 μm.

Figure 2. WNK1 ablation altered uterine morphology and microenvironment. (A) Three-dimensional reconstruction of Wnk1$^{f/f}$ and Wnk1$^{d/d}$ uteri on GD 4.5 using tissue clearing and confocal microscopy. The glands, myometrium and embryo were marked by FOXA2 (green), ACTA2 (red) and OCT4 (purple), respectively. Images were captured by tile-scanning and Z-stacking, and reassembled *in silico* using Imaris software. White arrow indicates position of the embryo. Scale bars = 500 μm. (B) Immunofluorescence of uterine cross section showing glands (FOXA2, green) and myometrium (ACTA2, red) from Wnk1$^{f/f}$ and Wnk1$^{d/d}$ uteri. White arrows indicate gland extension into myometrium. Scale bars = 50 μm. (C) Adenomyosis biomarker *Msn* mRNA expression as determined by qRT-PCR (n = 4). (D) Quantification of number of glands per cross section for Wnk1$^{f/f}$ and Wnk1$^{d/d}$ mice (n = 6), and (E) quantification of *Foxa2* mRNA expression as determined by qRT-PCR (n = 6). (F) Expression of mitotic markers CCND1 and H3S10P in the uterus of 26-week-old Wnk1$^{f/f}$ and Wnk1$^{d/d}$ mice, scale bars = 100 μm. (G) Masson's trichrome staining of uteri cross section from 26 and 50 week old Wnk1$^{f/f}$ and WNk1$^{d/d}$ mice, scale bars = 100 μm. All quantitative results shown are average ± SD, * *p* < 0.05.

Figure 3. Uterine WNK1 ablation compromised fertility and impaired implantation in mice.

(A) Numbers of litters and pups produced by Wnk1$^{f/f}$ and Wnk1$^{d/d}$ dams during the 6-month

breeding trial with wild-type males. (B) Average number of litters produced, (C) average number

of pups per litter produced and (D) time of last delivery, n = 7 for Wnk1$^{f/f}$ mice and n = 4 for Wnk1$^{d/d}$

mice. Results shown are mean ± SD, $*$ $p < 0.05$. (E) Percentage of mated Wnk1$^{f/f}$ and Wnk1$^{d/d}$

mice with implantation (green), without implantation (pink), and without implantation but presented

embryos in the uterus (blue) on GD 4.5 and GD 5.5, n = 19 and 12 for Wnk1$^{f/f}$ mice on GD 4.5

and GD 5.5, respectively; and n = 17 and 13 for Wnk1$^{d/d}$ mice on GD 4.5 and GD 5.5, respectively.

Red arrows indicate position of implantation sites. (F) Gross uterine morphology of Wnk1$^{f/f}$ and

Wnk1$^{d/d}$ mice on GD 4.5, scale bars = 1 cm. (G) Hematoxylin and eosin staining of uterine cross

sections at implantation site on GD 5.5 in Wnk1$^{f/f}$ and Wnk1$^{d/d}$ uteri, arrow indicates presence of

maternal epithelium, and E = embryo. Scale bars = 100 µm. (H) Implantation marker *Lif* mRNA

expression in the uteri as determined by qRT-PCR on GD 3.5 and GD 4.5 for Wnk1$^{f/f}$ and Wnk1$^{d/d}$

mice. Results shown are mean ± SD, $*$ $p < 0.05$. (I) Expression of proliferative marker Ki-67,

implantation markers PGR and FOXO1 on GD 4.5 in the stroma and epithelium of Wnk1$^{f/f}$ and

Wnk1$^{d/d}$ mice. LE = luminal epithelium and S = stroma, scale bars = 25 µm.

Figure 4. Delayed implantation negatively impacted pregnancy and embryo development

in the Wnk1$^{d/d}$ mice. (A – B) Number of implantation sites on GD 4.5 (A) and GD 5.5 (B) in Wnk1$^{f/f}$

and Wnk1$^{d/d}$ mice (n = 6). (C – D) Uterine gross morphology on GD 5.5 (C) and GD 8.5 (D), with

implantation sites on GD 5.5 marked by red arrows, and blue arrows indicate resorption and

abnormal decidualization on GD 8.5. Scale bars = 1 cm. (E) Hematoxylin and eosin staining of

cross section through the centre of decidual ball on GD 8.5 from Wnk1$^{f/f}$ and Wnk1$^{d/d}$ mice, with

black arrows and dashed line indicating vascularization and placentation, respectively. Scale bars

= 2 mm. (F) Ultrasound scans of uterus and embryo during mid-pregnancy at GD 8.5, 10.5 and

12.5. Scale bars = 2 mm. (G and H) Quantification of gestation sac size by length and width on

GD 8.5 (G), and embryo size by crown-rump length (CRL) on GD 8.5 and 10.5, as measured from

ultrasound scans (n = 6). Results shown are mean ± SD, $^*p < 0.05$.

A Activation Z-score

JAG2, HEY2, PTEN, SERPINE1, TGFB1, ERBB2, AKT, estrogen, ERK, MUC1, KLF5

B WNK1 RNA-seq 1,727 DEG — 313 — FOXO1 RNA-seq 631 DEG

C 90%

No. of genes

49% — 153 genes WNK1↑FOXO1↑
41% — 129 genes WNK1↓FOXO1↓
13 genes WNK1↑FOXO1↓
18 genes WNK1↓FOXO1↑

D Wnk1^f/f / Wnk1^d/d

E

No	Protein	Site	Change in Wnk1d/d mice	Fold change
1	TOR	S2448	Up	2.05
2	PRAS40	T246	Up	2.44
3	JNK 1/2/3	T183/Y185, T221/ Y223	Up	2.11
4	AMPKα1	T183	Up	3.85
5	GSK-3α/β	S21/S9	Up	2.78
6	AKT1/2/3	S473	Up	2.93
7	AKT1/2/3	T308	Up	2.03
8	WNK1	T60	Down	3.71

F Wnk1^f/f Wnk1^d/d — pAKT[S473], Total AKT, LE, S

G Wnk1^f/f Wnk1^d/d Wnk1^f/f Wnk1^d/d — pAKT [S473], pAKT [T308], AKT, GAPDH — GD 3.5 PPD 4.5

H Wnk1^f/f Wnk1^d/d — pFOXO1 [S256], FOXO1, GAPDH

Figure 5. Loss of uterine WNK1 altered the transcriptome and kinase phosphorylation

network during the receptive phase. (A) Activity of upstream regulators as predicted by

Ingenuity Pathway Analysis based on the altered uterine transcriptome of Wnk1$^{d/d}$ mice on PPD

4.5. See Table S4 for complete list. (B) Comparison of differentially expressed genes (DEGs)

between the uteri of WNK1 ablated (blue) and FOXO1 ablated (pink) mice identified 313 common

genes. (C) Percentage of the 313 genes categorized into commonly upregulated (pink),

commonly downregulated (green), or upregulated in one and downregulated in the other (yellow

and blue). (D and E) Kinome phosphorylation status in Wnk1$^{d/d}$ and Wnk1$^{f/f}$ uteri on

pseudopregnancy day (PPD) 4.5, with selected alterations shown in (E). All kinases with > 1.5

fold change in signal intensity as quantified by ImageJ is shown in Fig. S4. Results were acquired

using pooled uterine lysate from 6 mice in each group. (F and G) Expression of phosphorylated

and total AKT in Wnk1$^{f/f}$ and Wnk1$^{d/d}$ uteri on GD 4.5 as shown by immunohistochemistry ((F), LE

= luminal epithelium and S = stroma), and on GD 3.5 and PPD 4.5 as shown by western blotting

(G), scale bars = 25 μm. (H) Western blot analysis showing levels of phosphorylated and total

FOXO1 in Wnk1$^{f/f}$ and Wnk1$^{d/d}$ uteri on PPD 4.5.

1004

1005

Figure 6. WNK1 ablation led to FOXO1 nuclear exclusion via AKT phosphorylation which

is associated with decreased PP2A phosphatase expression. (A) Western blot showing levels

of phosphorylated and total AKT and FOXO1 in HEC1A and THESC cells transfected with 24 nM

siCTRL or siWNK1. (B) Immunofluorescence showing FOXO1 subcellular localization (green),

with nuclei presented in DAPI in HEC1A and THESC control cells (1, 4), siWNK1 transfected cells

(2, 5), and GDC0941 treated, siWNK1 transfected cells (3, 6), scale bars = 20 μm. (C) Expression

of FOXO1, AKT and PI3K members in HEC1A and THESC cells transfected with siCTRL,

siWNK1, and treated with AKT inhibitor GDC0941. (D and E) Expression of PI3K proteins (D),

and mTOR, PP2A subunits and PTEN (E) in Wnk1$^{f/f}$ and Wnk1$^{d/d}$ uteri on PPD 4.5.

1015

Figure 7. WNK1 regulates AKT signaling through direct interaction with PPP2R1A. (A)

WNK1 and YFP expression in HEC1A cells transfected the the YFP expressing control plasmid

(cYFP) or YFP-tagged WNK1 expression construct (c4161). (B) Co-immuniprecipitation of WNK1

and PPP2R1A with YFP in HEC1A whole cell lysate, as indicated by western blotting. (C)

Expression of PPP2R1A and PP2A subunits in HEC1A cells transfected with 24 nM siCTRL or

siWNK1 for 72 hours. (D) Expression of PP2A subunits A, B and C, AKT and FOXO1 in HEC1A

cells transfected with 72 nM siCTRL or siPPP2R1A for 72 hours. (E) Diagram illustrating the

WNK1-PP2A-AKT-FOXO1 signaling axis.

1025

1026

Supplemental Information Titles and Legends

Figure S1. Generation of uterine WNK1 ablation mouse model. (A) The Wnk1$^{f/f}$ mice harbours 2 loxP sites (red block "L") flanking the second exon of the *Wnk1* gene, which were crossed to mice carrying Cre under the control of PGR promoter. Resultant PGR$^{Cre/+}$; Wnk1$^{f/f}$ mice would have exon2 excised from the genome. (B and C) Decreased *Wnk1* gene and WNK1 protein expression were confirmed using qPCR (B) and western blotting (C), results shown are mean ± SD, * $p < 0.05$.

Figure S2. Delayed implantation in the Wnk1$^{f/f}$ mice was not due to aberrant ovarian-pituitary function. (A) Ovulation and fertilization was examined by inducing super-ovulation in the Wnk1$^{f/f}$ and Wnk1$^{d/d}$ mice, followed by mating to wild-type male mice. Number of 2-cell embryos were quantified by oviductal flushing and counting under brightfield microscope. (B and C) Serum progesterone (P4) and estradiol (E2) on GD 4.5 in Wnk1$^{f/f}$ and Wnk1$^{d/d}$ mice. Results shown are mean ± SD.

Figure S3. Principle component analysis of uterine transcriptome during receptivity from 4 Wnk1$^{f/f}$ mice (red) and 5 Wnk1$^{d/d}$ mice (blue).

Figure S4. Quantification and validation of the phosphokinase array. (A) The phosphokinase array was subjected to densitometrical quantification using ImageJ, and phosphorylation change which exceeded 1.5 fold are shown. (B) Phosphorylation and total levels of AKT, GSK-3α/β, and PRAS40 in the Wnk1$^{f/f}$ and Wnk1$^{d/d}$ uteri on PPD 4.5 were assayed by western blot, with each lane representing one mouse.

Figure S5. mTOR is activated in the Wnk1$^{d/d}$ uteri, but does not regulate AKT activity. (A)

Rapamycin, an mTOR inhibitor was unable to reverse WNK1 ablation induced FOXO1 nuclear

exclusion, indicating that AKT activity is maintained. (B) Rapamycin treatment did not rescue

WNK1 ablation induced phosphorylation of AKT and FOXO1 in both HEC1A (left) and THESC

(right) cells. (C) Expression and phosphorylation of FOXO1 and AKT in HEC1A cells transfected

with siCTRL, siWNK1, simTOR, or both siWNKT and simTOR. Results showed no rescue of

FOXO1 and AKT phosphorylation by mTOR knock-down.

1057

1058 **Figure S6. WNK1 immunoprecipitation confirmation.** HEC1A cells were subjected to

1059 immunoprecipitation (IP) using IgG (negative control) or WNK1 targeting antibody. The pulldown

1060 lysate showed enrichment of WNK1 in the WNK1 IP, but not in the IgG (black arrow indicates

1061 expected position of WNK1), as determined by western blotting.

1062

1063 **Fig. S7. Plasmid map of the WNK1 mammalian expression construct pcDNA-6.2-N-YFP-**

1064 **WNK1 (c4161).**

1065

1066 **Table S1. Serum progesterone levels in Wnk1$^{f/f}$ and Wnk1$^{d/d}$ mice on PPD 4.5 to confirm**

1067 **pseudopregnancy.**

1068

1069 **Table S2. List of differentially expressed genes in the Wnk1$^{d/d}$ uteri compared to Wnk1$^{f/f}$**

1070 **uteri on PPD 4.5.**

1071

1072 **Table S3. Functional annotation of altered transcriptome in Wnk1$^{d/d}$ uteri at PPD 4.5**

1073 **(DAVID).**

1074

1075 **Table S4. List of upstream regulators with altered activites in Wnk1$^{d/d}$ uteri at PPD 4.5 (IPA).**

1076

1077 **Table S5. List of potential WNK1 interacting partners identified by WNK1 IP-MS.**

1078

1079 **Table S6. List of antibodies, sources, and experimental conditions used in this study.**

1080 **Data Availability**

1081

1082 All data generated or analyzed during this study are included in this published article or
1083 in the data repositories listed in References.

1084

1085 Supplemental tables and figures can be found at:

1086

1087 Gene expression data can be found at: