Differential Effects of propofol and ketamine on Critical Brain Dynamics

Thomas Varley[1,2], Olaf Sporns[1], Aina Puce[1], John Beggs[3],

1 Psychological & Brain Sciences, Indiana University, Bloomington, Indiana, USA
2 School of Informatics, Indiana University, Bloomington, Indiana, USA
3 Department of Physics, Indiana University, Bloomington, Indiana, USA

¤Current Address: Psychological & Brain Sciences, Indiana University, Bloomington, Indiana, USA
* tvarley@iu.edu

Abstract

Whether the brain operates at a critical "tipping" point is a long standing scientific question, with evidence from both cellular and systems-scale studies suggesting that the brain does sit in, or near, a critical regime. Neuroimaging studies of humans in altered states of consciousness have prompted the suggestion that maintenance of critical dynamics is necessary for the emergence of consciousness and complex cognition, and that reduced or disorganized consciousness may be associated with deviations from criticality. Unfortunately, many of the cellular-level studies reporting signs of criticality were performed in non-conscious systems (in vitro neuronal cultures) or unconscious animals (e.g. anaesthetized rats). Here we attempted to address this knowledge gap by exploring critical brain dynamics in invasive ECoG recordings from multiple sessions with a single macaque as the animal transitioned from consciousness to unconsciousness under different anaesthetics (ketamine and propofol). We use a previously-validated test of criticality: avalanche dynamics to assess the differences in brain dynamics between normal consciousness and both drug-states. Propofol and ketamine were selected due to their differential effects on consciousness (ketamine, but not propofol, is known to induce an exotic state known as "dissociative anaesthesia"). Our analyses indicate that propofol dramatically restricted the size and duration of avalanches, while ketamine allowed for a more awake-like dynamic to persist. In addition, propofol, but not ketamine, triggered a large reduction in the complexity of brain dynamics. All states, however, showed some signs of persistent criticality when testing for exponent relations and universal shape-collapse. Further, maintenance of critical brain dynamics may be important for regulation and control of conscious awareness.

Author summary

Here we explore how different anaesthetic drugs change the nature of brain dynamics, using sub-dural electrophysiological arrays implanted in a macaque brain. Previous research has suggested that loss of consciousness under anaesthesia is associated with a movement away from critical brain dynamics, towards a less flexible regime. When comparing ketamine and propofol, two anaesthetics with largely different effects on consciousness, we find that propofol, but not ketamine, produces a dramatic reduction in the complexity of brain activity and restricts the range of scales where critical

dynamics are plausible. These results suggest that maintenance of critical dynamics may be important for regulation and control of conscious awareness.

1 Introduction

The hypothesis that the brain operates in a critical regime near a "tipping" point between different states (often, but not always, described as low and high entropy, respectively) is growing in popularity, both on neurophysiological evidence and due to appealing properties of critical, or near-critical, systems [1], that are thought to be key elements of optimal nervous system functioning. In both *in vivo* recordings and simulations, critical systems show the widest dynamic range [2–5], which indicates that critical systems can respond to, and amplify, a broad range of signals. For a system embedded in a complex environment, where salient signals may have a variety of intensities, a broad dynamic sensory range is a necessary adaptation. Critical systems also have optimized memory storage capacity from which complex information about states and patterns can be retrieved [6–9]. A related series of findings suggest that, in addition to optimal dynamic range and memory capacities, critical systems have optimized information transmission capabilities [10–12], and that the ability to integrate information is locally maximal near the critical zone as well [13]. Signs of critical dynamics have been found in the brains of a large number of different animals with different levels of CNS complexity, including rat brain cultures [18], leeches [17], turtles [16], zebrafish [15], freely-behaving rodents [19], non-human primates [20], and humans [14], suggesting that the evolutionary advantage conferred by critical neural dynamics is highly conserved between species. The "critical brain" hypothesis has not been universally accepted however [90], and disagreement remains within the field as to when it is acceptable to conclude data was produced by a critical or near-critical system [18].

Despite the considerable work that has been performed on identifying indicators of criticality at the level of neuronal circuits, the relationship between critical dynamics (or lack thereof) at the micro-scale and macro-scale phenomena such as cognition, sensation, and awareness is less clear. In human neuroimaging studies, deviations from criticality have been found to be associated with altered, or so-called "exotic" states of consciousness. "Exotic" states of consciousness refer to states that typically do not occur during normal brain functioning such as disorders of consciousness following traumatic brain injury, sleep deprivation, extreme psychiatric episodes, or drug-induced states such as those induced by psilocybin or ketamine. Long-term sleep deprivation reduces signatures of critical dynamics in human MEG activity [14], and the pathophysiology of epilepsy has been modelled as a failure to maintain healthy critical dynamics [21, 22]. Psychiatric disorders such as schizophrenia, obsessive-compulsive disorder and major depressive disorders have also been discussed as possible deviaitons away from "healthy" critical dynamics. Human neuroimaging studies have suggested that "classical" serotonergic psychedelic drugs (eg. psilocybin, lysergic acid diethylamide, dimethyltryptamine, etc) increase markers of criticality, both in fMRI [23, 24] and MEG/EEG/ECoG [25] studies. These findings have prompted some to hypothesize that normal consciousness emerges when brain activity is tuned near a critical regime, and that alterations in consciousness are reflective of transitions towards, or away from, the critical regime [26, 27].

However, signs of criticality in neural tissue are observed in animals (or cultured tissues) that could not be plausibly considered conscious. This includes dissociated cortical cultures [18], excised turtle brains [16], and animals anesthetized with a number of different anaesthetics [28]. Clearly criticality alone is not sufficient for conscious awareness, although the question of necessity remains open. Cortical recordings from

anaesthetized rats found loss of universal dynamics during the period of anaesthesia that re-emerged over the course of waking [29]. A similar study using cellular imaging found that anaesthesia reversibly altered signs of critical dynamics, as well as reducing the complexity of brain activity [30]. Computational models of brain dynamics informed by the pharmacology of anaesthetic agents have suggested that loss of consciousness induced by anaesthesia may be associated with a loss of critical dynamics [31] at a macro scale as well, although this remains an under-explored area of research.

To assess the relative differences in criticality between states of consciousness, we used publicly available invasive electrocorticography (ECoG) recordings from a *Macaca fuscata* monkey to compare the normal, awake resting-state (the monkey is awake, with eyes open, restrained in a primate chair) to two distinct states of anesthesia induced by two different drugs: propofol and ketamine (recordings were begun only after loss of consciousness had been diagnosed). These recordings are available as part of the NeuroTycho project's open data initiative [32, 33]. While both drugs induce surgical anaesthesia in high doses, and are used in clinical settings, they display markedly different pharmacologies and trigger different subjective experiences at low to moderate doses. Propofol is a commonly-used anaesthetic administered either by inhalation or intravenously. While its exact mechanism of action remains unknown, it is believed that it's primary action is through potentiating inhibitory $GABA_A$ receptors resulting in wide-spread decreases in neuronal activity [34]. Behaviourally, propofol induces sedation, atonia, and at high doses, cardiac arrest, respiratory depression, and hypotension [35]. ketamine acts as an antagonist of glutamatergic NMDA receptors and, in contrast to propofol, causes mild nervous system stimulation and has little effect on respiration [35, 36]. Furthermore, while propofol induces a state reminiscent of deep coma, at sub-anaesthetic doses, ketamine induces an atypical state known as "dissociative anaesthesia" [37, 38], in which a person appears unresponsive to sensory or physical stimuli, but will often experience dream-like states, including hallucinations, out-of-body experiences, and immersive visions [36]. In this way, ketamine models other exotic states of consciousness such as NREM sleep and locked-in syndrome, where phenomenological awareness can persist, despite an external appearance of unresponsive unconsciousness.

We chose electrophysiological recordings for this analysis for several reasons. Many previous studies that have found signs of criticality in anaesthetized animals have been performed while recording at the single-neuron micro-scale, while studies reporting alterations to critical dynamics following changes to consciousness have been performed using macro-scale neuroimaging methods (fMRI, MEG, etc). Intracranial ECoG provides a type of meso-scale signal: with coarse-graining of electrical activity in large numbers of neurons ($\approx 10^5$), but with finer localization than scalp EEG/MEG [39]. Consequently, these data present a bridging-scale, where the highly local preservation of critical dynamics in neuronal-level studies may interact with alterations to criticality seen in neuroimaging studies. Human ECoG studies have found indirect evidence of dynamical criticality [40, 41] - not using avalanche-based analyses such as we used here. Alonso et al (2014) reported changes high-frequency ECoG signal stability during the onset of clinical anaesthesia although signs of metastability persisted through all states.

Here we test two hypotheses concerning the relationship between consciousness and critical dynamics. Hypothesis 1: During the awake state, activity would express several hallmarks of criticality, including a power-law distribution of avalanche sizes and durations, exponent relations [18], as well as signs of universality and distribution shape-collapse when sub-sampling channels. Hypothesis 2: propofol would dramatically reduce signs of critical dynamics, but that criticality would persist under the influence of ketamine. We also included a related measure, complexity [42], which has been hypothesized to relate to consciousness and found to be associated with criticality in

Fig 1. Recording array map.

The placement of recording arrays for Chibi. Image taken from the NeuroTycho website, http://neurotycho.org/spatial-map-ecog-array-task

cortical cultures [13].

2 Materials and Methods

2.1 Data Set and Preprocessing

2.1.1 Data Set

We used the NeuroTycho dataset, an open-source set of 128-channel invasive ECoG recordings in two Macaca fuscata monkeys [32]. For this study, we analysed the resting-state scans from one monkey (Chibi), as the scans from the other monkey contained intractable artifacts that could not be eliminated during preprocessing. Arrays were placed on the left hemisphere only, recording from all major areas including the medial wall (for a map, see Fig. 1).

In both the propofol and ketamine conditions, the monkey was restrained in a primate chair. In the propofol condition, the monkey was given a single bolus of intra-venous propofol (5.2 mg/kg), until loss of consciousness was observed (defined as unresponsiveness to having their forepaws touched and/or unresponsiveness to having their nose tickled with a cotton swab). For the ketamine condition, a single bolus of intramuscular ketamine (5.1 mg/kg) was administered until loss of consciousness was observed using the above criteria. No maintenance anaesthesia was given following the initial induction. Following diagnosis of anaesthesia, 10-minute recordings were taken. We did not include any recordings from the transition period between wakefullness and anaesthesia. More detailed discussion of the drug administration protocols can be found in [33] and the Neurotycho data wiki (http://wiki.neurotycho.org/Anesthesia_and_Sleep_Task_Details).

2.1.2 Preprocessing

The data were initially examined visually in EEGLAB (v.14.1.2) [43] run in a MATLAB (v. 2018b) environment for major non-reoccurring artefacts. Some sample five-second plots of the selected ECoG channels can be seen in Figure 3. After inspection (no data

Fig 2. Best-fit normal distributions between conditions.
Representative examples of the histograms of instantaneous amplitudes across the three
conditions. **Left**: the awake condition. **Middle**: ketamine, and **right**: propofol. Note
that while ketamine allows for a heavy-tail to persist, the propofol condition collapses to
a tight Gaussian distribution.

were removed), we performed Independent Component Analysis (ICA) using the *runica*   127
function in EEGLAB to extract minor artefacts [39, 43]. We ran the ICA with the full   128
128 possible components and, across all scans, removed an average of 4 components,   129
corresponding to brief bursts of high-frequency noise. After the ICA and removal of   130
components, the data were filtered in MNE Python (v. 0.19.2) with a low-pass filter at   131
120Hz, high-pass filtered at 1Hz, and notch filtered at 50Hz and all subsequent   132
harmonics up to 250 Hz, to account for electrical line-noise in Japan. All filters were of 133
the FIR Overlap type (the default in MNE Python), [44])and all filters were run twice,   134
forwards and backwards to eliminate phase-shift artefacts. Finally, we normalized the   135
data by removing the mean of each channel and dividing it by its standard deviation.   136
Due to the need for long time-series in this analysis, we operated on the whole 10   137
minute time-series (there was no epoching). MNE Python (v. 0.19.2) was run in Python   138
3.7.4 using the Anaconda (v. 3.7) environment.   139

2.2   Point Process   140

For each condition, we followed the method for point-processing data described by [45]   141
and [14]. Briefly: after filtering, the probability distribution of signal amplitudes in   142
terms of their standard deviation from the mean, was plotted, along with the best-fit   143
Gaussian distribution, and the point at which the actual distribution and the best-fit   144
curve diverged were chosen as the threshold (σ). This ensured that diversions above the 145
threshold are unlikely to be due to noise inherent in the signal. While there were   146
differences in the point of divergence between conditions, we chose a threshold of 4, as   147
the most conservative likely value. We sampled a range of thresholds around 4 (3 to   148
4.25) and found that this did not substantively alter the results, although a too-low   149
threshold (< 3) allowed excessive noise through, while a too-high threshold (> 4.5) did 150
not allow enough events for analysis of avalanches. For visualization of the various   151
distributions, see Figure 2.   152

Once the threshold had been chosen, all instances where the absolute value of a   153
signal was less than the threshold where set to zero: 154

$$\forall t \in |X| : X(t) = 0 \ if \ |X(t)| < \sigma$$

For all excursions above σ, the global maxima of the excursion was set to 1 and all   155
other moments set to 0. We correlated the excursion, from t_{min} to t_{max} against the 156
same range in every other channel, and if $\rho \geq 0.75$, we also set the local maximum of 157
the interval in the associated channel to 1 (even if it did not cross our threshold σ) as 158
well (sending all other values to 0). For a more in-depth discussion of this, see [14].   159

Fig 3. Raw EEG timeseries visualized.
Visualization of five seconds of all 128 channels for each condition. Top: awake, middle: ketamine, bottom: propofol. Images taken from EEGLAB (v.14.1.2) [43]. Note that the time-series are un-normalized, with scales ranging from 464-515mV.

Fig 4. Coherence cascades in the awake condition.
An example of a section of the raster plot. Notice the banding effect of coherence
cascades (a few examples are highlighted by blue ovals). This banding effect shows the
cascades of events that include many distinct channels that "fire" in synchrony.

The resulting calculation produced a binary raster plot (channel $\times$ samples, for an
example see Fig. 4) where every event corresponds to the moment of maximal excursion
from the mean. This data structure matches the type used when analysing spiking
activity from cultured neurons [10], making it amenable to many of the tools already
developed for criticality analysis. We chose not to re-bin the data, instead maintaining
the 1000 Hz sampling rate across all scans due to the relative shortness of the
recordings: significant re-binning would reduce the available data to the point that not
enough individual avalanches could be identified.

2.3 Avalanches and Critical Exponents

"Avalanches", defined as transient periods of coordinated activity between elements of a
system, are a feature common to many complex, dynamic systems [46]. In the context
of the nervous system, avalanches typically refer to synchronized action potentials in a
neural network [10]. Coordinated avalanches of activity have also been found in human
EEG [45], MEG [14, 47], and fMRI [82] data. Avalanches are described by two values:
the avalanche size (S, the number of elements that participate in the avalanche) and the
avalanche duration (T, the lifetime of the avalanche from start to finish). In critical
systems, these two values are expected to follow power laws with exponents
τ, α and $1/\sigma\nu z$:

$$P(S) \propto S^{-\tau}$$
$$P(T) \propto T^{-\alpha}$$

S and T should be distributed such that, over all avalanches, the average size for a
given duration also follows a power law:

$$\langle S \rangle(T) \propto T^{1/\sigma\nu z}$$

τ, α and $1/\sigma\nu z$ are collectively known as the critical exponents of the system [18].
We note that in this context, $1/\sigma\nu z$ is treated as a single variable and should not be
considered a function of three distinct variables. Furthermore, all three exponents
should be related to each other such that:

$$\frac{\alpha - 1}{\tau - 1} - \frac{1}{\sigma\nu z} = 0$$

Fig 5. Synthetic and empirical doubly-truncated power law distributions for the data.
Left: an empirical, doubly-truncated power-law distribution taken from this dataset with an x_{min} of 16 and an x_{max} of 196. Within this range, the exponent has been estimated to be 2.7. Right: a synthetic doubly-truncated power-law distribution with the same minimum and maximum values as the empirical distribution as well as the same exponent. Note that the synthetic model has significant curves which begin well before the upper-bound cut-off. When modelling a system that produces data within a fixed range, even power-law distributed values may not follow the canonical straight line when plotted in log-log space [52, 53].

It is generally considered to be insufficient evidence of criticality if only the distributions of avalanche sizes and durations follow power laws: these exponents must be related to each other in specific ways [18].

To calculate the scaling exponents τ and α, we used the NCC Toolbox (v. 1.0) [49] to extract avalanches from our binary time series, and perform a maximum likelihood estimate (MLE) of the power-law exponent [49, 50]. The NCC Toolbox returns several values associated with each power-law inference: the MLE value of the exponent, the minimum and maximum values of x for which the power law estimate holds, and the p-value. Following the convention of Timme et al., (2016), we set our significance threshold such that we would only accept the power law hypothesis at $p \geq 0.2$ (this is the reverse of how significance estimation is usually performed, for discussion, see [50]. For the estimate of $1/\sigma\nu z$, we plotted $\langle S \rangle(T)$ against T and extracted an estimate of the exponent by linear regression in log-log space. While this is a much cruder method than the MLE power law fit described above, unfortunately the values of $\langle S \rangle(T)$ vs. T do not describe a probability distribution and so the usual methods of inference do not work.

When plotting the distribution of avalanche sizes and avalanche durations, we use complementary cumulative distribution functions (CCDFs) (defined as 1-CDF) instead of probability density functions (PDFs), following [51]. When dealing with power law (or plausibly power law) distributions with defined upper and lower bounds (x_{min}, x_{max}), the CDF and CCDFs may not display the characteristic straight line when plotted in log-log space [52, 53]. Consequently, distributions may look curved while still being plausibly drawn from a doubly-truncated power law. For a visualization of this, see Figure 5.

2.3.1 Avalanche Shape Collapse

In addition to having sizes and durations, avalanches also have *profiles*, which describe the number of channels active at each moment over the course of the avalanche's lifetime. In the critical state, the profiles of all avalanches should be self-similar, that is,

it should be possible to find some scaling exponent γ such that all avalanches can be ²¹² rescaled to lie on top of one another [13,18]. This is referred to as "shape collapse" and ²¹³ is a commonly used indicator of universality in dynamical systems. ²¹⁴

This scaling parameter γ can be defined as: ²¹⁵

$$\gamma = \frac{1}{\sigma \nu z} - 1$$

where $1/\sigma \nu z$ is the same exponent that describes the distribution of average ²¹⁶ avalanche sizes for a given duration. ²¹⁷

To calculate shape collapse, we first averaged together all avalanches of a given ²¹⁸ duration to create "average profiles". From there, we used the NCC Toolbox [49] to find ²¹⁹ the optimal rescaling value and extract an estimate of $1/\sigma \nu z$. ²²⁰

2.4 Universality and Shape Collapse ²²¹

A common issue with analysis of critical dynamics in complex systems is one of ²²² sub-sampling [54,55]. When a very large system is sub-sampled, the characteristic ²²³ power-law behaviour can be lost as large events are fragmented and perceived as ²²⁴ separate, smaller events. This is a particularly salient issue in electrophysiological ²²⁵ recordings, as the number of channels in an array is orders of magnitude less than the ²²⁶ number of functional units in the cortex that could be participating in critical avalanche ²²⁷ dynamics. ²²⁸

In a system in the critical regime, while subsampling destroys the power-law, it ²²⁹ should not destroy the scale-free nature of the distributions - that is, when ²³⁰ appropriately renormalized, the distributions (despite not following power laws), should ²³¹ display shape collapse. The logic here is the same as when performing shape collapse on ²³² individual avalanches, as above. ²³³

We tested for distribution shape-collapse following a procedure similar to [55]. For ²³⁴ each scan, we randomly sub-sampled half, a quarter, and an eighth of the channels, ten ²³⁵ times each, and from each, calculated the associated probability distribution of ²³⁶ avalanche events and sizes. In addition to plotting them to visualize the differences ²³⁷ between the four distributions (the original plus the three subsampled distributions), we ²³⁸ calculated the average pair-wise Wasserstein metric [80] to quantify the similarity ²³⁹ between all distributions. ²⁴⁰

We then performed the same sub-sampling again, this time rebinning the binary ²⁴¹ timeseries to reflect the sampling of elements: when taking half the channels, we ²⁴² rebinned by a factor of two, when taking a quarter of the channels, we rebinned by a ²⁴³ factor of four, etc. If the system is poised near the critical point, after rebinning, the ²⁴⁴ distributions of avalanche activities should collapse onto one-another. When ²⁴⁵ recalculating the Wasserstein metric, the average pairwise distance should be ²⁴⁶ significantly reduced. ²⁴⁷

2.5 Complexity ²⁴⁸

Here, "complexity" of an N-dimensional system X ($C_N(X)$) can be intuitively ²⁴⁹ understood as "the degree to which the whole is greater than the sum of it's parts" [42]. ²⁵⁰ To calculate the complexity, the entropy of the entire system is compared to the joint ²⁵¹ entropies of all possible subsets of the system. Each state i of an N-dimensional system ²⁵² can be described as an N-dimensional vector x_i, and the probability of state i (p_i) is the ²⁵³ number of occurrences of i divided by the number of samples. The Shannon entropy ²⁵⁴ ($H(X)$) of the system is given by: ²⁵⁵

$$H(X) = -\sum_i p(x_i) log_2(p(x_i))$$

We can measure the level of integration (or coordination) of a set of elements of the system by comparing the joint entropies of all the elements in the set to the sum of their individual entropies:

$$I(X)_j^k = \left(\sum_{j' \in k} H(X_{j'}^1) \right) - H(X)_j^k$$

Here, k is the number of neurons in a subset, and j is the index of a given subset of k neurons in the set of all sets of k neurons. This has also been called "multinformation" or "total correlation" [56]. For instance, X_j^1 refers to the j^{th} element from the set of N -choose-1 elements alone, while X_j^4 refers to the j^{th} set of 4 elements from the set of N-choose-4 sets of 4 elements.

The complexity is then found by comparing the integration of the entire system (X_1^N) to the average integration $(\langle I(k)_1^j \rangle j)$ over every possible subset, for every possible subset size of k.

$$C_N(X) = \frac{1}{N} \sum_{k=2}^{N} \left[\left(\frac{k-1}{N-1} \right) I(X) - \langle I(X_j^k) \rangle_j \right]$$

This value of $C_N(X)$ is a more intuitively meaningful representation of "complexity" than other commonly used measures, such as Lempel-Ziv Complexity [57], in that it is low for systems that are both perfectly ordered and systems that are perfectly random [13], while peaking in systems that combine elements of both. Calculating the complexity of a system is a non-trivial task, exploding to intractable levels as N gets even modestly large (for context, for a 128-channel system, it would take longer than the expected lifetime of the universe of exhaustively search all partitions). To avoid interminable run-times, the NCC Toolbox [49] includes several corrections for sub-sampling and heuristics for estimating integration in large systems.

As with the avalanche size and duration distributions, we used the same 30-second jitter null-model to explore the effect of randomizing the data. We hypothesized that jittering the data should significantly reduce the nonlinearity and total integrated information.

3   Results

3.1   Channel Activity

We found large differences between the channel rates following the administration of both ketamine and propofol. In the Awake condition, the average channel-wise firing rate was 9×10^{-4} spikes per millisecond (s/m), which dropped to 5×10^{-4} (s/m) in the ketamine condition and 8×10^{-5} (s/m) under propofol. We found that, within conditions, there was a high correlation between channel activity rates during different scans. Within the Awake condition (across all six possible pairwise comparisons), we found an average correlation of 0.89 ± 0.03. In the ketamine condition (which only allows a single comparison), the correlation was 0.76 and in the propofol condition, the correlation was 0.62). Interestingly, we did not find strong correlations when comparing channel activity rates within the same scan before and after induction of anaesthesia (data not shown). The correlation between channel activity rates before and after ketamine induction was 0.225 and the correlation between channel activity before and

Fig 6. Changes in channel activity levels between conditions.
A: For each condition, the correlation between channel activity (firing rate $\times$ 100) between two scans. In all conditions, it is apparent that those channels that are more activity in one scan are similarly active during different scans within the same condition. This suggests that brain dynamics are consistent within conditions. **B**: histograms of the the channel rate $\times$ 100 for all three conditions. It is apparent that propofol decreases the overall channel firing rate more that ketamine, although both have a lower firing rate than the Awake condition. **C-E**: normalized channel activity rates projected onto the recording array. It is clear that propofol and ketamine disrupt the activity-rate structure that exists in the Awake condition.

after propofol induction was 0.06. These results suggest a much higher degree of dynamical similarity within conditions scanned on different days than between conditions induced during a single scan. For visualization of these results, see Figure 6.

    Using the NeuroTycho channel labelling (see Figure 1), we found that, in the Awake condition, high levels of activity were seen occipito-temporal and parietal regions of the brain. This pattern was disrupted by both propofol and ketamine. In the ketamine condition, there was a decrease in the range of channel activities, and in both anaestheic conditions, there was a shift in activity towards the occipital lobe. This significance of this is unclear, as the monkeys were wearing eyeshades throughout the anaesthetic experience and consequently were not receiving visual input.

3.2   Avalanche Distributions

There are clear differences between the Awake, ketamine, and propofol distributions of avalanche sizes and durations (Figure 7). The Wasserstein distances between the conditions reveal that, in all cases, the Awake condition was far more similar to the ketamine condition than the propofol condition. The distance between the avalanche

size distribution in the Awake condition and the ketamine condition was 0.46 ± 0.02, while between the Awake condition and the propofol condition it was 1.77 ± 0.21. The same pattern was apparent between the avalanche duration distributions. The distance between the Awake and ketamine conditions was 0.05 ± 0.04, while for the Awake and propofol conditions it was 0.43 ± 0.06 (for the raw values, see Supplementary Table).

3.3 Scaling Exponents

For every distribution, there was a range between some x_{min} and x_{max} for which the power-law hypothesis held with $p \geq 0.2$. However, in the propofol condition, this range was dramatically restricted. In the Awake condition, the average value x_{min} for the avalanche size distribution was 16.25 ± 3.5 channels and the average value for x_{max} was 378.5 ± 160.15 channels. In the ketamine condition, the average value x_{min} for the avalanche size distribution was 14 ± 8 channels, and the average value for x_{max} was 347 ± 287 channels. Finally, in the propofol condition, the average value x_{min} for the avalanche size duration was 2 ± 0 channels and the average value x_{max} was 48 ± 8 channels. The same pattern was observed in the distributions of avalanche durations. The average x_{min} of the distribution of avalanche durations in the Awake condition is 8 ± 0.7 ms, while the average x_{max} is 42.5 ± 14.74 ms. In the ketamine condition, the average x_{min} was 6 ± 2 ms and the average x_{max} is 42 ± 25 ms. In the propofol condition, the range was much more constricted: the average x_{min} was 2 ± 1 ms and the average x_{max} was 12 ± 4 ms. These results indicate that, while all distributions had regions that could be plausibly modelled as following a power-law, in the propofol condition, this region was extremely narrow and typically restricted to very small values (excluding the tail, where the power-law distribution is most relevant). In contrast, ketamine maintained a range comparable to the Awake condition, suggesting that, unlike propofol, ketamine anaesthesia does not suppress the propagation of large bursts of coordinated activity.

For each range where the power-law fit held, we calculated the associated exponents, τ and α. The sample size is too small for significance testing, although on average, the Awake condition had the largest exponents for avalanche sizes (2.74 ± 0.19) and durations (4.23 ± 0.48), followed by ketamine, for both sizes (2.58 ± 0.5) and durations (3.52 ± 0.72) as well. Propofol had the smallest scaling exponents for both avalanche parameters (size: 2.3 ± 0.003, durations: 2.93 ± 0.68). The average calculated values of $1/\sigma \nu z$ were quite similar between all three (Awake: 1.89 ± 0.5, ketamine: 1.6 ± 0.05, propofol: 1.5 ± 0.53).

In all conditions, within the relevant ranges of x_{min} and x_{max}, we found a strong relationship between the average size for a given duration (see Figure 8). The calculated value of $1/\sigma \nu z$ was similar across all conditions (Awake: 1.33 ± 0.08, ketamine: 1.25 ± 0.005, propofol 1.34 ± 0.14). Having two separate estimations of $1/\sigma \nu z$ (one from the exponent relation, one from the average size for a given duration) allows us to see how well the scaling exponents relate. Surprisingly, the Awake condition had the greatest percentage difference between the two values, at 35.19%, followed by ketamine at 24.76% and finally propofol with 10.83%. We had hypothesized that the Awake and/or ketamine conditions would show the highest degree of concurrence between the measures (reflecting a greater degree of criticality), but instead, the Awake condition has the lowest degree of concurrence.

3.4 Avalanche Shape Collapse

The avalanche collapse results are largely consistent with the analysis of average size for a given duration. The estimated values of $1/\sigma \nu z$ calculated from the shape collapse between conditions were largely similar: Awake: 1.261 ± 0.12, ketamine: 1.24 ± 0.02,

Fig 7. CCDFs of Avalanche Size and Duration by Condition
Comparison CCDFs for Awake (purple) vs. ketamine (grey) and propofol (orange).
Right panels: Awake vs. ketamine. **Left** panels: Awake vs. Propofol. Avalanches sizes
(top four panels) and durations (bottom four panels). Visual inspection shows that the
ketamine condition tracks the Awake condition much more closely than the propofol
condition does. This indicates that ketamine supports larger, longer-lived avalanches
than propofol.

Fig 8. The average avalanche size for a given duration by condition
The color mapping is the same as in Figure 7: purple corresponds to the Awake
condition, grey to ketamine, and orange to propofol. In all cases there is a clear linear
relationship between the average avalanche size for a given duration. Such a relationship
is considered necessary, but not sufficient, to conclude a system is displaying critical
dynamics [18].

propofol 1.39 ± 0.17. Once again, the percentage difference between the these values   358

and the exponent relationship was highest in the Awake condition (40.2%), followed by the ketamine condition (25.95%), and with the greatest agreement in the propofol condition (6.7%). We also calculated the percentage difference between values of $1/\sigma\nu z$ derived from the shape collapse and the average size/given duration analysis. These showed a much higher degree of concurrence. The Awake condition again had the greatest percentage difference (5.19%), followed by propofol (3.48%), and ketamine had the lowest percentage difference (1.2%).

3.5 Universality and Distribution Collapse

In addition to power-law and exponent-relation based indicators of critical dynamics, we also tested whether the probability distributions of avalanche sizes and durations exhibited shape-collapse when subsampled [55]. We initially sub-sampled the system by randomly selecting subsets of channels (half the channels, a quarter, an eighth, etc) and comparing the distributions of avalanche sizes and durations by computing the average pairwise KS-distance between all four distributions. We then rebinned the binary time series with the inverse of the fraction of the sample selected (eg: when subsampling half the nodes, we rebinned the timeseries by two, when sampling a quarter, rebinned by four, etc).

When compared, all conditions showed visual indicators of universal shape-collapse (see Figure 9 for an illustration with the distributions of avalanche sizes). To quantify this, we calculated the fold-change between the average pairwise KS-distance before and after re-binning. For the distributions of avalanche sizes, ketamine showed the highest average fold change (-0.66 ± 0.05), indicating the greatest collapse. The Awake condition came second with a fold-change of -0.61 ± 0.04, and the propofol condition showed the weakest shape-collapse, with a fold change of -0.49 ± 0.05. For the avalanche duration distributions, the pattern was similar, with ketamine showing the greatest fold change (-8.1 ± 0.01), but in this case, propofol was in the middle with a fold change of -0.73 ± 0.01, and the Awake condition (-0.69 ± 0.03) at the end.

This suggests that all the conditions display some measurable shape collapse when renormalized, and in both avalanche shape and durations, the effect was most dramatic in the ketamine condition, compared to Awake and propofol.

3.6 Complexity

There were dramatic differences between the degree to which ketamine and propofol reduced the complexity of brain activity (Figure 10). Both anaesthetics reduced the total $C_N(X)$, however, propofol had a much stronger effect. On average, the fold-change between the Awake condition and the ketamine condition was -0.35 ± 0.03, compared to the Awake vs. Propofol condition, which showed an average fold-change of -0.91 ± 0.02. This is a strong indicator that, even at surgical doses, ketamine supports significantly more complex brain dynamics than propofol.

4 Discussion

In this manuscript we present several converging lines of evidence that propofol and ketamine produce different effects on markers of critical brain dynamics. Furthermore, ketamine, but not propofol, appears to supports dynamics similar to those observed in normal waking consciousness in ECoG activity recorded from a single macaque in different states of consciousness. Despite inducing states that appear similar to external observers (unresponsiveness to pain or sensory stimuli, decreased voluntary motor output, anaesthesia), propofol produces a near-total extinction of consciousness, while

Fig 9. Shape collapse and universality under rescaling

Shape collapse for the various subsampled avalanche sizes before (left column) and after (right column) renormalizing (rebinning) the binary timeseries. The color indicators are consistent with other figures: purple is Awake, grey is ketamine, yellow is propofol. In all conditions, renormalization resulted in noticeable shape collapse, which can be quantified by doing pairwise calculations of the Wasserstein distance metric.

ketamine frequently induces states of "dissociative" anaesthesia, which can include   405
dream-like or out-of-body experiences [37, 38]. To test for signs of criticality we explored   406
the dynamics of avalanches (coordinated bursts of high-intensity activity) across the   407
cortex, testing for scale-free distributions and exponent relations between them. We also   408

Fig 10. Differences in multi-scale complexity between conditions.
Integration of information across scales for Awake (purple) vs. ketamine (grey) and
Awake vs. Propofol (orange). Complexity was calculated using the NCC Toolbox [49],
using the measure first proposed by Tononi, Sporns & Edelman [42]. While the
ketamine condition (grey) was associated with a noticable decrease in complexity
compared the Awake condition (purple), it was a much smaller decrease than what was
observed in the propofol condition, which was an order of magnitude less complex. Note
that the total value of complexity is not the greatest value, but rather, the difference
between the area under the curves for the linear fit and the true non-linear integration.

examined universality under renormalization and an information-theoretic measure of
multi-dimensional complexity to further characterize the effects of these drugs on brain
dynamics.

We found that ketamine slightly reduced the rate of events across the cortex when
compared the Awake condition, while the propofol condition reduced activity rates by
an entire order of magnitude. Within conditions, the distributions of activities across
channels were consistent between scans, suggesting that these measures are stable across
multiple drug experiences, at least within one individual.

4.1 Avalanche Distributions

We found that, when compared to normal wakefulness, propofol dramatically attenuated
signs of power-law distributions in both the distributions of avalanche sizes and
durations. Heavy tailed distributions imply that avalanche dynamics are playing out
over a range of temporal and spatial scales; consequently, the finding of multiscale
dynamics in the Awake and ketamine conditions, but not the propofol condition
suggests that this kind of multiscale coordination is significant for the maintenance of
consciousness. This is consistent with previous work using fMRI neuroimaging

suggesting that loss of multiscale structure in brain networks occurs in patients with disorders of consciousness [66, 79] and decrease in the fractal dimension of EEG and fMRI BOLD signals [75–78].

The differences in the distributions of avalanche sizes and durations may be indicative of changes to the ability of the brain to coordinate activity in different states. In the Awake condition, we observed avalanches that included many distinct channels, as well as being comparatively longer lived, while those long, large avalanches were significantly impaired by propofol, but not by ketamine. The propofol result is consistent with previous work which found that propofol inhibited the ability of the brain to form integrated higher-level networks, while leaving local activity in tact [71, 72]. If avalanches indicate when signals are able to propagate across the cortex, we might expect to see alterations in avalanche activity being associated with fragmentation or disintegration of functional connectivity networks. Previous research using point-processing methods on fMRI time-series has found that rare, high-amplitude events can capture a significant information relevant to critical brain dynamics and states of consciousness [82, 83]. Using techniques from information theory such as mutual information and transfer entropy [84] it would be informative to construct functional connectivity networks from the binarized time-series to directly relate changes in critical dynamics to alterations to network topologies and computational capabilities [85, 86].

The relative differences in the frequency of large avalanches between the three conditions may also be relevant to work with the Global Workspace Theory (GWT). Previous work has proposed that large avalanches represent the "ignition" of information percolating through the neuronal global workspace [73, 81]. In the context of the GWT, this percolation is thought to define the difference between information processing that is "conscious", from processing that is "unconscious" [74], and so ketamine may allow consciousness to persist (in contrast to propofol) by failing to disrupt the ability of information to "ignite" and propagate into the Global Workspace.

4.2   Indicators of Critical Dynamics

The other indicators of criticality are harder to interpret. All conditions showed similar patterns of critical exponent relations, with the value of $1/\sigma\nu z$ calculated from the avalanche shape collapse showing a high degree of agreement when calculated from the regression of the average avalanche size for a given avalanche duration. In contrast, both values of $1/\sigma\nu z$ showed less consistency when compared to the exponent relation $\alpha - 1/\tau - 1$. Finally, when individual channels were subsampled and rescaled to test for signs of universality, all conditions showed reasonable signs of shape-collapse, with ketamine typically showing the most significant signs of universality.

None of the states showed consistently tighter exponent relations, although we should stress that all exponents were calculated *within the range of x_{min} and x_{max} for which the power-law hypothesis fit with p-values $\geq$ 0.2*. The propofol condition had a dramatically reduced range of x_{min} and x_{max} values compared to Awake and ketamine, so whatever inferences about the plausibility of critical dynamics we make from these data is only relevant within these ranges. Consequently, although all three conditions showed reasonably similar behaviour in terms of exponent relations, overall, the Awake and ketamine conditions showed this behaviour over 2 orders of magnitude, while propofol showed it over one tenth of that range. Based on these, we propose that the brain is able to support critical dynamics in all three states, but that propofol (but not ketamine) *reduces the scale over which critical dynamics can occur.* Critical dynamics are often described as being 'scale-free', so the notion of restricting critical dynamics to a range of scales may seen contradictory. However perfect scale-freeness of critical systems only occurs in infinite systems. In all finite systems, criticality can emerge in a restricted range.

This shrinking of the critical zone may provide an explanation for the original motivating question for this study: why do critical dynamics seemed preserved in unconscious systems at the micro-scale [16, 18, 28], but altered at the macroscale [23, 27]? It may be that at the local level, neural networks are able to maintain critical dynamics, but the ability of these local ensembles to coordinate is impaired [71, 72]. Relevant to this hypothesis is the finding that anaesthetics alter the functioning of pyramidal neurons in Layer IV and V of the cortex [87, 88]. As Layer V pyramidal neurons are thought to serve as "outputs" from one brain region to another [89] and so changes to pyramidal neural functioning may explain how local critical dynamics are able to propagate to macro-scale critical dynamics. Layer V pyramidal neurons expressing 5-HT_{2A} receptors have already been implicated in macro-scale critical dynamics [26] in the context of serotonergic drugs, so a possible involvement in anaesthesia is not totally out of the question.

The question about relative ranges is significant for the exponent relationship results, but not necessarily for the universality and sub-sampling results, which are probably the hardest to interpret. All three conditions showed roughly equivalent degrees of shape collapse upon time-series re-binning, suggesting that all three display self-similar dynamics across channels. One possible interpretation is that, despite the alteration in consciousness induced by propofol and ketamine, the brain is able to adaptively maintain at least some qualities of critical dynamics. Previous work has found that the brain appears to adapt to perturbations and return to the critical regime despite alterations to incoming sensory inputs [16, 91]. One possible explanation for the persistence of signs of criticality is that the brains are rapidly adapting to the drug state. If this is the case, it presents an intriguing window of possibility: might it be possible to break "criticality" writ-large down into separate phenomena and explore which ones are necessary (or sufficient) for complex consciousness and cognition, and which may be irrelevant?

4.3   Complexity

We also found that, using a multi-scale measure of complexity [42], the propofol condition was associated with dramatic decreases in the complexity of brain activity, while ketamine preserved higher levels of complexity.

The most dramatic difference is the effect of the anaesthetics on multi-scale complexity. This measure has been suggested to be relevant to the emergence of phenomenological consciousness [62]. Our findings are consistent with tendency of ketamine to preserve consciousness in states of dreamlike "dissociative anaesthesia" in contrast to propofol. It is unsurprising that highly complex behavioural states should be underpinned by complex dynamics, and evidence from studies of anaesthesia [63–65, 67], disorders of consciousness [66, 68], sleep [69], and psychedelic states [24, 65, 70] bears this out. Our results are consistent with, and extend these previous findings using a more principled measure of "complexity" than randomness-based measures.

4.4   Limitations

One of the most significant limitations of this work is that it rests on the assumption that, when in the ketamine condition, the monkey was experiencing dissociative anaesthesia, as opposed to true anaesthesia (as it presumably experienced under propofol). The dosage used in the original study (5.1mg/kg) were consistent with doses used for pre-surgical anaesthesia in primates [33] and it is unclear at what dose dissociative anaesthesia transitions into "true" anaesthesia (with no dream-like content at all). Presumably even in cases of "true" ketamine anaesthesia, the subject passes through hallucinatory states during induction and emergence [36–38]. It is also not

entirely clear that, even if the dose was appropriate to induce hallucinatory states of dissociative anaesthesia in humans, whether macaques experience such a state at all. Despite this limitation, the fact that ketamine maintained wakefulness-like dynamics on multiple measures implies that it is not inconceivable that whatever processes allow consciousness to persist under ketamine in humans are also playing out in the macaque brain.

Having only a single viable monkey to analyse is another significant limitation, although given the rarity of this dataset, we believe the results are still informative. This work should, however, be replicated in a larger cohort of animals undergoing similar anaesthesia and recording procedures. Concerns about the low power are at least somewhat ameliorated by the fact that, for our monkey, we have $2\times$ scans in both the propofol and ketamine conditions, and $4\times$ scans in the Awake condition. In general, the results were quite consistent within conditions, suggesting that, at least for this individual, the effects observed here are reasonably robust.

Finally, the analyses described here require binarizing continuous time-series data, which throws out a considerable amount of information (although point-processes like this have been found retain surprising amounts of information, see [82,83]). For this study, the point process was necessary as the indicators of critical dynamics we explored here were derived from models of binary branching processes [18]. Future analyses that incorporate the full, continuous time series may provide insights missing from this current body of work. The method that we used to define an "event" (an above-threshold excursion in one channel or a sequence in another channel that is significantly correlated with the excursion) partially addresses this concern by including temporal similarity as a criteria by which an event could be identified, but further improvements are no doubt possible.

5   Conclusions

These results show that, despite their shared status as anaesthetics and the similarity of the external appearance of effects, propofol and ketamine cause different states typified by dramatically different neural dynamics. Unlike propofol, ketamine allows for large, long-lasting avalanches of coordinated activity to persist in the cortex in a manner similar to normal consciousness, as well as maintaining a much higher degree of multi-scale complexity. These results may explain why the state induced by ketamine can often include complex, phenomenological consciousness to persist in a dissociated, dream-like state, while propofol extinguishes awareness completely. Understanding which brain dynamics are necessary or sufficient to support consciousness is of interest to both for those interested in the theoretical basis of conscious awareness as well as addressing clinical concerns related to identifying covert consciousness in patients who may not be able to directly communicate.

Acknowledgements

TFV is supported by the NSF-NRT grant 1735095, Interdisciplinary Training in Complex Networks and Systems. We would like to thank Dr. Yang-Yeol Ahn for advice on power-law inference and visualization, Dr. Filippo Radicci for insights and discussion, as well as Joshua Faskowitz and Dr. Alice Patania for support, friendship, and advice.

6 Supporting information

6.1 Glossary

- **Anaesthetic**: A drug that reversibly disrupts normal consciousness, inducing a state of coma-like unconsciousness.

- **Criticality**: Refers to a system on the boundary between two qualitatively distinct phases. A canonical example from physics is boiling water, which is at the critical boundary between liquid and gas.

- **Metastability**: Refers to a dynamical regime where the system orbits a set of competing attractors without any one "winning." Associated with the maintenance of complex, adaptive behaviour.

- **Power Law**: A statistical distribution where $P(X = x) \propto x^{-\alpha}$. Critical systems are expected to produce power law distributions of events, although non-critical systems can produce them as well.

- **Universality**: When a system has the same distributions across scales, usually a sign of critical dynamics. By appropriately rescaling events, distributions can be made to collapse onto each-other.

S1 Table   All of the results reported, as a .csv table.

References

1. Luca Cocchi, Leonardo L. Gollo, Andrew Zalesky, and Michael Breakspear. Criticality in the brain: A synthesis of neurobiology, models and cognition. *Progress in Neurobiology*, 158:132–152, November 2017.

2. Osame Kinouchi and Mauro Copelli. Optimal dynamical range of excitable networks at criticality. *Nature Physics*, 2(5):348–351, May 2006.

3. Woodrow L. Shew, Hongdian Yang, Thomas Petermann, Rajarshi Roy, and Dietmar Plenz. Neuronal Avalanches Imply Maximum Dynamic Range in Cortical Networks at Criticality. *Journal of Neuroscience*, 29(49):15595–15600, December 2009.

4. Daniel B. Larremore, Woodrow L. Shew, and Juan G. Restrepo. Predicting Criticality and Dynamic Range in Complex Networks: Effects of Topology. *Physical Review Letters*, 106(5):058101, January 2011.

5. Shree Hari Gautam, Thanh T. Hoang, Kylie McClanahan, Stephen K. Grady, and Woodrow L. Shew. Maximizing Sensory Dynamic Range by Tuning the Cortical State to Criticality. *PLOS Computational Biology*, 11(12):e1004576, December 2015.

6. M. Breakspear. Perception of odors by a nonlinear model of the olfactory bulb. *International Journal of Neural Systems*, 11(2):101–124, April 2001.

7. Craig V. Stewart and Dietmar Plenz. Inverted-U profile of dopamine-NMDA-mediated spontaneous avalanche recurrence in superficial layers of rat prefrontal cortex. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 26(31):8148–8159, August 2006.

8. Gustavo Deco and Viktor K. Jirsa. Ongoing cortical activity at rest: criticality, multistability, and ghost attractors. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 32(10):3366–3375, March 2012.

9. Dong-Ping Yang, Hai-Jun Zhou, and Changsong Zhou. Co-emergence of multi-scale cortical activities of irregular firing, oscillations and avalanches achieves cost-efficient information capacity. *PLOS Computational Biology*, 13(2):e1005384, February 2017.

10. John M. Beggs and Dietmar Plenz. Neuronal Avalanches in Neocortical Circuits. *Journal of Neuroscience*, 23(35):11167–11177, December 2003.

11. Woodrow L. Shew, Hongdian Yang, Shan Yu, Rajarshi Roy, and Dietmar Plenz. Information Capacity and Transmission Are Maximized in Balanced Cortical Networks with Neuronal Avalanches. *Journal of Neuroscience*, 31(1):55–63, January 2011.

12. Erik D. Fagerholm, Martin Dinov, Thomas Knöpfel, and Robert Leech. The characteristic patterns of neuronal avalanches in mice under anaesthesia and at rest: An investigation using constrained artificial neural networks. *PloS One*, 13(5):e0197893, 2018.

13. Nicholas M. Timme, Najja J. Marshall, Nicholas Bennett, Monica Ripp, Edward Lautzenhiser, and John M. Beggs. Criticality Maximizes Complexity in Neural Tissue. *Frontiers in Physiology*, 7, 2016.

14. Oren Shriki, Jeff Alstott, Frederick Carver, Tom Holroyd, Richard N.A. Henson, Marie L. Smith, Richard Coppola, Edward Bullmore, and Dietmar Plenz. Neuronal Avalanches in the Resting MEG of the Human Brain. *The Journal of neuroscience : the official journal of the Society for Neuroscience*, 33(16):7079–7090, April 2013.

15. Adrián Ponce-Alvarez, Adrien Jouary, Martin Privat, Gustavo Deco, and Germán Sumbre. Whole-Brain Neuronal Activity Displays Crackling Noise Dynamics. *Neuron*, 100(6):1446–1459.e6, December 2018.

16. Woodrow L. Shew, Wesley P. Clawson, Jeff Pobst, Yahya Karimipanah, Nathaniel C. Wright, and Ralf Wessel. Adaptation to sensory input tunes visual cortex to criticality. *Nature Physics*, 11(8):659–663, August 2015.

17. Alberto Mazzoni, Frédéric D. Broccard, Elizabeth Garcia-Perez, Paolo Bonifazi, Maria Elisabetta Ruaro, and Vincent Torre. On the Dynamics of the Spontaneous Activity in Neuronal Networks. *PLoS ONE*, 2(5), May 2007.

18. Nir Friedman, Shinya Ito, Braden A. W. Brinkman, Masanori Shimono, R. E. Lee DeVille, Karin A. Dahmen, John M. Beggs, and Thomas C. Butler. Universal critical dynamics in high resolution neuronal avalanche data. *Physical Review Letters*, 108(20):208102, May 2012.

19. Yuta Senzai, Antonio Fernandez-Ruiz, and György Buzsáki. Layer-Specific Physiological Features and Interlaminar Interactions in the Primary Visual Cortex of the Mouse. *Neuron*, 101(3):500–513.e5, February 2019.

20. Thomas Petermann, Tara C. Thiagarajan, Mikhail A. Lebedev, Miguel A. L. Nicolelis, Dante R. Chialvo, and Dietmar Plenz. Spontaneous cortical activity in awake monkeys composed of neuronal avalanches. *Proceedings of the National Academy of Sciences*, 106(37):15921–15926, September 2009.

21. David Hsu, Wei Chen, Murielle Hsu, and John M. Beggs. An open hypothesis: is epilepsy learned, and can it be unlearned? *Epilepsy & behavior : E&B*, 13(3):511–522, October 2008.

22. Christian Meisel, Alexander Storch, Susanne Hallmeyer-Elgner, Ed Bullmore, and Thilo Gross. Failure of Adaptive Self-Organized Criticality during Epileptic Seizure Attacks. *PLoS Computational Biology*, 8(1), January 2012.

23. Selen Atasoy, Leor Roseman, Mendel Kaelen, Morten L. Kringelbach, Gustavo Deco, and Robin L. Carhart-Harris. Connectome-harmonic decomposition of human brain activity reveals dynamical repertoire re-organization under LSD. *Scientific Reports*, 7(1):17661, December 2017.

24. T. F. Varley, R. Carhart-Harris, Leor Roseman, David K. Menon, and E. A. Stamatakis. Serotonergic Psychedelics LSD & Psilocybin Increase the Fractal Dimension of Cortical Brain Activity in Spatial and Temporal Domains. *bioRxiv*, page 517847, January 2019.

25. Suresh D. Muthukumaraswamy and David TJ. Liley. 1/f electrophysiological spectra in resting and drug-induced states can be explained by the dynamics of multiple oscillatory relaxation processes. *NeuroImage*, 179:582–595, October 2018.

26. Robin Lester Carhart-Harris, Robert Leech, Peter John Hellyer, Murray Shanahan, Amanda Feilding, Enzo Tagliazucchi, Dante R. Chialvo, and David Nutt. The entropic brain: a theory of conscious states informed by neuroimaging research with psychedelic drugs. *Frontiers in Human Neuroscience*, 8, 2014.

27. Robin L. Carhart-Harris. The entropic brain - revisited. *Neuropharmacology*, 142:167–178, November 2018.

28. Antonio J. Fontenele, Nivaldo A. P. de Vasconcelos, Thaís Feliciano, Leandro A. A. Aguiar, Carina Soares-Cunha, Bárbara Coimbra, Leonardo Dalla Porta, Sidarta Ribeiro, Ana João Rodrigues, Nuno Sousa, Pedro V. Carelli, and Mauro Copelli. Criticality between Cortical States. *Physical Review Letters*, 122(20):208101, May 2019.

29. Gregory Scott, Erik D. Fagerholm, Hiroki Mutoh, Robert Leech, David J. Sharp, Woodrow L. Shew, and Thomas Knöpfel. Voltage Imaging of Waking Mouse Cortex Reveals Emergence of Critical Neuronal Dynamics. *Journal of Neuroscience*, 34(50):16611–16620, December 2014.

30. Tomer Fekete, David B. Omer, Kazunori O'Hashi, Amiram Grinvald, Cees van Leeuwen, and Oren Shriki. Critical dynamics, anesthesia and information integration: Lessons from multi-scale criticality analysis of voltage imaging data. *NeuroImage*, 183:919–933, December 2018.

31. Heonsoo Lee, Daniel Golkowski, Denis Jordan, Sebastian Berger, Rüdiger Ilg, Joseph Lee, George A. Mashour, UnCheol Lee, Michael S. Avidan, Stefanie Blain-Moraes, Goodarz Golmirzaie, Randall Hardie, Rosemary Hogg, Ellen Janke, Max B. Kelz, Kaitlyn Maier, George A. Mashour, Hannah Maybrier, Andrew McKinstry-Wu, Maxwell Muench, Andrew Ochroch, Ben J. A. Palanca, Paul Picton, E. Marlon Schwarz, Vijay Tarnal, Giancarlo Vanini, and Phillip E. Vlisides. Relationship of critical dynamics, functional connectivity, and states of consciousness in large-scale human brain networks. *NeuroImage*, December 2018.

32. Yasuo Nagasaka, Kentaro Shimoda, and Naotaka Fujii. Multidimensional recording (MDR) and data sharing: an ecological open research and educational platform for neuroscience. *PloS One*, 6(7):e22561, 2011.

33. Toru Yanagawa, Zenas C. Chao, Naomi Hasegawa, and Naotaka Fujii. Large-Scale Information Flow in Conscious and Unconscious States: an ECoG Study in Monkeys. *PLoS ONE*, 8(11), November 2013.

34. Hugh C. Hemmings, Paul M. Riegelhaupt, Max B. Kelz, Ken Solt, Roderic G. Eckenhoff, Beverley A. Orser, and Peter A. Goldstein. Towards a Comprehensive Understanding of Anesthetic Mechanisms of Action: A Decade of Discovery. *Trends in Pharmacological Sciences*, 40(7):464–481, July 2019.

35. Khurram Saleem Khan, Ivan Hayes, and Donal J. Buggy. Pharmacology of anaesthetic agents I: intravenous anaesthetic agents. *Continuing Education in Anaesthesia Critical Care & Pain*, 14(3):100–105, June 2014.

36. Panos Zanos, Ruin Moaddel, Patrick J. Morris, Lace M. Riggs, Jaclyn N. Highland, Polymnia Georgiou, Edna F. R. Pereira, Edson X. Albuquerque, Craig J. Thomas, Carlos A. Zarate, and Todd D. Gould. ketamine and ketamine Metabolite Pharmacology: Insights into Therapeutic Mechanisms. *Pharmacological Reviews*, 70(3):621–660, July 2018.

37. E. F. Domino, P. Chodoff, and G. Corssen. Pharmacologic effects of CI-581, a new dissociative anesthetic, in man. *Clinical Pharmacology and Therapeutics*, 6:279–291, June 1965.

38. J. H. Krystal, L. P. Karper, J. P. Seibyl, G. K. Freeman, R. Delaney, J. D. Bremner, G. R. Heninger, M. B. Bowers, and D. S. Charney. Subanesthetic effects of the noncompetitive NMDA antagonist, ketamine, in humans. Psychotomimetic, perceptual, cognitive, and neuroendocrine responses. *Archives of General Psychiatry*, 51(3):199–214, March 1994.

39. Ritta Hari and Aina Puce. *MEG-EEG Primer*. Oxford University Press, 2017.

40. Guillermo Solovey, Kai J. Miller, Jeffrey G. Ojemann, Marcelo O. Magnasco, and Guillermo A. Cecchi. Self-Regulated Dynamical Criticality in Human ECoG. *Frontiers in Integrative Neuroscience*, 6, July 2012.

41. Leandro M. Alonso, Alex Proekt, Theodore H. Schwartz, Kane O. Pryor, Guillermo A. Cecchi, and Marcelo O. Magnasco. Dynamical criticality during induction of anesthesia in human ECoG recordings. *Frontiers in Neural Circuits*, 8, 2014.

42. G. Tononi, O. Sporns, and G. M. Edelman. A measure for brain complexity: relating functional segregation and integration in the nervous system. *Proceedings of the National Academy of Sciences*, 91(11):5033–5037, May 1994.

43. Arnaud Delorme and Scott Makeig. EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of Neuroscience Methods*, 134(1):9–21, March 2004.

44. Alexandre Gramfort, Martin Luessi, Eric Larson, Denis A. Engemann, Daniel Strohmeier, Christian Brodbeck, Roman Goj, Mainak Jas, Teon Brooks, Lauri Parkkonen, and Matti Hämäläinen. MEG and EEG data analysis with MNE-Python. *Frontiers in Neuroscience*, 7, 2013.

45. Christian Meisel, Eckehard Olbrich, Oren Shriki, and Peter Achermann. Fading Signatures of Critical Brain Dynamics during Sustained Wakefulness in Humans. *The Journal of Neuroscience*, 33(44):17363–17372, October 2013.

46. Per Bak and Kan Chen. Self-Organized Criticality. *Scientific American*, 264(1):46–53, 1991.

47. J. Matias Palva, Alexander Zhigalov, Jonni Hirvonen, Onerva Korhonen, Klaus Linkenkaer-Hansen, and Satu Palva. Neuronal long-range temporal correlations and avalanche dynamics are correlated with behavioral scaling laws. *Proceedings of the National Academy of Sciences of the United States of America*, 110(9):3585–3590, February 2013.

48. Enzo Tagliazucchi, Pablo Balenzuela, Daniel Fraiman, and Dante R. Chialvo. Criticality in Large-Scale Brain fMRI Dynamics Unveiled by a Novel Point Process Analysis. *Frontiers in Physiology*, 3, 2012.

49. Najja Marshall, Nicholas M. Timme, Nicholas Bennett, Monica Ripp, Edward Lautzenhiser, and John M. Beggs. Analysis of Power Laws, Shape Collapses, and Neural Complexity: New Techniques and MATLAB Support via the NCC Toolbox. *Frontiers in Physiology*, 7, 2016.

50. A. Clauset, C. Shalizi, and M. Newman. Power-Law Distributions in Empirical Data. *SIAM Review*, 51(4):661–703, November 2009.

51. M. E. J. Newman. Power laws, Pareto distributions and Zipf's law. *Contemporary Physics*, 46(5):323–351, September 2005. arXiv: cond-mat/0412004.

52. S.M. Burroughs and S.F. Tebbens. Upper-truncated Power Laws in Natural Systems. *pure and applied geophysics*, 158(4):741–757, April 2001.

53. Shan Yu, Andreas Klaus, Hongdian Yang, and Dietmar Plenz. Scale-Invariant Neuronal Avalanche Dynamics and the Cut-Off in Size Distributions. *PLOS ONE*, 9(6):e99761, June 2014.

54. Viola Priesemann, Matthias HJ Munk, and Michael Wibral. Subsampling effects in neuronal avalanche distributions recorded in vivo. *BMC Neuroscience*, 10(1):40, April 2009.

55. Tiago L. Ribeiro, Sidarta Ribeiro, Hindiael Belchior, Fábio Caixeta, and Mauro Copelli. Undersampled Critical Branching Processes on Small-World and Random Networks Fail to Reproduce the Statistics of Spike Avalanches. *PLOS ONE*, 9(4):e94992, April 2014.

56. S. Watanabe. Information Theoretical Analysis of Multivariate Correlation. *IBM Journal of Research and Development*, 4(1):66–82, January 1960.

57. J. Ziv and A. Lempel. Compression of individual sequences via variable-rate coding. *IEEE Transactions on Information Theory*, 24(5):530–536, September 1978.

58. S. Behnel, R. Bradshaw, C. Citro, L. Dalcin, D. S. Seljebotn, and K. Smith. Cython: The Best of Both Worlds. *Computing in Science Engineering*, 13(2):31–39, March 2011.

59. S. van der Walt, S. C. Colbert, and G. Varoquaux. The NumPy Array: A Structure for Efficient Numerical Computation. *Computing in Science Engineering*, 13(2):22–30, March 2011.

60. Eric Jones and Travis Oliphant. SciPy, Open source scientific tools for Python, 2001.

61. Wes McKinney. Data Structures for Statistical Computing in Python. In *Proceedings of the 9th Python in Science Conference*, pages 51 – 56, 2010.

62. Giulio Tononi. Consciousness as Integrated Information: a Provisional Manifesto. *The Biological Bulletin*, 215(3):216–242, December 2008.

63. Simone Sarasso, Melanie Boly, Martino Napolitani, Olivia Gosseries, Vanessa Charland-Verville, Silvia Casarotto, Mario Rosanova, Adenauer Girardi Casali, Jean-Francois Brichant, Pierre Boveroux, Steffen Rex, Giulio Tononi, Steven Laureys, and Marcello Massimini. Consciousness and Complexity during Unresponsiveness Induced by propofol, Xenon, and ketamine. *Current Biology*, 25(23):3099–3105, December 2015.

64. Michael Schartner, Anil Seth, Quentin Noirhomme, Melanie Boly, Marie-Aurelie Bruno, Steven Laureys, and Adam Barrett. Complexity of Multi-Dimensional Spontaneous EEG Decreases during propofol Induced General Anaesthesia. *PLOS ONE*, 10(8):e0133532, August 2015.

65. Jisung Wang, Gyu-Jeong Noh, Byung-Moon Choi, Seung-Woo Ku, Pangyu Joo, Woo-Sung Jung, Seunghwan Kim, and Heonsoo Lee. Suppressed neural complexity during ketamine- and propofol-induced unconsciousness. *Neuroscience Letters*, 653:320–325, July 2017.

66. Varley, T. F., Craig, M., Adapa, R., Finoia, P., Williams, G., Allanson, J., Pickard, J., Menon, D. K., Stamatakis, E. A. Fractal dimension of cortical functional connectivity networks & severity of disorders of consciousness. PLOS ONE, 15(2), e0223812. https://doi.org/10.1371/journal.pone.0223812 (2020)

67. Varley, T. F., Luppi, A. I., Pappas, I., Naci, L., Adapa, R., Owen, A. M., Menon, D. K., Stamatakis, E. A. Consciousness & Brain Functional Complexity in propofol Anaesthesia. Scientific Reports, 10(1), 1–13. https://doi.org/10.1038/s41598-020-57695-3 (2020)

68. Adenauer G. Casali, Olivia Gosseries, Mario Rosanova, Mélanie Boly, Simone Sarasso, Karina R. Casali, Silvia Casarotto, Marie-Aurélie Bruno, Steven Laureys, Giulio Tononi, and Marcello Massimini. A Theoretically Based Index of Consciousness Independent of Sensory Processing and Behavior. *Science Translational Medicine*, 5(198):198ra105–198ra105, August 2013.

69. Michael M. Schartner, Andrea Pigorini, Steve A. Gibbs, Gabriele Arnulfo, Simone Sarasso, Lionel Barnett, Lino Nobili, Marcello Massimini, Anil K. Seth, and Adam B. Barrett. Global and local complexity of intracranial EEG decreases during NREM sleep. *Neuroscience of Consciousness*, 2017(1), January 2017.

70. Michael M. Schartner, Robin L. Carhart-Harris, Adam B. Barrett, Anil K. Seth, and Suresh D. Muthukumaraswamy. Increased spontaneous MEG signal diversity for psychoactive doses of ketamine, LSD and psilocybin. *Scientific Reports*, 7:46421, April 2017.

71. Laura D. Lewis, Veronica S. Weiner, Eran A. Mukamel, Jacob A. Donoghue, Emad N. Eskandar, Joseph R. Madsen, William S. Anderson, Leigh R. Hochberg, Sydney S. Cash, Emery N. Brown, and Patrick L. Purdon. Rapid fragmentation of neuronal networks at the onset of propofol-induced unconsciousness. *Proceedings of the National Academy of Sciences of the United States of America*, 109(49):E3377–3386, December 2012.

72. George A. Mashour and Anthony G. Hudetz. Neural Correlates of Unconsciousness in Large-Scale Brain Networks. *Trends in Neurosciences*, 41(3):150–160, 2018.

73. Stanislas Dehaene. *Consciousness and the Brain: Deciphering How the Brain Codes Our Thoughts*. Penguin, January 2014. Google-Books-ID: CWw2AAAAQBAJ.

74. Richard Robinson. Exploring the "Global Workspace" of Consciousness. *PLOS Biology*, 7(3):e1000066, March 2009.

75. R. Ferenets, Tarmo Lipping, A. Anier, V. Jantti, S. Melto, and S. Hovilehto. Comparison of entropy and complexity measures for the assessment of depth of sedation. *IEEE Transactions on Biomedical Engineering*, 53(6):1067–1077, June 2006.

76. Goran Kekovic, Gordana Stojadinovic, Ljiljana Martac, Jelena Podgorac, Slobodan Sekulic, and Milka Culic. Spectral and fractal measures of cerebellar and cerebral activity in various types of anesthesia. *Acta Neurobiologiae Experimentalis*, 70(1):67–75, 2010.

77. Srdjan Kesić and Sladjana Z. Spasić. Application of Higuchi's fractal dimension from basic to clinical neurophysiology: A review. *Computer Methods and Programs in Biomedicine*, 133:55–70, September 2016.

78. J. Ruiz de Miras, F. Soler, S. Iglesias-Parro, A. J. Ibáñez-Molina, A. G. Casali, S. Laureys, M. Massimini, F. J. Esteban, J. Navas, and J. A. Langa. Fractal dimension analysis of states of consciousness and unconsciousness using transcranial magnetic stimulation. *Computer Methods and Programs in Biomedicine*, 175:129–137, July 2019.

79. Liu, X., Ward, B. D., Binder, J. R., Li, S.-J., Hudetz, A. G. Scale-Free Functional Connectivity of the Brain Is Maintained in Anesthetized Healthy Participants but Not in Patients with Unresponsive Wakefulness Syndrome. PLOS ONE, 9(3), e92182. https://doi.org/10.1371/journal.pone.0092182

80. Villani, C. (2009). Optimal Transport: Old and New. Springer-Verlag. https://doi.org/10.1007/978-3-540-71050-9

81. Mashour, G. A., Roelfsema, P., Changeux, J.-P., & Dehaene, S. (2020). Conscious Processing and the Global Neuronal Workspace Hypothesis. Neuron, 105(5), 776–798. https://doi.org/10.1016/j.neuron.2020.01.026

82. Enzo Tagliazucchi, Pablo Balenzuela, Daniel Fraiman, and Dante R. Chialvo. Criticality in Large-Scale Brain fMRI Dynamics Unveiled by a Novel Point Process Analysis. *Frontiers in Physiology*, 3, 2012.

83. Enzo Tagliazucchi, Robin Carhart-Harris, Robert Leech, David Nutt, and Dante R. Chialvo. Enhanced repertoire of brain dynamical states during the psychedelic experience. *Human Brain Mapping*, 35(11):5442–5456, November 2014. arXiv: 1405.6466.

84. Nicholas Timme, Wesley Alford, Benjamin Flecker, and John M. Beggs. Synergy, redundancy, and multivariate information measures: an experimentalist's perspective. *Journal of Computational Neuroscience*, 36(2):119–140, April 2014.

85. Nicholas M. Timme, Shinya Ito, Maxym Myroshnychenko, Sunny Nigam, Masanori Shimono, Fang-Chin Yeh, Pawel Hottowy, Alan M. Litke, and John M. Beggs. High-Degree Neurons Feed Cortical Computations. *PLoS computational biology*, 12(5):e1004858, 2016.

86. Samantha P. Faber, Nicholas M. Timme, John M. Beggs, and Ehren L. Newman. Computation is concentrated in rich clubs of local cortical networks. *Network Neuroscience*, pages 1–21, September 2018.

87. Sarah Potez and Matthew E. Larkum. Effect of common anesthetics on dendritic properties in layer 5 neocortical pyramidal neurons. *Journal of Neurophysiology*, 99(3):1394–1407, March 2008.

88. Mototaka Suzuki and Matthew E. Larkum. General Anesthesia Decouples Cortical Pyramidal Neurons. *Cell*, 180(4):666–676.e13, February 2020.

89. Nelson Spruston. Pyramidal neurons: dendritic structure and synaptic integration. *Nature Reviews. Neuroscience*, 9(3):206–221, March 2008.

90. Beggs, J. M. and Timme, N Being Critical of Criticality in the Brain. *Frontiers in Physiology* , 3. 2012

91. Ma, Zhengyu and Turrigiano, Gina G. and Wessel, Ralf and Hengen, Keith B. Cortical Circuit Dynamics Are Homeostatically Tuned to Criticality In Vivo Neuron, 2019 10.1016/j.neuron.2019.08.031,