The *Hoja loca1* Mutant and AUX/IAA Function in Phytomer Development

Richardson A. E. *[1,2,3#], Sluis A.[1,2,#], Hake S.[1,2],

(1) USDA Plant Gene Expression Center, 800 Buchanan Street, Albany, CA 94710.

(2) Department of Plant and Microbial Biology, University of California Berkeley, CA.

(3) [§]Institute of Molecular Plant Science, School of Biological Sciences, The University of Edinburgh, Daniel Rutherford Building, Max Born Crescent, Kings Buildings, Edinburgh, EH9 3BF

[#] these authors contributed equally to the article.

* corresponding author

[§] current address

Short Title: *Hoja loca1* and AUX/IAA Function

Abstract

Plant architecture is determined by the iterative building of phytomer units, each containing a portion of stem, an organ and an axillary meristem. Each phytomer can follow different developmental paths underpinning the complexity, and plasticity, of plant form. Auxin plays a central role in the coordination of phytomer development, regulating organ initiation and patterning across all axes. This diversity in auxin function results from changes in the activities and expression of auxin signaling components, including the AUX/IAA repressors. Higher land plants have multigene AUX/IAA families, which leads to functional redundancy. Dominant mutations, which prevent AUX/IAA degradation in response to auxin, have highlighted the importance of these proteins in fine-tuning organ development. Here we report a new dominant AUX/IAA mutant in maize, *Hoja loca1* (*Oja*). *Oja* has a mutation in the degron motif of ZmIAA28 and specifically affects aerial organ initiation and leaf medio-lateral patterning, but phytomer initiation remains unchanged. Mutant phenotypes are variable and transcriptional profiling of individual meristems identified clusters of genes that may underpin the phenotypic differences. The unique phenotype of *Oja* provides evidence of species-

specific sub-functionalization of the AUX/IAAs, and illustrates the crucial role of auxin

signaling in the tight coordination of phytomer unit development.

Introduction

The iterative post-embryonic development of plants has led to impressive

developmental plasticity, which allows plants to modify their architecture over time and

respond to environmental conditions. Organs initiate from meristems, groups of self-

organizing totipotent cells, in regular phyllotactic patterns. Initiation of each leaf includes

an axillary meristem and an internode. This developmental unit of leaf, axillary

meristem, and internode is known as a phytomer. Aerial plant development can

therefore be understood as the successive formation of phytomer units (Galinat, 1959),

with developmental plasticity manifesting in different amounts of growth in the three

components.

The fate of cells within the meristem is dependent on position. Cells in the center of

shoot meristems divide slowly, are isodiametric in shape, and are considered

indeterminate. As divisions push cells from the center outward, cells enter the peripheral

or morphogenetic zone, where organs initiate. It is here that the primordium is specified,

forming a leaf at plastochron 0 (P_0) (Bolduc et al., 2012a). The *KNOTTED1-LIKE*

HOMEOBOX (*KNOX*) genes mark indeterminate cell fates (Jackson et al., 1994; Smith

et al., 1992), and are down-regulated in P_0 cells. *KNOX* gene expression is maintained

within the other components of the phytomer, the internode and bud, until they elongate

(Tsuda et al., 2017). The downregulation of *KNOX* genes in the P_0 is preceded by the

formation of an auxin maximum, coordinated by a convergence point of the auxin

transporter PINFORMED1 (PIN1) in the epidermis (Bayer et al., 2009; Benková et al.,

2003; O'Connor et al., 2014; Reinhardt et al., 2003). Auxin acts by triggering the

degradation of AUX/IAA proteins, allowing AUXIN RESPONSE FACTORS (ARFs) to

regulate downstream genes (Lavy and Estelle, 2016; Salehin et al., 2015). These

coordinated events comprise the mechanism for organ initiation.

As organs develop and differentiate, asymmetries are established that determine tissue

identity across proximal/distal, abaxial/adaxial, and medial/lateral axes (Kuhlemeier and

Timmermans, 2016; Richardson and Hake, 2019). In grasses, the leaf initiates as a

ring-shaped primordium around the meristem with the separation of the lower leaf

(sheath) margins occurring later in development (Johnston et al., 2015; Sharman,

1942). Proximal-distal patterning in the grass leaf leads to the formation of a distal blade

and a proximal sheath. The ligule and auricle form at the boundary of blade and sheath.

For most grasses, the medial/lateral axes are marked by a midvein at the center which

is thickened in comparison with the margins of the leaf (referred to as the midrib).

Auxin signaling has been implicated in all stages and axes of grass leaf development.

Treatment of maize shoot apices, with the auxin inhibitor NPA (*N-1*-Naphthylphthalamic

Acid) limits organ initiation and prevents the separation of the sheath margins resulting

in a tube leaf (Scanlon, 2003). PIN proteins are also expressed at the position of a ligule

prior to its elaboration (Moon et al., 2013) and RNAseq analysis of leaf microdomains

has shown that ligule development involves the recapitulation of the organ initiation

mechanism (Johnston et al., 2014). The establishment of the abaxial/adaxial axis also

involves auxin signaling, utilizing opposing gradients of tasiR-ARFs and miR166

(Nogueira et al., 2007). Despite the role of auxin signaling throughout grass leaf

development, auxin signaling mutants (i.e. mutants in AUX/IAA or ARFs), which affect

leaf development, have not yet been reported in grasses.

Here we describe a dominant mutant, named *Hoja loca1* (*Oja*), with a range of leaf

defects. In its most severe expression, mutants fail to initiate leaves, although an

internode will be partitioned to each phytomer. In the least severe, midveins are missing

from otherwise correctly patterned leaves. Between these extremes, tube leaves are

formed, where the leaf margin is completely fused. Remarkably, leaves maintain

proximal distal and abaxial/adaxial patterning. Using positional cloning and whole

genome sequencing, we identify the gene as the AUX/IAA protein, ZmIAA28

(GRMZM2G035465). By analysing individual meristems, we were able to use the

variable phytomer phenotypes of *Oja* to dissect potential regulatory roles of ZmIAA28

that may be responsive to different auxin concentrations. Unlike other maize AUX/IAA

mutants that specifically affect roots (Woll et al., 2005) or inflorescences (Galli et al.,

2015), *Oja* provides a view into the unique role of AUX/IAA proteins in grass leaf

patterning. The ability of *Oja* to produce a leafless phytomer also hints at the role of

auxin signaling in coordinating development of the separate phytomer components.

Results

Hoja loca1

Hoja loca1 (*Oja*) is a dominant mutant that was discovered in a forward genetics EMS

screen. The phenotype is variable, ranging from mild to severe, and it exhibits

unpredictable expression through generations; crosses made with mildly affected plants

give rise to severe plants and vice versa.

***Hoja loca1* leaves have medial/lateral defects but normal proximal/distal and**

abaxial/adaxial patterning in aerial organs

Maize leaves are divided into a proximal sheath that wraps around the inner leaves and

a distal blade that bends away from the plant, optimized for photosynthesis (Figure 1Ai).

At the junction of the blade and sheath is the ligule, an epidermal fringe on the adaxial

surface and two auricles along the margin that allow the blade to tilt away from the

sheath (Figure 1Aii-iv). The blade contains a midrib, a thickened region with chlorophyll-

containing cells on the abaxial surface and clear cells on the adaxial surface (Figure

1Av), that supports the extended blade.

In contrast to wild-type seedlings, *Hoja loca1* (*Oja*) mutants have varied leaf

phenotypes present on the same plant, including normal leaves (Figure 1B,C). The

mildest leaf mutant phenotype has missing midribs (leaf 2 in Figure 1Bi). These leaves

droop backwards without the structural support of the midrib (Figure 1Bi-v). The

midribless leaves completely lack adaxial clear cells, although the flanking blade

regions appear normal (Figure 1Bv). Midribless leaves still have normal ligules and

auricles (Figure 1Bii-iv, arrowheads) suggesting that the midrib is not essential in

elaboration of proximal/distal patterning.

Severe mutant leaf phenotypes include tube leaves in which the margins are not

separated (Figure 1C). As in the case for midribless leaves, tube leaves also have

normal proximal/distal patterning, developing a normal ligule between the sheath and

blade (Figure Cii-iv). Cells in the tube leaves also have correct abaxial/adaxial polarity

with chlorophyll containing cells on the abaxial surface and clear cells on the adaxial

surface (Figure 1Cv). These results suggest that *Oja* does not influence proximal/distal

or abaxial/adaxial patterning.

To determine whether there was a pattern in initiation of different leaf

phenotypes, we recorded the occurrence of midribless and tube leaves in a population

of 150 plants (Figure 1D). Both midribless and tube leaves can be observed in the same

plant. In all cases, the first leaf is never affected. The second leaf may show one of

these defects and the third and fourth are most likely to be mutant (Figure 1D). The

frequency for both defects decreases during development but tube leaves could be

observed late in development. This variability of phytomer phenotype makes *Oja* difficult

to work with, but also provides a unique opportunity to dissect the different functions of

the underlying gene.

In many *Oja* plants, inflorescence branching and floral development was severely

affected. Both male and female inflorescences bore bare regions indicative of a failure

to initiate spikelet primordia, and the tassels were not branched. However, when flowers

did form, they appeared to be normal and fertile (Supplementary Figure 1). In contrast

to aerial development, root development in seedlings appeared to be normal with

initiation of primary, seminal and lateral roots and a normal gravitropic response

(Supplementary Figure 2). *Oja* therefore specifically affects aerial organ development.

Hoja loca1 leaves have defects in vein distribution, but normal vein density.

Given the midrib defect, we next questioned whether veins are altered in *Oja* mutants.

Seedling leaf cross-sections showed that the abaxial/adaxial patterning of vascular

bundles appears normal (Figure 2A,B), mirroring the normal abaxial/adaxial patterning

of the entire organ.

To assess whether the medio-lateral patterning of vascular bundles was altered

we measured leaf width and determined vein density and distribution in the third leaf of

3-week-old seedlings. Midribless leaves were narrower than wild type and tube leaves
were less than half the width of wild type (Figure 2C). Overall vein density was not
affected by *Oja* (Figure 2D) suggesting a reduction in total vein number to match the
decrease in size. To assess vein distribution, we counted the number of intermediate
veins between lateral veins in both wild-type and midribless leaves. In wild type, the
number of intermediate veins shows a symmetry across the leaf, with a dip at the mid
position (Figure 2Eii, blue). We did not identify a dip of intermediate veins in the midvein
position in *Oja* midribless leaves (Figure 2Eii, orange), supporting the hypothesis that
medio-lateral patterning is affected in the *Oja* mutant.
162

Hoja loca1 mutants fail to initiate leaves but still set aside internode cells
At each leaf initiation, cells are set aside for the leaf and the internode below (Johri and
Coe, 1983). A mature maize plant is thus composed of alternating internodes and
nodes, which is the position of leaf insertion. In mature *Oja* plants we observed bare
nodes, for example the plant in Figure 3A ended in a tube leaf after four nodes failed to
elaborate leaves (arrows). To assess whether these bare nodes were due to ectopic
nodes or a failure of leaf initiation, we counted the number of nodes in *Oja* versus wild-
type siblings and compared these to the number of leaves. In mature plants there is no
significant difference in number of nodes (Figure 3B), but there are significantly fewer
leaves in *Oja* mutants (Figure 3C). This alternating node, internode, node pattern
without accompanying leaves is also observed when slicing through developing tube
leaves (Figure 3D). These data suggest that *Oja* may be able to initiate new phytomers
consisting of just nodes and internodes.
To identify whether leaves failed to initiate, or whether they initiated but failed to
elaborate, we used scanning electron microscopy (SEM) to examine the earliest
primordial stages of leaf development. Wild-type maize leaves at the P_1 stage are a ring
of cells that surround the meristem. At P_2, the leaf begins to grow in the proximal distal
dimension, with growth highest at the midvein position. By P_3, the leaf primordium
covers the top of the meristem and the margins begin to expand (Figure 3E). Further
growth of the leaf margins at the P_4 stage envelopes the younger leaves and meristem
(Hay and Hake, 2004; Scanlon et al., 1996). Longitudinal growth of the internode and

sheath occur later in development. Compared to wild-type seedlings where developing leaf primordia always obscured the internodes, the internodes were often visible in severe *Oja* shoots (Figure 3F-I). The meristem in Figure 3F has a single ring primordium which may be the early stages of a tube leaf, and below this ring, no older leaves have initiated, leaving an elongated, exposed internode. The meristem in Figure 3G initiated five internodes without leaves, before more leaf primordia were initiated. In some *Oja* mutants, the leaf initiation defect was so severe that the shoot apex ended in pin-like barren meristems as in Figure 3H. In many mutants, axillary meristems formed which contained more normal leaf initiation (Figure 3I). In fact, the mature plant in Figure 3A had a fertile tiller although the main shoot ended in a tube leaf with skipped leaf initiation. The SEM data suggest that the discrepancy between node number and leaf number in *Oja* is due to patterning during the earliest stages of leaf primordium initiation. This finding supports the hypothesis that *Oja* mutants fail to initiate leaves, despite initiating new phytomers.

The skipped leaf initiation also explains some of the seedling architecture differences. For example, the normal seedling in Figure 1A has four visible leaves, while the seedling in Figure 1B has only three leaves visible, with the second leaf the size of a normal third leaf. In addition, the distance between the first and second leaf is much greater than in the wild type. This observation suggests that leaf two was skipped in the mutant.

Hoja loca1 encodes the AUX/IAA protein, ZmIAA28.

To identify the causal mutation, we mapped the Mo17 *Oja* region in a backcross population to A619. Bulk segregant mapping placed *Oja* on chromosome 8 (Figure 4). SSR primers were used to narrow the position to an interval of 1 Mb containing 19 annotated genes (Figure 4Aiii). Using published RNAseq data we narrowed the candidate gene list further, as only 10 genes in the interval were reported as expressed in the shoot apical meristem or leaf primordia (Figure 4Aiii, green).

To further map the interval, we used whole genome sequencing of two bulk pools, one wild type and one mutant pool, each with 40 individuals. A large interval was detected on chromosome 8 between 75 and 135 Mb. To generate a candidate SNP list,

we removed SNPs reported in the HapMapv3 from Panzea (www.panzea.org/data) and

limited the reported SNPs to EMS-type G to A, or C to T. Within the large interval, there

were nine moderate or high-effect SNPs within annotated gene models (Figure 4Ai), but

only one SNP detected within the fine SSR mapping interval (Figure 4Aiii, *). This

change was a C to T nucleotide transition in GRMZM2G035465, which encodes

ZmIAA28 (Figure 4B). The C to T transition causes a missense mutation from a Glycine

to Glutamic Acid in the center of the highly conserved degron motif (Figure 4C). Similar

mutations in other AUX/IAA proteins prevent degradation in response to high auxin

(Reed, 2001) suggesting that *Oja* is caused by a stabilizing mutation in the degron

motif, preventing degradation of ZmIAA28 in the presence of auxin.

***Hoja loca1* shows dysregulation of auxin signaling**.

To assess the effect of the *Oja1* mutation on gene expression in vegetative meristem

and developing leaf primordia, we sequenced mRNA extracted from dissected

meristems with P1/P2 primordia attached from 3-week-old seedlings (Figure 5A). Given

the variability in phenotype, each individual was used to make a single library, with 36

libraries sequenced in total. By scoring the libraries based on the seedling leaf

phenotype, we classified the libraries into two groups: wild type and mutant (orange and

blue annotations indicated on the principal component analysis plot in Figure 5B).

Differential gene expression analysis identified 2877 significantly differentially

expressed genes (padj<0.05) with a log2FC of >0.59 or <0.59, of which 689 were down

in the mutant-scored samples compared to the wild type. GO term analysis showed

enrichment for many terms, including those associated with auxin signaling, meristem

regulation and leaf development (Figure 5C). Consistent with the phenotype of *Oja*,

genes associated with leaf development are down, and those associated with meristem

regulation are up in the mutant-scored samples compared to the wild type-scored

samples. This suggests that ZmIAA28 has a role in regulation of auxin response, and

leaf development.

Transcriptomes of individual *Hoja loca1* meristems and leaf primordia can differentiate four distinct transcriptional profiles, correlated with the four phytomer phenotypes.

To further investigate how the different leaf phenotypes of *Oja* may arise, we interrogated the transcriptional profiles of each individual sample. We asked whether we could identify four transcriptional profiles representing the four potential phenotypes of an initiating *Oja* phytomer: normal leaf, midribless leaf, tube leaf and no leaf initiation.

We first filtered the expressed gene list to those shown to be expressed in the meristem and developing leaf primordia based on (Knaeur 2019) to reduce noise caused by differing amounts of stem and internode tissue harvested with each sample. Using the top 1500 most variable of these genes, we carried out hierarchical clustering and identified four clusters using K-means (Figure 6A). Overlaying these clusters on a PCA plot of the top 100 most variable of all expressed genes showed clear separation of the four groups along PC1 (Figure 6B). This indicated that the four clusters (C1-4) were transcriptionally distinct.

To assess which cluster may be associated with each phytomer phenotype, we generated self-organizing maps (SOMs) of all expressed genes across C1-4 (Figure 6D) and annotated the nodes with genes involved in leaf initiation and development. A defining feature of initiating leaf primordia is the specific downregulation of *KNOX* genes in the P_0 and the activation of leaf patterning genes. For the four phytomer phenotypes; normal leaf, midribless leaf, tube leaf and no leaf initiation; we predicted that these genes would show distinct profiles. In phytomers initiating normal leaves we would expect low *KNOX* gene expression, with high leaf patterning gene expression, and the converse for phytomers failing to initiate leaves. Phytomers initiating midribless leaves and tubes leaves would have an intermediate profile. We found that most *KNOX* genes were in node 14, (Figure 6D) where expression was at the lowest level in C2 (Figure 6E), consistent with *KNOX* genes being downregulated during normal leaf initiation. Genes associated with leaf development were found in nodes where expression was highest in C2 (e.g. node 7 contains *CUP-SHAPED-COTYLEDON* genes and *WUSCHEL-LIKE-HOMEOBOX* genes known to be involved in leaf development and

organ initiation). This supports the hypothesis that C2 represents meristems initiating
phytomers with normal leaves.

Conversely, higher *KNOX* (node 14) and lower leaf development gene
expression (e.g. nodes 7 and 8) associated with C4 (Figure 6D). This is consistent with
the hypothesis that C4 represents phytomers with a 'no leaf initiation' phenotype. This is
further supported by the expectation that phytomers failing to initiate leaves will be most
transcriptionally distinct from those initiating normal leaves, and the PCA and
hierarchical clustering both show C4 as most different to C2 (Figure 6B). Combined,
these data suggests that C2 represents phytomers initiating normal leaves, and C4
represents phytomers with no leaf initiation.

To identify which of the remaining two clusters may correlate with either the tube
leaf or midribless leaf phytomer phenotype we analysed the expression profiles of
medio-lateral leaf patterning genes. We predicted that these genes would be
misregulated in phytomers initiating midribless or tube leaves, and be downregulated in
C4 as these phytomers are not initiating leaves.

Tube leaves lack leaf margins, and this should be reflected by reduced
expression of genes like the *WOX* gene *NARROWSHEATH1 (NS1)* which defines the
leaf margin (Nardmann et al., 2004; Scanlon and Freeling, 1997). Many *WOX* genes,
including *NS1*, are found in node 7 where expression is lowest in C3 and C4. Low *WOX*
expression in C3 could indicate that this group represents phytomers initiating tube-
shaped leaves.

Few genes have been identified as responsible for midribless identity, but the
YABBY genes *DROPPING LEAF 1* (*DRL1/IFA1*) and 2 (*DRL2*)(Strable et al., 2017)
both have defects in midrib development. We would predict that genes involved in
midrib development would be downregulated in phytomers initiating midribless leaves.
DRL1 and *DRL2* are located in node 8, where expression is highest in C2 and low in all
other clusters, making it difficult to differentiate between C1 and C3. Given that the
midribless leaf phenotype appears most similar to a normal leaf, we predict that the
cluster most similar to C2 would represent phytomers initiating midribless leaves. Based
on the PCA and the differential gene expression analysis, C1 was most similar in
transcriptional profile to C2 (Figure 6B,C, pink). We therefore hypothesise that C1

represents phytomers initiating midribless leaves, and C3 represents phytomers
initiating tube leaves.

Each phytomer phenotype can be associated with specific gene expression
changes.

To further test the hypothesis that each cluster represents a different phytomer
phenotype (C1- midribless leaf, C2- normal leaf, C3- tube leaf, C4- no leaf initiation), we
carried out differential gene expression analysis for each 'mutant' cluster (C1, 3 and 4)
in comparison with the hypothesised normal leaf representative (C2). Overlap of the
significantly differentially expressed gene lists (padj <0.05, log2FC<-0.59 or>0.59,
Figure 6C, E-H) showed distinct transcriptional profiles for each cluster, suggesting that
each phytomer phenotype may be underpinned by different gene expression changes
triggered by variable effects of ZmIAA28 stabilization on auxin signaling.

To identify genes potentially important for each phenotype we overlaid the
significantly differentially expressed genes (log2 fold change >0.59 or <-0.59) for each
comparison, and identified 170 genes specific to C1, 601 to C3, and 2288 to C4. C4
showed the most unique significantly differentially expressed genes, many of which are
important components of leaf initiation. These genes included the boundary genes
CUC2 and *CUC3*, which were significantly downregulated, consistent with their roles in
specification of the boundary between the primordium and meristem (Hibara et al.,
2006; Aida et al., 1999). This further supports the hypothesis that C4 represents
phytomers which fail to initiate leaves, and suggests ZmIAA28 degradation in the
presence of auxin is involved in triggering leaf initiation.

We predict that C1 and C3 represent phytomers initiating leaves with medial-
lateral patterning defects. *WOX* genes exhibited distinct gene expression profiles across
the clusters, with opposite expression profiles in C3 compared to C2. *WOX3b, 4, 5a* and
13 were significantly upregulated in C3, whereas *WOX3a, 9a, 9b,* and *NS1* were all
significantly downregulated (Figure 6F). This finding suggests that medial-lateral
patterning and marginal domain development is severely perturbed in C3, correlating
with the tube leaf phenotype. The analysis demonstrates that auxin signalling via

ZmIAA28 degradation is important in regulation of *WOX* gene expression for medial-
lateral domain patterning.

Midribless leaves show defects in midrib formation and in vein distribution in the
leaf blade. We therefore analysed the expression profiles of genes with potential roles in
midrib and vein patterning. *YABBY1, 9, 15, DRL1 and DRL2* are all significantly
downregulated in C1 compared to C2 (Figure 6G). This finding suggests that ZmIAA28
may link auxin signalling with midrib formation via regulation of DRL1 and 2. C1 also
showed significant downregulation of multiple *SHORT INTERNODES (SHI) / STYLISH*
(STY) / SHI RELATED SEQUENCE (SRS) genes (Figure 6H), which in Arabidopsis are
involved in vascular patterning (Baylis et al., 2013) and have been shown to be
regulated by AUX/IAA proteins (Singh et al., 2020). ZmIAA28 may therefore also be
involved in auxin regulation of *SRS* gene expression during grass leaf vascular
patterning.

ZmIAA28 is involved in the auxin maxima maintenance preceding leaf initiation
The no leaf initiation phenotype of *Oja* led us to ask if we could use *Oja* to dissect the
link between organ initiation and auxin maxima formation. Using the comparison
between C4 and C2, we first interrogated the expression profile of genes known to be
involved in organ initiation. We then used *in situ* hybridisation and immunolocalisation to
test gene expression in a spatial manner.

In C4, *KNOX* genes are significantly upregulated, indicating that *KNOX* gene
downregulation is aberrant in *Oja* meristems in which leaf initiation fails. We used
antibodies to KNOTTED1 (KN1) (Smith et al., 1992) in order to follow leaf initiation
defects in the meristem. KN1 accumulates in the meristem and unexpanded stem and is
clearly down-regulated at the P_0 region, remaining absent in normal maize leaves
(Figure 7A). In *Oja* mutants we see strong accumulation at the tip of the meristem but,
unlike the normal meristem, KN1 down-regulation appears on both sides of the
meristem, suggesting that this P_0 encircles the meristem (Figure 7B). The encircling
downregulation could be indicative of tube leaf initiation. This particular meristem has
also failed to initiate at least two leaves, and KN1 localisation is still maintained in the
internode region below the meristem (Figure 7B). This variable downregulation of *KNOX*

genes supports the hypothesis that ZmIAA28 may be upstream of the downregulation of *KNOX* genes in initiating primordia.

Preceding *KNOX* downregulation, the auxin efflux transporter PIN1 moves auxin in the epidermis to the site of leaf initiation, where an auxin maxima forms. From the P_0, PIN1 proteins move auxin toward pre-existing vasculature, predicting the path of provascular traces (Reinhardt et al., 2003). In many species outside of the *Brassica*, PIN1 proteins have duplicated and diversified function. Sister of PIN1 (SoPIN1) localizes at the leaf initiation site and is also detected in vasculature in grasses (O'Connor et al., 2014). PIN1a and PIN1b have roles in vascular formation and stem growth (O'Connor et al., 2017). We found that all PIN gene expression was significantly downregulated in C4 compared to C2, suggesting that the feedback mechanism that maintains auxin maxima during organ initiation may be defective in *Oja*. As SoPIN1 is essential for organ initiation in the grasses (O'Connor et al., 2017) and is redundantly involved in vascular development, we chose to focus on SoPIN1 accumulation in the plant. To determine whether SoPIN1 patterning was defective in *Oja* meristems, we used an antibody to SoPIN1 to detect protein accumulation at the P_0 site and in the developing pre-vascular strand in wild-type plants (Figure 7C). In wild-type meristems, SoPIN1 convergence points form in the epidermis of the meristem at the P_0, which persists through to the tip of the P_1 leaf (Figure 7C). Strong internal traces of SoPIN1 are also observed highlighting the developing vascular strands that link the primordium apex to the stem vascular system (Figure 7C). In the *Oja* mutant shown, which has a number of skipped leaf initiations and appears as a pin, the SoPIN1 localization is throughout the tip of the meristem rather than localizing strongly to the P_0 region (Figure 7D). Normal accumulation and polarization can be seen in a vein trace within the mutant meristem, but this does not connect to the epidermis. This result suggests that SoPIN1 can polarize and be coordinated across the tissue. However, SoPIN1 fails to form a strong convergence point needed at the P_0 in *Oja*, implying that ZmIAA28 may be involved in the feedback regulation of SoPIN1 at lateral organ initiation sites, essential for organ initiation. In support of this hypothesis we observe changes in the expression of several genes involved in PIN protein localisation, including the PINOID protein BIF2 (McSteen and Hake, 2001), which is significantly downregulated in C4 compared to C2.

Downstream of auxin maxima formation and *KNOX* gene downregulation is

activation of leaf specific genes, such as *DRL1* and *2* (Strable et al., 2017). DRL genes

are early markers of leaf patterning and are expressed at the P_0 and at the midrib in leaf

primordia. To explore the effect of *Oja* on *DRL* expression, we used *in situ* hybridization.

In the normal seedling, expression was detected in the P_0 and each subsequent

midvein region (Figure 7E). In the *Oja* mutant shown, *DRL* expression was similar at the

midvein region in normal developing leaves, however, there was no detectable

expression in the P_0 and fewer leaves had initiated. As DRL is involved in midrib

specification (Strable et al., 2017), this lack of DRL in the P_0 could be due to the lack of

leaf initiation or due to the formation of a midribless leaf. Nevertheless, as an early

marker of leaf patterning this expression pattern supports the hypothesis that *Oja*

mutants fail to specify the P_0 leaf correctly despite defining the rest of the phytomer.

In summary, these findings suggest that *Oja* has a defect in leaf initiation and

patterning, not just leaf outgrowth, which leads to the reduced leaf number compared to

node number.

DISCUSSION

The dominant *Hoja loca1* (*Oja*) mutant, caused by a mutation in ZmIAA28,

specifically affects aerial organ initiation and medial/lateral patterning in the leaf,

illustrating highly specific AUX/IAA function essential for the fine-tuning of plant

development by auxin. By analysing individual meristems, we were able to use the

variable phytomer phenotypes of *Oja* to dissect potential regulatory roles of ZmIAA28

that may be responsive to different auxin concentrations.

AUX/IAA proteins function as repressors in low auxin conditions by interacting

with ARF transcription factors and preventing their activation of auxin responsive genes.

An increase in auxin promotes an interaction between AUX/IAA and Transport Inhibitor

Response1 (TIR1), an F-box component of the SCF complex that targets proteins for

degradation. Binding of AUX/IAA to TIR1 in the presence of auxin leads to AUX/IAA

degradation and thus releases ARF repression (Dharmasiri et al., 2005; Kepinski and

Leyser, 2005). As the AUX/IAA gene family is large, redundancy has mostly hidden

loss-of-function phenotypes (Overvoorde et al., 2005). Instead, genetic analysis has

identified dominant AUX/IAA mutations that stabilize these normally short-lived proteins
by disrupting their proteolysis or reducing their ability to bind to TIR1 (Ouellet et al.,
2001; Ramos et al., 2001; Tian et al., 2003; Worley et al., 2000; Yang et al., 2004). Most
mutations, including *Oja*, are located in domain II in the highly conserved core degron
motif (Fukaki et al., 2002; Nagpal et al., 2000; Rogg et al., 2001; Rouse et al., 1998;
Zenser et al., 2001) which interacts with TIR1 in the presence of auxin (Tan et al.,
2007).

Nine clades of AUX/IAA proteins have been identified (Matthes et al., 2019),
allowing for significant subfunctionalization during the evolution of the AUX/IAA family.
The majority of *Arabidopsis* AUX/IAA mutants were first described by their agravitropic
root, reduced lateral roots, and/or shorter root phenotypes (Fukaki et al., 2002; Nagpal
et al., 2000; Rouse et al., 1998). Mutants that were identified by a shoot phenotype such
as *bodenlos*, *Shy2*-2 and *axr3* have pleiotropic phenotypes, also affecting the roots. For
example, the semi-dominant *Shy2* mutant was identified as a suppressor of a
phytochrome elongated hypocotyl phenotype but also has root defects and curled
leaves (Reed et al., 1998; Tian and Reed, 1999). Embryonic patterning defects were
identified in *bodenlos* homozygotes but mature heterozygous plants also have root
defects (Hamann et al., 2002). *Axr3* mutants develop curled leaves (Leyser et al., 1996;
Pérez-Pérez et al., 2010) due to defects in coordination between the abaxial and
adaxial surfaces of the leaf. When these mutant phenotypes are mapped to the
AUX/IAA clades, no obvious pattern is observed, suggesting that function is not
conserved within a clade. This finding highlights the importance of studying the specific
function of AUX/IAA proteins in multiple species, as genetic similarity is unlikely to
correspond to related functionality.

Our work suggests that ZmIAA28 acts specifically during aerial organ initiation,
potentially upstream of *KNOX* downregulation. This contrasts with the function of the
tomato class VIII AUX/IAA, SlIAA9. Tomato leaf margin complexity is dependent on the
ectopic expression of *KNOX* genes in the leaf lamina. However, reduction of SlIAA9
expression causes a loss of leaf margin complexity, independent of *KNOX* gene
expression as determined by RT-PCR (Wang et al., 2005). This suggests that although
ZmIAA28 and SlIAA9 are in the same clade, they differ in their relationship with *KNOX*

gene regulation. This further argues for the importance for studying specific AUX/IAA
functions in different species.

AUX/IAA mutants in maize appear to be less pleiotropic than *Arabidopsis*; their
defects are confined to specific parts of the plant. For example, the *rootless with*
undetectable meristem (*rum1*) mutant is deficient in primary and lateral root initiation
with no defects in the aerial parts (Woll et al., 2005). *rum1* encodes *ZmIAA10* and,
unlike other described AUX/IAA mutants that have a missense mutation in the degron,
rum1 carries a deletion that includes the GWPPV degron sequence in domain II (Von
Behrens et al., 2011). The *Bif1* (ZmIAA27) and *Bif4* (ZmIAA20) AUX/IAA mutants
reduce the branching in male and female inflorescences (Barazesh et al., 2009; Galli et
al., 2015), however there is no effect on vegetative growth (Galli et al., 2015). *Oja*
mutants have striking defects in leaf development but have normal roots and flowers if
they make inflorescences. This suggests that ZmIAA10, ZmIAA20, ZmIAA27 and
ZmIAA28 have tissue-specific functions. These specificities do not appear to arise from
tissue specific expression as, based on published RNAseq data, ZmIAA27 and
ZmIAA28 have similar expression patterns. *RUM1* is expressed at very low levels
compared to a close paralog which is expressed at high levels in many root tissues as
well as the leaf (Zhang et al., 2016). Perhaps the tissue specificity results from the
interaction of each AUX/IAA with different ARFs or different TIR proteins and these are
responsible for the observed phenotypes in the dominant mutants (Salehin et al., 2015).

Through investigating leaf initiation, we found that SoPIN1 failed to converge at
the P_0 in *Oja* mutants suggesting that ZmIAA28 plays a role in the feedback signaling
required for forming the convergence point that triggers organ initiation. A role for the
TIR/ARF/AUX/IAA module in regulating cellular polarisation has also been observed in
Arabidopsis in which induction of the ARF, *MONOPTEROS* (MP) in the *mp* mutant
background triggered spontaneous PIN1 convergence point formation and primordium
outgrowth in the normally barren *mp* meristem (Bhatia et al., 2016). Interestingly,
although leaf initiation was severely affected in *Oja*, many other organ initiation events
occurred normally. For example, ligule development requires reactivation of the organ
initiation pathway (Johnston et al., 2014) but *Oja* mutants have no ligule defect,
suggesting that different AUX/IAAs may be involved in ligule development.

Given that formation of the SoPIN1 convergence point at the P_0 stimulates
midvein development, it is likely that ZmIAA28 is required for the signaling that
promotes midrib formation at the midvein, influencing medio-lateral patterning and
possibly subsequent *DRL* expression. Supporting the role of auxin signaling upstream
of *DRL* expression, DAPseq analyses in maize have identified ARF binding sites
upstream of both maize *DRL* genes (Galli et al., 2018). In midribless *Oja* mutant leaves,
the midrib domain appears to be completely lost, which is more severe than in the *drl*
double mutant (Strable et al., 2017), suggesting that other factors, also influenced by
auxin signaling, are required to specify midrib differentiation in addition to DRL.

Mediolateral development in the leaf is driven by *WOX* gene activity (Conklin et
al., 2021; Zhang et al., 2020). Our transcriptional analyses highlight wide-spread miss-
regulation of *WOX* gene expression in *Oja*, with most significant downregulation
associated with potential tube leaf formation, and failure of leaf initiation. The formation
of tube-shaped leaves in maize can be induced by incubation with the auxin inhibitor
NPA (Scanlon, 2003) which leads to a failure of sheath margin separation, and this
correlates with a high accumulation of ZmPIN1a in this region at the P4 stage of leaf
development (Johnston et al., 2015). Recent work has also shown that NS1 is induced
by auxin treatment (Conklin et al., 2021). This suggests that ZmIAA28 may be involved
in translation of the auxin concentration signal to leaf margin specification and later
margin development.

Analysis of proximal-distal patterning mutants gave rise to the hypothesis that the
midrib domain acts as a signaling centre to coordinate ligule development from the
midrib to the leaf margins (Becraft and Freeling, 1991; Harper and Freeling, 1996; Moon
et al., 2013). However, proximal-distal patterning is normal in *Oja* leaves lacking the
midrib. This suggests that it is not the midrib domain itself that links medial-lateral and
proximal-distal patterning but potentially medial-lateral patterning factors that are largely
independent of ZmIAA28 regulation of midrib formation.

The variability of the *Oja* phenotype is not unusual, having also been observed in
other AUX/IAA mutants such as in the *Arabidopsis* mutant *bodenlos,* with some mutants
missing hypocotyl and root and others just missing a root (Hamann et al., 2002). These
variable phenotypes could arise from stochastic over- and under- compensation of the

auxin signaling network. The sensitivity of different tissues to these stochastic changes

in auxin concentration could arise through different combinations of TIRs, ARFs and

other auxin signaling components (Powers and Strader, 2020; Parry et al., 2009;

Havens et al., 2012). In such a scenario, the four clusters of gene expression and

corresponding phytomer phenotypes identified in this work, may represent stable end-

states that arise from perturbations of the auxin pathway very early in development. As

such, when ZmIAA28 is not degraded in the presence of high auxin, as it is in wild type,

the three mutant phytomer phenotypes result from sub-optimal compensatory

mechanisms cascading to islands of stability. *Oja* therefore reveals three regions where

response to high auxin concentrations during the earliest phases of leaf development is

crucial in defining leaf shape: initiation, formation of a midrib/ vascular patterning, and

separation of the leaf margins.

The phytomer consists of a node, internode, axillary bud and organ. To maintain

correct plant architecture, the development of each component is tightly regulated. *Oja*

is unusual in that this coordination within the phytomer unit is apparently broken,

allowing the initiation of nodes and internodes without accompanying organs. This

finding suggests that auxin is not only involved in the initiation of organs, but also in the

regulation of phytomer development as a whole. The *Oja* mutant illustrates the species-

specific subfunctionalisation of the AUX/IAA proteins in regulating plant development. In

contrast to other known AUX/IAAs, ZmIAA28 specifically affects initiation of aerial

organs and impacts medio-lateral patterning. As further studies identify functions of

other grass AUX/IAAs it will be of interest to compare them to *Oja* to see how far this

subfunctionalisation holds. This will be particularly interesting to compare to *Arabidopsis*

aux/iaa mutants which appear to be more pleiotropic, and to determine the reasons for

the species-specific differences.

Methods

Isolation of *Hoja loca1*

Hoja loca1 (*Oja*) was discovered by a forward genetic approach. Pollen of the Mo17 maize inbred was treated with ethyl methanesulfonate (EMS) and crossed onto the B73 inbred. Hybrid F1 seed were planted in the field and screened. The single mutant plant was open pollinated and crossed to multiple inbreds the following season. Further crosses established the dominance of the mutant phenotype although often less than 50% of the plants showed the phenotype, indicating low phenotypic penetrance.

Each generation displayed unpredictable variability in phenotypic expression ranging from mild to severe. Crosses made with mildly affected plants gave rise to severe plants and vice versa. Due to the severity of the phenotype, homozygotes were rarely obtained, thus the observations are exclusively with heterozygotes. Analysis was carried out in plants with at least four backcrosses to B73.

Plant Growth Conditions

Analyses were carried out in families segregating for *Hoja loca1* after at least four backcrosses to the B73 inbred line. Plants were grown in Supersoil Brand Potting Soil (composted bark (60-85%), Sawdust (60-85%), Sphagnum Peat Moss (10-30%)) in standard greenhouse conditions. Seedlings were grown for 3 weeks in 30 by 45 by 2.5cm trays, and adult plants were grown in 12.5L pots until tassel emergence. Seedlings for the RNAseq analysis were the offspring of a cross between a normal sibling and a mutant *Oja* sibling, grown in M3 soil mix with insecticide in large trays equally spaced, in the Grodome facility at the University of Edinburgh with supplementary lighting day/night cycles of 21C, 16hours / 15C, 8 hours.

Root Analysis

Seeds were rehydrated overnight in sterile water, then sterilized with 10% bleach for 20 minutes, before washing with sterile water. Seeds were then germinated on damp chromotography paper in vertically fixed plastic wallets. After one week, the seedlings were photographed, then rotated 90°. The seedlings were left to grow for 24 hours and imaged every hour, for 12 hours. After 24 hours, the final root phenotype was imaged.

578 Primary root length was analysed using FIJI (Schindelin et al., 2012) and crown root
579 number was counted. The seedlings were transplanted to soil and three weeks after
580 germination the shoot phenotype was recorded.
581

582 **Histology Plastic Sections**
583 10µm plastic sections were made with Technovit® 9100 from Electron Microscopy
584 Services (Cat. #14655) according to manufacturer instructions. Sections were mounted
585 on glass slides and stained with toluidine blue. Imaging was performed on a Zeiss
586 Axiovert inverted compound microscope with transmitted light.
587

588 **Scanning Electron Microscopy**
589 SEMs were performed with material fixed in FAA (3.7% formaldehyde, 50% ethanol, 5%
590 acetic acid) overnight at 4°C and dehydrated in ethanol series to 100%. Samples were
591 critical point dried, sputter coated with palladium, and imaged on a Hitachi S-4700 with
592 2kV accelerating voltage.
593

Tissue for Immunolocalisation and *In Situ* Hybridisation
Plant tissue was collected from 3-week-old seedling shoot apices, dissected and fixed
overnight in 4% paraformaldehyde/ 0.1% DMSO/ 0.1% Triton-X100. Fixed samples
were transferred through an ethanol and histoclear series before embedding in
paraplast plus (McCormick Scientific #19216). 10µm sections of fixed tissue were
mounted on probe-on-probe-plus slides (FisherBrand #22-230-900) and dried overnight
on a 37°C hotplate before use in either immunolocalisation or *in situ* hybridization.
601

Immunolocalisation
Primary antibodies were guinea-pig anti-SoPIN1 as used in (Richardson et al., 2016)
and rabbit anti-KNOTTED1 (KN1) as used in (Bolduc et al., 2012b). Secondary
antibodies were commercial anti-guinea pig conjugated to Alexa-fluor594 (Jackson
ImmunoResearch, #706-585-148) and anti-rabbit alkaline phosphatase conjugate
(Jackson ImmunoResearch, #111-055-045). All antibodies were used at a 1:200 dilution
in 1% bovine serum albumin (BSA) in PBS.

Immunolocalisation used a protocol based on (Conti and Bradley, 2007), and
was as follows. Slides were de-paraffinised using histoclear, then rehydrated through an
ethanol series to water. Samples were then boiled in 10mM sodium citrate, pH6, for 10
minutes to retrieve the antigens. Blocking was carried out in 1% BSA/PBS/0.3% Triton-
X100 for 3hours. Slides were incubated overnight in primary antibody, before washing in
PBS/0.3% Triton-X100. They were then incubated with secondary antibody at room-
temperature for 2 hours.

Tissues incubated with anti-KN1 were incubated in a 1:50 dilution of NBT/BCIP
(Roche, #11681451001) in 0.05M $MgCl_2$/TBS, pH9.5 until a dark precipitate was
observed. These slides were then imaged on a Leica dissecting microscope in water.

Tissues incubated with anti-SoPIN1 were stained for 20 minutes in 0.1%
calcofluor (fluorescent brightener 28, Sigma-Aldrich #F3543), washed and mounted in
water and imaged on a Leica SP8 confocal microscope. The subcellular localization of
SoPIN1 was assessed in relation to the calcofluor cell wall stain using FIJI (Schindelin
et al., 2012).

***In situ* Hybridisation**
Antisense probes targeted to *DROOPLING LEAF1* (GRMZM2G168893) mRNA were
designed as used in (Strable et al., 2017). *In situ* hybridization was carried out as in
(Coen et al., 1990) outlined here in brief. Tissue was deparaffinised using histoclear,
then rehydrated through an ethanol series. Samples were digested using 0.125mg/mL
pronase (Sigma-Aldrich #P6611) in 50mMTris/5mM EDTA, pH7.5, for 15minutes at
room temperature. Digestion was stopped by the addition of 0.2% glycine before
washing and re-fixing in 4% PFA/PBS for 10minutes. Slides were then treated in 0.1M
Triethanolamine-HCl / 0.5% Acetic Anhydride (Sigma-Aldrich #320102) for 10 minutes
before dehydrating through an ethanol series. Tissue was incubated overnight with the
probes at a 1:50 dilution in hybridisation buffer (0.375M NaCl, 12.5mM Tris-HCl pH8,
12.5mM Sodium Phosphate pH6.8, 6.25mM EDTA, 50% deionized formamide, 12.5%
dextran sulfate, 1.25x Denhardt solution, 0.0125mg/mL tRNA) at 50°C. Slides were
washed in 0.2% SSC at 50°C, then treated with RNAseA (in 0.5M NaCl/10mM Tris/
1mM EDTA, pH7.5) at 37°C for 15 mins, before repeating the SSC washes. Slides were

blocked in Roche Blocking Reagent (#11-096-176-001) for 30 minutes, washed in1%

BSA/0.3 % Triton-X100/TBS, and incubated at room-temperature with a 1:500 dilution

of anti-Dioxigenin-AP Fab fragments (Roche, #11093274910) in 1% BSA/0.3 % Triton-

X100/TBS for 1hour. Slides were then washed with 1% BSA/0.3 % Triton-X100/TBS,

then in 0.05M MgCl$_2$/TBS, pH9.5. To visualize the probe localisation, the tissue was

incubated in 1:50 dilution of NBT/BCIP in 0.05M MgCl$_2$/TBS, pH9.5, until dark

precipitate formed. The staining reaction was stopped by transferring to water and the

slides were imaged on a Leica MZ16F microscope with a Qimaging Micropublisher

camera under water and brightfield conditions.

Mapping with SSRs

651 The *Oja* mutation was mapped to chromosome 3 using short sequence repeat (SSR)

652 PCR mapping. Known SSRs were identified using Maize gdb and are listed in Figure 3.

653 Custom SSRs were identified by using the Gramene SSR prediction tool (Temnykh et

654 al., 2001), and specific primers were designed to amplify them.

Whole Genome Sequencing

10 mm^2 area leaf discs were collected from 30 wild type and 30 mutant siblings in an

advanced B73 introgressed population. Genomic DNA was extracted using a UREA

based extraction method (Chen and Dellaporta, 1994) which is as follows. Tissue was

ground in UREA extraction buffer (4M urea, 4M NaCl, 1M Tris pH8, 0.5M EDTA, 34mM

n-lauroyl sarcosine). Then an equal volume of phenol:chloroform:isoamyl alcohol

(25:24:1), was added and vortexed to mix. After centrifugation, the supernatant was

decanted and mixed with an equal volume of chloroform. After centrifugation, the

supernatant was decanted and mixed with 1/10th volume of 4.4 M NH$_4$OAc pH 5.2 and

0.7 volume of isopropanol. Strands of DNA were then collected and washed with 70%

ethanol in a separate tube. The dried pellet was resuspended in 100ul of TE.

DNA was quantified using Qubit dsDNA HS Assay Kit (Invitrogen, Q32854), and

sent to Novogene for library preparation and sequencing (350bp insert, 150PE).

Sequencing was at 10 times coverage of the maize genome. Sequence analysis was

carried out using the CYVERSE Atmosphere computing environment. The sequence

data was quality checked and the adapters trimmed using fastp, then aligned to the

B73_RefGen_v3 maize genome using bowtie2, variants were called using samtools

mileup. The variant list was filtered using R, by removing variants recorded in the

Panzea HapMapv3 (https://hapmap.ncbi.nlm.nih.gov/) and those present in the wild

type pool as well as restricting to EMS type SNPs. Moderate and high effect SNPs

specific to the mutant were calculated using SNPEff2 with the B73 RefGenv3 5b+ gene

models.

RNASeq

Seedlings were imaged using a Canon EOS250D camera, then dissected and imaged

using a Leica MZ16F stereoscope with a Canon EOS250D camera attached via an LM-

Scope DSLR adapter. Seedlings were dissected at 21 and 28 days down to the

meristem and the P1/ P2 primordium, and individually flash frozen in liquid nitrogen.

RNA was extracted from each individual using TRIzol (Invitrogen), precipitated (0.5M

NaCl + 2-propyl alcohol), and washed with 70% ethanol, then resuspended in nuclease-

free water. RNA was quantified using a QuBit BR-RNA kit. RNA of sufficient quality was

sent to Novogene for 150bp paired-end sequencing, and 36 libraries passed quality

checks.

The CYVERSE Atmosphere virtual computing environment was used to carry out

sequence quality checks, genome alignments and count extraction. (4CPUs, 32G RAM,

with a ubuntu operating system). Sequenced libraries were checked for quality and

adapters were trimmed using fastp (Chen et al., 2018), then aligned to the B73

RefGenv3 genome using HiSAT2 (Kim et al., 2019). Counts were called using

FeatureCounts (Liao et al., 2014). All downstream analysis was carried out locally

(3.6GHz 8-Core Intel Core i9 processor, 32GB RAM) using R packages combined with

Jupyter notebooks. Differential gene expression analysis was carried out using DESeq2

(Love et al., 2014) using all genes defined as expressed (in more than 3 samples, with

more than 5 counts). Significantly expressed genes were defined as those with a padj

value of <0.05, and filtered to a log2 foldchange value of <-0.59 or >0.59. GO term

analysis used GOSeq (Young et al., 2010) with the maize-GAMER v3 annotations

(Wimalanathan et al., 2018), and significantly enriched GO terms had a Benjamini and
Hochberg cutoff padj value of <0.05.

To carry out hierarchical clustering we stringently filtered the expressed gene list
to those genes with >5 counts in >10 samples, then filtered for those genes listed as
expressed in the meristem and leaf primordia from (Knauer et al., 2019). The top 1500
most variable of these genes were normalised and used to plot the 1 minus the absolute
value of the Pearson correlation matrix, and the resulting plot was clustered
hierarchically and cut using Kmeans =4. The samples in each cluster were then labelled
with cluster identity.

To generate self-organising maps, expressed genes (in more than 3 samples,
with more than 5 counts) were normalised, and the average variance stablised value
was calculated for each cluster. 4 x 4 node SOM grids were then generated using the
kohonen package (Wehrens and Kruisselbrink, 2018).

Vein Analysis
Cross-sections through the third leaf of 3-week-old seedlings, 1.5cm above the ligule
were taken and imaged under water on a Leica DM4000 microscope. Cross-section
images were analysed using FIJI (Schindelin et al., 2012). The width of the leaf was
recorded, then the number of lateral and intermediate veins were counted. Vein density
was calculated as the total number of veins (lateral + intermediate veins) over the width
of the leaf.

Vein distribution was assessed by counting the number of intermediate veins
between lateral veins. To align the wild type and midribless leaves, the lateral vein
closest to the midpoint of the leaf was designated at position 0, the "midvein". Each
lateral vein was then allocated a position number in reference to position 0.

Acknowledgements: With thanks to Samuel Leiboff and China Lunde for extensive help and advice, ARS microscopy staff, UC Berkeley greenhouse staff, and the University of Edinburgh Plant Growth Facilities. **Author Contributions:** SH conceived the original project, developed the mapping population and secured funding. AR and AS designed and carried out the experiments. AR and SH wrote the manuscript. AR agrees to serve as the author responsible for contact and ensures communication. **Funding:** This work was funded by NSF BIO/BBSRC1547062, ARS 5335-21000-013-00D and AR's University of Edinburgh Start-Up funds. This work made use of material supported by the National Science Foundation under Award Numbers DBI-0735191, DBI-1265383, and DBI-1743442 (www.cyverse.org) **Materials and Data Availability:** All microscopy data is available via the University of Edinburgh DataShare (https://datashare.ed.ac.uk/handle/10283/3987) in The Plant Shape Lab collection. All next generation sequencing data is available via MaizeGDB or the University of Edinburgh DataShare in The Plant Shape Lab collection. Jupyter notebooks for the bioinformatic analyses are available via Github (https://github.com/ThePlantShapeLab/Hojaloca1).

All materials are available on request from the corresponding author. All other data are in the main paper or the supplement.

Figures 1-7

Supplementary Table 1: Custom SSR mapping primers

Supplementary Figures 1-3

Supplementary Datasets 1-5

References

Aida, M., Ishida, T., and Tasaka, M. (1999). Shoot apical meristem and cotyledon

formation during Arabidopsis embryogenesis: interaction among the CUP-SHAPED

COTYLEDON and SHOOT MERISTEMLESS genes. Development **126**: 1563–

1570.

Barazesh, S., Nowbakht, C., and McSteen, P. (2009). sparse inflorescence1, barren

inflorescence1 and barren stalk1 promote cell elongation in maize inflorescence

development. Genetics **182**: 403–406.

Bayer, E.M., Smith, R.S., Mandel, T., Nakayama, N., Sauer, M., Prusinkiewicz, P.,

and Kuhlemeier, C. (2009). Integration of transport-based models for phyllotaxis

and midvein formation. Genes Dev. **23**: 373–384.

Baylis, T., Cierlik, I., Sundberg, E., and Mattsson, J. (2013). SHORT

INTERNODES/STYLISH genes, regulators of auxin biosynthesis, are involved in

leaf vein development in Arabidopsis thaliana. New Phytol. **197**: 737–750.

Becraft, P.W. and Freeling, M. (1991). Sectors of liguleless-1 Tissue Interrupt an

Inductive Signal during Maize Leaf Development. Plant Cell.

Von Behrens, I., Komatsu, M., Zhang, Y., Berendzen, K.W., Niu, X., Sakai, H.,

Taramino, G., and Hochholdinger, F. (2011). Rootless with undetectable

meristem 1 encodes a monocot-specific AUX/IAA protein that controls embryonic

seminal and post-embryonic lateral root initiation in maize. Plant J.

Benková, E., Michniewicz, M., Sauer, M., Teichmann, T., Seifertová, D., Jürgens,

G., and Friml, J. (2003). Local, efflux-dependent auxin gradients as a common

module for plant organ formation. Cell **115**: 591–602.

Bhatia, N., Bozorg, B., Larsson, A., Ohno, C., Jönsson, H., and Heisler, M.G.

(2016). Auxin acts through MONOPTEROS to regulate plant cell polarity and

pattern phyllotaxis. Curr. Biol. **26**: 3202–3208.

Bolduc, N., O'Connor, D., Moon, J., Lewis, M., and Hake, S. (2012a). How to pattern

a leaf. In Cold Spring Harbor symposia on quantitative biology (Cold Spring Harbor

Laboratory Press), pp. 47–51.

Bolduc, N., Yilmaz, A., Mejia-Guerra, M.K., Morohashi, K., O'Connor, D.,

Grotewold, E., and Hake, S. (2012b). Unraveling the KNOTTED1 regulatory

network in maize meristems. Genes Dev. **26**: 1685–1690.

Chen, J. and Dellaporta, S. (1994). Urea-based plant DNA miniprep. In The maize

handbook (Springer), pp. 526–527.

Chen, S., Zhou, Y., Chen, Y., and Gu, J. (2018). Fastp: An ultra-fast all-in-one FASTQ

preprocessor. Fastp An ultra-fast all-in-one FASTQ preprocessor.

Coen, E.S., Romero, J., Doyle, S., Elliott, R., Murphy, G., and Carpenter, R. (1990).

floricaula: a homeotic gene required for flower development in Antirrhinum majus.

Cell **63**: 1311–1322.

Conklin, P.A., Johnston, R., Conlon, B.R., Shimizu, R., and Scanlon, M.J. (2021).

Plant homeodomain proteins provide a mechanism for how leaves grow wide. Dev.

Conti, L. and Bradley, D. (2007). TERMINAL FLOWER1 is a mobile signal controlling

Arabidopsis architecture. Plant Cell **19**: 767–778.

Dharmasiri, N., Dharmasiri, S., and Estelle, M. (2005). The F-box protein TIR1 is an

auxin receptor. Nature **435**: 441–445.

Fukaki, H., Tameda, S., Masuda, H., and Tasaka, M. (2002). Lateral root formation is

blocked by a gain-of-function mutation in the SOLITARY-ROOT/IAA14 gene of

Arabidopsis. Plant J. **29**: 153–168.

Galinat, W.C. (1959). The phytomer in relation to floral homologies in the American

Maydeae. Bot. Mus. Lealf. Harv. Univ. **19**: 1–xvi.

Galli, M., Khakhar, A., Lu, Z., Chen, Z., Sen, S., Joshi, T., Nemhauser, J.L.,

Schmitz, R.J., and Gallavotti, A. (2018). The DNA binding landscape of the maize

AUXIN RESPONSE FACTOR family. Nat. Commun. **9**: 1–14.

Galli, M., Liu, Q., Moss, B.L., Malcomber, S., Li, W., Gaines, C., Federici, S.,

Roshkovan, J., Meeley, R., and Nemhauser, J.L. (2015). Auxin signaling

modules regulate maize inflorescence architecture. Proc. Natl. Acad. Sci. **112**:

13372–13377.

Hamann, T., Benkova, E., Bäurle, I., Kientz, M., and Jürgens, G. (2002). The

Arabidopsis BODENLOS gene encodes an auxin response protein inhibiting

MONOPTEROS-mediated embryo patterning. Genes Dev. **16**: 1610–1615.

Harper, L. and Freeling, M. (1996). Interactions of liguleless1 and liguleless2 function

during ligule induction in maize. Genetics **144**: 1871–1882.

Havens, K.A., Guseman, J.M., Jang, S.S., Pierre-Jerome, E., Bolten, N., Klavins, E.,
and Nemhauser, J.L. (2012). A synthetic approach reveals extensive tunability of
auxin signaling. Plant Physiol. **160**: 135–142.

Hibara, K., Karim, M.R., Takada, S., Taoka, K., Furutani, M., Aida, M., and Tasaka,
M. (2006). Arabidopsis CUP-SHAPED COTYLEDON3 regulates postembryonic
shoot meristem and organ boundary formation. Plant Cell **18**: 2946–2957.

Jackson, D., Veit, B., and Hake, S. (1994). Expression of maize KNOTTED1 related
homeobox genes in the shoot apical meristem predicts patterns of morphogenesis
in the vegetative shoot. Development **120**: 405–413.

Johnston, R., Leiboff, S., and Scanlon, M.J. (2015). Ontogeny of the sheathing leaf
base in maize (Zea mays). New Phytol. **205**: 306–315.

Johnston, R., Wang, M., Sun, Q., Sylvester, A.W., Hake, S., and Scanlon, M.J.
(2014). Transcriptomic analyses indicate that maize ligule development
recapitulates gene expression patterns that occur during lateral organ initiation.
Plant Cell **26**: 4718–4732.

Kepinski, S. and Leyser, O. (2005). The Arabidopsis F-box protein TIR1 is an auxin
receptor. Nature **435**: 446–451.

Kim, D., Paggi, J.M., Park, C., Bennett, C., and Salzberg, S.L. (2019). Graph-based
genome alignment and genotyping with HISAT2 and HISAT-genotype. Nat.
Biotechnol.

Knauer, S. et al. (2019). A high-resolution gene expression atlas links dedicated
meristem genes to key architectural traits. Genome Res.

Kuhlemeier, C. and Timmermans, M.C.P. (2016). The Sussex signal: insights into leaf
dorsiventrality. Development **143**: 3230–3237.

Lavy, M. and Estelle, M. (2016). Mechanisms of auxin signaling. Development **143**:
3226–3229.

Leyser, H.M.O., Pickett, F.B., Dharmasiri, S., and Estelle, M. (1996). Mutations in the
AXR3 gene of Arabidopsis result in altered auxin response including ectopic
expression from the SAUR-AC1 promoter. Plant J. **10**: 403–413.

Liao, Y., Smyth, G.K., and Shi, W. (2014). FeatureCounts: An efficient general
purpose program for assigning sequence reads to genomic features.

Bioinformatics.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change

and dispersion for RNA-seq data with DESeq2. Genome Biol.

Matthes, M.S., Best, N.B., Robil, J.M., Malcomber, S., Gallavotti, A., and McSteen,

P. (2019). Auxin EvoDevo: conservation and diversification of genes regulating

auxin biosynthesis, transport, and signaling. Mol. Plant **12**: 298–320.

McSteen, P. and Hake, S. (2001). Barren inflorescence2 regulates axillary meristem

development in the maize inflorescence. Development.

Moon, J., Candela, H., and Hake, S. (2013). The Liguleless narrow mutation affects

proximal-distal signaling and leaf growth. Development **140**: 405–412.

Nagpal, P., Walker, L.M., Young, J.C., Sonawala, A., Timpte, C., Estelle, M., and

Reed, J.W. (2000). AXR2 encodes a member of the Aux/IAA protein family. Plant

Physiol. **123**: 563–574.

Nardmann, J., Ji, J., Werr, W., and Scanlon, M.J. (2004). The maize duplicate genes

narrow sheath1 and narrow sheath2 encode a conserved homeobox gene function

in a lateral domain of shoot apical meristems. Development **131**: 2827–2839.

Nogueira, F.T.S., Madi, S., Chitwood, D.H., Juarez, M.T., and Timmermans, M.C.P.

(2007). Two small regulatory RNAs establish opposing fates of a developmental

axis. Genes Dev. **21**: 750–755.

O'Connor, D.L., Elton, S., Ticchiarelli, F., Hsia, M.M., Vogel, J.P., and Leyser, T.

(2017). Cross-species functional diversity within the PIN auxin efflux protein family.

Elife.

O'Connor, D.L., Runions, A., Sluis, A., Bragg, J., Vogel, J.P., Prusinkiewicz, P.,

and Hake, S. (2014). A division in PIN-mediated auxin patterning during organ

initiation in grasses. PLoS Comput Biol **10**: e1003447.

Ouellet, F., Overvoorde, P.J., and Theologis, A. (2001). IAA17/AXR3: biochemical

insight into an auxin mutant phenotype. Plant Cell **13**: 829–841.

Overvoorde, P.J., Okushima, Y., Alonso, J.M., Chan, A., Chang, C., Ecker, J.R.,

Hughes, B., Liu, A., Onodera, C., and Quach, H. (2005). Functional genomic

analysis of the AUXIN/INDOLE-3-ACETIC ACID gene family members in

Arabidopsis thaliana. Plant Cell **17**: 3282–3300.

Parry, G., Calderon-Villalobos, L.I., Prigge, M., Peret, B., Dharmasiri, S., Itoh, H.,
Lechner, E., Gray, W.M., Bennett, M., and Estelle, M. (2009). Complex regulation
of the TIR1/AFB family of auxin receptors. Proc. Natl. Acad. Sci. **106**: 22540 LP –
22545.

Pérez-Pérez, J.M., Candela, H., Robles, P., López-Torrejón, G., Del Pozo, J.C., and
Micol, J.L. (2010). A role for AUXIN RESISTANT3 in the coordination of leaf
growth. Plant cell Physiol. **51**: 1661–1673.

Powers, S.K. and Strader, L.C. (2020). Regulation of auxin transcriptional responses.
Dev. Dyn. **249**: 483–495.

Ramos, J.A., Zenser, N., Leyser, O., and Callis, J. (2001). Rapid degradation of
auxin/indoleacetic acid proteins requires conserved amino acids of domain II and is
proteasome dependent. Plant Cell **13**: 2349–2360.

Reed, J.W. (2001). Roles and activities of Aux/IAA proteins in Arabidopsis. Trends
Plant Sci. **6**: 420–425.

Reed, J.W., Elumalai, R.P., and Chory, J. (1998). Suppressors of an Arabidopsis
thaliana phyB mutation identify genes that control light signaling and hypocotyl
elongation. Genetics **148**: 1295–1310.

Reinhardt, D., Pesce, E.-R., Stieger, P., Mandel, T., Baltensperger, K., Bennett, M.,
Traas, J., Friml, J., and Kuhlemeier, C. (2003). Regulation of phyllotaxis by polar
auxin transport. Nature **426**: 255–260.

Richardson, A.E. and Hake, S. (2019). Drawing a line: Grasses and boundaries.
Plants **8**.

Richardson, A.E., Rebocho, A.B.A.B., and Coen, E.S. (2016). Ectopic KNOX
expression affects plant development by altering tissue cell polarity and identity.
Plant Cell **28**: tpc-00284.

Rogg, L.E., Lasswell, J., and Bartel, B. (2001). A gain-of-function mutation in IAA28
suppresses lateral root development. Plant Cell **13**: 465–480.

Rouse, D., Mackay, P., Stirnberg, P., Estelle, M., and Leyser, O. (1998). Changes in
auxin response from mutations in an AUX/IAA gene. Science (80-.). **279**: 1371–
1373.

Salehin, M., Bagchi, R., and Estelle, M. (2015). SCFTIR1/AFB-based auxin

perception: mechanism and role in plant growth and development. Plant Cell **27**: 9–
19.

Scanlon, M.J. (2003). The polar auxin transport inhibitor N-1-naphthylphthalamic acid
disrupts leaf initiation, KNOX protein regulation, and formation of leaf margins in
maize. Plant Physiol. **133**: 597–605.

Scanlon, M.J. and Freeling, M. (1997). Clonal sectors reveal that a specific
meristematic domain is not utilized in the maize mutantnarrow sheath. Dev. Biol.
182: 52–66.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., and Schmid, B. (2012). Fiji: an open-
source platform for biological-image analysis. Nat. Methods **9**: 676–682.

Sharman, B.C. (1942). Developmental anatomy of the shoot of Zea mays L. Ann. Bot.
6: 245–282.

Singh, S., Yadav, S., Singh, A., Mahima, M., Singh, A., Gautam, V., and Sarkar,
A.K. (2020). Auxin signaling modulates LATERAL ROOT PRIMORDIUM1 (LRP1)
expression during lateral root development in Arabidopsis. Plant J. **101**: 87–100.

Smith, L.G., Greene, B., Veit, B., and Hake, S. (1992). A dominant mutation in the
maize homeobox gene, Knotted-1, causes its ectopic expression in leaf cells with
altered fates. Development **116**: 21–30.

Strable, J., Wallace, J.G., Unger-Wallace, E., Briggs, S., Bradbury, P.J., Buckler,
E.S., and Vollbrecht, E. (2017). Maize YABBY genes drooping leaf1 and drooping
leaf2 regulate plant architecture. Plant Cell **29**: 1622–1641.

Tan, X., Calderon-Villalobos, L.I.A., Sharon, M., Zheng, C., Robinson, C. V, Estelle,
M., and Zheng, N. (2007). Mechanism of auxin perception by the TIR1 ubiquitin
ligase. Nature **446**: 640–645.

Temnykh, S., DeClerck, G., Lukashova, A., Lipovich, L., Cartinhour, S., and
McCouch, S. (2001). Computational and experimental analysis of microsatellites in
rice (Oryza sativa L.): frequency, length variation, transposon associations, and
genetic marker potential. Genome Res. **11**: 1441–1452.

Tian, Q., Nagpal, P., and Reed, J.W. (2003). Regulation of Arabidopsis SHY2/IAA3
protein turnover. Plant J. **36**: 643–651.

Tian, Q. and Reed, J.W. (1999). Control of auxin-regulated root development by the
Arabidopsis thaliana SHY2/AA3 gene. Development.

Tsuda, K., Abraham-Juarez, M.-J., Maeno, A., Dong, Z., Aromdee, D., Meeley, R.,
Shiroishi, T., Nonomura, K., and Hake, S. (2017). KNOTTED1 cofactors, BLH12
and BLH14, regulate internode patterning and vein anastomosis in maize. Plant
Cell **29**: 1105–1118.

Wang, H., Jones, B., Li, Z., Frasse, P., Delalande, C., Regad, F., Chaabouni, S.,
Latché, A., Pech, J.C., and Bouzayen, M. (2005). The tomato Aux/IAA
transcription factor IAA9 is involved in fruit development and leaf morphogenesis.
Plant Cell.

Wehrens, R. and Kruisselbrink, J. (2018). Flexible self-organizing maps in kohonen
3.0. J. Stat. Softw.

Wimalanathan, K., Friedberg, I., Andorf, C.M., and Lawrence-Dill, C.J. (2018).
Maize GO Annotation—Methods, Evaluation, and Review (maize-GAMER). Plant
Direct.

Woll, K., Borsuk, L.A., Stransky, H., Nettleton, D., Schnable, P.S., and
Hochholdinger, F. (2005). Isolation, characterization, and pericycle-specific
transcriptome analyses of the novel maize lateral and seminal root initiation mutant
rum1. Plant Physiol.

Worley, C.K., Zenser, N., Ramos, J., Rouse, D., Leyser, O., Theologis, A., and
Callis, J. (2000). Degradation of Aux/IAA proteins is essential for normal auxin
signalling. Plant J.

Yang, X., Lee, S., So, J.H., Dharmasiri, S., Dharmasiri, N., Ge, L., Jensen, C.,
Hangarter, R., Hobbie, L., and Estelle, M. (2004). The IAA1 protein is encoded by
AXR5 and is a substrate of SCF TIR1. Plant J.

Young, M.D., Wakefield, M.J., Smyth, G.K., and Oshlack, A. (2010). Gene ontology
analysis for RNA-seq: accounting for selection bias. Genome Biol.

Zenser, N., Ellsmore, A., Leasure, C., and Callis, J. (2001). Auxin modulates the
degradation rate of Aux/IAA proteins. Proc. Natl. Acad. Sci. U. S. A.

Zhang, Y., Marcon, C., Tai, H., Von Behrens, I., Ludwig, Y., Hey, S., Berendzen,
K.W., and Hochholdinger, F. (2016). Conserved and unique features of the

homeologous maize Aux/IAA proteins ROOTLESS with UNDETECTABLE

MERISTEM 1 and RUM1-like 1. J. Exp. Bot.

Zhang, Z., Runions, A., Mentink, R.A., Kierzkowski, D., Karady, M., Hashemi, B.,

Huijser, P., Strauss, S., Gan, X., and Ljung, K. (2020). A WOX/Auxin

Biosynthesis Module Controls Growth to Shape Leaf Form. Curr. Biol.

Figure 1. *Hoja loca1* mutants have leaf defects.
The phenotype of wild type (Wt) (A), mild (B) and severe (C) *Oja* mutants including (i)
whole-seedling, and (ii-iv) magnified images of the ligule/ auricle regions, showing the
side (ii) and abaxial (iii) and the adaxial (iv) views. (C iv) is a cross-section through the
tube leaf showing the intact ligule (arrowhead in Cii). (v) Cross-sections through

988 seedling leaves 1cm above the ligule. Scale bars are 1cm, arrowheads indicate the
989 ligule or midrib regions. (D) The distribution of midribless and tube leaves in *Oja* plants
990 (n=150), node 1 is at the base of the plant.
991

Figure 2. *Hoja loca1* mutants have a defect in vein distribution.
(A-B) Plastic sections through wild type (A) and midribless (B) leaves. The midrib is
indicated by the M, the lateral veins by the black arrowheads, and intermediate veins by
the cyan arrowheads. Scale bars are 1mm. (C) Seedling leaf width varies significantly
between wild type and midribless (n=10 of each phenotype, ANOVA and Tukeys HSD,

df=2, * p= 0.0032) and between wild type and tube leaves (**, p=0.000083). (D) Total
leaf vein density does not vary significantly. (E) Leaf vein distribution is perturbed in
midribless mutants. (Ei) Cartoons depicting the alignment of midribless and wild-type
leaves. Wild type and mutant leaves were aligned by allocating the position of "0" to the
wild type midvein, and the lateral vein closest to the midpoint of the leaf in the
midribless mutant. Each successive lateral vein was then allocated a position relative to
0 (Ei) and the number of intermediate veins between each position was recorded. (Eii)
The mean number of intermediate veins near the midvein were significantly different in
the midribless mutant (orange) when compared to wild type (blue) (n=10 for each
phenotype, ANOVA and Tukeys HSD, * p = 0.0051 and p=0.00089).
1008

1009

Figure 3. *Hoja loca1* mutants initiate new phytomers but fail to initiate leaves.
(A-C) Mature *Oja* plants are often short, and despite having a similar number of visible nodes (B, p=0.2537, Welch's 2-Sample T-Test, df= 39.8, n *Oja*=29, Wt=14), they have fewer leaves (C, ** p= 4.17e-11, Welch's 2-Sample T-Test, df=14.194, n= 10 of each phenotype) than wild type. In particular, the nodes below the tassel are often bare (Aii, arrowheads). Bare nodes can also be identified in cross-sections through developing shoots (D). The lack of leaf initiation is clear at the earliest stages of development. Scanning electron micrographs (SEMs) of wild type (E) and mutant (F-I) vegetative meristems with attached leaf primordia show severe defects in *Oja*. Some leaf primordia fail to arch over the meristem, instead forming a ring-shaped primordium (F). Some have clear ridges where internodes have formed below the meristem (G), which would normally be concealed by leaves in wild-type plants. Other mutants form pin-shaped meristems (H). Often the tiller buds in *Oja* are normal (I) allowing the plant to produce progeny. The internodes are indicated by the *, meristems by 'M', axillary meristems by 'AM'. Scale bars are 100µm.

Figure 4. *Hoja loca1* maps to ZmlAA28 on Chr8.

Using short sequence repeat (SSR) PCR markers, *Hoja* was mapped to Chr8 between 109.7-110.6Mb (A). Custom SSR markers are labeled as 'AS#'. Of the 19 genes in the fine mapping interval (Aiii), 10 are expressed in shoot meristems and leaf primordia (green), 9 are not (magenta) based on published RNAseq datasets. Whole genome sequencing (WGS) data identified 9 high to moderate effect SNPs in gene bodies in the larger mapping region (Ai, blue ovals, Chr8:75-135Mb). Of these, only one is an EMS-type SNP that falls within the fine mapping interval (Aiii, *). (B)The SNP causes a C to T change in the second exon of GRMZM2G035465, *ZmlAA28* which results in a glycine (G) to glutamic acid (E) amino acid change in the highly conserved core degron motif (C, cyan box).

1038

Figure 5. *Hoja loca1* mutants show changes in auxin, leaf development, and meristem regulation genes.

(A) Examples of the dissected shoot apical meristems harvested for the RNAseq experiment. Scalebar:100μm. (B) Principal component analysis plot (PCA) of all RNAseq libraries, coloured by seeding phenotype at time of harvest; mutant (orange) or wild-type (blue) (n=100). (C) Significantly enriched GO terms (–log(padj) >1) associated with auxin, meristem regulation and leaf development for all genes that are significantly differentially expressed between mutant and wild type scored samples (padj<0.05, log2foldchange >0.59 or <-0.59). Z-score indicates whether the genes in the GO term are on average upregulated (green) or downregulated (purple).

1049

Figure 6. Transcriptional profiles of individual vegetative meristems enable identification of four transcriptional profiles associated with mutant leaf phenotypes.

(A) Hierarchical clustering of 1 minus the Pearsons correlation of sample distances (dark blue most similar) based on the top 1500 most variable leaf and meristem associated expressed genes. K=4. (B) Principal component analysis plot (PCA) coloured by the 4 clusters identified in the hierarchical clustering (n=100), cluster 1 (red), cluster 2 (green), cluster 3 (blue), cluster 4 (purple). (C) Venn diagram showing the overlap in significantly differentially expressed (sigDE) genes in comparisons with cluster 2 (C2) which have a log2foldchange of >0.59 or <-0.59. C#vsC# indicates cluster comparison. (#) indicate the total number of sigDE genes, numbers indicate the number

of genes in each category. (D) 4x4 self-organizing maps built from all expressed genes
(defined as more than 5 rpm in more than 3 samples) in each cluster. Nodes 1-16 are
shown. N indicates number of genes in the node. (E-H) Heatmaps of average FPKM
values of expressed *KNOX* (E), WOX (F), *YABBY* (G) *and SRS* (H) genes in clusters 1-
4. Colouring is relative average FPKM per row, green is positive, purple is negative.
Genes that are significantly differentially expressed in one or more pairwise
comparisons with cluster 2 are underlined (padj<0.05), "*" indicates genes with a
log2foldchange >0.59 or <-0.59, "**" indicate genes with log2foldchange >1.0 or <-1.0.
1069

Figure 7. *Hoja loca1* have defects in organ initiation markers localization.
KNOTTED1 (KN1) immunolocalisation (dark brown/black, A-B) and SISTER OF
PINFORMED1 (SoPIN1, green, C-D) in longitudinal sections through 3-week-old
seedlings in WT (A,C) and mutant (B,D) plants. The inset in D shows a magnified image
of the boxed region, the arrowheads indicate the orientation of the SoPIN1 localization
(green) within the cell outline (magenta). Large arrowheads indicate the P_0 and P_1
domains. Dotted lines indicate the outline of the meristem region. The white dotted line
in A indicate the P_0. Grey dashed lines indicate vascular traces. (E-F) *In situ*
hybridization of *drooping leaf* (dark brown, DL) in transverse cross-sections through 3-
week-old seedlings, in wild type (E) and mutant (F) plants. Sections are 100µm from the
tip of the meristem. * indicate the position of the midvein in successive leaves, the white
dotted line outlines the meristem. Scale bars are 100µm.

1084

Primer Name	Sequence (Forward/Reverse)
AS_1	TGGGCCATGCTAGGCATTAACT/ TCTGGCTCCTCTCTACCGGA
AS_2	CGTCCTCCAGCTTAGTGTTG/ CTTCGACGACTTCTCCTTCC
AS_3	GGGCTGGTTTGGTGGCAGGG/ TGTCATGCGTCCGTCACTGCT
AS_4	GGTTTCCCGGAGGAGGAATG/ AGTTAGTCGCCTCTTTCGCC
AS_5	ATGCCAGGAAATCACGGAGG/ GAGTGTTTCCGAGGCTTTGC

1085 **Supplementary Table 1: Custom SSR mapping primers**
1086 Primers designed using the Gramene SSR prediction tool (Temnykh et al., 2001) to
1087 map the *Oja* mutation. Primer locations are indicated in Figure. 4.
1088
1089

Supplementary Figures

Supplementary Figure 1. *Hoja loca1* **Mature Phenotype.**
The mature phenotype of a wild type (A) and mutant (B) plant, with leaves attached (i) and removed (ii), scale bar is 15cm. The positions of nodes are indicated by the white arrowheads, those that have a leaf are indicated by "L" and the base of the tassel is indicated by "T". The phenotype of tassels and ears in wild type (C) and mutant (D) plants. (i) Whole tassel, (ii) zoomed-in image of the boxed region, (iii) an individual spikelet pair, (iv) unfertilized ear. Scale bars are 1cm.

1099

Supplementary Figure 2. *Hoja loca1* mutants have a normal root phenotype.
(A) Oja seedling roots have a normal gravitropic response, showing a change in root tip
orientation by 1 hour after turning (*) (scale bars are 2cm). (B-C) Root architecture in
1.5-week-old seedlings is not significantly different between wild-type and *Oja* plants, as
shown by primary root length (B, Welch 2-sample t-test, df=24, n *Oja*=13, Wt=24,
p=0.853) and number of initiating crown roots (C, Welch 2-sample t-test, df=19, n
Oja=13, Wt=24, p=0.1452). (D-E) Root architecture in 3-week-old seedlings appears to
be similar between mutant and normal sibling (Wt) (scale bars are 2cm).

1108

1109

Supplementary Figure 3. *Hoja loca1* mutants show changes in auxin-signaling related genes.

Heatmap of average FPKM values of Auxin related genes in clusters 1-4. Colouring is relative average FPKM per row. (* & **) indicate genes that are significantly differentially expressed (padj<0.05), (**) log2foldchange >1 or <-1, foldchange of 2 or 0.5

1115 respectively, (*) log2foldchange <0.59<1 or >-1<0.59, foldchange of 1.5 or 0.5
1116 respectively between the indicated cluster and cluster 2.
1117

Supplementary Datasets:

Supplementary Dataset 1: Broad RNAseq differential gene expression comparison between individuals scored as normal or mutant based on seedling phenotype.

DeSeq2 output of comparisons between samples with seedling phenotypes scored as wild-type or mutant. This data relates to the results shown in Figure 5. All significantly expressed genes between the pools of mutant and wildtype scored samples (padj <0.05) are listed by their gene identifiers (GID) based on the maize B73 RefGenv3 genome, with matched annotations from Maize, and the best BLAST hit for Arabidopsis and rice. Negative log2FoldChange values indicate down-regulation in mutants.

Supplementary Dataset 2: RNAseq differential gene expression comparison between individuals in cluster 1 versus cluster 2.

DeSeq2 output of comparisons between samples defined as cluster 1 or 2 based on the hierarchical clustering. This data relates to the results shown in Figure 6. All significantly expressed genes between the cluster 1 and cluster 2 scored samples (padj <0.05) are listed by their gene identifiers (GID) based on the maize B73 RefGenv3 genome, with matched annotations from Maize, and the best BLAST hit for Arabidopsis and rice.

Supplementary Dataset 3: RNAseq differential gene expression comparison between individuals in cluster 3 versus cluster 2.

DeSeq2 output of comparisons between samples defined as cluster 3 or 2 based on the hierarchical clustering. This data relates to the results shown in Figure 6. All significantly expressed genes between the cluster 3 and cluster 2 scored samples (padj <0.05) are listed by their gene identifiers (GID) based on the maize B73 RefGenv3 genome, with matched annotations from Maize, and the best BLAST hit for Arabidopsis and rice

Supplementary Dataset 4: RNAseq differential gene expression comparison between individuals in cluster 4 versus cluster 2.

1148   DeSeq2 output of comparisons between samples defined as cluster 4 or 2 based on the

1149   hierarchical clustering. This data relates to the results shown in Figure 6. All significantly

1150   expressed genes between the cluster 4 and cluster 2 scored samples (padj <0.05) are

1151   listed by their gene identifiers (GID) based on the maize B73 RefGenv3 genome, with

1152   matched annotations from Maize, and the best BLAST hit for Arabidopsis and rice

1153

Supplementary Dataset 5: Self-organising map information for all expressed

genes

List of all expressed genes (defined as more than 5 counts in more than 3 samples) with

their associated gene identifiers (GID) based on the maize B73 RefGenv3 genome, with

matched annotations from Maize, and the best BLAST hit for Arabidopsis and rice, and

their node defined in the 4x4 SOM analysis shown in Figure 6D.

1160