Evaluating the transcriptional fidelity of cancer models

Da Peng[1*], Rachel Gleyzer[2*], Wen-Hsin Tai[2], Pavithra Kumar[2], Qin Bian[2], Bradley Issacs[2],
Edroaldo Lummertz da Rocha[3], Stephanie Cai[1], Kathleen DiNapoli[4,5], Franklin W Huang[6],
Patrick Cahan[1,2,7]

[1]Department of Biomedical Engineering, Johns Hopkins University School of Medicine,
Baltimore MD 21205 USA

[2]Institute for Cell Engineering, Johns Hopkins University School of Medicine,
Baltimore MD 21205 USA

[3]Department of Microbiology, Immunology and Parasitology,
Federal University of Santa Catarina, Florianópolis SC, Brazil

[4]Department of Cell Biology, Johns Hopkins University School of Medicine,
Baltimore, MD 21205 USA

[5]Department of Electrical and Computer Engineering, Johns Hopkins University,
Baltimore MD 21218 USA

[6]Division of Hematology/Oncology, Department of Medicine; Helen Diller Family Cancer Center;
Bakar Computational Health Sciences Institute; Institute for Human Genetics;
University of California, San Francisco, San Francisco, CA

[7]Department of Molecular Biology and Genetics, Johns Hopkins University School of Medicine,
Baltimore MD 21205 USA

* These authors made equal contributions.

Correspondence to: patrick.cahan@jhmi.edu

Article type: Analysis

Website: http://www.cahanlab.org/resources/cancerCellNet_web

Code: https://github.com/pcahan1/cancerCellNet

ABSTRACT

Cancer researchers use cell lines, patient derived xenografts, and genetically engineered mice as models to investigate tumor biology and to identify therapies. The generalizability and power of a model derives from the fidelity with which it represents the tumor type of investigation, however, the extent to which this is true is often unclear. The preponderance of models and the ability to readily generate new ones has created a demand for tools that can measure the extent and ways in which cancer models resemble or diverge from native tumors. Here, we present a computational tool, CancerCellNet, that measures the similarity of cancer models to 22 naturally occurring tumor types and 36 subtypes, in a platform and species agnostic manner. We applied this tool to 657 cancer cell lines, 415 patient derived xenografts, and 26 distinct genetically engineered mouse models, documenting the most faithful models, identifying cancers underserved by adequate models, and finding models with annotations that do not match their classification. By comparing models across modalities, we find that genetically engineered mice have higher transcriptional fidelity than patient derived xenografts and cell lines in four out of five tumor types. We have made CancerCellNet available as freely downloadable software and as a web application that can be applied to new cancer models.

INTRODUCTION

Models are widely used to investigate cancer biology and to identify potential therapeutics. Popular modeling modalities are cancer cell lines (CCLs)[1], genetically engineered mouse models (GEMMs)[2], and patient derived xenografts (PDXs)[3].These classes of models differ in the types of questions that they are designed to address. CCLs are often used to address cell intrinsic mechanistic questions[4], GEMMs to chart progression of molecularly defined-disease[5], and PDXs to explore patient-specific response to therapy in a physiologically relevant context[6]. Models also differ in the extent to which the they represent specific aspects of a cancer type[7]. Even with this intra- and inter-class model variation, all models should represent the tumor type or subtype under investigation, and not another type of tumor, and not a non-cancerous tissue. Therefore, cancer-models should be selected not only based on the specific biological question but also based on the similarity of the model to the cancer type under investigation[8,9].

Various methods have been proposed to determine the similarity of cancer models to their intended subjects. Domcke et al devised a 'suitability score' as a metric of the molecular similarity of CCLs to high grade serous ovarian carcinoma based on a heuristic weighting of copy number alterations, mutation status of several genes that distinguish ovarian cancer subtypes, and hypermutation status[10]. Other studies have taken analogous approaches by either focusing on transcriptomic or ensemble molecular profiles (e.g. transcriptomic and copy number alterations) to quantify the similarity of cell lines to tumors[11–13]. These studies were tumor-type specific, focusing on CCLs that model, for example, hepatocellular carcinoma or breast cancer. More recently, Yu et al compared the transcriptomes of CCLs to the Cancer Genome Atlas (TCGA) by correlation analysis, resulting in a panel of CCLs recommended as most representative of 22 tumor types[14]. While all of these studies have provided valuable information, they leave two major challenges unmet. The first challenge is to determine the fidelity of GEMMs and PDXs and whether there are stark differences between these classes of models and CCLs. The other major unmet challenge is to enable the rapid assessment of new,

emerging cancer models. This challenge is especially relevant now as technical barriers to

generating models have been substantially lowered[15,16], and because each PDX can be

considered a distinct entity requiring individual validation[17].

To address these challenges, we developed CancerCellNet (CCN), a computational tool

that uses transcriptomic data to quantitatively assess the similarity between cancer models and

22 naturally occurring tumor types and 36 subtypes in a platform- and species-agnostic manner.

Here, we describe CCN's performance, and the results of applying it to assess 657 cancer cell

lines, 415 patient derived xenografts, and 26 distinct genetically engineered mouse models.

This has allowed us to identify the most faithful models currently available, to document cancers

underserved by adequate models, and to find models with inaccurate tumor type annotation.

Moreover, because CCN is open-source and easy to use, it can be readily applied to newly

generated cancer models as a means to assess their fidelity.

RESULTS

CancerCellNet classifies samples accurately across species and technologies

Previously, we had developed a computational tool using the Random Forest

classification method to measure the similarity of engineered cell populations to their *in vivo*

counterparts based on transcriptional profiles[18,19]. More recently, we elaborated on this

approach to allow for classification of single cell RNA-seq data in a manner that allows for

cross-platform and cross-species analysis[20]. Here, we used an analogous approach to

quantitatively compare cancer models to naturally occurring patient tumors (**Fig 1A**). In brief, we

used TCGA RNA-seq expression data from 22 solid tumor types to train a top-pair multi-class

Random forest classifier. We combined training data from Rectal Adenocarcinoma (READ) and

Colon Adenocarcinoma (COAD) into one COAD_READ category because READ and COAD

are considered to be virtually indistinguishable at a molecular level[21]. We included an 'Unknown'

category trained using randomly shuffled gene-pair profiles generated from the training data of

22 tumor types to identify query samples that are not reflective of any of the training data.

We assessed the performance of this approach by computing the area under the

precision recall (AUPR) curves derived by 50 iterations of cross validation (**Fig 1B, Supp Fig**

1A). In the cross validations, the mean AUPR exceeded 0.95 in most of the tumor types. In

addition to achieving high mean AUPRs on held-out TCGA data, we found that CCN also

achieved high AUPR (above 0.9) when we applied it to independent testing data from the

International Cancer Genome Consortium (ICGC) consisting of RNA-Seq data from 886 tumors

across 5 tumor types (**Supp Fig 1B**)[22]**.**

As one of the central aims of our study is to compare distinct cancer models, including

GEMMs, our method needed to be able to classify samples from mouse and human samples

equivalently. We used the Top-Pair transform[20] to achieve this and we tested the feasibility of

this approach by assessing the performance of a normal (i.e non-tumor) cell and tissue classifier

trained on human data as applied to mouse samples. Consistent with prior applications[23], we

found that the cross-species classifier performed well, achieving mean AUPR of 0.96 when

applied to mouse data (**Supp Fig 1C**)**.**

To evaluate cancer models at a finer resolution, we also developed an approach to

perform tumor subtype classifications (**Supp Fig 1D**). We constructed 11 different cancer

subtype classifiers based on the availability of expression or histological subtype

information[21,24–34]. We also included non-cancerous, normal tissues as categories for several

subtype classifiers when sufficient data was available: breast invasive carcinoma (BRCA),

COAD_READ, head and neck squamous cell carcinoma (HNSC), kidney renal clear cell

carcinoma (KIRC) and uterine corpus endometrial carcinoma (UCEC). The 11 subtype

classifiers all achieved high overall average AUPRs ranging from 0.75 to 0.99 (**Supp Fig 1E**)**.**

Fidelity of cancer cell lines

Having validated the performance of CCN, we then used it to determine the fidelity of

CCLs. We mined RNA-Seq expression data of 657 different cell lines across 20 cancer types

from the Cancer Cell Line Encyclopedia (CCLE) and applied CCN to them, finding a wide

classification range for cell lines of each tumor type (**Fig 2A, Supp Tab 1**). To verify the

classification results, we applied CCN to CCLE expression profiles generated through

microarray expression profiling[35]. To ensure that CCN would function on microarray data, we

first tested it by applying a CCN classifier created to test microarray data to 720 expression

profiles of 12 tumor types. The cross-platform CCN classifier performed well, based on the

comparison to study-provided annotation, achieving a mean AUPR of 0.944 (**Supp Fig 2A**).

Next, we applied this cross-platform classifier to microarray expression profiles of CCLE (**Supp**

Fig 2B). From the classification results of 571 cell lines that have both RNA-seq and microarray

expression profiles, we found a strong positive association between the classification scores

from RNA-seq and those from microarray (**Supp Fig 2C**). This comparison supports the notion

that the classification scores for each cell line are not artifacts of profiling methodology.

Moreover, this comparison shows that the scores are consistent between the times that the cell

lines were first assayed by microarray expression profiling in 2012 and by RNA-Seq in 2019.

We also observed high level of correlation between our analysis and the analysis done by Yu et

al[14](**Supp Fig 2D**), further validating the robustness of the CCN results.

Next, we assessed the extent to which CCN classifications agreed with their nominal

tumor type of origin. We annotated cell lines based their CCN score profile as follows. 'Correct'

Cell lines with CCN score > 0.3 for the tumor type of origin were annotated 'correct'. Those with

CCN scores > 0.3 in the tumor type of origin and at least one other tumor type were annotated

as 'mixed'. Cell lines with CCN scores > 0.3 for tumor types other than that of the cell lines origin

were annotated as 'other', and those lines that did not received a CCN score > 0.3 for any tumor

type were annotated as 'none' (**Fig 2B**). We selected a decision threshold of 0.3 based on the

average of the thresholds that produced the highest Macro F1 measure, harmonic mean of

precision and recall, across 50 cross validations. We found that majority of cell lines originally

annotated as Breast invasive carcinoma (BRCA), Cervical squamous cell carcinoma and

endocervical adenocarcinoma (CESC), Skin Cutaneous Melanoma (SKCM), Colorectal Cancer

(COAD_READ) and Sarcoma (SARC) fell into the 'correct' category (**Fig 2B**). On the other

hand, no Esophageal carcinoma (ESCA) or Brain Lower Grade Glioma (LGG) were classified as

'correct', demonstrating the need for more transcriptionally faithful cell lines that model those

general cancer types.

There are several possible explanations for cell lines not receiving a 'correct'

classification. One possibility is that the sample was incorrectly labeled in the study from which

we harvest the expression data. Consistent with this explanation, we found that colorectal

cancer line NCI-H684[36,37], a cell line labelled as liver hepatocellular carcinoma (LIHC) by CCLE,

was classified strongly as COAD_READ (**Supp Tab 1**). Another possibility to explain low CCN

score is that cell lines were derived from subtypes of tumors that are not well-represented in

TCGA. To explore this hypothesis, we first performed tumor subtype classification on the CCLE

lines from 11 tumor types for which we had trained subtype classifiers (**Supp Tab 2**). We

reasoned that if a cell was a good model for a rarer subtype, then it would receive a poor

general classification but a high classification for the subtype that it models well. Therefore, we

counted the number of lines that fit this pattern. We found that of the 223 lines with no general

classification, 54 (24%) were classified as a specific subtype, suggesting that derivation from

rare subtypes is not the major contributor to the poor overall fidelity of CCLs.

Another potential contributor to low scoring cell lines is intra-tumor stromal and immune

cell impurity in the training data. If impurity were a confounder of CCN scoring, then we would

expect a strong positive correlation between mean purity and mean CCN classification scores of

CCLs per general tumor type. However, the Pearson correlation coefficient between the mean

purity of general tumor type and mean CCN classification scores of CCLs in the corresponding

general tumor type was low (0.059), suggesting that tumor purity is not a major contributor to

the low CCN scores across CCLE (**Supp Fig 2E**).

To more directly assess the impact of intra-tumor heterogeneity in the training data on

evaluating cell lines, we constructed a classifier using cell types found in human melanoma and

glioblastoma scRNA-seq data[38,39]. Previously, we have demonstrated the feasibility of using our

classification approach on scRNA-seq data[23]. Our scRNA-seq classifier achieved a high

average AUPR (0.95) when applied to held-out data (**Supp Fig 3A-B**). Comparing the CCN

score from bulk RNA-seq general classifier and scRNA-seq classifier, we observed a high level

of correlation (Pearson correlation of 0.89) between the SKCM CCN classification scores and

scRNA-seq SKCM malignant CCN classification scores for SKCM cell lines (**Fig 2C, Supp Fig**

3C). Among the 37 SKCM cell lines that were classified as SKCM in general classification, 36

SKCM cell lines were also classified as SKCM malignant cells in scRNA-seq classifier.

Interestingly, we also observed a high correlation between the SARC CCN classification score

and scRNA-seq cancer associated fibroblast (CAF) CCN classification scores (Pearson

correlation of 0.89). Six of the 10 SKCM cell lines that had been classified as SARC by CCN

were classified as CAF by the scRNAseq classifier (**Fig 2D, Supp Fig 3C**), which suggests the

possibility that these cell lines were derived from CAF or other mesenchymal populations, or

that they have acquired a mesenchymal character through their derivation. The high level of

agreement between scRNA-seq and bulk RNA-seq classification results shows that

heterogeneity in the training data of general CCN classifier has little impact in the classification

of SKCM cell lines.

In contrast, we observed a weaker correlation between GBM CCN classification scores

and scRNA-seq GBM neoplastic CCN classification scores (Pearson correlation of 0.58) for

GBM cell lines (**Fig 2E, Supp Fig 3D**). Of the 32 GBM lines that were not classified as GBM

with CCN, 26 were classified as GBM neoplastic cells with the scRNA-seq classifier. Among the

27 GBM lines that were classified as SARC with CCN, 15 cell lines were classified as CAF (**Fig**

2F), and 10 of 15 lines were classified as both GBM neoplastic and CAF in the scRNA-seq

classifier. Similar to the situation with SKCM lines that classify as CAF, this result is consistent

with the possibility that some GBM lines classified as SARC by CCN could be derived from

mesenchymal subtypes exhibiting both strong mesenchymal signatures and glioblastoma

signatures or that they have acquired a mesenchymal character through their derivation[40]. The

lower level of agreement between scRNA-seq and bulk RNA-seq classification results for GBM

models suggests that the heterogeneity of glioblastomas[41] can impact the classification of GBM

cell lines, and that the use of scRNA-seq classifier can resolve this deficiency.

To experimentally explore some of our computational analyses, we performed

immunofluorescence on three cell lines that were not classified as their labelled categories: the

ovarian cancer line SK-OV-3 had a high UCEC CCN score (0.2345), the ovarian cancer line

A2780 had a high Testicular Germ Cell Tumors (TGCT) CCN score (0.2255), and the prostate

cancer line PC-3 had a high bladder cancer (BLCA) score (0.3895) (**Supp Tab 1**). We reasoned

that if SK-OV-3 and A2780 were classified most strongly as UCEC, TGCT and BLCA,

respectively, then they would express proteins that are indicative of these cancer types.

First, we measured the expression of the uterine-associated transcription factor

HOXB6[42,43], and the UCEC serous ovarian tumor biomarker WT1[44] in SK-OV-3, in the OV cell

line Caov-4, and in the UCEC cell line HEC-59. We chose Coav-4 as our positive control for OV

biomarker expression because it was determined by our analysis and others[10,14] to be a good

model of OV. Likewise, we chose HEC-59 to be positive controls for UCEC. We found that SK-

OV-3 has a small percentage (5%) of cells expressing the UCEC marker HOXB6 and a large

portion (73%) of cells expressing WT1 (**Fig 3A**). In contrast, no Coav-4 cells expressing

HOXB6, and 85% of cells expressing WT1. This suggests that SK-OV-3 exhibits both

biomarkers of ovarian tumor and uterine tissue. From our computational analysis and

experimental validation, SK-OV-3 is most likely an endometrioid subtype of ovarian cancer. This

result is also consistent with prior classification of SK-OV-3[45], and the fact that SK-OV-3 lacks

p53 mutations, which is prevalent in high-grade serous ovarian cancer[46], but it does possess an

endometrioid-associated ARID1A mutation[10,45,47]. Next, we measured the expression of markers

of OV and germ cell cancers (LIN28A[48]) in the OV-annotated cell line A2780, which received a

high TCGT CCN score. We found that 54% of A2780 cells expressed LIN28A whereas it was

not detected in Coav-4 (**Fig 3B**). The OV marker WT1 was also expressed in fewer A2780 cells

as compared to Caov-4 (48% vs 85%), which suggests that A2780 is a germ cell derived

ovarian tumor. Taken together, our results suggest that SK-OV-3 and A2780 could represent

OV subtypes of that are not well represented in TCGA training, which resulted in a low OV score

and higher (yet still below our classification threshold) CCN score in other categories.

Lastly, we examined PC-3, annotated as a PRAD cell line but classified to be most

similar to BLCA. We found that 30% of the PC-3 cells expressed PPARG, a contributor to

urothelial differentiation[50] that is not detected in the PRAD Vcap cell line but is highly expressed

in the BLCA RT4 cell line (**Fig 3C**). PC-3 cells also expressed the PRAD biomarker FOLH1[49]

suggesting that PC-3 has an PRAD origin and gained urothelial or luminal characteristics

through the derivation process. In short, our limited experimental data support the CCN

classification results.

Next, we explored the subtype classification of CCLs from three general tumor types in

more depth, focusing first on UCEC. The histologically defined subtypes of UCEC, endometrioid

and serous, differ in prevalence, molecular properties, prognosis, and treatment. For instance,

the endometrioid subtype, which accounts for approximately 80% of uterine cancers, retains

estrogen receptor and progesterone receptor status and is responsive towards progestin

therapy[51,52]. Serous, a more aggressive subtype, is characterized by the loss of estrogen and

progesterone receptor and is not responsive to progestin therapy[51,52]. CCN classified the

majority of the UCEC cell lines as serous except for JHUEM-1 which is classified as

endometrioid (**Fig 4A**). The preponderance CCLE lines of serous versus endometroid character

may be due to properties of serous cancer cells that promote their *in vitro* propagation, such as

upregulation in cell adhesion[53]. Some of our subtype classification results are consistent with

prior observations. For example, HEC-1A, HEC-1B, and KLE were previously characterized as

type II endometrial[54]. On the other hand, our subtype classification results contradict prior

observations in at least one case. For instance, the Ishikawa cell line was derived from type I

endometrial cancer (endometrioid histological subtype)[54,55], however CCN classified a derivative

of this line, Ishikawa 02 ER-, as serous. The high serous CCN score could result from a shift in

phenotype of the line concomitant with its loss of estrogen receptor (ER) as this is a

distinguishing feature of type II endometrial cancer (serous histological subtype)[51]. Taken

together, these results indicate a need for more endometroid-like CCLs.

Next, we examined the subtype classification of Lung Squamous Cell Carcinoma

(LUSC) cell lines (**Fig 4C**). We found that of the 22 lines unclassified or misclassified in the

general classifier, 6 (27%) were classified as a subtype. Among the LUSC cell lines that were

classified as unknown in the general classifier and classified with a subtype, several cell lines

had general classification scores modestly below the threshold. All the LUSC lines with at least

one subtype classification had an underlying primitive subtype classification. This is consistent

either with the ease of deriving lines from tumors with a primitive character, or with a process by

which cell line derivation promotes similarity to more the primitive subtype, which is marked by

increased cellular proliferation[26]. Some of our results are consistent with prior reports that have

investigated the resemblance of some lines to LUSC subtypes. For example, HCC-95, classified

as classical and primitive subtype, has previously been characterized as classical[26,56]. Further,

LUDLU-1, classified as a primitive subtype, has classification signal in classical subtype which

was previously characterized as resembling classical[56]. Likewise, although EPLC-272H was

also classified as primitive subtype, it has relatively high CCN score in the basal subtype, which

corresponds to its previous characterization as basal[56]. Lung Adenocarcinoma (LUAD) cell lines

had classification results similar to LUSC: most lines did not classify as LUAD in the general

classifier (66 of 76) (**Fig 4B**). The cell lines that were classified as a subtype were either

classified as proximal inflammation, proximal proliferation or a mix of the two. RERF-LC-Ad1

had the highest general classification score and the highest proximal inflammation subtype

classification score. Taken together, these subtype classification results have revealed an

absence of cell lines models for basal, classical, and secretory LUSC, and for the Terminal

respiratory unit (TRU) LUAD subtype.

Finally, we sought to measure the extent to which cell line transcriptional fidelity related

to model use. We used the number of papers in which a model was mentioned, normalized by

the number of years since the cell line was documented, as a rough approximation of model

usage. To explore this metric, we plotted the normalized citation count versus general

classification score, labeling the highest cited and highest classified cell lines from each general

tumor type (**Fig 4D**). For most of the general tumor types, the highest cited cell line is not the

highest classified cell line except for Hep G2 and ML-1, representing liver hepatocellular

carcinoma (LIHC) and thyroid carcinoma (THCA), respectively. On the other hand, the general

scores of the highest cited cell lines representing BLCA, SKCM, BRCA, PRAD and

COAD_READ fall below the classification threshold of 0.3. Notably, each of these tumor types

have other lines with scores exceeding 0.7, which should be considered as more faithful

transcriptional models when selecting lines for a study (**Supp Tab 1 and**

http://www.cahanlab.org/resources/cancerCellNet_results/).

Evaluation of patient derived xenografts

Next, we sought to evaluate a more recent class of cancer models: PDX. To do so, we

subjected the RNA-Seq expression profiles of 415 PDX models from 13 different types of

cancer types generated previously[17] to CCN. Similar to the results of CCLE, the PDXs exhibited

a wide range of classification scores (**Fig 5A, Supp Tab 3**). By categorizing the CCN scores of

PDX based on the proportion of samples associated with each tumor type that were correctly

classified**,** we found that SARC, SKCM, COAD_READ and BRCA have higher proportion of

correctly classified PDX than those of other cancer categories (**Fig 5B**). In contrast to CCLE, we

found a higher proportion of correctly classified PDX in Stomach adenocarcinoma (STAD) and

KIRC (**Fig 5B**)**.** However, similar to CCLE, no ESCA PDXs were classified as such. This held

true when we performed subtype classification on PDX samples: none of the PDX in ESCA

were classified as any of the ESCA subtypes (**Supp Tab 4**). UCEC PDXs had both

endometrioid subtypes, serous subtypes, and mixed subtypes, which provided a broader

representation than in CCLE (**Fig 5C**). Many LUSC PDXs that were classified as a subtype

were also classified as Head and Neck squamous cell carcinoma (HNSC) (**Fig 5D**). This could

be due to the similarity in expression profiles of basal and classical subtypes of HNSC and

LUSC[26,57], which is consistent with the observation that these PDXs were also subtyped as

basal and classical. No LUSC PDXs were classified as the secretory subtype. While eight of the

LUAD PDX samples were classified as the unknown subtype class classification, the remaining

six classified as proximal proliferative or proximal inflammatory (**Fig 5E**). Finally, similar to the

CCLE, there were no TRU subtypes in the PDX cohort. In summary, we found that while

individual PDXs can reach extremely high transcriptional fidelity to both general tumor types and

subtypes, many PDXs were not classified as the general tumor type from which they originated.

Evaluation of GEMMs

Next, we used CCN to evaluate GEMMs of six general tumor types from nine studies for

which expression data was publicly available[58–66]. As was true for CCLs and PDXs, GEMMs

also had a wide range of CCN scores (**Fig 6A, Supp Tab 5**). We next categorized the CCN

scores based on the proportion of samples associated with each tumor type that were correctly

classified (**Fig 6B**)**.** In contrast to CCLs and PDXs, the GEMM dataset included multiple

replicates per model, which allowed us to examine intra-GEMM variability. Both at the level of

CCN score and at the level of categorization, GEMMs were highly invariant. For example,

replicate of UCEC GEMMs driven by Prg(cre/+)Pten(lox/lox) received almost identical general

and subtype classification profiles (**Supp Fig 4, Supp Tab 6**). GEMMs sharing genotypes

across studies such as LUAD GEMMs driven by Kras mutation and loss of p53[58,64,66] received

similar general and subtype classification scores (**Fig. 6A,B,D**). Even GEMMs with mixed

classifications received consistent CCN scores. For example, LGG GEMMs, generated by Nf1

mutations expressed in different neural progenitors in combination with Pten deletion[65],

consistently received mixed classification as both LGG and GBM (**Fig 6A**).

To explore the extent to which driver genotype impacted subtype classification, we

examined two general tumor types in which there were GEMMs with different tumor drivers:

LUSC and LUAD. The LUSC GEMMs were generated using loss of Lkb1 and either

overexpression of Sox2 (via two distinct mechanisms) or loss of Pten[64]. Although most of the

lenti-Sox2-Cre-infected;Lkb1$^{fl/fl}$ and Rosa26LSL-Sox2-IRES-GFP;Lkb1$^{fl/fl}$ samples were

classified as unknown, their general LUSC CCN scores were only modestly lower than the

decision threshold and consistently throughout (**Fig 6C**). Those two models also classified

mostly as secretory subtype of LUSC. The consistency is not surprising given both models

overexpress Sox2 and lose Lkb1. Most of the Lkb1$^{fl/fl}$;Pten$^{fl/fl}$ GEMMs received unknown general

classifications with general LUSC CCN scores substantially lower than those of lenti-Sox2-Cre-

infected;Lkb1$^{fl/fl}$ samples and Rosa26LSL-Sox2-IRES-GFP;Lkb1$^{fl/fl}$ samples. Moreover, our

subtype classification indicated that this GEMM was mostly classified as unknown, in contrast to

prior reports suggesting that it is most similar to a basal subtype[67]. The lenti-Sox2-Cre-

infected;Lkb1$^{fl/fl}$ samples received high secretory subtype scores, whereas the Rosa26LSL-

Sox2-IRES-GFP;Lkb1$^{fl/fl}$ samples were classified as a more balanced mix of secretory and

primitive subtypes. None of the three LUSC GEMMs have strong classical or basal sub-type

CCN scores.

All of the LUAD GEMMs, which were generated using various combinations of activating

Kras mutation, loss of Trp53, and loss of Smarca4L[58,64,66], were correctly classified (**Fig 6D**).

There were no substantial differences in general, or subtype classification across driver

genotypes. Notably, the subtypes tended to have CCN scores in mixture of proximal

proliferation, proximal inflammation and TRU. Taken together, this analysis suggests that there

is a degree of similarity, and perhaps plasticity between the primitive and secretory (but not

basal or classical) subtypes of LUSC. On the other hand, while the LUAD GEMMs classify

strongly as LUAD, do not have strong particular subtype classification -- a result that does not

vary by genotype.

Comparison of CCLs, PDXs, and GEMMs

Finally, we sought to estimate the comparative transcriptional fidelity of the three cancer

models modalities, limiting our comparison to those five general tumor types for which there

were at least two examples per modality: UCEC, Pancreatic adenocarcinoma (PAAD), LUSC,

LUAD, and LIHC. We compared the general CCN scores of each model on a per tumor type

basis (**Fig 7A**). In the case of GEMMs, we used the mean classification score of all samples

with shared genotypes. We found that GEMMs had the highest median general classification

scores in four out of the five tumor types. However, some PDXs achieved the highest

classification scores. In UCEC, LUAD and LIHC, the maximum classification score of PDXs

exceeded 0.75 and were thus comparable to the majority of scores on held out TCGA data,

highlighting the potential for PDXs to mirror the transcriptional state of natural tumors (**Fig 7A**).

Because the CCN score is based on a moderate number of gene pairs (i.e. 1647) relative to the

total number of protein-coding genes, it is possible that a cancer model with a high CCN score

might not have a high global similarity to a naturally occurring tumor. Therefore, we also

calculated the GRN status, a metric of the extent to which tumor-type specific gene regulatory

network is established[18], for all models (**Supp Fig 5**). We observed high level of correlation

between the two similarity metrics, which suggests that although CCN classifies on a selected

set of genes, its scores are highly correlated with global assessment of transcriptional similarity.

417 We also sought to compare model modalities in terms of the diversity of subtypes that

418 they represent (**Supp Fig 6**). As a reference, we also included in this analysis the overall

419 subtype incidence, as approximated by incidence in TCGA. In models of UCEC, there is a

420 notable difference in endometroid incidence, and the proportion of models classified as

421 endometroid, with only PDX having any representatives (**Fig 7B**). The vast majority of CCLs

422 and all of the GEMM models of PAAD have an unknown subtype classification. However, the

423 majority of PDXs are subtyped as either a mixture of basal and classical, or basal and classical

424 alone. LUSC have proximal inflammation and proximal proliferation subtypes modelled by CCLs

425 and PDX, and TRU subtype modelled by GEMMs exclusively (**Fig 7B**). Likewise, LUAD have

426 basal, classical and primitive subtypes modelled by CCLs and PDXs, and secretory subtype

427 modelled by GEMMs exclusively (**Fig 7B**). Taken together, these results demonstrate the need

428 to carefully select different model systems to more suitably model certain cancer subtypes.

430 **DISCUSSION**

431 A major goal in the field of cancer biology is to develop models that mimic naturally occurring

432 tumors with enough fidelity to enable therapeutic discoveries. However, methods to measure

433 the extent to which cancer models resemble or diverge from native tumors are lacking. This is

434 especially problematic now because there are many existing models from which to choose, and

435 it has become easier to generate new models. Here, we present CancerCellNet (CCN), a

436 computational tool that measures the similarity of cancer models to 22 naturally occurring tumor

437 types and 36 subtypes. Because CCN is platform- and species-agnostic, it can be applied

438 across many model modalities, including CCLs, PDXs, and GEMMs, and thus it represents a

439 consistent platform to compare models across modalities. Here, we applied CCN to 657 cancer

440 cell lines, 415 patient derived xenografts, and 26 distinct genetically engineered mouse models.

441 Several lessons emerged from our computational analyses that have implications for the field of

442 cancer biology.

First, CancerCellNet indicates that GEMMs are transcriptionally the most faithful models

of four out of five general tumor types for which data from all modalities was available. This is

consistent with the fact that GEMMs are typically derived by recapitulating well-defined driver

mutations of natural tumors, and thus this observation corroborates the importance of genetics

in the etiology of cancer[68]. Moreover, in contrast to most PDXs, GEMMs are typically generated

in immune replete hosts. Therefore, the higher fidelity of GEMMs may also be a result of the

influence of a native immune system on GEMM tumors[69]. Second, PDXs and CCLs have lower

scores that are comparable to each other. This is consistent with the observation that PDXs can

undergo selective pressures in the host that distort the progression of genomic alterations away

from what is observed in natural tumors[70]. Furthermore, the observation that a few PDXs have

very high classification scores, approaching a level that is indistinguishable from held out TCGA

data, suggests that under certain conditions, PDX can almost perfectly mimic natural tumors

transcriptionally. It is unclear what are these conditions; it may be that these few PDXs were

profiled prior to the acquisition of non-typical genomic alterations. Third, we have found that

none of the samples that we evaluated here are transcriptionally adequate models of ESCA,

and therefore this tumor type requires further attention to derive new models. Fourth, we found

that in several tumor types, GEMMs tend to reflect mixtures of subtypes rather than conforming

strongly to single subtypes. The reasons for this are not clear but it is possible that in the cases

that we examined the histologically defined subtypes have a degree of plasticity that is

exacerbated in the murine host environment. We have made the results of our analyses

available online so that researchers can easily explore the performance of selected models or

identify the best models for any of the 22 general tumor types and the 36 subtypes presented

here.

Currently, there are several limitations to our CCN tool, and caveats to our analyses

which indicate areas for future work and improvement. First, CCN is based on transcriptomic

data but other molecular readouts of tumor state, such as profiles of the proteome[71],

epigenome[72], non-coding RNA-ome[72], and genome[68] would be equally, if not more important, to

mimic in a model system. Therefore, it is possible that some models reflect tumor behavior well,

and because this behavior is not well predicted by transcriptome alone, these models have

lower CCN scores. To both measure the extent that such situations exist, and to correct for

them, we plan in the future to incorporate other omic data into CCN so as to make more

accurate and integrated model evaluation possible. A second limitation is that in the cross-

species analysis, CCN implicitly assumes that homologs are functionally equivalent. The extent

to which they are not functionally equivalent determines how confounded the CCN results will

be. This possibility seems to be of limited consequence based on the high performance of the

normal tissue cross-species classifier and based on the fact that GEMMs have the highest

median CCN scores. Finally, the TCGA training data is made up of RNA-Seq from bulk tumor

samples, which necessarily includes non-tumor cells, whereas the CCLs are by definition cell

lines of tumor origin. Therefore, CCLs theoretically could have artificially low CCN scores due to

the presence of non-tumor cells in the training data. This problem appears to be limited as we

found no correlation between tumor purity and CCN score in the CCLE samples. However, this

problem is related to the question of intra-tumor heterogeneity. We demonstrated the feasibility

of using CCN and single cell RNA-seq data to refine the evaluation of cancer cell lines

contingent upon availability of scRNA-seq training data. As more sufficient training single cell

RNA-seq data accrues, CCN would be able to not only evaluate models on a per cell type basis,

but also based on cellular composition.

To ensure that CCN is widely available we have developed a free web application,

which performs CCN analysis on user-uploaded data and allows for direct comparison of their

data to the cancer models evaluated here. We have also made the CCN code freely available

under an Open Source license and as an easily installed R package, and we are actively

supporting its further development. The documentation describes how to analyze model(s) and

compare the results to the panel of models that we evaluated here, thereby allowing

researchers to immediately compare their models to the broader field in a comprehensive and

standard fashion.

Online Methods

Training General CancerCellNet Classifier

To generate training data sets, we downloaded 8991 patient tumor RNA-seq expression

count matrix and their corresponding sample table across 22 different tumor types from TCGA

using TCGAWorkflowData, TCGAbiolinks[73] and SummarizedExperiment[74] packages. We used

all the patient tumor samples for training the general CCN classifier. Later, we found the

intersecting genes between TCGA dataset and all the query samples (CCLs, PDXs, GEMMs),

and used TCGA dataset with intersecting genes as raw input for CCN. To train the top pair

Random Forest classifier, we used a method similar to our previous method[23]. CCN first

normalized the training counts matrix by down-sampling the counts to 500,000 counts per

sample. To significantly reduce the time and resource of generating gene pairs for all possible

genes, CCN then selected 30 up-regulated genes, 30 down-regulated genes and 30 least

differentially expressed genes for each of the 22 cancer categories using template matching[75]

as the genes to generate top scoring gene pairs. In short, for each tumor type, CCN defined a

template vector that labelled the training tumor samples in cancer type of interest as 1 and all

other tumor samples as 0 CCN then calculated the Pearson correlation coefficient between

template vector and gene expressions for all genes. The genes with strong match to template

as either upregulated or downregulated had large absolute Pearson correlation coefficient. CCN

chose the upregulated, downregulated and least differentially expressed genes based on the

magnitude of Pearson correlation coefficient.

After CCN selected the genes for each cancer type, CCN generated gene pairs among

those genes. Gene pair transformation was a method inspired by the top-scoring pair classifier[76]

to allow compatibility of classifier with query expression profiles that were collected through

different platforms (e.g. microarray query data applied to RNA-seq training data). In brief, the

gene pair transformation compares 2 genes within an expression sample and encodes the

"gene1_gene2" gene-pair as 1 if the first gene has higher expression than the second gene.

Otherwise, gene pair transformation would encode the gene-pair as 0. Using all the gene pair

combinations generated through the gene sets across all cancer types, CCN then selected top

75 discriminative gene pairs for each category using template matching (with large absolute

Pearson correlation coefficient) described above. To prevent any single gene from dictating the

gene pair list, we allowed each gene to appear at maximum of three times among the gene

pairs selected as features.

After the top discriminative gene pairs were selected for each cancer categories, CCN

concatenate all the gene pairs into a vector and gene pair transformed the training samples into

a binary matrix with all the discriminative gene pairs as row names and all the training samples

as column names. Using the binary gene pair matrix, CCN randomly shuffled the binary values

across rows then across column generating random profiles that should not resemble training

data from any of the cancer categories. CCN then sampled 70 random profiles, annotated them

as "Unknown" and used them as training data for the "Unknown" category.

Using gene pair binary training matrix, CCN constructed a multi-class Random Forest

classifier of 2000 trees and used stratified sampling of 60 sample size to ensure balance of

training data in constructing the decision trees. The specific parameters for the final CCN

classifier using the function "broadClass_train" in the package cancerCellNet are in **Supp Tab**

7. The gene-pairs are in **Supp Tab 8.**

Validating General CancerCellNet Classifier

Two thirds of patient tumor data from each cancer type was randomly sampled as

training data to construct a CCN classifier. After the classifier was built, 35 held-out samples

from each cancer category were sampled and 40 "Unknown" profiles were generated for

validation. CCN gene pair transformed the held-out data for assessment based on the top gene

pairs selected to construct the classifier. The process of randomly sample training set from 2/3

of all patient tumor data, train classifier and validate using validation set was repeated 50 times

to have a more comprehensive assessment of the classifier. We used precision-recall curve and

area under the precision-recall curve (AUPR) as our metric of assessing the classifiers.

Classifying Query Data into General Cancer Categories

We downloaded the RNA-seq cancer cell lines expression profiles and sample table

from (https://portals.broadinstitute.org/ccle/data), and microarray cancer cell lines expression

profiles and sample table from Barretina et al [35]. We extracted 2 WT control NCCIT RNA-seq

expression profiles from Grow et al[77]. We received PDX expression estimates and sample

annotations from the authors of Gao et al [17]. We gathered GEMM expression profiles from 9

different studies[58–66]. To use CCN classifier on GEMM data, the mouse genes from GEMM

expression profiles were converted into their human homologs. The query samples were gene

pair transformed using gene pairs selected from the training step, and then inputted into

Random Forest classifier for classification. Each query classification profile was labelled as one

of the four classification categories: "correct", "mixed", "none" and "other" based on classification

profiles. If a sample has a CCN score higher than the decision threshold (0.3) in the labelled

cancer category, we assigned that as "correct". If a sample has CCN score higher than the

decision threshold in labelled cancer category and in other cancer categories, we assigned that

as "mixed". If a sample has no CCN score higher than the decision threshold in any cancer

category or has the highest CCN score in 'Unknown' category, then we assigned it as "none". If

a sample has CCN score higher than the decision threshold in a cancer category or categories

not including the labelled cancer category, we assigned it as "other". We analyzed and

visualized the results using R and R packages pheatmap[78] and ggplot2[79].

Cross-Species Assessment

To assess the performance of cross-species classification, we downloaded 1003

labelled human tissue/cell type and 1993 labelled mouse tissue/cell type RNA-seq expression

profiles from Github (https://github.com/pcahan1/CellNet). We first converted the mouse genes

into human homologous genes. Then we found the intersecting genes between mouse

tissue/cell expression profiles and human tissue/cell expression profiles. Limiting the input of

human tissue RNA-seq profiles to the intersecting genes, we trained a CCN classifier with all

the human tissue/cell expression profiles. The parameters used for the function

"broadClass_train" in the package cancerCellNet are in **Supp Tab 7.** After the classifier was

trained, we randomly sampled 75 samples from each tissue category in mouse tissue/cell data

and applied the classifier on those samples to assess performance.

Cross-Technology Assessment

To assess the performance of CCN in applications to microarray data, we gathered

6,219 patient tumor microarray profiles across 12 different cancer types from more than 100

different projects (**Supp Tab 9**). We found the intersecting genes between the microarray

profiles and TCGA patient RNA-seq profiles. Limiting the input of RNA-seq profiles to the

intersecting genes, we created a CCN classifier with all the TCGA patient profiles using

parameters for the function "broadClass_train" listed in **Supp Tab 7**. After the microarray

specific classifier was trained, we randomly sampled 60 microarray patient samples from each

cancer category, and applied CCN classifier on them as assessment of the cross-technology

performance in **Supp Fig 3A**. The same CCN classifier was used to assess microarray CCL

samples **Supp Fig 3B**.

Training and validating scRNA-seq Classifier

We extracted labelled human melanoma and glioblastoma scRNA-seq expression

profiles[38,39], and compiled the two datasets excluding 3 cell types T.CD4, T.CD8 and Meyloid

due to low number of cells for training. 60 cells from each of the 11 cell types were sampled for

training a scRNA-seq classifier. The parameters for training a general scRNA-seq classifier

using the function "broadClass_train" are in **Supp Tab 7.** 25 cells from each of the 11 cell types

from the held-out data were selected to assess the single cell classifier**.** Using the PR curve and

maximizing Macro F1 measure, we selected the decision threshold of 0.255. We then applied

the scRNA-seq classifier on SKCM CCLs and GBM CCLs.

Training Subtype CancerCellNet

We found 11 cancer types (BRCA, COAD, ESCA, HNSC, KIRC, LGG, PAAD, UCEC,

STAD, LUAD, LUSC) which have meaningful subtypes based on either histology or molecular

profile and have sufficient samples to train a subtype classifier with high AUPR. We also

included normal tissues samples from BRCA, COAD, HNSC, KIRC, UCEC to create a normal

tissue category in the construction of their subtype classifiers. Training samples were either

labelled as a cancer subtype for the cancer of interest or as "Unknown" if they belong to other

cancer types. Similar to general classifier training, CCN performed gene pair transformation and

selected the most discriminate gene pairs for each cancer subtype. In addition to the gene pairs

selected to discriminate cancer subtypes, CCN also performed general classification of all

training data and appended the classification profiles of training data with gene pair binary

matrix as additional features. The reason behind using general classification profile as additional

features is that many general cancer types may share similar subtypes, and general

classification profile could be important features to discriminate the general cancer type of

interest from other cancer types before performing finer subtype classification. The specific

parameters used to train individual subtype classifiers using "subClass_train" function of

CancerCellNet package can be found in **Supp Tab 7** and the gene pairs are in **Supp Tab 8**.

Validating Subtype CancerCellNet

Similar to validating general class classifier, we randomly sampled 2/3 of all samples in

each cancer subtype as training data and sampled an equal amount across subtypes in the

held-out data for assessing subtype classifiers. We repeated the process 20 times for more

comprehensive assessment of subtype classifiers.

Cells culture, Immunohistochemistry and histomorphometry

Caov-4 (ATCC® HTB-76™), SK-OV-3(ATCC® HTB-77™), RT4 (ATCC® HTB-2™), and

NCCIT(ATCC® CRL-2073™) cell lines were purchased from ATCC. HEC-59 (C0026001) and

A2780 (93112519-1VL) were obtained from Addexbio Technologies and Sigma-Aldrich. Vcap

and PC-3 (generously provided by Franklin W Huang's lab). SK-OV-3, Vcap, and RT4 were

cultured in Dulbecco's Modified Eagle Medium (DMEM, high glucose, 11960069, Gibco) with

1% Penicillin-Streptomycin-Glutamine (10378016, Life Technologies); Caov-4, PC-3, NCCIT,

and A2780 were cultured using RPMI-1640 medium (11875093, Gibco) while HEC-59 was in

Iscove's Modified Dulbecco's Medium (IMDM, 12440053, Gibco). Both medium were

supplemented with 1% Penicillin-Streptomycin (15140122, Gibco). All medium were added with

10% Fetal Bovine Serum (FBS).

Cells cultured in 48-well plate were washed twice with PBS and fixed in 10% buffered

formalin for 24□hrs at 4 °C. Immunostaining was performed using a standard protocol. Cells

were incubated with primary antibodies to goat HOXB6 (10 µg/mL, PA5-37867, Invitrogen),

mouse WT1(10 µg/mL, MA1-46028, Invitrogen), rabbit PPARG (1:50, ABN1445, Millipore),

mouse FOLH1(10 µg/mL, UM570025, Origene), and rabbit LIN28A (1:50, #3978, Cell Signaling)

in Antibody Diluent (S080981-2, DAKO), at 4□°C overnight followed with three 5 min washes in

TBST. The slides were then incubated with secondary antibodies conjugated with fluorescence

at room temperature for 1□h while avoiding light followed with three 5 min washes in TBST and

nuclear stained with mounting medium containing DAPI. Images were captured by Nikon

EcLipse Ti-S, DS-U3 and DS-Qi2.

Histomorphometry was performed using ImageJ (Version 2.0.0-rc-69/1.52i). % N.postive

cells was calculated by the percentage of the number of positive stained cells divided by the

number of DAPI-positive nucleus within three of randomly chosen areas. The data were

expressed as means ± SD.

Classifying Query Data into Subtypes

We assigned subtype to query sample if the query sample has CCN score higher than

the decision threshold. If a query sample has CCN scores higher than decision threshold, which

was chosen through maximizing Macro F1 measure. The table of decision threshold for subtype

classifiers are in **Supp Tab 10**. If a query sample with no CCN score higher than decision

threshold in any subtype or has the highest CCN score in 'Unknown' category, then we

assigned that sample as 'Unknown'. Analysis and visualizations were done in R and

ComplexHeatmap package[80].

Tumor Purity Analysis

We used the R package ESTIMATE[81] to calculate the ESTIMATE scores from TCGA

tumor expression profiles that we used as training data for CCN classifier. To calculate tumor

purity we used the equation described in YoshiHara et al., 2013[81]:

$$\text{Tumour purity} = \cos\left(0.6049872018 + 0.0001467884 \times \text{ESTIMATE score}\right)$$

671   **Extracting Citation Counts**

672   We used the R package RISmed[82] to extract the number of citations for each cell line

673   through query search of "*cell line name*[Text Word] AND cancer[Text Word]" on PubMed. The

674 citation counts were normalized by dividing the citation counts with the number of years since

675 first documented.

$$Normalized\ citation\ counts = \frac{citation\ counts}{\#\ years\ since\ first\ documented}$$

GRN construction and GRN Status

GRN construction was extended from our previous method[18]. 80 samples per cancer

type were randomly sampled and normalized through down sampling as training data for the

CLR GRN construction algorithm. Cancer type specific GRNs were identified by determining the

differentially expressed genes per each cancer type and extracting the subnetwork using those

genes.

To extend the original GRN status algorithm[18] across different platforms and species, we

devised a rank-based GRN status algorithm. Like the original GRN status, rank based GRN

status is a metric of assessing the similarity of cancer type specific GRN between training data

in the cancer type of interest and query samples. Hence, high GRN status represents high level

of establishment or similarity of the cancer specific GRN in the query sample compared to those

of the training data. The expression profiles of training data and query data were transformed

into rank expression profiles by replacing the expression values with the rank of the expression

values within a sample (highest expressed gene would have the highest rank and lowest

expressed genes would have a rank of 1). Cancer type specific mean and standard deviation of

every gene's rank expression were calculated using training data. The modified Z-score values

for genes within cancer type specific GRN were calculated for query sample's rank expression

profiles to quantify how dissimilar the expression values of genes in query sample's cancer type

specific GRN compared to those of the reference training data:

$$Zscore(gene\ i)_{mod} = \begin{cases} 0, & if\ Zscore\ is\ positive\ and\ the\ gene\ is\ found\ to\ be\ upregulated \\ 0, & if\ Zscore\ is\ negative\ and\ the\ gene\ is\ found\ to\ be\ downregulated \\ & abs(Zscore),\ otherwise \end{cases}$$

If a gene in the cancer type specific GRN is found to be upregulated in the specific

cancer type relative to other cancer types, then we would consider query sample's gene to be

similar if the ranking of the query sample's gene is equal to or greater than the mean ranking of

the gene in training sample. As a result of similarity, we assign that gene of a Z-score of 0. The

same principle applies to cases where the gene is downregulated in cancer specific subnetwork.

GRN status for query sample is calculated as the weighted mean of the $(1000 -$

$Zscore(gene\ i)_{mod})$ across genes in cancer type specific GRN. 1000 is an arbitrary large

number, and larger dissimilarity between query's cancer type specific GRN indicate high Z-

scores for the GRN genes and low GRN status.

$$RGS = \sum_{i=1}^{n} (1000 - Zscore(gene\ i)_{mod}) weight_{gene\ i}$$

$$GRN\ Status = \frac{RGS}{\sum_{i=1}^{n} weight_{gene\ i}}$$

706   The weight of individual genes in the cancer specific network is determined by the

707   importance of the gene in the Random Forest classifier. We later normalize the GRN status in

708   respect to the GRN status of the cancer type of interest and the cancer type with the lowest

709   mean GRN status.

$$Normalized\ GRN\ status = \frac{GRN\ status_{query} - avg(GRN\ status_{min\ cancer})}{avg(GRN\ status_{cancer\ type\ interest})}$$

Where "min cancer" represents the cancer type where its training data have the lowest

mean GRN status in the cancer type of interest, and $avg(GRN\ status_{min\ cancer})$ represents the

average GRN status of cancer type with the lowest average GRN status in the "min cancer".

$avg(GRN\ status_{cancer\ type\ interest})$ represents average GRN status of the cancer type of interest

in the training data.

Code availability

CancerCellNet code and documentation is available at GitHub:

https://github.com/pcahan1/cancerCellNet

Acknowledgements

This work was supported by the National Institutes of Health NCI Ovarian Cancer SPORE

P50CA228991 via a Development Research Program award to PC. FWH was supported by a

Prostate Cancer Foundation Young Investigator Award, Department of Defense W81XWH-17-

PCRP-HD (F.W.H.), the National Institutes of Health/National Cancer Institute P20 CA233255-

01 (F.W.H.) U19 CA214253 (F.W.H.). We would like to thank John Powers, Hao Zhu, Tian-Li

Wang, and Kaloyan Tsanov for comments on the manuscript and helpful discussions.

FIGURE LEGENDS

Fig. 1 CancerCellNet (CCN) workflow and performance. **(A)** Schematic of CCN training (top)

and usage (bottom). CCN was designed to assess and compare the expression profiles of

cancer models such as CCLs, PDXs, and GEMMs with native patient tumors. First, CCN takes

patient tumor expression profiles of 22 different cancer types from TCGA to train a multi-class

Random Forest classifier and performs gene pair transformation on tumor expression profiles.

Then CCN selects the most discriminative gene pairs for each cancer type as features. Lastly,

CCN trains a multi-class Random Forest classifier using gene-pair transformed training data

and feature gene pairs. To use trained classifier, CCN inputs the query samples (e.g.

expression profiles from CCLs, PDXs, GEMMs) and generates a classification profile for the

query samples. The column names of the classification heatmap represent sample annotation

and the row names of the classification heatmap represent different cancer types. Each grid is

colored from black to yellow representing the lowest classification score (e.g. 0) to highest

classification score (e.g. 1). **(B)** Mean and standard deviation of area under the precision recall

curve (AUPR) of classifiers based on 50 iterations of cross-validation: random sampling of

training data, training CCN classifiers using training data and testing the classifiers on held-out

data.

Fig. 2 Evaluation of cancer cell lines. **(A)** General classification heatmap of CCLs extracted from

CCLE. Column annotations of the heatmap represent the labelled cancer category of the CCLs

given by CCLE and the row names of the heatmap represent different cancer categories. CCLs'

general classification profiles are categorized into 4 categories: correct (red), correct mixed

(pink), no classification (light green) and other classification (dark green) based on the decision

threshold of 0.3. **(B)** Bar plot represents the proportion of each classification category in CCLs

across cancer types ordered from the cancer types with the highest proportion of correct and

correct mixed CCLs to lowest proportion. **(C)** Comparison between SKCM general CCN scores

from bulk RNA-seq classifier and SKCM malignant CCN scores from scRNA-seq classifier for

SKCM CCLs. **(D)** Comparison between SARC general CCN scores from bulk RNA-seq

classifier and CAF CCN scores from scRNA-seq classifier for SKCM CCLs. **(E)** Comparison

between GBM general CCN scores from bulk RNA-seq classifier and GBM neoplastic CCN

scores from scRNA-seq classifier for GBM CCLs. **(F)** Comparison between SARC general CCN

scores and CAF CCN scores from scRNA-seq classifier for GBM CCLs. The green lines

indicate the decision threshold for scRNA-seq classifier and general classifier.

Fig. 3 Immunofluorescence of selected cell lines. **(A)** Classification profiles (left) and IF

expression (middle) of Coav-4 (OV positive control), HEC-59 (UCEC positive control) and SK-

OV-3 for WT1 (OV biomarker) and HOXB6 (uterine biomarker). The bar plots quantify the

average percentage of positive cells for WT1 (top-right) and HOXB6 (bottom-right). **(B)**

Classification profiles (left) and IF expression (middle) of Coav-4, NCCIT (germ cell tumor

positive control) and A2780 for WT1 and LIN28A (germ cell tumor biomarker). Classification of

NCCIT were performed using RNA-seq profiles of WT control NCCIT duplicate from Grow et

al[77]. The bar plots quantify the average percentage of positive cells for WT1 (top-right) and

LIN28A (bottom-right) **(C)** Classification profiles (left) and IF expression (middle) of Vcap (PRAD

positive control), RT4 (BLCA positive control) and PC-3 for FOLH1 (prostate biomarker) and

PPARG (urothelial biomarker). The bar plots quantify the average percentage of positive cells

for FOLH1 (top-right) and PPARG (bottom-right).

Fig. 4 Subtype classification of CCLs. The heatmap visualizations represent subtype

classification of UCEC CCLs **(A),** LUAD CCLs **(B)** and LUSC CCLs **(C)**. The row names

represent CCLs and column names represent cancer subtypes of UCEC, LUSC and LUAD. The

bar plots to the right of the subclass heatmaps represent the general classification scores in cell

lines' annotated cancer category, and the color strips to the right of the subclass heatmaps

represent subclass classification (left) and general classification (right). **(D)** Comparison of

normalized citation counts and general CCN classification scores of CCLs. Labelled cell lines

either have the highest CCN classification score in their labelled cancer category or highest

normalized citation count. Each citation count was normalized by number of years since first

documented on PubMed.

Fig. 5 Evaluation of patient derived xenografts. **(A)** General classification heatmap of PDXs.

Column annotations represent annotated cancer type of the PDXs, and row names represent

cancer categories. **(B)** Proportion of classification categories in PDXs across cancer types is

visualized in the bar plot and ordered from the cancer type with highest proportion of correct and

mixed correct classified PDXs to the lowest. Subtype classification heatmaps of UCEC PDXs

(C), LUSC PDXs **(D)** and LUAD PDXs **(E)**.

Fig. 6 Evaluation of genetically engineered mouse models. **(A)** General classification heatmap

of GEMMs. Column annotations represent annotated cancer type of the GEMMs, and row

names represent cancer categories. **(B)** Proportion of classification categories in GEMMs

across cancer types is visualized in the bar plot and ordered from the cancer type with highest

proportion of correct and mixed correct classified GEMMs to the lowest. Subtype classification

heatmap of LUSC GEMMs **(C)** and LUAD GEMMs **(D)**.

Fig. 7 Comparison of CCLs, PDXs, and GEMMs. **(A)** Box-and-whiskers plot comparing general

CCN scores across CCLs, GEMMs, PDXs of five general tumor types. **(B)** Proportion of UCEC

(top-left), PAAD (top-right), LUAD (bottom-left) and LUSC (bottom-right) subtypes across cancer

model modalities and TCGA patient data. For GEMMs, all classification profiles of replicates

with the same genotype from the same study are averaged into one classification profile when

calculating the proportion.

Supplementary Information

Supplementary Figure 1 Assessment of CCN general classifier and subtype classifier. **(A)**

Mean and range of CCN classifier's PR curves from 50 cross validations. **(B)** AUPR of CCN

classifier when applied to independent patient tumor data from ICGC. **(C)** AUPR of CCN human

tissue classifier when applied to mouse tissue data. **(D)** The schematic of training a subtype

classifier in CCN. CCN uses patient tumor expression profiles from cancer of interest as training

data. CCN performs gene-pair transformation and selects the most discriminative gene pairs

among the cancer subtypes from training data as features. CCN then applies the general

classification on training data and uses the general classification profile as features in addition

to gene pairs for training a Random Forest classifier. The weight of the general classification

profiles as features can be tuned to maximize AUPR. **(E)** The mean and standard deviation of

AUPR for 11 subtype classifiers based on 20 iterations of random sampling of training and held-

out data, training subtype classifier using training data, classification of held-out data, and

calculation of recall and precision.

Supplementary Figure 2 Further validation of CCN and classification results. To validate the

cross-platform classification performance of CCN, a new classifier specifically trained to classify

microarray data was trained using RNA-seq data from TCGA as training data and intersecting

genes between RNA-seq data and microarray data. **(A)** AUPR of CCN classifier when applied to

primary microarray testing tumor data. **(B)** Classification heatmap of CCLs using microarray

expression data. **(C)** Pearson correlation between CCN scores of CCLE lines generated from

RNA-seq data and microarray data. **(D)** Comparison between CCLs' CCN scores and the

similarity metric from Yu et al[14], median correlations of transcriptional profiles between CCLs

and TCGA tumors from CCLs' labelled cancer category. **(E)** Comparison of mean tumor purity

of training data and mean CCN scores of CCLs for each cancer category.

Supplementary Figure 3 Single-cell classification of SKCM and GBM cell lines. **(A)**

Classification heatmap of held-out scRNA-seq data. **(B)** AUPR of the scRNA-seq classifier

when applied to scRNA-seq held-out data. **(C)** Single-cell classification of SKCM CCLs. Red

bar-plot (top) represents general CCN scores in SARC and blue bar-plot (bottom) represents

general CCN scores in SKCM. **(D)** Single-cell classification of GBM CCLs. Red bar-plot (top)

represents general CCN scores in SARC and yellow bar-plot (bottom) represents general CCN

scores in GBM.

Supplementary Figure 4 Cancer subtype classification heatmap of UCEC GEMMs.

Supplementary Figure 5 Correlation between cancer type specific network GRN status and
general CCN scores.

Supplementary Figure 6 Proportion of cancer subtypes in different cancer models and TCGA
tumor data across 11 general cancer types.

850
851 **Supplementary Table 1** General classification profiles of CCLs.

852
853 **Supplementary Table 2** Subtype classification profiles of CCLs.

854
855 **Supplementary Table 3** General classification profiles of PDXs.

856
857 **Supplementary Table 4** Subtype classification profiles of PDXs.

858
859 **Supplementary Table 5** General classification profiles of GEMMs

860
861 **Supplementary Table 6** Subtype classification profiles of GEMMs.

862
863 **Supplementary Table 7** Specific parameters used for training of all classifiers.

864
865 **Supplementary Table 8** Gene-pairs selected for final training of CCN general and subtype
866 classifiers.

867
868 **Supplementary Table 9** Accessions of tumor microarray data used in validation.

869
Supplementary Table 10 Decision thresholds for subtype classifiers.

REFERENCES

1. Sharma, S. V., Haber, D. A. & Settleman, J. Cell line-based platforms to evaluate
the therapeutic efficacy of candidate anticancer agents. *Nat. Rev. Cancer* **10,** 241–
253 (2010).
2. Kersten, K., de Visser, K. E., van Miltenburg, M. H. & Jonkers, J. Genetically
engineered mouse models in oncology research and cancer medicine. *EMBO Mol.*
Med. **9,** 137–153 (2017).
3. Hidalgo, M. *et al.* Patient-derived xenograft models: an emerging platform for
translational cancer research. *Cancer Discov.* **4,** 998–1013 (2014).
4. Klijn, C. *et al.* A comprehensive transcriptional portrait of human cancer cell lines.
Nat. Biotechnol. **33,** 306–312 (2015).
5. Koren, S. *et al.* PIK3CA(H1047R) induces multipotency and multi-lineage mammary
tumours. *Nature* **525,** 114–118 (2015).
6. DeRose, Y. S. *et al.* Tumor grafts derived from women with breast cancer
authentically reflect tumor pathology, growth, metastasis and disease outcomes.
Nat. Med. **17,** 1514–1520 (2011).
7. Sharpless, N. E. & Depinho, R. A. The mighty mouse: genetically engineered
mouse models in cancer drug development. *Nat. Rev. Drug Discov.* **5,** 741–754
(2006).
8. Mouradov, D. *et al.* Colorectal cancer cell lines are representative models of the
main molecular subtypes of primary cancer. *Cancer Res.* **74,** 3238–3247 (2014).
9. Stuckelberger, S. & Drapkin, R. Precious GEMMs: emergence of faithful models for
ovarian cancer research. *J. Pathol.* **245,** 129–131 (2018).

10. Domcke, S., Sinha, R., Levine, D. A., Sander, C. & Schultz, N. Evaluating cell lines
as tumour models by comparison of genomic profiles. *Nat. Commun.* **4,** 2126
(2013).
11. Jiang, G. *et al.* Comprehensive comparison of molecular portraits between cell lines
and tumors in breast cancer. *BMC Genomics* **17 Suppl 7,** 525 (2016).
12. Chen, B., Sirota, M., Fan-Minogue, H., Hadley, D. & Butte, A. J. Relating
hepatocellular carcinoma tumor samples and cell lines using gene expression data
in translational research. *BMC Med. Genomics* **8 Suppl 2,** S5 (2015).
13. Vincent, K. M., Findlay, S. D. & Postovit, L. M. Assessing breast cancer cell lines as
tumour models by comparison of mRNA expression profiles. *Breast Cancer Res.*
17, 114 (2015).
14. Yu, K. *et al.* Comprehensive transcriptomic analysis of cell lines as models of
primary tumors across 22 tumor types. *Nat. Commun.* **10,** 3574 (2019).
15. Guernet, A. & Grumolato, L. CRISPR/Cas9 editing of the genome for cancer
modeling. *Methods* **121-122,** 130–137 (2017).
16. Gargiulo, G. Next-Generation in vivo Modeling of Human Cancers. *Front. Oncol.* **8,**
429 (2018).
17. Gao, H. *et al.* High-throughput screening using patient-derived tumor xenografts to
predict clinical trial drug response. *Nat. Med.* **21,** 1318–1325 (2015).
18. Cahan, P. *et al.* CellNet: network biology applied to stem cell engineering. *Cell* **158,**
903–915 (2014).
19. Radley, A. H. *et al.* Assessment of engineered cells using CellNet and RNA-seq.
Nat. Protoc. **12,** 1089–1102 (2017).
20. Tan, Y. & Cahan, P. SingleCellNet: A Computational Tool to Classify Single Cell
RNA-Seq Data Across Platforms and Across Species. *Cell Syst.* **9,** 207–213.e2
(2019).
21. Cancer Genome Atlas Network. Comprehensive molecular characterization of
human colon and rectal cancer. *Nature* **487,** 330–337 (2012).
22. Zhang, J. *et al.* International Cancer Genome Consortium Data Portal--a one-stop
shop for cancer genomics data. *Database (Oxford)* **2011,** bar026 (2011).
23. Tan, Y. & Cahan, P. SingleCellNet: a computational tool to classify single cell RNA-
Seq data across platforms and across species. *BioRxiv* (2018). doi:10.1101/508085
24. Cancer Genome Atlas Network. Comprehensive molecular portraits of human
breast tumours. *Nature* **490,** 61–70 (2012).
25. Parker, J. S. *et al.* Supervised risk predictor of breast cancer based on intrinsic
subtypes. *J. Clin. Oncol.* **27,** 1160–1167 (2009).
26. Wilkerson, M. D. *et al.* Lung squamous cell carcinoma mRNA expression subtypes
are reproducible, clinically important, and correspond to normal cell types. *Clin.*
Cancer Res. **16,** 4864–4875 (2010).
27. Cancer Genome Atlas Research Network. Electronic address:
andrew_aguirre@dfci.harvard.edu & Cancer Genome Atlas Research Network.
Integrated genomic characterization of pancreatic ductal adenocarcinoma. *Cancer*
Cell **32,** 185–203.e13 (2017).
28. Cancer Genome Atlas Research Network *et al.* Integrated genomic characterization
of endometrial carcinoma. *Nature* **497,** 67–73 (2013).

29. Cancer Genome Atlas Research Network *et al.* Integrated genomic characterization
of oesophageal carcinoma. *Nature* **541,** 169–175 (2017).
30. Cancer Genome Atlas Network. Comprehensive genomic characterization of head
and neck squamous cell carcinomas. *Nature* **517,** 576–582 (2015).
31. Cancer Genome Atlas Research Network. Comprehensive molecular
characterization of clear cell renal cell carcinoma. *Nature* **499,** 43–49 (2013).
32. Verhaak, R. G. W. *et al.* Integrated genomic analysis identifies clinically relevant
subtypes of glioblastoma characterized by abnormalities in PDGFRA, IDH1, EGFR,
and NF1. *Cancer Cell* **17,** 98–110 (2010).
33. Cancer Genome Atlas Research Network. Comprehensive molecular profiling of
lung adenocarcinoma. *Nature* **511,** 543–550 (2014).
34. Hu, B. *et al.* Gastric cancer: Classification, histology and application of molecular
pathology. *J. Gastrointest. Oncol.* **3,** 251–261 (2012).
35. Barretina, J. *et al.* The Cancer Cell Line Encyclopedia enables predictive modelling
of anticancer drug sensitivity. *Nature* **483,** 603–607 (2012).
36. Medico, E. *et al.* The molecular landscape of colorectal cancer cell lines unveils
clinically actionable kinase targets. *Nat. Commun.* **6,** 7002 (2015).
37. Park, J.-G. *et al.* Characteristics of Cell Lines Established from Human Colorectal
Carcinoma. *Cancer Res.* (1987).
38. Jerby-Arnon, L. *et al.* A cancer cell program promotes T cell exclusion and
resistance to checkpoint blockade. *Cell* **175,** 984–997.e24 (2018).
39. Darmanis, S. *et al.* Single-Cell RNA-Seq Analysis of Infiltrating Neoplastic Cells at
the Migrating Front of Human Glioblastoma. *Cell Rep.* **21,** 1399–1410 (2017).
40. Lee, J. *et al.* Tumor stem cells derived from glioblastomas cultured in bFGF and
EGF more closely mirror the phenotype and genotype of primary tumors than do
serum-cultured cell lines. *Cancer Cell* **9,** 391–403 (2006).
41. Patel, A. P. *et al.* Single-cell RNA-seq highlights intratumoral heterogeneity in
primary glioblastoma. *Science* **344,** 1396–1401 (2014).
42. Xu, B. *et al.* Regulation of endometrial receptivity by the highly expressed HOXA9,
HOXA11 and HOXD10 HOX-class homeobox genes. *Hum. Reprod.* **29,** 781–790
(2014).
43. Raines, A. M. *et al.* Recombineering-based dissection of flanking and paralogous
Hox gene functions in mouse reproductive tracts. *Development* **140,** 2942–2952
(2013).
44. Netinatsunthorn, W., Hanprasertpong, J., Dechsukhum, C., Leetanaporn, R. &
Geater, A. WT1 gene expression as a prognostic marker in advanced serous
epithelial ovarian carcinoma: an immunohistochemical study. *BMC Cancer* **6,** 90
(2006).
45. Kelly, Z. *et al.* The prognostic significance of specific HOX gene expression patterns
in ovarian cancer. *Int. J. Cancer* **139,** 1608–1617 (2016).
46. Cancer Genome Atlas Research Network. Integrated genomic analyses of ovarian
carcinoma. *Nature* **474,** 609–615 (2011).
47. Wiegand, K. C. *et al.* ARID1A mutations in endometriosis-associated ovarian
carcinomas. *N. Engl. J. Med.* **363,** 1532–1543 (2010).
48. Murray, M. J. *et al.* LIN28 Expression in malignant germ cell tumors downregulates
let-7 and increases oncogene levels. *Cancer Res.* **73,** 4872–4884 (2013).

49. Fair, W. R., Israeli, R. S. & Heston, W. D. Prostate-specific membrane antigen.
Prostate **32,** 140–148 (1997).
50. Biton, A. *et al.* Independent component analysis uncovers the landscape of the
bladder tumor transcriptome and reveals insights into luminal and basal subtypes.
Cell Rep. **9,** 1235–1245 (2014).
51. Black, J. D., English, D. P., Roque, D. M. & Santin, A. D. Targeted therapy in
uterine serous carcinoma: an aggressive variant of endometrial cancer. *Womens*
Health (Lond. Engl.) **10,** 45–57 (2014).
52. Yang, S., Thiel, K. W. & Leslie, K. K. Progesterone: the ultimate endometrial tumor
suppressor. *Trends Endocrinol. Metab.* **22,** 145–152 (2011).
53. Huszar, M. *et al.* Up-regulation of L1CAM is linked to loss of hormone receptors and
E-cadherin in aggressive subtypes of endometrial carcinomas. *J. Pathol.* **220,** 551–
561 (2010).
54. Kozak, J., Wdowiak, P., Maciejewski, R. & Torres, A. A guide for endometrial
cancer cell lines functional assays using the measurements of electronic
impedance. *Cytotechnology* **70,** 339–350 (2018).
55. Korch, C. *et al.* DNA profiling analysis of endometrial and ovarian cell lines reveals
misidentification, redundancy and contamination. *Gynecol. Oncol.* **127,** 241–248
(2012).
56. Wu, D. *et al.* Gene-expression data integration to squamous cell lung cancer
subtypes reveals drug sensitivity. *Br. J. Cancer* **109,** 1599–1608 (2013).
57. Walter, V. *et al.* Molecular subtypes in head and neck cancer exhibit distinct
patterns of chromosomal gain and loss of canonical cancer genes. *PLoS One* **8,**
e56823 (2013).
58. Adeegbe, D. O. *et al.* BET Bromodomain Inhibition Cooperates with PD-1 Blockade
to Facilitate Antitumor Response in Kras-Mutant Non-Small Cell Lung Cancer.
Cancer Immunol Res **6,** 1234–1245 (2018).
59. Blaisdell, A. *et al.* Neutrophils oppose uterine epithelial carcinogenesis via
debridement of hypoxic tumor cells. *Cancer Cell* **28,** 785–799 (2015).
60. Fitamant, J. *et al.* YAP inhibition restores hepatocyte differentiation in advanced
HCC, leading to tumor regression. *Cell Rep.* **10,** 1692–1707 (2015).
61. Jia, D. *et al.* Crebbp loss drives small cell lung cancer and increases sensitivity to
HDAC inhibition. *Cancer Discov.* **8,** 1422–1437 (2018).
62. Kress, T. R. *et al.* Identification of MYC-Dependent Transcriptional Programs in
Oncogene-Addicted Liver Tumors. *Cancer Res.* **76,** 3463–3472 (2016).
63. Li, L. *et al.* GKAP acts as a genetic modulator of NMDAR signaling to govern
invasive tumor growth. *Cancer Cell* **33,** 736–751.e5 (2018).
64. Mollaoglu, G. *et al.* The Lineage-Defining Transcription Factors SOX2 and NKX2-1
Determine Lung Cancer Cell Fate and Shape the Tumor Immune
Microenvironment. *Immunity* **49,** 764–779.e9 (2018).
65. Pan, Y. *et al.* Whole tumor RNA-sequencing and deconvolution reveal a clinically-
prognostic PTEN/PI3K-regulated glioma transcriptional signature. *Oncotarget* **8,**
52474–52487 (2017).
66. Lissanu Deribe, Y. *et al.* Mutations in the SWI/SNF complex induce a targetable
dependence on oxidative phosphorylation in lung cancer. *Nat. Med.* **24,** 1047–1057
(2018).

67. Xu, C. *et al.* Loss of Lkb1 and Pten leads to lung squamous cell carcinoma with
elevated PD-L1 expression. *Cancer Cell* **25,** 590–604 (2014).
68. Stratton, M. R., Campbell, P. J. & Futreal, P. A. The cancer genome. *Nature* **458,**
719–724 (2009).
69. Balkwill, F. R., Capasso, M. & Hagemann, T. The tumor microenvironment at a
glance. *J. Cell Sci.* **125,** 5591–5596 (2012).
70. Ben-David, U. *et al.* Patient-derived xenografts undergo mouse-specific tumor
evolution. *Nat. Genet.* **49,** 1567–1575 (2017).
71. Hristova, V. A. & Chan, D. W. Cancer biomarker discovery and translation:
proteomics and beyond. *Expert Rev Proteomics* **16,** 93–103 (2019).
72. Dawson, M. A. & Kouzarides, T. Cancer epigenetics: from mechanism to therapy.
Cell **150,** 12–27 (2012).
73. Silva, T. C. *et al.* TCGA Workflow: Analyze cancer genomics and epigenomics data
using Bioconductor packages. [version 2; peer review: 1 approved, 2 approved with
reservations]. *F1000Res.* **5,** 1542 (2016).
74. Morgan, M., Obenchain, V., Hester, J. & Pag`es, H. *SummarizedExperiment:*
SummarizedExperiment container. (2018).
75. Pavlidis, P. & Noble, W. S. Analysis of strain and regional variation in gene
expression in mouse brain. *Genome Biol.* **2,** RESEARCH0042 (2001).
76. Geman, D., d Avignon, C., Naiman, D. Q. & Winslow, R. L. Classifying gene
expression profiles from pairwise mRNA comparisons. *Stat Appl Genet Mol Biol* **3,**
Article19 (2004).
77. Grow, E. J. *et al.* Intrinsic retroviral reactivation in human preimplantation embryos
and pluripotent cells. *Nature* **522,** 221–225 (2015).
78. Kolde, R. *pheatmap: Pretty Heatmaps.* (CRAN, 2019).
79. Wickham, H. *ggplot2 - Elegant Graphics for Data Analysis* . (Springer-Verlag New
York, 2016). doi:10.1007/978-0-387-98141-3
80. Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations
in multidimensional genomic data. *Bioinformatics* **32,** 2847–2849 (2016).
81. Yoshihara, K. *et al.* Inferring tumour purity and stromal and immune cell admixture
from expression data. *Nat. Commun.* **4,** 2612 (2013).
82. Kovalchik, S. *RISmed: Download Content from NCBI Databases.* (CRAN.R-project,
2017).
1065

Figure 1

A

B

Figure 2

Figure 3

Figure 4

Figure 5

A

PDX Classification

BLCA
BRCA
CESC
COAD_READ
ESCA
GBM
HNSC
KIRC
KIRP
LGG
LIHC
LUAD
LUSC
OV
PAAD
PRAD
SARC
SKCM
STAD
TGCT
THCA
UCEC
Unknown

group
BRCA
COAD_READ
ESCA
KIRC
LIHC
LUAD
LUSC
OV
PAAD
SARC
SKCM
STAD
UCEC

B

COAD_READ
SARC
BRCA
SKCM
PAAD
OV
KIRC
STAD
UCEC
LIHC
LUAD
LUSC
ESCA

Correct
Mixed
None
Other

C

General Classification
General CCN Scores (UCEC)
Subclass Classification
Endometrioid
Serous
Solid Tissue Normal
Unknown

General Classification
UCEC
Not Classified
SARC
CESC
Subclass Classification
Endometrioid
Serous
Unknown

D

General Classification
General CCN Scores (LUSC)
Subclass Classification
basal
classical
primitive
secretory
Unknown

General Classification
CESC
HNSC
LUAD
LUSC
Not Classified
Subclass Classification
basal
classical
primitive
Unknown

E

General Classification
General CCN Scores (LUAD)
Subclass Classification
prox.-inflam
prox.-prolif.
TRU
Unknown

General Classification
LUAD
Not Classified
SARC
COAD_READ
Subclass Classification
prox.-inflam
prox.-prolif.
Unknown

Figure 6

A

B

C

D

Figure 7

A

B

Supplemental Figure 1

A

Classifier performance

B

C

D

E

Supplemental Figure 2

Supplemental Figure 3

Supplemental Figure 4

Supplemental Figure 5

Supplemental Figure 6