7 **Evaluating the transcriptional fidelity of cancer models**

9
10 Da Peng[1*], Rachel Gleyzer[2*], Wen-Hsin Tai[2], Pavithra Kumar[2], Qin Bian[2], Bradley Issacs[2],
11 Edroaldo Lummertz da Rocha[3], Stephanie Cai[1], Kathleen DiNapoli[4,5], Franklin W Huang[6],
12 Patrick Cahan[1,2,7]

13
14 [1]Department of Biomedical Engineering, Johns Hopkins University School of Medicine,
15 Baltimore MD 21205 USA

16
17 [2]Institute for Cell Engineering, Johns Hopkins University School of Medicine,
18 Baltimore MD 21205 USA

19
20 [3]Department of Microbiology, Immunology and Parasitology,
21 Federal University of Santa Catarina, Florianópolis SC, Brazil

22
23 [4]Department of Cell Biology, Johns Hopkins University School of Medicine,
24 Baltimore, MD 21205 USA

25
26 [5]Department of Electrical and Computer Engineering, Johns Hopkins University,
27 Baltimore MD 21218 USA

28
29 [6]Division of Hematology/Oncology, Department of Medicine; Helen Diller Family Cancer Center;
30 Bakar Computational Health Sciences Institute; Institute for Human Genetics;
31 University of California, San Francisco, San Francisco, CA

32
33 [7]Department of Molecular Biology and Genetics, Johns Hopkins University School of Medicine,
34 Baltimore MD 21205 USA

35
36
37 * These authors made equal contributions.

38
39
40 **Correspondence to: patrick.cahan@jhmi.edu**

42 Article type: Research

43
44 Website: http://www.cahanlab.org/resources/cancerCellNet_web

45
46 Code: https://github.com/pcahan1/cancerCellNet

47
48
49
50

ABSTRACT

Background: Cancer researchers use cell lines, patient derived xenografts, engineered mice, and tumoroids as models to investigate tumor biology and to identify therapies. The generalizability and power of a model derives from the fidelity with which it represents the tumor type under investigation, however, the extent to which this is true is often unclear. The preponderance of models and the ability to readily generate new ones has created a demand for tools that can measure the extent and ways in which cancer models resemble or diverge from native tumors.

Methods: We developed a machine learning based computational tool, CancerCellNet, that measures the similarity of cancer models to 22 naturally occurring tumor types and 36 subtypes, in a platform and species agnostic manner. We applied this tool to 657 cancer cell lines, 415 patient derived xenografts, 26 distinct genetically engineered mouse models, and 131 tumoroids. We validated CancerCellNet by application to independent data, and we tested several predictions with immunofluorescence.

Results: We have documented the cancer models with the greatest transcriptional fidelity to natural tumors, we have identified cancers underserved by adequate models, and we have found models with annotations that do not match their classification. By comparing models across modalities, we report that, on average, genetically engineered mice and tumoroids have higher transcriptional fidelity than patient derived xenografts and cell lines in four out of five tumor types. However, several patient derived xenografts and tumoroids have classification scores that are on par with native tumors, highlighting both their potential as faithful model classes and their heterogeneity.

Conclusions: CancerCellNet enables the rapid assessment of transcriptional fidelity of tumor models. We have made CancerCellNet available as freely downloadable software and as a web application that can be applied to new cancer models that allows for direct comparison to the cancer models evaluated here.

INTRODUCTION

Models are widely used to investigate cancer biology and to identify potential therapeutics. Popular modeling modalities are cancer cell lines (CCLs)[1], genetically engineered mouse models (GEMMs)[2], patient derived xenografts (PDXs)[3], and tumoroids[4]. These classes of models differ in the types of questions that they are designed to address. CCLs are often used to address cell intrinsic mechanistic questions[5], GEMMs to chart progression of molecularly defined-disease[6], and PDXs to explore patient-specific response to therapy in a physiologically relevant context[7]. More recently, tumoroids have emerged as relatively inexpensive, physiological, in vitro 3D models of tumor epithelium with applications ranging from measuring drug responsiveness to exploring tumor dependence on cancer stem cells. Models also differ in the extent to which the they represent specific aspects of a cancer type[8]. Even with this intra- and inter-class model variation, all models should represent the tumor type or subtype under investigation, and not another type of tumor, and not a non-cancerous tissue. Therefore, cancer-models should be selected not only based on the specific biological question but also based on the similarity of the model to the cancer type under investigation[9,10].

Various methods have been proposed to determine the similarity of cancer models to their intended subjects. Domcke et al devised a 'suitability score' as a metric of the molecular similarity of CCLs to high grade serous ovarian carcinoma based on a heuristic weighting of copy number alterations, mutation status of several genes that distinguish ovarian cancer subtypes, and hypermutation status[11]. Other studies have taken analogous approaches by either focusing on transcriptomic or ensemble molecular profiles (e.g. transcriptomic and copy number alterations) to quantify the similarity of cell lines to tumors[12–14]. These studies were tumor-type specific, focusing on CCLs that model, for example, hepatocellular carcinoma or breast cancer. Notably, Yu et al compared the transcriptomes of CCLs to The Cancer Genome Atlas (TCGA) by correlation analysis, resulting in a panel of CCLs recommended as most representative of 22 tumor types[15]. Most recently, Najgebauer et al[16] and Salvadores et al[17]

have developed methods to assess CCLs using molecular traits such as copy number

alterations (CNA), somatic mutations, DNA methylation and transcriptomics. While all of these

studies have provided valuable information, they leave two major challenges unmet. The first

challenge is to determine the fidelity of GEMMs, PDXs, and tumoroids, and whether there are

stark differences between these classes of models and CCLs. The other major unmet challenge

is to enable the rapid assessment of new, emerging cancer models. This challenge is especially

relevant now as technical barriers to generating models have been substantially lowered[18,19],

and because new models such as PDXs and tumoroids can be derived on patient-specific basis

therefore should be considered a distinct entity requiring individual validation[4,20].

To address these challenges, we developed CancerCellNet (CCN), a computational tool

that uses transcriptomic data to quantitatively assess the similarity between cancer models and

22 naturally occurring tumor types and 36 subtypes in a platform- and species-agnostic manner.

Here, we describe CCN's performance, and the results of applying it to assess 657 CCLs, 415

PDXs, 26 GEMMs, and 131 tumoroids. This has allowed us to identify the most faithful models

currently available, to document cancers underserved by adequate models, and to find models

with inaccurate tumor type annotation. Moreover, because CCN is open-source and easy to

use, it can be readily applied to newly generated cancer models as a means to assess their

fidelity.

RESULTS

CancerCellNet classifies samples accurately across species and technologies

Previously, we had developed a computational tool using the Random Forest

classification method to measure the similarity of engineered cell populations to their *in vivo*

counterparts based on transcriptional profiles[21,22]. More recently, we elaborated on this

approach to allow for classification of single cell RNA-seq data in a manner that allows for

cross-platform and cross-species analysis[23]. Here, we used an analogous approach to build a

platform that would allow us to quantitatively compare cancer models to naturally occurring

patient tumors (**Fig 1A**). In brief, we used TCGA RNA-seq expression data from 22 solid tumor

types to train a top-pair multi-class Random forest classifier (**Fig 1B**). We combined training

data from Rectal Adenocarcinoma (READ) and Colon Adenocarcinoma (COAD) into one

COAD_READ category because READ and COAD are considered to be virtually

indistinguishable at a molecular level[24]. We included an 'Unknown' category trained using

randomly shuffled gene-pair profiles generated from the training data of 22 tumor types to

identify query samples that are not reflective of any of the training data. To estimate the

performance of CCN and how it is impacted by parameter variation, we performed a parameter

sweep with a 5-fold 2/3 cross-validation strategy (i.e. 2/3 of the data sampled across each

cancer type was used to train, 1/3 was used to validate) (**Fig 1C**). The performance of CCN, as

measured by the mean area under the precision recall curve (AUPRC), did not fall below 0.945

and remained relatively stable across parameter sets (**Supp Fig 1A**). The optimal parameters

resulted in 1,979 features. The mean AUPRCs exceeded 0.95 in most tumor types with this

optimal parameter set (**Fig 1D, Supp Fig 1B**). The AUPRCs of CCN applied to independent

data RNA-Seq data from 725 tumors across five tumor types from the International Cancer

Genome Consortium (ICGC)[25] ranged from 0.93 to 0.99, supporting the notion that the platform

is able to accurately classify tumor samples from diverse sources (**Fig 1E**).

As one of the central aims of our study is to compare distinct cancer models, including

GEMMs, our method needed to be able to classify samples from mouse and human samples

equivalently. We used the Top-Pair transform[23] to achieve this and we tested the feasibility of

this approach by assessing the performance of a normal (i.e. non-tumor) cell and tissue

classifier trained on human data as applied to mouse samples. Consistent with prior

applications[23], we found that the cross-species classifier performed well, achieving mean

AUPRC of 0.97 when applied to mouse data (**Supp Fig 1C**)**.**

To evaluate cancer models at a finer resolution, we also developed an approach to

perform tumor subtype classifications (**Supp Fig 1D**). We constructed 11 different cancer

subtype classifiers based on the availability of expression or histological subtype

information[24,26–36]. We also included non-cancerous, normal tissues as categories for several

subtype classifiers when sufficient data was available: breast invasive carcinoma (BRCA),

COAD_READ, head and neck squamous cell carcinoma (HNSC), kidney renal clear cell

carcinoma (KIRC) and uterine corpus endometrial carcinoma (UCEC). The 11 subtype

classifiers all achieved high overall average AUPRs ranging from 0.80 to 0.99 (**Supp Fig 1E**).

Fidelity of cancer cell lines

Having validated the performance of CCN, we then used it to determine the fidelity of

CCLs. We mined RNA-seq expression data of 657 different cell lines across 20 cancer types

from the Cancer Cell Line Encyclopedia (CCLE) and applied CCN to them, finding a wide

classification range for cell lines of each tumor type (**Fig 2A, Supp Tab 1**). To verify the

classification results, we applied CCN to expression profiles from CCLE generated through

microarray expression profiling[37]. To ensure that CCN would function on microarray data, we

first tested it by applying a CCN classifier created to test microarray data to 720 expression

profiles of 12 tumor types. The cross-platform CCN classifier performed well, based on the

comparison to study-provided annotation, achieving a mean AUPRC of 0.91 (**Supp Fig 2A**).

Next, we applied this cross-platform classifier to microarray expression profiles from CCLE

(**Supp Fig 2B**). From the classification results of 571 cell lines that have both RNA-seq and

microarray expression profiles, we found a strong overall positive association between the

classification scores from RNA-seq and those from microarray (**Supp Fig 2C**). This comparison

supports the notion that the classification scores for each cell line are not artifacts of profiling

methodology. Moreover, this comparison shows that the scores are consistent between the

times that the cell lines were first assayed by microarray expression profiling in 2012 and by

RNA-Seq in 2019. We also observed high level of correlation between our analysis and the

analysis done by Yu et al[15](**Supp Fig 2D**), further validating the robustness of the CCN results.

Next, we assessed the extent to which CCN classifications agreed with their nominal

tumor type of origin, which entailed translating quantitative CCN scores to classification labels.

To achieve this, we selected a decision threshold that maximized the Macro F1 measure,

harmonic mean of precision and recall, across 50 cross validations. Then, we annotated cell

lines based their CCN score profile as follows. Cell lines with CCN scores > threshold for the

tumor type of origin were annotated as 'correct'. Cell lines with CCN scores > threshold in the

tumor type of origin and at least one other tumor type were annotated as 'mixed'. Cell lines with

CCN scores > threshold for tumor types other than that of the cell line's origin were annotated

as 'other'. Cell lines that did not receive a CCN score > threshold for any tumor type were

annotated as 'none' (**Fig 2B**). We found that majority of cell lines originally annotated as Breast

invasive carcinoma (BRCA), Cervical squamous cell carcinoma and endocervical

adenocarcinoma (CESC), Skin Cutaneous Melanoma (SKCM), Colorectal Cancer

(COAD_READ) and Sarcoma (SARC) fell into the 'correct' category (**Fig 2B**). On the other

hand, no Esophageal carcinoma (ESCA), Pancreatic adenocarcinoma (PAAD) or Brain Lower

Grade Glioma (LGG) were classified as 'correct', demonstrating the need for more

transcriptionally faithful cell lines that model those general cancer types.

There are several possible explanations for cell lines not receiving a 'correct'

classification. One possibility is that the sample was incorrectly labeled in the study from which

we harvested the expression data. Consistent with this explanation, we found that colorectal

cancer line NCI-H684[38,39], a cell line labelled as liver hepatocellular carcinoma (LIHC) by CCLE,

was classified strongly as COAD_READ (**Supp Tab 1**). Another possibility to explain low CCN

score is that cell lines were derived from subtypes of tumors that are not well-represented in

TCGA. To explore this hypothesis, we first performed tumor subtype classification on CCLs from

11 tumor types for which we had trained subtype classifiers (**Supp Tab 2**). We reasoned that if

a cell was a good model for a rarer subtype, then it would receive a poor general classification

but a high classification for the subtype that it models well. Therefore, we counted the number of

lines that fit this pattern. We found that of the 188 lines with no general classification, 25 (13%)

were classified as a specific subtype, suggesting that derivation from rare subtypes is not the

major contributor to the poor overall fidelity of CCLs.

Another potential contributor to low scoring cell lines is intra-tumor stromal and immune

cell impurity in the training data. If impurity were a confounder of CCN scoring, then we would

expect a strong positive correlation between mean purity and mean CCN classification scores of

CCLs per general tumor type. However, the Pearson correlation coefficient between the mean

purity of general tumor type and mean CCN classification scores of CCLs in the corresponding

general tumor type was low (0.14), suggesting that tumor purity is not a major contributor to the

low CCN scores across CCLs (**Supp Fig 2E**).

Comparison of SKCM and GBM CCLs to scRNA-seq

To more directly assess the impact of intra-tumor heterogeneity in the training data on

evaluating cell lines, we constructed a classifier using cell types found in human melanoma and

glioblastoma scRNA-seq data[40,41]. Previously, we have demonstrated the feasibility of using our

classification approach on scRNA-seq data[23]. Our scRNA-seq classifier achieved a high

average AUPRC (0.95) when applied to held-out data and high mean AUPRC (0.99) when

applied to few purified bulk testing samples (**Supp Fig 3A-B**). Comparing the CCN score from

bulk RNA-seq general classifier and scRNA-seq classifier, we observed a high level of

correlation (Pearson correlation of 0.89) between the SKCM CCN classification scores and

scRNA-seq SKCM malignant CCN classification scores for SKCM cell lines (**Fig 2C, Supp Fig**

3C). Of the 41 SKCM cell lines that were classified as SKCM by the bulk classifier, 37 were also

classified as SKCM malignant cells by the scRNA-seq classifier. Interestingly, we also observed

a high correlation between the SARC CCN classification score and scRNA-seq cancer

associated fibroblast (CAF) CCN classification scores (Pearson correlation of 0.92). Six of the

seven SKCM cell lines that had been classified as exclusively SARC by CCN were classified as

CAF by the scRNA-seq classifier (**Fig 2D, Supp Fig 3C**), which suggests the possibility that

these cell lines were derived from CAF or other mesenchymal populations, or that they have

acquired a mesenchymal character through their derivation. The high level of agreement

between scRNA-seq and bulk RNA-seq classification results shows that heterogeneity in the

training data of general CCN classifier has little impact in the classification of SKCM cell lines.

In contrast, we observed a weaker correlation between GBM CCN classification scores

and scRNA-seq GBM neoplastic CCN classification scores (Pearson correlation of 0.72) for

GBM cell lines (**Fig 2E, Supp Fig 3D**). Of the 31 GBM lines that were not classified as GBM

with CCN, 25 were classified as GBM neoplastic cells with the scRNA-seq classifier. Among the

22 GBM lines that were classified as SARC with CCN, 15 cell lines were classified as CAF (**Fig**

2F), 10 which were classified as both GBM neoplastic and CAF in the scRNA-seq classifier.

Similar to the situation with SKCM lines that classify as CAF, this result is consistent with the

possibility that some GBM lines classified as SARC by CCN could be derived from

mesenchymal subtypes exhibiting both strong mesenchymal signatures and glioblastoma

signatures or that they have acquired a mesenchymal character through their derivation. The

lower level of agreement between scRNA-seq and bulk RNA-seq classification results for GBM

models suggests that the heterogeneity of glioblastomas[42] can impact the classification of GBM

cell lines, and that the use of scRNA-seq classifier can resolve this deficiency.

Immunofluorescence confirmation of CCN predictions

To experimentally explore some of our computational analyses, we performed

immunofluorescence on three cell lines that were not classified as their labelled categories: the

ovarian cancer line SK-OV-3 had a high UCEC CCN score (0.246), the ovarian cancer line

A2780 had a high Testicular Germ Cell Tumors (TGCT) CCN score (0.327), and the prostate

cancer line PC-3 had a high bladder cancer (BLCA) score (0.307) (**Supp Tab 1**). We reasoned

that if SK-OV-3, A2780 and PC-3 were classified most strongly as UCEC, TGCT and BLCA,

respectively, then they would express proteins that are indicative of these cancer types.

First, we measured the expression of the uterine-associated transcription factor

HOXB6[43,44], and the UCEC serous ovarian tumor biomarker WT1[45] in SK-OV-3, in the OV cell

line Caov-4, and in the UCEC cell line HEC-59. We chose Caov-4 as our positive control for OV

biomarker expression because it was determined by our analysis and others[11,15] to be a good

model of OV. Likewise, we chose HEC-59 to be a positive control for UCEC. We found that SK-

OV-3 has a small percentage (5%) of cells that expressed the uterine marker HOXB6 and a

large proportion (73%) of cells that expressed WT1 (**Fig 3A**). In contrast, no Caov-4 cells

expressed HOXB6, whereas 85% of cells expressed WT1. This suggests that SK-OV-3 exhibits

both biomarkers of ovarian tumor and uterine tissue. From our computational analysis and

experimental validation, SK-OV-3 is most likely an endometrioid subtype of ovarian cancer. This

result is also consistent with prior classification of SK-OV-3[46], and the fact that SK-OV-3 lacks

p53 mutations, which is prevalent in high-grade serous ovarian cancer[47], and it harbors an

endometrioid-associated mutation in ARID1A[11,46,48]. Next, we measured the expression of

markers of OV and germ cell cancers (LIN28A[49]) in the OV-annotated cell line A2780, which

received a high TCGT CCN score. We found that 54% of A2780 cells expressed LIN28A

whereas it was not detected in Caov-4 (**Fig 3B**). The OV marker WT1 was also expressed in

fewer A2780 cells as compared to Caov-4 (48% vs 85%), which suggests that A2780 could be a

germ cell derived ovarian tumor. Taken together, our results suggest that SK-OV-3 and A2780

could represent OV subtypes of that are not well represented in TCGA training data, which

resulted in a low OV score and higher CCN score in other categories.

Lastly, we examined PC-3, annotated as a PRAD cell line but classified to be most

similar to BLCA. We found that 30% of the PC-3 cells expressed PPARG, a contributor to

urothelial differentiation[50] that is not detected in the PRAD Vcap cell line but is highly expressed

in the BLCA RT4 cell line (**Fig 3C**). PC-3 cells also expressed the PRAD biomarker FOLH1[51]

suggesting that PC-3 has an PRAD origin and gained urothelial or luminal characteristics

through the derivation process. In short, our limited experimental data support the CCN

classification results.

Subtype classification of cancer cell lines

Next, we explored the subtype classification of CCLs from three general tumor types in

more depth. We focused our subtype visualization (**Fig 4A-C**) on CCL models with general CCN

score above 0.1 in their nominal cancer type as this allowed us to analyze those models that fell

below the general threshold but were classified as a specific sub-type (**Supp Tab 1-2**).

Focusing first on UCEC, the histologically defined subtypes of UCEC, endometrioid and serous,

differ in prevalence, molecular properties, prognosis, and treatment. For instance, the

endometrioid subtype, which accounts for approximately 80% of uterine cancers, retains

estrogen receptor and progesterone receptor status and is responsive towards progestin

therapy[52,53]. Serous, a more aggressive subtype, is characterized by the loss of estrogen and

progesterone receptor and is not responsive to progestin therapy[52,53]. CCN classified the

majority of the UCEC cell lines as serous except for JHUEM-1 which is classified as mixed, with

similarities to both endometrioid and serous (**Fig 4A**). The preponderance CCLE lines of serous

versus endometroid character may be due to properties of serous cancer cells that promote

their *in vitro* propagation, such as upregulation of cell adhesion transcriptional programs[54].

Some of our subtype classification results are consistent with prior observations. For example,

HEC-1A, HEC-1B, and KLE were previously characterized as type II endometrial cancer, which

includes a serous histological subtype[55]. On the other hand, our subtype classification results

contradict prior observations in at least one case. For instance, the Ishikawa cell line was

derived from type I endometrial cancer (endometrioid histological subtype)[55,56], however CCN

classified a derivative of this line, Ishikawa 02 ER-, as serous. The high serous CCN score

could result from a shift in phenotype of the line concomitant with its loss of estrogen receptor

(ER) as this is a distinguishing feature of type II endometrial cancer (serous histological

subtype)[52]. Taken together, these results indicate a need for more endometroid-like CCLs.

Next, we examined the subtype classification of Lung Squamous Cell Carcinoma

(LUSC) and Lung adenocarcinoma (LUAD) cell lines (Fig 4B-C). All the LUSC lines with at least

one subtype classification had an underlying primitive subtype classification. This is consistent

either with the ease of deriving lines from tumors with a primitive character, or with a process by

which cell line derivation promotes similarity to more primitive subtype, which is marked by

increased cellular proliferation[28]. Some of our results are consistent with prior reports that have

investigated the resemblance of some lines to LUSC subtypes. For example, HCC-95,

previously been characterized as classical[28,57], had a maximum CCN score in the classical

subtype (0.429) . Similarly, LUDLU-1 and EPLC-272H, previously reported as classical[57] and

basal[57] respectively, had maximal tumor subtype CCN scores for these sub-types (0.323 and

0.256) (**Fig 4B, Supp Tab 2**) despite classified as Unknown**.** Lastly, the LUAD cell lines that

were classified as a subtype were either classified as proximal inflammation or proximal

proliferation (**Fig 4C**). RERF-LC-Ad1 had the highest general classification score and the

highest proximal inflammation subtype classification score. Taken together, these subtype

classification results have revealed an absence of cell lines models for basal and secretory

LUSC, and for the Terminal respiratory unit (TRU) LUAD subtype.

Cancer cell lines' popularity and transcriptional fidelity

Finally, we sought to measure the extent to which cell line transcriptional fidelity related

to model prevalence. We used the number of papers in which a model was mentioned,

normalized by the number of years since the cell line was documented, as a rough

approximation of model prevalence. To explore this relationship, we plotted the normalized

citation count versus general classification score, labeling the highest cited and highest

classified cell lines from each general tumor type (**Fig 4D**). For most of the general tumor types,

the highest cited cell line is not the highest classified cell line except for Hep G2, AGS and ML-

1, representing liver hepatocellular carcinoma (LIHC), stomach adenocarcinoma (STAD), and

thyroid carcinoma (THCA), respectively. On the other hand, the general scores of the highest

cited cell lines representing BLCA (T24), BRCA (MDA-MB-231), and PRAD (PC-3) fall below

the classification threshold of 0.25. Notably, each of these tumor types have other lines with

scores exceeding 0.5, which should be considered as more faithful transcriptional models when

selecting lines for a study (**Supp Tab 1 and**

http://www.cahanlab.org/resources/cancerCellNet_results/).

Evaluation of patient derived xenografts

Next, we sought to evaluate a more recent class of cancer models: PDX. To do so, we

subjected the RNA-seq expression profiles of 415 PDX models from 13 different types of cancer

types generated previously[20] to CCN. Similar to the results of CCLs, the PDXs exhibited a wide

range of classification scores (**Fig 5A, Supp Tab 3**). By categorizing the CCN scores of PDX

based on the proportion of samples associated with each tumor type that were correctly

classified**,** we found that SARC, SKCM, COAD_READ and BRCA have higher proportion of

correctly classified PDX than those of other cancer categories (**Fig 5B**). In contrast to CCLs, we

found a higher proportion of correctly classified PDX in STAD, PAAD and KIRC (**Fig 5B**)**.**

However, similar to CCLs, no ESCA PDXs were classified as such. This held true when we

performed subtype classification on PDX samples: none of the PDX in ESCA were classified as

any of the ESCA subtypes (**Supp Tab 4**). UCEC PDXs had both endometrioid subtypes, serous

subtypes, and mixed subtypes, which provided a broader representation than CCLs (**Fig 5C**).

Several LUSC PDXs that were classified as a subtype were also classified as Head and Neck

squamous cell carcinoma (HNSC) or mix HNSC and LUSC (**Fig 5D**). This could be due to the

similarity in expression profiles of basal and classical subtypes of HNSC and LUSC[28,58], which is

consistent with the observation that these PDXs were also subtyped as classical. No LUSC

PDXs were classified as the secretory subtype. In contrast to LUAD CCLs, four of the five LUAD

PDXs with a discernible sub-type were classified as proximal inflammatory (**Fig 5E**). On the

other hand, similar to the CCLs, there were no TRU subtypes in the LUAD PDX cohort. In

summary, we found that while individual PDXs can reach extremely high transcriptional fidelity

to both general tumor types and subtypes, many PDXs were not classified as the general tumor

type from which they originated.

Evaluation of GEMMs

Next, we used CCN to evaluate GEMMs of six general tumor types from nine studies for

which expression data was publicly available[59–67]. As was true for CCLs and PDXs, GEMMs

also had a wide range of CCN scores (**Fig 6A, Supp Tab 5**). We next categorized the CCN

scores based on the proportion of samples associated with each tumor type that were correctly

classified (**Fig 6B**). In contrast to LGG CCLs, LGG GEMMs, generated by Nf1 mutations

expressed in different neural progenitors in combination with Pten deletion[66], consistently were

classified as LGG (**Fig 6A-B**). The GEMM dataset included multiple replicates per model, which

allowed us to examine intra-GEMM variability. Both at the level of CCN score and at the level of

categorization, GEMMs were invariant. For example, replicates of UCEC GEMMs driven by

Prg(cre/+)Pten(lox/lox) received almost identical general CCN scores (**Fig 6C, Supp Tab 6**).

GEMMs sharing genotypes across studies, such as LUAD GEMMs driven by Kras mutation and

loss of p53[59,65,67], also received similar general and subtype classification scores (**Fig 6A,B,E**).

Next, we explored the extent to which genotype impacted subtype classification in

UCEC, LUSC, and LUAD. Prg(cre/+)Pten(lox/lox) GEMMs had a mixed subtype classification of

both serous and endometrioid, consistent with the fact that Pten loss occurs in both subtypes

(albeit more frequently in endometrioid). We also analyzed Prg(cre/+)Pten(lox/lox)Csf3r-/-

GEMMs. Polymorphonuclear neutrophils (PMNs), which play anti-tumor roles in endometrioid

cancer progression, are depleted in these animals. Interestingly, Prg(cre/+)Pten(lox/lox)Csf3r-/-

GEMMs had a serous subtype classification, which could be explained by differences in PMN

involvement in endometrioid versus serous uterine tumor development that are reflected in the

respective transcriptomes of the TCGA UCEC training data. We note that the tumor cells were

sorted prior to RNA-seq and thus the shift in subtype classification is not due to contamination of

GEMMs with non-tumor components. In short, this analysis supports the argument that tumor-

cell extrinsic factors, in this case a reduction in anti-tumor PMNs, can shift the transcriptome of

a GEMM so that it more closely resembles a serous rather than endometrioid subtype.

The LUSC GEMMs that we analyzed were Lkb1$^{fl/fl}$ and they either overexpressed of

Sox2 (via two distinct mechanisms) or were also Pten$^{fl/fl}$ [65]. We note that the eight lenti-Sox2-

Cre-infected;Lkb1$^{fl/fl}$ and Rosa26LSL-Sox2-IRES-GFP;Lkb1$^{fl/fl}$ samples that classified as

'Unknown' had LUSC CCN scores only modestly lower than the decision threshold (**Fig 6D**)

(mean CCN score = 0.217). Thirteen out of the 17 of the Sox2 GEMMs classified as the

secretory subtype of LUSC. The consistency is not surprising given both models overexpress

Sox2 and lose Lkb1. On the other hand, the Lkb1$^{fl/fl}$;Pten$^{fl/fl}$ GEMMs had substantially lower

general LUSC CCN scores and our subtype classification indicated that this GEMM was mostly

classified as 'Unknown', in contrast to prior reports suggesting that it is most similar to a basal

subtype[68]. None of the three LUSC GEMMs have strong classical CCN scores. Most of the

LUAD GEMMs, which were generated using various combinations of activating Kras mutation,

loss of Trp53, and loss of Smarca4L[59,65,67], were correctly classified (**Fig 6E**). Those that were

not classified have modestly lower CCN score than the decision threshold (mean CCN score =

0.214) . There were no substantial differences in general or subtype classification across driver

genotypes. Although the sub-type of all LUAD GEMMs was 'Unknown', the subtypes tended to

have a mixture of high CCN proximal proliferation, proximal inflammation and TRU scores.

Taken together, this analysis suggests that there is a degree of similarity, and perhaps plasticity

between the primitive and secretory (but not basal or classical) subtypes of LUSC. On the other

hand, while the LUAD GEMMs classify strongly as LUAD, they do not have strong particular

subtype classification -- a result that does not vary by genotype.

Evaluation of Tumoroids

Lastly, we used CCN to assess a relatively novel cancer model: tumoroids. We

downloaded and assessed 131 distinct tumoroid expression profiles spanning 13 cancer

categories from The NCI Patient-Derived Models Repository (PDMR)[69] and from three individual

studies[70–72] (**Fig 7A, Supp Tab 7**). We note that several categories have three or fewer samples

(BRCA, CESC, KIRP, OV, LIHC, and BLCA from PDMR). Among the cancer categories

represented by more than three samples, only LUSC and PAAD have fewer than 50% classified

as their annotated label (**Fig 7B**). In contrast to GBM CCLs, all three induced pluripotent stem

cell-derived GBM tumoroids[72] were classified as GBM with high CCN scores (mean = 0.53). To

further characterize the tumoroids, we performed subtype classification on them (**Supp Tab 8**).

UCEC tumoroids from PDMR contains a wide range of subtypes with two endometrioid, two

serous and one mixed type (**Fig 7C**). On the other hand, LUSC tumoroids appear to be

predominantly of classical subtypes with one tumoroid classified as a mix between classical and

primitive (**Fig 7D**). Lastly, similar to the CCL and PDX counterparts, LUAD tumoroids are

classified as proximal inflammatory and proximal proliferation with no tumoroids classified as

TRU subtype (**Fig 7E**).

Comparison of CCLs, PDXs, GEMMs and tumoroids

Finally, we sought to estimate the comparative transcriptional fidelity of the four cancer

models modalities. We compared the general CCN scores of each model on a per tumor type

basis (**Fig 8**). In the case of GEMMs, we used the mean classification score of all samples with

shared genotypes. We also used mean classification of technical replicates found in LIHC

tumoroids[70]. We evaluated models based on both the maximum CCN score, as this represents

the potential for a model class, and the median CCN score, as this indicates the current overall

transcriptional fidelity of a model class. PDXs achieved the highest CCN scores in three (UCEC,

PAAD, LUAD) out of the five cancer categories in which all four modalities were available (**Fig**

8), despite having low median CCN scores. Notably, PDXs have a median CCN score above

the 0.25 threshold in PAAD while none of the other three modalities have any samples above

the threshold. In LIHC, the highest CCN score for PDX (0.9) is only slightly lower than the

highest CCN score for tumoroid (0.91). This suggest that certain individual PDXs most closely

mimic the transcriptional state of native patient tumors despite a portion of the PDXs having low

CCN scores. Similarly, while the majority of the CCLs have low CCN scores, several lines

achieve high transcriptional fidelity in LUSC, LUAD and LIHC (**Fig 8**). Collectively, GEMMs and

tumoroids had the highest median CCN scores in four of the five model classes (LUSC and

LUAD for GEMMs and UCEC and LIHC for tumoroids). Notably, both of the LIHC tumoroids

achieved CCN scores on par with patient tumors (**Fig 8**). In brief, this analysis indicates that

PDXs and CCLs are heterogenous in terms of transcriptional fidelity, with a portion of the

models highly mimicking native tumors and the majority of the models having low transcriptional

fidelity (with the exception of PAAD for PDXs). On the other hand, GEMMs and tumoroids

displayed a consistently high fidelity across different models.

Because the CCN score is based on a moderate number of gene features (i.e. 1,979

gene pairs consisting of 1,689 unique genes) relative to the total number of protein-coding

genes in the genome, it is possible that a cancer model with a high CCN score might not have a

high global similarity to a naturally occurring tumor. Therefore, we also calculated the GRN

status, a metric of the extent to which tumor-type specific gene regulatory network is

established[21], for all models (**Supp Fig 4**). We observed high level of correlation between the

two similarity metrics, which suggests that although CCN classifies on a selected set of genes,

its scores are highly correlated with global assessment of transcriptional similarity.

We also sought to compare model modalities in terms of the diversity of subtypes that

they represent (**Supp Fig 5**). As a reference, we also included in this analysis the overall

subtype incidence, as approximated by incidence in TCGA. Replicates in GEMMs and

tumoroids were averaged into one classification profile. In models of UCEC, there is a notable

difference in endometroid incidence, and the proportion of models classified as endometroid,

with PDX and tumoroids having any representatives (**Supp Fig 5**). All of the CCL, GEMM, and

tumoroid models of PAAD have an unknown subtype classification and no correct general

classification. However, the majority of PDXs are subtyped as either a mixture of basal and

classical, or classical alone. LUAD have proximal inflammation and proximal proliferation

subtypes modelled by CCLs and PDX (**Supp Fig 5**). Likewise, LUSC have basal, classical and

primitive subtypes modelled by CCLs and PDXs, and secretory subtype modelled by GEMMs

exclusively (**Supp Fig 5**). Taken together, these results demonstrate the need to carefully select

different model systems to more suitably model certain cancer subtypes.

DISCUSSION

A major goal in the field of cancer biology is to develop models that mimic naturally occurring

tumors with enough fidelity to enable therapeutic discoveries. However, methods to measure

the extent to which cancer models resemble or diverge from native tumors are lacking. This is

especially problematic now because there are many existing models from which to choose, and

it has become easier to generate new models. Here, we present CancerCellNet (CCN), a

computational tool that measures the similarity of cancer models to 22 naturally occurring tumor

types and 36 subtypes. While the similarity of CCLs to patient tumors has already been

explored in previous work, our tool introduces the capability to assess the transcriptional fidelity

of PDXs, GEMMs, and tumoroids. Because CCN is platform- and species-agnostic, it

represents a consistent platform to compare models across modalities including CCLs, PDXs,

GEMMs and tumoroids. Here, we applied CCN to 657 cancer cell lines, 415 patient derived

xenografts, 26 distinct genetically engineered mouse models and 131 tumoroids. Several

insights emerged from our computational analyses that have implications for the field of cancer

biology.

First, PDXs have the greatest potential to achieve transcriptional fidelity with three out of

five general tumor types for which data from all modalities was available, as indicated by the

high scores of individual PDXs. Notably PDXs are the only modality with samples classified as

PAAD. At the same time, the median CCN scores of PDXs were lower than that of GEMMs and

tumoroids in the other four tumor types. It is unclear what causes such a wide range of CCN

scores within PDXs. We suspect that some PDXs might have undergone selective pressures in

the host that distort the progression of genomic alterations away from what is observed in

natural tumor[73]. Future work to understand this heterogeneity is important so as to yield

consistently high fidelity PDXs, and to identify intrinsic and host-specific factors that so

powerfully shape the PDX transcriptome.

Second, in general GEMMs and tumoroids have higher median CCN scores than those

of PDXs and CCLs. This is also consistent with that fact that GEMMs are typically derived by

recapitulating well-defined driver mutations of natural tumors, and thus this observation

corroborates the importance of genetics in the etiology of cancer[74]. Moreover, in contrast to

most PDXs, GEMMs are typically generated in immune replete hosts. Therefore, the higher

overall fidelity of GEMMs may also be a result of the influence of a native immune system on

GEMM tumors[75]. The high median CCN scores of tumoroids can be attributed to several factors

including the increased mechanical stimuli and cell-cell interactions that come from 3D self-

organizing cultures[76,77].

Third, we have found that none of the samples that we evaluated here are

transcriptionally adequate models of ESCA. This may be due to an inherent lability of the ESCA

transcriptome that is often preceded by a metaplasia that has obscured determining its cell

type(s) of origin[78]. Therefore, this tumor type requires further attention to derive new models.

Fourth, we found that in several tumor types, GEMMs tend to reflect mixtures of

subtypes rather than conforming strongly to single subtypes. The reasons for this are not clear

but it is possible that in the cases that we examined the histologically defined subtypes have a

degree of plasticity that is exacerbated in the murine host environment.

Lastly, we recognize that many CCLs are not classified as their annotated labels. While

we have suggested that the lack of immune component is not a major confounder, we suspect

that the CCLs could undergo genetic divergence due to high number of passages,

chemotherapy before biopsy, culture condition and genetic instability[79–82], which could all be

factors that drive CCLs away from their labelled tumors.

Currently, there are several limitations to our CCN tool, and caveats to our analyses

which indicate areas for future work and improvement. First, CCN is based on transcriptomic

data but other molecular readouts of tumor state, such as profiles of the proteome[83],

epigenome[84], non-coding RNA-ome[84], and genome[74] would be equally, if not more important, to

mimic in a model system. Therefore, it is possible that some models reflect tumor behavior well,

and because this behavior is not well predicted by transcriptome alone, these models have

lower CCN scores. To both measure the extent that such situations exist, and to correct for

them, we plan in the future to incorporate other omic data into CCN so as to make more

accurate and integrated model evaluation possible. As a first step in this direction, we plan to

incorporate DNA methylation and genomic sequencing data as additional features for our

Random forest classifier as this data is becoming more readily available for both training and

cancer models. We expect that this will allow us to both refine our tumor subtype categories and

it will enable more accurate predictions of how models respond to perturbations such as drug

treatment.

A second limitation is that in the cross-species analysis, CCN implicitly assumes that

homologs are functionally equivalent. The extent to which they are not functionally equivalent

determines how confounded the CCN results will be. This possibility seems to be of limited

consequence based on the high performance of the normal tissue cross-species classifier and

based on the fact that GEMMs have the highest median CCN scores (in addition to tumoroids).

A third caveat to our analysis is that there were many fewer distinct GEMMs and

tumoroids than CCLs and PDXs. As more transcriptional profiles for GEMMs and tumoroids

emerge, this comparative analysis should be revisited to assess the generality of our results.

Finally, the TCGA training data is made up of RNA-Seq from bulk tumor samples, which

necessarily includes non-tumor cells, whereas the CCLs are by definition cell lines of tumor

origin. Therefore, CCLs theoretically could have artificially low CCN scores due to the presence

of non-tumor cells in the training data. This problem appears to be limited as we found no

correlation between tumor purity and CCN score in the CCLE samples. However, this problem

is related to the question of intra-tumor heterogeneity. We demonstrated the feasibility of using

CCN and single cell RNA-seq data to refine the evaluation of cancer cell lines contingent upon

availability of scRNA-seq training data. As more training single cell RNA-seq data accrues, CCN

would be able to not only evaluate models on a per cell type basis, but also based on cellular

composition.

We have made the results of our analyses available online so that researchers can

easily explore the performance of selected models or identify the best models for any of the 22

general tumor types and the 36 subtypes presented here. To ensure that CCN is widely

available we have developed a free web application, which performs CCN analysis on user-

uploaded data and allows for direct comparison of their data to the cancer models evaluated

here. We have also made the CCN code freely available under an Open Source license and as

an easily installed R package, and we are actively supporting its further development. Included

in the web application are instructions for training CCN and reproducing our analysis. The

documentation describes how to analyze models and compare the results to the panel of

models that we evaluated here, thereby allowing researchers to immediately compare their

models to the broader field in a comprehensive and standard fashion.

Online Methods

Training General CancerCellNet Classifier

To generate training data sets, we downloaded 8,991 patient tumor RNA-seq expression count matrix and their corresponding sample table across 22 different tumor types from TCGA using TCGAWorkflowData, TCGAbiolinks[85] and SummarizedExperiment[86] packages. We used all the patient tumor samples for training the general CCN classifier. We limited training and analysis of RNA-seq data to the 13,142 genes in common between the TCGA dataset and all the query samples (CCLs, PDXs, GEMMs, and tumoroids). To train the top pair Random forest classifier, we used a method similar to our previous method[23]. CCN first normalized the training counts matrix by down-sampling the counts to 500,000 counts per sample. To significantly reduce the execution time and memory of generating gene pairs for all possible genes, CCN then selected n up-regulated genes, n down-regulated genes and n least differentially expressed genes (CCN training parameter nTopGenes = n) for each of the 22 cancer categories using template matching[87] as the genes to generate top scoring gene pairs. In short, for each tumor type, CCN defined a template vector that labelled the training tumor samples in cancer type of interest as 1 and all other tumor samples as 0 CCN then calculated the Pearson correlation coefficient between template vector and gene expressions for all genes. The genes with strong match to template as either upregulated or downregulated had large absolute Pearson correlation coefficient. CCN chose the upregulated, downregulated and least differentially expressed genes based on the magnitude of Pearson correlation coefficient.

After CCN selected the genes for each cancer type, CCN generated gene pairs among those genes. Gene pair transformation was a method inspired by the top-scoring pair classifier[88] to allow compatibility of classifier with query expression profiles that were collected through different platforms (e.g. microarray query data applied to RNA-seq training data). In brief, the gene pair transformation compares 2 genes within an expression sample and encodes the

"gene1_gene2" gene-pair as 1 if the first gene has higher expression than the second gene.

Otherwise, gene pair transformation would encode the gene-pair as 0. Using all the gene pair

combinations generated through the gene sets per cancer type, CCN then selected top m

discriminative gene pairs (CCN training parameter nTopGenePairs = m) for each category using

template matching (with large absolute Pearson correlation coefficient) described above. To

prevent any single gene from dominating the gene pair list, we allowed each gene to appear at

maximum of three times among the gene pairs selected as features per cancer type.

After the top discriminative gene pairs were selected for each cancer category, CCN

grouped all the gene pairs together and gene pair transformed the training samples into a binary

matrix with all the discriminative gene pairs as row names and all the training samples as

column names. Using the binary gene pair matrix, CCN randomly shuffled the binary values

across rows then across columns to generate random profiles that should not resemble training

data from any of the cancer categories. CCN then sampled 70 random profiles, annotated them

as "Unknown" and used them as training data for the "Unknown" category. Using gene pair

binary training matrix, CCN constructed a multi-class Random Forest classifier of 2000 trees

and used stratified sampling of 60 sample size to ensure balance of training data in constructing

the decision trees.

To identify the best set of genes and gene-pair parameters (n and m), we used a grid-

search cross-validation[89] strategy with 5 cross-validations at each parameter set. The specific

parameters for the final CCN classifier using the function "broadClass_train" in the package

cancerCellNet are in **Supp Tab 9**. The gene-pairs are in **Supp Tab 10.**

Validating General CancerCellNet Classifier

Two thirds of patient tumor data from each cancer type were randomly sampled as

training data to construct a CCN classifier. Based on the training data, CCN selected the

classification genes and gene-pairs and trained a classifier. After the classifier was built, 35

held-out samples from each cancer category were sampled and 40 "Unknown" profiles were

generated for validation. The process of randomly sampling training set from 2/3 of all patient

tumor data, selecting features based on the training set, training classifier and validating was

repeated 50 times to have a more comprehensive assessment of the classifier trained with the

optimal parameter set. To test the performance of final CCN on independent testing data, we

applied it to 725 profiles from ICGC spanning 6 projects that do not overlap with TCGA (BRCA-

KR, LIRI-JP, OV-AU, PACA-AU, PACA-CA, PRAD-FR).

Selecting Decision Thresholds

Our strategy for selecting a decision threshold was to find the value that maximizes the

average Macro F1 measure[90] for each of the 50 cross-validations that were performed with the

optimal parameter set, testing thresholds between 0 and 1 with a 0.01 increment. The F1

measure is defined as:

643
$$Macro\ F1 = \frac{2 \times precision \times recall}{precision + recall}$$

We selected the most commonly occurring threshold above 0.2 that maximized the average

Macro F1 measure across the 50 cross-validations as the decision threshold for the final

classifier (threshold = 0.25). The same approach was applied for the subtype classifiers. The

thresholds and the corresponding average precision, recall and F1 measures are recorded in

(**Supp Tab 11**).

Classifying Query Data into General Cancer Categories

We downloaded the RNA-seq cancer cell lines expression profiles and sample table

from (https://portals.broadinstitute.org/ccle/data), and microarray cancer cell lines expression

profiles and sample table from Barretina et al [37]. We extracted two WT control NCCIT RNA-seq

expression profiles from Grow et al[91]. We received PDX expression estimates and sample

annotations from the authors of Gao et al [20]. We gathered GEMM expression profiles from nine

different studies[59–67]. We downloaded tumoroid expression profiles from The NCI Patient-

Derived Models Repository (PDMR)[69] and from three individual studies[70–72]. To use CCN

classifier on GEMM data, the mouse genes from GEMM expression profiles were converted into

their human homologs. The query samples were classified using the final CCN classifier. Each

query classification profile was labelled as one of the four classification categories: "correct",

"mixed", "none" and "other" based on classification profiles. If a sample has a CCN score higher

than the decision threshold in the labelled cancer category, we assigned that as "correct". If a

sample has CCN score higher than the decision threshold in labelled cancer category and in

other cancer categories, we assigned that as "mixed". If a sample has no CCN score higher

than the decision threshold in any cancer category or has the highest CCN score in 'Unknown'

category, then we assigned it as "none". If a sample has CCN score higher than the decision

threshold in a cancer category or categories not including the labelled cancer category, we

assigned it as "other". We analyzed and visualized the results using R and R packages

pheatmap[92] and ggplot2[93].

Cross-Species Assessment

To assess the performance of cross-species classification, we downloaded 1003

labelled human tissue/cell type and 1993 labelled mouse tissue/cell type RNA-seq expression

profiles from Github (https://github.com/pcahan1/CellNet). We first converted the mouse genes

into human homologous genes. Then we found the intersecting genes between mouse

tissue/cell expression profiles and human tissue/cell expression profiles. Limiting the input of

human tissue RNA-seq profiles to the intersecting genes, we trained a CCN classifier with all

the human tissue/cell expression profiles. The parameters used for the function

"broadClass_train" in the package cancerCellNet are in **Supp Tab 9.** We randomly sampled 75

samples from each tissue category in mouse tissue/cell data and applied the classifier on those

samples to assess performance.

Cross-Technology Assessment

To assess the performance of CCN in applications to microarray data, we gathered

6,219 patient tumor microarray profiles across 12 different cancer types from more than 100

different projects (**Supp Tab 12**). We found the intersecting genes between the microarray

profiles and TCGA patient RNA-seq profiles. Limiting the input of RNA-seq profiles to the

intersecting genes, we created a CCN classifier with all the TCGA patient profiles using

parameters for the function "broadClass_train" listed in **Supp Tab 9**. After the microarray

specific classifier was trained, we randomly sampled 60 microarray patient samples from each

cancer category and applied CCN classifier on them as assessment of the cross-technology

performance in **Supp Fig 2A**. The same CCN classifier was used to assess microarray CCL

samples **Supp Fig 2B**.

Training and validating scRNA-seq Classifier

We extracted labelled human melanoma and glioblastoma scRNA-seq expression

profiles[40,41], and compiled the two datasets excluding 3 cell types T.CD4, T.CD8 and Myeloid

due to low number of cells for training. 60 cells from each of the 11 cell types were sampled for

training a scRNA-seq classifier. The parameters for training a general scRNA-seq classifier

using the function "broadClass_train" are in **Supp Tab 9.** 25 cells from each of the 11 cell types

from the held-out data were selected to assess the single cell classifier**.** Using maximization of

average Macro F1 measure, we selected the decision threshold of 0.255. The gene-pairs that

were selected to construct the classifier are in **Supp Tab 10**. To assess the cross-technology

capability of applying scRNA-seq classifier to bulk RNA-seq, we downloaded 305 expression

profiles spanning 4 purified cell types (B cells, endothelial cells, monocyte/macrophage,

fibroblast) from https://github.com/pcahan1/CellNet.

Training Subtype CancerCellNet

We found 11 cancer types (BRCA, COAD, ESCA, HNSC, KIRC, LGG, PAAD, UCEC,

STAD, LUAD, LUSC) which have meaningful subtypes based on either histology or molecular

profile and have sufficient samples to train a subtype classifier with high AUPR. We also

included normal tissues samples from BRCA, COAD, HNSC, KIRC, UCEC to create a normal

tissue category in the construction of their subtype classifiers. Training samples were either

labelled as a cancer subtype for the cancer of interest or as "Unknown" if they belong to other

cancer types. Similar to general classifier training, CCN performed gene pair transformation and

selected the most discriminate gene pairs for each cancer subtype. In addition to the gene pairs

selected to discriminate cancer subtypes, CCN also performed general classification of all

training data and appended the classification profiles of training data with gene pair binary

matrix as additional features. The reason behind using general classification profile as additional

features is that many general cancer types may share similar subtypes, and general

classification profile could be important features to discriminate the general cancer type of

interest from other cancer types before performing finer subtype classification. The specific

parameters used to train individual subtype classifiers using "subClass_train" function of

CancerCellNet package can be found in **Supp Tab 9** and the gene pairs are in **Supp Tab 10**.

Validating Subtype CancerCellNet

Similar to validating general class classifier, we randomly sampled 2/3 of all samples in

each cancer subtype as training data and sampled an equal amount across subtypes in the 1/3

held-out data for assessing subtype classifiers. We repeated the process 20 times for more

comprehensive assessment of subtype classifiers.

Classifying Query Data into Subtypes

We assigned subtype to query sample if the query sample has CCN score higher than the decision threshold. The table of decision threshold for subtype classifiers are in **Supp Tab 11**. If no CCN scores exceed the decision threshold in any subtype or if the highest CCN score is in 'Unknown' category, then we assigned that sample as 'Unknown'. Analysis was performed in R and visualizations were generated with the ComplexHeatmap package[94].

Cells culture, Immunohistochemistry and histomorphometry

Caov-4 (ATCC® HTB-76™), SK-OV-3(ATCC® HTB-77™), RT4 (ATCC® HTB-2™), and NCCIT(ATCC® CRL-2073™) cell lines were purchased from ATCC. HEC-59 (C0026001) and A2780 (93112519-1VL) were obtained from Addexbio Technologies and Sigma-Aldrich. Vcap and PC-3. SK-OV-3, Vcap, and RT4 were cultured in Dulbecco's Modified Eagle Medium (DMEM, high glucose, 11960069, Gibco) with 1% Penicillin-Streptomycin-Glutamine (10378016, Life Technologies); Caov-4, PC-3, NCCIT, and A2780 were cultured using RPMI-1640 medium (11875093, Gibco) while HEC-59 was in Iscove's Modified Dulbecco's Medium (IMDM, 12440053, Gibco). Both media were supplemented with 1% Penicillin-Streptomycin (15140122, Gibco). All medium included 10% Fetal Bovine Serum (FBS).

Cells cultured in 48-well plate were washed twice with PBS and fixed in 10% buffered formalin for 24 hrs at 4 °C. Immunostaining was performed using a standard protocol. Cells were incubated with primary antibodies to goat HOXB6 (10 µg/mL, PA5-37867, Invitrogen), mouse WT1(10 µg/mL, MA1-46028, Invitrogen), rabbit PPARG (1:50, ABN1445, Millipore), mouse FOLH1(10 µg/mL, UM570025, Origene), and rabbit LIN28A (1:50, #3978, Cell Signaling) in Antibody Diluent (S080981-2, DAKO), at 4 °C overnight followed with three 5 min washes in TBST. The slides were then incubated with secondary antibodies conjugated with fluorescence at room temperature for 1 h while avoiding light followed with three 5 min washes in TBST and

nuclear stained with mounting medium containing DAPI. Images were captured by Nikon

EcLipse Ti-S, DS-U3 and DS-Qi2.

Histomorphometry was performed using ImageJ (Version 2.0.0-rc-69/1.52i). %

N.positive cells was calculated by the percentage of the number of positive stained cells divided

by the number of DAPI-positive nucleus within three of randomly chosen areas. The data were

expressed as means ± SD.

Tumor Purity Analysis

We used the R package ESTIMATE[95] to calculate the ESTIMATE scores from TCGA

tumor expression profiles that we used as training data for CCN classifier. To calculate tumor

purity we used the equation described in YoshiHara et al., 2013[95]:

$$\text{Tumour purity} = \cos (0.6049872018 + 0.0001467884 \times \text{ESTIMATE score})$$

Extracting Citation Counts

We used the R package RISmed[96] to extract the number of citations for each cell line

through query search of "*cell line name*[Text Word] AND *cancer*[Text Word]" on PubMed. The

citation counts were normalized by dividing the citation counts with the number of years since

first documented.

$$Normalized\ citation\ counts = \frac{citation\ counts}{\#\ years\ since\ first\ documented}$$

GRN construction and GRN Status

GRN construction was extended from our previous method[21]. 80 samples per cancer

type were randomly sampled and normalized through down sampling as training data for the

CLR GRN construction algorithm. Cancer type specific GRNs were identified by determining the

differentially expressed genes per each cancer type and extracting the subnetwork using those

genes.

To extend the original GRN status algorithm[21] across different platforms and species, we

devised a rank-based GRN status algorithm. Like the original GRN status, rank based GRN

status is a metric of assessing the similarity of cancer type specific GRN between training data

in the cancer type of interest and query samples. Hence, high GRN status represents high level

of establishment or similarity of the cancer specific GRN in the query sample compared to those

of the training data. The expression profiles of training data and query data were transformed

into rank expression profiles by replacing the expression values with the rank of the expression

values within a sample (highest expressed gene would have the highest rank and lowest

expressed genes would have a rank of 1). Cancer type specific mean and standard deviation of

every gene's rank expression were learned from training data. The modified Z-score values for

genes within cancer type specific GRN were calculated for query sample's rank expression

profiles to quantify how dissimilar the expression values of genes in query sample's cancer type

specific GRN compared to those of the reference training data:

$$Zscore(gene\ i)_{mod} = \begin{cases} 0,\ if\ Zscore\ is\ positive\ and\ the\ gene\ is\ found\ to\ be\ upregulated \\ 0,\ if\ Zscore\ is\ negative\ and\ the\ gene\ is\ found\ to\ be\ downregulated \\ \qquad\qquad abs(Zscore),\ otherwise \end{cases}$$

If a gene in the cancer type specific GRN is found to be upregulated in the specific

cancer type relative to other cancer types, then we would consider query sample's gene to be

similar if the ranking of the query sample's gene is equal to or greater than the mean ranking of

the gene in training sample. As a result of similarity, we assign that gene of a Z-score of 0. The

same principle applies to cases where the gene is downregulated in cancer specific subnetwork.

GRN status for query sample is calculated as the weighted mean of the

$(1000 - Zscore(gene\ i)_{mod})$ across genes in cancer type specific GRN. 1000 is an arbitrary

large number, and larger dissimilarity between query's cancer type specific GRN indicate high

Z-scores for the GRN genes and low GRN status.

$$RGS = \sum_{i=1}^{n} (1000 - Zscore(gene\ i)_{mod})weight_{gene\ i}$$

$$GRN\ Status = \frac{RGS}{\sum_{i=1}^{n} weight_{gene\ i}}$$

The weight of individual genes in the cancer specific network is determined by the

importance of the gene in the Random Forest classifier. Finally, the GRN status gets normalized

with respect to the GRN status of the cancer type of interest and the cancer type with the lowest

mean GRN status.

$$Normalized\ GRN\ status = \frac{GRN\ status_{query} - avg(GRN\ status_{min\ cancer})}{avg(GRN\ status_{cancer\ type\ interest})}$$

Where "min cancer" represents the cancer type where its training data have the lowest

mean GRN status in the cancer type of interest, and $avg(GRN\ status_{min\ cancer})$ represents the

lowest average GRN status in the cancer type of interest. $avg(GRN\ status_{cancer\ type\ interest})$

represents average GRN status of the cancer type of interest in the training data.

Code availability

CancerCellNet code and documentation is available at GitHub:

https://github.com/pcahan1/cancerCellNet

Acknowledgements

This work was supported by the National Institutes of Health NCI Ovarian Cancer SPORE

P50CA228991 via a Development Research Program award to PC. FWH was supported by a

Prostate Cancer Foundation Young Investigator Award, Department of Defense W81XWH-17-

PCRP-HD (F.W.H.), the National Institutes of Health/National Cancer Institute P20 CA233255-

01 (F.W.H.) U19 CA214253 (F.W.H.). We would like to thank John Powers, Hao Zhu, Tian-Li

Wang, Charles Eberhart, and Kaloyan Tsanov for comments on the manuscript and helpful

discussions. Some figures were created in part with Biorender.com.

FIGURE LEGENDS

Fig. 1 CancerCellNet (CCN) workflow, training, and performance. (A) Schematic of CCN

usage. CCN was designed to assess and compare the expression profiles of cancer models

such as CCLs, PDXs, GEMMs, and tumoroids with native patient tumors. To use trained

classifier, CCN inputs the query samples (e.g. expression profiles from CCLs, PDXs, GEMMs,

tumoroids) and generates a classification profile for the query samples. The column names of

the classification heatmap represent sample annotation and the row names of the classification

heatmap represent different cancer types. Each grid is colored from black to yellow representing

the lowest classification score (e.g. 0) to highest classification score (e.g. 1). **(B)** Schematic of

CCN training process. CCN uses patient tumor expression profiles of 22 different cancer types

from TCGA as training data. First, CCN identifies n genes that are upregulated, n that are

downregulated, and n that are relatively invariant in each tumor type versus all of the others.

Then, CCN performs a pair transform on these genes and subsequently selects the most

discriminative set of m gene pairs for each cancer type as features (or predictors) for the

Random forest classifier. Lastly, CCN trains a multi-class Random Forest classifier using gene-

pair transformed training data. **(C)** Parameter optimization strategy. 5 cross-validations of each

parameter set in which 2/3 of TCGA data was used to train and 1/3 to validate was used search

for the values of n and m that maximized performance of the classifier as measured by area

under the precision recall curve (AUPRC). **(D)** Mean and standard deviation of classifiers based

on 50 cross-validations with the optimal parameter set. **(E)** AUPRC of the final CCN classifier

when applied to independent patient tumor data from ICGC.

Fig. 2 Evaluation of cancer cell lines. (A) General classification heatmap of CCLs extracted

from CCLE. Column annotations of the heatmap represent the labelled cancer category of the

CCLs given by CCLE and the row names of the heatmap represent different cancer categories.

CCLs' general classification profiles are categorized into 4 categories: correct (red), correct

mixed (pink), no classification (light green) and other classification (dark green) based on the

decision threshold of 0.25. **(B)** Bar plot represents the proportion of each classification category

in CCLs across cancer types ordered from the cancer types with the highest proportion of

correct and correct mixed CCLs to lowest proportion. **(C)** Comparison between SKCM general

CCN scores from bulk RNA-seq classifier and SKCM malignant CCN scores from scRNA-seq

classifier for SKCM CCLs. **(D)** Comparison between SARC general CCN scores from bulk RNA-

seq classifier and CAF CCN scores from scRNA-seq classifier for SKCM CCLs. **(E)** Comparison

between GBM general CCN scores from bulk RNA-seq classifier and GBM neoplastic CCN

scores from scRNA-seq classifier for GBM CCLs. **(F)** Comparison between SARC general CCN

scores and CAF CCN scores from scRNA-seq classifier for GBM CCLs. The green lines

indicate the decision threshold for scRNA-seq classifier and general classifier.

Fig. 3 Immunofluorescence of selected cell lines. (A) Classification profiles (left) and IF

expression (middle) of Caov-4 (OV positive control), HEC-59 (UCEC positive control) and SK-

OV-3 for WT1 (OV biomarker) and HOXB6 (uterine biomarker). The bar plots quantify the

average percentage of positive cells for WT1 (top-right) and HOXB6 (bottom-right). **(B)**

Classification profiles (left) and IF expression (middle) of Caov-4, NCCIT (germ cell tumor

positive control) and A2780 for WT1 and LIN28A (germ cell tumor biomarker). Classification of

NCCIT were performed using RNA-seq profiles of WT control NCCIT duplicate from Grow et

al[91]. The bar plots quantify the average percentage of positive cells for WT1 (top-right) and

LIN28A (bottom-right). **(C)** Classification profiles (left) and IF expression (middle) of Vcap

(PRAD positive control), RT4 (BLCA positive control) and PC-3 for FOLH1 (prostate biomarker)

and PPARG (urothelial biomarker). The bar plots quantify the average percentage of positive

cells for FOLH1 (top-right) and PPARG (bottom-right).

Fig. 4 Subtype classification of CCLs and CCL prevalence. The heatmap visualizations

represent subtype classification of **(A)** UCEC CCLs**, (B)** LUSC CCLs and **(C)** LUAD CCLs. Only

samples with CCN scores > 0.1 in their nominal tumor type are displayed. **(D)** Comparison of

normalized citation counts and general CCN classification scores of CCLs. Labelled cell lines

either have the highest CCN classification score in their labelled cancer category or highest

normalized citation count. Each citation count was normalized by number of years since first

documented on PubMed.

Fig. 5 Evaluation of patient derived xenografts. (A) General classification heatmap of PDXs.

Column annotations represent annotated cancer type of the PDXs, and row names represent

cancer categories. **(B)** Proportion of classification categories in PDXs across cancer types is

visualized in the bar plot and ordered from the cancer type with highest proportion of correct and

mixed correct classified PDXs to the lowest. Subtype classification heatmaps of **(C)** UCEC

PDXs, **(D)** LUSC PDXs and **(E)** LUAD PDXs. Only samples with CCN scores > 0.1 in their

nominal tumor type are displayed.

Fig. 6 Evaluation of genetically engineered mouse models. (A) General classification

heatmap of GEMMs. Column annotations represent annotated cancer type of the GEMMs, and

row names represent cancer categories. **(B)** Proportion of classification categories in GEMMs

across cancer types is visualized in the bar plot and ordered from the cancer type with highest

proportion of correct and mixed correct classified GEMMs to the lowest. Subtype classification

heatmap of **(C)** UCEC GEMMs, **(D)** LUSC GEMMs and **(E)** LUAD GEMMs. Only samples with

CCN scores > 0.1 in their nominal tumor type are displayed.

Fig. 7 Evaluation of tumoroid models. (A) General classification heatmap of tumoroids.

Column annotations represent annotated cancer type of the tumoroids, and row names

represent cancer categories. **(B)** Proportion of classification categories in tumoroids across

cancer types is visualized in the bar plot and ordered from the cancer type with highest

proportion of correct and mixed correct classified tumoroids to the lowest. Subtype classification

heatmap of **(C)** UCEC tumoroids, **(D)** LUSC tumoroids and **(E)** LUAD tumoroids. Only samples

with CCN scores > 0.1 in their nominal tumor type are displayed.

Fig. 8 Comparison of CCLs, PDXs, and GEMMs. Box-and-whiskers plot comparing general

CCN scores across CCLs, GEMMs, PDXs of five general tumor types (UCEC, PAAD, LUSC,

LUAD, LIHC).

Supplementary Information

Supplementary Figure 1 Assessment of CCN general classifier and subtype classifier. **(A)**

Mean AUPRC of repeated grid-search cross-validation for each parameter grid. **(B)** Mean and

range of CCN classifier's PR curves from 50 cross validations based on the optimal feature

selection parameters n and m. **(C)** AUPRC of CCN human tissue classifier when applied to

mouse tissue data. **(D)** The schematic of training a subtype classifier in CCN. CCN uses patient

tumor expression profiles from cancer of interest as training data. CCN performs gene-pair

transformation and selects the most discriminative gene pairs among the cancer subtypes from

training data as features. CCN then applies the general classification on training data and uses

the general classification profile as features in addition to gene pairs for training a Random

Forest classifier. The weight of the general classification profiles as features can be tuned to

improve AUPRC. **(E)** The mean and standard deviation of AUPRC for 11 subtype classifiers

based on 20 iterations of random sampling of training and held-out data, training subtype

classifier using training data, classification of held-out data, and calculation of recall and

precision.

Supplementary Figure 2 Further validation of CCN and classification results. To validate the

cross-platform classification performance of CCN, a new classifier specifically trained to classify

microarray data was trained using RNA-seq data from TCGA as training data and intersecting

genes between RNA-seq data and microarray data. **(A)** AUPRC of CCN classifier when applied

to tumor profiles assayed on microarrays. **(B)** Classification heatmap of CCLs using microarray

expression data. **(C)** Pearson correlation between CCN scores of CCLE lines generated from

RNA-seq data and microarray data. **(D)** Comparison between CCLs' CCN scores and the

similarity metric from Yu et al[15], median correlations of transcriptional profiles between CCLs

and TCGA tumors from CCLs' labelled cancer category. **(E)** Comparison of mean tumor purity

of training data and mean CCN scores of CCLs for each cancer category.

Supplementary Figure 3 Single-cell classification of SKCM and GBM cell lines. **(A)** AUPRC of

the single-cell classifier when applied to scRNA-seq held-out data. **(B)** AUPRC of the scRNA-

seq classifier when applied to purified bulk RNA samples. **(C)** Single-cell classification of SKCM

CCLs. Red bar-plot (top) represents general CCN scores in SARC and blue bar-plot (bottom)

represents general CCN scores in SKCM. **(D)** Single-cell classification of GBM CCLs. Red bar-

plot (top) represents general CCN scores in SARC and yellow bar-plot (bottom) represents

general CCN scores in GBM.

Supplementary Figure 4 Correlation between cancer type specific network GRN status and
general CCN scores.
954

956　**Supplementary Figure 5** Proportion of cancer subtypes in different cancer models and TCGA
957　tumor data across 11 general cancer types.
958

Supplementary Table 1 General classification profiles of CCLs.

Supplementary Table 2 Subtype classification profiles of CCLs.

Supplementary Table 3 General classification profiles of PDXs.

Supplementary Table 4 Subtype classification profiles of PDXs.

Supplementary Table 5 General classification profiles of GEMMs

Supplementary Table 6 Subtype classification profiles of GEMMs.

Supplementary Table 7 General classification profiles of tumoroids.

Supplementary Table 8 Subtype classification profiles of tumoroids.

Supplementary Table 9 Specific parameters used for training of all classifiers.

Supplementary Table 10 Gene-pairs selected for final training of CCN general, subtype
classifiers and single-cell classifier.

Supplementary Table 11 Decision thresholds and the corresponding precision and recall for
the general classifier and subtype classifier.

Supplementary Table 12 Accessions of tumor microarray data used in validation.

REFERENCES

1. Sharma, S. V., Haber, D. A. & Settleman, J. Cell line-based platforms to evaluate
the therapeutic efficacy of candidate anticancer agents. *Nat. Rev. Cancer* **10,** 241–
253 (2010).
2. Kersten, K., de Visser, K. E., van Miltenburg, M. H. & Jonkers, J. Genetically
engineered mouse models in oncology research and cancer medicine. *EMBO Mol.*
Med. **9,** 137–153 (2017).
3. Hidalgo, M. *et al.* Patient-derived xenograft models: an emerging platform for
translational cancer research. *Cancer Discov.* **4,** 998–1013 (2014).
4. Drost, J. & Clevers, H. Organoids in cancer research. *Nat. Rev. Cancer* **18,** 407–
418 (2018).
5. Klijn, C. *et al.* A comprehensive transcriptional portrait of human cancer cell lines.
Nat. Biotechnol. **33,** 306–312 (2015).
6. Koren, S. *et al.* PIK3CA(H1047R) induces multipotency and multi-lineage mammary
tumours. *Nature* **525,** 114–118 (2015).
7. DeRose, Y. S. *et al.* Tumor grafts derived from women with breast cancer
authentically reflect tumor pathology, growth, metastasis and disease outcomes.
Nat. Med. **17,** 1514–1520 (2011).

8.   Sharpless, N. E. & Depinho, R. A. The mighty mouse: genetically engineered
mouse models in cancer drug development. *Nat. Rev. Drug Discov.* **5,** 741–754
(2006).
9.   Mouradov, D. *et al.* Colorectal cancer cell lines are representative models of the
main molecular subtypes of primary cancer. *Cancer Res.* **74,** 3238–3247 (2014).
10. Stuckelberger, S. & Drapkin, R. Precious GEMMs: emergence of faithful models for
ovarian cancer research. *J. Pathol.* **245,** 129–131 (2018).
11. Domcke, S., Sinha, R., Levine, D. A., Sander, C. & Schultz, N. Evaluating cell lines
as tumour models by comparison of genomic profiles. *Nat. Commun.* **4,** 2126
(2013).
12. Jiang, G. *et al.* Comprehensive comparison of molecular portraits between cell lines
and tumors in breast cancer. *BMC Genomics* **17 Suppl 7,** 525 (2016).
13. Chen, B., Sirota, M., Fan-Minogue, H., Hadley, D. & Butte, A. J. Relating
hepatocellular carcinoma tumor samples and cell lines using gene expression data
in translational research. *BMC Med. Genomics* **8 Suppl 2,** S5 (2015).
14. Vincent, K. M., Findlay, S. D. & Postovit, L. M. Assessing breast cancer cell lines as
tumour models by comparison of mRNA expression profiles. *Breast Cancer Res.*
17, 114 (2015).
15. Yu, K. *et al.* Comprehensive transcriptomic analysis of cell lines as models of
primary tumors across 22 tumor types. *Nat. Commun.* **10,** 3574 (2019).
16. Najgebauer, H. *et al.* CELLector: Genomics-Guided Selection of Cancer In Vitro
Models. *Cell Syst.* **10,** 424–432.e6 (2020).
17. Salvadores, M., Fuster-Tormo, F. & Supek, F. Matching cell lines with cancer type
and subtype of origin via mutational, epigenomic, and transcriptomic patterns. *Sci.*
Adv. **6,** (2020).
18. Guernet, A. & Grumolato, L. CRISPR/Cas9 editing of the genome for cancer
modeling. *Methods* **121-122,** 130–137 (2017).
19. Gargiulo, G. Next-Generation in vivo Modeling of Human Cancers. *Front. Oncol.* **8,**
429 (2018).
20. Gao, H. *et al.* High-throughput screening using patient-derived tumor xenografts to
predict clinical trial drug response. *Nat. Med.* **21,** 1318–1325 (2015).
21. Cahan, P. *et al.* CellNet: network biology applied to stem cell engineering. *Cell* **158,**
903–915 (2014).
22. Radley, A. H. *et al.* Assessment of engineered cells using CellNet and RNA-seq.
Nat. Protoc. **12,** 1089–1102 (2017).
23. Tan, Y. & Cahan, P. SingleCellNet: A Computational Tool to Classify Single Cell
RNA-Seq Data Across Platforms and Across Species. *Cell Syst.* **9,** 207–213.e2
(2019).
24. Cancer Genome Atlas Network. Comprehensive molecular characterization of
human colon and rectal cancer. *Nature* **487,** 330–337 (2012).
25. Zhang, J. *et al.* International Cancer Genome Consortium Data Portal--a one-stop
shop for cancer genomics data. *Database (Oxford)* **2011,** bar026 (2011).
26. Cancer Genome Atlas Network. Comprehensive molecular portraits of human
breast tumours. *Nature* **490,** 61–70 (2012).
27. Parker, J. S. *et al.* Supervised risk predictor of breast cancer based on intrinsic
subtypes. *J. Clin. Oncol.* **27,** 1160–1167 (2009).

28. Wilkerson, M. D. *et al.* Lung squamous cell carcinoma mRNA expression subtypes
are reproducible, clinically important, and correspond to normal cell types. *Clin.*
Cancer Res. **16,** 4864–4875 (2010).
29. Cancer Genome Atlas Research Network. Electronic address:
andrew_aguirre@dfci.harvard.edu & Cancer Genome Atlas Research Network.
Integrated genomic characterization of pancreatic ductal adenocarcinoma. *Cancer*
Cell **32,** 185–203.e13 (2017).
30. Cancer Genome Atlas Research Network *et al.* Integrated genomic characterization
of endometrial carcinoma. *Nature* **497,** 67–73 (2013).
31. Cancer Genome Atlas Research Network *et al.* Integrated genomic characterization
of oesophageal carcinoma. *Nature* **541,** 169–175 (2017).
32. Cancer Genome Atlas Network. Comprehensive genomic characterization of head
and neck squamous cell carcinomas. *Nature* **517,** 576–582 (2015).
33. Cancer Genome Atlas Research Network. Comprehensive molecular
characterization of clear cell renal cell carcinoma. *Nature* **499,** 43–49 (2013).
34. Verhaak, R. G. W. *et al.* Integrated genomic analysis identifies clinically relevant
subtypes of glioblastoma characterized by abnormalities in PDGFRA, IDH1, EGFR,
and NF1. *Cancer Cell* **17,** 98–110 (2010).
35. Cancer Genome Atlas Research Network. Comprehensive molecular profiling of
lung adenocarcinoma. *Nature* **511,** 543–550 (2014).
36. Hu, B. *et al.* Gastric cancer: Classification, histology and application of molecular
pathology. *J. Gastrointest. Oncol.* **3,** 251–261 (2012).
37. Barretina, J. *et al.* The Cancer Cell Line Encyclopedia enables predictive modelling
of anticancer drug sensitivity. *Nature* **483,** 603–607 (2012).
38. Medico, E. *et al.* The molecular landscape of colorectal cancer cell lines unveils
clinically actionable kinase targets. *Nat. Commun.* **6,** 7002 (2015).
39. Park, J.-G. *et al.* Characteristics of Cell Lines Established from Human Colorectal
Carcinoma. *Cancer Res.* (1987).
40. Jerby-Arnon, L. *et al.* A cancer cell program promotes T cell exclusion and
resistance to checkpoint blockade. *Cell* **175,** 984–997.e24 (2018).
41. Darmanis, S. *et al.* Single-Cell RNA-Seq Analysis of Infiltrating Neoplastic Cells at
the Migrating Front of Human Glioblastoma. *Cell Rep.* **21,** 1399–1410 (2017).
42. Patel, A. P. *et al.* Single-cell RNA-seq highlights intratumoral heterogeneity in
primary glioblastoma. *Science* **344,** 1396–1401 (2014).
43. Xu, B. *et al.* Regulation of endometrial receptivity by the highly expressed HOXA9,
HOXA11 and HOXD10 HOX-class homeobox genes. *Hum. Reprod.* **29,** 781–790
(2014).
44. Raines, A. M. *et al.* Recombineering-based dissection of flanking and paralogous
Hox gene functions in mouse reproductive tracts. *Development* **140,** 2942–2952
(2013).
45. Netinatsunthorn, W., Hanprasertpong, J., Dechsukhum, C., Leetanaporn, R. &
Geater, A. WT1 gene expression as a prognostic marker in advanced serous
epithelial ovarian carcinoma: an immunohistochemical study. *BMC Cancer* **6,** 90
(2006).
46. Kelly, Z. *et al.* The prognostic significance of specific HOX gene expression patterns
in ovarian cancer. *Int. J. Cancer* **139,** 1608–1617 (2016).

47. Cancer Genome Atlas Research Network. Integrated genomic analyses of ovarian
carcinoma. *Nature* **474,** 609–615 (2011).
48. Wiegand, K. C. *et al.* ARID1A mutations in endometriosis-associated ovarian
carcinomas. *N. Engl. J. Med.* **363,** 1532–1543 (2010).
49. Murray, M. J. *et al.* LIN28 Expression in malignant germ cell tumors downregulates
let-7 and increases oncogene levels. *Cancer Res.* **73,** 4872–4884 (2013).
50. Biton, A. *et al.* Independent component analysis uncovers the landscape of the
bladder tumor transcriptome and reveals insights into luminal and basal subtypes.
Cell Rep. **9,** 1235–1245 (2014).
51. Fair, W. R., Israeli, R. S. & Heston, W. D. Prostate-specific membrane antigen.
Prostate **32,** 140–148 (1997).
52. Black, J. D., English, D. P., Roque, D. M. & Santin, A. D. Targeted therapy in
uterine serous carcinoma: an aggressive variant of endometrial cancer. *Womens*
Health (Lond. Engl.) **10,** 45–57 (2014).
53. Yang, S., Thiel, K. W. & Leslie, K. K. Progesterone: the ultimate endometrial tumor
suppressor. *Trends Endocrinol. Metab.* **22,** 145–152 (2011).
54. Huszar, M. *et al.* Up-regulation of L1CAM is linked to loss of hormone receptors and
E-cadherin in aggressive subtypes of endometrial carcinomas. *J. Pathol.* **220,** 551–
561 (2010).
55. Kozak, J., Wdowiak, P., Maciejewski, R. & Torres, A. A guide for endometrial
cancer cell lines functional assays using the measurements of electronic
impedance. *Cytotechnology* **70,** 339–350 (2018).
56. Korch, C. *et al.* DNA profiling analysis of endometrial and ovarian cell lines reveals
misidentification, redundancy and contamination. *Gynecol. Oncol.* **127,** 241–248
(2012).
57. Wu, D. *et al.* Gene-expression data integration to squamous cell lung cancer
subtypes reveals drug sensitivity. *Br. J. Cancer* **109,** 1599–1608 (2013).
58. Walter, V. *et al.* Molecular subtypes in head and neck cancer exhibit distinct
patterns of chromosomal gain and loss of canonical cancer genes. *PLoS One* **8,**
e56823 (2013).
59. Adeegbe, D. O. *et al.* BET Bromodomain Inhibition Cooperates with PD-1 Blockade
to Facilitate Antitumor Response in Kras-Mutant Non-Small Cell Lung Cancer.
Cancer Immunol Res **6,** 1234–1245 (2018).
60. Blaisdell, A. *et al.* Neutrophils oppose uterine epithelial carcinogenesis via
debridement of hypoxic tumor cells. *Cancer Cell* **28,** 785–799 (2015).
61. Fitamant, J. *et al.* YAP inhibition restores hepatocyte differentiation in advanced
HCC, leading to tumor regression. *Cell Rep.* **10,** 1692–1707 (2015).
62. Jia, D. *et al.* Crebbp loss drives small cell lung cancer and increases sensitivity to
HDAC inhibition. *Cancer Discov.* **8,** 1422–1437 (2018).
63. Kress, T. R. *et al.* Identification of MYC-Dependent Transcriptional Programs in
Oncogene-Addicted Liver Tumors. *Cancer Res.* **76,** 3463–3472 (2016).
64. Li, L. *et al.* GKAP acts as a genetic modulator of NMDAR signaling to govern
invasive tumor growth. *Cancer Cell* **33,** 736–751.e5 (2018).
65. Mollaoglu, G. *et al.* The Lineage-Defining Transcription Factors SOX2 and NKX2-1
Determine Lung Cancer Cell Fate and Shape the Tumor Immune
Microenvironment. *Immunity* **49,** 764–779.e9 (2018).

66. Pan, Y. *et al.* Whole tumor RNA-sequencing and deconvolution reveal a clinically-prognostic PTEN/PI3K-regulated glioma transcriptional signature. *Oncotarget* **8,** 52474–52487 (2017).

67. Lissanu Deribe, Y. *et al.* Mutations in the SWI/SNF complex induce a targetable dependence on oxidative phosphorylation in lung cancer. *Nat. Med.* **24,** 1047–1057 (2018).

68. Xu, C. *et al.* Loss of Lkb1 and Pten leads to lung squamous cell carcinoma with elevated PD-L1 expression. *Cancer Cell* **25,** 590–604 (2014).

69. NCI-Frederick, Frederick, MD. National Laboratory for Cancer Research. The NCI Patient-Derived Models Repository (PDMR). (2019). at <https://pdmr.cancer.gov/>

70. Broutier, L. *et al.* Human primary liver cancer-derived organoid cultures for disease modeling and drug screening. *Nat. Med.* **23,** 1424–1435 (2017).

71. Lee, S. H. *et al.* Tumor Evolution and Drug Response in Patient-Derived Organoid Models of Bladder Cancer. *Cell* **173,** 515–528.e17 (2018).

72. Ogawa, J., Pao, G. M., Shokhirev, M. N. & Verma, I. M. Glioblastoma model using human cerebral organoids. *Cell Rep.* **23,** 1220–1229 (2018).

73. Ben-David, U. *et al.* Patient-derived xenografts undergo mouse-specific tumor evolution. *Nat. Genet.* **49,** 1567–1575 (2017).

74. Stratton, M. R., Campbell, P. J. & Futreal, P. A. The cancer genome. *Nature* **458,** 719–724 (2009).

75. Balkwill, F. R., Capasso, M. & Hagemann, T. The tumor microenvironment at a glance. *J. Cell Sci.* **125,** 5591–5596 (2012).

76. Lancaster, M. A. & Knoblich, J. A. Organogenesis in a dish: modeling development and disease using organoid technologies. *Science* **345,** 1247125 (2014).

77. Bregenzer, M. E. *et al.* Integrated cancer tissue engineering models for precision medicine. *PLoS One* **14,** e0216564 (2019).

78. Wang, D. H. & Souza, R. F. Biology of Barrett's esophagus and esophageal adenocarcinoma. *Gastrointest Endosc Clin N Am* **21,** 25–38 (2011).

79. Lee, J. *et al.* Tumor stem cells derived from glioblastomas cultured in bFGF and EGF more closely mirror the phenotype and genotype of primary tumors than do serum-cultured cell lines. *Cancer Cell* **9,** 391–403 (2006).

80. Wenger, S. L. *et al.* Comparison of established cell lines at different passages by karyotype and comparative genomic hybridization. *Biosci. Rep.* **24,** 631–639 (2004).

81. Ben-David, U. *et al.* Genetic and transcriptional evolution alters cancer cell line drug response. *Nature* **560,** 325–330 (2018).

82. Cooke, S. L. *et al.* Genomic analysis of genetic heterogeneity and evolution in high-grade serous ovarian carcinoma. *Oncogene* **29,** 4905–4913 (2010).

83. Hristova, V. A. & Chan, D. W. Cancer biomarker discovery and translation: proteomics and beyond. *Expert Rev Proteomics* **16,** 93–103 (2019).

84. Dawson, M. A. & Kouzarides, T. Cancer epigenetics: from mechanism to therapy. *Cell* **150,** 12–27 (2012).

85. Silva, T. C. *et al.* TCGA Workflow: Analyze cancer genomics and epigenomics data using Bioconductor packages. [version 2; peer review: 1 approved, 2 approved with reservations]. *F1000Res.* **5,** 1542 (2016).

86. Morgan, M., Obenchain, V., Hester, J. & Pag`es, H. *SummarizedExperiment: SummarizedExperiment container.* (2018).

87. Pavlidis, P. & Noble, W. S. Analysis of strain and regional variation in gene
expression in mouse brain. *Genome Biol.* **2,** RESEARCH0042 (2001).
88. Geman, D., d Avignon, C., Naiman, D. Q. & Winslow, R. L. Classifying gene
expression profiles from pairwise mRNA comparisons. *Stat Appl Genet Mol Biol* **3,**
Article19 (2004).
89. Krstajic, D., Buturovic, L. J., Leahy, D. E. & Thomas, S. Cross-validation pitfalls
when selecting and assessing regression and classification models. *J. Cheminform.*
6, 10 (2014).
90. Lipton, Z. C., Elkan, C. & Naryanaswamy, B. Optimal Thresholding of Classifiers to
Maximize F1 Measure. *Mach. Learn. Knowl. Discov. Databases* **8725,** 225–239
(2014).
91. Grow, E. J. *et al.* Intrinsic retroviral reactivation in human preimplantation embryos
and pluripotent cells. *Nature* **522,** 221–225 (2015).
92. Kolde, R. *pheatmap: Pretty Heatmaps*. (CRAN, 2019).
93. Wickham, H. *ggplot2 - Elegant Graphics for Data Analysis* . (Springer-Verlag New
York, 2016). doi:10.1007/978-0-387-98141-3
94. Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations
in multidimensional genomic data. *Bioinformatics* **32,** 2847–2849 (2016).
95. Yoshihara, K. *et al.* Inferring tumour purity and stromal and immune cell admixture
from expression data. *Nat. Commun.* **4,** 2612 (2013).
96. Kovalchik, S. *RISmed: Download Content from NCBI Databases*. (CRAN.R-project,
2017).
1211

Figure 1

A

RNA-seq from …

B

C

(1) Set parameters *n, m*

(2) Randomly select 2/3 TCGA data; run training process

(3) Assess performance on 1/3 held out data

(4) Repeat steps (2-3) 5 times

(5) Repeat steps (1-4) for each parameter set

Select parameter set with maximum mean AUPRC. Train on all TCGA data

D

E

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

A

B

C

D

E

Figure 7

Figure 8

Supplemental Figure 1

Supplemental Figure 2

Supplemental Figure 3

Supplemental Figure 4

Supplemental Figure 5