Analysis of female song provides insight into the evolution of sex differences in a widely studied songbird

Matthew R. Wilkins[1,2]*, Karan J. Odom[3], Lauryn Benedict[4], Rebecca J. Safran[1]

Keywords: sexual selection, female song, complex signaling, multimodal signaling, *Hirundo rustica*, animal communication

1-University of Colorado at Boulder, Department of Ecology and Evolutionary Biology

2-Vanderbilt University, Collaborative for STEM Education and Outreach

3-Cornell University, Cornell Lab of Ornithology and Department of Neurobiology and Behavior

4-University of Northern Colorado, School of Biological Sciences

*-Corresponding Author
m.r.wilkins06@gmail.com (preferred)
matthew.r.wilkins@vanderbilt.edu (institutional)

Word Count for Main Text: ~6,100

Author Contributions:

MRW collected data, performed analysis, and wrote initial manuscript draft; KJO measured and extracted acoustic metrics from audio files; LB undertook major restructuring of the manuscript; and RJS provided project administration and resources for carrying out the work. All authors contributed significantly to manuscript writing and revision.

ABSTRACT:

Understanding the patterns and processes related to sexual dimorphism and sex differences in
diverse animal taxa is a foundational research topic in ecology and evolution. Within the realm
of animal communication, studies have traditionally focused on male signals, assuming that
female choice and male-male competition have promoted sex differences via elaboration of
male traits, but selection on females also has the potential to drive sexual differentiation in
signals. Here, we describe female song in barn swallows (*Hirundo rustica erythrogaster*) for the
first time, report rates of female song production, and couple song data with plumage data to
explore the relative degree to which sex differences in phenotypic traits are consistent with
contemporary selection on males versus females. During previous intensive study of male song
over two years, we opportunistically recorded songs for 15 females, with matched phenotypic
and fitness data. We randomly selected 15 high-quality samples from our larger male dataset to
test whether sex differences in song and plumage are more strongly associated with fledgling
success for females or genetic paternity for males. Analyses included 35 potential sexual
signals including 22 song parameters and 13 plumage traits. Outcomes indicate that: female
songs were used in multiple contexts, restricted primarily to the beginning of the breeding
season; song traits showed greater sexual differentiation than visual plumage traits; and trait
correlations with reproductive success in females, rather than males, predicted sex-based
differences in song and plumage. These results are consistent with phylogenetic studies
showing that sex-based phenotypic differences are driven by changes in females, highlighting
the potential role of female trait evolution in explaining patterns of sexual differentiation. To
achieve a better understanding of sex differences and dimorphism, we require comprehensive
studies that measure the same traits in males and females and their fitness consequences.

*KEYWORDS***:**
sexual selection; female song; sexual dimorphism; multimodal signaling; barn swallow; Hirundo
rustica; animal communication

INTRODUCTION:

Ecologists and evolutionary biologists have long sought to understand the processes driving
dimorphism and other sex-based phenotypic differences (Andersson, 1994; Badyaev & Hill,
2003; Burns, 1998; Darwin, 1859, p. 94; Darwin, 1801, p. 396; Hedrick & Temeles, 1989;
Lande, 1980; Ng et al., 2019). However, owing to historical biases, studies of the drivers of
differentiation in sexual signaling traits have traditionally focused on male signals, and most
approaches assume that sexual selection has promoted sex differences via elaboration of male
traits (Badyaev & Hill, 2003; Freed, 2000; Langmore, 1998; Riebel, 2016; Riebel et al., 2005,
2019; Rosvall, 2011). Yet, meta-analyses of the strength of sexual selection on male traits
report moderate effect sizes (Jennions et al., 2012), with evidence that males are often under
variable selection pressures within and across breeding seasons (Chaine & Lyon, 2008;
Kingsolver et al., 2012; Robinson et al., 2008; Steele et al., 2011), have trait values near optima
(Evans, 1998; Rodríguez et al., 2006), or that the magnitude of trait differences may not reflect
the strength of current selection on males (Miller et al., 2016). Indeed, sex differences in signals
can be caused by a range of selection pressures on both males and females, resulting in
exaggeration or reduction of a variety of sex-specific signals (Bell & Zamudio, 2012; Dunn,
Armenta, and Whittingham 2015; Price, 2015; Shultz & Burns, 2017; Wiens, 2001). To better
understand how divergent selection operating between the sexes drives sex differences, we
require holistic approaches that fully describe the traits of both males and females (Hare et al.,
2019; Riebel et al., 2019). Ideally, studies should include multiple signaling traits that mediate
inter- and intra-sexual interactions (Bro-Jørgensen, 2010; Hebets et al., 2016; Hebets & Papaj,
2004; Partan & Marler, 2005), as well as their fitness consequences.

Bird song and plumage color offer excellent examples for how female, as well as male,
signal evolution can lead to sex differences. Females sang in the ancestor of modern songbirds
and still sing in many tropical and temperate oscines (Garamszegi, et al., 2006; Odom et al.,

2014; Price et al., 2009), suggesting that losses of female song have most often created sex-
based differences in song. Similarly, in species in which females have dull plumage, there has
been repeated, independent selection on feather color in both males and females (Dale et al.,
2015; Hofmann et al., 2008; Price et al., 2014), with females experiencing more evolutionary
plumage change than males in some lineages (Price et al., 2014). Even though losses of female
song and elaborate plumage appear to contribute considerably to sex differences in signaling
traits, the processes and conditions leading to these changes within individual species have not
received much attention, particularly in species where female song is rare (Brunton et al., 2016;
Kleindorfer et al., 2016).

Large-scale evolutionary change in dimorphic signaling traits is generally assumed to
result because some signals increase fitness. Numerous studies have identified male plumage
traits associated with strong female preferences and increased fitness (Byers et al., 2010;
Møller, 1988; Rodríguez et al., 2006; Ryan et al., 2019; Searcy & Andersson, 1986). In contrast,
relatively few studies have looked for or uncovered relationships between female visual signals
and fitness proxies, but see: (Amundsen et al., 1997; Bulluck et al., 2017; Cain & Ketterson,
2011; Jawor et al., 2004; Pap et al., 2019). Likewise, many studies have examined the fitness
correlates of male song (Catchpole & Slater, 2003; Gil & Gahr, 2002), but only a handful have
looked at how female song impacts fitness (Brunton et al., 2016; Cain et al., 2015; Kleindorfer et
al., 2016; Krieg & Getty, 2016). Although inclusive, systems-based approaches can best
untangle the relative contributions of selection on males and females to signal evolution across
modalities (Hebets et al., 2016; Riebel et al., 2019), researchers have seldom included both
visual and acoustic phenotypes from both sexes within the same studies (Hebets et al., 2016;
Riebel et al., 2019), but see (Webb et al. 2016). Moreover, to our knowledge, no study has
directly tested whether the pattern of sex differences in vocal and visual sexual signaling traits is
better predicted by contemporary selection on males (as is often assumed) or females. This
likely reflects the difficulty of collecting reproductive success in the field, a lack of study in
females, and a resulting paucity of datasets where the same traits and associated reproductive
success are measured for both sexes. This is a particular challenge in species with infrequent
female song, although such species provide a fertile testing ground for asking questions about
why female song is reduced and differs from male song.

We propose that North American barn swallows (*Hirundo rustica erythrogaster*) offer a
valuable system in which to investigate the selection pressures governing the evolution of
sexual signaling traits and associated sex differences. We explored patterns of plumage and
song in a population of male and female barn swallows by: 1) Describing female song in this
species, 2) exploring the form and seasonal timing of song production in both sexes, 3)
determining which acoustic and visual traits are robustly dimorphic, and 4) testing whether sex
differences are more strongly associated with contemporary selection (i.e. trait correlations with
reproductive success) in males or females. Because the differences between several measured
male and female traits were not categorical, we conservatively discuss them as sex differences
rather than dimorphism although other studies of this system may choose to call them
dimorphic.

Much previous research is predicated on the idea that sex differences and dimorphism in
communicative traits result from selection for male trait exaggeration (Badyaev & Hill, 2003;
Freed, 2000; Langmore, 1998; Riebel, 2016; Riebel et al., 2005, 2019; Rosvall, 2011). Here, we
test that assumption by including parallel data for both sexes and testing for linear associations
between a measure of current selection and the magnitude of trait sexual differentiation in a
variety of visual and acoustic traits (similar to Badyaev & Martin, 2000). If sex differences are
driven/maintained by directional selection on males (for showier or more elaborate traits), we
expect that trait correlations with male reproductive success will predict degree of sexual
differentiation. However, if differences are driven/maintained by directional selection on females
(for more cryptic or energy-efficient communication), we expect that trait associations with
female reproductive success will predict levels of sexual differentiation. While the combined

effect of differential selection on the sexes is ultimately responsible for the overall pattern of sex
differences (Hedrick & Temeles, 1989; T. D. Price, 1984), we are chiefly concerned with testing
the common assumption that it is sexual selection on males, rather than females, that drives this
pattern in signaling traits. Alternatively, if trait sex differences arose in the past, but are currently
under stabilizing selection, we would expect to find that the magnitude of trait sex differences
have no contemporary associations with reproductive performance.

METHODS:

Background

As the subject of hundreds of studies on sexual signal evolution since the late 1980s, barn
swallows offer an excellent system for testing questions related to sex differences. Barn
swallows are a weakly dimorphic species, in which males have darker ventral plumage and
longer tail feathers on average. They comprise six Holarctic subspecies that have rapidly
diversified from a common ancestor as recently as 7,700 years ago (Smith et al., 2018). In
North American barn swallows (*H. r. erythrogaster*), males with darker ventral feathers have
higher reproductive success, while tail feather length is not a preferred trait in males (Eikenaar
et al., 2011; Safran & McGraw, 2004; Safran et al., 2016), or females (Safran & McGraw, 2004).
There is limited evidence of selection for darker plumage in females in a New York population of
H. r. erythrogaster (Safran & McGraw, 2004).

Previous barn swallow studies indicate that different components of male song (e.g.
song rate and duration, rattle frequency and length) may provide receivers with information on
the age, viability, condition, motivation and/or overall quality of singers (Dreiss et al., 2008;
Galeotti et al., 1997, 2001; Garamszegi et al., 2005; Garamszegi, Hegyi, et al., 2006; Saino et
al., 2003; Wilkins et al., 2015). Despite intensive study of this species, female signals have
received relatively less attention. While there is a modest body of research into male song, there
are no indexed papers describing female barn swallow vocalizations in any detail. Females
have been reported to produce a "twitter-warble" song (Brown & Brown, 2020), but no formal
descriptions or quantitative analyses of that song exist (see Supplementary Methods and
Background for clarifying notes on female song).

General Field Methods

Barn swallows in this study were part of a long-term study population in Boulder County, CO,
which was set up in 2008 by Author 1 by contacting local birders and equestrian clubs,
surveying the county for culverts and structures with old nests, and, after arrival, looking for
signs of flying or perched swallows nearby. The five sites included in this study covered roughly
65 km^2 and ranged from 8 to 64 banded individuals. Persistent mist netting effort throughout the
breeding season resulted in the capture and unique marking of every or nearly every bird
present at each study site (using color bands, and application of Sharpie marker ink
combinations to left and right rectrices). These individual markings were used to identify
individuals during sound recording. At the time of capture, tail streamer length was measured
and a small sample (<90μL) of blood was taken via brachial venipuncture and stored in 2% lysis
buffer for later parentage analysis. Additionally, a set of ~5 contour feathers were sampled
along a ventral transect (throat, breast, belly, and vent), and attached to index cards with tape
for color analysis using a spectrophotometer in the lab (Hubbard et al., 2017; Wilkins et al.,
2015). Nests were closely monitored for egg-laying activity, with checks conducted at least
every four days to determine clutch initiation date, and date of hatching. On approximately day
12 post-hatching, all nestlings in a nest were banded and blood samples taken for parentage
analysis.

Ethical Note

We made every possible attempt to minimize handling time and other sources of stress on
swallows in our study population. Capture typically involved closing off all but one exit from a

barn or other structure, in front of which a fine mesh nylon mist net was held or attached in
place with bungee cords. Nets were constantly monitored and birds were usually removed
within one minute of landing in the net and placed into custom-sewn cloth bags (closed with a
ribbon) to calm them until they could be transported to a temporary banding station a short
distance away (usually <30m), sampled, and released. Birds held temporarily in bags were kept
in a quiet, cool location until they were removed for processing. Typically, birds are quiet and
fairly still both in the net and while in the bag. Obvious signs of stress during handling were
attended to constantly. In the rare instances where a bird showed signs of distress during
netting, banding, measurement, or blood sampling, it was immediately released. All field
methods were approved by the University of Colorado Institutional Animal Care and Use
Committee (Protocols 07-07-SAF-01 and 1004.01).

Visual Trait Measures

Consistent with a previous study on complex male signaling in this population (Wilkins et al.
2015), we measured maximum tail streamer length (rather than mean, as it is more normally
distributed; See Table S1 for trait definitions). Additionally, we measured a total of 12 traits to
characterize color variation: three axes of color (average brightness, hue, and red chroma)
along a four-patch ventral transect (throat, breast, belly, and vent). We opted to use these hue-
saturation-brightness (HSB) measures to make interpretation and comparison to other studies in
this system as easy as possible. We are also confident about the biological relevance of these
trait measures, since they strongly correlate with eumelanin and phaeomelanin concentrations
in feathers (McGraw et al., 2005), ventral feathers do not show an ultraviolet reflectance peak
(Safran & McGraw, 2004), and color manipulations (measured using HSB) to mimic natural
melanization profiles have shown predictable impacts on physiology and reproductive
performance (Safran et al., 2005, 2008, 2016). However, as a check, we calculated color
measures from raw spectra in the tetrahedral color space (TCS) that incorporates a model of
avian visual sensitivity (Endler & Mielke, 2005; Stoddard & Prum, 2008). See Supplementary
Methods for details on how we calculated both HSB and TCS color traits. As shown in Figure
S2, average brightness is exactly equivalent to the TCS measure 'brilliance' described in
Stoddard and Prum (2008). Our measures of chroma showed a strong positive (0.84)
Spearman's correlation with the equivalent TCS measure of saturation (rA, achieved chroma).
Hue showed a moderate (-0.35) correlation with brilliance, though direct comparison with TCS
measures of hue is challenging because TCS θ and Φ jointly describe stimulation of avian
cones and a single correlation between hue and one of these measures in not meaningful.
Overall, given biochemical and experimental evidence, and a lack of significant UV reflectance
in barn swallow ventral coloration (Hubbard et al., 2017, McGraw et al., 2005), we are very
confident that our HSB measures of color capture biologically meaningful variation in this
system.

Fitness Measures

We used female seasonal (i.e. annual) fledging success and male within-pair genetic paternity
as fitness proxies for females and males, respectively. Female fledging success was calculated
as the number of offspring leaving the nest across all broods. Genetic paternity was calculated
as the number of fledged genetic offspring within the social nest across all broods, determined
through genetic paternity exclusions. Paternity assignments (and total extra-pair paternity) were
not feasible in our study area, due to the presence of known, unmonitored breeding sites within
easy flying distance from monitored sites. To get a representative sample of males' EPY would
also require genotyping thousands of individuals in any given year. Briefly, genotypes were
derived from fluorescent-labeled PCR products of six microsatellite loci. Paternity was
conducted using CERVUS v3.03 (Kalinowski et al., 2007) and an offspring was considered
extra-pair when the offspring-mother-father trio confidence did not reach the 95% level. Detailed
paternity exclusion methods are in the Supplementary Methods. Four of the 12 fully-sampled

males and females were social pairs, though we did not treat this statistically as the fitness
correlational analysis was conducted separately for each sex.

Recordings

Female barn swallows were recorded opportunistically as part of a study of male song (Wilkins
et al., 2015) conducted in 2011 and 2012. Though males were the target of the prior study, this
rarer, unpredictable female vocalization was of growing interest to Author 1. When a female
vocalization was heard, Author 1 immediately redirected the microphone at the source of the
sound, capturing as many vocalizations as possible. Given the complex acoustic environment in
which barn swallows live, active, annotated recording is necessary for confident singer IDs.
Thus, our samples by no means capture comprehensive singing outputs for individuals;
however, our estimates of relative singing outputs should reflect realistic activity patterns. Songs
were recorded in 16-bit WAV format, with 48kHz sampling rate using a Marantz PMD 660 paired
with a Sennheiser MKH 20 and Telinga parabola (2011), or a Marantz PMD 661 paired with a
Sennheiser ME62/k6 microphone and Telinga parabola (2012). Total recording time was
approximately 57 hours (May 2-May 31) in 2011 and 48 hours (May 1-Aug 21) in 2012, all
between 5am and noon (Wilkins et al., 2015).

Acoustic Trait Measures

Using Raven v1.5 (Bioacoustics Research Program, 2011), we extracted acoustic parameters of
all recorded songs from the 15 females (n=64) and a comparable number of songs (n=66) for 15
males (chosen randomly from among those with high quality song recordings) to create two
acoustic datasets: (1) an element-level dataset of acoustic parameters measured for every
element in all songs, and (2) a song-level dataset of acoustic parameters averaged for all
elements in a song or calculated for the whole song. The element-level dataset was used to
estimate an element diversity score (Keen et al., in review), which estimates the variability, or
diversity, of elements for a given song within an acoustic feature space of all barn swallow
element measurements (see full explanation in Supplementary Methods). This value was then
appended to the song-level dataset, which was used for all subsequent statistical analyses
involving song.

Of the 18 song-level variables extracted (see Supplementary Methods), we selected 11
that were likely to be robust to noise in recordings from barn swallow habitats and relevant for
sexual signaling, based on previous studies of male song (Galeotti et al., 1997; Garamszegi,
Hegyi, et al., 2006; Wilkins et al., 2015). The 11 selected acoustic variables were: song entropy
(a measure of tonality), dominant frequency range, mean peak frequency, number of elements
per song, element duration, element rate, song duration, gap duration, frequency interquartile
range, and element diversity (see Table S1 for definitions). We then calculated individual means
and coefficients of variation for each acoustic variable. This resulted in 22 variables describing
differences in the magnitude and variability of different aspects of song frequency, timing, and
elemental diversity for both sexes. The number of song traits was further reduced to avoid
redundancy in analysis of sex differences and selection, as described below.

Statistical Analysis

Our sample included 15 females with at least one song measured. To make female-male
comparisons that are not skewed by uneven sample sizes, we randomly selected 15 males with
high quality recordings from our data set. Our subsample of males was temporally equivalent to
that for females, as the sexes did not differ in recording date relative to first clutch initiation:
mean ± (SD) for females: -6.6 ± (10.7) and for males: -2.6 ± (21.1); $t_{unpaired}$= -0.65, df= 20.8, p=
0.52. For both sexes, due to missing data, we had 12 individuals with acoustic, visual, and
fitness data. Because three females only had a single song recorded and coefficient of variation
could not be calculated, only nine females had data for all 36 variables (22 acoustic traits,13
visual traits, and a fitness proxy). Thus, correlations between a particular trait and fitness metric
in females were calculated from sample sizes ranging from nine to 15 individuals. Although our

sample sizes are limited, the data set represents extensive population sampling time, and
provides a starting point and methodology for more concerted effort.

All analyses were conducted using R v3.6.0 (R Core Team, 2019). To measure the
magnitude of the sex difference for each trait, we used Cohen's d, calculated as the difference
of male from female population means divided by the pooled standard deviation (Cohen, 1988,
p. 21). Thus, negative difference values indicate male-biased differences and positive values
indicate female-biased differences. For example, in barn swallows, tail feathers show male-
biased differentiation (as males have longer tails than females on average), resulting in a
negative value on our sex difference scale. To determine which traits were robustly distinct, we
created 10,000 bootstraps using the 'bootstraps' function in the {rsample} package v0.0.5 (Kuhn
et al., 2019), resampling sets of 15 males and 15 females with replacement. We then calculated
95% confidence intervals from this posterior distribution using the base function 'quantile.'

To avoid collinearity in further analysis of dimorphic traits, we identified variables with
Spearman's rank correlations >|0.7| and retained the most biologically intuitive variables for
analysis. For example, average number of elements, song duration, and our measure of
element diversity showed pairwise correlations >0.8. As element diversity is a higher level
measure of complexity, for which there is a wide literature suggesting it to be a common target
of selection (Benedict & Najar, 2019; Snyder & Creanza, 2019), we retained mean element
diversity in our set of dimorphic traits, and discarded mean song duration and mean number of
elements. After filtering out redundant and non-dimorphic traits, we were left with 10 visual and
acoustic traits for exploring the connection between sex differences and contemporary selection
pressures.

Testing trait-wise associations between reproductive success and sexual differentiation

As a measure of contemporary selection pressures, we calculated each trait's Spearman's rank
correlation with a fitness surrogate (seasonal number of fledged offspring for females and
seasonal number of within-pair genetic offspring for males). Finally, in order to test the linear
relationship between our surrogate measures of selection and observed trait sex differences,
while accounting for nonindependence of traits, we used the 'crunch' function in the {caper}
package v1.0.1 (Orme et al., 2018), which implements the CAIC (Comparative Analysis by
Independent Contrasts) algorithm (Purvis & Rambaut, 1995). Following a similar approach to
Garamszegi et al. (2006), we treated traits as species with known phylogenetic relationships
(i.e. correlation structure). To build the tree used by 'crunch,' we calculated trait distances as 1 -
abs(Spearman's correlation), and constructed the tree using a single-linkage clustering method
with the base 'hclust' function.

Thus, we made a linear model with a trait's magnitude of sex difference as a response
and a trait's correlation with reproductive success as a predictor, controlling for correlation
structure among traits. Given our metric of sexual differentiations, we would expect a significant
positive association between a trait's correlation with reproductive success and the magnitude
of sexual differentiation if female fitness is maintaining sex-based trait differences. That is, traits
that confer fitness benefits to females should have a high degree of differentiation for our metric,
and traits that reduce fecundity in females should have low differentiation values. We would
expect the opposite (a negative slope) if male fitness is driving/maintaining sexual
differentiation. That is, traits associated with higher paternity would be more exaggerated in
males (having more negative differentiation values on our metric), and traits that reduce
paternity would be less exaggerated in males (higher differentiation values), resulting in a
negative slope.

Data availability

Male and female song clips, HSB and TCS color, and other phenotypic and fitness data are
available in our GitHub repository at https://github.com/drwilkins/femaleSongInBARS.

Spectrograms in Figure 1 were generated with R code, available at
https://github.com/drwilkins/rspect

RESULTS

Characterizing female song

Recordings and observations revealed that females used a distinctive vocalization in
spontaneous solo renditions, and in response to hearing other females, in a manner
synonymous to male counter-singing. They were also observed using the vocalization to
interrupt the songs of their mates (see supplementary video 1–vimeo.com/424642268). This
vocalization is relatively complex, including many syllables with similar frequency modulation
and pace to male song (Figure 1). Collectively, this evidence confirms that female barn
swallows do produce facultative song, rather than just simple calls (Langmore, 1998).

338
Figure 1. Spectrograms of female and male barn swallows. We show two song renditions for 3 females and 3 males
to demonstrate variability. Photos by Author 1.

Female song bouts were often produced within the nest, but also on perches near the
nest or outside the barn/structure and were usually short and infrequent. As such, only 78
songs were recorded from 18 identifiable females over the course of two years, compared to

753 (865% more) songs from 40 males, given the same recording effort (~105 hours). This
amounts to a recording efficiency of about 7.17 clear, identifiable songs/hr for males and 0.74
songs/hr for females over the course of the study. The sexes also differed strikingly in the
phenology of song production. Male songs were observed and recorded between May 1st and
August 21st, while female songs were only observed and recorded between May 10th and May
29th (pooling years) (Figure 2a). Restricting to only the active singing period for females during
May of both years, a cumulative 91 hours of recording documented 8.9 individually identifiable
songs/hr for males and 0.86 songs/hr for females. We could not calculate exact song rates per
individual due to the complexities of recordings with shifting focal singers in dynamic colonial
environments. To explore effects of breeding phenology on singing output, we also calculated
relative recording dates by subtracting the ordinal date of first clutch initiation (i.e. breeding
onset) from the ordinal recording date. For males, these relative recording dates ranged from
59 days before to 109 days after the first egg was laid by a males' mate, with an average date
(weighted by number of songs recorded each day) of 5.43 days *after* breeding onset; Figure
2b). For females the range was 29 days before to 14 days after breeding onset, with a weighted
average of 3.90 days *before* breeding onset. Due to dropouts resulting from low recording
quality, the total number of songs analyzed was 48 for females (mean=4.0, range= [1,11],
sd=3.02) and 66 for males (mean=4.4, range= [2,5], sd= 0.99).

362
Figure 2. Seasonal variation in song production, as demonstrated by opportunistic recording of spontaneous male
and female songs. A) Shows number of songs recorded for both sexes, as a function of recording date. In B) the
ordinal date of breeding onset (first clutch initiation) has been subtracted from the ordinal recording date; thus,
negative x-values represent recordings taken before breeding onset. The dashed vertical line marks Day 4 post
clutch initiation–i.e. the last day of female fertility before laying of the ultimate egg of a modal 5-egg clutch. Lines
connect points for individuals with recordings on multiple days.

Visual and acoustic differentiation in males and females

Our bootstrap analysis of 35 acoustic and visual traits showed that 16 were robustly distinct in the two sexes–i.e. had bootstrap 95% confidence intervals that did not overlap zero (Figure 3). These include a mixture of song and visual traits, with song traits being most distinctive (13/22=59% for acoustic traits, compared to 3/13=23% for visual traits). Traits that had higher values in females (showing female-biased differences) included: entropy, CV_Mean Peak Frequency, CV_Dom Frequency Range, CV_Frequency IQR, CV_Element Diversity, Belly Brightness, and Element Duration. Male-biased dimorphic traits included: Vent Chroma, Frequency IQR, Mean Peak Frequency, CV_Entropy, Frequency Range, Tail Length, Element Diversity, Song Duration, and Element Count. Most notably, female songs are brief in duration, include few elements, include elements with high entropy (i.e. noise), and have low element diversity across songs, compared to males (Figure 3). Consistent with previous results for visual traits (Safran & McGraw, 2004), females had shorter tail streamers and lighter ventral plumage (higher brightness scores for belly and lower chroma scores for the vent). Trait means, sample sizes, and sex difference confidence intervals are reported in Table S2.

Figure 3. Sex differences in 35 potential sexual signaling traits. Empirical point estimates for Cohen's d are shown, with 95% bootstrapped confidence intervals not overlapping zero bolded in purple. Symbols for points indicate trait modality (circle=acoustic, t triangle=visual). Traits are means, except where preceded by CV (coefficient of variation). Trait names beginning with a star were the chosen set of robustly sexually distinct traits used for further analysis, after removing highly redundant traits with Spearman's correlations ≥|0.7|.

Connecting contemporary selection to phenotypic sex differences

Our final objective was to test whether sexual differentiation is more strongly associated with
contemporary selection in males or females (here, using surrogate measures of lifetime fitness:
within-pair reproductive performance for a single breeding season). After controlling for
intercorrelations between the 10 nonredundant, sexually distinct traits using independent
contrasts, we found a significant positive relationship between trait sex differences and our
measure of current selection pressures (i.e. correlation with either seasonal fledging success or
genetic paternity) in females (df=8, slope=6.04, t=2.92, p=0.019), but not males (df=8,
slope=1.06, t=0.547, p=0.599) (Figure 4). That is, traits that had a negative association with
female reproductive success (see Figure S1) showed male-biased sex differences (i.e., greater
trait expression in males), while traits that positively correlated with female reproductive success
showed female-biased sex differences. The relationship remained significant for females when
including all 16 robustly distinct traits (df=14, slope=4.93, t=3.53, p=0.003), as well as for the
subset of distinct acoustic traits (df=11, slope=5.28, t=3.70, p=0.003). In contrast, the
relationship between male traits and within-pair genetic paternity did not predict levels of sex
differences for either of these subsets (all p≥0.260).

Figure 4. Relationship between trait sexual differentiation and trait association with reproductive output for A) females
and B) males. Sexual differentiation was measured as the difference of scaled female and male trait values in pooled
standard deviation units (Cohen's d). Reproductive output was measured as seasonal fledging success for females
and fledged within-pair seasonal genetic paternity for males. Females showed a much stronger (and significant)
positive relationship between how traits impacted reproductive performance and the observed amount of sexual
differentiation.

DISCUSSION:

Female song in barn swallows

Despite intensive study of this species for decades, female song in barn swallows has been
reported as absent by some authors (Sibley, 2014; Stokes & Stokes, 2010), the same as male
song by others (Brown & Brown, 2020), or left ambiguous (Pieplow, 2017) (see Supplementary
Methods for notes on previous descriptions). Our data and videos confirm that female barn
swallows sing a quantitatively distinct song from males. We report that while female song is
approximately ten times less frequent than male song and relegated largely to the first month of
the breeding season, it has an overall similar structure to male song and seems to be used
similarly in counter-singing, as well as mate-interruption contexts. The use of female song in

counter-singing (see Supplementary Video 1) just prior to clutch initiation is consistent with
singing patterns and use of song for territory/nest defense or intra-sexual competition in other
Northern temperate breeding species (Cain et al., 2015; Krieg & Getty, 2016; Levin, 1996;
Magoolagan et al., 2019; Rose et al., 2019; Yasukawa, 1989, 1990). Specifically, female barn
swallow song could be actively maintained and used by females during this short period of the
breeding season when females are establishing and competing for nest sites (Krieg & Getty,
2016; Rosvall, 2011). Further, the use of female song to interrupt male song fits with theories
that describe song functionality in mate attraction, pair bonding, and mate signal jamming (Grafe
et al., 2004; Tobias & Seddon, 2009).

Although female and male barn swallow songs have a generally similar form, female
songs lack the terminal trill which is universal for male song among different barn swallow
subspecies (Wilkins et al., 2018). Our results further indicate that female songs are shorter and
noisier, with fewer, more variable elements, reduced frequency ranges, and lower element
diversity (Figures 3, 4, S1). Thus, barn swallow song is categorically dimorphic in some aspects
and shows variable sex differences in others. Overall, in this species song traits are more
distinct by sex (59% of measured traits) than plumage traits (23% of measured traits) and the
direction of sex differences is predicted by both the trait correlations with female fledging
success and qualitative expectations from selection for efficient, cryptic signaling in females. For
example, element diversity (our measure of average syllable complexity, which was retained
after removing the redundant measure "song duration") is much lower in females and showed a
strong negative relationship with fledging success. In contrast, element duration (the length of
individual song elements) was significantly longer in females and this trait showed the strongest
positive correlation with reproductive success of any trait. Together, these results are consistent
with selection for shorter songs with longer individual elements. Shorter songs may be favored
to avoid attracting nest predators, an effect previously shown for song rate (Kleindorfer et al.,
2016). Shorter songs also offer the cognitive benefit of simpler comparison due to easier
discriminability of proportional differences (Akre et al., 2011; Akre & Johnsen, 2014).

The fact that mean element diversity is male-biased and negatively correlates with
female fledging success, but CV_element diversity (i.e. song versatility: variability of syllable
complexity across songs) is female-biased and positively correlates with fledging success was
unexpected. One possible explanation for this may stem from variable functions (and
audiences) for female song. That is, signals in competitive contexts tend to be shorter and more
repetitive, which would select for low element diversity if this is the primary function of song
(Collins et al., 2009; Galeotti et al., 1997). On the other hand, song versatility and the ability to
adaptively change element diversity across competitive and mating-related signaling contexts
could be especially important for females if they are constrained to shorter signal rate and
duration, and there is a premium on efficient information transfer about motivation or quality
within a shorter time window than males. This is, of course, speculative and close study of the
signaling context of song production is necessary to better understand these patterns of female
song variation.

Taken together, our results provide a mechanism supporting the evolution of female
song in many North Temperate bird species from complex songs comparable to male songs
toward highly condensed, context-dependent songs, or even toward the loss of songs altogether
(Odom et al., 2014). This is significant because, while the vast literature on male birdsong (and
acoustic signal evolution more broadly) has informed many aspects of sexual selection theory
(Gil & Gahr, 2002; Nowicki et al., 2002), and male acoustic divergence has been shown to play
a key role in premating isolation and speciation across diverse taxa (Alcaide et al., 2014;
Blankers et al., 2019; Hasiniaina et al., 2020; Irwin et al., 2001; Lee et al., 2016; Sosa-López et
al., 2016; Wilkins et al., 2013, 2018), we know relatively little about female vocal signaling or its
implications for broader evolutionary processes, even in classic study systems like barn
swallows.

Overall patterns of sexual differentiation in signals

In contrast to a common assumption in the sexual selection literature, we found that
overall patterns of sex differences are better explained by (surrogate measures of) selection on
females, rather than selection for more elaborate males. That is, the overall magnitudes of sex-
based differences in song form, tail streamer length, and color were predicted by how those
traits correlated with female, but not male reproductive performance. This does not mean that
selection on males (i.e. through female choice) is not important or relevant in explaining sex
differences. For example, as shown in Figures 4B and S1, belly brightness–a known sexual
signal in males within this population (Safran et al., 2016; Wilkins et al., 2015)–showed a strong
negative association with genetic paternity (selecting for darker males with lower brightness). In
turn, belly brightness in females showed a slight positive association with fledging success, and
sex difference was low (0.8 SD), but significantly biased towards lighter females. That is, higher
reproductive performance among darker males, combined with negligibly higher performance in
lighter females correlates with an overall pattern of slightly darker males. In contrast, a strong
positive correlation between element diversity and reproductive success in males and a strong
negative correlation for females (i.e. divergent selection in the sexes) corresponds with a highly
male-biased sex difference (-3.11 SD) toward more complex syllable composition of songs.
While an additional analysis testing the combined impact of trait-wise reproductive performance
on trait-wise sex differences for both sexes is desirable, it would likely be similar to the result
shown for females (Figure 4A) and is not possible using our current comparative analysis by
independent contrasts approach, as we are unaware of a way to control for known differences in
trait correlation structure between the sexes in a single model.

Broader implications

Although our small sample size limits our ability to rigorously estimate selection
gradients or account for error in our estimate of the correlation between traits and fitness
proxies, our findings offer baseline evidence that selection on females may be more consistent
and therefore more important in maintaining phenotypic distinctions between the sexes within
populations. This is perhaps unsurprising, given previous studies showing variability in female
preferences for male traits (Chaine & Lyon, 2008; Kingsolver et al., 2012; Robinson et al., 2008;
Steele et al., 2011). Nevertheless, it has important implications for comparative studies which
rely on the degree of plumage differences as a surrogate for the strength of sexual selection. It
is widely recognized that there are limitations to using sex differences, including dimorphism, as
a surrogate for sexual selection (Bell & Zamudio, 2012; Huang & Rabosky, 2014; Kraaijeveld et
al., 2011; Price, 2015). We would add the caveat that researchers must consider the action of
sexual selection on both males and females, and include the traits of both sexes, even if traits
are reduced or difficult to sample. Many studies assume that sex differences largely result from
sexual selection for exaggerated male traits in conjunction with ecological selection for crypsis
on females. This may be a valid assumption in some cases, such as in sexual dichromatism in
damselflies (Svensson & Waller, 2013). However, our results highlight the need (especially in
taxa with more complex communication and/or cognition) to consider how female competition
and mutual mate choice affect the evolution of sex differences within the context of constraining
ecological selection on females from increased costs of predation, migration, and/or coloniality
(Jawor et al., 2004; Price, 2015; Tobias et al., 2011). Outcomes of this study suggest that
following initial evolution of a mutually ornamented ancestor, sex differences and dimorphism in
a broad suite of visual and acoustic traits was most likely created and maintained in barn
swallows through countervailing selection on females, rather than on males (Kraaijeveld et al.,
2007).

CONCLUSIONS:

With accumulating evidence for mutual mate choice and mutual ornamentation as the
ancestral state across diverse taxa (Edward & Chapman, 2011; Kraaijeveld et al., 2007; Odom
et al., 2014; Hofmann et al., 2008), we offer a method for testing whether selection on females
may promote sex differences to a much greater extent than is currently appreciated.
Collectively, our results highlight the importance of studying rare behavioral phenomena and, in
particular, call for better documentation and targeted research on female song (Karan J. Odom
& Benedict, 2018). There are growing accounts of temperate-zone female birds that sing for
short time periods during the breeding season, particularly during settlement of breeding
territories (Hathcock & Benedict, 2018; Krieg & Getty, 2016; Taff et al., 2012). Thus, more
focused research effort during this time period may very well show that complete loss of song in
female Passerines is the exception, rather than the widely purported rule, and that similar
selective pressures shape female song in multiple species.

Though there is increasing evidence that female signaling traits can evolve as fast or
faster than male traits across species (Johnson et al., 2013; Dunn, Armenta, and Whittingham
2015; J. J. Price, 2019), we have few studies within species to guide our understanding of the
processes driving these patterns. In contrast to the vast literature on within-species geographic
divergence in male signals (McLean & Stuart-Fox, 2014; Podos & Warren, 2007; Slabbekoorn &
Smith, 2002; Velásquez, 2014), there are few studies for females in the visual modality (McCoy
et al., 1997; McLean & Stuart-Fox, 2014; Obara et al., 2008; Roulin, 2003; Tuomaala et al.,
2012) and fewer still in the acoustic modality (Graham et al., 2017, 2018; Mennill & Rogers,
2006; Odom & Mennill, 2012). Yet, it is worth noting that three studies (in butterflies and birds)
that considered intraspecific geographic variation in both sexes found greater signal divergence
in females than males across populations (Graham et al., 2018; Mennill & Rogers, 2006;
Tuomaala et al., 2012). The implications of female signal evolution, male mate
choice/recognition, and their impact on speciation have received little attention in the literature,
with a few relevant studies focusing primarily on fish and insects (Chung et al., 2014; Edward &
Chapman, 2011; Jiggins et al., 2004; Roberts & Mendelson, 2017, 2020). Targeted study of
both sexes in concert across taxa is necessary to gain a more holistic understanding of how
signals evolve within and among populations and how these trait changes feed into larger
ecological, evolutionary, and eco-evolutionary processes (Bondurianksy, 2011; Cole & Endler,
2015; Endler & Basolo, 1998; Fryxell et al., 2019).

ACKNOWLEDGEMENTS:

MRW thanks the Macaulay Library for loaning him audio equipment in 2011, and numerous
undergraduate and graduate students in the Safran Lab who helped collect the mountain of
phenotypic and fitness data that enabled this study. He would also like to thank the many site
owners who allowed us to traipse around their property at odd hours of the day (in particular, the
Colorado Horse Rescue). We also thank Megan Barkdull for help with sound analysis and Joey
Hubbard for sage counsel on color analysis.

FUNDING:

564   This work was supported by an American Ornithological Society: Joseph Grinnell Award and an
565   NSF Graduate Research Fellowship to MRW. KJO was supported by a U.S. National Science
566   Foundation Postdoctoral Research Fellowship in Biology (grant no. 1612861).

567
568

REFERENCES*:*

Akre, K. L., Farris, H. E., Lea, A. M., Page, R. A., & Ryan, M. J. (2011). Signal perception in
frogs and bats and the evolution of mating signals. *Science*, 333(6043), 751–752.

Akre, K. L., & Johnsen, S. (2014). Psychophysics and the evolution of behavior. Trends in
Ecology & Evolution, 29(5), 291–300.

Alcaide, M., Scordato, E. S. C., Price, T. D., & Irwin, D. E. (2014). Genomic divergence in a ring
species complex. *Nature*, 511(7507), 83–85.

Amundsen, T., Forsgren, E., & Hansen, L. T. T. (1997). On the function of female ornaments:
male bluethroats prefer colourful females. *Proceedings of the Royal Society B: Biological*
Sciences, 264(1388), 1579–1586.

Andersson, M. B. (1994). Sexual Selection. Princeton University Press.

Badyaev, A. V., & Hill, G. E. (2003). Avian Sexual Dichromatism in Relation to Phylogeny and
Ecology. In *Annual Review of Ecology, Evolution, and Systematics* (Vol. 34, Issue 1, pp.
27–49). https://doi.org/10.1146/annurev.ecolsys.34.011802.132441

Badyaev, A. V., & Martin, T. E. (2000). Sexual dimorphism in relation to current selection in the
house finch. *Evolution; International Journal of Organic Evolution*, 54(3), 987–997.

Bell, R. C., & Zamudio, K. R. (2012a). Sexual dichromatism in frogs: natural selection, sexual
selection and unexpected diversity. *Proceedings. Biological Sciences / The Royal*
Society, 279(1748), 4687–4693.

Benedict, L., & Najar, N. A. (2019). Are commonly used metrics of bird song complexity
concordant? *The Auk*, 136(1), 201.

Bioacoustics Research Program. (2011). *Raven Pro: Interactive Sound Analysis Software*
(Version 1.5) [Computer software]. http://www.birds.cornell.edu/raven

Blankers, T., Oh, K. P., & Shaw, K. L. (2019). Parallel genomic architecture underlies repeated
sexual signal divergence in Hawaiian crickets. *Proceedings. Biological Sciences / The*
Royal Society, 286(1912), 20191479.

Bonduriansky, R. (2011). Sexual selection and conflict as engines of ecological diversification.
The American Naturalist, 178(6), 729–745.

Bro-Jørgensen, J. (2010). Dynamics of multiple signalling systems: animal communication in a
world in flux. *Trends in Ecology & Evolution*, 25(5), 292–300.

Brown, M. B., & Brown, C. R. (2020). Barn Swallow - Hirundo rustica. *Birds of the World*.
https://doi.org/10.2173/bow.barswa.01

Brunton, D. H., Roper, M. M., & Harmer, A. M. T. (2016). Female Song Rate and Structure
Predict Reproductive Success in a Socially Monogamous Bird. Frontiers in Ecology and
Evolution, 4. https://doi.org/10.3389/fevo.2016.00013

Bulluck, L. P., Foster, M. J., Kay, S., Cox, D. E., Viverette, C., & Huber, S. (2017). Feather
carotenoid content is correlated with reproductive success and provisioning rate in
female Prothonotary Warblers. *The Auk*, 134(1), 229–239.

Burns, K. J. (1998). A Phylogenetic Perspective on the Evolution of Sexual Dichromatism in
Tanagers (Thraupidae): The Role of Female versus Male Plumage. *Evolution;*
International Journal of Organic Evolution, 52(4), 1219.

Byers, J., Hebets, E., & Podos, J. (2010). Female mate choice based upon male motor
performance. In *Animal Behaviour* (Vol. 79, Issue 4, pp. 771–778).
https://doi.org/10.1016/j.anbehav.2010.01.009

Cain, K. E., Cockburn, A., & Langmore, N. E. (2015). Female song rates in response to
simulated intruder are positively related to reproductive success. *Frontiers in Ecology*
and Evolution, 3. https://doi.org/10.3389/fevo.2015.00119

Cain, K. E., & Ketterson, E. D. (2011). Competitive females are successful females; phenotype,
mechanism, and selection in a common songbird. *Behavioral Ecology and Sociobiology*,
66(2), 241–252.

Catchpole, C. K., & Slater, P. J. B. (2003). *Bird Song: Biological Themes and Variations*.
Cambridge University Press.

Chaine, A. S., & Lyon, B. E. (2008). Adaptive plasticity in female mate choice dampens sexual
selection on male ornaments in the lark bunting. *Science*, 319(5862), 459–462.
Chung, H., Loehlin, D. W., Dufour, H. D., Vaccarro, K., Millar, J. G., & Carroll, S. B. (2014). A
single gene affects both ecological divergence and mate choice in Drosophila. *Science*,
343(6175), 1148–1151.
Cohen, J. (1988). *Statistical Power Analysis for the Behavioral Sciences* (2nd ed.). Lawrence
Erlbaum Associates.
Cole, G. L., & Endler, J. A. (2015). Variable environmental effects on a multicomponent sexually
selected trait. *The American Naturalist*, 185(4), 452–468.
Collins, S. A., de Kort, S. R., Pérez-Tris, J., & Tellería, J. L. (2009). Migration strategy and
divergent sexual selection on bird song. *Proceedings. Biological Sciences / The Royal
Society*, 276(1656), 585–590.
Dale, J., Dey, C. J., Delhey, K., Kempenaers, B., & Valcu, M. (2015). The effects of life history
and sexual selection on male and female plumage colouration. *Nature*, 527(7578), 367–
370.
Darwin, C. (1859). *On the origin of species by means of natural selection, or, The preservation
of favoured races in the struggle for life /*. https://doi.org/10.5962/bhl.title.68064
Darwin, E. (1801). *Zoonomia Or the Laws of Organic Life*.
Dreiss, A. N., Navarro, C., de Lope, F., & Møller, A. P. (2008). Effects of an Immune Challenge
on Multiple Components of Song Display in Barn Swallows Hirundo rustica: Implications
for Sexual Selection. Ethology: Formerly Zeitschrift Fur Tierpsychologie, 114(10), 955–
964.
Dunn, Peter O., Jessica K. Armenta, and Linda A. Whittingham. (2015). "Natural and Sexual
Selection Act on Different Axes of Variation in Avian Plumage Color." *Science Advances*.
https://doi.org/10.1126/sciadv.1400155.
Edward, D. A., & Chapman, T. (2011). The evolution and significance of male mate choice. In
Trends in Ecology & Evolution (Vol. 26, Issue 12, pp. 647–654).
https://doi.org/10.1016/j.tree.2011.07.012
Eikenaar, C., Whitham, M., Komdeur, J., van der Velde, M., & Moore, I. T. (2011). Testosterone,
plumage colouration and extra-pair paternity in male North-American barn swallows.
PloS One, 6(8), e23288.
Endler, J. A., & Basolo, A. L. (1998). Sensory ecology, receiver biases and sexual selection.
Trends in Ecology & Evolution, 13(10), 415–420.
Endler, J.A. & Mielke, P. (2005). Comparing entire colour patterns as birds see them. *Biol. J.
Linn. Soc*. 86: 405–431.
Evans, M. R. (1998). Selection on swallow tail streamers. In *Nature* (Vol. 394, Issue 6690, pp.
233–234). https://doi.org/10.1038/28297
Freed, L. A. (2000). Female ornamentation, mate choice and sexual selection. *Trends in
Ecology & Evolution*, 15(11), 471.
Fryxell, D. C., Weiler, D. E., Kinnison, M. T., & Palkovacs, E. P. (2019). Eco-Evolutionary
Dynamics of Sexual Dimorphism. *Trends in Ecology & Evolution*, 34(7), 591–594.
Galeotti, P., Saino, N., Sacchi, R., & Møller, A. P. (1997). Song correlates with social context,
testosterone and body condition in male barn swallows. *Animal Behaviour*, 53(4), 687–
700.
Galeotti, P., Saino, N., Perani, E., Sacchi, R., & Møller, A. R. (2001). Age-related song variation
in male barn swallows. In *Italian Journal of Zoology* (Vol. 68, Issue 4, pp. 305–310).
https://doi.org/10.1080/11250000109356423
Garamszegi, L. Z., Hegyi, G., Heylen, D., Ninni, P., de Lope, F., Eens, M., & Møller, A. P.
(2006). The design of complex sexual traits in male barn swallows: associations
between signal attributes. *Journal of Evolutionary Biology*, 19(6), 2052–2066.
Garamszegi, L. Z., Heylen, D., Møller, A. P., Eens, M., & de Lope, F. (2005). Age-dependent
health status and song characteristics in the barn swallow. *Behavioral Ecology: Official
Journal of the International Society for Behavioral Ecology*, 16(3), 580–591.

Garamszegi, L. Z., Pavlova, D. Z., Eens, M., & Møller, A. P. (2006). The evolution of song in
675   female birds in Europe. *Behavioral Ecology: Official Journal of the International Society*
676   *for Behavioral Ecology*, 18(1), 86–96.
677   Gil, D., & Gahr, M. (2002). The honesty of bird song: multiple constraints for multiple traits.
678   *Trends in Ecology & Evolution*, 17(3), 133–141.
679   Grafe, T. U., Ulmar Grafe, T., & Bitz, J. H. (2004). Functions of duetting in the tropical boubou,
680   Laniarius aethiopicus: territorial defence and mutual mate guarding. In *Animal Behaviour*
681   (Vol. 68, Issue 1, pp. 193–201). https://doi.org/10.1016/j.anbehav.2003.11.003
682   Graham, B. A., Heath, D. D., Walter, R. P., Mark, M. M., & Mennill, D. J. (2018). Parallel
683   evolutionary forces influence the evolution of male and female songs in a tropical
684   songbird. In *Journal of Evolutionary Biology* (Vol. 31, Issue 7, pp. 979–994).
685   https://doi.org/10.1111/jeb.13279
686   Graham, B. A., Sandoval, L., Dabelsteen, T., & Mennill, D. J. (2017). A test of the Acoustic
687   Adaptation Hypothesis in three types of tropical forest: degradation of male and female
688   Rufous-and-white Wren songs. In *Bioacoustics* (Vol. 26, Issue 1, pp. 37–61).
689   https://doi.org/10.1080/09524622.2016.1181574
690   Hare, Robin M., and Leigh W. Simmons. 2019. "Sexual Selection and Its Evolutionary
691   Consequences in Female Animals." *Biological Reviews of the Cambridge Philosophical*
692   *Society* 94 (3): 929–56.
693   Hasiniaina, A. F., Radespiel, U., Kessler, S. E., Rina Evasoa, M., Rasoloharijaona, S.,
694   Randrianambinina, B., Zimmermann, E., Schmidt, S., & Scheumann, M. (2020).
695   Evolutionary significance of the variation in acoustic communication of a cryptic
696   nocturnal primate radiation (*Microcebus spp.*). *Ecology and Evolution*, 10(8), 3784–3797.
697   Hathcock, T. J., & Benedict, L. (2018). Conspecific challenges provoke female canyon wrens to
698   sing but not to duet. In *Behavioral Ecology and Sociobiology* (Vol. 72, Issue 12).
699   https://doi.org/10.1007/s00265-018-2625-0
700   Hebets, E. A., Barron, A. B., Balakrishnan, C. N., Hauber, M. E., Mason, P. H., & Hoke, K. L.
701   (2016). A systems approach to animal communication. *Proceedings. Biological*
702   *Sciences/ The Royal Society*, 283(1826), 20152889.
703   Hebets, E. A., & Papaj, D. R. (2004). Complex signal function: developing a framework of
704   testable hypotheses. *Behavioral Ecology and Sociobiology*, 57(3), 197–214.
705   Hedrick, A. V., & Temeles, E. J. (1989). The evolution of sexual differences in animals:
706   Hypotheses and tests. *Trends in Ecology & Evolution*, 4(5), 136–138.
707   Hofmann, C. M., Cronin, T. W., & Omland, K. E. (2008). Evolution of sexual dichromatism. 1.
708   Convergent losses of elaborate female coloration in new world orioles (Icterus spp.). In
709   *The Auk* (Vol. 125, Issue 4, pp. 778–789). https://doi.org/10.1525/auk.2008.07112
710   Huang, H., & Rabosky, D. L. (2014). Sexual selection and diversification: reexamining the
711   correlation between dichromatism and speciation rate in birds. *The American Naturalist*,
712   184(5), E101–E114.
713   Hubbard, J. K., Hund, A. K., Levin, I. I., McGraw, K. J., Wilkins, M. R., & Safran, R. J. (2017).
714   The importance of cross-validation, accuracy, and precision for measuring plumage
715   color: A comment on Vaquero-Alba et al. (2016). *The Auk*, 134(1), 34–38.
716   Irwin, D. E., Bensch, S., & Price, T. D. (2001). Speciation in a ring. *Nature*, 409(6818), 333–337.
717   Jawor, J. M., Gray, N., Beall, S. M., & Breitwisch, R. (2004). Multiple ornaments correlate with
718   aspects of condition and behaviour in female northern cardinals, Cardinalis cardinalis.
719   *Animal Behaviour*, 67(5), 875–882.
720   Jennions, M. D., Kahn, A. T., Kelly, C. D., & Kokko, H. (2012). Meta-analysis and sexual
721   selection: past studies and future possibilities. Evolutionary Ecology, 26(5), 1119–1151.
722   Jiggins, C. D., Estrada, C., & Rodrigues, A. (2004). Mimicry and the evolution of premating
723   isolation in Heliconius melpomene Linnaeus. *Journal of Evolutionary Biology*, 17(3),
724   680–691.

Johnson, A. E., Jordan Price, J., & Pruett-Jones, S. (2013). Different modes of evolution in
males and females generate dichromatism in fairy-wrens (Maluridae). *Ecology and*
Evolution, 3(9), 3030–3046.
Kalinowski, S. T., Taper, M. L., & Marshall, T. C. (2007). Revising how the computer program
CERVUS accommodates genotyping error increases success in paternity assignment.
Molecular Ecology, 16(5), 1099–1106.
Keen, S. C., Odom, K. J., Kohn, G., Wright, T., Webster, M. S. and Araya-Salas, M. (In review).
A machine-learning approach for quantifying element diversity in animal vocalizations.
Methods in Ecology and Evolution
Kingsolver, J. G., Diamond, S. E., Siepielski, A. M., & Carlson, S. M. (2012). Synthetic analyses
of phenotypic selection in natural populations: lessons, limitations and future directions.
In *Evolutionary Ecology* (Vol. 26, Issue 5, pp. 1101–1118).
https://doi.org/10.1007/s10682-012-9563-5
Kleindorfer, S., Evans, C., & Mahr, K. (2016). Female in-nest chatter song increases predation.
Biology Letters, 12(1), 20150513.
Kraaijeveld, K., Kraaijeveld-Smit, F. J. L., & Komdeur, J. (2007). The evolution of mutual
ornamentation. In *Animal Behaviour* (Vol. 74, Issue 4, pp. 657–677).
https://doi.org/10.1016/j.anbehav.2006.12.027
Kraaijeveld, K., Kraaijeveld-Smit, F. J. L., & Maan, M. E. (2011). Sexual selection and
speciation: the comparative evidence revisited. *Biological Reviews of the Cambridge*
Philosophical Society, 86(2), 367–377.
Krieg, C. A., & Getty, T. (2016). Not just for males: females use song against male and female
rivals in a temperate zone songbird. *Animal Behaviour*, 113, 39–47.
Kuhn, M., Chow, F., & Wickham, H. (2019). rsample: General Resampling Infrastructure
(Version 0.0.5) [Computer software]. https://CRAN.R-project.org/package=rsample
Lande, R. (1980). Sexual differences, sexual selection, and adaptation in polygenic characters.
Evolution; International Journal of Organic Evolution, 34(2), 292–305.
Langmore, N. E. (1998). Functions of duet and solo songs of female birds. *Trends in Ecology &*
Evolution, 13(4), 136–140.
Lee, K.-H., Shaner, P.-J. L., Lin, Y.-P., & Lin, S.-M. (2016). Geographic variation in
advertisement calls of a Microhylid frog - testing the role of drift and ecology. *Ecology*
and Evolution, 6(10), 3289–3298.
Levin, R. N. (1996). Song behaviour and reproductive strategies in a duetting wren, *Thryothorus*
nigricapillus: I. Removal experiments. In *Animal Behaviour* (Vol. 52, Issue 6, pp. 1093–
1106). https://doi.org/10.1006/anbe.1996.0257
Magoolagan, L., Mawby, P. J., Whitehead, F. A., & Sharp, S. P. (2019). The structure and
context of male and female song in White-throated Dippers. In *Journal of Ornithology*
(Vol. 160, Issue 1, pp. 195–205). https://doi.org/10.1007/s10336-018-1599-z
McCoy, J. K., Kelly McCoy, J., James Harmon, H., Baird, T. A., & Fox, S. F. (1997). Geographic
Variation in Sexual Dichromatism in the Collared Lizard, *Crotaphytus collaris* (Sauria:
Crotaphytidae). In *Copeia* (Vol. 1997, Issue 3, p. 565). https://doi.org/10.2307/1447560
McGraw, K. J., Safran, R. J., & Wakamatsu, K. (2005). How feather colour reflects its melanin
content. In *Functional Ecology* (Vol. 19, Issue 5, pp. 816–821).
https://doi.org/10.1111/j.1365-2435.2005.01032.x
McLean, C. A., & Stuart-Fox, D. (2014). Geographic variation in animal colour polymorphisms
and its role in speciation. *Biological Reviews of the Cambridge Philosophical Society*,
89(4), 860–873.
Mennill, D. J., & Rogers, A. C. (2006). Whip it good! Geographic consistency in male songs and
variability in female songs of the duetting eastern whipbird Psophodes olivaceus. In
Journal of Avian Biology (Vol. 37, Issue 1, pp. 93–100).
https://doi.org/10.1111/j.2006.0908-8857.03548.x
Miller, C. W., McDonald, G. C., & Moore, A. J. (2016). The tale of the shrinking weapon:
seasonal changes in nutrition affect weapon size and sexual differences, but not

778   contemporary evolution. In *Journal of Evolutionary Biology* (Vol. 29, Issue 11, pp. 2266–
779   2275). https://doi.org/10.1111/jeb.12954
780 Møller, A. P. (1988). Female choice selects for male sexual tail ornaments in the monogamous
781   swallow. In *Nature* (Vol. 332, Issue 6165, pp. 640–642).
782   https://doi.org/10.1038/332640a0
783 Ng, T. P. T., Rolán-Alvarez, E., Dahlén, S. S., Davies, M. S., Estévez, D., Stafford, R., &
784   Williams, G. A. (2019). The causal relationship between sexual selection and sexual size
785   differences in marine gastropods. In *Animal Behaviour* (Vol. 148, pp. 53–62).
786   https://doi.org/10.1016/j.anbehav.2018.12.005
787 Nowicki, S., Searcy, W. A., & Peters, S. (2002). Brain development, song learning and mate
788   choice in birds: a review and experimental test of the "nutritional stress hypothesis."
789   *Journal of Comparative Physiology A: Neuroethology, Sensory, Neural, and Behavioral*
790   *Physiology*, 188(11-12), 1003–1014.
791 Obara, Y., Ozawa, G., & Fukano, Y. (2008). Geographic variation in ultraviolet reflectance of the
792   wings of the female cabbage butterfly, *Pieris rapae*. *Zoological Science*, 25(11), 1106–
793   1110.
794 Odom, K. J., & Benedict, L. (2018). A call to document female bird songs: Applications for
795   diverse fields. In *The Auk* (Vol. 135, Issue 2, pp. 314–325). https://doi.org/10.1642/auk-
796   17-183.1
797 Odom, K. J., Hall, M. L., Riebel, K., Omland, K. E., & Langmore, N. E. (2014). Female song is
798   widespread and ancestral in songbirds. *Nature Communications*, 5, 3379.
799 Odom, K. J., & Mennill, D. J. (2012). Inconsistent geographic variation in the calls and duets of
800   Barred Owls (Strix varia) across an area of genetic introgression. *The Auk*, 129(3), 387–
801   398.
802 Orme, D., Freckleton, R., Thomas, G., Petzoldt, T., Fritz, S., Isaac, N., & and Pearse, W.
803   (2018). *caper: Comparative Analyses of Phylogenetics and Evolution in R (Version*
804   *1.0.1)* [Computer software]. https://CRAN.R-project.org/package=caper
805 Pap, P. L., Fülöp, A., Adamkova, M., Cepak, J., Michalkova, R., Safran, R. J., Stermin, A. N.,
806   Tomasek, O., Vágási, C. I., Vincze, O., Wilkins, M. R., & Albrecht, T. (2019). Selection
807   on multiple sexual signals in two Central and Eastern European populations of the barn
808   swallow. In *Ecology and Evolution*. https://doi.org/10.1002/ece3.5629
809 Partan, S. R., & Marler, P. (2005). Issues in the classification of multimodal communication
810   signals. *The American Naturalist*, 166(2), 231–245.
811 Pieplow, N. (2017). *Peterson Field Guide to Bird Sounds of Eastern North America*. Houghton
812   Mifflin Harcourt.
813 Podos, J., & Warren, P. S. (2007). The Evolution of Geographic Variation in Birdsong. In
814   *Advances in the Study of Behavior* (pp. 403–458). https://doi.org/10.1016/s0065-
815   3454(07)37009-5
816 Price, J. J. & Eaton, M. D. (2014). Reconstructing the evolution of sexual dichromatism: current
817   color diversity does not reflect past rates of male and female change. In *Evolution* (Vol.
818   68, Issue 7, pp. 2026–2037). https://doi.org/10.1111/evo.12417
819 Price, J. J. (2015). Rethinking our assumptions about the evolution of bird song and other
820   sexually dimorphic signals. In *Frontiers in Ecology and Evolution* (Vol. 3).
821   https://doi.org/10.3389/fevo.2015.00040
822 Price, J. J. (2019). Sex differences in song and plumage color do not evolve through sexual
823   selection alone: new insights from recent research. In *Journal of Ornithology* (Vol. 160,
824   Issue 4, pp. 1213–1219). https://doi.org/10.1007/s10336-019-01681-8
825 Price, J. J., Lanyon, S. M., & Omland, K. E. (2009). Losses of female song with changes from
826   tropical to temperate breeding in the New World blackbirds. *Proceedings. Biological*
827   *Sciences / The Royal Society*, 276(1664), 1971–1980.
828 Price, T. D. (1984). The Evolution of Sexual Size differences in Darwin's Finches. The American
829   Naturalist, 123(4), 500–518.

Purvis, A., & Rambaut, A. (1995). Comparative analysis by independent contrasts (CAIC): an Apple Macintosh application for analysing comparative data. *Computer Applications in the Biosciences: CABIOS*, 11(3), 247–251.

R Core Team. (2019). *R: A Language and Environment for Statistical Computing (Version 3.6.0)* [Computer software]. https://www.R-project.org/

Riebel, K. (2016). Understanding Sex Differences in Form and Function of Bird Song: The Importance of Studying Song Learning Processes. *Frontiers in Ecology and Evolution*, 4. https://doi.org/10.3389/fevo.2016.00062

Riebel, K., Hall, M. L., & Langmore, N. E. (2005). Female songbirds still struggling to be heard [Review of Female songbirds still struggling to be heard]. *Trends in Ecology & Evolution*, 20(8), 419–420.

Riebel, K., Odom, K. J., Langmore, N. E., & Hall, M. L. (2019). New insights from female bird song: towards an integrated approach to studying male and female communication roles. *Biology Letters*, 15(4), 20190059.

Roberts, N. S., & Mendelson, T. C. (2017). Male mate choice contributes to behavioural isolation in sexually dimorphic fish with traditional sex roles. *Animal Behaviour*, 130, 1–7.

Roberts, N. S., & Mendelson, T. C. (2020). Identifying female phenotypes that promote behavioral isolation in a sexually dimorphic species of fish (*Etheostoma zonale*). In *bioRxiv* (No. biorxiv;2020.04.20.051714v1; p. 224).

Robinson, M. R., Pilkington, J. G., Clutton-Brock, T. H., Pemberton, J. M., & Kruuk, L. E. B. (2008). Environmental heterogeneity generates fluctuating selection on a secondary sexual trait. *Current Biology: CB*, 18(10), 751–757.

Rodríguez, R. L., Ramaswamy, K., & Cocroft, R. B. (2006). Evidence that female preferences have shaped male signal evolution in a clade of specialized plant-feeding insects. *Proceedings. Biological Sciences / The Royal Society*, 273(1601), 2585–2593.

Rose, E. M., Coss, D. A., Haines, C. D., Danquah, S. A., Studds, C. E., & Omland, K. E. (2019). Why do females sing?—pair communication and other song functions in eastern bluebirds. In *Behavioral Ecology* (Vol. 30, Issue 6, pp. 1653–1661). https://doi.org/10.1093/beheco/arz130

Rosvall, K. A. (2011). Intrasexual competition in females: evidence for sexual selection? *Behavioral Ecology: Official Journal of the International Society for Behavioral Ecology*, 22(6), 1131–1140.

Roulin, A. (2003). Geographic variation in sexual differences in the barn owl Tyto alba: a role for direct selection or genetic correlation? *In Journal of Avian Biology* (Vol. 34, Issue 3, pp. 251–258). https://doi.org/10.1034/j.1600-048x.2003.03022.x

Ryan, M. J., Akre, K. L., Baugh, A. T., Bernal, X. E., Lea, A. M., Leslie, C., Still, M. B., Wylie, D. C., & Rand, A. S. (2019). Nineteen Years of Consistently Positive and Strong Female Mate Preferences despite Individual Variation. *The American Naturalist*, 194(2), 125–134.

Safran, R. J., McGraw, K. (2004). Plumage coloration, not length or symmetry of tail-streamers, is a sexually selected trait in North American barn swallows. In *Behavioral Ecology* (Vol. 15, Issue 3, pp. 455–461). https://doi.org/10.1093/beheco/arh035

Safran, R. J., Neuman, C. R., McGraw, K. J., & Lovette, I. J. (2005). Dynamic paternity allocation as a function of male plumage color in barn swallows. *Science*, 309(5744), 2210–2212.

Safran, R. J., Adelman, J. S., McGraw, K. J., & Hau, M. (2008). Sexual signal exaggeration affects physiological state in male barn swallows. In *Current Biology* (Vol. 18, Issue 11, pp. R461–R462). https://doi.org/10.1016/j.cub.2008.03.031

Safran, R. J., Vortman, Y., Jenkins, B. R., Hubbard, J. K., Wilkins, M. R., Bradley, R. J., & Lotem, A. (2016). The maintenance of phenotypic divergence through sexual selection: An experimental study in barn swallows Hirundo rustica. *Evolution; International Journal of Organic Evolution*, 70(9), 2074–2084.

Saino, N., Romano, M., Sacchi, R., Ninni, P., Galeotti, P., & Møller, A. P. (2003). Do male barn
swallows (Hirundo rustica) experience a trade-off between the expression of multiple
sexual signals? *Behavioral Ecology and Sociobiology*, 54(5), 465–471.
Searcy, W. A., & Andersson, M. (1986). Sexual Selection and the Evolution of Song. *Annual
Review of Ecology and Systematics*, 17(1), 507–533.
Shultz, Allison J., and Kevin J. Burns. 2017. "The Role of Sexual and Natural Selection in
Shaping Patterns of Sexual Dichromatism in the Largest Family of Songbirds (Aves:
Thraupidae)." *Evolution; International Journal of Organic Evolution* 71 (4): 1061–74.
Sibley, D. A. (2014). *The Sibley Guide to Birds*. Knopf.
Slabbekoorn, H., & Smith, T. B. (2002). Bird song, ecology and speciation. In *Philosophical
Transactions of the Royal Society of London. Series B: Biological Sciences* (Vol. 357,
Issue 1420, pp. 493–503). https://doi.org/10.1098/rstb.2001.1056
Smith, C. C. R., Flaxman, S. M., Scordato, E. S. C., Kane, N. C., Hund, A. K., Sheta, B. M., &
Safran, R. J. (2018). Demographic inference in barn swallows using whole-genome data
shows signal for bottleneck and subspecies differentiation during the Holocene.
Molecular Ecology, 27(21), 4200–4212.
Snyder, K. T., & Creanza, N. (2019). Polygyny is linked to accelerated birdsong evolution but
not to larger song repertoires. *Nature Communications*, 10(1), 884.
Sosa-López, J. R., Martínez Gómez, J. E., & Mennill, D. J. (2016). Divergence in mating signals
correlates with genetic distance and behavioural responses to playback. *Journal of
Evolutionary Biology*, 29(2), 306–318.
Steele, D. B., Siepielski, A. M., & McPEEK, M. A. (2011). Sexual selection and temporal
phenotypic variation in a damselfly population. In *Journal of Evolutionary Biology* (Vol.
24, Issue 7, pp. 1517–1532). https://doi.org/10.1111/j.1420-9101.2011.02284.x
Stoddard, M.C. & Prum, R.O. 2008. Evolution of Avian Plumage Color in a Tetrahedral Color
Space: A Phylogenetic Analysis of New World Buntings. *The American Naturalist*. 171:
755–776.
Stokes, D., & Stokes, L. (2010). *The Stokes Field Guide to the Birds of North America*. Little,
Brown.
Svensson, E. I., & Waller, J. T. (2013). Ecology and sexual selection: evolution of wing
pigmentation in calopterygid damselflies in relation to latitude, sexual differences, and
speciation. *The American Naturalist*, 182(5), E174–E195.
Taff, C. C., Littrell, K. A., & Freeman-Gallant, C. R. (2012). Female Song in the Common
Yellowthroat. In *The Wilson Journal of Ornithology* (Vol. 124, Issue 2, pp. 370–374).
https://doi.org/10.1676/11-182.1
Tobias, J. A., Gamarra-Toledo, V., García-Olaechea, D., Pulgarín, P. C., & Seddon, N. (2011).
Year-round resource defence and the evolution of male and female song in suboscine
birds: social armaments are mutual ornaments. In *Journal of Evolutionary Biology* (Vol.
24, Issue 10, pp. 2118–2138). https://doi.org/10.1111/j.1420-9101.2011.02345.x
Tobias, J. A., & Seddon, N. (2009). Signal jamming mediates sexual conflict in a duetting bird.
Current Biology: CB, 19(7), 577–582.
Tuomaala, M., Kaitala, A., & Rutowski, R. L. (2012). Females show greater changes in wing
colour with latitude than males in the green-veined white butterfly, *Pieris napi*
(Lepidoptera: Pieridae). In *Biological Journal of the Linnean Society* (Vol. 107, Issue 4,
pp. 899–909). https://doi.org/10.1111/j.1095-8312.2012.01996.x
Velásquez, N. A. (2014). Geographic variation in acoustic communication in anurans and its
neuroethological implications. *Journal of Physiology*, Paris, 108(2-3), 167–173.
Webb, Wesley H., Dianne H. Brunton, J. David Aguirre, Daniel B. Thomas, Mihai Valcu, and
James Dale. 2016. "Female Song Occurs in Songbirds with More Elaborate Female
Coloration and Reduced Sexual Dichromatism." *Frontiers in Ecology and Evolution*.
https://doi.org/10.3389/fevo.2016.00022.

Wiens, J. J. (2001). Widespread loss of sexually selected traits: how the peacock lost its spots.
In *Trends in Ecology & Evolution* (Vol. 16, Issue 9, pp. 517–523).
https://doi.org/10.1016/s0169-5347(01)02217-0
Wilkins, M. R., Scordato, E. S. C., Semenov, G. A., Karaardiç, H., Shizuka, D., Rubtsov, A.,
Pap, P. L., Shen, S.-F., & Safran, R. J. (2018). Global song divergence in barn swallows
(Hirundo rustica): exploring the roles of genetic, geographical and climatic distance in
sympatry and allopatry. In *Biological Journal of the Linnean Society* (Vol. 123, Issue 4,
pp. 825–849). https://doi.org/10.1093/biolinnean/bly012
Wilkins, M. R., Seddon, N., & Safran, R. J. (2013). Evolutionary divergence in acoustic signals:
causes and consequences. *Trends in Ecology & Evolution*, 28(3), 156–166.
Wilkins, M. R., Shizuka, D., Joseph, M. B., Hubbard, J. K., & Safran, R. J. (2015). Multimodal
signalling in the North American barn swallow: a phenotype network approach.
Proceedings. Biological Sciences / The Royal Society, 282(1816), 20151574.
Yasukawa, K. (1989). The costs and benefits of a vocal signal: the nest-associated "Chit" of the
female red-winged blackbird, Agelaius phoeniceus. In *Animal Behaviour* (Vol. 38, Issue
5, pp. 866–874). https://doi.org/10.1016/s0003-3472(89)80118-6
Yasukawa, K. (1990). Does the "teer" vocalization deter prospecting female red-winged
blackbirds? *Behavioral Ecology and Sociobiology*, 26(6), 421–426.
951

Electronic Supplement

953 **Video S1.** "Female Barn Swallows Sing!": provides detail and examples about timing and
954 potential functions of female song. Accessible at: https://vimeo.com/424642268

Supplementary Methods and Background

Acoustics

Clarifying notes on female song

Popular field guide entries for barn swallows do not specify which sex sings, and it can be
assumed the song description refers to males (Sibley, 2014; Stokes & Stokes, 2010). In
contrast, the *Peterson Field Guide to Bird Sounds of Eastern North America* states "females
also reported to sing" (Pieplow, 2017) and the entry in *Birds of the World* explicitly states that
both sexes sing the "twitter-warble" song, i.e. typical male song (Brown & Brown, 2020). These
latter interpretations may stem from Samuel's description of the barn swallow vocal repertoire,
in which he states that the courtship song "is occasionally given by the female" (Samuel, 1971).
He also describes a "whine call," which may be the same as the copulation call ("waeae-
waeae") referred to in the *Birds of the World* (Brown & Brown, 2020), which may refer to what
we call song in this study. In lieu of the original recordings or a high-quality spectrogram, it is
difficult to assess which female vocalizations are being described in previous studies. In Møller's
Sexual Selection and the Barn Swallow (1994) there is no mention of female song or
vocalizations at all. In fact the copulation call is ascribed to the male (p78). We believe this
vocalization, the main phrase of which we transliterate as "dwidle dwight," is more accurately
described as a song rather than a call (sensu Marler, 2004), since it is multisyllabic, structurally
complex, and seems to have a function in mate attraction, pair bonding, and competition, based
on observations (see supplementary video).

General acoustic analysis details

Prior to song analysis, all songs were standardized to a sample rate of 44.1 kHz and a bit depth
of 16. We used Raven 1.5 (Bioacoustics Research Program, 2011) to select and extract
measurements for each element in each song. We defined an element as a single, continuous
trace on a spectrogram, separated from other elements by a visible break in time or an extreme
shift in overall harmonic structure (occasionally, structurally distinct barn swallow elements were
separated by very short breaks in time, so this additional criterion helped us distinguish among
elements). For pure tone elements, we selected and compared the fundamental frequency of
the element, whereas for harmonically rich elements, we selected the fundamental to the
dominant frequencies. We did substantial visual checking to be sure that elements were
separated and measured consistently across all songs. We used the following Hann window
settings with a sample size of 512 (DFT), hop size of 256, and 50 percent overlap in Raven v1.5
for selecting and measuring elements (Bioacoustics Research Program, 2011).

Element-level acoustic parameters

For every element in all songs we extracted a large number of energy-based measurements
calculated from the cumulative amplitude of the spectrum or envelope using Raven Bioacoustic
Analysis Software v1.5 and the warbleR package in R (Araya-Salas & Smith-Vidaurre, 2017;
Bioacoustics Research Program, 2011). We removed any highly correlated variables with a
Pearson correlation coefficient greater than 0.95. This resulted in the following acoustic
variables measured for each element using each bioacoustics program: Raven measurements –
peak frequency, frequency 5%, delta frequency, interquartile range bandwidth, time 5%,
interquartile range duration, duration 90%, aggregate entropy, average entropy, average power,
energy; warbleR measurements – standard deviation of frequency, mean dominant frequency,
mean peak frequency, minimum dominant frequency, maximum dominant frequency, 1st
quartile frequency, interquartile frequency range, dominant frequency range, dominant
frequency slope, 1st quartile time, spectral entropy, time entropy, entropy, spectral flatness,
modulation index, skewness, and kurtosis. See Raven and warbleR manuals for definitions of
each parameter (Araya-Salas & Smith-Vidaurre, 2017; Bioacoustics Research Program, 2011).

Song-level acoustic parameters – Using the warbleR package in R, we extracted the following
acoustic parameters for each song: mean frequency, mean peak frequency, median frequency,
frequency range, dominant frequency range, 1st quartile frequency, 3rd quartile frequency,
interquartile frequency range, median time, 1st quartile time, 3rd quartile time, entropy, average
element duration per song, song duration, number of elements, element rate, and gap duration.
See the warbleR vignette for descriptions of each parameter (Araya-Salas & Smith-Vidaurre,
2017).

Element diversity calculations – Using the element-level acoustic parameters, we estimated
element diversity for each song as the area encompassing the elements for that song in a multi-
dimensional feature space. To do this, we first created an element-level acoustic space by
inputting the element-level acoustic parameters for every element for all songs into an
unsupervised random forest. We used the 'randomForest' function in the R package
{randomForest} with the following specifications: 10000 trees, minimal node size of 1, Gini
impurity index as split rule, 5 randomly sampled variables at each split and out-of-bag proximity
(Liaw & Wiener, 2002). This produced a proximity matrix, which we transformed into a set of five
vectors using classic multidimensional scaling, using the 'cmdscale' function in the {stats} R
package. We then calculated and extracted minimum convex polygons to calculate the area
surrounding all elements for each song within this multidimensional space using the function
'mcp' in the {adehabitatHR} R package (Calenge, 2006). Therefore, each area is an estimate of
the variability, or diversity, of elements for a given song within an acoustic feature space of all
barn swallow element measurements.

Color Analysis

Hue Saturation Brightness (HSB) Color Measures

Methods reprinted from Wilkins et al. (2015), Online Appendix S1.

Feather samples from four ventral patches (throat, breast, belly, vent) were taped to a standard
white card background so that they overlapped as they do on the body of a bird. The color of
each patch was measured using a spectrometer (USB 4000, Ocean Optics), pulsed xenon light
(PX-2, Ocean Optics) and SpectraSuite software (v2.0.151). The probe was held at 90 degrees
to the feather surface at a distance such that a 2.5 mm diameter of the surface was illuminated
and measured. Each sample was measured three times, lifting the probe between

measurements, and averaged. Each measurement was an average of 20 scans of the
spectrometer. From the generated spectra, we calculated standard color descriptors: i) average
brightness, ii) hue, and iii) red chroma. Average brightness was calculated as the average
percent reflectance between 300 and 700 nm, hue was calculated as the wavelength that
corresponds to where the slope of the curve is steepest between 550 and 700 nm, and red
chroma was calculated as the proportion of light reflected in the red range (600 to 700 nm)
relative to the entire range (300 to 700 nm).

Comparison of HSB to TCS color metrics

Though we have opted to use HSB methods for the current manuscript for easier comparison to
most published barn swallow studies, we also calculated color measures in the tetrahedral color
space (TCS) for comparison. TCS measurements follow Wilkins, et al. (2016). Briefly, we used
the 'tcs' function in the R package {pavo} v. 0.5-5 (Maia et al. 2013). These color metrics
describe the relative stimulation of the four bird cone channels (u, s, m, and l), relative to the
achromatic origin. Thus, each color can be described by a vector and represented by θ
(horizontal angle), Φ (vertical angle), and r (vector length), with θ and Φ reflecting hue. Because
the color space is a tetrahedron, rather than a sphere, the maximum value of r varies by hue. As
such, we use rA (achieved chroma, r/rmax) as a relative measure of saturation. Additionally, we
calculated brilliance, or the averaged reflectance values for each measured wavelength
between 300 and 700 nm. Collectively, our color metrics, θ, Φ, rA, and brilliance describe the
color of each ventral plumage patch, quantified using the average avian UV visual model, as
defined in {pavo}.

To assess the match between traditional HSB color metrics and the avian visual model color
measures, we explored Spearman's rho correlations between all color metrics calculated for the
breast patch for the 27 individuals in our study with complete feather samples. TCS brilliance is
synonymous with HSB average brightness. HSB chroma is very strongly correlated with the
most relevant TCS metric (rA: achieved chroma). Hue shows a moderate negative correlation
with TCS brilliance, though a direct comparison to TCS hue is complicated, due to two angular
measures (θ and Φ) of hue in TCS space. Overall, the HSB and TCS show strong
correspondence, suggesting that our reported measures of color represent discernible color
variation for live birds, which is supported by experimental and biochemical evidence.

1064

1065

Paternity Analyses

Methods reprinted from Wilkins et al. (2015), Online Appendix S1.

The six microsatellite markers used for paternity analyses were Escu6: (Hanotte et al. 1994);
Ltr6: (McDonald & Potts 1994); Pocc6: (Bensch et al. 1997); and Hir11, Hir19, and Hir20:
(Tsyusko et al. 2007)). Reaction conditions for pooled Escu6, Ltr6, Hir20, and Hir11 primers
consisted of a 10 ul solution with 50-100 ng DNA, 0.12 mM of each labeled forward primer, 0.12
mM of each reverse primer, 200 M each dNTP, 3.25 mM MgCl2, 1x PCR Buffer, 0.15 units Taq
polymerase (New England Biolabs, Massachusetts, U.S.A.), and were amplified with the
following protocol: initial denaturation step of 94°C for 1 minute, followed by 10 cycles of 94°C
for 30 s, 55°C for 30 s, and 72°C for 45 s, with an additional 25 cycles starting at 87°C for 30 s
instead of 94°C, and completed with a final extension at 72 °C for 3 min. The Pocc6 reaction

was modified from the above conditions by using 1.25 mM $MgCl_2$, and modified for the Hir19
reaction with 3 mM MgCl2 and 0.2 mM of each forward and reverse primer. The PCR
amplification protocol for Pocc6 and Hir19 was similar to the pooled loci protocol with the
exception that 60°C was used for the annealing temperature. Amplified PCR products
containing the fluorescently-labeled forward primer were detected using an ABI3730 DNA
analyzer (ABI, Inc.). Allele peaks were manually called using Peak Scanner v1.0 (Applied
Biosystems) in order to minimize genotyping error associated with irregular PCR products. For
the paternity analysis simulation in CERVUS, we left the proportion of loci typed at 0.959
(determined from our data) and the proportion of loci mistyped at the default 0.01, and ran the
program for 10,000 iterations. For each male, we set the female seen brooding eggs as the
known mother and considered an offspring as extra-pair when offspring-mother-father trio
confidence did not reach the 95% level. We adopted this approach, rather than a threshold
number of parent-offspring mismatches for paternity exclusions because confidence levels are
derived from likelihood equations which take into account potential genotyping errors.

1091

1092

Supplemental References:

Araya-Salas, M., & Smith-Vidaurre, G. (2017). warbleR: an R package to streamline analysis of

    animal acoustic signals. In *Methods in Ecology and Evolution* (Vol. 8, Issue 2, pp. 184–

    191). https://doi.org/10.1111/2041-210x.12624

Bensch, S., Price, T., Kohn, J. & others. (1997). Isolation and characterization of microsatellite

    loci in a Phylloscopus warbler. Mol. Ecol., 6, 91–92.

Bioacoustics Research Program. (2011). *Raven Pro: Interactive Sound Analysis Software*

    (Version 1.5) [Computer software]. http://www.birds.cornell.edu/raven

Brown, M. B., & Brown, C. R. (2020). *Barn Swallow - Hirundo rustica*. Birds of the World.

    https://doi.org/10.2173/bow.barswa.01

Calenge, C. (2006). The package "adehabitat" for the R software: A tool for the analysis of

    space and habitat use by animals. In *Ecological Modelling* (Vol. 197, Issues 3-4, pp. 516–

    519). https://doi.org/10.1016/j.ecolmodel.2006.03.017

Hanotte, O., Zanon, C., Pugh, A., Greig, C., Dixon, A. & Burke, T. (1994). Isolation and

    characterization of microsatellite loci in a passerine bird: the reed bunting Emberiza

    schoeniclus. Mol. Ecol., 3, 529–530.

Liaw, A. L., & Wiener, M. (2002). Classification and regression by randomForest. *R News*, 2,

    18–22.

Maia, R., Eliason, C. M., Bitton, P.-P., Doucet, S. M., & Shawkey, M. D. (2013). pavo: an R

    package for the analysis, visualization and organization of spectral data. In Methods in

    Ecology and Evolution. https://doi.org/10.1111/2041-210x.12069

Marler, P. (2004). Bird calls: a cornucopia for communication. In *Nature's Music* (pp. 132–177).

    https://doi.org/10.1016/b978-012473070-0/50008-6

McDonald, D.B. & Potts, W.K. (1994). Cooperative display and relatedness among males in a

    lek-mating bird. Science (80-.)., 266, 1030–1032.

Møller, A. P. (1994). *Sexual Selection and the Barn Swallow*. Oxford University Press.

Pieplow, N. (2017). *Peterson Field Guide to Bird Sounds of Eastern North America*. Houghton

Mifflin Harcourt.

Samuel, D. E. (1971). Vocal Repertoires of Sympatric Barn and Cliff Swallows. In *The Auk* (Vol.

88, Issue 4, pp. 839–855). https://doi.org/10.2307/4083842

Sibley, D. A. (2014). *The Sibley Guide to Birds*. Knopf.

Stokes, D., & Stokes, L. (2010). *The Stokes Field Guide to the Birds of North America*. Little,

Brown.

Tsyusko, O. V., Peters, M.B., Hagen, C., Tuberville, T.D., Mousseau, T.A., Møller, A.P., et al.

(2007). Microsatellite markers isolated from barn swallows (Hirundo rustica). Mol. Ecol.

Notes, 7, 833–835.

Wilkins, M. R., Karaardıç, H., Vortman, Y., Parchman, T. L., Albrecht, T., Petrželková, A.,

Özkan, L., Pap, P. L., Hubbard, J. K., Hund, A. K., & Safran, R. J. (2016). Phenotypic

differentiation is associated with divergent sexual selection among closely related barn

swallow populations. In Journal of Evolutionary Biology (Vol. 29, Issue 12, pp. 2410–2421).

https://doi.org/10.1111/jeb.12965

1134
1135

Figure S1. Spearman's rank correlations between robustly dimorphic traits shown in Figure 4
and a relevant fitness metric–our measure of contemporary selection. The fitness metric for
males was seasonal within pair genetic paternity, and for females it was seasonal fledging
success. Orange 95% confidence ellipses and points are for females; blue ellipses and points
are for males. Plots are ordered (top left to bottom right) from lowest to highest correlation for
females.

1142

1143 **Figure S2. Spearman's correlations between our reported Hue-Saturation-Brightness**
1144 **(HSB) color measures and tetrahedral color space (TCS) color metrics. All measures are**
1145 **for the 27 individuals (both sexes) in our dataset with complete feather color measures.**
1146 **Only the breast patch is shown, since correlations**

1147

1148

1149

 Table S1. Trait definitions for all per individual visual and acoustic variables calculated for this
1151 study.

Trait Type	Trait	Definition
Acoustic	Entropy	Average spectrographic entropy: the product of time and energy distribution of the frequency spectrum; unitless, ranging from 0 (pure tone) to 1 (white noise)
Acoustic	Dominant Frequency Range	Average range of the dominant frequency measured across the song
Acoustic	Frequency Range	Average range of raw frequencies measured across song
Acoustic	Mean Peak Frequency	Average frequency with the highest energy from the mean frequency spectrum
Acoustic	Element Count	Number of song elements (syllables) per song
Acoustic	Element Duration	Average duration of elements per individual (in s)
Acoustic	Element Rate	Average rate of element production (number of elements divided by duration) per song (in elements/s)
Acoustic	Song Duration	Average duration of songs (in s)
Acoustic	Gap Duration	Average length of intervals between elements (in s)
Acoustic	Element Diversity	Average minimum convex polygon area surrounding the element-level acoustic space for each song
Acoustic	Frequency IQR	Average interquartile range for frequency (in kHz)
Acoustic	CV_Entropy	Coefficient of variation for spectrographic entropy
Acoustic	CV_Dom Frequency Range	Coefficient of variation for the range of the dominant frequency measured across the song
Acoustic	CV_Frequency Range	Coefficient of variation for raw frequencies measured across song
Acoustic	CV_Mean Peak Frequency	Coefficient of variation for frequency with the highest energy from the mean frequency spectrum
Acoustic	CV_Element Count	Coefficient of variation for the number of song elements (syllables) per song
Acoustic	CV_Element Duration	Coefficient of variation for duration of elements (in s)
Acoustic	CV_Element Rate	Coefficient of variation for rate of element production per song (in elements/s)
Acoustic	CV_Song Duration	Coefficient of variation for duration of songs (in s)
Acoustic	CV_Gap Duration	Coefficient of variation for length of intervals between

		elements (in s)
Acoustic	CV_Element Diversity	Coefficient of variation for the average minimum convex polygon area surrounding the element-level acoustic space for each song
Acoustic	CV_Frequency IQR	Coefficient of variation for interquartile range for frequency (in kHz)
Visual	Tail Length	Maximum length of two longest tail feathers (streamers), each of which was the average of 3 repeated measures (in mm)
Visual	Brightness	The average percent reflectance between 300 and 700 nm; lower values are darker; measured for Throat, Breast, Belly, and Vent
Visual	Hue	The wavelength at the maximum slope of reflectance spectrum (in nm); low values pale/yellowish; high values dark/reddish; measured for Throat, Breast, Belly, and Vent
Visual	Chroma	The proportion of light reflected in the red color range (600-700nm); higher values darker; measured for Throat, Breast, Belly, and Vent

1152
1153

Table S2. Female and male, sample sizes, means, standard deviations, sex difference (i.e.
1155 Cohen's D), and dimorphism 95% confidence intervals for 10,000 bootstraps. Sample sizes for
1156 coefficients of variation are lower for females, as three individuals only had a single song
1157 recorded. Bolded traits had dimorphism confidence intervals that did not overlap zero. Starred
1158 traits were selected for the final analysis in Figure 4, after eliminating the less biologically
1159 intuitive trait for redundant trait pairs with >|0.7| correlations.

1160

Trait	F_n	F_Mean (SD)	M_n	M_mean (SD)	Dimorphism	95% Conf Int
*Belly Brightness	**13**	**35.5 (7.62)**	**14**	**29.3 (7.62)**	**0.799**	**(0.0694, 1.31)**
Belly Chroma	13	0.421 (0.0434)	14	0.441 (0.0434)	-0.425	(-1.09, 0.333)
Belly Hue	13	608 (29)	14	626 (29)	-0.554	(-1.21, 0.19)
Breast Brightness	15	35.7 (9.39)	14	30.8 (9.39)	0.589	(-0.114, 1.1)
Breast Chroma	15	0.424 (0.0356)	14	0.446 (0.0356)	-0.577	(-1.1, 0.129)
Breast Hue	15	616 (23.3)	14	631 (23.3)	-0.62	(-1.16, 0.0884)
CV_Dom Frequency Range	**9**	**0.0902 (0.0483)**	**15**	**0.0363 (0.0483)**	**1.27**	**(0.362, 1.79)**
CV_Element Count	9	1.5 (2.24)	15	1.41 (2.24)	0.0468	(-0.962, 0.875)
*CV_Element Diversity	**9**	**2.34e-06 (2.01e-06)**	**15**	**8.61e-07 (2.01e-06)**	**0.954**	**(0.181, 1.57)**
CV_Element Duration	9	0.000904 (0.000889)	15	0.000465 (0.000889)	0.626	(-0.264, 1.34)
CV_Element Rate	9	0.149 (0.188)	15	0.0899 (0.188)	0.432	(-0.903, 1.15)
CV_Entropy	**9**	**0.000127 (0.000117)**	**15**	**0.000359 (0.000117)**	**-1.02**	**(-1.29, -0.454)**
CV_Frequency IQR	**9**	**0.069 (0.066)**	**15**	**0.0189 (0.066)**	**1.04**	**(0.517, 1.58)**
CV_Frequency Range	9	0.25 (0.573)	15	0.497 (0.573)	-0.319	(-0.82, 0.582)
CV_Gap Duration	9	0.00152 (0.0015)	15	0.000986 (0.0015)	0.454	(-0.546, 1.15)
CV_Mean Peak Frequency	**9**	**0.0499 (0.0408)**	**15**	**0.0099 (0.0408)**	**1.37**	**(0.801, 1.77)**
CV_Song Duration	9	0.117 (0.139)	15	0.151 (0.139)	-0.199	(-0.829, 0.774)
Dom Frequency Range	12	2.21 (0.303)	15	2.16 (0.303)	0.172	(-0.608, 0.958)

Trait	F_n	F_Mean (SD)	M_n	M_mean (SD)	Dimorphism	95% Conf Int
Element Count	**12**	**15.4 (6.28)**	**15**	**43.6 (6.28)**	**-3.79**	**(-1.88, -1.65)**
***Element Diversity**	**12**	**1.53e-05 (8.21e-06)**	**15**	**3.44e-05 (8.21e-06)**	**-3.11**	**(-1.88, -1.53)**
***Element Duration**	**12**	**0.0667 (0.00862)**	**15**	**0.0615 (0.00862)**	**0.778**	**(0.0664, 1.28)**
Element Rate	12	10.3 (0.914)	15	10.7 (0.914)	-0.516	(-1.16, 0.27)
***Entropy**	**12**	**0.831 (0.0121)**	**15**	**0.792 (0.0121)**	**3.27**	**(1.54, 1.85)**
***Frequency IQR**	**12**	**0.916 (0.189)**	**15**	**1.04 (0.189)**	**-0.816**	**(-1.45, -0.0324)**
***Frequency Range**	**12**	**7.11 (2.49)**	**15**	**9.46 (2.49)**	**-1.2**	**(-1.77, -0.264)**
Gap Duration	12	0.0347 (0.00477)	15	0.0338 (0.00477)	0.21	(-0.583, 0.994)
***Mean Peak Frequency**	**12**	**3.37 (0.299)**	**15**	**3.64 (0.299)**	**-0.881**	**(-1.45, -0.125)**
Song Duration	**12**	**1.49 (0.543)**	**15**	**4.12 (0.543)**	**-3.45**	**(-1.85, -1.62)**
***Tail Length**	**15**	**80.2 (5.36)**	**15**	**90.5 (5.36)**	**-1.21**	**(-1.45, -0.563)**
Throat Brightness	13	22.2 (5.81)	14	20.5 (5.81)	0.3	(-0.438, 1)
Throat Chroma	13	0.482 (0.0443)	14	0.486 (0.0443)	-0.0779	(-0.825, 0.644)
Throat Hue	13	641 (33.9)	14	654 (33.9)	-0.458	(-1.09, 0.327)
Vent Brightness	15	27.7 (7.2)	14	23.5 (7.2)	0.651	(-0.0368, 1.19)
***Vent Chroma**	**15**	**0.454 (0.0413)**	**14**	**0.487 (0.0413)**	**-0.725**	**(-1.21, -0.0361)**
Vent Hue	15	628 (23.4)	14	645 (23.4)	-0.642	(-1.28, 0.0747)

1161
1162