Structural analysis of SARS-CoV-2 and prediction of the human interactome

Andrea Vandelli[1,2], Michele Monti[1,3], Edoardo Milanetti[4,5], Riccardo Delli Ponti[6,*] and Gian Gaetano Tartaglia [1,3,5,7,*]

[1] Centre for Genomic Regulation (CRG), The Barcelona Institute for Science and Technology, Dr. Aiguader 88, 08003 Barcelona, Spain and Universitat Pompeu Fabra (UPF), 08003 Barcelona, Spain

[2] Systems Biology of Infection Lab, Department of Biochemistry and Molecular Biology, Biosciences Faculty, Universitat Autònoma de Barcelona, 08193 Cerdanyola del Vallès, Spain

[3] RNA System Biology Lab, department of Neuroscience and Brain Technologies, Istituto Italiano di Tecnologia, Via Morego 30, 16163, Genoa, Italy.

[4] Department of Physics, Sapienza University, Piazzale Aldo Moro 5, 00185, Rome, Italy

[5] Center for Life Nanoscience, Istituto Italiano di Tecnologia, Viale Regina Elena 291, 00161, Rome, Italy

[4] Department of Biology 'Charles Darwin', Sapienza University of Rome, P.le A. Moro 5, Rome 00185, Italy

[6] School of Biological Sciences, Nanyang Technological University, 60 Nanyang Drive, Singapore, 637551, Singapore

[7] Institucio Catalana de Recerca i Estudis Avançats (ICREA), 23 Passeig Lluis Companys, 08010 Barcelona, Spain

*to whom correspondence should be addressed to: riccardo.ponti@ntu.edu.sg (RDP) and giangaetano.tartaglia@uniroma1.it or gian.tartaglia@iit.it (GGT)

ABSTRACT

We calculated the structural properties of >2500 coronaviruses and computed >100000 human protein interactions with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). Using the *CROSS* method, we found that the SARS-CoV-2 region encompassing nucleotides 23000 and 24000 is highly conserved at the structural level, while the region 1000 nucleotides up-stream varies significantly. The two sequences code for a domain of the spike S protein that binds to the host receptor angiotensin-converting enzyme 2 (ACE2) that mediates human infection and in the homologue from Middle East respiratory syndrome coronavirus (MERS-CoV) interacts with sialic acids. Highly structured regions are predicted at the 5' and 3' where our calculations also indicate strong propensity to bind to human proteins. Using the *cat*RAPID method, we calculated 3500 interactions with the 5' and identified Cyclin T1 CCNT1, ATP-dependent RNA helicase DDX1,

Zinc Finger Protein ZNF175 and A-kinase anchor protein 8-like AKAP8L, among 20 high-

confidence candidate partners. We propose these proteins, also implicated in HIV replication, to be

further investigated for a better understanding of host-virus interaction mechanisms.

INTRODUCTION

A novel disease named Covid-19 by the World Health Organization and caused by the severe acute

respiratory syndrome coronavirus 2 (SARS-CoV-2) has been recognized as responsible for the

pneumonia outbreak that started in December, 2019 in Wuhan City, Hubei, China [1] and spread in

February to Milan, Lombardy, Italy [2] becoming pandemic. As of April 2020, the virus infected

>900'000 people in more than 200 countries.

SARS-CoV-2 shares similarities with other beta-coronavirus such as severe acute

respiratory syndrome coronavirus (SARS-CoV) and Middle East respiratory syndrome coronavirus

(MERS-CoV) [3]. Bats have been identified as the primary host for SARS-CoV and SARS-CoV-2 [4,5]

but the intermediate host linking SARS-CoV-2 to humans is still unknown, although a recent report

indicates that pangolins could be involved [6].

The coronaviruses use species-specific regions to mediate the entry in the host cell and SARS-CoV,

MERS-CoV and SARS-CoV-2, the spike S protein activates the infection in human respiratory

epithelial cells [7]. Spike S is assembled as a trimer and contains around 1,300 amino acids within

each unit [8]. In the S' region of the protein, the receptor binding domain (RBD), which contains

around 300 amino acids, mediates the binding with angiotensin-converting enzyme, (ACE2)

attacking respiratory cells. Another region upstream of the RBD, present in MERS-CoV but not in

SARS-CoV, is involved in the adhesion to sialic acid and could play a key role in regulating viral

infection [7,9].

At present, very few molecular details are available on SARS-CoV-2 and its interactions with

human host, which are mediated by specific RNA elements [10]. To study the RNA structural

content, we used *CROSS* [11] that was previously developed to investigate large transcripts such as the

human immunodeficiency virus HIV-1 [12]. *CROSS* predicts the structural profile (single- and double-

stranded state) at single-nucleotide resolution using sequence information only. We performed

sequence and structural alignments among 62 SARS-CoV-2 strains and identified the conservation

of specific elements in the spike S region, which provide clues on the evolution of domains

involved in the binding to ACE2 and sialic acid.

As highly structured regions of RNA molecules have strong propensity to form stable contacts with

proteins [13] and promote assembly of complexes [14,15], SARS-CoV-2 domains containing large

amount of double-stranded content are expected to establish specific interactions in host cells. To

investigate the interactions with human proteins, we employed *cat*RAPID [16,17]. *cat*RAPID [18]

estimates the binding potential or protein and RNA molecules through van der Waals, hydrogen

bonding and secondary structure propensities of allowing identification of interaction partners with

high confidence [19]. Our analysis revealed that the 5' of SARS-CoV-2 has strong propensity to

attract human proteins, especially those associated with viral infection, among which we found a

group linked to HIV infection. Intriguingly, a previous study reported similarities of viral proteins

in SARS-CoV and HIV-1 [20]. In HIV and SARS-CoV-2, but not   SARS-CoV nor MERS-CoV, a

furin-cleavage site occurs in the spike S protein [21]. This unique feature could explain the spread

velocity of SARS-CoV-2 compared to SARS-CoV and MERS-CoV.

We hope that our large-scale calculations of structural properties and binding partners of SARS-

CoV-2 can be useful to identify the mechanisms of virus interactions with the human host.

RESULTS

SARS-CoV-2 contains highly structured elements

Structural elements within RNA molecules attract proteins [13] and reveal regions important for

interactions with the host [22].

To analyze SARS-CoV-2 (reference Wuhan strain MN908947), we employed *CROSS* [11] that

predicts the double- and single-stranded content of large transcripts such as *Xist* and HIV-1 [12]. We

found the highest density of double-stranded regions in the 5' (nucleotides 1-253), membrane M

protein (nucleotides 26523-27191), spike S protein (nucleotides 23000-24000), and nucleocapsid N

protein (nucleotides 2874-29533; **Fig. 1**) [23]. The lowest density of double-stranded regions were

observed at nucleotides 6000-6250 and 20000-21500 and correspond to the regions between the

non-structural proteins nsp14 and nsp15 and the upstream region of the spike surface protein S (**Fig.**

1) [23]. In addition to the maximum corresponding to nucleotides 23000-24000, the structural content

of spike S protein shows minima at around nucleotides 20500 and 24500 (**Fig. 1**).

We used the *Vienna* method [24] to further investigate the RNA secondary structure of specific

regions identified with *CROSS* [12]. Employing a 100 nucleotide window centered around *CROSS*

maxima and minima, we found good match between *CROSS* scores and Vienna free energies (**Fig.**

1). Strong agreement is also observed between *CROSS* and *Vienna* positional entropy, indicating

that regions with the highest structural content have also the lowest structural diversity.

Our analysis suggests presence of structural elements in SARS-CoV-2 that have evolved to interact

with specific human proteins [10]. Our observation is based on the assumption that structured regions

have an intrinsic propensity to recruit proteins [13], which is supported by the fact that structured

transcripts act as scaffolds for protein assembly [14,15].

Structural comparisons reveal that the spike S region of SARS-CoV-2 is conserved among

coronaviruses

We employed *CROSS*align [12] to study the structural conservation of SARS-CoV-2 in different

strains.

In this analysis, we compared the Wuhan strain MN908947 with around 2800 other coronaviruses

(data from NCBI) having as host human (**Fig. 2**) or other species (**Supp. Fig. 1**). When comparing

SARS-CoV-2 with human coronaviruses (1387 strains, including SARS-CoV and MERS-CoV), we

found that the most conserved region falls inside the spike S genomic locus (**Fig. 2**). More

precisely, the conserved region is between nucleotides 23000 and 24000 and exhibits an intricate

and stable secondary structure (RNA*fold* minimum free energy= -269 kcal/mol)[24]. High

conservation of a structured regions suggests a functional activity that might be relevant for host

infection.

While the 3' and 5' of SARS-CoV-2 were shown to be relatively conserved in some beta-

coronavirus [10], they are highly variable in the entire set. However, the 3' and 5' are more structured

in SARS-CoV-2 than other coronaviruses (average structural content for SARS-CoV-2 = 0.56 in the

5' and 0.49 in the 3'; other coronaviruses 0.49 in the 5' and 0.42 in the 3').

Sequence and structural comparisons among SARS-CoV-2 strains indicate conservation of the ACE2 binding site and high variability in the region interacting with sialic acids.

To better investigate the sequence conservation of SARS-CoV-2, we compared 62 strains isolated

form different countries during the pandemic (including China, USA, Japan, Taiwan, India, Brazil,

Sweden, and Australia; data from NCBI and in VIPR www.viprbrc.org). Our analysis aims to

determine the relationship between structural content and sequence conservation.

Using *Clustal W* for multiple sequence alignments [25], we observed general conservation of the

coding regions with several *minima* in correspondence to areas between genes (**Fig. 3A**). One

highly conserved region is between nucleotides 23000 and 24000 in the spike S genomic locus,

while sequences up- and down-stream are variable (**Fig. 3A**). We then used CROSS*align* [12] to

compare the structural content. High variability of structure is observed for both the 5' and 3' and

for nucleotides between 21000 and 22000 as well as 24000 and 25000, associated with the S region

(see coloured bars in **Fig. 3A**). The rest of the regions are significantly conserved at a structural

level (p-value < 0.0001; Fisher's test).

We then compared protein sequences coded by the spike S genomic locus (NCBI reference

QHD43416) and found that both the sequence (**Fig. 3A**) and structure (**Fig. 2**) of nucleotides 23000

and 24000 are highly conserved. The region corresponds to amino acids 330-500 that contact the

host receptor angiotensin-converting enzyme 2 (ACE2) [26] provoking lung injury [27,28]. By contrast,

the region upstream of the binding site receptor ACE2 and located in correspondence to the

minimum of the structural profile at around nucleotides 22500-23000 (**Fig. 1**) is highly variable [52],

as calculated with *Tcoffee* multiple sequence alignments [52] (**Fig. 3A**). This part of the spike S

region corresponds to amino acids 243-302 that in MERS-CoV bind to sialic acids regulating

infection through cell-cell membrane fusion (**Fig. 3B;** see related manuscript by E. Milanetti *et al*.

"In-Silico evidence for two receptors based strategy of SARS-CoV-2") [9,29,30].

Our analysis suggests that the structural region between nucleotides 23000 and 24000 of Spike S

region is conserved among coronaviruses (**Fig. 2**) and the binding site for ACE2 has poor variation

in human SARS-CoV-2 strains (**Fig. 3B**). By contrast, the region upstream, potentially involved in

adhesion to sialic acids, has almost poor structural content and varies significantly in the human

population (**Fig. 3B**).

Analysis of human interactions with SARS-CoV-2 identifies proteins involved in viral

replication and HIV infection

In order to obtain insights on how the virus is replicated in human cells, we analysed protein-RNA

interactions with SARS-CoV-2 against the whole RNA-binding human proteome. Following a

protocol to study structural conservation in viruses [12], we divided the Wuhan sequence in 30

fragments of 1000 nucleotides moving from the 5' to 3' and calculated the protein-RNA

interactions of each fragment with the human proteome using *cat*RAPID *omics* (105000

interactions, consisting of around 3500 protein candidates for each of the 30 fragments. The list

includes canonical and non-canonical RNA-binding proteins, RBPs) [16]. For each fragment, we

identified the most significant interactions by filtering according to the Z interaction propensity. We

used three different thresholds in ascending order of stringency: Z greater or equal than 1.50, 1.75

and 2 respectively. Importantly, we removed from the list proteins that were predicted to interact

promiscuously with different fragments.

Fragment 1 corresponds to the 5' and is the most contacted by RBPs (around 120 with Z>2 high-

confidence interactions; **Fig. 4A**), which is in agreement with the observation that highly structured

regions attract a large number of proteins [13]. the 5' contains a leader sequence and the untranslated

region with multiple stem loop structures that control RNA replication and transcription [31,32].

The interactome of each fragment was then analysed using *clever*GO, a tool for GO enrichment

analysis [33]. Proteins interacting with fragments 1, 2 and 29 were associated with annotations

related to viral processes (**Fig. 4B; Supp. Table 1**). Considering the three thresholds applied

(**Materials and Methods**), we found 22 viral proteins for fragment 1, 2 proteins for fragment 2 and

11 proteins for fragment 29 (**Fig. 4C**).

Among the high-confidence interactors of fragment 1, we discovered RBPs involved in positive

regulation of viral processes and viral genome replication, such as Cyclin-T1 CCNT1 (Uniprot

code O60563 [34]), Double-stranded RNA-specific editase 1 ADARB1 (P78563) and 2-5A-

dependent ribonuclease RNASEL (Q05823). We also identified proteins related to the

establishment of integrated proviral latency, including X-ray repair cross-complementing protein 5

XRCC5 (P13010) and X-ray repair cross-complementing protein 6 XRCC6 (P12956; **Fig. 4D**).

Importantly, we found proteins related to defence response to viruses, such as ATP-dependent RNA

helicase DDX1 (Q92499) and Zinc finger protein 175 ZNF175 (Q9Y473), while Prospero

homeobox protein 1 PROX1 (Q92786) is involved in the negative regulation of viral genome

replication. Some of the remaining proteins are listed as DNA binding proteins and were included

because they could have potential RNA-binding ability (Fig. 4D)[35]. As for fragment 2, we found

two viral proteins: E3 ubiquitin-protein ligase TRIM32 (Q13049) and E3 ubiquitin-protein ligase

TRIM21 (P19474), which are listed as negative regulators of viral release from host cell, negative

regulators of viral transcription and positive regulators of viral entry into host cells. Finally, for

fragment 29, 10 of the 11 viral proteins found are members of the *endogenous retrovirus group K*

Gag polyprotein family, that perform different tasks during virus assembly, budding, maturation

(**Supp. Table 1**).

Analysis of functional annotations carried out with *GeneMania* [36] reveals that proteins interacting

with the 5' of SARS-CoV-2 RNA are associated with regulatory pathways involving NOTCH2,

MYC and MAX that have been previously connected to viral infection processes (**Fig. 4B**) [37,38].

Interestingly, some of the proteins, including CCNT1, DDX1, ZNF175 for fragment 1 and TRIM32

for fragment 2, are reported to be necessary for HIV functions and replications inside the cells.

More specifically, in the case of HIV infection, CCNT1 binds to the transactivation domain of the

viral nuclear transcriptional activator, Tat, increasing Tat's affinity for the transactivation response

RNA element; by doing so, it becomes an essential cofactor for Tat, promoting RNA Pol II

activation and allowing transcription of viral genes [39,40]. DDX1 is required for HIV-1 Rev function

as well as for HIV-1 and coronavirus IBV replication and it binds to the RRE sequence of HIV-1

RNAs [41,42]. ZNF175 is reported to interfere with HIV-1 replication by suppressing Tat-induced viral

LTR promoter activity [43]. Finally, TRIM32 is a well-defined Tat binding protein and, more

specifically, it binds to the activation domain of HIV-1 Tat and can also interact with the HIV-2 and

EIAV Tat proteins *in vivo* [44].

Recently, Gordon *et al*. reported a list of human proteins binding to Open Reading Frames (ORF)

translated from SARS-CoV-2 [45]. Identified through affinity purification followed by mass

spectrometry quantification, 332 proteins from HEK-293T cells interact with viral ORF peptides.

By selecting 266 proteins binding at the 5' with Z score >1.5 (**Supp. Table 1**), of which 140 are

exclusively interacting with fragment 1 (**Fig. 4B**), we found that 8 are also reported in the list by

Gordon *et al*. [45], which indicates significant enrichment (representation factor of 2.5; p-value of

0.02; hypergeometric test with the human proteome in background). Specifically, we note that out

of 332 interactions, 60 are RBPs (as reported in Uniprot [34]), which is a considerable fraction (20%),

considering that there are around 1500 RBPs in the human proteome (6%) [46]. Comparing the RBPs

present in Gordon *et al.* [45] and those present in our list (79 as reported in Uniprot), we found an

overlap of 6 proteins (representation factor = 26.5; p-value $< 10^{-8}$; hypergeometric test), including:

Janus kinase and microtubule-interacting protein 1 JAKMIP1 (Q96N16), A-kinase anchor protein 8

AKAP8 (O43823) and A-kinase anchor protein 8-like AKAP8L (Q9ULX6), which in case of HIV-

1 infection is involved in the DHX9-promoted annealing of human tRNA to viral genomic RNA[47],

Signal recognition particle subunit SRP72 (O76094), binding to the 7S RNA in presence of SRP68,

La-related protein 7, LARP7 (Q4G0J3) and La-related protein 4B LARP4B (Q92615), which are

part of a system for transcriptional regulation of polymerase II genes acting by means of the 7SK

RNP system [48] (**Fig. 4E; Supp. Table 2**).

Moreover, by analysing the RNA interaction potential of all the 332 proteins by Gordon *et al.* [45],

*cat*RAPID identified 38 putative binders at the 5' (Z score > 1.5; 27 occurring in the 5' and not in

other regions of the viral RNA) [16], including Serine/threonine-protein kinase TBK1 (Q9UHD2),

among which 10 RBPs (as reported in Uniprot) such as: Splicing elements U3 small nucleolar

ribonucleoprotein protein MPP10 (O00566) and Pre-mRNA-splicing factor SLU7 (O95391),

snRNA methylphosphate capping enzyme MEPCE involved in negative regulation of transcription

by RNA polymerase II 7SK (Q7L2J0) [49], Nucleolar protein 10 NOL10 (Q9BSC4) and protein

kinase A Radixin RDX (P35241; in addition to those mentioned above; **Supp. Table 2**).

CONCLUSIONS

Our study is motivated by the need to identify interactions involved in Covid-19 spreading. Using

advanced computational approaches, we investigated the structural content of the virus and

predicted its binding to human proteins.

We employed *CROSS* [12,50] to compare the structural properties of 2800 coronaviruses and identified

elements conserved in SARS-CoV-2 strains. The regions containing the highest amount of structure

are the 5' as well as glycoproteins spike S and membrane M.

We found that the spike S protein domain encompassing amino acids 330-500 is highly conserved

across SARS-CoV-2 strains. This result suggests that spike S has evolved to specifically interact

with its host partner ACE2 [26] and mutations increasing the binding affinity are infrequent. As the

nucleic acids encoding for this region are enriched in double-stranded content, we speculate that the

structure might attract host regulatory proteins, which further constrains its variability. The fact that

the ACE2 receptor binding site is conserved among the SARS-CoV-2 strains suggests that a

specific drug can be designed to prevent host interaction and thus infection, which could work for a

large number of coronaviruses.

By contrast, the highly variable region at amino acids 243-302 in spike S protein corresponds to the

binding site of sialic acid in MERS-CoV (see related manuscript by E. Milanetti *et al*. "In-Silico

evidence for two receptors based strategy of SARS-CoV-2") [7,9,30] and regulates host cell infection

[29]. The fact that the binding region change in the different strains might indicate different binding

affinities, which could provide clues on the different levels of contagion in the human population.

Interestingly, the sialic acid binding is absent in SARS-CoV but present in MERS-CoV, which

indicates that it must have evolved recently.

Both our sequence and structural analyses of spike S protein indicate that human engineering of

SARS-CoV-2 is highly unlikely.

Using *cat*RAPID [16,17] we predicted that the highly structured region at the 5' is the region with

largest number or protein partners, including the helicase DDX1, which has been previously

reported to be essential for HIV-1 and coronavirus IBV [41,42], the non-canonical RNA-binding

proteins CCNT1 [39,40] and ZNF175 [43] involved in polymerase II recruitment (among others).

Connections to regulatory pathways involving NOTCH2, MYC and MAX have also been

identified. A significant overlap exists with the list of protein interactions reported by Gordon *et al*.

[45], and among the candidate binding partners we found AKAP8L, involved in the DHX9 helicase-

promoted annealing of host tRNA to HIV genomic RNA [47]. The link between HIV and these

proteins could motivate analysing HIV drugs for treatment of SARS-CoV infection [51].

We hope that our analysis would be useful to the scientific community to identify virus-host

interactions and block SARS-CoV-2 spreading.

Acknowledgements

The authors would like to thank Jakob Rupert, Dr. Mattia Miotto, Dr Lorenzo Di Rienzo, Dr. Alexandros Armaos, Dr. Alessandro Dasti, Dr. Elias Bechara, Dr. Claudia Giambartolomei and Dr. Elsa Zacco for discussions.

The research leading to these results has been supported by European Research Council (RIBOMYLOME_309545 to GGT, ASTRA_855923 to GGT), Spanish Ministry of Economy and Competitiveness BFU2017-86970-P, the H2020 projects INFORE_825080 and IASIS_727658, as well as the collaboration with Peter St. George-Hyslop financed by the Wellcome Trust.

Contributions. AV carried out the *cat*RAPID analysis of protein interactions, RDP calculated *CROSS* structures of coronaviruses, GGT, MM and EM performed and analysed sequence alignments. GGT and RDP conceived the study. AV, RDP and GGT wrote the paper.

MATERIALS AND METHODS

Structure prediction

We predicted the secondary structure of transcripts using *CROSS* (Computational Recognition of Secondary Structure [12,50]. *CROSS* was developed to perform high-throughput RNA profiling. The algorithm predicts the structural profile (single- and double-stranded state) at single-nucleotide resolution using sequence information only and without sequence length restrictions (scores > 0 indicate double stranded regions). We used the *Vienna* method [24] to further investigate the RNA secondary structure of minima and maxima identified with *CROSS* [12].

Structural conservation

We used *CROSS*align [12,50] an algorithm based on Dynamic Time Warping (DTW), to check and evaluate the structural conservation between different viral genomes [12]. CROSS*align* was previously employed to study the structural conservation of ~5000 HIV genomes. SARS-CoV-2 fragments (1000 nt, not overlapping) were searched inside other complete genomes using the OBE (open begin and end) module, in order to search a small profile inside a larger one. The lower the structural distance, the higher the structural similarities (with a minimum of 0 for almost identical secondary structure profiles). The significance is assessed as in the original publication [12].

Sequence collection

The fasta sequences of the complete genomes of SARS-CoV-2 were downloaded from Virus Pathogen Resource (VIPR; www.viprbrc.org), for a total of 62 strains. Regarding the overall coronaviruses, the sequences were downloaded from NCBI selecting only complete genomes, for a total of 2862 genomes. The reference Wuhan sequence with available annotation (EPI_ISL_402119) was downloaded from Global Initiative on Sharing All Influenza Data. (GISAID https://www.gisaid.org/).

Protein-RNA interaction prediction

Interactions between each fragment of target sequence and the human proteome were predicted

using *cat*RAPID *omics* [16,17] , an algorithm that estimates the binding propensity of protein-RNA

pairs by combining secondary structure, hydrogen bonding and van der Waals contributions. The

complete list of interactions between the 30 fragments and the human proteome is available at

http://crg-webservice.s3.amazonaws.com/submissions/2020-

03/252523/output/index.html?unlock=f6ca306af0. The output then is filtered according to the Z-

score column. We tried three different thresholds in ascending order of stringency: Z greater or

equal than 1.50, 1.75 and 2 respectively and for each threshold we then selected the proteins that

were unique for each fragment for each threshold.

GO terms analysis

*clever*GO [33], an algorithm for the analysis of Gene Ontology annotations, was used to determine

which fragments present enrichment in GO terms related to viral processes. Analysis of functional

annotations was performed in parallel with *GeneMania* [36].

RNA and protein alignments

We sued *Clustal W* [25] for 62 SARS-CoV-2 strains alignments and *Tcoffee* [52] for spike S proteins

alignments. The variability in the spike S region was measured by computing Shannon entropy on

translated RNA sequences. The Shannon entropy is computed as follows:

$S(a) = - \text{Sum\_i } P(a,i) \log P(a,i)$

Where *a* correspond to the amino acid at the position *i* and $P(a,i)$ is the frequency of a certain

amino-acid *a* at position *i* of the sequence. Low entropy indicates poorly variability: if $P(a,x) = 1$ for

one *a* and 0 for the rest, then $S(x) = 0$. By contrast, if the frequencies of all amino acids are equally

distributed, the entropy reaches its maximum possible value.

1. Zhu, N. *et al.* A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N. Engl. J.*

Med. **382**, 727–733 (2020).

2. D'Antiga, L. Coronaviruses and immunosuppressed patients. The facts during the third

epidemic. *Liver Transplant. Off. Publ. Am. Assoc. Study Liver Dis. Int. Liver Transplant. Soc.*

(2020) doi:10.1002/lt.25756.

3. Cascella, M., Rajnik, M., Cuomo, A., Dulebohn, S. C. & Di Napoli, R. Features, Evaluation and

Treatment Coronavirus (COVID-19). in *StatPearls* (StatPearls Publishing, 2020).

4. Ge, X.-Y. *et al.* Isolation and characterization of a bat SARS-like coronavirus that uses the

ACE2 receptor. *Nature* **503**, 535–538 (2013).

5. Follis, K. E., York, J. & Nunberg, J. H. Furin cleavage of the SARS coronavirus spike

glycoprotein enhances cell-cell fusion but does not affect virion entry. *Virology* **350**, 358–369

(2006).

6. Xiao, K. *et al.* Isolation and Characterization of 2019-nCoV-like Coronavirus from Malayan

Pangolins. *bioRxiv* 2020.02.17.951335 (2020) doi:10.1101/2020.02.17.951335.

7. Park, Y.-J. *et al.* Structures of MERS-CoV spike glycoprotein in complex with sialoside

attachment receptors. *Nat. Struct. Mol. Biol.* **26**, 1151–1157 (2019).

8. Walls, A. C. *et al.* Cryo-electron microscopy structure of a coronavirus spike glycoprotein

trimer. *Nature* **531**, 114–117 (2016).

9. Li, W. *et al.* Identification of sialic acid-binding function for the Middle East respiratory

syndrome coronavirus spike glycoprotein. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E8508–E8517

(2017).

10. Yang, D. & Leibowitz, J. L. The Structure and Functions of Coronavirus Genomic 3' and 5'

Ends. *Virus Res.* **206**, 120–133 (2015).

11. Delli Ponti, R., Marti, S., Armaos, A. & Tartaglia, G. G. A high-throughput approach to profile

RNA structure. *Nucleic Acids Res.* **45**, e35–e35 (2017).

12. Delli Ponti, R., Armaos, A., Marti, S. & Gian Gaetano Tartaglia. A Method for RNA Structure

Prediction Shows Evidence for Structure in lncRNAs. *Front. Mol. Biosci.* **5**, 111 (2018).

13. Sanchez de Groot, N. *et al.* RNA structure drives interaction with proteins. *Nat. Commun.* **10**,

3246 (2019).

14. Cid-Samper, F. *et al.* An Integrative Study of Protein-RNA Condensates Identifies Scaffolding

RNAs and Reveals Players in Fragile X-Associated Tremor/Ataxia Syndrome. *Cell Rep.* **25**,

3422-3434.e7 (2018).

15. Cerase, A. *et al.* Phase separation drives X-chromosome inactivation: a hypothesis. *Nat. Struct.*

Mol. Biol. **26**, 331 (2019).

16. Agostini, F. *et al.* catRAPID omics: a web server for large-scale prediction of protein-RNA

interactions. *Bioinforma. Oxf. Engl.* **29**, 2928–2930 (2013).

17. Cirillo, D. *et al.* Quantitative predictions of protein interactions with long noncoding RNAs.

Nat. Methods **14**, 5–6 (2017).

18. Bellucci, M., Agostini, F., Masin, M. & Tartaglia, G. G. Predicting protein associations with

long noncoding RNAs. *Nat. Methods* **8**, 444–445 (2011).

19. Lang, B., Armaos, A. & Tartaglia, G. G. RNAct: Protein–RNA interaction predictions for

model organisms with supporting experimental data. *Nucleic Acids Res.*

doi:10.1093/nar/gky967.

20. Kliger, Y. & Levanon, E. Y. Cloaked similarity between HIV-1 and SARS-CoV suggests an

anti-SARS strategy. *BMC Microbiol.* **3**, 20 (2003).

21. Hallenberger, S. *et al.* Inhibition of furin-mediated cleavage activation of HIV-1 glycoprotein

gp160. *Nature* **360**, 358–361 (1992).

22. Gultyaev, A. P., Richard, M., Spronken, M. I., Olsthoorn, R. C. L. & Fouchier, R. A. M.

Conserved structural RNA domains in regions coding for cleavage site motifs in hemagglutinin

genes of influenza viruses. *Virus Evol.* **5**, (2019).

23. Wu, A. *et al.* Genome Composition and Divergence of the Novel Coronavirus (2019-nCoV)

Originating in China. *Cell Host Microbe* **27**, 325–328 (2020).

24. Lorenz, R. *et al.* ViennaRNA Package 2.0. *Algorithms Mol. Biol.* **6**, 26 (2011).

25. Madeira, F. *et al.* The EMBL-EBI search and sequence analysis tools APIs in 2019. *Nucleic*

Acids Res. **47**, W636–W641 (2019).

26. Andersen, K. G., Rambaut, A., Lipkin, W. I., Holmes, E. C. & Garry, R. F. The proximal origin

of SARS-CoV-2. *Nat. Med.* 1–3 (2020) doi:10.1038/s41591-020-0820-9.

27. Glowacka, I. *et al.* Differential downregulation of ACE2 by the spike proteins of severe acute

respiratory syndrome coronavirus and human coronavirus NL63. *J. Virol.* **84**, 1198–1205

(2010).

28. Zhou, P. *et al.* A pneumonia outbreak associated with a new coronavirus of probable bat origin.

Nature **579**, 270–273 (2020).

29. Qing, E., Hantak, M., Perlman, S. & Gallagher, T. Distinct Roles for Sialoside and Protein

Receptors in Coronavirus Infection. *mBio* **11**, (2020).

30. Milanetti, E. *et al.* In-Silico evidence for two receptors based strategy of SARS-CoV-2.

ArXiv200311107 Phys. Q-Bio (2020).

31. Lu, K., Heng, X. & Summers, M. F. Structural determinants and mechanism of HIV-1 genome

packaging. *J. Mol. Biol.* **410**, 609–633 (2011).

32. Fehr, A. R. & Perlman, S. Coronaviruses: An Overview of Their Replication and Pathogenesis.

Methods Mol. Biol. Clifton NJ **1282**, 1–23 (2015).

33. Klus, P., Ponti, R. D., Livi, C. M. & Tartaglia, G. G. Protein aggregation, structural disorder

and RNA-binding ability: a new approach for physico-chemical and gene ontology

classification of multiple datasets. *BMC Genomics* **16**, 1071 (2015).

34. UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Res.* **47**, D506–D515 (2019).

35. Castello, A. *et al.* Insights into RNA biology from an atlas of mammalian mRNA-binding

proteins. *Cell* **149**, 1393–1406 (2012).

36. Warde-Farley, D. *et al.* The GeneMANIA prediction server: biological network integration for

gene prioritization and predicting gene function. *Nucleic Acids Res.* **38**, W214–W220 (2010).

37. Hayward, S. D. Viral interactions with the Notch pathway. *Semin. Cancer Biol.* **14**, 387–396

(2004).

38. Dudley, J. P., Mertz, J. A., Rajan, L., Lozano, M. & Broussard, D. R. What retroviruses teach

us about the involvement of c- Myc in leukemias and lymphomas. *Leukemia* **16**, 1086–1098

(2002).

39. Ivanov, D. *et al.* Cyclin T1 domains involved in complex formation with Tat and TAR RNA are

critical for tat-activation. *J. Mol. Biol.* **288**, 41–56 (1999).

40. Kwak, Y. T., Ivanov, D., Guo, J., Nee, E. & Gaynor, R. B. Role of the human and murine cyclin

T proteins in regulating HIV-1 tat-activation. *J. Mol. Biol.* **288**, 57–69 (1999).

41. Fang, J. *et al.* A DEAD box protein facilitates HIV-1 replication as a cellular co-factor of Rev.

Virology **330**, 471–480 (2004).

42. Xu, L. *et al.* The cellular RNA helicase DDX1 interacts with coronavirus nonstructural protein

14 and enhances viral replication. *J. Virol.* **84**, 8571–8583 (2010).

43. Carlson, K. A. *et al.* Molecular characterization of a putative antiretroviral transcriptional

factor, OTK18. *J. Immunol. Baltim. Md 1950* **172**, 381–391 (2004).

44. Locke, M., Tinsley, C. L., Benson, M. A. & Blake, D. J. TRIM32 is an E3 ubiquitin ligase for

dysbindin. *Hum. Mol. Genet.* **18**, 2344–2358 (2009).

45. Gordon, D. E. *et al.* A SARS-CoV-2-Human Protein-Protein Interaction Map Reveals Drug

Targets and Potential Drug-Repurposing. *bioRxiv* 2020.03.22.002386 (2020)

doi:10.1101/2020.03.22.002386.

46. Castello, A. *et al.* Comprehensive Identification of RNA-Binding Domains in Human Cells.

Mol. Cell **63**, 696–710 (2016).

47. Xing, L., Zhao, X., Guo, F. & Kleiman, L. The role of A-kinase anchoring protein 95-like

protein in annealing of tRNALys3 to HIV-1 RNA. *Retrovirology* **11**, 58 (2014).

48. Markert, A. *et al.* The La-related protein LARP7 is a component of the 7SK ribonucleoprotein

and affects transcription of cellular and viral polymerase II genes. *EMBO Rep.* **9**, 569–575

(2008).

49. C, J. *et al.* Systematic Analysis of the Protein Interaction Network for the Human Transcription

Machinery Reveals the Identity of the 7SK Capping Enzyme. *Molecular cell* vol. 27

https://pubmed.ncbi.nlm.nih.gov/17643375/ (2007).

50. Delli Ponti, R., Marti, S., Armaos, A. & Tartaglia, G. G. A high-throughput approach to profile

RNA structure. *Nucleic Acids Res.* **45**, e35–e35 (2017).

51. Arabi, Y. M. *et al.* Treatment of Middle East respiratory syndrome with a combination of

lopinavir/ritonavir and interferon-β1b (MIRACLE trial): statistical analysis plan for a recursive

two-stage group sequential randomized controlled trial. *Trials* **21**, 8 (2020).

52. Di Tommaso, P. *et al.* T-Coffee: a web server for the multiple sequence alignment of protein

and RNA sequences using structural information and homology extension. *Nucleic Acids Res.*

39, W13–W17 (2011).

FIGURES LEGENDS

Fig. 1 *Using the CROSS approach [12,50], we studied the structural content of SARS-CoV-2. We*

found the highest density of double-stranded regions in the 5' (nucleotides 1-253), membrane M

protein (nucleotides 26523-27191), and the spike S protein (nucleotides 23000-24000). Strong

match is observed between CROSS and Vienna analyses (centroid structures shown, indicating that

regions with the highest structural content have the lowest free energies.

Fig. 2 *We employed the CROSSalign approach [12,50] to compare the Wuhan strain MN908947 with*

other coronaviruses (1387 strains, including SARS-CoV and MERS-CoV) indicates that the most

conserved region falls inside the spike S genomic locus. The inset shows thermodynamic structural

variability (positional entropy) within regions encompassing nucleotides 23000-24000 along with

the centroid structure and free energy.

Fig. 3 *Sequence and structural comparison of human SARS-CoV-2 strains. (**A**) Strong sequence*

conservation (Clustal W multiple sequence alignments [33]) is observed in coding regions, including

the region between nucleotides 23000 and 24000 of spike S protein. High structural variability

(colour bars on top) is observed for both the UTRs and for nucleotides between 21000 and 22000

as well as 24000 and 25000, associated with the S region. The rest of the regions are significantly

*conserved at a structural level. (**B**) The sequence variability (Shannon entropy computed on Tcoffee*

multiple sequence alignments [52]) in the spike S protein indicate conservation between amino-acids

460 and 520 (blue box) binding to the host receptor angiotensin-converting enzyme 2 ACE2. The

region encompassing amino-acids 243 and 302 is highly variable and is implicated in sialic acids

in MERS-CoV (red box). The S1 and S2 domains of Spike S protein are displayed.

Fig. 4. *Characterization of protein interactions with SARS-CoV-2 RNA, (**A**) Number of RBP*

interactions for different SARS-CoV-2 regions (colours indicate different catRAPID [16,17] confidence

levels: Z=1.5 or low Z=1.75 or medium and Z=2.0 or high; regions with scores lower than Z=1.5

*are omitted); (**B**) Enrichment of viral processes in the 5' of SARS-CoV-2 (precision = term*

precision calculated from the GO graph structure lvl = depth of the term; go_term = GO term

identifier, with link to term description at AmiGO website ; description = Textual label for the term;

e/d = e signifies enrichment of the term, d signifies depletion compared to the population; %_set =

coverage on the provided set - how much of the set is annotated with the GO?; %_pop = coverage

of the same term on the population; p_bonf = p-value of the enrichment. To correct for multiple

testing bias, we are applying Bonferroni correction) [33]*; (**C**) Viral processes are the third largest*

*cluster identified in our analysis; (**D**) Protein interactions with the 5' of SARS-CoV-2 RNA (inner*

circle) and associations with other human genes retrieved from literature (green: genetic

*associations; pink: physical associations); (**E**) Number of RBP interactions identified by Gordon et*

al. [45] *for different SARS-CoV-2 regions (see panel A for reference).*

SUPPLEMENTARY MATERIAL

Supp. Table 1. *1) catRAPID* [16,17] *score for interactions with fragment 1; 2) GO* [33] *and Uniprot*

annotations of viral proteins interacting with fragment 1 and ; 3) catRAPID score for interactions

with fragment 2; 4) GO annotations of viral proteins interacting with fragment 2; 5) catRAPID

score for interactions with fragment 29; 6) GO annotations of viral proteins interacting with

fragment 29;

Supp. Table 2. *RBP interactions from Gordon et al.* [45] *classified according to catRAPID scores.*

GO [33] *and Uniprot* [34] *annotations are reported.*

Supp. Figure 1. *We employed CROSSalign* [12,50] *was to compare the Wuhan strain MN908947*

with other coronaviruses (2800 strains, including SARS-CoV, MERS-CoV and coronaviruses

having as host other species, such as bats). The result highlights that the most conserved region

falls inside the spike S genomic locus.

A

Number of protein interactions (y-axis), Genomic Region (KB) (x-axis)

Legend:
CCNT1
DDX1
ZNF175
TRIM32

Z>1.5
Z>1.75
Z>2

B

precision	lvl	go_term	description	e/d	%_set	%_pop	p_bonf
0.657	4	GO:0050792	regulation of viral process	e	4.918	0.205	4.59e-4
0.775	5	GO:0045069	regulation of viral genome replication	e	3.279	0.083	7.33e-3
0.794	5	GO:0048524	positive regulation of viral process	e	3.279	0.091	1.05e-2
0.748	3	GO:0009615	response to virus	e	4.918	0.362	1.23e-2
0.510	3	GO:0016032	viral process	e	6.557	0.791	1.3e-2
0.903	3	GO:0051607	defense response to virus	e	4.098	0.216	1.53e-2
0.927	6	GO:0045071	negative regulation of viral genome replication	e	2.459	0.053	7.93e-2
1.000	5	GO:0075713	establishment of integrated proviral latency	e	1.639	0.009	1.05e-1
0.835	4	GO:0019043	establishment of viral latency	e	1.639	0.011	1.69e-1
0.794	5	GO:0048525	negative regulation of viral process	e	2.459	0.098	4.98e-1

C

mutualism, multiorganism, replication, viral, positive, regulation, genome, parasitism, negative, encompassing, symbiosis, process

D

E

Number of protein interactions (y-axis), Genomic Region (KB) (x-axis)

Legend:
AKAP8
AKAP8L
LARP7
LARP4B

Z>1.5
Z>1.75
Z>2