NAD$^+$ prevents septic shock-induced death by non-canonical inflammasome

blockade and IL-10 cytokine production in macrophages

Jasper Iske[1,2†], Rachid El Fatimy[3,4†], Yeqi Nian[1,5†], Amina Ghouzlani[6], Siawosh K. Eskandari[7], Hector Rodriguez Cetina Biefer[1,8], Anju Vasudevan[9] and Abdallah Elkhal[1,6]∗

[1]Division of Transplant Surgery, Department of Surgery, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA

[2]Department of Cardiothoracic and Vascular Surgery, Germany Heart Center Berlin, Berlin, Germany

[3]Department of Neurology, Ann Romney Center for Neurologic Diseases, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA

[4]Institute of Biological Sciences (ISSB-P), Mohammed VI Polytechnic University (UM6P), 43150, Benguerir, Morocco

[5]Institute of Transplant Medicine, Tianjin First Central Hospital, Nankai University, Tianjin 300192, China

[6]NAD$^+$ Immunology Laboratory, Huntington Medical Research Institutes, Pasadena, CA, USA

[7]Department of Internal Medicine, University of Groningen, Groningen, Netherlands

[8]Department of Cardiac Surgery, Stadtspital Zurich Triemli, Zurich, Switzerland

[9]Angiogenesis and Brain Development Laboratory, Huntington Medical Research Institutes, Pasadena, CA, USA

[†]These authors contributed equally to this work.

Corresponding author:

Abdallah Elkhal, Ph.D.,

Cardiovascular Department

Huntington Medical Research Institutes

Pasadena, CA

E-mail: Abdala.elkhal@hmri.org

Abstract

Septic shock is characterized by an excessive inflammatory response depicted in a cytokine

storm that results from invasive bacterial, fungi, protozoa, and viral infections. Non-canonical

inflammasome activation is crucial in the development of septic shock promoting pyroptosis and pro-

inflammatory cytokine production via caspase-11 and Gasdermin-D (GSDMD). Here, we show

that NAD^+ treatment protected mice towards bacterial and LPS induced endotoxic shock by

blocking the non-canonical inflammasome specifically. NAD^+ administration impeded systemic

IL-1β and IL-18 production and GSDMD-mediated pyroptosis of macrophages via the IFN-

β/STAT-1 signaling machinery. More importantly, NAD^+ administration not only improved

casp-11 KO (knockout) survival but rendered WT mice completely resistant to septic shock

via the IL-10 signaling pathway that was independent from the non-canonical inflammasome.

Here, we delineated a two-sided effect of NAD^+ blocking septic shock through a specific inhibition of

the non-canonical inflammasome and promoting immune homeostasis via IL-10, underscoring its

unique therapeutic potential.

Introduction

Sepsis is characterized by a systemic inflammatory response syndrome (SIRS)[1] driven by host cells following systemic bacterial[2] and viral infections. The excessive inflammatory response can derail into septic shock resulting in multiple organ failure (MOF), the leading cause of death in intensive care units. Inflammasome activation, which downstream pathways cause the release of proinflammatory cytokines and the induction of an inflammatory cell death termed pyroptosis[3], has been pointed out as the major driver of septic shock. Hereby, a two-armed LPS derived induction of the NLRP3-canonical inflammasome, the major source of IL-1β and IL-18 cytokine production[4] and the caspase-11 mediated non-canonical inflammasome leading to pyroptosis in monocytes[5] was determined as the underlying mechanism. Mechanistically, Caspase-11 acts as a pattern recognition receptor for intracellular bacteria[6] that cleaves gasdermin-D, a membrane-pore forming protein subsequently inducing pyroptotic cell death[7]. The NLRP3-canonical inflammasome in turn, was found to be indispensable[8] for septic shock induced death. However cross-activation through Caspase-11 promoting cytokine release has been described[9, 10, 11], assigning the non-canonical inflammasome a cardinal role[12].

Recent approaches such as anti-proinflammatory cytokine strategies, blocking downstream targets of inflammasomes have been ineffective[13] while inhibiting inflammatory key regulators such as NF-κB may promote adverse side-effects[14]. Hence, contemporary clinical therapy of septic shock is based on symptomatic treatment rather than curative approaches that clear the cause of the disease itself.

In our previous studies, we have underscored the immunosuppressive properties of NAD^+ in autoimmune diseases and allo-immunity via the regulation of $CD4^+$ T cell fate[15, 16]. More recently, we have shown that NAD^+ administration protected mice from lethal doses of *Listeria*

monocytogenes (*L. m.*) via mast cells (MCs) exclusively and independently of major antigen

presenting cells (APCs)[17]. However, the underlying mechanism that allows NAD^+, to

concomitantly protect against autoimmune diseases, via its immunosuppressive properties[15, 16],

and against lethal bacterial infection remains unclear.

Therefore, in the current study we investigated whether NAD^+ protects against bacterial infection

by dampening the systemic inflammatory response associated with sepsis or through enhanced

bacterial clearance. Although, wild type (WT) mice subjected to NAD^+ or PBS and lethal doses

of pathogenic *Escherichia coli* (*E. coli*) exhibited similar bacterial load in various tissues, mice

treated with NAD^+ displayed a robust survival. Moreover, NAD^+ protected against LPS-induced

death that was associated with a dramatic decrease of systemic IL-1β and IL-18 levels, two major

cytokines involved in the inflammasome signaling machinery. More importantly, we show that

NAD^+ protected from LPS-induced death by targeting specifically the non-canonical

inflammasome via a blockade of the STAT-1/IFN-β signaling pathway. Moreover, NAD^+

treatment rendered not only Caspase-11 KO but WT mice fully resistant to poly(I:C) + LPS

induced septic shock, via an inflammasome independent pathway mediated by a systemic IL-10

cytokine production.

Results

NAD[+] protects mice against septic shock not via bacterial clearance but via inflammasome blockade

Our previous studies have underscored the role of NAD[+] in regulating CD4[+] T cell fate and its immunosuppressive properties via IL-10 cytokine production[15, 16, 17]. More recently, we have shown that NAD[+] protected mice against lethal doses of *L. m.* independently of major APCs[15]. However, it remained unclear whether NAD[+] protected mice against lethal doses of *L. m.*, a gram-positive bacterium, via a clearance mechanism or by dampening the inflammatory response. Since *L. m.* is known to be an intracellular pathogen, we tested if NAD[+] protects as well against *E. coli*, a gram-negative bacterium that is well known to induce septic shock[18]. Wild type mice were treated with NAD[+] or PBS for 2 consecutive days followed by a lethal dose (1×10^9) of *E. coli*. or PBS. Notably, mice treated with PBS died within 5 hours after infection, while mice treated with NAD[+] exhibited an impressive survival (**Fig. 1A**). Moreover, when assessing the bacterial load in liver and kidney (**Fig. 1B**), organs exposed to the infection, by counting CFU in both, NAD[+] and PBS groups, revealed no significant difference, suggesting that NAD[+] does not promote bacterial clearance**.** More importantly, these data suggest that NAD[+] may reduce the inflammatory response towards bacterial infection. It is well established that the bacterial lipopolysaccharide (LPS) abundant on the outer membrane exhibits a key role in the pathology of *E. coli* derived septic shock[13]. Thus, we further characterize the impact of NAD[+] on septic shock by subjecting mice to a lethal dose (54mg/kg) of two different LPS serotypes (O111:B4 and O55:B5) described to vary in the antigen lipid A content and to promote distinct hypothermia kinetics[19]. Following LPS (O111:B4 and O55:B5) administration, PBS treated control mice displayed severe symptoms of endotoxic shock with a dramatical body temperature

117 decrease (<23˚C) within 15 hours. In contrast, mice subjected to NAD[+] exhibited highly distinct

118 kinetics with a recovery of body temperatures after 15 hours (**Fig. 1C**). When monitoring

119 survival, 100% of PBS treated mice succumbed to LPS after 24 hours while NAD[+] treated

120 animals exhibited an overall survival >85% (**Fig. 1D**), which was consistent with our bacterial

121 infection model. Mice infected and treated with NAD[+] survived for several months and recovered fully

after 10 days. Of note, mice survived for over a year following infection and died of aging.

LPS-induced death derives from multi-organ failure[20]. Therefore, lung, kidney, liver, ileum and

spleen were harvested 15 hours after LPS administration and tissue damage was assessed by

Hematoxylin and Eosin (H&E) staining. Tissue evaluation indicated severe pulmonary

hemorrhage, excessive tubular fibrin deposition, hepatocyte cell swelling, disseminated

intravascular coagulation (DIC) and ileal villi destruction consistent with a multi-organ-

dysfunction syndrome (MODS)[21] in mice treated with PBS. In contrast, NAD[+] administration

dramatically attenuated signs of organ failure with significantly less pulmonary hemorrhage and

DIC, intact liver and kidney tissue architecture and preserved ileal villi (**Fig. 1E, Fig. 1 Supplement 1**

and Fig. 1 Supplement 2).

To elucidate the protective effects of NAD[+] systemic levels of IL-1β and IL-18, two major cytokines

implicated in inflammasome activation, were measured 10 and 15 hours after intraperitoneal

injection of LPS (**Fig. 1F**). Of, note IL-6 and TNFα systemic levels were measured as well (**Fig. 1F**).

Our findings indicated that LPS injection resulted in a robust

systemic increase of IL-1β, IL-6, TNFα and IL-18 in the PBS group, which was almost abolished in

NAD[+] treated mice. Taken together, our results suggest that NAD[+] protects mice against septic shock

not via bacterial clearance but rather via inflammasome blockade.

NAD$^+$ specifically inhibits the non-canonical inflammasome

Our data suggest that NAD$^+$ protects against septic shock via inflammasome blockade.

Monocytes, especially macrophages, have been described as major drivers of inflammasome

derived cytokine secretion in the context of septic shock[22]. Thus, to test the effect of NAD$^+$ on

inflammasome function, bone marrow derived macrophages (BMDMs) were obtained and both,

canonical and non-canonical inflammasomes were stimulated in in the presence or absence of

100 µmol/ml NAD$^+$. Activation of the canonical pathway was achieved through LPS priming

(1µg/ml) followed by ATP stimulation (5 mmol/l). Notably, BMDMs subjected to NAD$^+$ or PBS

treatment followed by canonical inflammasome activation did not exhibit any significant

difference in IL-1β secretion or pyroptosis that was assessed by LDH release measurement, a

marker for cell death[23] (**Fig. 2A**).

To trigger the non-canonical inflammasome pathway, BMDMs were primed with Pam3CSK4, a

TLR1/2 agonist, followed by cholera toxin B (CTB) and LPS (2µg/ml) administration. The data

showed that NAD$^+$ treatment resulted in a robust reduction of IL-1β release and cell death when

compared to the PBS control group (**Fig. 2A**). Furthermore, Western blotting revealed that

BMDMs cultured in presence of NAD$^+$ exhibited a dramatic decrease of casp-11 expression and

its downstream targets including casp-1, IL-1β and cleaved gasdermin-D (GSDMD) (**Fig 2B**).

Moreover, we observed a prominent decrease in casp-1 expression under NAD$^+$ treatment that

was constant over the time course of 16 hours. In contrast, BMDMs treated with PBS exhibited

excessive casp-1 expression at 4 hours that was found to be absent after 16 hours (**Fig. 2C**),

which is consistent with the strong cytotoxicity leading to membrane permeabilization and

release of Casp-1 into the supernatant. Noteworthy, Pam3CK4 derived BMDM priming was not

affected by NAD$^+$ since NF-κB as well as pro-caspase-1 levels had not been altered (**Fig. 2A**

and Fig. 2 Supplement 1) underlining the specific inhibition of casp-11. To visualize NAD$^+$

mediated blockade of pyroptotic macrophage death, BMDMs were treated with PBS or NAD$^+$,

primed with Pam3CSK4, then stimulated with LPS and CTB and cell viability and apoptosis were

monitored using the IncuCyte$^{®}$ live microscopy system. Hereby, we observed distinct

longitudinal kinetics over 100 hours with complete disaggregation of cell integrity in the PBS

group contrary to overall preserved cell structure in NAD$^+$ treated BMDMs (**Fig. 2D, Fig. 2**

Supplement 2, Mov. 1). To rule out, that NAD$^+$ impairs LPS internalization into cells, BMDMs

were stimulated with CTB and LPS that was coupled to a fluorescent reporter (FITC) and transfection

effectivity was assessed by fluorescence microscopy and flow cytometry.

Our data indicated no significant difference between the PBS and NAD$^+$ treated group (**Fig. 2E**),

suggesting that NAD$^+$ does not alter LPS internalization. Notably, BMDMs only stimulated with LPS

showed no internalization of LPS consistent with previous reports[12].

Casp-4 and 5 have been delineated as the human homolog of casp-11 in mice carrying out the

same effector functions including pyroptosis induction and IL-1β secretion[24]. As clinical

relevance, we therefore tested whether NAD$^+$ was also able to block the non-canonical pathway

in human macrophages. Hence, human macrophages were differentiated from PBMC and treated

with NAD$^+$ followed by intracellular LPS transfection (Fugene) and IL-1β secretion and

cytotoxicity were quantified. The results indicated that NAD$^+$ treatment significantly dampened

both, IL-1β secretion and pyroptosis (**Fig. 2F**), underscoring its therapeutic potential.

Collectively, our results suggest that NAD$^+$ acts directly on macrophages by targeting

specifically the non-canonical inflammasome signaling machinery.

NAD$^+$ inhibits the non-canonical inflammasome via STAT-1/IFN-β pathway blockade

Although our data emphasized that NAD$^+$ blocks the non-canonical inflammasome pathway, the

underlying mechanisms remained yet to be determined. Therefore, we performed RNA-

sequencing of Pam3CSK4 primed BMDMs that were treated with PBS or NAD$^+$ and

subsequently stimulated with CTB + LPS O111:B4. Interestingly, when blotting gene expression

differences in a Venn diagram, we found strikingly more genes commonly expressed in the

NAD$^+$ and control group when compared to the PBS treated group (**Fig. 3A**). Gene ontology

enrichment analysis revealed a significant downregulation of genes involved in the antiviral

response in addition to the cellular response to the type-I-IFN, IFN-β, when comparing NAD$^+$

and PBS treated groups (**Fig. 3B**). Type-I-IFN are known to promote the expression of over

2,000 IFN-stimulated genes (ISGs), translated into ISGs-induced proteins which have been

shown to act by enhancing pathogen detection and restrict their replication[25]. Recently, it was

reported that type-I-IFNs are required for casp-11 expression contributing to non-canonical

inflammasome activation[26, 27]. Consistently, LPS-stimulated macrophages from TRIF-deficient

mice displayed impaired casp-11 expression, implying a context-dependent role for type-I-IFN in

the regulation of caspase-11 activity[26]. Indeed, when comparing expression of genes involved in

IFN-β signaling through cluster analysis we found a significant decrease in a broad range in

genes in the NAD$^+$ treated group (**Fig. 3C**). Most strikingly, GTPases and guanylate binding

proteins involved in the downstream signaling of IFN-β were significantly downregulated (**Fig.**

3C and Fig. 3D) while IFN-β-receptor 2 expression remained unaffected (**Fig. 3C**). Recently,

IFN-inducible GTPases and guanylate binding proteins have been assigned a crucial role for the

intracellular recognition of LPS and linked caspase-11 activation[27, 28]. Thus, to test if NAD$^+$

mediated non-canonical inflammasome blockade via IFN-β, NAD$^+$ or PBS treated BMDMs were

primed with Pam3CSK4 and subsequently stimulated with LPS O111:B4 + CTB and 1000 U/ml

of recombinant IFN-β. Strikingly, administration of recombinant IFN-β resulted in a complete

reversal of NAD$^+$ mediated blockade of IL-1β secretion and pyroptosis (**Fig. 3E**). Moreover,

IFN-β administration restored casp-11, NLRP3 and GSDMD expression in the NAD$^+$ treated

group (**Fig. 3F**). It is well established that STAT-1 phosphorylation constitutes the link between

intracellular type-I-IFN signaling and the transcription of ISGs through nuclear translocation[29,]

[30]. Notably, our RNA-seq data indicated a significant downregulation of STAT-1 (**Fig. 3C**).

Moreover, we have previously shown that NAD$^+$ administration dampens the expression and

activation of transcription factors such as STAT-5[16]. To test, whether NAD$^+$ blocks IFN-β

signaling via STAT-1, BMDMs were subjected to NAD$^+$ or PBS followed by non-canonical

inflammasome stimulation and recombinant IFN-β. After 16 hours STAT-1 expression and

phosphorylation were assessed by Western blotting. Consistent with our previous results, NAD$^+$

treatment downregulated expression levels of STAT-1 and phospho-STAT-1. In contrast,

addition of recombinant IFN-β treatment to NAD$^+$ treated BMDMs restored STAT-1 and

phospho-STAT-1 expression that was equivalent to the PBS treated group (**Fig. 3G**). Taken

together, our data indicate that NAD$^+$ impedes non-canonical inflammasome activation via IFN-

β /STAT-1 blockade (**Fig. 4**).

NAD$^+$ increases Caspase-1 KO mice resistance to endotoxic shock via systemic IL-10

production

Caspase-11 KO mice have been reported to be resistant towards lethal doses of LPS inducing septic

shock[12]. However, upon priming with TLR3 instead of a TLR4 ligand, Casp-11 KO mice merely

exhibit partial resistance towards LPS-induced shock with a 50–60% survival rate[12, 31]. Our data

indicate that NAD$^+$ prevents LPS-induced cell death via the non-canonical inflammasome

pathway and casp-11 blockade. We thus tested whether NAD$^+$ could achieve similar protection

against septic shock in WT vs casp-11 KO mice. Casp-11 KO mice were intraperitoneally injected

with NAD$^+$ and PBS and treated with 6 mg/kg poly(I:C) 6 hours prior LPS administration.

Consistent with previous studies the results indicated a modest resistance of casp-11 KO mice (40%

survival). In high contrast, both WT and casp-11 KO mice subjected to NAD$^+$ exhibited 85%-100%

survival, respectively when compared to casp-11 KO mice that were treated with PBS, suggesting

the existence of an alternative protective pathway against septic shock that is casp-11

independent. WT mice, treated with 6 mg/kg poly(I:C) followed by LPS (54mg/kg)

administration not only survived but fully recovered 7 days later, underscoring the unique and

robust therapeutic effect of NAD$^+$ in septic shock.

Previous studies have reported inferior outcomes of IL-10$^{-/-}$ mice in septic shock[32, 33] pointing

out a 20-fold lower lethal dose of LPS compared to WT mice[33]. Moreover, IL-10 itself has been

shown to prevent mice from septic shock induced death after a single administration[34]. We have

previously delineated immunosuppressive properties of NAD$^+$ via a systemic production of IL-

10, a robust immunosuppressive cytokine. In addition, we have described the pivotal role of

NAD$^+$ protecting towards EAE and allograft rejection via an increased frequency of IL-10

producing CD4$^+$ T cells[15, 16]. To test if IL-10 plays an additional protective role in the context of

NAD$^+$ mediated protection towards LPS-induced death, WT mice treated with NAD$^+$ or PBS

were subjected to intraperitoneal LPS injection (54mg/kg) and IL-10 expression by

macrophages, dendritic cells and T cells was assessed 15 hours after LPS administration.

Consistent with our previous studies[15, 16], we found significantly augmented frequencies of IL-10

producing CD4$^+$ and CD8$^+$ T cells (**Fig. 5C**). Moreover, we detected a dramatic increase of IL-

10 production by macrophages, but not the DC population (**Fig. 5B**). Interestingly, IL-10 has

been described to inhibit macrophage function and pro-inflammatory cytokine production in

both, human[35] and mice[36]. Moreover, autocrine IL-10 secretion of macrophages was found to

decrease pro-IL-1β concentration by promoting signal transducer activator of transcription-3

(STAT-3) expression[37]. To investigate the potential autocrine impact of an augmented IL-10

production on macrophage self-regulation, we administered combined IL-10 neutralizing

antibody and IL-10 receptor antagonist to BMDMs primed with Pam3CSK4 and stimulated with

CTB and LPS O111:B4. The results showed that neutralization of the autocrine IL-10 signaling

pathway dampened NAD$^+$ mediated decrease of IL-1β secretion and reversed pyroptotic cell

death partially (**Fig. 5D**). To further investigate the relevance of our in vitro findings, IL-10$^{-/-}$

mice were treated with NAD$^+$ or PBS, subjected to LPS (54mg/kg) and survival was monitored.

Consistent with previous reports[32, 33], mice lacking IL-10 exhibited an inferior protection against

septic shock when compared to WT animals. More importantly, IL-10$^{-/-}$ mice subjected to NAD$^+$

exhibited a compromised survival (**Fig. 5E**) suggesting that systemic production of IL-10

following NAD$^+$ administration plays a pivotal role in NAD$^+$-mediated protection against septic

shock.

Discussion

Previously, we have delineated the protective role of NAD^+ in the context of *L. m.* infection, a

gram-positive bacterium[17]. However, it remained unclear whether NAD^+ conveyed resistance

towards *L. m.* by an augmented bacterial clearance or rather through its immunosuppressive

effects dampening pathological systemic inflammation. Although the cell membrane of *L. m.* has

been shown to bear lipoteichoic acids, which resemble the endotoxin LPS from gram-negative

bacteria in both, structure and function, it is widely considered as an intracellular bacterium[38]. In

our current study, we administered a lethal dose of pathogenic *E. coli*, that is well known to

promote septic shock, and showed that NAD^+ also protected towards a lethal dose of this gram-

negative bacterium. More importantly, we demonstrate that NAD^+ conveys protection towards

septic shock by specifically inhibiting the non-canonical inflammasome but not via bacterial

clearance. Mechanistically, NAD^+ impedes pro-casp-11 and casp-11 expression in macrophages

blocking non-canonical derived GSDMD cleavage and NLRP3 inflammasome activation, thus

inhibiting pyroptotic cell death and pro-inflammatory cytokine release. The resistance of NAD^+

treated WT mice against *E. coli* and LPS induced septic shock reflected the robust inhibitory

effect observed in vitro of NAD^+ on the non-canonical inflammasome signaling machinery.

Until now, the exact mechanism how pro-casp-11 expression and its maturation to casp-11 is

regulated remains unclear. Given the low basal expression of both pro-casp-11 and casp-11[39], a

priming signal is required for initiating the non-canonical inflammasome pathway and

macrophage sensing of intracellular LPS[40]. Previous work has demonstrated that transcriptional

induction of the pro-casp-11 isoforms p42 and p38 in macrophages requires type I IFN

stimulation[39, 41] while IFN-β has been shown to promote transcriptional induction and processing

of caspase-11[26]. In line with these findings, CTB treatment of macrophages primed with

Pam3csk4 failed to elicit IL1-β release compared to LPS primed controls while exogenous

administration of IFN-β in turn restored CTB-induced IL-1β production[26] underscoring the

transcriptional role of type I IFN. Our RNA-seq results indicated a dampened cellular response

towards IFN-β while western blotting revealed a significant downregulation of both, pro-casp-11

and casp-11 suggesting a transcriptional downregulation of both enzymes. Consistently, NAD^+

decreased STAT-1 expression and phosphorylation, which constitutes the mechanistic link

between extracellular type I IFN stimulation and transcriptional effects through translocation of

phosphorylated STAT-1 to the nucleus inducing ISGs[30]. Thus, treatment of stimulated

macrophages subjected to NAD^+ with recombinant IFN-β restored STAT-1 signaling, caspase-11

expression and GSDM cleavage which translated into reconstituted IL-1β production and LDH

release. Collectively, NAD^+ mitigates the intracellular response to IFN-β that leads to non-

canonical inflammasome induction by suppressing macrophage derived STAT-1 expression and

phosphorylation.

Furthermore, we showed that NAD^+ treatment improved resistance of casp-11 KO mice towards

poly I:C primed septic shock. More importantly, WT mice treated with NAD^+ exhibited 100%

survival while casp-11 KO mice treated with PBS exhibited a modest 40% survival, suggesting that

NAD^+ promotes survival beyond non-canonical inflammasome blockade. Our previous studies

have delineated the effects of NAD^+ on various immune cells such as dendritic cells and $CD4^+$ T

cells including Th1, Th17, regulatory type 1 (Tr1) and Treg cells communicated exclusively

through mast cells (MCs)[15, 16, 17]. Thereby, NAD^+ treatment promoted MC derived induction of

TR1 cells that resulted into increased systemic levels of IL-10. Latter one was found to play a

cardinal feature during bacterial infection as $MC^{-/-}$ mice were more susceptible to *L. m.* infection

than WT animals when treated with NAD^+. Here, we found a direct effect of NAD^+ on

macrophages by specifically inhibiting the non-canonical inflammasome and promoting IL-10

production. Polymorphisms in the IL-10 locus or IL-10R deficiencies have been linked to severe

intestinal inflammatory diseases in both, human and mice[42, 43, 44, 45]. More importantly, mice

deficient for IL-10 have been shown to display elevated inflammasome activation and IL-1β

production resulting in severe colitis[46] or Ag-induced arthritis[47]. When inhibiting the autocrine

pathway for IL-10 through combined receptor antagonization and IL-10 neutralization, we found

a pronounced increase of IL-1β production of NAD$^+$ treated macrophages stimulated with CTB

and LPS (Fig. 4D). This is consistent with previous reports showing that autocrine IL-10

signaling interferes with the transcription of pro-IL-1β[37]. LDH release in turn, was only restored

partly possibly due to missing effects of second party leucocytes secreting IL-10 in vivo such as

Tr1 cells which have been shown to inhibit the transcription of IL-1β and inflammasome

mediated activation of caspase-1[48].

More recently, casp-8, that plays a central role in apoptosis, has been reported as an important

mediator of endotoxemia resistance and LPS-driven systemic inflammation. Since our RNA-

sequencing results revealed a dramatically attenuated cellular response towards type-I-IFN with

downregulation of various interferon regulatory factors, that have been reported as major

regulators of casp-8[49, 50] it is possible that NAD$^+$ may exert protection against septic shock by

altering caspase-8 expression as well. Although, we have previously reported the protective

effect of NAD$^+$ against apoptosis of activated CD4$^+$ T cells[15], it remains yet to be determined

how NAD$^+$ impacts executioner proteins of other cell death processes such as apoptosis and

necroptosis.

Notably, both casp-8 and casp-11 have been found dispensable in the hematopoietic

compartment that produces the pro-inflammatory cytokines necessary to initiate shock[51]. Thus,

NAD$^+$ treatment may improve resistance of casp-11 KO mice to septic shock by also dampening the

initiating pro-inflammatory cytokine cascade via its systemic IL-10 cytokine production.

Importantly, while inhibiting macrophage derived inflammasome function, NAD$^+$ does not

interfere with NF-κB signaling which has been shown to promote various inflammatory and

autoimmune diseases when dysregulated[52].

Taken together, we dissected the dichotomous capacity of NAD$^+$ to dampen auto- and allo-

immunity while concomitantly protecting towards severe bacterial infection, outlining its unique

effects in the context of septic shock.

Material and Methods

Animals

Young (8-12 weeks) C57BL/6, B6.129P2-IL10tm1Cgn/J and B6.129S4(D2)-Casp4tm1Yuan/J

mice were purchased from Charles River Laboratory, Wilmington, MA. All mice were male,

age-matched and experimental and control animals were housed separately. Animals and

samples were randomly assigned to either the control or treatment group to ensure biological

diversity. The study protocol was approved by the Brigham and Women´s Hospital Institutional

Animal Care and use Committee (IACUC) (animal protocol #2018N000049). All mice were

male, age-matched and experimental and control animals were housed separately. Owing to the

exploratory nature of our study, we did not use randomization and blinding. No statistical

methods were used to predetermine sample size. All animals were maintained in specific

pathogen-free (SPF) conditions at the Brigham and Women´s Hospital animal facility in

accordance with federal, state, and institutional guidelines. Animals were maintained on 12-h

light, 12-h dark cycle in facilities with an ambient temperature of 19–22 °C and 40–60%

humidity and were allowed free access to water and standard chow. Euthanasia was performed

by cervical dislocation following anesthesia with isoflurane (Patterson Veterinary, Devens, MA,

USA).

Murine Bone Marrow Derived Macrophage Differentiation and Culture

8-12-week-old C57BL/6 mice were euthanized by cervical dislocation, sprayed with alcohol and

skin was removed to expose femurs. The femur was flushed with ice cold PBS and the obtained

bone marrow was filtered through 70um Nylon cell strainer. After washing with PBS, red blood

cell lysis was performed using ammonium-chloride-potassium-solution (Fisher scientific) and

the reaction was blocked with complete Dulbecco's modified eagle medium (DMEM) (Fisher

Scientific) supplemented with 10% endotoxin-free bovine serum and PS. To minimize fibroblast

contamination cells were cultured in complete DMEM at 37°C, 5% CO_2 and non-adherent cells

were collected after 30 minutes.

Bone marrow cells were then differentiated into macrophages in DMEM supplemented with 10%

endotoxin-free bovine serum, PS and 40ng/ml murine GM-CSF (Abcam) for 8 days. Medium

was changed every 2 days to remove non-adherent cells.

Canonical and Non-canonical Inflammasome Activation in Murine Macrophages

After 8 days of culture the medium was replaced by 40ng/ml GM-CSF containing 100µmol

NAD^+ culture medium. For 2 following days NAD^+ was added daily until stimulation. To induce

canonical and non-canonical inflammasome activation in murine macrophages, NAD^+ treated

and control BMDMs were cultured overnight in a 24 well plate at $1x10^6$ cells/ml and afterwards

primed with 1µg/ml Pam3CSK4 or 1µg/ml LPS O111:B4 (Sigma) for 5-6 hours. Primed

BMDMs were then stimulated for 16 hours with either 5 mmol ATP (canonical inflammasome

stimulation) or 2µg/ml LPS O111:B4 and 20µg/ml CTB (Sigma) to allow LPS entry (non-

canonical inflammasome stimulation) where indicated. To test the effect of NAD^+ on type 1 IFN

and STAT1 signaling, BMDMs were cultured overnight in a 24 well plate at $1x10^6$ cells/ml and

afterwards primed with 1µg/ml Pam3CSK4 or 1µg/ml LPS O111:B4 (Sigma) for 5-6 hours.

Primed BMDMs were then stimulated for 16 hours 2µg/ml LPS O111:B4, 20µg/ml CTB and

U/ml recombinant IFN-β.

ELISA

Expression of macrophage derived murine IL-1β, IL-18 and human IL-1β was analyzed in the cell culture supernatant by commercial ELISA kits (Invitrogen) following the manufacturer's recommended procedures.

Pyroptosis Assay

Pyroptotic cell death was measured by assessing LDH-release in the cell culture supernatant of human and murine macrophages using a CytoTox 96 Non-radioactive Cytotoxic Assay (Promega) following the manufacturer's recommended procedures.

RNA Extraction and RNA-Sequencing

BMDMs were harvested and differentiated as outlined in the particular section. After 8 days of culture the medium was replaced by 40ng/ml GM-CSF containing culture medium (control group) or 40ng/ml GM-CSF containing 100µmol NAD^+ culture medium (NAD^+ treated group). For 2 following days NAD^+ was added daily. Subsequently, NAD^+ treated and control BMDMs were cultured overnight in a 24 well plate at $1x10^6$ cells/ml and afterwards primed with 1µg/ml Pam3CSK4 or 1µg/ml LPS O111:B4 (Sigma) for 5-6 hours. Primed BMDMs were then stimulated for 16 hours with 2µg/ml LPS O111:B4 and 20µg/ml CTB (Sigma) to allow LPS entry. Another set of BMDMs were differentiated without additional treatment serving as naive controls.

Subsequently, RNA was extracted with the RNAqueous extraction kit (Applied Biosystems),

according to the manufacturer's protocols. Briefly, cells were homogenized in lysis buffer (total

volume of 0.5 mL) and passed through a column. After successive washes, RNA was eluted.

RNA sequencing was commercially performed by Novogene CO., Ltd. In brief, mRNA was

enriched from total RNA using oligo(dT) beads and subsequently fragmented randomly in

fragmentation buffer, followed by cDNA synthesis using random hexamers and reverse

transcriptase. After first-strand synthesis, a custom second-strand synthesis buffer (Illumina) was

added with dNTPs, RNase H and Escherichia coli polymerase I to generate the second strand by

nick-translation. The final cDNA library is ready after a round of purification, terminal repair, A-

tailing, ligation of sequencing adapters, size selection and PCR enrichment.

Isolation and Differentiation of Human Macrophages from PBMCs

Blood was obtained from healthy adult volunteers with the only purpose to isolate PBMCs in order to

create a basis for macrophage cultures. Blood withdrawal was performed in accordance with guidelines

of and approved by the Institutional Review Board of the Charité Universitätsmedizin Berlin

(EA4/006/22). Informed consent and consent to publish was obtained from each volunteer in

accordance with the Declaration of Helsinki. All personal information collected from volunteers were

treated with utmost confidentiality. For experiments on human macrophages, PBMC were isolated by

performing a density centrifugation in SepMate tubes (Stem cell) using lymphoprep (Stem cell) density

gradient medium. PBMC were then plated in DMEM culture medium supplemented with standard

antibiotics, 10% FCS and human 50ng/ml GM-CSF (peprotech) at a density of $1x10^6$ cells/ml. The

medium was changed every 2-3 days until the cells reached a full confluence.

453 Non-canonical Inflammasome Induction in Human Macrophages

After 8 days of culture the medium was replaced by 50ng/ml GM-CSF containing 100µmol

NAD^+ culture medium. For 2 following days NAD^+ was added daily until stimulation. To induce

non-canonical inflammasome activation in human macrophages, cells were primed with 1µg/ml

Pam3CSK4 for 5-6 hours. Subsequently the medium was replaced, and cells were treated with

3µg/ml LPS O111:B4 and 0.25% (vol/vol) Fugene HD Plus (Promega) to cause transfection.

Finally, plates were centrifuged at 805 x g for 2 minutes and subsequently cultured for 20 hours

at 37°C, 5% CO_2.

462 Western blot

For Western blot analysis, proteins were extracted using RIPA buffer and the concentrations

determined using Pierce™ BCA Protein Assay Kit. Subsequently, proteins were resolved in SDS-

PAGE, transferred to 0.45 μm nitrocellulose membranes (BioRad), blocked with 5% non-fat dry

milk in PBS with 0.1% Tween 20, and processed for immunodetection. The following primary

antibodies were used according to manufacturer's instructions: Pro-Caspase-1 (#ab179515,

Abcam), Caspase-1 (#14-9832-82, eBioscience) IL-1β (AF-401-NA, RD Systems), NLRP3

(#768319, RD Systems), Caspase-11 (#mab8648, RD Systems), Gasdermin D (ab209845,

Abcam), P-STAT-1 (#9167S, Cell Signaling), STAT-1 (#9172S, Cell Signaling), NF-κB-p65

(#49445S, Cell Signaling), NF-κB-p52 (#4882S, Cell Signaling), β-Actin (ab3280, Abcam).

Antibody detection was performed with HRP-coupled goat secondary anti-mouse or anti-rabbit

antibodies (Immunoresearch), followed by ECL reaction (Perkin Elmer) and exposure to Fuji X-

ray films. Finally, films were scanned, and signals quantified using the web-based image

processing software ImageJ (NIH).

Analysis of LPS-Transfection Efficiency

For intracellular detection of LPS, primed BMDMs were stimulated with 20µg/ml CTB and

FITC-conjugated LPS O111:B4 for 16 hours, washed twice with PBS, fixed in 4% PFA

containing PBS and DAPI for 10 minutes and subsequently analyzed using a confocal

microscope and flow cytometry. To determine transfection efficiency using confocal

microscopy, FITC-stained pixels per image were quantified using the web-based image

processing software ImageJ (NIH).

Caspase-1 assay

To determine the expression of Caspase-1, primed BMDMs were stimulated with 20µg/ml CTB

and 2µg/ml LPS O111:B4 for 4 and 16 hours respectively, washed twice with PBS, stained using

a caspase-1 active staining kit (Abcam) including caspase-1 staining (fluorescent green) and

DAPI staining (fluorescent blue) according to the manufacturer's protocol and analyzed using a

confocal microscope.

Endotoxic shock model

8-12-week-old C57BL/6 mice were treated with 40mg NAD$^+$ for 2 following days before

intraperitoneal injection of 54 mg/kg LPS O111:B4 or LPS O55:B5. Where indicated mice were

administered 6mg/kg poly(I:C) 6 hours prior to LPS administration. Consequently, survival and

body temperature were monitored every 2-4 hours for up to 100 hours. To assess the amount of

systemic IL-1β and IL-18 by Elisa (both Invitrogen), mice were euthanized by decapitation 10

hours and 15 hours after LPS injection serum was isolated from collected blood.

Flow cytometric analysis

To analyze splenic lymphocytes for the intracellular expression of IL-10 mice were challenged

with 54 mg/kg LPS O111:B4 for 10 hours and euthanized by cervical dislocation subsequently.

Spleens were harvested in a sterile environment and single cell suspensions were obtained using

a 70um Nylon cell strainer.

Then, 1×10^6 splenocytes per animal per condition were cultured in RPMI 1640 (#10-040-CV,

Corning) supplemented with 10% BenchMark Fetal Bovine Serum (#100-106, Gemini), 1%

penicillin/streptomycin (#30-002-CI, Corning), 2 mM L-glutamine (#25-005-CI, Corning), 20

ng/mL phorbol 12-myristate 13-acetate (PMA, #P8139-1MG, Sigma-Aldrich), 1 μg/mL

ionomycin (#I9657-1MG, Sigma-Aldrich), and 0.67 μL/mL BD GolgiStop (#554724, BD

Biosciences) for 4 hours at 37°C and 5% CO2 in 1 mL-volumes in a 12-well plate. After 4 hours,

the cells were collected from each 12-well plate well and prepared for flow cytometry by

staining the surface epitopes in flow staining buffer consisting of $1\times$ DPBS supplemented with

1.0% (w/v) bovine serum albumin (#A2153, Sigma-Aldrich) and 0.020% sodium azide (#S8032,

Sigma-Aldrich) for 25 min at 4°C. Then, the cells were fixed and permeabilized with the

eBioscience Foxp3 Fixation/Permeabilization concentrate and diluent cocktail (#00-5523-00,

Invitrogen) for 30 min at 4°C. Finally, the intracellular cytokine target was stained in $1\times$

permeabilization buffer diluted from $10\times$ eBioscience Foxp3 Permeabilization Buffer (#00-5523-

00, Invitrogen) with deionized water. Finally, the stained samples were analyzed on a FACS

Canto II (BD Biosciences, San Jose, CA, United States) flow cytometer, and the resultant flow

cytometry standard (FCS) files were analyzed with FlowJo version 10 (Flowjo LLC, Ashland,

OR, United States).

Bacterial infection model

Frozen stock suspensions of Escherichia coli (Migul) (ATCC, 700928) were obtained from ATCC and cultured in 5ml Luria-Bertani medium at 37°C. Bacterial concentration was determined by plating 100ul, 10-fold serial diluted bacterial samples and counting the colony-forming units (CFU) after overnight incubation at 37°C. One day prior to injection 1ml of culture was reinoculated into 5ml of medium and incubated for 3-5 hours using a 37°C shaker at 250rpm agitation. Bacterial cultures were then centrifuged for 10 minutes at 3000rpm and washed twice with PBS. Mice were previously treated with NAD^+ for 2 serial days and subsequently infected with E. coli by injecting 1ml of $1x10^9$ CFU/ml bacterial suspension intraperitoneally. The survival was monitored. In another set of experiments mice were sacrificed 5h hours after infection and kidneys and liver were harvested. The collected tissues were homogenized in 1ml of sterile PBS and 10-fold serial dilutions plated overnight at 37°C on LB agar plates to determine bacterial load per gram.

Data Availability Statement

All data generated and analyzed during this study are provided with the manuscript as source data files for each figure. Data generated from RNA sequencing of BMDMs have been made publicly available in Dryad (DOI: 10.5061/dryad.zw3r228fj).

Acknowledgements

J.I. was supported by the Berlin Institutes of Health Junior Clinician Scientist Program. Y.N. was supported by the Chinese Scholarship Council (201606370196) and Central South University. H.R.C.B. was supported by the Swiss Society of Cardiac Surgery. A.V. was supported by awards from the National Institute of Mental Health (R01MH110438) and National Institute of

557   Neurological Disorders and Stroke (R01NS100808).

556   **Competing interests**

557   The authors declare no conflict of interest

References

568
569   1.   Kaukonen KM, Bailey M, Pilcher D, Cooper DJ, Bellomo R. Systemic inflammatory response
570   syndrome criteria in defining severe sepsis. *N Engl J Med* **372**, 1629-1638 (2015).

571
572   2.   Ramachandran G. Gram-positive and gram-negative bacterial toxins in sepsis: a brief review.
573   *Virulence* **5**, 213-218 (2014).

574
575   3.   Kumar V. Inflammasomes: Pandora's box for sepsis. *Journal of inflammation research* **11**, 477-
576   502 (2018).

577
578   4.   Lopez-Castejon G, Brough D. Understanding the mechanism of IL-1beta secretion. *Cytokine*
579   *Growth Factor Rev* **22**, 189-195 (2011).

580
581   5.   Yi YS. Caspase-11 non-canonical inflammasome: a critical sensor of intracellular
582   lipopolysaccharide in macrophage-mediated inflammatory responses. *Immunology* **152**, 207-217
583   (2017).

584
585   6.   Ding J, Shao F. SnapShot: The Noncanonical Inflammasome. *Cell* **168**, 544-544.e541 (2017).

586
587   7.   Shi J, *et al.* Cleavage of GSDMD by inflammatory caspases determines pyroptotic cell death.
588   *Nature* **526**, 660-665 (2015).

589
590   8.   Man SM, *et al.* Differential roles of caspase-1 and caspase-11 in infection and inflammation. *Sci*
591   *Rep* **7**, 45126 (2017).

592
593   9.   Kayagaki N, *et al.* Caspase-11 cleaves gasdermin D for non-canonical inflammasome signalling.
594   *Nature* **526**, 666-671 (2015).

595
596   10.   Kayagaki N, *et al.* Non-canonical inflammasome activation targets caspase-11. *Nature* **479**, 117-
597   121 (2011).

598
599   11.   Yang D, He Y, Munoz-Planillo R, Liu Q, Nunez G. Caspase-11 Requires the Pannexin-1 Channel
600   and the Purinergic P2X7 Pore to Mediate Pyroptosis and Endotoxic Shock. *Immunity* **43**, 923-932
601   (2015).

603 12. Kayagaki N, *et al.* Noncanonical inflammasome activation by intracellular LPS independent of
604   TLR4. *Science* **341**, 1246-1249 (2013).

605
606 13. Angus DC, van der Poll T. Severe sepsis and septic shock. *N Engl J Med* **369**, 840-851 (2013).

607
608 14. Fraser CC. Exploring the positive and negative consequences of NF-kappaB inhibition for the
609   treatment of human disease. *Cell cycle (Georgetown, Tex)* **5**, 1160-1163 (2006).

610
611 15. Tullius SG, *et al.* NAD+ protects against EAE by regulating CD4+ T-cell differentiation. *Nat*
612   *Commun* **5**, 5101 (2014).

613
614 16. Elkhal A, *et al.* NAD(+) regulates Treg cell fate and promotes allograft survival via a systemic IL-
615   10 production that is CD4(+) CD25(+) Foxp3(+) T cells independent. *Sci Rep* **6**, 22325 (2016).

616
617 17. Rodriguez Cetina Biefer H, *et al.* Mast cells regulate CD4(+) T-cell differentiation in the absence
618   of antigen presentation. *J Allergy Clin Immunol* **142**, 1894-1908.e1897 (2018).

619
620 18. Mellata M, Mitchell NM, Schodel F, Curtiss RR, Pier GB. Novel vaccine antigen combinations
621   elicit protective immune responses against Escherichia coli sepsis. *Vaccine* **34**, 656-662 (2016).

622
623 19. Dogan MD, Ataoglu H, Akarsu ES. Effects of different serotypes of Escherichia coli
624   lipopolysaccharides on body temperature in rats. *Life Sci* **67**, 2319-2329 (2000).

625
626 20. Bullock B, Benham MD. Bacterial Sepsis. In: *StatPearls*). StatPearls Publishing

627   StatPearls Publishing LLC. (2019).

628
629 21. Rossaint J, Zarbock A. Pathogenesis of Multiple Organ Failure in Sepsis. *Crit Rev Immunol* **35**,
630   277-291 (2015).

631
632 22. Evans TJ. The role of macrophages in septic shock. *Immunobiology* **195**, 655-659 (1996).

633
634 23. Chan FK, Moriwaki K, De Rosa MJ. Detection of necrosis by release of lactate dehydrogenase
635   activity. *Methods Mol Biol* **979**, 65-70 (2013).

636
637 24. Shi J, *et al.* Inflammatory caspases are innate immune receptors for intracellular LPS. *Nature*
638   **514**, 187-192 (2014).

639
25. Schneider WM, Chevillotte MD, Rice CM. Interferon-stimulated genes: a complex web of host
defenses. *Annu Rev Immunol* **32**, 513-545 (2014).

26. Rathinam VA, *et al.* TRIF licenses caspase-11-dependent NLRP3 inflammasome activation by
gram-negative bacteria. *Cell* **150**, 606-619 (2012).

27. Tang Y, *et al.* TRIF signaling is required for caspase-11-dependent immune responses and
lethality in sepsis. *Mol Med* **24**, 66 (2018).

28. Pilla DM, *et al.* Guanylate binding proteins promote caspase-11-dependent pyroptosis in
response to cytoplasmic LPS. *Proc Natl Acad Sci U S A* **111**, 6046-6051 (2014).

29. Stark GR, Darnell JE, Jr. The JAK-STAT pathway at twenty. *Immunity* **36**, 503-514 (2012).

30. Ivashkiv LB, Donlin LT. Regulation of type I interferon responses. *Nat Rev Immunol* **14**, 36-49
(2014).

31. Hagar JA, *et al.* Lipopolysaccharide Potentiates Insulin-Driven Hypoglycemic Shock. *J Immunol*
199, 3634-3643 (2017).

32. Latifi SQ, O'Riordan MA, Levine AD. Interleukin-10 controls the onset of irreversible septic shock.
Infect Immun **70**, 4441-4446 (2002).

33. Berg DJ, *et al.* Interleukin-10 is a central regulator of the response to LPS in murine models of
endotoxic shock and the Shwartzman reaction but not endotoxin tolerance. *J Clin Invest* **96**,
2339-2347 (1995).

34. Howard M, Muchamuel T, Andrade S, Menon S. Interleukin 10 protects mice from lethal
endotoxemia. *J Exp Med* **177**, 1205-1208 (1993).

35. de Waal Malefyt R, Abrams J, Bennett B, Figdor CG, de Vries JE. Interleukin 10(IL-10) inhibits
cytokine synthesis by human monocytes: an autoregulatory role of IL-10 produced by
monocytes. *J Exp Med* **174**, 1209-1220 (1991).

36. Fiorentino DF, Zlotnik A, Mosmann TR, Howard M, O'Garra A. IL-10 inhibits cytokine production
by activated macrophages. *J Immunol* **147**, 3815-3822 (1991).

37. Sun Y, *et al.* Interleukin-10 inhibits interleukin-1beta production and inflammasome activation of
microglia in epileptic seizures. *J Neuroinflammation* **16**, 66 (2019).

38. Farber JM, Peterkin PI. Listeria monocytogenes, a food-borne pathogen. *Microbiol Rev* **55**, 476-
511 (1991).

39.   Schauvliege R, Vanrobaeys J, Schotte P, Beyaert R. Caspase-11 gene expression in response to
684   lipopolysaccharide and interferon-gamma requires nuclear factor-kappa B and signal transducer
685   and activator of transcription (STAT) 1. *J Biol Chem* **277**, 41624-41630 (2002).

687   40.   Yang J, Zhao Y, Shao F. Non-canonical activation of inflammatory caspases by cytosolic LPS in
688   innate immunity. *Curr Opin Immunol* **32**, 78-83 (2015).

690   41.   Yen JH, Ganea D. Interferon beta induces mature dendritic cell apoptosis through caspase-
691   11/caspase-3 activation. *Blood* **114**, 1344-1354 (2009).

693   42.   Franke A, *et al.* Sequence variants in IL10, ARPC2 and multiple other loci contribute to ulcerative
694   colitis susceptibility. *Nat Genet* **40**, 1319-1323 (2008).

696   43.   Franke A, *et al.* Genome-wide meta-analysis increases to 71 the number of confirmed Crohn's
697   disease susceptibility loci. *Nat Genet* **42**, 1118-1125 (2010).

699   44.   Kuhn R, Lohler J, Rennick D, Rajewsky K, Muller W. Interleukin-10-deficient mice develop chronic
700   enterocolitis. *Cell* **75**, 263-274 (1993).

702   45.   Begue B, *et al.* Defective IL10 signaling defining a subgroup of patients with inflammatory bowel
703   disease. *Am J Gastroenterol* **106**, 1544-1555 (2011).

705   46.   Zhang J, Fu S, Sun S, Li Z, Guo B. Inflammasome activation has an important role in the
706   development of spontaneous colitis. *Mucosal Immunol* **7**, 1139-1150 (2014).

708   47.   Greenhill CJ, *et al.* Interleukin-10 regulates the inflammasome-driven augmentation of
709   inflammatory arthritis and joint destruction. *Arthritis Res Ther* **16**, 419 (2014).

711   48.   Yao Y, *et al.* Tr1 Cells, but Not Foxp3+ Regulatory T Cells, Suppress NLRP3 Inflammasome
712   Activation via an IL-10-Dependent Mechanism. *J Immunol* **195**, 488-497 (2015).

714   49.   Apelbaum A, Yarden G, Warszawski S, Harari D, Schreiber G. Type I interferons induce apoptosis
715   by balancing cFLIP and caspase-8 independent of death ligands. *Mol Cell Biol* **33**, 800-814 (2013).

717   50.   Newton K, *et al.* Cleavage of RIPK1 by caspase-8 is crucial for limiting apoptosis and necroptosis.
718   *Nature*, (2019).

720   51.   Mandal P, *et al.* Caspase-8 Collaborates with Caspase-11 to Drive Tissue Damage and Execution
721   of Endotoxic Shock. *Immunity* **49**, 42-55.e46 (2018).

722
723   52.   Liu T, Zhang L, Joo D, Sun SC. NF-kappaB signaling in inflammation. *Signal transduction and*
724   *targeted therapy* **2**, (2017).
725
726

Figure Legends

Figure 1. *NAD+ protects mice from lethal bacterial infection and endotoxic shock by dampening systemic inflammation*

(**A**) mice were treated with PBS or NAD+ prior to administration of a lethal dose of either pathogenic *E. coli* or LPS by intraperitoneal injection. (**B**) Kidneys and Livers were removed after and bacterial load was determined by counting CFU (**C**) Survival was monitored over 48 hours after bacterial infection and (**D**) LPS injection of both serotypes (n=6, 3 independent survival experiments). In addition, body temperature was monitored in the kinetics of up to 100 hours (**E**) Lungs, Kidneys and Livers were removed IHC was performed for H&E staining (**F**) Systemic levels (serum) of IL-6, TNFα, IL-1β and IL-18 were assessed by ELISA. Column plots display mean with standard deviation (n=5). Statistical significance was determined by using Student's T-test or One-Way-ANOVA while survival data were compared using log-rank Mantel-Cox test. Asterisks indicate p-values * = p<0.05, **= p<0.01 and *** = p<0.001, only significant values are shown. All data depicted in this figure are provided as source data.

Figure 1 Source Data 1 – Raw Data for **Fig. 1B:** Bacterial Load

Figure 1 Source Data 2 – Raw Data for **Fig. 1C:** E. coli Infection

Figure 1 Source Data 3 – Raw Data for **Fig. 1D:** LPS Infection

Figure 1 Source Data 4 – Raw Data for **Fig. 1E:** Histology

Figure 1 Source Data 5 – Raw Data for **Fig. 1F:** ELISA

Figure 2. *NAD⁺ specifically inhibits the non-canonical inflammasome by targeting caspase 11*

Bone marrow was isolated from mice and BMDM were differentiated in vitro. Subsequently, BMDM were cultured in the presence of NAD⁺ or PBS. BMDMs were then primed with either Pam3CSK4 or LPS O111:B4. Next primed BMDMs were stimulated with ATP or LPS and CTB. **(A)** Pro-casp-1, Pro-casp-11, Casp-11, NLRP3, Casp-1, IL1β and GSDMD expression were determined using Western blot and **(B)** IL-1β secretion and LDH release were assessed in the supernatant. **(C)** Time dependent Caspase-1 expression was determined via active staining and assessed using a confocal microscope. **(D)** Cell viability and apoptosis were monitored using the IncuCyte® live microscopy system. **(E)** LPS transfection with CTB was visualized by using FITC-coupled LPS and DAPI staining and quantified by confocal microscopy and flow cytometry. **(F)** For human experiments macrophages were differentiated from PBMC, primed with Pam3CSK4 and subsequently transfected with LPS and 0.25% Fugene HD Plus. Column plots display mean with standard deviation (n=6). Statistical significance was determined by using Student's T-test or One-Way-ANOVA. Asterisks indicate p-values * = p<0.05, **= p<0.01 and ***= p<0.001, only significant values are shown. All data depicted in this figure are provided as source data.

Figure 2 Source Data 1 – Raw Data for **Fig. 2A:** Original Western Blots

Figure 2 Source Data 2 – Raw Data for **Fig. 2A:** Western Blots with highlighted Bands and Sample Labels

Figure 2 Source Data 3 – Raw Data for **Fig. 2B:** ELISA Mouse BMDMs

Figure 2 Source Data 4 – Raw Data for **Fig. 2C:** Caspase 1 Staining

Figure 2 Source Data 5 – Raw Data for **Fig. 2D**: Incucyte Live Microscopy

Figure 2 Source Data 6 – Raw Data for **Fig. 2E:** LPS Transfection Staining

Figure 2 Source Data 7 – Raw Data for **Fig. 2F:** ELISA Human Macrophages

Figure 3. *NAD⁺ mediated inhibition of the non-canonical inflammasome is based on an impaired response to IFN-β*

Differentiated BMDMs were cultured in the presence of NAD⁺ or PBS. BMDMs were then primed with Pam3CSK4, subsequently stimulated with LPS and CTB and RNA sequencing was performed. Unstimulated BMDMs served as controls. **(A)** Venn diagram plotting common gene expression between all 3 groups **(B)** Gene ontology enrichment analysis displaying the highest significant pathways differing when comparing NAD⁺ and PBS treated BMDMs **(C)** Expression cluster analysis of genes involved in IFN-β signaling through cluster analysis depicted in a heat map **(D)** Vulcano plot displaying the most significant genes up or downregulated comparing NAD⁺ and PBS treated BMDMs **(E)** Stimulated BMDMs were additionally treated with recombinant INF-β, and IL-1β and LDH release were measured. **(F)** Moreover, pro-Casp-1, Casp-11, NLRP3, GSDMD, **(G)** STAT-1 and Phospho-STAT-1 expression were assessed by Western blot. Column plots display mean with standard deviation (n=6). Statistical significance was determined by using Student's T-test or One-Way-ANOVA. Asterisks indicate p-values * = p<0.05, **= p<0.01 and ***= p<0.001, only significant values are shown. All data depicted in this figure are provided as source data.

Figure 3 Source Data 1 – Raw Data for **Fig. 3E:** ELISA BMDM
Figure 3 Source Data 2 – Raw Data for **Fig. 3F:** Original Western Blots
Figure 3 Source Data 3 – Raw Data for **Fig. 3F**: Western Blots Highlighted Bands and Sample Labels
Figure 3 Source Data 4 – Raw Data for **Fig. 3G:** Original Western Blots
Figure 3 Source Data 5 – Raw Data for **Fig. 3G**: Western Blots Highlighted Bands and Sample Labels

Figure 4. *Inhibitory effects of NAD$^+$ on IFN-ß downstream signaling and inflammasome activation*

NAD$^+$ inhibits STAT-1 expression and phosphorylation, thus compromising the intracellular response to IFN-β. Subsequently, stimulation of the IFNAR receptor by IFN-β leads to a decreased transcription of pro-caspase-11 as well as ISGs (IFN-inducible GTPases and GBPs). Due to diminished caspase-11 levels, non-canonical inflammasome activation through intracellular, gram-negative bacteria opsonization by GBPs is significantly inhibited.

Figure 5. *IL-10 constitutes an additional mechanism mediating the protective capacities of NAD⁺ in the context of septic shock*

(A) Caspase-11 KO (knockout) mice were treated with NAD⁺ or PBS. Subsequently mice were subjected to Poly(I:C) prior to LPS injection and survival was monitored (n=5, 2 independent survival experiments*)*. Mice treated with either NAD⁺ or PBS were injected with LPS and after 10 hours, Splenic frequencies of IL-10 producing **(B)** Macrophages and Dendritic cells **(C)** and CD4⁺ and CD8⁺ T cells were assessed by flow cytometry. **(D)** BMDMs treated with NAD⁺ or PBS were stimulated with LPS and CTB in the presence of IL-10 neutralizing antibodies and IL-10 receptor antagonists. Subsequently IL-1β and LDH release were assessed. **(E)** IL-10⁻ᐟ⁻ mice treated with NAD⁺ or PBS were challenged with LPS and survival was monitored (n=5-7, 2 independent survival experiments*)*. Column plots display mean with standard deviation (n=6). Statistical significance was determined by using Student's T-test or One-Way-ANOVA while survival data were compared using log-rank Mantel-Cox test. Asterisks indicate p-values * = p<0.05, **= p<0.01 and *** = p<0.001, only significant values are shown. All data depicted in this figure are provided as source data.

Figure 5 Source Data 1 – Raw Data for **Fig. 5A:** Casp11 KO Survival
Figure 5 Source Data 2 – Raw Data for **Fig. 5B:** FACS Macrophages and DCs
Figure 5 Source Data 3 – Raw Data for **Fig. 5C:** FACS CD4+ and CD8+ T cells
Figure 5 Source Data 4 – Raw Data for **Fig. 5D:** ELISA BMDM
Figure 5 Source Data 5 – Raw Data for **Fig. 5E:** IL-10$^{-/-}$ Survival

Figure 1 Supplement 1. NAD$^+$ preserves ileal villi structure and reduces splenic hemorrhage during LPS induced septic shock

C57BL/6 mice were treated with PBS or NAD$^+$ for 2 days prior to administration of a lethal dose of LPS (O55:B5 /O111:B4) by intraperitoneal injection. Ileum and Spleen were removed after 15 hours and subsequently IHC for H&E staining was performed. All data depicted in this figure are provided as source data.

Figure 1 Supplement 1 Source Data 1 – Raw Data for **Fig.1 Supplement 1**: Histology

Figure 1 Supplement 2. **Neutrophils per mm^2 infiltrating mice Ileum, Kidney Lung and Liver in the IHC stains.**

C57BL/6 mice were treated with PBS or NAD$^+$ for 2 days prior to administration of a lethal dose of LPS (O55:B5 /O111:B4) by intraperitoneal injection. After a 15-hour interval, the ileums, lungs, kidneys, and livers were extracted and subjected to immunohistochemical staining with hematoxylin and eosin (H&E). Neutrophil quantification was performed subsequent to the immunohistochemical staining in various tissue samples. Notably, in the Kidney, Lung, and Liver tissues, the number of neutrophils was observed to be significantly higher in the PBS-treated mice group as compared to the NAD$^+$ treated mice group. Column plots display mean with standard deviation (n=4-5). Statistical significance was determined by using Student's T-test or One-Way-ANOVA. Asterisks indicate p-values * = $p<0.05$, **= $p<0.01$. All data depicted in this figure are provided as source data.

Figure 1 Supplement 2 Source Data 1 – Raw Data for **Fig.1 Supplement 2**: Neutrophil Count

Figure 2 Supplement 1. NAD⁺ does not alter BMDM derived NF-κB expression or phosphorylation

Differentiated BMDM were cultured in the presence of 100µmol NAD⁺ or PBS for 2 following days. BMDMs were then primed with 1µg/ml Pam3CSK4 and subsequently stimulated with 2µg/ml LPS O111:B4 and 20µg/ml CTB. Unstimulated BMDMs served as controls. (**A**) P52 and p65 expression was determined using Western blot. (**B**) stimulated BMDMs were stained with p52, p65 and phospho-p65 and expression levels assessed using confocal microscopy. All data depicted in this figure are provided as source data.

Figure 2 Supplement 1 Source Data 1 – Raw Data for **Fig.2 Supplement 1 A**: Western Blot
Figure 2 Supplement 1 Source Data 2 – Raw Data for **Fig.2 Supplement 1A**: Western Blots Bands Highlighted and Sample Labels
Figure 2 Supplement 1 Source Data 3 – Raw Data for **Fig.2 Supplement 1 B**: Immunofluorescence

Figure 2 Supplement 2. Unstimulated BMDM cell viability and apoptosis

Differentiated BMDMs were cultured in the presence of 100µmol NAD⁺ or PBS for 2 following days. BMDMs were then primed with 1µg/ml Pam3CSK4, subsequently stimulated with 2µg/ml LPS O111:B4 and 20µg/ml CTB and cell viability and apoptosis were monitored for 100 hours using the IncuCyte® live microscopy system. All data depicted in this figure are provided as source data.

Figure 2 Supplement 2 Source Data 1 – Raw Data for **Fig.2 Supplement 2**: Incucyte Live Microscopy