The names of the authors

Aru Toyoda[1,2], Kazunari Matsudaira[3], Tamaki Maruhashi[4], Suchinda Malaivijitnond[3,5] and Yoshi

Kawamoto[6]

Title

Highly Versatile, Non-Invasive Method for Collecting Buccal DNA from Free-Ranging Non-Human

Primates

The affiliations and addresses of the authors

[1] Primate Research Institute, Kyoto University, Inuyama, Aichi, Japan

[2] Chubu University Academy of Emerging Sciences, Kasugai, Aichi, Japan

[3] Department of Biology, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

[4] Department of Human Cultures, Musashi University, Tokyo, Japan

[5] National Primate Research Center of Thailand, Chulalongkorn University, Saraburi, Thailand

[6] Graduate School of Veterinary Medicine and Life Science, Nippon Veterinary and Life Science

University, Tokyo, Japan

The e-mail address, and telephone number of the corresponding author

Aru Toyoda : atoyoda.pri.kyoto-u@outlook.com

ABSTRACT

Non-invasive techniques for collection of DNA samples of suitable quality and quantity are important for

improving the efficiency of genetic wildlife research. The development of a non-invasive method for

collection of DNA samples from wild stump-tailed macaques (*Macaca arctoides*) is described herein.

Polyester rope was cut into 10 cm pieces, which were then soaked in a 20% sugar solution to bait

individuals. Rope swabs were immediately collected and transferred to a lysis buffer solution after subjects had picked up, chewed, and discarded them. DNA was later extracted from the buffer. Quantitative real-time PCR and both allelic dropout and genotype failure rates were used to compare the quantity and quality of the buccal DNA samples to those of intestinal slough cell DNA samples collected from freshly dropped feces. The buccal samples yielded significantly more DNA (27.1 ± 33.8 ng/μL) than did the fecal samples (11.4 ± 15.4 ng/μL) and exhibited lower allelic dropout and genotyping failure rates for the 10 autosomal microsatellites investigated. Buccal cell collection was also simple, inexpensive, reliable, and less time-consuming compared to fecal sampling. Thus, this method should facilitate genome-wide studies of non-human primates and other wildlife species.

Keywords

Non-invasive DNA collection, Microsatellite markers, Quantitative real-time PCR, Allelic dropout

Acknowledgments

We are grateful to Warayut Nilpaung, Chuchat Choklap, the superintendent of the Khao Krapuk Khao Taomor Non-Hunting Area. Phanlerd Inprasoet, Wanchai Inprasoet, and Napatchaya Techaatiwatkun for providing a great deal of support in the continued success of this fieldwork; Dr. Yuzuru Hamada, Dr. Hiroki Koda, Dr. Ikuma Adachi, and Dr. Takeshi Nishimura for their supports in our research. This study was funded by the Japan Society for the Promotion of Science KAKENHI (#16J0098 to AT), a Young Science Explorer Grant from the National Geographic Foundation for Science and Exploration - Asia (#Asia-22-15 to AT), Kyoto University Foundation (to AT), and the Cooperation Research Program of the Wildlife Research Center, Kyoto University (#2015-Jiyuu-8 to AT).

Introduction

Wildlife, including non-human primates, has been subject to genetic analyses in a wide variety of research fields, such as evolutionary biology (e.g., Blancher et al. 2008, Osada et al. 2015, Tosi et al. 2002), population genetics (e.g., Evans et al. 2001, Kanthaswamy et al. 2013, Trask et al. 2013), phylogeography (e.g., Bunlungsup et al. 2016), pedigree analysis (e.g., Barelli et al. 2013), and conservation biology (e.g., Minhós et al. 2013), using a variety of DNA markers. Mitochondrial DNA (mtDNA), for example, is generally used for investigating maternal relationships and phylogeography (Avise 2004), whereas Y-chromosome genes of mammals are used to investigate paternal relationships and male dispersal (Tosi et al. 2000, Tosi et al. 2002). Meanwhile, autosomal markers, such as microsatellite and single-nucleotide polymorphism (SNP) markers, are often used to investigate population genetics and genomic diversity (Bruford and Wayne 1993, Brumfield et al. 2003).

As a result of recent advances in DNA analysis technology and growing concerns over animal welfare (Russell and Burch 1959), genetic studies of wildlife frequently use DNA samples that have been collected by non-invasive means (e.g. Woodruff 1993, Taberlet et al. 1999). For example, DNA samples have been collected from egg shells (herring gull, *Larus argentatus*; Egloff et al. 2009), blood-fed mosquitos

(Ejiri et al. 2011), koala feces (*Phascolarctos cinereus*; Wedrowicz et al. 2013), and

bug-bite blood (Sumatran rhinoceros, *Dicerorhinus sumatrensis*; Rovie-Ryan et al.

2013). DNA has similarly been collected non-invasively in genetic studies of wild,

non-human primates, for example, from trapped hairs (white-headed langur,

Trachypithecus leucocephalus; Wang et al. 2016), semen (Japanese macaques, *Macaca*

fuscata; Domingo-Roura et al. 2004), urine (Japanese macaques; Hayakawa and

Takenaka 1999), and saliva (mountain gorillas, *Gorilla beringei beringei*, and Grauer's

gorillas, *Gorilla beringei graueri*; Smiley et al. 2010, Chimpanzee, *Pan troglodytes*,

Inoue et al. 2007). Among these DNA resources, fecal samples have been most

commonly used (Blair and Melnick 2011, Buchan et al. 2003, Nietlisbach et al. 2012,

Städele et al. 2015, Vigilant et al. 2001). However, fecal samples generally yield low

quantities of low-quality DNA, and even though the markers used in some studies (e.g.,

mtDNA markers) can be amplified successfully owing to their high copy numbers

(Bunlungsup et al. 2016), enormous efforts are required when examining nuclear markers

(Navidi et al. 1992, Taberlet et al. 1996). One major problem with using fecal DNA

samples for nuclear genotyping is allelic dropout, a phenomenon in which one of two

autosomal alleles is not amplified by PCR, causing heterozygous genotypes to be

misinterpreted as homozygous (Pompanon et al. 2005, Tebbutt and Ruan 2008). Allelic

dropout is problematic in paternity and kinship analyses using autosomal microsatellites

(Vigilant et al. 2001).

As such, development of non-invasive DNA sampling methods that allow

researchers to obtain large quantities of high-quality DNA samples with low levels of

contamination is needed. Buccal cell collection methods, such as collecting sugarcane

wedges or pith of terrestrial herbaceous vegetation after their chewing by wild bonobos

(*Pan paniscus*, Hashimoto et al. 1996, Ishizuka et al. 2018), taking oral swabs from

anesthetized mountain and Grauer's gorillas (Smiley et al. 2010), and attaching ropes to

saliva-collecting devices near free-ranging Tibetan macaques (*Macaca thibetana*,

Simons et al. 2012), have been reported previously. Collecting DNA from wedges of

sugar cane or other plants is a non-invasive method that does not require manipulation

of animals and is thus applicable to other study sites with appropriate modification

according to certain factors, such as the environment of the study and the feeding

patterns of the subjects. However, methods that require anesthesia are impossible in

most of field studies, and methods that require specialized equipment takes time and

cost to produce the device. Especially, in the wild condition, using specific devices is

less flexible to collect multiple samples from several monkeys at once due to mobilities.

Such methods were inapplicable to the stump-tailed macaques at our study site because

of the difficulty in preparation and storage of the bite materials. Thus, we designed

another method for collecting buccal cells reporting here.

Herein, a non-invasive method for collecting buccal DNA samples using rope

swabs is described as simple, reliable, inexpensive, and less time-consuming than other

commonly used methods. To test the effectiveness of this method, two experiments

were conducted. The first was a quantitative comparative test of host DNA in 41 fecal

and 41 buccal DNA samples randomly selected using real-time PCR. In addition, gel

electrophoresis ("gel tests") were also used to quantitatively test DNA samples cheaply

and conveniently, and their results were compared with those of costlier real-time PCR

to verify their accuracy. The second experiment was a qualitative comparison based on

allelic dropout and genotype failure rates in 30 fecal and 30 buccal DNA samples

selected using gel tests.

Materials and methods

Study site

The present study was conducted at the Khao Krapuk Khao Taomor Non-Hunting Area,

Phetchaburi Province, Thailand (12°47'59.2″ N, 99°44'31.1″ E), which harbors five

free-ranging groups of stump-tailed macaques (*Macaca arctoides*). There are five

groups: Ting group, 115 individuals; Nadam group, 91 individuals; Third group, 71

individuals; Fourth group, 75 individuals; Wngklm group, 43 individuals (Toyoda and

Malaivijitnond 2018). This survey area is mainly a mountainous area composed of

secondary forests and bamboo forests, and open areas coexist including temple and

houses of local people. The moving area of monkeys was divided between north and

south by large roads, and food provisioning by locals or visitors was occasionally

observed along the road or at temple ground. As information on environmental

conditions, mean annual temperature and annual rainfall are 27°C and 1070 mm,

respectively, in the data of the nearby national park, named Keang Krachan National

Park, about 30km far from this study site (Wijitkosum 2012). This site consists

primarily of secondary forest, including stands of bamboo and agricultural areas. Food

provisioning by locals or visitors was occasionally observed.

Collection and extraction of DNA samples

Buccal cells were collected using baited ropes (hereafter *rope swabs)*. Polyester ropes (6

mm in diameter; Takagi Corporation, Kagawa, Japan, JAN code: 4943 956 261 513)

were cut into approximately 10 cm pieces, autoclaved, and dried (Figure 1). To bait

individuals, the rope swabs were soaked in a 20% sugar solution (70 g cane sugar

dissolved in 350 mL distilled water) for at least 30 min, and then scattered on the open

ground where the monkeys were found. After being chewed (Figure 2) and discarded by

a monkey, the rope swab was quickly collected and transferred to a 5 mL carrying tube

containing 3 mL lysis buffer (0.5 % (w/v) in SDS, 100 mM EDTA pH 8.0, 100 mM

Tris-HCl pH 8.0, and 10 mM NaCl) (Hayaishi and Kawamoto 2006). To compare the

quantity and quality of the buccal DNA with that of other commonly used DNA sources,

intestinal slough cells from freshly dropped fecal samples were also collected. A sterile

cotton bud, which was soaked in 2 mL lysis buffer, was used to swab the surfaces of

feces, following the protocol of Bunlungsup et al. (2016). To increase DNA yields, the

surfaces of the feces were swabbed at least three times.

149   The buccal and intestinal cells that were transferred to the lysis buffer were

kept at room temperature for at least five months until DNA extraction. DNA was

extracted following the procedure of Kawamoto et al. (2013). Potential PCR inhibitors

were removed by adding 600 mg of hydrolyzed starch (Wako, Osaka, Japan) to 1.5 mL

of lysis buffer per sample. The samples were incubated at 36 °C for 10 min, and then

centrifuged at 1000 $\times g$ for 15 min. Finally, 750 μL of each supernatant was processed

using a commercially available DNA clean-up system (Wizard SV Gel and PCR

Clean-Up System; Promega, Madison, WI, USA), and the DNA was finally eluted with

50 µL pure water. Study had been conducted from September 25[th], 2015 to June 15[th],

2017, and 74 fecal samples and 579 buccal samples were collected.

DNA quantification

The amount of host DNA was quantified by quantitative real-time PCR (Morin

et al. 2001). Forty-one DNA samples extracted from buccal and 41 from fecal samples

were selected randomly from all of the extracted DNA samples. The real-time PCR

method was used because both the buccal and intestinal DNA samples were

contaminated with other exotic DNA sources, such as bacteria, eukaryotic parasites, and

dietary materials (e.g., plants, insects, or small animals), which could not be

differentiated using conventional spectrophotometry. The sequences of the real-time

PCR primers and c-*myc* probe were 5'-GCCAGAGGAGGAACGAGCT-3'

(CMYC_E3_F1U1), 5'-GGGCCTTTTCATTGTTTTCCA-3' (CMYC_E3_R1U1), and

5'-FAM-TGCCCTGCGTGACCAGATCC-TAMRA-3'   (CMYC_E3_TMV),

respectively (Morin et al. 2001). Real-time PCR was performed using a StepOnePlus

real-time PCR System (Thermo Fisher Scientific, Waltham, MA, USA), and each 20 µL

reaction contained 2 µL DNA template, 1× TaqMan Fast Advanced Master Mix

(Thermo Fisher Scientific), 250 nM probe, and 900 nM of each primer. In addition, the

PCR amplification conditions included an initial denaturation step of 95 °C for 20 s,

followed by 45 cycles of 95 °C for 1 s and 60 °C for 20 s. Host DNA quantity

(concentration) was determined using a standard curve made by a duplicate set of DNA

with known quantity. The standard set was made from DNA extracted from the blood of

a northern pig-tailed macaque (*Macaca leonina*) reared in the Primate Research Unit,

Chulalongkorn University (Bangkok, Thailand). The DNA was quantified using a

spectrophotometer and diluted to 10 ng/μL, 2.5 ng/μL, 625 pg/μL, 156 pg/μL, 39.1

pg/μL, and 9.8 pg/μL with deionized water. The mean DNA yields obtained from the

buccal and fecal samples were compared using the Wilcoxon rank sum test in R Ver.

3.4.2 (R Core Team 2016).

Real-time PCR provides an accurate host DNA concentration for each DNA

sample, and thus was appropriate for comparing the DNA yields of the buccal and fecal

samples. However, real-time PCR analysis is expensive. Therefore, to select suitable

samples for microsatellite genotyping, the usability of the 82 DNA samples was roughly

screened using conventional PCR and agarose gel electrophoresis following the

procedure of Kawamoto et al. (2013) and Ball et al. (2007) (gel electrophoresis). For the

gel test, the c-*myc* gene was PCR-amplified in 12.5 μL reactions of 1 μL template DNA,

1× PCR Buffer for KOD FX, 400 μM dNTPs, 0.25 U KOD FX (Toyobo, Osaka, Japan),

and 0.015 pM of both the forward and reverse real-time PCR primers, using the

following conditions: initial denaturation step of 94 °C for 2 min, 45 cycles of 98 °C for

10 s, 58 °C for 30 s, and 68 °C for 30 s. The resulting amplicons were electrophoresed

on 2% agarose-TAE gels, stained with SYBR Safe DNA Gel Stain (Thermo Fisher

Scientific), and visualized using UV transilluminators to determine the intensity of the

target band. To estimate the amount of buccal and intestinal DNA, a series of human

placental DNA (Sigma-Aldrich, St. Louis, MO, USA) at concentrations of 500, 300, and

100 pg/µL were used as reference controls. When the luminous intensity of a PCR

product was > 300 pg/µL of the control, the sample was considered to have sufficient

yield and was used in the next step for microsatellite amplification. We used human

placental DNA as a reference as following Kawamoto et al. (2013), that was different

from the *Macaca leonina*'s DNA used in the real-time PCR. This was because of the

difference of availability of the DNA standard in Japan and Thailand, and the difference

of the species was considered not to affect the substantial results (Smith et al. 2002).

The accuracy of the real-time PCR analysis and gel test screening were compared using

the Wilcoxon rank sum test with continuity correction.

DNA quality analysis

To determine DNA quality, the 30 paired buccal and intestinal DNA samples that passed

the gel test were randomly selected for microsatellite genotyping. Ten microsatellite loci

(D3S1768, D6S2793, D7S2004, D8S1106, D11S2002, D13S765, D14S306, D17S1290,

D18S537, and D19S582; Aarnink et al. 2011, Barelli et al. 2013, Kawamoto et al. 2008,

Nair et al. 2000, Smith et al. 2000) were amplified using a modified version of the

two-step multiplex method described by Arandjelovic et al. (2009). During the first step

of PCR, all microsatellite loci were amplified in a single 20 μL reaction that included 1

μL template DNA. During the second step, the 10 loci were divided into three subsets

and were amplified in 12.5 μL multiplex PCR reactions that each included 1 μL of

non-diluted amplicon from the first multiplex PCR reaction. The PCR thermocycling

conditions were the same as those from the gel test, except that 35 cycles were used for

the first PCR, and 45 for the second PCR. Allelic dropout rates and false allele rates

were calculated using PEDANT Ver.1 (Johnson and Haydon 2007, available from

http://sites.google.com/site/pcdjohnson/home/pedant). In the program, the results of two

independent PCR products per sample per locus were used to estimate the allelic

dropout and false allele rates. The allelic dropout and false allele rates of the buccal and

fecal sample DNA were compared using the Wilcoxon signed-rank test ($p < 0.05$) in R.

In addition, the genotype failure rate (a phenomenon in which the peak of an allele is

detected at extremely low levels or is not detected) of each locus was calculated based

on the duplicated PCR results, and the genotype failure rates of the buccal and fecal

DNA samples were compared using the Wilcoxon signed-rank test (p < 0.05) in R.

Results

DNA quantity

Analysis of the 82 DNA samples (41 buccal and 41 intestinal DNA samples) revealed

that the buccal samples yielded significantly more host DNA (27.1 ± 33.8 ng/µL) than

did the fecal samples (11.4 ± 15.4 ng/µL; $W = 473$, $P < 0.001$). Although 68% (28/41)

of intestinal samples yielded concentrations less than 10 ng/µL, only 29% of buccal

samples produced such low concentrations (12/41) (Figure 3).

The determination by the gel test was possibly made the presence/absence of

the band (Figure 5, as an example of gel-test judgment). Of the 41 fecal and buccal

DNA samples tested, 22 (53.7%) and 35 (85.4%) met the criterion for sufficient yield ($\geq$

300 pg/µl), respectively. The concentration of host DNA that passed and failed the gel

tests as measured by real-time PCR was a significant difference (W = 991, p < 0.01),

indicating that either real-time PCR or the gel test can be used for DNA quantification.

DNA quality

For the 30 monkeys whose samples passed the gel test, the allelic dropout rate of the 10

microsatellite loci was significantly lower for the buccal (0.00%, range: $0.00 - 6 \times$

10^{-6} %) than for the fecal DNA samples (18.12 $\pm$ 16.12%, range: 0.00–55.96%;

Wilcoxon signed-rank test, V = 44, p < 0.01; Table 1). Estimated dropout rates were

used to calculate the amount of repetition necessary for accurate results at the 99.99%

certainty level (Morin et al. 2001). At least 6 repetitions were needed for fecal sample

analysis to produce reliable genotype data, whereas one repetition was sufficient for

buccal samples.

Similarly, the genotyping failure rate was significantly lower for buccal DNA

samples (2.70% $\pm$ 3.88, range: 0.0–13.3 %) than for fecal DNA samples (35.67% $\pm$

15.35, range: 18.3–65.0%; Wilcoxon signed-rank test, V = 55, p < 0.01), although the

rate was variable among the loci examined (Table 1).

Discussion

Advantages from sampling point of view

When fecal sample are used as genetic resources, the success depends largely on the

skill of the collector, the state of the feces, and other factors. Using the rope swab

method in our study, the collection of high-quantity and quality DNA samples would be

possible without much training, providing a more versatile option that is not dependent

heavily on the level of experience of the sample collector. Our rope swab method may

also be useful for collecting samples from infants. Indeed, our method was capable of

collecting samples from infants aged 2–3 weeks, even though the feces of infants were

often soft, diarrhea-like or very small and often difficult or almost impossible to collect.

Thus, we strongly believe that our method would be a powerful alternative to overcome

the difficulty of collecting fecal samples from infants which are indispensable for

genetic analysis such as paternity test. The rope swab method is also less time

consuming than fecal collection. Since the quality of DNA samples cannot be checked

in-situ study, multiple fecal samples must be collected to ensure collection of an

adequate sample from the target animals. On the other hand, most of buccal samples

provided usable DNA, and thus, fewer specimens need to be collected from each animal.

Additionally, to collect fecal samples, researchers must patiently follow the targeted

animals until they defecate, which is time-consuming. Therefore, the rope swab method

presented in this study has great potential to save time and mitigate these factors.

Advantages from analysis point of view

Our study showed that the rope swab method is more effective, in terms of both

quantity and quality of recovered DNA, compared to extraction from fecal samples. The

rope swab method yielded up to 2.4 times more host DNA than did fecal samples and

exhibited much lower allelic dropout and genotype failure rates, indicating that our

method possibly facilitates genotyping analyses with fewer repetitive PCR trials, which

could save time, labor, and money. This is because low DNA quantity increases

genotyping errors that affect the reliability of genotyping in microsatellite analysis

(Taberlet et al. 1999), and thus repeating experiments for each locus and extract is

recommended (Goossens et al. 1998).

Important notice using rope swab method

Although our method would be useful, there are several cautionary notes while

collecting samples. Firstly, in the initial phase, monkeys may not chew on the rope

swabs. In this case, a habituation period using fruit juice instead of sugar water to

increase the attractiveness of the swab rope is recommended. From experience, however,

it seems better to switch to sugar water during the sample-collection phase. Genotyping

results were not stable when using DNA samples collected with orange juice, probably

due to the acid or other chemical compounds present in the fruit juice.

Secondly, the collection of samples shortly after monkeys have consumed food

should be avoided, especially at provisioned sites or when targeting captive animals, as

fruits are the main food items given and contain acids or other chemical compounds that

may inhibit PCR. Complex polysaccharides possibly originating from vegetable

material in the diet are also considered potential PCR inhibitors (Monteiro et al. 1997).

Thus, time of sampling may affect the quality of the sample rather than the duration for

which the monkey chews the rope.

Thirdly, adjustments to the soaking time of the rope swab in the sugar-water

solution and the concentration of sugar according to the condition of the subject animals

or study site may be needed. Extended soaking times or high sugar concentrations could

encourage monkeys to chew the rope swabs for longer periods, which may lead to

greater DNA yields. However, the potential downside of a longer chewing period is that

the target monkeys may move while chewing, making retrieval of the rope swabs more

difficult for the researcher. Although some individuals spent significant time chewing

the swabs and occasionally broke them into small fragments, no monkeys accidentally

ate the rope swabs during this study period, demonstrating their safety in application.

Fourth, the rope swabs should be well-distributed among the troop, otherwise

higher-ranking males will take multiple ropes at once. When samples from subordinate

individuals are needed, spreading the rope swabs over a wide area to attract

high-ranking individuals, and then casting some swabs to the target individual may be

an effective strategy.

Lastly, because this method requires that the rope swabs be provided to the

animals, it may not be suitable for use with non-habituated, wild animals. This method

also cannot be used in research sites where access to wildlife or provisioning is

prohibited. Since this method involves material once contained in the mouths of animals,

researchers must be aware of the possibility of touching saliva to prevent zoonosis (e.g.,

Kelesidis and Tsiodras 2010). When conducting behavioral observation at the same time,

the possibility of influencing the behavior of the target animals must also be considered.

Ultimately, the applicability of this method will depend on the specific needs and

conditions of the research.

Furthermore, we must note about the standard range of quantitative real-time

PCR. In this study, the standard range of quantitative real-time PCR could not cover the

sample concentration range due to the fact that the quantity of DNA was extracted at a

higher concentration than our assumption. We followed the protocol of Wizard SV Gel

and PCR Clean-Up System and used 50 µL of water for the final elution step, though

200 µL is used in Morin et al. (2001). This difference of the final elution volume should

have resulted in the higher concentration of DNA both from buccal and fecal samples in

our study.

Future possibility of application

The successful DNA collection and genotyping of *M. arctoides* using our method can be

further applied to different conditions, for example, for populations kept in captive

conditions at research institutions or individuals kept in cages in laboratories but also

for provisioned or well-habituated free-ranging primates such as populations living near

temples which are widely seen in most Southeast Asian countries, as long as researchers

pay attention to risks and take precautions. This is a very useful method for researchers

who have to obtain samples from specific individuals in a limited research period in the

wild. Furthermore, with some modifications, this method can be applied for hormone

and veterinary analysis (e.g., detecting a specific virus in the saliva; Musso et al. 2015,

Huff et al. 2003). The non-invasive buccal cell collection method described by this

study may further facilitate animal population genomic studies in both captive and field

environments. Further integration of genetic information with behavioral and ecological

data is expected to provide more insights into *M. arctoides*, including genetic structure

and socioecological characteristics such as reproductive strategy and kinship structure.

References

Aarnink A, Dereuddre-Bosquet N, Vaslin B, Le Grand R, Winterton P, Apoil PA,

Blancher A (2011) Influence of the MHC genotype on the progression of

experimental SIV infection in the Mauritian cynomolgus macaque. Immunogenetics

63:267-274

Arandjelovic M, Guschanski K, Schubert G, Harris TR Thalmann O, Siedel H, Vigilant

L (2009) Two-step multiplex polymerase chain reaction improves the speed and

accuracy of genotyping using DNA from noninvasive and museum samples.

Molecular Ecology Resources 9:28-36

Avise JC (2004) Molecular markers, natural history, and evolution, Second Edition.

Sinauer Associates, Inc., MA, USA.

Ball MC, Pither R, Manseau M, Clark J, Petersen SD, Kingston S, Morrill N, and

Wilson P (2007) Characterization of target nuclear DNA from faeces reduces

technical issues associated with the assumptions of low-quality and quantity

template. Conservation Genetics 8:577-586

Barelli C, Matsudaira K, Wolf T, Roos C, Heistermann M, Hodges K, Ishida T,

Malaivijitnond S, Reichard UH (2013) Extra-pair paternity confirmed in wild

white-handed gibbons. American Journal of Primatology 75:1185-1195

Blair ME, Melnick DJ (2011) Genetic evidence for dispersal by both sexes in the central

American squirrel monkey, *Saimiri oerstedii citrinellus*. American Journal of

Primatology 74:37-47

Blancher A, Bonhomme M, Crouau-Roy B, Terao K, Kitano T, Saitou N (2008)

Mitochondrial DNA sequence phylogeny of 4 populations of the widely distributed

cynomolgus macaque (*Macaca fascicularis fascicularis*). Journal of Heredity

99:254-264

Bruford MW, Wayne RK (1993) Microsatellites and their application to population

genetic studies. Current Opinion in Genetics and Development 3:939-943

Brumfield RT, Beerli P, Nickerson DA, Edwards SV (2003) The utility of single

nucleotide polymorphisms in inferences of population history. Trends in

Ecology and Evolution 18:249-256

Buchan JC, Alberts SC, Silk JB, Altmann J (2003) True paternal care in a multi-male

primate society. Nature 425:179-181

Bunlungsup S, Imai H, Hamada Y, Gumert MD, San AM, Malaivijitnond S (2016)

Morphological characteristics and genetic diversity of Burmese longtailed

macaques (*Macaca fascicularis aurea*). American Journal of Primatology

74:441-455

Domingo-Roura X, Marmi J, Andrés O, Yamagiwa J, Terradas J (2004) Genotyping

from semen of wild Japanese macaques (*Macaca fuscata*). American Journal of

Primatology 62:31-42

Egloff C, Labrosse A, Hebert C, Crump D (2009) A nondestructive method for obtaining

maternal DNA from avian eggshells and its application to embryonic viability

determination in herring gulls (*Larus argentatus*). Molecular Ecology Resources

9:19-27

Ejiri H, Sato Y, Kim KS, Hara T, Tsuda Y, Imura T, Murata K, Yukawa M (2011)

Entomological study on transmission of avian malaria parasites in a zoological garden

in Japan: bloodmeal identification and detection of avian malaria parasite DNA from

blood-fed mosquitoes. Journal of Medical Entomology. 48:600-607

Evans BJ, Supriatna J, Melnick DJ (2001) Hybridization and population genetics of two

macaque species in Sulawesi, Indonesia. Evolution 55:1686-1702

Goossens B, Waits LP, Taberlet P (1998) Plucked hair samples as a source of DNA:

reliability of dinucleotide microsatellite genotyping. Molecular Ecology

7:1237-1241

Hashimoto C, Furuichi T, Takenaka O (1996) Matrilineal kin relationship and social

behavior of wild bonobos (*Pan paniscus*): sequencing the D-loop region of

mitochondrial DNA. Primates 37:305-318

Hayakawa S, Takenaka O (1999) Urine as another potential source for template DNA in

polymerase chain reaction (PCR). American Journal of Primatology 48:299-304

Hayaishi H, Kawamoto Y (2006) Low genetic diversity and biased distribution of

mitochondrial DNA haplotypes in the Japanese macaque (*Macaca fuscata yakui*) on

Yakushima Island. Primates 47:158-164

Huff JL, Eberle R, Capitanio J, Zhou SS, Barry PS (2003) Differential detection of B

virus and rhesus cytomegalovirus in rhesus macaques. Journal of General Virology

84:83-92

Inoue E, Inoue-Murayama M, Takenaka O, Nishida T (2007) Wild chimpanzee infant

urine and saliva sampled noninvasively usable for DNA analyses. Primates

48:156-159

Ishizuka S, Kawamoto Y, Toda K, Furuichi T (2018) Bonobos' saliva remaining on the

pith of terrestrial herbaceous vegetation can serve as non-invasive wild genetic

resources. Primates 60:7-13

Johnson PCD, Haydon DT (2007) Maximum-likelihood estimation of allelic dropout

and false allele error rates from microsatellite genotypes in the absence of reference

data. Genetics 175:827-842

Kanthaswarry S, Ng J, Trask JS, George DA, Kou AJ, Hoffman LN, Doherty TB,

Houghton P, Smith DG (2013) The genetic composition of populations of

cynomolgus macaques (*Macaca fascicularis*) used in biomedical research. Journal

of Medical Primatology 42:120-131

Kawamoto Y, Kawamoto S, Matsubayashi K, Nozawa K, Watanabe T, Stanley MA,

Perwitasari-Farajallah D (2008) Genetic diversity of longtail macaques (*Macaca

fascicularis*) on the island of Mauritius: an assessment of nuclear and mitochondrial

DNA polymorphisms. Journal of Medical Primatology 37:45-54

Kawamoto Y, Takemoto H, Higuchi S, Sakamaki Tart JA, Hart TB, Tokuyama N,

Reinartz GE, Guislain P, Dupain J, Cobden AK, Mulavwa MN, Yangozene K,

Darroze S, Devos C, Furuichi T (2013) Genetic structure of wild bonobo

populations: diversity of mitochondrial DNA and geographical distribution. PLoS

ONE 8:e59660

Kelesidis T, Tsiodras S (2010) *Staphylococcus intermedius* is not only a zoonotic

pathogen, but may also cause skin abscesses in humans after exposure to saliva.

International Journal of Infectious Diseases 14:838-841

Minhós T, Wallace E, Ferreira da Silva MJ, Sa RM, Carmo M, Barata A, Bruford MW

(2013) DNA identification of primate bushmeat from urban markets in Guinea-Bissau

and its implications for conservation. Biological Conservation 167:43-49

Monteiro L, Bonnemaison D, Vekris A, Petry KG, Bonnet J, Vidal R, Cabrita J,

Mégraud F (1997) Complex polysaccharides as PCR inhibitors in feces *Helicobacter*

pylori model. Journal of Clinical Microbiology 35:995-998

Morin PA, Chambers KE, Boesch C, Vigilant L (2001) Quantitative polymerase chain

reaction analysis of DNA from noninvasive samples for accurate microsatellite

genotyping of wild chimpanzees (*Pan troglodytes verus*). Molecular Ecology

10:1835-1844

Musso D, Roche C, Nhan TX, Robin E, Teissier A, Cao-Lormeau VM (2015) Detection

of Zika virus in saliva. Journal of Clinical Virology 68:53-55

Nair S, Ha J, Rogers J (2000) Nineteen new microsatellite DNA polymorphisms in

pigtailed macaques (*Macaca nemestrina*). Primates 41:343-350

Navidi W, Arnheim N, Waterman MS (1992) A multiple-tubes approach for accurate

genotyping of very small DNA samples by using PCR: statistical considerations.

The American Journal of Human Genetics 50:347-359

Nietlisbach P, Arora N, Nater A, Goossens B, Van Schaik CP, Krützen M (2012)

Heavily male-biased long-distance dispersal of orang-utans (genus: *Pongo*), as

revealed by Y-chromosomal and mitochondrial genetic markers. Molecular Ecology

21:3173-3186

Osada N, Hettiarachchi N, Babarinde IA, Saitou N, Blancher A (2015) Whole-genome

sequencing of six Mauritian cynomolgus macaques (*Macaca fascicularis*) reveals a

genome-wide pattern of polymorphisms under extreme population bottleneck.

Genome Biology and Evolution 7:821-830

Pidancier N, Miquel C, Miaud C (2003) Buccal swabs as a non-destructive tissue

sampling method for DNA analysis in amphibians. Herpetological Journal

13:175-178

Pompanon F, Bonin A, Bellemain E, Taberlet P (2005) Genotyping errors: causes,

consequences and solutions. Nature Reviews Genetics 6:847-846

R Core Team (2016) R: a language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. URL

htpps://ww,R-project.org/.

Rovie-Ryan JJ, Zainuddin ZZ, Marni W, Ahmad AH, Ambu LN, Payne J (2013) Blood

meal analysis of tabanid fly after it biting the rare Sumatran rhinoceros. Asian

Pacific Journal of Tropical Biomedicine 3:95-99

Russell WMS, Burch RL (1959) The Principles of Humane Experimental Technique.

Methuen, London. ((http://altweb.jhsph.edu/pubs/books/humane_exp/het-toc)

Accessed December 18, 2017)

Simons ND, Lorenz JG, Sheeran LK, Li JH, Xia DP, Wargner RS (2012) Noninvasive

saliva collection for DNA analyses from free-ranging Tibetan macaques (*Macaca*

thibetana). American Journal of Primatology 74:1064-1070

Smiley T, Spelman L, Lukasik-Braum M, Mukherjee J, Kaufman G, Akiyoshi DE,

Cranfield M (2010) Noninvasive saliva collection techniques for free-ranging

mountain gorillas and captive eastern gorillas. Journal of Zoo and Wildlife Medicine

1:201-209

Smith DG, Kanthaswamy S, Viray J, Cody L (2000) Additional highly polymorphic

microsatellite (STR) loci for estimating kinship in rhesus macaques (*Macaca*

mulatta). American Journal of Primatology 50:1-7

Smith S, Vigilant L, Morin PA (2002) The effect of sequence length and oligonucleotide

mismatches on 5' exonuclease assay efficiency. Nucleic Acids Research 30:e111

Städele V, van Doren V, Pines M, Swedell L, Vigilant L (2015) Fine-scale genetic

assessment of sex-specific dispersal patterns in a multilevel primate society. Journal

of Human Evolution 78:103-113

Taberlet P, Griffin S, Goossens B, Questiau S, Manceau V, Escaravage N, Waits LP,

Bouvet J (1996) Reliable genotyping of samples with very low DNA quantities

using PCR. Nucleic Acids Research 24:3189-3194

Taberlet P, Waits L.P., Luikart G (1999) Noninvasive genetic sampling: look before you

leap. Trends in Ecology and Evolution 14:323-327.

Tebbutt S, Ruan J (2008) Combining multiple PCR primer pairs for each amplicon can

improve SNP genotyping accuracy by reducing allelic drop-out. Biotechniques

45:637-646

Tosi AJ, Morales JC, Melnick DJ (2000) Comparison of Y-chromosome and mtDNA

phylogenies leads to unique inferences of macaque evolutionary history. Molecular

Phylogenetics and Evolution 17:133-144

Tosi AJ, Morales JC, Melnick DJ (2002) Y-chromosome and mitochondrial markers in

Macaca fascicularis indicate introgression with Indochinese *M. mulatta* and a

biogeographic barrier in the Isthmus of Kra. International Journal of Primatology

23:161-178

Trask JS, George D, Houghton P, Kanthaswarry S, Smith DG (2013) Population and

landscape genetics of an introduced species (*M. fascicularis*) on the island of

Mauritius. PLoS One 8:e53001-10.1371/journal.pone.0053001

Vigilant L, Hofreiter M, Siedel H, Boesch C (2001) Paternity and relatedness in wild

chimpanzee communities. Proceedings of the National Academy of Sciences of the

United States of America 98:12890-12895

Wang W, Qiao Y, Zheng Y, Yao M (2016) Isolation of microsatellite loci and reliable

genotyping using noninvasive samples of a critically endangered primate,

Trachypithecus leucocephalus. Integrative Zoology 11:250-262

Wedrowicz F, Karsa M, Mosse J, Hogan FE (2013) Reliable genotyping of the koala

(*Phascolarctos cinereus*) using DNA isolated from a single faecal pellet. Molecular

Ecology Resources 13:634-641

Woodruff DS (1993) Non-invasive genotyping of primates. Primates 33:333-346

Table 1. Allelic dropout and genotype failure rates of 10 microsatellite loci for fecal and
buccal DNA samples of stump-tailed macaques in Khao Krapuk Khao Taomor.

Loci	Allelic dropout rate (%)		Genotype failure rate (%)	
	Fecal	Buccal	Fecal	Buccal
D3S1768	10.54	0.00	38.33	0.00
D6S2793	25.00	0.00	58.33	2.00
D7S2204	8.80	0.00	31.67	13.33
D8S1106	13.04	0.00	45.00	0.00
D11S2002	0.00	0.00	65.00	1.67
D13S765	29.20	0.00	23.33	0.00
D14S306	0.00	0.00	18.33	1.67
D17S1290	28.45	0.00	33.33	5.00
D18S537	55.97	0.00	25.00	3.33
D19S582	10.21	0.00	18.33	0.00

530   Figure

532   **Fig 1** Rope swabs cut into 10cm length and 3ml of lysis buffer in 5ml tube

535　**Fig 2** Monkey chewing a rope swab

Fig 3 Frequency of buccal and fecal DNA in each DNA concentration zone. Although

many fecal samples are dense in the low concentration zone, meaning that the sampling

efficiency is not good, buccal samples shows a gentle peak overall, indicating that

samples with high concentration can be more easily obtained.