Title Page

Title: HIFα regulates developmental myelination independent of autocrine Wnt signaling

Running Title: Mechanisms of HIFα-regulated myelination

Author name: Sheng Zhang[1,3], Yan Wang[1,3], Jie Xu[3], Wenbin Deng[2,3], and Fuzheng Guo[1,3]

Author affiliation:

1 Department of Neurology, 2 Department of Neurology Biochemistry and Molecular Medicine,

School of Medicine, University of California, Davis; 3 Institute for Pediatric Regenerative

Medicine (IPRM), Shriners Hospitals for Children, Northern California.

Corresponding author: Fuzheng Guo, PhD, 2425 Stockton Blvd, Sacramento, CA 95817.

Email: fzguo@ucdavis.edu   ORCID: 0000-0003-3410-8389

Competing interests: The authors declare no competing financial interests

Acknowledgements: This study was supported by the grants funded by NIH/NINDS

(R21NS109790 and R01NS094559 to F.G.) and Shriners Hospitals for Children (86100, 85200-

NCA16, 85107-NCA-19 to F.G., 84307-NCAL to S.Z.).

Abstract

The developing CNS is exposed to physiological hypoxia, under which hypoxia inducible

factor alpha (HIFα) is stabilized and plays a crucial role in regulating neural development. The

cellular and molecular mechanisms of HIFα in developmental myelination remain incompletely

understood. Previous concept proposes that HIFα regulates CNS developmental myelination by

activating the autocrine Wnt/β-catenin signaling in oligodendrocyte progenitor cells (OPCs).

Here, by analyzing a battery of genetic mice of both sexes, we presented *in vivo* evidences

supporting an alternative understanding of oligodendroglial HIFα-regulated developmental

myelination. At the cellular level, we found that HIFα was required for developmental

myelination by transiently controlling upstream OPC differentiation but not downstream

oligodendrocyte maturation and that HIFα dysregulation in OPCs but not oligodendrocytes

disturbed normal developmental myelination. We demonstrated that HIFα played a minor, if any,

role in regulating canonical Wnt signaling in the oligodendroglial lineage or in the CNS. At the

molecular level, blocking autocrine Wnt signaling did not affect HIFα-regulated OPC

differentiation and myelination. We further identified HIFα-Sox9 regulatory axis as an underlying

molecular mechanism in HIFα-regulated OPC differentiation. Our findings support a concept

shift in our mechanistic understanding of HIFα-regulated CNS myelination from the previous

Wnt-dependent view to a Wnt-independent one and unveil a previously unappreciated HIFα-

Sox9 pathway in regulating OPC differentiation.

Introduction

HIFα is a master transcriptional regulator of the adaptive response to hypoxia (Semenza,

2012). The oxygen concentration in the developing CNS was reported ranging from 0.5 to 7%

(Ivanovic, 2009; Zhang et al., 2011). HIFα protein (HIF1α and HIF2α) is constitutively translated

but subjected to rapid turnover (half-life<5 min) by the proteasome-mediated degradation in

which the von Hippel-Lindau (VHL) protein plays an essential role (Semenza, 2012). Under

physiological or pathological hypoxia or VHL mutation, HIFα accumulates in the nucleus where

it complexes with the stable HIFβ subunit and other co-activators to activate target gene

expression.

The role of HIFα, particularly the representative HIF1α, in neural precursor cells under

physiological conditions has been extensively investigated (Cunningham et al., 2012; Li et al.,

2014; Milosevic et al., 2007; Tomita et al., 2003). Aggravated hypoxia and/or ischemia causes

severe disturbance of normal oligodendroglial myelination in the preterm infants born between

28[th] and 37[th] week gestational ages (Volpe, 2009). Animal studies demonstrate that pathological

hypoxia insult delays developmental myelination in preterm-equivalent early postnatal murine

CNS (Jablonska et al., 2012; Liu et al., 2011; Liu et al., 2013). The role of HIFα in

developmental myelination has not been defined until an important study reported that HIFα

plays a major role in hypoxia-elicited myelination disturbance in cell/brain slice culture systems

(Yuen et al., 2014). However, the cellular and molecular mechanisms underlying

oligodendroglial HIFα-regulated myelination remain incompletely defined. The current popular

hypothesis states that HIFα dysregulation disturbs developmental myelination by activating

autocrine Wnt/β-catenin signaling (Yuen et al., 2014). This "Wnt-dependent" mechanistic

hypothesis is in line with the inhibitory effect of intracellular Wnt/β-catenin activation on CNS

myelination (Guo et al., 2015), but it was only tested in the cell / slice culture systems in

combination with pharmacological manipulations. Given the intrinsic caveats of non-cellular

specificity and/or off-target effects of pharmacological application, it is imperative to determine,

by using *in vivo* genetic approaches, the involvement of oligodendroglial HIFα-derived Wnt

signaling in developmental myelination.

CNS developmental myelination consists of at least two major sequential steps: OPC
differentiation into oligodendrocytes (OLs, referred to as OPC differentiation) and
oligodendrocyte maturation and myelination (Guo et al., 2015; Huang et al., 2013). In humans,
developmental myelination starts in the third trimester of gestational ages, which is equivalent to
the perinatal and early postnatal ages in rodents (Semple et al., 2013). Therefore, we used the
murine CNS of perinatal and early postnatal ages as a third trimester-equivalent *in vivo* model to
dissect the cellular and molecular mechanisms underlying development myelination. By
analyzing a series of cell-specific HIFα and Wnt genetic mutant mice, we provided convincing
evidence supporting an alternative HIFα model of CNS myelination: HIFα transiently regulates
developmental myelination by controlling OPC differentiation but subsequent OL maturation and
its hyperactivation disturbs OPC differentiation in a manner independent of autocrine Wnt/β-
catenin signaling. Our results further demonstrate that sustained Sox9 activation is a
downstream mechanism underlying disturbed OPC differentiation elicited by HIFα
hyperactivation.

Materials and Methods
Transgenic animals: The following transgenic mice were used in our study: *Cnp-Cre* mice
(RRID: MGI_3051754) kindly provided by Dr. Nave (Lappe-Siefke et al., 2003), *Sox10-Cre*
(RRID: IMSR_JAX:025807), *Sox10-CreER^{T2}* (RRID:IMSR_JAX: 027651) , *Pdgfrα-CreER^{T2}*
(RRID: IMSR_JAX:018280) , *Hif1α*-floxed (RRID: IMSR_JAX:007561), *Hif2α*-floxed (RRID:
IMSR_JAX:008407), *Vhl*-floxed (RRID: IMSR_JAX:012933), and *Wls*-floxed (RRID:
IMSR_JAX:012888). In our study, we found no difference in oligodendroglial phenotypes
between non-Cre mice carrying floxed alleles and Cre transgenic mice carrying no floxed alleles.
Therefore, we used non-Cre and/or Cre transgenic mice from same litters of cKO as littermate
control mice. All mice were maintained on the C57BL/6 background and housed in 12 h
light/dark cycle with free access to water and food. Animal protocols were approved by the
Institutional Animal Care and Use Committee at the University of California, Davis.

Tamoxifen treatment: Tamoxifen (TM) (T5648, Sigma) was prepared at a concentration of 30
mg/ml in a mixture of ethanol and sunflower seed oil (1:9, v/v). All study mice including
littermate control and conditional knockout were received tamoxifen intraperitoneally at a dose
of 200 μg/g body weight.

Hypoxia/ischemia (H/I)-induced brain white matter injury: We used our established

protocols (Chung et al., 2013; Shen et al., 2010) to induce H/I injury in C57BL/6J background

mice at postnatal day 6 (P6), a time point when OPC differentiation barely occurs in the brain. In

brief, P6 mice were anesthetized with indirect cooling by wet ice, which were immediately

subjected to permanent occlusion (by cauterizers) of the right side common carotid artery

(ischemia). One hour after the artery occlusion, mice were exposed to 6% O_2 for 45min

(hypoxia). After hypoxia exposure, the mice were returned to the dam. Sham operation was

performed in the same way as H/I injury including anesthesia, neck skin incision, and exposure

of the right common carotid artery but without artery cauterization or hypoxia. Previous study

demonstrates that H/I injury results in widespread periventricular white matter injury including

arrested OPC differentiation and microglial/astroglial activation (Liu et al., 2011).

Tissue processing and immunohistochemistry: After anesthetization by ketamine/xylazine

mixture, mice were perfused with cold PBS. Tissues were collected, fixed with fresh 4% PFA for

2 h at room temperature, cryopreserved in 30% sucrose overnight at 4°C and embedded in

OCT (VWR International, Wetzlar, Germany). Serial coronal sections (14 µm) were cut by a

Leica Cryostat (CM 1900-3-1). Tissue sections were incubated with primary antibody overnight

at 4°C after blocking with 10% donkey serum at room temperature for at least 1 h. After 3 times

wash, the fluorescence conjugated secondary antibody (Jackson ImmuneResearch) was

applied and incubated for 1.5 h. Then the slice was protected by mounting medium after DAPI

nucleus staining. The following primary antibodies were used in immunohistochemical study:

HIF-1α (RRID:AB_409037,1:200,Cayman Chemical #100642,), Sox10 (RRID:

AB_2195374,1:100,Santa Cruz Biotechnology, sc-17342; RRID: AB_778021, 1:500, Abcam,

ab27655), Olig2 (RRID: AB_570666,1:200, Millipore, AB9610), Mbp (RRID:AB_2564741, 1:500,

Biolegend, SMI-99,), SMI312 (RRID:AB_2135329, 1:1000, Covance, SMI-312R), CC1

(RRID:AB_213434, 1:200, Calbiochem, OP80,), PDGFRα (RRID:AB_2236897,1:200, R&D

system, AF1062), SMI32 (RRID:AB_2314912, 1:1000, Biolegend, SMI-32P), and Sox9

(RRID:AB_2194160,1:200, R&D system, AF3075).

In situ hybridization and transmission electron microscopy (TEM): Briefly, the slices were

treated with Proteinase K (Thermo Fisher, AM2548) and acetylated by triethanolamine (Sigma,

90279) and acetic anhydride (Sigma, 320102). The tissue was incubated with 100 ng/ul DIG

labelled c-RNA at 65^{0}C overnight. Then the slices were incubated with Alkaline phosphatase

(AP) anti-DIG secondary antibody (1:100,11093274910, Sigma) overnight after blocking with 10%

donkey serum. The signal was determined by nitroblue tetrazolium (NBT) and 5-bromo-4-

chloro-3-indolyl phosphate (toluidine salt) (BCIP) (Sigma, 72091).Semithin sections for toluidine

blue myelin staining and ultrathin section for TEM imaging were prepared according to our

previous protocols (Zhang et al., 2018a; Zhang et al., 2018b). TEM imaging was conduction by

the professionals at UC Davis Core Facility who were blinded to the mouse genotypes.

Enzyme-linked immunosorbent assay (ELISA): The Wnt3a protein level was measured by

the mouse Wnt3a ELISA kit (M13146365, CSB-EL026136MO, Cusabio) according to the

manufacturer's instruction.

Primary OPC culture and gene manipulation: The mouse cortex between P0 and P2 was

collected and dissociated with papain kit and DNase I (#LK003176, Worthington; 250U/ml;

#D5025, Sigma). OPCs were purified by immnopanning approach according to our previous

protocol (Lang et al., 2013). Purified OPCs were used for experiments of Wls knockdown,

Wnt3a overexpression, and Sox9 overexpression. Wls-shRNA (TRCN 0000 234932, Mission

ShRNA bacterial Glycerol stock NM_026582) and scramble control (Mission TRC2 PlkO.5-

PURO Non-Mammalian shRNA control Plasmid) were purchased from Sigma-Aldrich. Wnt3a

plasmid pLCN-Wnt3a-HA (Addgene, #18030) and empty pLCN-exp (Addgene, #64865) were

used for Wnt3a overexpression experiment. The transfection of Wls-shRNA and pLCN-Wnt3a-

HA and controls was performed by using HiPerFect transfection reagent (#301704, Qiagen)

according to the manual. p-WPXL-Sox9 (Addgene, #36979) and control plasmid p-WPXL

(Addgene, #12257) were used for Sox9 overexpression and the transfection was performed by

using FuGENE6 Transfection reagent (Promega, #E2691, lot#000371257).

Primary rat OPCs were used in Sox9 knockdown and DMOG treatment. DMOG (0.5 mM

dissolved in DMSO, D3695, Sigma), and Sox9 siRNA (SASI_Rn02_00372739, Sigma) or

negative control siRNA (SIC001-5X1NMOL, Sigma) were applied according to the experimental

design in Figure 12H. Transfection of Sox9 siRNA was conducted by using HiPerFect

transfection reagent (#301704, Qiagen).

Purification of brain OPCs by magnetic activated cell sorting (MACS): MACS purification of

OPCs was performed according to the protocols in the kit instruction from Miltenyi Biotec. The

mouse forebrain was dissociated using the papain dissociation system (LK003176, Worthington)

and gentleMACS^{TM} Dissociator (Miltenyi Biotec, #130-092-235). Astrocytes and microglia in the

cell suspension were removed by using anti-ACSA-2 micro beads (130-097-679, Miltenyi Biotec)

and anti-CD11b micro beads (130-049-601, Miltenyi Biotec), respectively. After astroglial and

microglial removal, the cell suspension was then incubated with anti-O4 micro beads (130-094-

543, Miltenyi Biotec) for OPC purification. O4$^+$ cells were collected for RNA preparation.

RNA preparation, reverse transcription and real-time quantitative PCR (RT-qPCR): Total

RNA was obtained using Qiagen RNeasy for lipid tissue kit (74804, Qiagen) according to the

manual. We used on-column DNase I digestion to eliminate DNA contamination.

Complementary DNA (cDNA) was prepared using Qiagen Omniscript RT Kit (205111, Qiagen).

Mouse and rat gene expression was normalized to the internal control Hsp90 and β-actin,

respectively, and calculated by the equation of $2^{\wedge(\text{Ct(cycle threshold) of Hsp90 - Ct of indicated genes})}$. The value

of control groups was normalized to 1 throughout the study. Real time qPCR was done by using

QuantiTect SYBR® Green PCR Kit (204145, QIAGEN) and Stratagene MP3005P thermocycler.

The qPCR primers used in the study were shown as below.

Mouse Wls: AGGGCAAGGAAGAAGGAGAG / ATCCCTCCAACAATGCAGAG

Mouse Glut1: CAGTTCGGCTATAACACTGGTG / GCCCCCGACAGAGAAGATG

Mouse Ldha: CATTGTCAAGTACAGTCCACACT / TTCCAATTACTCGGTTTTTGGGA

Mouse Hk2: TGATCGCCTGCTTATTCACGG / AACCGCCTAGAAATCTCCAGA

Mouse Axin2: AACCTATGCCCGTTTCCTCTA / GAGTGTAAAGACTTGGTCCACC

Mouse Naked1: CAGCTTGCTGCATACCATCTAT / GTTGAAAAGGACGCTCCTCTTA

Mouse Wnt7a: CGACTGTGGCTGCGACAAG / CTTCATGTTCTCCTCCAGGATCTTC

Mouse Wnt7b: CTTCACCTATGCCATCACGG / TGGTTGTAGTAGCCTTGCTTCT

Mouse Vegfa: GCACATAGAGAGAATGAGCTTCC / CTCCGCTCTGAACAAGGCT

Mouse Mbp: ACACGAGAACTACCCATTATGGC / CCAGCTAAATCTGCTGAGGGA

Mouse Plp (exon3b-specific): CCAGAATGTATGGTGTTCTCCC /

GGCCCATGAGTTTAAGGACG

Mouse Cnp: TTTACCCGCAAAAGCCACACA / CACCGTGTCCTCATCTTGAAG

Mouse Mobp: AACTCCAAGCGTGAGATCGT / CAGAGGCTGTCCATTCACAA

Mouse Opalin: CTGCCTCTCACTCAACATCA / GCTGGATCAAAGTAAACAGC

Mouse Mog: AGCTGCTTCCTCTCCCTTCTC / ACTAAAGCCCGGATGGGATAC

Mosue Qk: CTGGACGAAGAAATTAGCAGAGT / ACTGCCATTTAACGTGTCATTGT

Mouse Myrf: CAGACCCAGGTGCTACAC / TCCTGCTTGATCATTCCGTTC

Mouse Sox9: AGTACCCGCATCTGCACAAC / ACGAAGGGTCTCTTCTCGCT

Mouse Hsp90: AAACAAGGAGATTTTCCTCCGC / CCGTCAGGCTCTCATATCGAAT

Rat Bnip3: GCTCCCAGACACCACAAGAT / TGAGAGTAGCTGTGCGCTTC

Rat Mbp: TTGACTCCATCGGGCGCTTCTTTA / TTCATCTTGGGTCCTCTGCGACTT

Rat Sox9: CTGAAGGGCTACGACTGGAC / TACTGGTCTGCCAGCTTCCT

Rat β-actin: CGTCTTCCCCTCCATCGT / GGAGTCCTTCTGACCCATACC

Chromatin immunoprecipitation (ChIP) by HIFα antibody and qPCR verification: ChIP was

performed using the SimpleChIP plus Enzymatic Chromatin IP kit (Cell Signaling Technology,

#9005) following the manufacturer's protocol with some modifications. Briefly, primary cultured

rat OPCs (4 x 10^6 cells) treated with 1 mM DMOG for 7 hours were crosslinked in culture

medium containing 1% formaldehyde at room temperature for 10 min. After addition of glycine

solution to stop the reaction, the cells were then collected, centrifuged and lysed. Nuclei were

collected and treated with 0.5 ul micrococcal nuclease per IP for 20 min at 37 °C. The reaction

was stopped using 0.05 M EDTA, and nuclear membranes were destroyed by sonication. Then

the supernatant was collected by centrifugation. Chromatin solutions were incubated with 10 ug

of anti-HIF1α (Cayman, Rabbit, #10006421) antibody or anti-rabbit IgG (Cell Signaling #2729)

overnight at 4 °C, and immunoprecipitated with protein G magnetic beads for 2 hours at 4 °C

with rotation. Bound DNA-protein complexes were washed and eluted, and crosslinks were

reversed according to kit instructions. DNA was purified using spin columns and used for ChIP-

qPCR analysis. The fold enrichment of specific genomic regions was assessed relative to the

qPCR data from the IPs normalized to the control IgG values. Fold enrichment = $2^{\wedge\text{-}[(Ct\ IP) - (Ct\ IgG)]}$.

qPCR primer sets for ChIP binding site verification are:

Negative site: forward TTCCTCTTGGGATGGTTGTC / reverse CCACCTCTGGGGAGTATGAA

-75bp site: forward TTTCAAAATCCGGTCCAATC / reverse CCCCCTTCACCTTAGAGACC

-828bp site: forward CGGGGAAGGACTTGTCAGT / reverse ATGAAAACCAAAGCCAAGCA

-1211bp site: forward CGGCTCCAAGCACTCTTAAA /reverse GCGTCCTTTTAGACCTGCAC

-1928 site: forward GTCGTTCTCGCTGCCTTTAG / reverse TTGAAGAGACCAGGGACCAC

-2098 site: forward CTCTGGATGTTGCCGAAAAT / reverse CTCCACGCAAGCGTTTTTAT

Western blot and antibodies: Twenty microgram protein was loaded into AnykD Mini-

PROTEAN gel (#4569035, BIO-RAD). After blocking, the membrane was incubated with primary

antibodies overnight. The membrane was washed 3 times by TBST (TBS with 1% tween 20).

Then the HRP-conjugated secondary antibody was reacted with membrane for 1.5 hours.

Protein signals were developed by using ECL kit (NEL103001EA, Perkin Elmer). Quantification

of interested protein bands were conducted by using NIH Image J software. Antibodies used in

western blot are MBP (SMI-99, RRID:AB_2564741, 1:2000, Biolegend), Active β-catenin (#05-

665, RRID: AB_309887, 1:1000, Millipore), Axin-2(#6163, RRID: AB_10904353, 1:1000, Prosci)

and β-actin (3700, RRID:AB_2242334, 1:2000, Cell signaling Technology).

Experimental design and statistical analyses: Data collection was conducted by lab

members who were blinded to mouse genotypes. We used Shapiro-Wilk approach to test data

normality. F test and Browne-Forsythe test were used to variance equality of two groups and

three or more groups, respectively. We used scatter graphs to present the quantification data

throughout the current manuscript. Both male and female mice were included in our data

analysis. All data were presented as mean ± standard effort of mean (s.e.m.) where each dot

represents one mouse. Unpaired two-tailed Student's t test was used for statistically analyzing

two groups of data. The *t*-values and the degree of freedom (df) were shown as $t_{(df)}$ in each

graph. One-way ANOVA followed by Tukey's post-hoc test was used for statistically analyzing

three or more groups of data. The F ratio, and DFn and DFd was presented as $F_{(DFn, DFd)}$ in the

figure legends where DFn stands for degree of freedom numerator and DFd for degree of

freedom of denominator. The exact P-value of one-way ANOVA test was also described in the

figure legends. All data graphing and statistical analyses were performed using GraphPad Prism

version 8.0. The P-value was designated as * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, ns, not

significant $p > 0.05$.

Results

Transient HIFα stabilization during early CNS development

HIF1α and HIF2α share high sequence homology and function in a similar manner

through partnering with HIF1β and subsequently activating HIFα target genes (Semenza, 2012).

Therefore, we used antibodies against the representative HIF1α to determine whether HIFα

protein is stabilized in the perinatal and early postnatal CNS, where the vasculature is still in

immature state (Harb et al., 2013; Yuen et al., 2014). We found that HIF1α protein was detected

by immunohistochemistry (IHC) in Sox10[+] oligodendroglial lineage cells in the white matter

areas of the spinal cord (**Fig. 1A**) and the forebrain (**Fig. 1C**) at postnatal day 5 (P5). As positive

controls, HIF1α immunoreactive signals were markedly increased in age-matched transgenic

Sox10-Cre:Vhl[fl/fl] mice (**Fig. 1B, D**), in which *Sox10-Cre*-mediated VHL disruption prevents HIFα

from degradation and causes HIFα accumulation specifically in oligodendroglial lineage cells.

HIF1α signals became barely detectable in the forebrain white matter by P10 (**Fig. 1E**), which

temporally approximates to the peak of CNS angiogenesis (Harb et al., 2013; Yuen et al., 2014),

and remarkably upregulated in mice that had been challenged by hypoxia/ischemia injury at P6

(**Fig. 1F**) (Chung et al., 2013; Liu et al., 2011; Shen et al., 2010). Our quantification data

showed that the density of HIF1α$^+$Sox10$^+$ cells were significantly increased in P10 H/I-injured

forebrain white matter compared with sham controls (4.2 ± 4.2 sham, 312 ± 9.7 H/I, mean ±

s.e.m., n=3, $t_{(4)}$ = 29.3 P < 0.0001). Consistent with HIF1α histological upregulation, Western

blot assays of micro-dissected subcortical white matter demonstrated a substantial elevation of

HIF1α and its canonical target gene PKM2 in H/I-injured brain (**Fig. 1G**) which were also

confirmed by PKM2 IHC (**Fig. 1H**). Concomitant with HIFα activation, myelin staining by myelin

basic protein (MBP) showed severe disturbance of developmental myelination in H/I-injured

white matter compared with sham control (**Fig. 1I1-3**). These results demonstrate that HIFα is

transiently stabilized in the early postnatal CNS but rapidly downregulated in the second

postnatal week and that hypoxia/ischemia injury results in sustained HIFα stabilization in the

brain white matter.

HIFα hyperactivity impairs OPC differentiation but not oligodendrocyte maturation

Pathological hypoxia-ischemia insults caused aberrant HIFα activation in the white

matter (**Fig. 1F-H**) and hypomyelination (**Fig. 1I1-3**). To determine whether aberrant HIFα

stabilization disturbs normal developmental myelination, we induced HIFα hyperactivity by

conditionally disrupting the negative regulator VHL (cf **Fig. 1B, D**). Constitutive *Sox10-Cre:Vhl$^{fl/fl}$*

(Sox10:VHL cKO) pups were much smaller than littermate controls (*Vhl$^{fl/+}$*, or *Vhl$^{fl/fl}$*) at birth and

rarely survived beyond P7. Our prior data show that *Sox10-Cre*-mediated gene disruption

initiates in early OPCs during embryonic CNS development (Zhang et al., 2018a). *Sox10-Cre*-

elicited HIFα hyperactivity (cf **Fig. 6A**) caused severe hypomyelination in the developing

postnatal spinal cord (**Fig. 2A**). The number of CC1$^+$ mature OLs (**Fig. 2B, C**) and the mRNA

level of major myelin protein (**Fig. 2D**) were reduced by ~40-60% in Sox10:VHL cKO mutants

compared with littermate controls. However, the number of PDGFRα$^+$ OPCs was unaffected in

Sox10:VHL cKO mutants (**Fig. 2C**). These data suggest that constitutive HIFα activation in

OPCs inhibits OPC differentiation and disturbs developmental myelination.

To determine the effect of HIFα hyperactivity on subsequent oligodendrocyte maturation,

we analyzed *Cnp-Cre:Vhl$^{fl/fl}$* (Cnp:VHL cKO) mutants. Previous studies have demonstrated that,

compared with *Sox10-Cre*, *Cnp-Cre* drives gene disruption in later stages of oligodendrocyte

development (Moyon et al., 2016; Zhang et al., 2018a; Zhang et al., 2018b). Cnp:VHL cKO mice

were viable and indistinguishable from their littermate controls (*Vhl*$^{fl/fl}$, *Vhl*$^{fl/+}$, or *Cnp-Cre:Vhl*$^{fl/+}$)

throughout the early postnatal and adult ages. In sharp contrast to Sox10:VHL cKO mice,

Cnp:VHL cKO mice had normal levels of myelin gene expression (**Fig. 2E, F, H**), unperturbed

myelination evidenced by toluidine blue myelin staining (**Fig. 2G,** left) and transmission electron

microscopic (TEM) myelination assay (**Fig. 2G**, right), and normal numbers of OLs and OPCs

(**Fig. 2I**) despite the elevated density of blood vessels in the CNS we recently reported (Guo et

al., 2015; Zhang et al., 2020). These data indicate that HIFα hyperactivity in OLs does not affect

their maturation.   Taken together, our genetic evidences convincingly demonstrate elevated

HIFα activity perturbs developmental myelination by inhibiting OPC differentiation but not OL

maturation, suggesting a lineage stage-dependent role of HIFα in developmental myelination.

We next used time-conditional and OPC-specific genetic model to circumvent the early

death of Sox10:VHL cKO mice and interrogate whether time-conditional HIFα activation in

postnatal OPCs impacts their differentiation. Towards this end, we generated *Pdgfra-*

CreER^{T2}*:Vhl*$^{fl/fl}$ (PDGFRα:VHL cKO) mutants. Our data showed that OPC-specific HIFα

activation during neonatal ages (**Fig. 3A-B**) significantly reduced the expression of genes

encoding major myelin proteins (**Fig. 3C**) and decreased the densities of mature OLs but not

OPCs in the spinal cord (**Fig. 3D**). IHC of myelin protein PLP demonstrated a marked

hypomyelination (**Fig. 3E**). Further analysis of myelination by transmission electron microscopy

(**Fig. 3F**) demonstrated that the density of myelinated axons was reduced by ~42% in the

ventral spinal white matter of PDGFRα:VHL cKO mice compared with littermate control (**Fig.**

3G), which is consistent with histological myelin protein assays (**Fig. 3F**) . To assess the long-

term effect of HIFα activation, we induced HIFα activation by tamoxifen injections to neonatal

mice (**Fig. 3H-I**) and analyzed oligodendroglial phenotypes in the adult CNS (P60-P70). We

found that the number of OLs but not OPCs was persistently reduced in the corpus callosum of

PDGFRα:VHL cKO compared with control mice (**Fig. 3J**), suggesting that neonatal HIFα

activation inhibits OPC differentiation in the adult CNS. Given that virtually all axons in the optic

nerve are myelinated in the adult rodents (Bercury and Macklin, 2015), we then analyzed

myelinated axons visualized by toluidine blue myelin staining of semithin resin sections (**Fig.**

3K1-K4). Our quantification demonstrated a ~44% reduction in the density of myelinated axons

in the optic nerve of adult PDGFRα:VHL cKO mutants compared with age-matched fully

myelinated control optic nerves (**Fig. 3K5**). Together, our data suggest that HIFα activation

inhibits long-term OPC differentiation and results in hypomyelination throughout postnatal CNS

development.

Taken together, our *in vivo* data derived from three different genetic mouse models
suggest that HIFα hyperactivity disturbs developmental myelination by inhibiting OPC
differentiation.

Disrupting HIFα in OPCs but not OLs delays developmental myelination

Transient HIFα stabilization (**Fig. 1A-E**) suggests that HIFα may regulate OPC
differentiation and/or myelination. To define the role of physiological HIFα stabilization in
developmental myelination, we analyzed oligodendroglial phenotypes in HIFα cKO genetic mice.
Our extensive analysis of HIF1α or HIF2α conditionally knockout mice driven by *Pdgfra-CreER^{T2}*
and *Cnp-Cre* revealed no phenotypic abnormalities of the oligodendroglial lineage, suggesting
functional redundancy between HIF1α and HIF2α. We therefore generated HIF1α and HIF2α
double conditional knockout (referred to as HIFα cKO).

To determine the role of HIFα in OPC differentiation, we used *Pdgfra-*
CreER^{T2}:Hif1α^{fl/fl}:Hif2α^{fl/fl} (PDGFRα:HIFα cKO) transgenic mice. Tamoxifen was administered to
PDGFRα:HIFα cKO and littermate control (*Hif1α^{fl/fl}:Hif2α^{fl/fl}*) pups at P1, P2, and P3 and the CNS
tissue was analyzed at P8. Our previous data show that this tamoxifen paradigm induces >85%
of gene cKO efficiency in OPCs (Zhang et al., 2018a; Zhang et al., 2018b). The density of
differentiated OLs (CC1$^+$Olig2$^+$) but not OPCs (PDGFRα$^+$Olig2$^+$) was significantly decreased in
the spinal cord white matter of PDGFRα:HIFα cKO mice compared with HIFα controls (**Fig. 4A-**
B), suggesting that HIFα regulates OPC differentiation and is dispensable for OPC population
expansion. In agreement of impaired OPC differentiation, the mRNA levels of major myelin
protein (*Mbp* and *Plp*) and mature OL-enriched protein (Opalin and QK, aka CC1) were
significantly reduced in PDGFRα:HIFα cKO mice compared with littermate controls (**Fig. 4C**). In
the forebrain, developmental myelination in the periventricular white matter was delayed in
PDGFRα:HIFα cKO mice at P8 (**Fig. 4D**) and became indistinguishable from that in littermate
control mice at P14 (**Fig. 4E**). These data suggest that HIFα regulates the timing of
developmental myelination.

To determine the role of HIFα in oligodendroglial maturation, we used *Cnp-*
Cre:Hif1α^{fl/fl}:Hif2α^{fl/fl} (Cnp:HIFα cKO) and littermate control (*Hif1α^{fl/fl}:Hif2α^{fl/fl}*) mice. Cnp:HIFα
cKO mice were born at expected Mendelian ratios and displayed no behavioral abnormalities
during postnatal development and throughout adult ages. The expression of canonical HIFα
target gene *Ldha* and *Vegfa* (Sharp and Bernaudin, 2004) was significantly reduced in
Cnp:HIFα cKO mice (**Fig. 4F**), thus validating the disruption of HIFα function. Unexpectedly, we
found no significant difference in developmental myelination (**Fig. 4G**), the number of

CC1$^+$OLig2$^+$ mature OLs (**Fig. 4H**) and PDGFRα$^+$Olig2$^+$ OPCs (**Fig. 4I**), and myelin gene

expression (**Fig. 4J**) between Cnp:HIFα cKO and littermate controls mice at different ages (P2,

P8, and P14). Double IHC showed that some CC1$^+$ OLs were positive for HIF1α in the

periventricular white matter of the brain at P8 (**Fig. 4K,** arrowheads) but not at P14 (not shown).

These data indicate that HIFα plays a minor role in oligodendrocyte maturation. Collectively, the

contrast phenotypes derived from PDGFRα:HIFα cKO vs Cnp:HIFα cKO mice support a working

model that HIFα controls the timing of developmental myelination by regulating the

differentiation of OPCs into OLs but not subsequent oligodendrocyte maturation.

Disrupting oligodendroglial HIFα does not affect axonal integrity or cell survival.

HIFα is required for neuronal survival, as HIFα deficiency in Nestin-expressing neural

cells results in massive neuronal death and neurodegeneration (Tomita et al., 2003). Recent

data reported that HIFα deficiency in Olig1-expressing neural cells leads to neuronal death and

widespread axonal injury in the corpus callosum at the perinatal ages (Yuen et al., 2014). To

determine whether OPC or OL-specific HIFα deficiency causes axonal damage and cell death,

we employed IHC of SMI32 monoclonal antibody and cleaved Caspase 3 (CC3) antibody.

SMI32 labels functionally compromised axons (Soulika et al., 2009; Yuen et al., 2014) and a

subset of pyramidal neuronal cell bodies and dendrites during normal development (Campbell

and Morrison, 1989; Haynes et al., 2005; Voelker et al., 2004) (**Fig. 5A**, Ctx). We did not

observe SMI32 immunoreactive signals in the corpus callosum of Cnp:HIFα cKO (**Fig. 5A**) and

PDGFRα:HIFα cKO (**Fig. 5C**) mice compared with respective littermate controls. In contrast,

many tightly packed axons running through the corpus callosum were clearly labeled by the

pan-axonal marker SMI312 (**Fig. 5B, D**). Furthermore, the densities of CC3$^+$ apoptotic cells and

CC3$^+$/Sox10$^+$ apoptotic oligodendroglial lineage cells were statistically indistinguishable

between PDGFRα (or Cnp):HIFα cKO mutants and their respective littermate controls (**Fig. 5E-**

G). We employed TEM to assess the structural integrity of axons in the corpus callosum of

PDGFRα:HIFα cKO mice at P8, a time point that few axons are myelinated yet. The density of

axons with diameter ≥ 400nm, which will be myelinated by oligodendrocyte during postnatal

development (Hines et al., 2015; Lee et al., 2012), was similar between PDGFRα:HIFα cKO

(458,401 ± 59,957 axons/mm^2) and littermate control (485,558 ± 38,780 axons/mm^2, $t_{(4)} = 0.4$, p

= 0.72), suggesting that axons in HIFα cKO mice are not subjected to degeneration. We did not

find visible morphological differences in axonal structure (**Fig. 5H-I**, Ax) and axonal

mitochondria (**Fig. 5H-I**, Mt) between PDGFRα:HIFα cKO and littermate control mice. Thus,

disrupting oligodendroglial HIFα does not affect neuronal cell survival and axonal integrity in the

white matter in the early postnatal CNS.

Oligodendroglial HIFα plays a minor role, if any, in regulating Wnt/β-catenin activity.

Previous study reported that aberrant HIFα stabilization in OPCs activates Wnt7a/Wnt7b

expression and canonical Wnt signaling (i.e. Wnt/β-catenin) (Yuen et al., 2014). The unbiased

RNA-seq analysis of *Sox10-Cre:Vhl*$^{fl/fl}$ mice prompted us to revisit HIFα-Wnt regulation in

oligodendroglial lineage cells (**Fig. 6A-E**). Sox10-Cre-mediated VHL cKO markedly elevated

HIFα activity, as evidenced by increased expression of canonical HIFα target genes *HK2, Ldha,*

Glut1, Bnip3, and *Vegfa* (**Fig. 6A**). In stark contrast to HIFα activation, there were no significant

changes in the expression of Wnt genes *Wnt7a* and *Wnt7b* and canonical Wnt signaling target

genes *Axin2* and *Naked1* (**Fig. 6B**). The KEGG pathway of HIF-1 signaling and the GO term of

response to hypoxia were significantly overrepresented amongst 255 upregulated genes (**Fig.**

6C, D) whereas the GO terms of oligodendrocyte differentiation and sterol biosynthesis were

significantly overrepresented amongst 351 downregulated genes (**Fig. 6E**), which is in

agreement with elevated HIFα signaling and diminished OPC differentiation in the CNS of

Sox10:VHL cKO mice we documented in **Fig. 2A-D**. However, we did not find

overrepresentation of Wnt-related KEGG pathways or GO terms amongst the differentially

regulated genes. The unbiased analysis of gene regulation at the transcriptomic level strongly

suggests that oligodendroglial HIFα stabilization plays a minor role in regulating canonical Wnt

signaling.

To strengthen our conclusion derived from RNA-seq, we analyzed OPCs isolated from

PDGFRα:VHL cKO mice. Tamoxifen was injected to PDGFRα:VHL cKO and littermate control

mice at P5 and brain OPCs were acutely purified at P6 (**Fig. 6F**) by magnetic activated cell

sorting (MACS). As shown in **Fig. 6G**, HIFα was indeed functionally stabilized in purified VHL-

disrupted OPCs, as evidenced by the elevated expression of HIFα target gene *Hk2, Ldha*, and

Glut1. However, there were no significant changes in the expression of canonical Wnt/β-catenin

signaling target genes *Axin2* and *Naked1* and Wnt ligands *Wnt7a* and *Wnt7b* in VHL-disrupted

OPCs compared with VHL-intact OPCs (**Fig. 6H**), suggesting that genetically stabilizing HIFα

does not perturb the activity of autocrine Wnt/β-catenin or the expression of Wnt7a and Wnt7b

in OPCs.

To corroborate our findings, stabilizing HIFα in PDGFRα$^+$ OPCs did not affect the activity

of Wnt/β-catenin signaling in the spinal cord at P8 (tamoxifen at P1, P2, and P3) and in the

forebrain at P14 (tamoxifen at P6 and P7) of PDGFRα:VHL cKO mice compared with littermate

controls, which was supported by the unaltered expression of canonical Wnt target genes at the

mRNA and protein levels (**Fig. 6I-J**). We also found that the expression of Axin2, Naked1,

Wnt7a, and Wnt7b was not perturbed in PDGFRα:HIFα cKO mice compared with littermate

controls (**Fig. 6K**). Thus, HIFα does not regulate canonical Wnt signaling activity or

Wnt7a/Wnt7b expression in the oligodendroglial lineage and in the CNS.

Blocking oligodendroglial-derived Wnt signaling does not affect HIFα hyperactivation-

elicited inhibition of OPC differentiation and myelination.

That non-perturbation of Wnt signaling in HIFα-stabilized OPCs led us to hypothesize

that HIFα hyperactivation inhibits myelination in a manner independent of autocrine Wnt

activation in OPCs. To test our hypothesis, we stabilized HIFα function and simultaneously

blocked oligodendroglial-derived autocrine Wnt signaling.

Previous studies demonstrated that disrupting WLS blocks the secretion of Wnt ligands

from Wnt-producing cells and inhibits intracellular Wnt signaling activity in Wnt-receiving cells

(Banziger et al., 2006; Bartscherer et al., 2006). To determine whether WLS deficiency blocks

autocrine Wnt/β-catenin activity in OPCs, we knocked down WLS in Wnt3a-expressing primary

OPCs (**Fig. 7A-C**). We chose Wnt3a because it is a typical Wnt ligand that activates the

canonical Wnt signaling in Wnt-receiving cells. Wnt3a expression in OPCs significantly

increased Wnt3a concentration and simultaneous WLS knockdown abrogated Wnt3a elevation

in the culture medium of primary OPCs (**Fig. 7D**), suggesting that WLS is required for Wnt3a

secretion from OPCs. We found that the activity of the canonical Wnt signaling, as assessed by

Wnt target genes *Axin2* and *Sp5*, was upregulated in Wnt3a-expressing OPCs but blocked in

Wnt3a-expressing/WLS-deficient OPCs (**Fig. 7E, F**). These results demonstrate that disrupting

WLS blocks OPC-derived autocrine Wnt/β-catenin signaling activity.

Genetically disrupting WLS in PDGFRα$^+$ OPCs had no rescuing effects on HIFα

hyperactivation-elicited hypomyelination (**Fig. 8A-C**), impaired OPC differentiation (**Fig. 8D-E**),

and diminished myelin gene expression (**Fig. 8F**) in PDGFRα:VHL/WLS cKO mice, suggesting

that autocrine Wnt signaling is unlikely a downstream pathway in mediating HIFα stabilization-

induced inhibition of OPC differentiation. To corroborate this conclusion, we used a different

inducible genetic model, *Sox10-CreER*^{T2}:*Vhl*$^{fl/fl}$:*Wls*$^{fl/fl}$ (iSox10:VHL/WLS cKO) (**Fig. 9A**).

Conditionally stabilizing HIFα in Sox10-expressing oligodendroglial lineage cells resulted in

impaired OPC differentiation and hypomyelination in iSox10:VHL cKO mice, however,

simultaneous WLS cKO did not affect the degree of inhibited OPC differentiation and

hypomyelination in iSox10:VHL/WLS cKO mice (**Fig. 9B-E**). We further demonstrated that

disrupting WLS alone in Sox10-expressing oligodendroglial lineage cells had no detectable

effect on normal OPC differentiation and developmental myelination in *Sox10-CreER*[T2]:*Wls*[fl/fl]

(iSox10:WLS cKO) mice compared with non-Cre littermate control mice (**Fig. 10**). Collectively,

our results derived from cell-specific genetic approaches suggest that autocrine Wnt/β-catenin

signaling plays a dispensable role in HIFα-regulated OPC differentiation and myelination.

Sustained Sox9 activation mediates HIFα stabilization-elicited inhibition of OPC

differentiation

The mRNA level of Sox9 was significantly upregulated in primary OPCs maintained in

growth medium that had been treated with dimethyloxalylglycine (DMOG), a potent HIFα

stabilizer (Yuen et al., 2014), and this upregulation was abrogated upon simultaneous

treatment of chetomin (**Fig. 11A**), an inhibitor of HIFα transcriptional activity (Kung et al., 2004;

Viziteu et al., 2016). To support the functional regulation, our assay of chromatin

immunoprecipitation followed by qPCR assay (ChIP-qPCR) showed that HIFα physically bound

to the promoter region (-828 bp) of the rat *Sox9* gene (**Fig. 11B**). We found that Sox9 mRNA

expression was significantly upregulated in the spinal cord of PDGFRα:VHL cKO mice

compared with littermate controls at P8 (**Fig. 11C**) and P14 (**Fig. 11D**). At the cellular level,

sustained Sox9 expression was observed in HIFα-stabilized Sox10-expesing oligodendroglial

lineage cells in Sox10:VHL cKO mice (**Fig. 11E**, right, arrowheads), in sharp contrast to control

mice in which Sox9 was expressed at much lower level or barely detectable in Sox10-

expressing cells (**Fig. 11E**, left, arrowheads). Our quantification showed that the percentage of

Sox10[+] cells expressing Sox9 (**Fig. 11F**) and the level of Sox9 mRNA (**Fig. 11G**) were

significantly increased in Sox10:VHL cKO mice. Furthermore, Sox9 expression was significantly

elevated in primary OPCs, which were purified from Sox10:VHL cKO neonatal brain (**Fig. 11H**).

Sox9 is expressed in virtually all astrocytes in the postnatal CNS (Sun et al., 2017; Zhang et al.,

2018b). We found that the density of Sox9[+]GFAP[+] astrocytes was not significantly different in

the spinal cord of Sox10:VHL cKO mutants from that in control mice (**Fig. 11I**). We also found

that the percent of Sox10[+] oligodendroglial cells expressing Sox9 (**Fig. 11J1-J2**) was

unchanged in PDGFRα:HIFα cKO mice (**Fig. 11J3**).

Recent data demonstrate that Sox9 is rapidly downregulated upon OPC differentiation

(Reiprich et al., 2017), prompting us to hypothesize that sustained Sox9 expression may inhibits

OPC differentiation. To test this hypothesis, we overexpressed Sox9 in primary OPCs in growth

medium and assess OPC differentiation in differentiating medium (**Fig. 12A**). Sox9 mRNA and

protein were significantly increased in primary OPCs transfected with Sox9-expressing plasmid

compared with those transfected with empty vector (EV) (**Fig. 12B-D**). Sox9 overexpression

reduced the mRNA expression of the most abundant myelin protein Plp and the potent pro-

differentiation factor Myrf (Emery et al., 2009) assessed by RT-qPCR (**Fig. 12 E)**, the number of

ramified MBP$^+$/Sox10$^+$ oligodendrocytes assessed by immunocytochemistry (**Fig. 12F**), and the

protein level of MBP assessed by Western blot (**Fig. 12G**). To define the role of Sox9 activation

in HIFα-regulated OPC differentiation, we stabilized HIFα in OPCs by DMOG treatment and

simultaneously knocked down Sox9 expression by siRNA (**Fig. 12H-J**). Our data showed that

Sox9 downregulation rescued the degree of myelin gene inhibition elicited DMOG treatment, as

evidenced by the density of MBP$^+$/Sox10$^+$ differentiated OLs with ramified morphology (**Fig.**

12K-L) and the expression of myelin gene *Mbp* (**Fig. 12M**). Collectively, our results suggest that

sustained Sox9 expression is one of the downstream molecular pathways in mediating HIFα-

regulated OPC differentiation.

Discussion

Summary of key findings and conclusions

Through the analysis of a series of HIFα genetic models (stage-dependent HIFα loss or

gain-of-function in OPCs and OLs), our study provides compelling evidences supporting an

alternative understanding of the cellular and molecular mechanisms underlying HIFα-regulated

CNS myelination (**Fig. 12N**). At the cellular level, our in vivo genetic data support a new working

model in which physiological HIFα transiently regulates OPC differentiation whereas persistent

HIFα activation directly causes arrested OPC differentiation, leading to developmental

hypomyelination. At the molecular level, HIFα regulates OPC differentiation in a manner

independent of the autocrine Wnt/β-catenin signaling. Instead, dysregulated Sox9 expression is

one of the downstream pathways in mediating HIFα activation-elicited inhibition of OPC

differentiation (**Fig. 12N**).

HIFα regulates developmental myelination by controlling OPC differentiation but not

subsequent oligodendrocyte maturation

One of the unexpected observations is the differential responses of OPCs and OLs to

HIFα disruption and hyperactivation. Disrupting or stabilizing HIFα function results in severe

disturbance of myelin gene expression and developmental myelination in the CNS of OPC-

specific PDGFRα:HIFα (VHL) cKO mice. In sharp contrast, HIFα disruption or stabilization in

Cnp:HIFα (VHL) cKO mice does not perturb normal myelin gene expression and myelination.

Previous genetic studies suggest that *Cnp-Cre* elicits gene disruption primarily in the later

stages of oligodendrocyte development. For example, disrupting DNA methyltransferase 1

(DNMT1) by *Cnp-Cre* line shows no phenotypes in OPC proliferation and expansion, in sharp

contrast to DMNT1 disruption by *Olig1-Cre* which targets the progenitor stages of the

oligodendroglial lineage (Moyon et al., 2016). Our recent studies also demonstrated that

disrupting Sox2 by *Cnp-Cre* shows no phenotypes in OPC proliferation and expansion, in sharp

contrast to Sox2 disruption by *Sox10-Cre* or *Pdgfrα-CreER^{T2}* which targets OPCs (Zhang et al.,

2018a; Zhang et al., 2018b). Therefore, the discrepant phenotypes in HIFα (or VHL) deletion

driven by Cnp-Cre versus *Sox10-Cre* or *Pdgfrα-CreER^{T2}* (**Fig. 2, 4, 5**) support the interpretation

that HIFα play a major role in OPC differentiation into OLs and is dispensable for

oligodendrocyte maturation.

We observed impaired OPC differentiation in HIFα activation (Fig. 2A-C, Fig. 3) and

inactivation (Fig. 4A-D) genetic models. These observations suggest delicate downstream

pathways regulated by HIFα. We recently reported that HIFα depletion reduces the canonical

HIFα target genes involving in glucose metabolism and CNS angiogenesis (Zhang et al., 2020),

both of which entails energy supply for OPC differentiation(Yuen et al., 2014). It is possible that

HIFα inactivation may transiently affect OPC differentiation through limiting the energy supply

regulated by physiological HIFα whereas sustained HIFα activation might stimulate additional

non-canonical HIFα target genes which are otherwise non-responsive to physiological HIFα

during normal development. Our experimental data identified one of such non-canonical targets

as Sox9 whose expression is upregulated in HIFα activation mice and unaltered in HIFα

inactivation mice. Furthermore, the unaltered developmental myelination in OL-specific HIFα

activation and inactivation models indicates that OLs are resistant to HIFα dysregulation, which

is in line with the previous concept that OLs are remarkably resistant to hypoxia/ischemia injury

(Back et al., 2002; Back et al., 2007).

Molecular mechanisms underlying HIFα-regulated OPC differentiation

Previous study employed cerebellar slice/cell culture as a studying model and proposed

that the autocrine activation of Wnt/β-catenin signaling in OPCs by Wnt7a/Wnt7b plays a crucial

role in HIFα activation-induced hypomyelination (Yuen et al., 2014). Our data derived from *in*

vivo genetic models do not support this concept. Firstly, unbiased RNA-seq analysis did not

reveal dysregulation of the Wnt/β-catenin signaling activity and Wnt7a/Wnt7b expression.

Secondly, neither the activity of Wnt/β-catenin signaling nor the expression of Wnt7a/Wnt7b is

perturbed in HIFα-stabilized OPCs. Thirdly, disrupting the Wnt secretion mediator WLS blocks

OPC-derived autocrine Wnt/β-catenin signaling, but it does not affect the degree of inhibited

OPC differentiation elicited by HIFα hyperactivation. Our findings indicate that Wnt7a/Wnt7b or

Wnt/β-catenin signaling is unlikely a direct target of HIFα in the oligodendroglial lineage cells

(**Fig. 12N**), which is supported by a recently submitted study from an independent group (Allan

et al., 2020). The reasons for the different observations between our current study and prior

study (Yuen et al., 2014) are unclear. It is possible that non-cellular-specificity and/or off-target

of pharmacological compounds may underlie the discrepancy. HIFα stabilization induced by

hypoxia treatment or by pharmacological DMOG application occurs not only in oligodendroglial

lineage cells but also in other lineages of neural cells or vascular cells. Similarly, Wnt/β-catenin

inhibition in cultured slices by pharmacological XAV939 or IWP2 treatment may also happen in

other lineage cells (Yuen et al., 2014). We used cell-specific *Cre-loxP* genetic approaches to

circumvent the potential caveats of pharmacological application and convincingly demonstrated

that OPC-derived autocrine Wnt/β-catenin plays a minor, if any, role in mediating HIFα-

regulated OPC differentiation and myelination.

Another novel finding of this study is the identification of Sox9 as a potential target that

mediates HIFα stabilization-elicited inhibition of OPC differentiation (**Fig. 12N,** right panel). Sox9

is highly expressed in neural stem cells (NSCs) (Scott et al., 2010), downregulated in OPCs

(Stolt et al., 2003), and completely absent from differentiated OLs (Sun et al., 2017). Sox9

disruption affects OPC specification but not OPC terminal differentiation (Finzsch et al., 2008;

Stolt et al., 2003). HIFα stabilization activated Sox9 expression in OPCs both *in vitro* and *in vivo*,

which is in agreement with previous data showing that Sox9 is transcriptionally activated by

HIFα in chondrocytes (Amarilio et al., 2007) and pancreatic β-cell precursor cells (Puri et al.,

2013). We found that elevated Sox9 expression perturbed OPC differentiation and that Sox9

knockdown rescued the extent of inhibited OPC differentiation elicited by pharmacological HIFα

stabilization in primary OPC culture system.

It should be pointed out that the non-regulation of HIFα-Wnt axis does not negate the

crucial role of Wnt/β-catenin in oligodendroglial lineage cells. Previous studies have revealed an

inhibitory effect of Wnt/β-catenin activation on OPC differentiation (Guo et al., 2015). However,

these studies modulated the signaling activity through manipulating the intracellular components

or relevant regulatory molecules within OPCs , for instance, by disrupting intracellular β-catenin

(Fancy et al., 2009; Feigenson et al., 2009; Ye et al., 2009), Axin2 (Fancy et al., 2011), APC

(Fancy et al., 2014; Lang et al., 2013), APCDD1 (Lee et al., 2015b), and Daam2 (Lee et al.,

2015a). The Wnt-producing cells, which play a crucial role in activating the intracellular Wnt/β-

catenin signaling axis in OPCs remain unknown (Guo et al., 2015). Our WLS cKO data

demonstrate that oligodendroglial lineage-derived autocrine Wnt plays a minor role in OPC

differentiation and myelination in the early postnatal CNS under physiological (**Fig. 10**) or HIFα-

stabilized (**Figs. 8, 9**) conditions, suggesting that Wnt ligands derived from other lineage cells

may be responsible for control OPC differentiation through paracrine Wnt signaling. Previous

RNA-seq data (Zhang et al., 2014) indicate that astrocytes may be the major cellular source of

Wnt proteins, for example Wnt7a and Wnt5a. Future studies are needed to test the tempting

hypothesis that astrocyte-derived paracrine Wnt signaling may play major a role in controlling

OPC differentiation under normal or HIFα-stabilized conditions.

Implications of our findings in hypoxia-ischemia-induced oligodendroglial pathology in

preterm white matter injury (WMI)

Disturbance of normal developmental myelination is one of the established pathological

hallmarks in preterm (or premature) infants affected by diffuse WMI (Back, 2017; Deng, 2010).

Hypoxia/ischemia-induced diffuse WMI frequently occurs in the brain of preterm infants because

of the anatomical and physiological immaturity of the respiratory system and the brain white

matter vasculature. We demonstrated that, compared with healthy controls, HIF1α was

markedly elevated in the brain white matter of mice subjected to hypoxia-ischemia injury at the

time equivalent to developmental myelination in human brains (**Fig. 1F-I**). Our findings -

sustained HIFα activation disturbs OPC differentiation and results in hypomyelination - suggest

that oligodendroglial HIFα upregulation may play a crucial role in hypoxia-ischemia-induced

myelination disturbance. The function and molecular mechanisms of oligodendroglial HIFα in

hypoxia-ischemia-induced hypomyelination (Liu et al., 2011) have yet to be determined in

genetic animal models. Our *in vivo* data indicate that sustained HIFα activation in response to

hypoxia-ischemia injury may inhibit OPC differentiation and cause hypomyelination and that the

underlying mechanisms may be independent of OPC autocrine canonical Wnt signaling as

previously proposed. Future studies using cell-specific genetic HIFα models (HIFα loss and

gain-of-function) and pre-term equivalent mouse models of diffuse white matter injury are

needed to address these important questions.

Figures and Legends

Figure 1 – HIFα stabilization in the early postnatal murine CNS. **A-B**, immunohistochemistry (IHC) of HIF1α and oligodendroglial lineage marker Sox10 in the spinal cord ventral white matter (Spc-WM, marked by dotted lines) of HIFα intact mice (**A**) and HIFα stabilized (*Sox10-Cre:Vhl*[fl/fl]) mice (**B**) at postnatal day 5 (P5). **C-D**, IHC of HIF1α and Sox10 in the forebrain periventricular white matter (Fbr-PWM, marked by dotted lines) of HIFα intact mice (**C**) and HIFα stabilized (*Sox10-Cre:Vhl*[fl/fl]) mice (**D**) at P5. **E-F**, IHC of HIF1α and Sox10 in the Fbr-PWM of P10 mice that had been subjected to sham operation (**E**) and hypoxia/ischemia (H/I) injury on P6, i.e. 4 days post-H/I or sham (see Materials and Methods for details). Arrowheads in **A-F** point to HIF1α⁺Sox10⁺ cells. **G**, Western blot of micro-dissected subcortical white matter for HIF1α and canonical HIFα target PKM2 at 6 hours post-H/I or sham. GAPDH, internal loading

control. **H**, PKM2 IHC in the Fbr-PWM at P10, 4 days post-H/I or sham. **I1-I3**, IHC and

quantification of myelin basic protein (MBP) staining in the Fbr-PWM at P10, 4 days post-H/I or

sham. MBP intensity was normalized to that of the contralateral brain hemispheres to the

occluded artery of the same mouse. Scale bars, 20 µm.

Figure 2 – HIFα activation impairs developmental myelination by inhibiting OPC

differentiation but not subsequent OL maturation. A, Myelin staining of MBP in the spinal

cord of *Sox10-Cre:Vhl*$^{fl/fl}$ (Sox10:VHL cKO) and littermate control (*Vhl*$^{fl/+}$ or *Vhl*$^{fl/fl}$) pups at P5.

Most Sox10:VHL cKO pups died by the first postnatal week. **B**, representative confocal images

of Sox10 (red) and differentiated OL marker CC1 (green) in P5 spinal cord (n=5 Ctrl, 3 cKO).

Blue, DAPI nuclear counterstaining. **C**, quantification of Sox10$^+$CC1$^+$ OLs and Sox10$^+$PDGFRa$^+$

OPCs (right) in P5 spinal cord (n=5 Ctrl, 3 cKO). **D**, RT-qPCR assay for myelin-specific genes in

the spinal cord at P5. *Plp*, proteolipid protein; *Cnp*, 2',3'-Cyclic-nucleotide 3'-phosphodiesterase;

Mobp, myelin oligodendrocyte basic protein; *Mog*, myelin oligodendrocyte glycoprotein. **E-F**,

mRNA in situ hybridization (ISH) of *Plp* and *Mbp* in the spinal cord (**E**) and forebrain (**F**) of *Cnp-*

Cre:Vhl$^{fl/fl}$ (Cnp:VHL cKO) and littermate control (*Cnp-Cre:Vhl*$^{fl/+}$, *Vhl*$^{fl/+}$, or *Vhl*$^{fl/fl}$) mice at P14. **G**,

myelination in the adult spinal cord indicated by toluidine blue staining of semithin sections (left)

and transmission electron microscopy (TEM) of ultrathin section (right). Red asterisks in **G**

denotes blood vessels, the density of which is elevated in Cnp:VHL cKO mice (Guo et al., 2015).

H, *Mbp* mRNA levels quantified by RT-qPCR in Cnp:VHL cKO and Ctrl mice at different time

points. Spc, spinal cord; Fbr, forebrain. **I**, densities of Sox10$^+$CC1$^+$ OLs and Sox10$^+$PDGFRα$^+$

OPCs in P8 spinal cord of Cnp:VHL cKO and Ctrl mice. Scale bars: A, 100μm, B, 20μm, E-F,

200μm, G, 10μm (left), 1μm (right). Data are presented as mean ± s.e.m.; gray bars are from

Ctrl, white bars cKO; two-tailed Student's *t* test; * *p*<0.05, ** *p*<0.01, *** *p*<0.001, ns, not

significant. The above data presentation and statistics are applied to all figures except indicated

otherwise).

Figure 3 – Inducible HIFα stabilization in postnatal OPCs inhibits OPC differentiation. A,
experimental design for B-**F**. PDGFRα:VHL cKO (*Pdgfra-CreER*[T2]:*Vhl*[fl/fl]) and littermate control
(*Vhl*[fl/fl]) mice were treated with tamoxifen intraperitoneally at P1, P2, and P3, and analyzed at P8.
B, RT-qPCR quantification of HIFα target gene hexokinase 2 *Hk2* in the spinal cord. **C**,
expression of major myelin proteins in the spinal cord quantified by RT-qPCR. **D**, representative
confocal images and the density of Sox10[+]/CC1[+] OLs and Sox10[+]/PDGFRα[+] OPCs. **E**, IHC of
PLP showing myelination in the spinal cord ventral white matter. **F-G**, representative TEM
images (**F**) and the density of myelinated axons (**G**) in the spinal cord ventral white mater. **H**,
experimental design for **I-K**. PDGFRα:VHL cKO and littermate control mice were treated with
tamoxifen intraperitoneally at P1, P2, and P3, and analyzed at P60-70. **I**, representative
confocal images showing HIF1α stabilization in PDGFRα[+] OPCs (arrowheads) of PDGFRα:VHL
cKO mice. **J**, densities of CC1[+] mature OLs and PDGFRα[+] OPCs in the corpus callosum. K1-K4,
toluidine blue staining for myelin on semithin (500nm) resin sections showing hypomyelination in
the optic nerve of adult PDGFRα:VHL cKO mice. Boxed areas in **K1** and **K3** were shown at
higher magnification images in **K2** and **K4**, respectively. **K5**, density of myelinated axons in the
adult optic nerves. Scale bars, D, 25μm; E, 100μm; F, 0.5 μm; I-K, 10 μm.

Figure 4 –HIFα inactivation transiently delays developmental myelination by controlling OPC differentiation but not subsequent OL maturation. A-E, PDGFRα:HIFα cKO and littermate control (Hif1α$^{fl/fl}$:Hif2α$^{fl/fl}$) mice were treated with tamoxifen at P1, P2, and P3, and analyzed at P8 (n=4 each group) and P14 (n=3 each group). **A-B**, IHC (**A**) and quantification (**B**) of Olig2$^+$CC1$^+$ OLs and Olig2$^+$PDGFRα$^+$ OPCs in the spinal cord. **C**, relative expression of myelin protein genes of *Mbp* and *Plp* and mature OL-specific gene of *Opalin* and quake (*Qk,* aka CC1)* in the forebrain measured by real-time quantitative PCR (RT-qPCR). **D**, IHC of myelin marker MBP and pan-axonal marker SMI312 in forebrain periventricular white matter (left, marked by dotted area) and quantification (right) at P8. EC, external capsule, IC, internal capsule, Ctx, cortex. **E**, IHC of MBP in forebrain periventricular white matter at P14 (tamoxifen at P1, P2, and P3). **F-J**, Cnp:HIFα cKO and littermate control (Hif1α$^{fl/fl}$:Hif2α$^{fl/fl}$) mice were analyzed at P2, P8, and P14. **F**, RT-qPCR quantification of HIFα target genes *Ldha* and *Vegfa*

in P8 spinal cord. **G**, myelin staining by MBP in the forebrain. CC, corpus callosum, Ctx, cortex,

Hip, hippocampus. **H-I**, densities (#/mm^2) of Olig2$^+$CC1$^+$ OLs (**H**) and Olig2$^+$PDGFRα$^+$ OPCs (**I**)

at P14. Spc-WM: spinal cord white matter, Fbr-CC: forebrain corpus callosum. **J**, RT-qPCR

assay of exon3b-containing Plp mRNA, which is specific to mature OLs, in the spinal cord at

different time points. **K,** IHC of CC1 and HIF1α in P8 forebrain periventricular white matter (Fbr-

PWM). Arrowheads point to double positive cells. LV, lateral ventricle. Scale bar, A, K, 10um; D,

20um; E, 100um; G, 250um.

Figure 5 – Oligodendroglial HIFα is dispensable for axonal integrity or cell death. **A**, immunostaining images of P14 forebrain using SMI32, a monoclonal antibody recognizing the non-phosphorylated neurofilament proteins of heavy chain (NFH), which has been reported labeling injured axons and a subset of pyramidal neuron bodies and dendrites. Boxed areas show the corpus callosum (CC) at higher magnification images of SMI32 and MBP in the right

panels. Ctx, cortex, Hip, hippocampus. Note the absence of SMI32-positive signals in the CC of

both Cnp:HIFα cKO and Ctrl mice. **B**, immunostaining images of P14 CC using SMI312, a

monoclonal antibody recognizing both phosphorylated and non-phosphorylated NFH (a pan-

axonal marker). **C-D**, IHC of SMI32/MBP (**C**) and SMI312 (**D**) in the CC of P8 non-Cre Ctrl and

PDGFRα:HIFα cKO and littermate control (Hif1α$^{fl/fl}$:Hif2α$^{fl/fl}$) mice (tamoxifen injection at P1, P2,

and P3). **E-F**, representative confocal images and quantification of cells positive for cleaved

caspase 3 (CC3) and/or pan-oligodendroglial lineage marker Sox10 in P8 CC of PDGFRα:HIFα

cKO and control mice. Arrowheads point to CC3$^+$/Sox10$^+$ cells. **G**, quantification of CC3$^+$ and

CC3$^+$/Sox10$^+$ cells in P14 CC of Cnp:HIFα cKO and control mice. **H-I**, representative TEM

images showing the cross-sections of axons in the corpus callosum of HIFα Ctrl (**I**) and cKO (**H**)

mice at P8 (tamoxifen injections at P1, P2, and P3) when myelination is barely detectable at the

TEM level. Note that the morphology of axons (Ax) and axonal mitochondria (Mt) were

indistinguishable between HIFα Ctrl and cKO mice. Scale bar, A, 100μm (low magnification),

20μm (high magnification); B-E, 20μm, H-I, 400nm.

Figure 6 – Stabilizing HIFα activates HIFα-mediated signaling but does not perturb Wnt/β-catenin signaling in the oligodendroglial lineage and in the CNS. A-B, relative expression of HIFα and Wnt target genes from unbiased RNA-seq analysis. FPKM, fragments per kilobase of transcript per million mapped reads, a normalized measurement of gene expression. The RNA was prepared from the spinal cord of P5 *Sox10-Cre:Vhl*^fl/fl^ mice (n=3, white bars) and littermate control (*Vhl*^fl/+^ or *Vhl*^fl/fl^) (n=3, gray bars) mice. **C-E**, Top 3 significant enriched KEGG pathway and gene ontology (GO) terms among 255 upregulated and 351 downregulated genes derived from unbiased RNA-seq. Note that HIFα signaling pathway (**C**) and response to hypoxia (**D**) were significantly upregulated, validating HIFα hyperactivation in *Sox10-Cre:Vhl*^fl/fl^ mice and that oligodendroglial cell differentiation was significantly downregulated (**E**), which is agreement with the data of **Fig. 2A-D**. **F**, experimental design for **G** and **H**. PDGFRα:VHL cKO (n=3, white bars) and littermate control (*Vhl*^fl/fl^, n=3, gray bars) mice were treated with tamoxifen (TM) at P5,

and forebrain OPCs at P6 were acutely purified by MACS. **G-H**, RT-qPCR quantification of HIFα

target gene (**G**) and Wnt target gene (**H**) expression in purified OPCs. **I**, expression of Wnt

target gene (*Axin2* and *Naked1*) and *Wnt7a/7b* in P8 spinal cord (n=3 Ctrl, gray bars, 5

PDGFRα cKO, white bars) quantified by RT-qPCR. **J**, Western blot images and relative

expression (normalized to the internal loading control β-actin) of the active form of β-catenin

(ABC) and Axin2 in P14 forebrain (n=3 each group). **K**, expression of Wnt target genes (*Axin2*

and *Naked1*) and *Wnt7a/7b* in P8 spinal cord measured by RT-qPCR (n=4 each group).

Figure 7- WLS deficiency inhibits Wnt3a secretion and blocks autocrine Wnt/β-catenin

signaling activity in OPCs. A, experimental design of cell transfection of Wls shRNA and

Wnt3a-expressing plasmids to primary brain OPC maintained in the growth medium. **B-C**, RT-

qPCR assay of *Wls* (**B**) and *Wnt3a* (**C**) expression in transfected primary OPCs. One-way

ANOVA followed by Tukey's multiple comparisons, $F_{(3, 8)}$ = 15.39, P = 0.001 *Wls*, $F_{(3, 8)}$ = 32.81,

P < 0.0001 *Wnt3a*. **D**, ELISA quantification of Wnt3a in the growth medium in the absence or

presence of OPCs transfected by Wls shRNA and/or Wnt3a. One-way ANOVA followed by

Tukey's multiple comparisons, *** P < 0.001, ns not significant. $F_{(4, 10)}$ = 6.70, P < 0.0069. Note

that Wnt3a concentration in the GM in the presence of primary OPCs is statistically

indistinguishable from that in the GM alone, suggesting that intact primary OPCs do not secrete

Wnt3a. **E-F**, RT-qPCR quantification of Wnt target genes *Axin2* and *Sp5* in OPCs transfected

with Wls shRNA and/or Wnt3a. One-way ANOVA followed by Tukey's multiple comparisons, $F_{(3, 8)}$

= 11.06, P = 0.0032 *Axin2*, $F_{(3, 8)}$ = 13.25, P = 0.0018 *Sp5*.

Figure 8 – Disrupting WLS in PDGFRα-expressing OPCs does not affect the degree of inhibited OPC differentiation and hypomyelination elicited by HIFα hyperactivation. A, experimental design for **B-F**. PDGFRα:VHL cKO, *Pdgfrα-CreER*[T2]:*Vhl*[fl/fl] (n=5); PDGFRα:VHL/WLS cKO, *Pdgfrα-CreER*[T2]:*Vhl*[fl/fl]:Wls[fl/fl] (n=4); non-Cre Ctrl carrying *Vhl*[fl/fl] and/or Wls[fl/fl] (n=8). **B-C**, representative confocal images (**B**) and quantification (**C**) of myelination by MBP staining. The signal of SMI312, pan-axonal marker is indistinguishable among each group and used as an internal control of MBP quantification. One-way ANOVA followed by Tukey's multiple comparisons, $F_{(2, 14)}$=31.93 $P < 0.0001$. **D-E**, representative confocal images (**D**) and quantification (**E**) of Sox10[+]/CC1[+] differentiated OLs (**D**, arrowheads), Sox10[+]/PDGFRα[+] OPCs (**D**, arrows), and Sox10[+] oligodendroglial lineage cells in the spinal cord. One-way ANOVA followed by Tukey's multiple comparisons, $F_{(2, 14)} = 17.34$ $P = 0.0002$ Sox10[+], $F_{(2, 14)} = 18.44$ $P = 0.0001$ Sox10[+]/CC1[+], $F_{(2, 14)} = 2.967$ $P = 0.0843$ Sox10[+]/PDGFRα[+]. **F**, RT-qPCR quantification of mature OL-enriched genes in the spinal cord. One-way ANOVA followed by Tukey's multiple

comparisons, $F_{(2, 14)} = 11.53$ $P = 0.0011$ Mbp, $F_{(2, 14)} = 39.88$ $P < 0.0001$ Mag, $F_{(2, 14)} = 38.78$ $P <$

0.0001 Cnp. Scale bars=20μm in **B** and **D**.

Figure 9 - Disrupting WLS in Sox10-expressing oligodendroglial lineage cells does not

affect HIFα hyperactivation-elicited inhibition of OPC differentiation and hypomyelination.

A, experimental design for **B-E**. tamoxifen inducible $Sox10\text{-}CreER^{T2}$:$Vhl^{fl/fl}$ (iSox10:VHL cKO

n=3); $Sox10\text{-}CreER^{T2}$:$Vhl^{fl/fl}$:Wls$^{fl/fl}$ (iSox10:VHL/WLS cKO, n=3); non-Cre Ctrl carrying $Vhl^{fl/fl}$

and/or Wls$^{fl/fl}$ (n=5). **B-C**, representative confocal images (**B**) and quantification (**C**) of

myelination by MBP staining of the spinal cord. SMI312 signal was used as an internal control of

MBP quantification. One-way ANOVA followed by Tukey's multiple comparisons, $F_{(2, 8)}= 8.484$ P

$= 0.0105$. **D-E**, representative confocal images (**D**) and quantification (**E**) of Sox10$^+$/CC1$^+$

differentiated OLs (**D**, arrowheads) and Sox10$^+$/PDGFRα$^+$ OPCs (**D**, arrows) in the spinal cord.

One-way ANOVA followed by Tukey's multiple comparisons, $F_{(2, 8)} = 11.90$ $P = 0.004$

Sox10$^+$/CC1$^+$, $F_{(2, 14)} = 2.880$ $P = 0.1143$ Sox10$^+$/PDGFRα$^+$. Scale bars=20μm in **B** and **D**.

Figure 10 – Disrupting WLS in Sox10-expressing oligodendroglial lineage cells does not perturb normal OPC differentiation and myelination. A, experimental design for **B-G**. Tamoxifen inducible *Sox10-CreER*[T2]:*Wls*[fl/fl] (iSox10:WLS cKO, n=4); littermate non-Cre control Wls[fl/fl] mice (n=3). **B-C**, RT-qPCR quantification of Wls (**B**) and differentiated OL-enriched gene Cnp and Ugt8a (**C**) in the spinal cord. **D-E**, representative confocal images (**D**) and quantification (**E**) of Sox10[+]/CC1[+] differentiated OLs (**D**, arrowheads), Sox10[+]/PDGFRα[+] OPCs (**D**, arrows), and Sox10[+] oligodendroglial lineage cells in the spinal cord. **F-G**, representative confocal images (**B**) and quantification (**C**) of myelination by MBP staining of the spinal cord. SMI312 signal was used as an internal control of MBP quantification. Scale bars, D, 20µm, E, 50µm.

Figure 11 – HIFα stabilization activates Sox9 expression. A, Sox9 expression in the purified primary OPCs assessed by RT-qPCR. Purified cortical OPCs from neonatal rat brains were treated with HIFα stabilizer dimethyloxalylglycine (DMOG, 1mM) in the absence and presence of HIFα signaling blocker chetomin (100nM) for 7 h. n=4 each group. One-way ANOVA followed by Tukey's multiple comparisons, $F_{(2, 9)} = 62.53$ $P < 0.0001$. **B**, potential HIFα binding sites (CGTG) in the upstream sequence of rat Sox9 genes (+1bp is defined as the transcription start site, TSS) and ChIP-qPCR verification of physical binding of HIFα to the promoter region of -828bp in primary OPCs treated with DMOG. One way ANOVA with Dunnett post-hoc test comparing each group with negative control. $F_{(2,12)} = 69.21$ $P < 0.0001$. **C**, RT-qPCR quantification of Sox9 expression in P8 spinal cord of PDGFRα:VHL cKO (n=5) and littermate control(*Vhl*^fl/fl, n=3) mice that had been treated with tamoxifen at P1, P2, and P3. **D**, RT-qPCR quantification of Sox9 expression in P14 spinal cord of PDGFRα:VHL cKO (n=3) and littermate

control(*Vhl*^fl/fl^, n=3) that had received tamoxifen at P6 and P7. **E**, confocal images showing that

Sox9 is barely expressed in Sox10$^+$ oligodendroglial lineage cells in the control spinal cord but

upregulated in Sox10:VHL cKO mice at P5. WM, white matter, GM, gray matter. The boxed

areas of #1 and #2 were shown at higher magnification at the right. **F**, percentage of Sox10$^+$

oligodendroglial lineage cells expressing Sox9 in the spinal cord (n=3 each group). **G**, RT-qPCR

quantification of Sox9 mRNA in the spinal cord of Sox10:VHL cKO (n=5) and littermate control

(*Vhl*^fl/+^ and/or *Vhl*^fl/fl^, n=8) mice at P5. **H**, RT-qPCR assay of Sox9 expression in primary OPCs

isolated from the neonatal brain of *Sox10-Cre:Vhl*^fl/fl^ (n=4) and littermate control (n=6) pups. **I**,

representative confocal images and quantification of Sox9+GFAP+ astrocytes in the ventral

spinal cord of P5 *Sox10-Cre:Vhl*^fl/fl^ (n=3) and littermate control (n=6) mice. J1-J3, representative

confocal images and quantification of Sox10 and Sox9 in the ventral spinal cord of P8

PDGFRα:HIFα cKO (n=4) and littermate control (n=4) mice (tamoxifen at P1, P2, and P3).

Arrowheads point to Sox10$^+$Sox9$^+$ cells. Scale bars, 50µm in E, 10 µm in I, 10 µm in boxed

areas of panel E. 20 µm in J1-J2.

Figure 12 – Sustained Sox9 expression inhibits OPC differentiation. **A**, experimental design for **B-G**. Cortical primary OPCs in growth medium were transfected with Sox9-expressing plasmid (Sox9, n=3) or empty vector (EV, n=3), then differentiate for 24 h in serum-free differentiating medium. **B**, RT-qPCR quantification of Sox9 mRNA expression. **C,** representative immunocytochemical images of Sox9/Sox10. Arrowheads point to Sox9$^+$/Sox10$^+$ oligodendrocytes and arrows point to Sox9$^+$/Sox10$^-$ cells, which are presumably astrocytes.

Scale bar=20μm. **D,** percentage of Sox10[+] oligodendroglial lineage cells that express Sox9 (top)

and Sox9 immunoreactive intensity normalized to Sox10 (bottom). **E**, RT-qPCR assay of

exon3b-containing *Plp* (*Plp-E3b*), which specifically expresses in differentiated OLs, and myelin

regulatory factor (*Myrf*), a potent pro-differentiation gene. **F**, representative immunocytochemical

images of MBP/Sox10 and density (#/mm^2) of Sox10[+] oligodendroglial cells of complex ramified

morphology indicated by MBP staining (arrowheads in **F**), Scale bar=10μm. **G**, Western blot

images and quantification of MBP and internal loading control β-actin. **H**, experimental designs

for panels **I-M**, primary OPCs were treated with HIFα stabilizer DMOG and transfected with

Sox9 siRNA twice with 24 h apart. **I-J**, expression of *Sox9* (**I**) and HIFα target gene *Bnip3* (**J**)

quantified by RT-qPCR. One way ANOVA followed by Tukey's test; $F_{(3, 8)}$ = 10.33, P = 0.004

Sox9; $F_{(3, 8)}$ = 20.83, P = 0.0004 *Bnip3.* **K**, representative immunostaining images of

differentiated OLs labeled by MBP and Sox10. Scale bar=10μm. **L-M**, density (#/mm^2) of

MBP[+]/Sox10[+] differentiated OLs (**L**) and RT-qPCR assay of *Mbp* mRNA expression (**M**). One

way ANOVA followed by Tukey's test; $F_{(3, 8)}$ = 33.29, P < 0.0001 MBP[+]/Sox10[+]*;* $F_{(3, 8)}$ = 97.96, P

< 0.0001 *Mbp.* **N**, previously proposed and our alternative working model explaining the role of

HIFα in OPC differentiation. Endogenous HIFα regulates CNS myelination by controlling

upstream OPC differentiation but not downstream OL maturation in perinatal and early postnatal

CNS whereas aberrant HIFα activation (via VHL mutation of hypoxia-ischemia insult) inhibits

OPC differentiation through the Wnt-independent and Sox9-dependent pathway.

Reference

Allan, K., L. Hu, A. Morton, M. Scavuzzo, A. Gevorgyan, C. B., B. I., S. Hung, C. Bartels, M. Madhavan, and

P.J. Tesar. 2020. Non-Canonical Targets of HIF1a Drive Cell-Type-Specific Dysfunction. *BioRxiv*

[Preprint].

Amarilio, R., S.V. Viukov, A. Sharir, I. Eshkar-Oren, R.S. Johnson, and E. Zelzer. 2007. HIF1alpha regulation

of Sox9 is necessary to maintain differentiation of hypoxic prechondrogenic cells during early

skeletogenesis. *Development*. 134:3917-3928.

Back, S.A. 2017. White matter injury in the preterm infant: pathology and mechanisms. *Acta*

Neuropathol. 134:331-349.

Back, S.A., B.H. Han, N.L. Luo, C.A. Chricton, S. Xanthoudakis, J. Tam, K.L. Arvin, and D.M. Holtzman. 2002.

Selective vulnerability of late oligodendrocyte progenitors to hypoxia-ischemia. *J Neurosci*.

22:455-463.

Back, S.A., A. Riddle, and M.M. McClure. 2007. Maturation-dependent vulnerability of perinatal white
matter in premature birth. *Stroke*. 38:724-730.

Banziger, C., D. Soldini, C. Schutt, P. Zipperlen, G. Hausmann, and K. Basler. 2006. Wntless, a conserved
membrane protein dedicated to the secretion of Wnt proteins from signaling cells. *Cell*.
125:509-522.

Bartscherer, K., N. Pelte, D. Ingelfinger, and M. Boutros. 2006. Secretion of Wnt ligands requires Evi, a
conserved transmembrane protein. *Cell*. 125:523-533.

Bercury, K.K., and W.B. Macklin. 2015. Dynamics and mechanisms of CNS myelination. *Dev Cell*. 32:447-
458.

Campbell, M.J., and J.H. Morrison. 1989. Monoclonal antibody to neurofilament protein (SMI-32) labels
a subpopulation of pyramidal neurons in the human and monkey neocortex. *J Comp Neurol*.
282:191-205.

Chung, S.H., F. Guo, P. Jiang, D.E. Pleasure, and W. Deng. 2013. Olig2/Plp-positive progenitor cells give
rise to Bergmann glia in the cerebellum. *Cell Death Dis*. 4:e546.

Cunningham, L.A., K. Candelario, and L. Li. 2012. Roles for HIF-1alpha in neural stem cell function and
the regenerative response to stroke. *Behav Brain Res*. 227:410-417.

Deng, W. 2010. Neurobiology of injury to the developing brain. *Nat Rev Neurol*. 6:328-336.

Emery, B., D. Agalliu, J.D. Cahoy, T.A. Watkins, J.C. Dugas, S.B. Mulinyawe, A. Ibrahim, K.L. Ligon, D.H.
Rowitch, and B.A. Barres. 2009. Myelin gene regulatory factor is a critical transcriptional
regulator required for CNS myelination. *Cell*. 138:172-185.

Fancy, S.P., S.E. Baranzini, C. Zhao, D.I. Yuk, K.A. Irvine, S. Kaing, N. Sanai, R.J. Franklin, and D.H. Rowitch.
2009. Dysregulation of the Wnt pathway inhibits timely myelination and remyelination in the
mammalian CNS. *Genes Dev*. 23:1571-1585.

Fancy, S.P., E.P. Harrington, S.E. Baranzini, J.C. Silbereis, L.R. Shiow, T.J. Yuen, E.J. Huang, S. Lomvardas,
and D.H. Rowitch. 2014. Parallel states of pathological Wnt signaling in neonatal brain injury and
colon cancer. *Nat Neurosci*.

Fancy, S.P., E.P. Harrington, T.J. Yuen, J.C. Silbereis, C. Zhao, S.E. Baranzini, C.C. Bruce, J.J. Otero, E.J.
Huang, R. Nusse, R.J. Franklin, and D.H. Rowitch. 2011. Axin2 as regulatory and therapeutic
target in newborn brain injury and remyelination. *Nat Neurosci*. 14:1009-1016.

Feigenson, K., M. Reid, J. See, E.B. Crenshaw, 3rd, and J.B. Grinspan. 2009. Wnt signaling is sufficient to
perturb oligodendrocyte maturation. *Mol Cell Neurosci*. 42:255-265.

Finzsch, M., C.C. Stolt, P. Lommes, and M. Wegner. 2008. Sox9 and Sox10 influence survival and

migration of oligodendrocyte precursors in the spinal cord by regulating PDGF receptor alpha

expression. *Development*. 135:637-646.

Guo, F., J. Lang, J. Sohn, E. Hammond, M. Chang, and D. Pleasure. 2015. Canonical Wnt signaling in the

oligodendroglial lineage-puzzles remain. *Glia*.

Harb, R., C. Whiteus, C. Freitas, and J. Grutzendler. 2013. In vivo imaging of cerebral microvascular

plasticity from birth to death. *Journal of cerebral blood flow and metabolism : official journal of*

the International Society of Cerebral Blood Flow and Metabolism. 33:146-156.

Haynes, R.L., N.S. Borenstein, T.M. Desilva, R.D. Folkerth, L.G. Liu, J.J. Volpe, and H.C. Kinney. 2005.

Axonal development in the cerebral white matter of the human fetus and infant. *J Comp Neurol*.

484:156-167.

Hines, J.H., A.M. Ravanelli, R. Schwindt, E.K. Scott, and B. Appel. 2015. Neuronal activity biases axon

selection for myelination in vivo. *Nat Neurosci*. 18:683-689.

Huang, H., X.F. Zhao, K. Zheng, and M. Qiu. 2013. Regulation of the timing of oligodendrocyte

differentiation: mechanisms and perspectives. *Neuroscience bulletin*. 29:155-164.

Ivanovic, Z. 2009. Hypoxia or in situ normoxia: The stem cell paradigm. *J Cell Physiol*. 219:271-275.

Jablonska, B., J. Scafidi, A. Aguirre, F. Vaccarino, V. Nguyen, E. Borok, T.L. Horvath, D.H. Rowitch, and V.

Gallo. 2012. Oligodendrocyte regeneration after neonatal hypoxia requires FoxO1-mediated

p27Kip1 expression. *J Neurosci*. 32:14775-14793.

Kung, A.L., S.D. Zabludoff, D.S. France, S.J. Freedman, E.A. Tanner, A. Vieira, S. Cornell-Kennon, J. Lee, B.

Wang, J. Wang, K. Memmert, H.U. Naegeli, F. Petersen, M.J. Eck, K.W. Bair, A.W. Wood, and D.M.

Livingston. 2004. Small molecule blockade of transcriptional coactivation of the hypoxia-

inducible factor pathway. *Cancer Cell*. 6:33-43.

Lang, J., Y. Maeda, P. Bannerman, J. Xu, M. Horiuchi, D. Pleasure, and F. Guo. 2013. Adenomatous

polyposis coli regulates oligodendroglial development. *The Journal of neuroscience : the official*

journal of the Society for Neuroscience. 33:3113-3130.

Lappe-Siefke, C., S. Goebbels, M. Gravel, E. Nicksch, J. Lee, P.E. Braun, I.R. Griffiths, and K.A. Nave. 2003.

Disruption of Cnp1 uncouples oligodendroglial functions in axonal support and myelination. *Nat*

Genet. 33:366-374.

Lee, H.K., L.S. Chaboub, W. Zhu, D. Zollinger, M.N. Rasband, S.P. Fancy, and B. Deneen. 2015a. Daam2-

PIP5K Is a Regulatory Pathway for Wnt Signaling and Therapeutic Target for Remyelination in the

CNS. *Neuron*. 85:1227-1243.

Lee, H.K., D. Laug, W. Zhu, J.M. Patel, K. Ung, B.R. Arenkiel, S.P. Fancy, C. Mohila, and B. Deneen. 2015b.
Apcdd1 stimulates oligodendrocyte differentiation after white matter injury. *Glia*. 63:1840-1849.

Lee, S., M.K. Leach, S.A. Redmond, S.Y. Chong, S.H. Mellon, S.J. Tuck, Z.Q. Feng, J.M. Corey, and J.R. Chan.
2012. A culture system to study oligodendrocyte myelination processes using engineered
nanofibers. *Nat Methods*. 9:917-922.

Li, L., K.M. Candelario, K. Thomas, R. Wang, K. Wright, A. Messier, and L.A. Cunningham. 2014. Hypoxia
inducible factor-1alpha (HIF-1alpha) is required for neural stem cell maintenance and vascular
stability in the adult mouse SVZ. *J Neurosci*. 34:16713-16719.

Liu, W., Y. Shen, J.M. Plane, D.E. Pleasure, and W. Deng. 2011. Neuroprotective potential of
erythropoietin and its derivative carbamylated erythropoietin in periventricular leukomalacia.
Exp Neurol. 230:227-239.

Liu, X.B., Y. Shen, J.M. Plane, and W. Deng. 2013. Vulnerability of premyelinating oligodendrocytes to
white-matter damage in neonatal brain injury. *Neurosci Bull*. 29:229-238.

Milosevic, J., M. Maisel, F. Wegner, J. Leuchtenberger, R.H. Wenger, M. Gerlach, A. Storch, and J.
Schwarz. 2007. Lack of hypoxia-inducible factor-1 alpha impairs midbrain neural precursor cells
involving vascular endothelial growth factor signaling. *J Neurosci*. 27:412-421.

Moyon, S., J.L. Huynh, D. Dutta, F. Zhang, D. Ma, S. Yoo, R. Lawrence, M. Wegner, G.R. John, B. Emery, C.
Lubetzki, R.J. Franklin, G. Fan, J. Zhu, J.L. Dupree, and P. Casaccia. 2016. Functional
Characterization of DNA Methylation in the Oligodendrocyte Lineage. *Cell Rep*.

Puri, S., H. Akiyama, and M. Hebrok. 2013. VHL-mediated disruption of Sox9 activity compromises beta-
cell identity and results in diabetes mellitus. *Genes Dev*. 27:2563-2575.

Reiprich, S., M. Cantone, M. Weider, T. Baroti, J. Wittstatt, C. Schmitt, M. Kuspert, J. Vera, and M.
Wegner. 2017. Transcription factor Sox10 regulates oligodendroglial Sox9 levels via microRNAs.
Glia. 65:1089-1102.

Scott, C.E., S.L. Wynn, A. Sesay, C. Cruz, M. Cheung, M.V. Gomez Gaviro, S. Booth, B. Gao, K.S. Cheah, R.
Lovell-Badge, and J. Briscoe. 2010. SOX9 induces and maintains neural stem cells. *Nat Neurosci*.
13:1181-1189.

Semenza, G.L. 2012. Hypoxia-inducible factors in physiology and medicine. *Cell*. 148:399-408.

Semple, B.D., K. Blomgren, K. Gimlin, D.M. Ferriero, and L.J. Noble-Haeusslein. 2013. Brain development
in rodents and humans: Identifying benchmarks of maturation and vulnerability to injury across
species. *Prog Neurobiol*. 106-107:1-16.

Sharp, F.R., and M. Bernaudin. 2004. HIF1 and oxygen sensing in the brain. *Nat Rev Neurosci*. 5:437-448.

Shen, Y., J.M. Plane, and W. Deng. 2010. Mouse models of periventricular leukomalacia. *J Vis Exp*.

Soulika, A.M., E. Lee, E. McCauley, L. Miers, P. Bannerman, and D. Pleasure. 2009. Initiation and
progression of axonopathy in experimental autoimmune encephalomyelitis. *J Neurosci*.
29:14965-14979.

Stolt, C.C., P. Lommes, E. Sock, M.C. Chaboissier, A. Schedl, and M. Wegner. 2003. The Sox9 transcription
factor determines glial fate choice in the developing spinal cord. *Genes Dev*. 17:1677-1689.

Sun, W., A. Cornwell, J. Li, S. Peng, M.J. Osorio, N. Aalling, S. Wang, A. Benraiss, N. Lou, S.A. Goldman,
and M. Nedergaard. 2017. SOX9 Is an Astrocyte-Specific Nuclear Marker in the Adult Brain
Outside the Neurogenic Regions. *J Neurosci*. 37:4493-4507.

Tomita, S., M. Ueno, M. Sakamoto, Y. Kitahama, M. Ueki, N. Maekawa, H. Sakamoto, M. Gassmann, R.
Kageyama, N. Ueda, F.J. Gonzalez, and Y. Takahama. 2003. Defective brain development in mice
lacking the Hif-1alpha gene in neural cells. *Mol Cell Biol*. 23:6739-6749.

Viziteu, E., C. Grandmougin, H. Goldschmidt, A. Seckinger, D. Hose, B. Klein, and J. Moreaux. 2016.
Chetomin, targeting HIF-1 alpha/p300 complex, exhibits antitumour activity in multiple
myeloma. *Brit J Cancer*. 114:519-523.

Voelker, C.C., N. Garin, J.S. Taylor, B.H. Gahwiler, J.P. Hornung, and Z. Molnar. 2004. Selective
neurofilament (SMI-32, FNP-7 and N200) expression in subpopulations of layer V pyramidal
neurons in vivo and in vitro. *Cereb Cortex*. 14:1276-1286.

Volpe, J.J. 2009. Brain injury in premature infants: a complex amalgam of destructive and developmental
disturbances. *Lancet Neurol*. 8:110-124.

Ye, F., Y. Chen, T. Hoang, R.L. Montgomery, X.H. Zhao, H. Bu, T. Hu, M.M. Taketo, J.H. van Es, H. Clevers,
J. Hsieh, R. Bassel-Duby, E.N. Olson, and Q.R. Lu. 2009. HDAC1 and HDAC2 regulate
oligodendrocyte differentiation by disrupting the beta-catenin-TCF interaction. *Nat Neurosci*.
12:829-838.

Yuen, T.J., J.C. Silbereis, A. Griveau, S.M. Chang, R. Daneman, S.P. Fancy, H. Zahed, E. Maltepe, and D.H.
Rowitch. 2014. Oligodendrocyte-Encoded HIF Function Couples Postnatal Myelination and
White Matter Angiogenesis. *Cell*. 158:383-396.

Zhang, K., L. Zhu, and M. Fan. 2011. Oxygen, a Key Factor Regulating Cell Behavior during Neurogenesis
and Cerebral Diseases. *Front Mol Neurosci*. 4:5.

Zhang, S., B. Kim, X. Zhu, X. Gui, Y. Wang, Z. Lan, P. Prabhu, K. Fond, A. Wang, and F. Guo. 2020. Glial
type specific regulation of CNS angiogenesis by HIFalpha-activated different signaling pathways.
Nat Commun. 11:2027.

Zhang, S., A. Rasai, Y. Wang, J. Xu, P. Bannerman, D. Erol, D. Tsegaye, A. Wang, A. Soulika, X. Zhan, and F.

Guo. 2018a. The Stem Cell Factor Sox2 Is a Positive Timer of Oligodendrocyte Development in

the Postnatal Murine Spinal Cord. *Mol Neurobiol.*

Zhang, S., X. Zhu, X. Gui, C. Croteau, L. Song, J. Xu, A. Wang, P. Bannerman, and F. Guo. 2018b. Sox2 is

essential for oligodendroglial proliferation and differentiation during postnatal brain

myelination and CNS remyelination. *J Neurosci.*

Zhang, Y., K. Chen, S.A. Sloan, M.L. Bennett, A.R. Scholze, S. O'Keeffe, H.P. Phatnani, P. Guarnieri, C.

Caneda, N. Ruderisch, S. Deng, S.A. Liddelow, C. Zhang, R. Daneman, T. Maniatis, B.A. Barres,

and J.Q. Wu. 2014. An RNA-sequencing transcriptome and splicing database of glia, neurons,

and vascular cells of the cerebral cortex. *The Journal of neuroscience : the official journal of the*

Society for Neuroscience. 34:11929-11947.

1014