# Antimicrobial peptide induced-stress renders *Staphylococcus aureus* susceptible to toxic nucleoside analogues

Alexandro Rodríguez-Rojas[1]*, Arpita Nath[1], Baydaa El Shazely[1,] Greta Santi[1], Joshua Jay Kim[1], Benno Kuropka[2], Christoph Weise[2] and Jens Rolff[1].

[1]Freie Universität Berlin, Institut für Biologie, Evolutionary Biology, Königin-Luise-Strasse 1-3, 14195 Berlin, Germany.

[2]Freie Universität Berlin, Institute of Chemistry and Biochemistry, für Chemie und Biochemie, Thielallee 63, 14195 Berlin, Germany.

*Corresponding author: a.rojas@fu-berlin.de

**-Abstract-**

Cationic antimicrobial peptides (AMPs) are active immune effectors of multicellular organisms and also considered as new antimicrobial drug candidates. One of the problems encountered when developing AMPs as drugs is the difficulty to reach sufficient killing concentrations under physiological conditions. Here, using pexiganan, a cationic peptide derived from a host defence peptide of the African clawed frog and the first AMP developed into a drug, we studied if sub-lethal effects of AMPs can be harnessed to devise treatment combinations. We studied the pexiganan stress response of *Staphylococcus aureus* at sub-lethal concentrations using a whole proteomics approach by liquid chromatography coupled with mass spectrometry. Several proteins involved in nucleotide metabolism were elevated, suggesting a metabolic demand. We then show that *S. aureus* is highly susceptible to antimetabolite nucleoside analogues when exposed to pexiganan, even at sub-inhibitory concentrations. These findings could be used to enhance pexiganan potency while decreasing the risk of resistance emergence, and our findings can likely be extended to other antimicrobial peptides.

**Introduction.**

Antimicrobial peptides (AMPs, we use AMPs here as synonymous with host defence peptides) are immune effector molecules used by multicellular organisms to control infections (Nicolas and Mor, 1995; Zasloff, 2002). These peptides are usually active against a broad spectrum of bacterial pathogens and some display activity against antibiotic-resistant bacteria. Thus, antimicrobial peptides are considered a promising source of new antibacterial drugs (Hancock and Sahl, 2006; Czaplewski et al., 2016) to tackle the current antibiotic crisis (Baker, 2015).

Reasons that make AMPs attractive are their high diversity across the tree of life (Wang and Wang, 2004) and the finding that albeit drug resistance evolves against AMPs (Perron et al., 2006; Habets and Brockhurst, 2012; Lofton et al., 2013; Johnston et al., 2016; Makarova et al., 2018), it evolves at a much lower probability in comparison to conventional antibiotics (Yu et al., 2018; Spohn et al., 2019). One common problem with the development of AMPs as drugs is that, under physiological conditions, their antimicrobial activity cannot easily recaptured and the required dosage is extremely high (Mookherjee et al., 2020). This dosage issue can be addressed by making use of synergistic combinations of AMPs (Yu et al., 2016), a property common in natural defense cocktails (Westerhoff et al., 1989; Yan and Hancock, 2001).

While the mode of action of AMPs on bacterial membranes has been worked out for some (Zerweck et al 2015) the consequence of AMP induced stress on bacterial physiology is less studied. The first goal of this study therefore is to understand how the pathogen *Staphylococcus aureus* reacts to different doses of a pexiganan at the minimum inhibitory concentration (MIC). Pexiganan is a drug that was mostly developed against this bacterium (Ge et al., 1999). Pexiganan is a 22-amino-acid antimicrobial peptide, an analogue of the magainin peptides isolated from the skin of the African clawed frog. Pexiganan exhibited broad-spectrum antibacterial activity *in vitro* when tested against 3,109 clinical isolates of gram-positive and gram-negative, anaerobic and aerobic bacteria (Ge et al., 1999).

Using pexiganan as an example, we found that different concentrations induce the upregulation of several genes depending on nucleotides or related to nucleotide metabolism. Based on these results, we hypothesized that this leads a way to identify phenotypic collateral sensitivity. We hypothesised that the response to pexiganan sensitizes *S. aureus* against certain nucleoside antimetabolites or toxic nucleoside analogues. Interestingly, these analogues have been proposed as an alternative to antibiotics as a consequence of resistance emergence (Thomson and Lamont, 2019). Nucleoside analogues have the advantage of being clinically approved for cancer therapies, but also as antiviral and antifungal treatments (Thomson and Lamont, 2019). Pyrimidine and purine analogues, as we use here, show potent antimicrobial activity against *S. aureus* in the

past (Rogers and Perkins, 1960; Stickgold and Neuhaus, 1967; Jordheim et al., 2012a; Thomson and Lamont, 2019).

In this study, we show how the understanding of *S. aureus* response to low-dose pexiganan uncovers cellular soft spots and helps to identify intervention opportunities. Secondly, it could also contribute to our understanding of the early stages of resistance evolution to antimicrobial peptides. Here, we first study the global proteomic response of *S. aureus* to the cationic antimicrobial peptide pexiganan at concentrations similar to and below MIC to detect the possible metabolic changes that open the path to collateral sensitivity to nucleoside analogues. We then find that these treatments sensitize *S. aureus* to antimetabolite purine and pyrimidines analogues.

**Results.**

**Changes in protein profiles after pexiganan treatment.** We examined *S. aureus* exposed to pexiganan by studying proteome-wide changes after 30-minute treatment to different concentrations (0.125, 0.25, 0.5 and 1x MIC, Table S1). Overall, 1160 proteins were identified at 1 % or less false discovery rate (FDR) among which 968 proteins were quantified and used for downstream analysis. All identified proteins, their quantification and statistical tests are provided in supplementary Table S2. A global overview shows a proteome-wide perturbation induced by pexiganan stress for all concentrations. Hundreds of proteins were significantly differentially expressed (Figure 1, Figure S1). It is noticeable that as long as the dose increases, the level of expression (fold-change) of both overexpressed and suppressed genes, decreases, making the dot scattering of the volcano plot less disperse (Figure S1). This indicates a decrease in the ability of the cell to react with increasing peptide concentration.

Within the upregulated proteome fraction (Figure 1, Figure S1), a group of proteins related to osmotic stress response shows up. The multi-peptide resistance factor MprF, a protein associated with cationic peptide resistance, which is conserved among many bacterial species (Kristian et al., 2003; Weidenmaier et al., 2005). MprF catalyses the transfer of a lysyl group from L-lysyl-tRNA(Lys) to membrane-bound phosphatidylglycerol producing lysylphosphatidylglycerol, a major component of the bacterial membrane with a net positive charge, hence modification of anionic phosphatidylglycerol with positively charged L-lysine results in repulsion of the peptides. The changes on the membrane charge is *per se* resistance mechanism against cationic antimicrobial peptides. Thus, MprF increases resistance to moenomycin and vancomycin, resistance to human defensins (HNP1-3) and evasion of oxygen-independent neutrophil killing and other AMPs and antibiotics (Oku et al., 2004; Staubitz et al., 2004). Another highly expressed protein is CapFis, involved in the pathway capsule polysaccharide biosynthesis, a mucous layer on the surface of bacterium that facilitates immune evasion and infection. CapF is an important virulence factor during infections by *S. aureus*. The enzyme CapF is considered a therapeutic candidate to disrupt the capsule polysaccharide biosynthesis (Miyafusa et al., 2013). TagG upregulates the protein which is part of the wall teichoic acid synthesis during the final steps of the pathway. Wall teichoic acid are important in pathogenesis and play key roles in antimicrobial resistance (Weidenmaier et al., 2005; Brown et al., 2013). The chaperons/proteases ClpL and TreP are among the most fifty upregulated genes for the MIC dose. ClpL is an ATP-dependent Clp protease. Clp proteases play a central role in stress survival, virulence and antibiotic resistance of *S. aureus* (Frees et al., 2014).

In addition to the above virulence factors, pexiganan also induced the expression PepT, also known as Staphopain A. This enzyme is a cysteine protease that plays an important role in the inhibition of host innate immune response. It cleaves host elastins from connective tissues, pulmonary surfactant protein A in the lungs, and the chemokine receptor CXCR2 on leukocytes (Kantyka et al., 2013). Proteolytic cleavage of surfactant protein A impairs bacterial phagocytosis by neutrophils while CXCR2 degradation blocks neutrophil activation and chemotaxis (Potempa et al., 1988; Kantyka et al., 2013). Additionally, PepT promotes vascular leakage by activating the plasma kallikerin/kinin system, resulting in patient hypotension (Imamura et al., 2005). Another important virulence factor is coded by the Gene SAOUHSC_02980, a protein containing an isochorismatase domain. This enzyme participates in the biosynthesis of siderophore groups that in *S. aureus* are redundant systems and varies even across different vertebrates hosts (Perry et al., 2019).

NptA, a phosphate transporter, usually induced by phosphate limitation, is highly abundant. Inorganic phosphate acquisition via NptA is particularly important for the pathogenesis of *S. aureus*. NptA homologs are widely distributed among bacteria and closely related less pathogenic staphylococcal species do not possess this importer. Another phosphate metabolism related gene with high expression is SAOUHSC_00480, that codes for a putative nucleoside triphosphate pyrophosphohydrolase (Pundir et al., 2016). Another two proteins, CarA and CarB, participate in the L-arginine biosynthesis. They are involved in the first step of the sub-pathway that synthesizes carbamoyl phosphate from bicarbonate. The elevation of these enzymes could indicate that pexiganan stress may be involved in amino acid depletion. Also related with phosphate metabolism, we observed a high level of FruA in different pexiganan concentrations. This protein a phosphoenolpyruvate-dependent sugar phosphotransferase system (a PTS system) is a major carbohydrate active transport system, which catalyses the phosphorylation of incoming sugar substrates concomitantly with their translocation across the cell membrane and potentially important for survival in respiratory tract of the host (Garnett et al., 2014). GlcB, another PTS system is a phosphoenolpyruvate-dependent sugar phosphotransferase system. This protein is another major carbohydrate active -transport system and catalyses the phosphorylation of incoming sugar substrates and their translocation across the cell membrane (Vitko et al., 2016).

The gene SAOUHSC_00456 that codes for YabA is significantly increased as well. YabA is involved in initiation of chromosome replication, and is a negative controller of DNA replication initiation in *Bacillus subtillis*. YabA and DnaD inhibit helix assembly of the DNA replication initiation protein DnaA (Scholefield and Murray, 2013). The elevated concentration within the cell of YabA could stall the cell division while the bacteria is under severe stress. *S. aureus* upregulates Spermidine/putrescine import ATP-binding protein

PotA. This protein is part of the ABC transporter complex PotABCD and responsible for energy coupling to the transport system. Spermidine and putrescine are polyamines which role in *S. aureus* is not well defined (Di Martino et al., 2013). There are also a set of up-regulated proteins coded by the genes SAOUHSC_01717, SAOUHSC_02581 and SAOUHSC_02581 which function remains unknown as described in Uniprot database and showed no homology with any known sequence (Pundir et al., 2016).

One of the hallmarks of our proteomic dataset is that we found higher level of expression, compared to control, for proteins related with nucleotide metabolism (Figure 1), which is directly connected to the upregulation of phosphate metabolism proteins described above. GmK for example, an essential protein for recycling GMP and indirectly, cGMP Guanylate kinase is highly upregulated. GMK is an essential enzyme and a potential antimicrobial drug target owing to its role in supplying DNA and RNA precursors (Omari et al., 2006). Another nucleobase metabolism related protein having or exhibiting a higher expression for the 1x MIC treated cells is PyrG. This enzyme catalyses the ATP-dependent amination of UTP to CTP with either L-glutamine or ammonia as the source of nitrogen. It also regulates intracellular CTP levels through interactions with the four ribonucleotide triphosphates.

Pexiganan also impacted negatively the level of expression of many proteins proteome-wide (Figure S1, Table S2). Among the most affected gene expressions throughout all concentrations are genes such as *dps* (coding for a known iron storage protein), *hld*, *copZ*, *cspC*, *metQ*, *sceD*, *isaA csoB*/*C*, *dltC*, adsA and *sasG*, SAOUHSCA_01134 and SAOUHSCA_02576. The protein cspB codes for the downregulated protein CspD, a cold shock protein that accumulates during low temperature or cold shock. This gene is also a component of the stringent response, indicating that it could be a general stress response gene (Anderson et al., 2006). Other genes showing differentially low level of expression are SAOUHSC_01986, SAOUHSC_01986, SAOUHSC_008020, SAOUHSC_02093, SAOUHSC_02535 and SAOUHSC_01414 which code for uncharacterized proteins (Pundir et al., 2016). SAOUHSC_01030 is a putative glutaredoxin domain-containing protein but it is not characterized either. The gene SAOUHSC_02576 codes for a putative secretory antigen SsaA, identified in *S. epidermidis* but its function is also unknown (Pundir et al., 2016).

In contrast to the upregulation of peptidoglycan synthesis, we observe that putative peptidoglycan hydrolases and probable lytic transglycosylases IsaA and SceD were downregulated. Interestingly, the *isaA sceD* double mutant is attenuated for virulence, while SceD is essential for nasal colonization in cotton rats (Stapleton et al., 2007). The gene *moaD* shows also a reduced level of expression and it codes for a molybdopterin

converting factor subunit 1. Molybdopterins are a class of cofactors found in most molybdenum-containing and all tungsten-containing enzymes. Molybdopterin pathway reactions consume guanosine triphosphate that is converted into the cyclic phosphate of pyranopterin (Mendel and Leimkühler, 2015). Another metabolic enzyme, AldA, aldehyde dehydrogenase central carbohydrate metabolism is downregulated in all doses of pexiganan. This is also the case of CopZ, a chaperone that serves for the intracellular sequestration and transport of copper, delivering it to the copper-exporting P-type ATPase A (CopA) (Sitthisak et al., 2007).

**Pexiganan stress has strong impact on essential proteome.** We visualised the global impact of pexiganan stress (at 1x MIC) on bacterial physiology by a network analysis based on protein-protein interactions and function (Szklarczyk et al., 2015) of *S. aureus* essential genes (Figure S3). This network analysis provides a global view information on protein level alterations and integrates protein-protein interactions, including indirect functional and direct physical associations (Szklarczyk et al., 2015). At this concentration, it is noticeable that the majority of the essential genes are downregulated and it is possible that this pattern has strong influence on pexiganan lethality. The majority of unregulated proteins are ribosomal components.

**Gene ontology analysis points to an upregulation of nucleotide metabolism.** The signature of pexiganan stress on *S. aureus* in the upregulated fraction points to nucleotide metabolism related genes. GO annotation allows enrichment analysis providing global information based on the gene expression levels by proteomics or transcriptomics or other gene expression datasets (Mi et al., 2019). We focus this comparative analysis on the protein expression levels of the most 100 most upregulated proteins of every pexiganan dosage. We focussed on categorizing by pathways. Some of the upregulated pathways involved genes related with oxidative stress, peptidoglycan synthesis and N-acetylglucosamine that are expected from cationic antimicrobial peptides since they attack the cell envelopes. In addition, there is a reactivation of the central metabolism by the upregulation of genes from glycolysis, TCA cycle, arginine and thiamine synthesis. However, the most enriched pathways in the GO analysis for all pexiganan concentrations were related with nucleotide metabolism (Figure 2). The nucleotide upregulated pathways include ATP synthesis, Adenine and hypoxanthine salvage pathways, de novo synthesis of purines and pyrimidines and S-adenosylmethionine. This result confirms that pexiganan stress induces a scarcity of these metabolites within the cell. Taking into account the previous results, we hypothesised that upregulation of nucleotide dependent and related genes could create a collateral sensitivity.

**Nucleoside analogues antimetabolites act synergistically or additively with pexiganan.** We designed a simple drug interaction experiment between pexiganan and some nucleoside analogues including the purine and pyrimidines antimetabolites: 6-azauracil, gemcitabine, 5-fluorouracil

and 6-thioguanine (Figure S4, Figure S5, Table S4). This experiment is the classic isobologram, also known as checkerboard assay (Tallarida, 2006). All analogues showed additive (5-fluorouracil, 6-azauracil and 6-thiogunine) or synergistic (gemcitabine) activity when combined with pexiganan (Table S4). The most active ones were gemcitabine and 5-fluorouracil, while the 6-azauracil and 6-thiogunine showed a milder effect. However, all of them managed a decrease all cases of the minimal inhibitory concentration for each drug when compared to the respective drug alone. These results indicate pexiganan induces a collateral sensitivity to nucleoside analogues.

To study the influence of nucleoside analogues on the killing by pexiganan, we carried out a time-kill experiment combining each of 6-azauracil, gemcitabine, 5-fluorouracil and 6-thioguanine with pexiganan. We assayed all drugs using half of the minimal inhibitory concentration. We exposed mid-exponential phase *S. aureus* cells to these combinations and sampled the viability of the cultures every 20 minutes (Figure 3). All compounds significantly increased the killing ability of pexiganan, being gemcitabine and 5-fluorouracil the most active drugs. The killing rate was increased by some order of magnitudes in all combinations. The killing by the combination with gemcitabine or 5-fluorouracil with pexiganan, at their corresponding half a MIC values, were more efficient killing bacteria than 8x MIC concentration of pexiganan alone. The viability was assessed not only by the absence of growth but also by the addition of resazurin, a reagent that turn from blue to purple when it is reduced by microbial enzymes that only work within living bacteria (Elshikh et al., 2016).

Figure 1. Heatmap of relative protein expression based on label-free quantification by liquid chromatography/mass spectrometry (LC-MS). Only the 50 most significantly up-regulated proteins at 1XMIC. Red rectangle highlights proteins that participate in or depend on nucleotide metabolism. Proteins were extracted after 30 minutes of pexiganan addition (0.125, 0.25, 0.5 and 1 fractions of the minimal inhibitory concentration). Intensity ranges of the log2 fold-changes are given from highest intensity (green) to lowest (red).

## Panther pathway analysis (gene ontology)

Identified genes: 93, pathway hits: 14

Indentified genes: 91, pathway hits: 12

1/8x MIC

1/4x MIC

- ATP synthesis
- Adenine and hypoxantine salvage pathway
- De novo purine biosynthesis
- De novo pyrimidine ribonucleoides biosynthesis
- Xanthine and guanine salvage pathway
- S-adenosylmethionine biosynthesis
- Hypoxia response via HIF activation
- Oxidative stress
- Parkinson disease
- Peptidoglycan biosynthesis
- N-acetylglucosamine metabolism
- O-antigen biosynthesis
- 5-Hydroxytryptamine degradation
- Pyruvate metabolism
- Fructose galactose metabolism
- Glycolysis
- TCA cycle
- Arginine biosynthesis
- Cholesterol biosynthesis
- Thiamin biosynthesis

Indentified genes: 93, pathway hits: 16

Indentified genes: 94, pathway hits: 18

1/2x MIC

1x MIC

Figure 2. Functional characterisation of pathways of up-regulated proteins in *S. aureus* SH1000 at different concentrations of pexiganan (0.125, 0.25, 0.5 and 1 fractions of the minimal inhibitory concentration). For this analysis only the 100 most highly significant expressed proteins for each concentration of pexiganan were used. The analysis was carried out using the online gene ontology analysis software PANTHER (Mi et al., 2019).

Figure 3. Isobolographical response of pexiganan combination with different antimetabolite nucleosides. Blue squares represent the minimal inhibitory concentrations for pexiganan and each of the tested analogues (see also supplementary Figure S4). The crosses indicate the presence of bacterial growth in the unique concentration combinations of each well.

Figure 4. Pexiganan-nucleoside antimetabolite combination induced drastically increases the killing capacity of pexiganan. (A) Killing dynamic of *S. aureus* SH100 at different concentrations of pexiganan using the MIC as the starting point. (B) Example data of time-kill experiment exposing mid-exponential phase bacterial cultures to pexiganan-nucleoside antimetabolite combinations (both at 1/2x MIC concentrations). The combination has a dramatic effect on killing ability of pexiganan. Data points were determined by counting colony-forming units (CFU) at different time points. Mean ± SDM, n=5. Asterisks represent significant differences (R package nparLD, one asterisk for $p < 0.05$ and two asterisks for $p < 0.01$ and three asterisks for $p < 0.001$). Only comparisons between pexiganan (1/2x MIC) and pexiganan-analogues combinations are shown.

## Discussion.

We have found that pexiganan, a cationic antimicrobial peptide, can induce a stress response in *S. aureus* that results in a proteome-wide impact. Pexiganan treatment upregulates known virulence factors such as MprF, the capsule synthesis protein CapF, a wall teichoic acid TagG, the proteases ClpL and PepT and other important proteins for the interactions with the hosts. This could lead to a phenotypic cross-tolerance of other immune effectors of hosts and possibly complicate the bacterial infection in case of inefficient treatment where bacteria could be exposed to sub-lethal concentrations. This is a legitimate concern that have been reported for antimicrobial peptide evolved resistant variants that have shown some cross-resistance with immune system effectors (Bell and Gouyon, 2003; Fleitas and Franco, 2016). This risk has been shown for pexiganan (Habets and Brockhurst, 2012). Our data also provides input about possible induced physiological changes that would help *S. aureus* to adapt to the intra-host environment during it interaction with specific immune system effectors.

It is important to note that, given coverage of the proteomic data and range of pexiganan doses, we did not find evidence of activation of mutagenic stress pathways or recombination. This indicates that the mode of killing by cationic antimicrobial peptide does not increase genome instability as is typical for classic antibiotics (Blázquez et al., 2012). Along these lines, we

have previously shown and proposed that antimicrobial peptides, including pexiganan, do not increase mutagenesis (Rodríguez-Rojas et al., 2014) neither recombination (Rodríguez-Rojas et al., 2018) in Gram-negative bacteria. Our findings here are consistent with these observations in the Gram-positive model bacterium *S. aureus*.

The elevated level of expression of proteins such as GmK, PyrG, NptA and some amino acids related enzymes such as CarA and CarB that participate in the biosynthesis of L-arginine could be explained by changes in permeability. Amino acids, nucleobase and nucleotides are small molecules that could easily escape from the cellular compartment in case of membrane damage. This is a well know property of cationic agents, including AMPs (Asthana et al., 2004; Brogden, 2005; Rodríguez-Rojas et al., 2015). The fact that only a few proteins from the amino acids biosynthesis pathways are upregulated could be explained because the experiments were carried out in a complex medium like MHB that contains several amino acids and bacteria would upregulate only necessary pathways. A similar situation maybe expected within a host.

The upregulation of the phosphate and nucleotide-related proteins provides a direction to investigate drug susceptibilities created by pexiganan stress. Although the antimetabolites used in this work have good antibacterial activity, if they are used in monotherapy they are also prone to generate resistance (Jordheim et al., 2012b; Thomson and Lamont, 2019). Thus, their use in combination could possibly help to prevent resistance (Yu et al., 2016; Tyers and Wright, 2019).

The combined action of pexiganan with nucleoside antimetabolites that we report here to be either additive or synergistic could have two underlying reasons. First, pexiganan stress forces a response by *S. aureus* that upregulates nucleobase salvage pathways and other nucleotide-dependent metabolic pathways. Second, pexiganan has the potential to change membrane permeability and induce the uptake of such metabolites even at sub-lethal concentrations possibly leading to much higher intracellular concentrations (Figure 5). We have shown previously that cationic antimicrobial peptides can mediate the uptake of small molecules due to changes in permeability at sub-lethal concentrations (Rodríguez-Rojas et al., 2015). The more potent activity of gemcitabine and 5-fluorouracil could be explained because they act on the cell walls as previously reported (Gieringer et al., 1986; Jordheim et al., 2012b).

An additional advantage of the potential therapeutic advantage of the nucleoside analogues studied here is that all the clinical properties of these drugs are well known, including toxicological profile, pharmacological activities and metabolising properties (Cheng et al., 2018; Thomson and Lamont, 2019). All of them are approved drugs, which should facilitate the introduction of such combinations in clinical practices.

We have shown recently that antimicrobial peptides, including pexiganan, can induce priming in bacteria, an enhanced response to the peptides when bacteria are pre-exposed to low concentrations. The consequence of priming is not only survival but increases the tolerance and persistence (Rodríguez-Rojas et al., 2019). The use of antimetabolites could potentially abolish this property in therapeutic usage.

Figure 5. A general model illustrating the positive interaction between pexiganan and nucleoside antimetabolites against *S. aureus*. The interactions of pexiganan with the membrane at sub-inhibitory concentrations lead to transient permeability changes in the envelope that promote small molecules leakage such as nucleotides, nucleobase or nucleoside. Simultaneously, other small molecules such as toxic nucleoside analogues can increase the diffusion rate toward intracellular compartment. This stress is sensed by the cell that responds activating nucleoside metabolism creating an intervention opportunity. In this situation, toxic nucleoside antimetabolites are more efficiently incorporated into RNA, DNA other nucleotide depending reactions such as envelope synthesis, enhancing toxicity and leading to faster cell killing, even below MIC for both compounds.

**Conclusions.**

The analysis of the pexiganan stress response by *S. aureus* has shown a global response involving several proteins known for their role on antimicrobial peptides and other immune system effector resistance. Pexiganan has also shown an increased antibacterial activity when it is combined with nucleoside antimetabolites. Taken together, our results suggest that pexiganan renders *S. aureus* susceptible to purine and pyrimidine analogues which are traditionally used for cancer treatment. These analogues can enhance the bactericidal activity of pexiganan against *S. aureus* under the tested conditions. The significant potentiation of the

pexiganan bactericidal activity and the decrease of minimal inhibitory concentrations when compared with pexiganan alone could be the basis for new formulations of pexiganan. These results are probably extendable to other antimicrobial peptides and other bacterial pathogens. Thus, the leakage of nucleotides and intermediate small metabolites or cofactors caused by cationic peptides and nucleotide metabolic pathways are common traits of bacteria-peptide interactions as proposed for the symbiont–host interface (Mergaert et al., 2017). Our results also show that understanding how antimicrobials work and how pathogens respond to them is important to increase effectivity of the therapy. Physiological response by bacteria is informative or suggestive about additional combinations that can limit the chances of pathogens to evolve resistance. This approach should be exploited to rationally design new antimicrobial combinations.

**Material and methods.**

**Bacteria and growth conditions.** We used *S. aureus* SH1000 (Horsburgh et al., 2002) for all experiments. Bacteria were cultured in non-cation-adjusted (unsupplemented) Mueller–Hinton broth (MHB) as recommended for antimicrobial peptides susceptibility testing (Giacometti et al., 2000).

**Global proteomics by LC-mass spectrometry.** *Staphylococcus aureus* strain SH1000 was grown in non-cation-adjusted MHB to the mid-exponential-phase ($OD_{600}$ 0.5) at 37°C with vigorous shaking. The cultures were diluted 100 times in fresh MHB in a separate tube with a final volume of 5 ml. Pexiganan was added to tubes for a final concentration of 0.5, 1, 2 and 4 µg/ml (1/8,1/4, 1/2, 1x MIC respectively) in a final culture volume of 10 ml per tube. Non-treated samples were used as controls. After the addition of the treatments, all tubes were incubated for 30 minutes with moderate shaking at 37°C. The pellets were collected by centrifugation at 10 000 x g for 5 minutes and the supernatant was removed by aspiration using a sterile vacuum line. The pellets then added 50 µl of denaturation urea buffer (6 M urea/2 M thiourea/10 mM HEPES, pH 8.0). The resulting suspensions were transferred to new 1.5 ml Eppendorf tubes and exposed to 5 freeze-thawing cycles alternating liquid nitrogen and a 37°C water baths. The tubes were centrifuged at 20 000 x g for 10 minutes and the supernatants were transferred to fresh tubes and used as starting protein material for digestion. Each experimental condition had six independent biological replications. Approximately 50 µg proteins were processed per sample and were in-solution digested as described elsewhere (Rappsilber et al., 2007). Denaturation buffer-containing protein solutions were reduced adding 1 µl of 10 mM DTT (final concentration) and incubated for 30 minutes. The reactions were then alkylated by adding 1 µl of 55 mM iodoacetamide and incubated for 20 minutes in the dark. Lysyl

endopeptidase (LysC, Wako, Japan) resuspended in 50 mM ABC was added to digestion reaction in a proportion of 1 µg per 50 µg of total sample protein and incubated for 3 hours. The samples were diluted with four volumes of 50 mM ABC buffer and digested overnight with trypsin using 1 µg per tube of sequencing grade modified trypsin (Promega, USA) diluted in the buffer provided by the manufacturer. All digestion steps were incubated at room temperature. Next day, the digestions were stopped by adding 5% acetonitrile and 0.3% trifluoroacetic acid (final concentrations). The samples were processed using the Stage-tip protocol as described previously (Rappsilber et al., 2007), and the eluates were vacuum-dried. Re-dissolved samples were loaded on a ReprosilPur C18 reverse phase column and peptides were analysed using a nano-HPLC Dionex Ultimate 3000 system (Thermo Scientific, Germany) coupled to an Orbitrap Velos mass spectrometer (Thermo Scientific, Germany). MS and MS/MS data from each LC/MS run were analysed using the MaxQuant software (Tyanova et al., 2016a). Identification of proteins was performed using the MaxQuant implemented Andromeda peptide search engine and statistical analysis was carried out using the software Perseus (Tyanova et al., 2016b).

**Antimetabolite nucleosides.** In this study we used four nucleoside analogues. We used the pyrimidine analogues 6-azauracil, gemcitabine, 5-fluorouracil and the purine analogue 6-thioguanine. All drugs were purchased from Sigma Aldrich (Germany).  6-Azauracil is used as a growth inhibitor of various microorganisms via depletion of intracellular GTP and UTP nucleotide pools (Habermann, 1961). Gemcitabine is a chemotherapy medication used to treat a different types of cancer. Gemcitabine is a synthetic pyrimidine nucleoside analogue in which the hydrogen atoms on the 2' carbon of deoxycytidine are replaced by fluorine atoms and competitively takes part and disrupts several pathways where pyrimidines are needed (Jordheim et al., 2012c). 5-Fluorouracil is also used as an anticancer treatment and it works by inhibiting cell metabolism by blocking many pathways, but its major action is the inhibition of the thymidylate synthase. By doing so, the synthesis of the pyrimidine thymidine is stalled, which is an essential nucleoside required for DNA replication (Singh et al., 2015). 5-Fluorouracil causes a drop on dTMP and then cells undergo cell death via thymineless death (Khodursky et al., 2015; Singh et al., 2015).

**Pexiganan and antimetabolite nucleosides susceptibility testing.** Minimal inhibitory concentration (MIC) was determined by broth micro-dilution method modified for cationic antimicrobial peptides (Wiegand et al., 2008). Briefly, 2 µl of the mid-exponential phase culture diluted 1:100 (around $10^5$ bacteria) were inoculated into each well of a polypropylene V-bottom 96-well plates with anti-evaporation ring lids (Greiner Bio-One GmbH, Germany). Prior to inoculation, pexiganan and the analogues (a kind gift from Dr Michael A. Zasloff, Georgetown University) were two-fold serially diluted in a final volume of 100 µl MHB per well using 32 µg/ml as starting concentration. Each assay was performed with eight replications

and the plate was incubated at 37°C in a humid chamber. The MIC was defined as the lowest concentration where no visible bacterial growth was observed after 24 hours.

**Isobologram Assay.** The combined activity and interactions between peptides and pexiganan and purine and pyrimidine analogues against *S. aureus* in MHB was determined using isobolographic combinations, also called checkerboard assay method, (8×8 matrix of concentrations combinations) (Tallarida, 2006). In a 96-well plate, 50 μl of pexiganan at 4x MIC concentration was two-fold serially diluted ranging from 32 to 0.25 μg/ml in the direction of the columns from 1 to 8. In another 96-well plate 100 μl of nucleoside analogues at 8x MIC concentrations were prepared in an identical way as the previous plate, but diluted in the direction of the rows from A to H. Half of the content (50 μl) of each well from the analogue drug plate was transferred to the corresponding well of plate containing pexiganan in an equal 1:1 mix fashion, halving the concentration of both compounds. In the same plate, the columns 9 and 10 were used to serially dilute both, the peptide and the analogue drug in the same concentrations that were present in the combination to compare single compounds vs combination. Columns 11 and 12 were used as a control, by inoculating column 11 wells with bacteria without any drug and leaving columns 12 only with the same volume of MHB as a media contamination control. Each plate was prepared in triplicates to check for consistency. The bacterial suspension was prepared by growing *S. aureus* SH1000 to mid-exponential phase (2.5 hours, with moderate shaking at 37°C) in MHB to an $OD_{600}$ between 0.3 to 0.5. The bacterial suspension was diluted in MHB and approximately $1 \times 10^6$ bacteria were inoculated in each well. After 24 h of incubation at 37°C in a humid chamber, the plates were visually examined for growth. The FIC index for a combination of pexiganan and each antimetabolite drug was calculated as [(MIC of the peptide in combination with a given analogue)/(MIC of peptide alone)] + [(MIC of analogue in combination with peptide)/(MIC of analogue alone)]. The interpretation of the results was as follow: FIC ≤ 0.5, synergistic; 0.5 < FIC ≤ 1, additive; 1 < FIC ≤ 4, indifferent; FIC > 4, antagonistic (Ng et al., 2018). To ensure that bacteria lost viability while reading MIC values for pexiganan-analogue combinations, we used the resazurin colorimetric assay as described previously with minor modifications (Elshikh et al., 2016). Resazurin (THK, Germany) was prepared at 0.015 % in in distilled water and sterilised by filtration. It was stored at 4 °C for a maximum of 1 weeks after preparation. Resazurin (0.015 %) was added each well (10 μl per well, 1/3 of the original described quantity) and further incubated for 3 h for the observation of colour change. Columns with no colour change (blue resazurin) were scored as dead culture. In contrast, colour change to purple (reduced resazurin) was considered as a sign of viability.

**Time-kill experiments.** Starting from early mid-exponential phase cultures ($1 \times 10^7$ CFU/ml), bacteria were exposed to growing concentrations

of pexiganan ranging from 1 to 8x MIC or pexiganan combined with the nucleoside analogues 6-azauracil, gemcitabine, 5-fluorouracil and 6-thioguanine at their respective 1/2x MIC values. The cultures were incubated with soft shaking at 37°C for 2 hours. Samples from each cultures (1ml) were taken at 20-minute time-point intervals. The samples were diluted and plated to determine cell viability. The experiments consisted of five independent replications. Non treated cells were used as a control.

**Statistical analysis**. The effect of treatments on bacterial killing were analysed using R package nparLD (Noguchi et al., 2012). *P* values less than or equal to 0.05, after correction, if needed, were considered statistically significant. All tests were performed with the statistic software R (R Core Team, 2017).

**Funding.** ARR and JR were supported SFB 973 (Deutsche Forschungsgemeinschaft, project C5). We acknowledge support by the German Research Foundation and the Open Access Publication Fund of Freie Universität Berlin. For mass spectrometry (B.K. and C.W.) we would like to acknowledge the assistance of the Core Facility BioSupraMol supported by the Deutsche Forschungsgemeinschaft (DFG).

**Acknowledgements.** We would like to thank Dr Dan Roizman from Freie Universität Berlin for help with Resazurin assay. We would like to thank also Dr Michael A. Zasloff from Georgetown University for kindly providing the pexiganan.

# References

1. Anderson, K. L., Roberts, C., Disz, T., Vonstein, V., Hwang, K., Overbeek, R., et al. (2006). Characterization of the Staphylococcus aureus heat shock, cold shock, stringent, and SOS responses and their effects on log-phase mRNA turnover. *J. Bacteriol.* 188, 6739–6756. doi:10.1128/JB.00609-06.

2. Asthana, N., Yadav, S. P., and Ghosh, J. K. (2004). Dissection of antibacterial and toxic activity of melittin: a leucine zipper motif plays a crucial role in determining its hemolytic activity but not antibacterial activity. *J. Biol. Chem.* 279, 55042–50. doi:10.1074/jbc.M408881200.

3. Baker, S. (2015). A return to the pre-antimicrobial era? *Science (80-. ).* 347, 1064–1066. doi:10.1126/science.aaa2868.

4. Bell, G., and Gouyon, P. H. (2003). Arming the enemy: The evolution of resistance to self-proteins. *Microbiology* 149, 1367–1375. doi:10.1099/mic.0.26265-0.

5. Blázquez, J., Couce, A., Rodríguez-Beltrán, J., Rodríguez-Rojas, A., Blazquez, J., Couce, A., et al. (2012). Antimicrobials as promoters of genetic variation. *Curr Opin Microbiol* 15, 561–569. doi:S1369-5274(12)00102-6 [pii]10.1016/j.mib.2012.07.007.

6. Brogden, K. A. (2005). Antimicrobial peptides: pore formers or metabolic inhibitors in bacteria? *Nat. Rev. Microbiol.* 3, 238–50. doi:10.1038/nrmicro1098.

7. Brown, S., Santa Maria, J. P., and Walker, S. (2013). Wall Teichoic Acids of Gram-Positive Bacteria. *Annu. Rev. Microbiol.* 67, 313–336. doi:10.1146/annurev-micro-092412-155620.

8. Cheng, Y.-S., Sun, W., Xu, M., Shen, M., Khraiwesh, M., Sciotti, R. J., et al. (2018). Repurposing Screen Identifies Unconventional Drugs With Activity Against Multidrug Resistant Acinetobacter baumannii. *Front. Cell. Infect. Microbiol.* 8, 438. doi:10.3389/fcimb.2018.00438.

9. Czaplewski, L., Bax, R., Clokie, M., Dawson, M., Fairhead, H., Fischetti, V. A., et al. (2016). Alternatives to antibiotics-a pipeline portfolio review. *Lancet Infect. Dis.* 16, 239–251. doi:10.1016/S1473-3099(15)00466-1.

10. Di Martino, M. L., Campilongo, R., Casalino, M., Micheli, G., Colonna, B., and Prosseda, G. (2013). Polyamines: Emerging players in bacteria-host interactions. *Int. J. Med. Microbiol.* 303, 484–491. doi:10.1016/j.ijmm.2013.06.008.

11. Elshikh, M., Ahmed, S., Funston, S., Dunlop, P., McGaw, M., Marchant, R., et al. (2016). Resazurin-based 96-well plate microdilution method for the determination of minimum inhibitory concentration of biosurfactants. *Biotechnol. Lett.* 38, 1015–1019. doi:10.1007/s10529-016-2079-2.

12. Fleitas, O., and Franco, O. L. (2016). Induced Bacterial Cross-Resistance toward Host Antimicrobial Peptides: A Worrying Phenomenon. *Front. Microbiol.* 7, 381. doi:10.3389/fmicb.2016.00381.

13. Frees, D., Gerth, U., and Ingmer, H. (2014). Clp chaperones and proteases are central in stress survival, virulence and antibiotic resistance of Staphylococcus aureus. *Int. J. Med. Microbiol.* 304, 142–149. doi:10.1016/j.ijmm.2013.11.009.

14. Garnett, J. P., Braun, D., McCarthy, A. J., Farrant, M. R., Baker, E. H., Lindsay, J. A., et al. (2014). Fructose transport-deficient Staphylococcus aureus reveals important role of epithelial glucose transporters in limiting sugar-driven bacterial growth in airway surface liquid. *Cell. Mol. Life Sci.* 71, 4665–73. doi:10.1007/s00018-014-1635-y.

15. Ge, Y., MacDonald, D. L., Holroyd, K. J., Thornsberry, C., Wexler, H., and Zasloff, M. (1999). In Vitro Antibacterial Properties of Pexiganan, an Analog of Magainin. *Antimicrob. Agents Chemother.* 43, 782–788.

16. Giacometti, A., Cirioni, O., Barchiesi, F., Del Prete, M. S., Fortuna, M., Caselli, F., et al. (2000). In Vitro Susceptibility Tests for Cationic Peptides: Comparison of Broth Microdilution Methods for Bacteria That Grow Aerobically. *Antimicrob. Agents Chemother.* 44, 1694–1696.

doi:10.1128/AAC.44.6.1694-1696.2000.

17. Gieringer, J. H., Wenz, A. F., Just, H.-M., and Daschner, F. D. (1986). Effect of 5-Fluorouracil, Mitoxantrone, Methotrexate, and Vincristine on the Antibacterial Activity of Ceftriaxone, Ceftazidime, Cefotiam, Piperacillin, and Netilmicin. *Chemotherapy* 32, 418–424. doi:10.1159/000238445.

18. Habermann, V. (1961). The effect of 6-azauracil on microorganisms inhibited by chloramphenicol. *BBA - Biochim. Biophys. Acta* 49, 204–211. doi:10.1016/0006-3002(61)90884-8.

19. Habets, M. G. J. L., and Brockhurst, M. a (2012). Therapeutic antimicrobial peptides may compromise natural immunity. *Biol. Lett.* 8, 416–8. doi:10.1098/rsbl.2011.1203.

20. Hancock, R. E. W., and Sahl, H.-G. (2006). Antimicrobial and host-defense peptides as new anti-infective therapeutic strategies. *Nat. Biotechnol.* 24, 1551–7. doi:10.1038/nbt1267.

21. Horsburgh, M. J., Aish, J. L., White, I. J., Shaw, L., Lithgow, J. K., and Foster, S. J. (2002). δb modulates virulence determinant expression and stress resistance: Characterization of a functional rsbU strain derived from Staphylococcus aureus 8325-4. *J. Bacteriol.* 184, 5457–5467. doi:10.1128/JB.184.19.5457-5467.2002.

22. Imamura, T., Tanase, S., Szmyd, G., Kozik, A., Travis, J., and Potempa, J. (2005). Induction of vascular leakage through release of bradykinin and a novel kinin by cysteine proteinases from Staphylococcus aureus. *J. Exp. Med.* 201, 1669–76. doi:10.1084/jem.20042041.

23. Johnston, P. R., Dobson, A. J., and Rolff, J. (2016). Genomic Signatures of Experimental Adaptation to Antimicrobial Peptides in Staphylococcus aureus. *G3 (Bethesda).* 6, 1535–9. doi:10.1534/g3.115.023622.

24. Jordheim, L. P., Ben Larbi, S., Fendrich, O., Ducrot, C., Bergeron, E., Dumontet, C., et al. (2012a). Gemcitabine is active against clinical multiresistant Staphylococcus aureus strains and is synergistic with gentamicin. *Int. J. Antimicrob. Agents* 39, 444–447. doi:10.1016/J.IJANTIMICAG.2012.01.019.

25. Jordheim, L. P., Ben Larbi, S., Fendrich, O., Ducrot, C., Bergeron, E., Dumontet, C., et al. (2012b). Gemcitabine is active against clinical multiresistant Staphylococcus aureus strains and is synergistic with gentamicin. *Int. J. Antimicrob. Agents* 39, 444–447. doi:10.1016/j.ijantimicag.2012.01.019.

26. Jordheim, L. P., Ben Larbi, S., Fendrich, O., Ducrot, C., Bergeron, E., Dumontet, C., et al. (2012c). Gemcitabine is active against clinical multiresistant Staphylococcus aureus strains and is synergistic with gentamicin. *Int. J. Antimicrob. Agents* 39, 444–447. doi:10.1016/j.ijantimicag.2012.01.019.

27. Kantyka, T., Pyrc, K., Gruca, M., Smagur, J., Plaza, K., Guzik, K., et al. (2013). Staphylococcus aureus proteases degrade lung surfactant protein A potentially impairing innate immunity of the lung. *J. Innate Immun.* 5, 251–60. doi:10.1159/000345417.

28. Khodursky, A., Guzmán, E. C., and Hanawalt, P. C. (2015). Thymineless Death Lives On: New Insights into a Classic Phenomenon. *Annu. Rev. Microbiol.* 69, 247–263. doi:10.1146/annurev-micro-092412-155749.

29. Kristian, S. A., Dürr, M., Van Strijp, J. A. G., Neumeister, B., and Peschel, A. (2003). MprF-mediated lysinylation of phospholipids in Staphylococcus aureus leads to protection against oxygen-independent neutrophil killing. *Infect. Immun.* 71, 546–9.

30. Lofton, H., Pränting, M., Thulin, E., and Andersson, D. I. (2013). Mechanisms and fitness costs of resistance to antimicrobial peptides LL-37, CNY100HL and wheat germ histones. *PLoS One* 8, e68875. doi:10.1371/journal.pone.0068875.

31. Makarova, O., Johnston, P., Rodriguez-Rojas, A., El Shazely, B., Morales, J. M., and Rolff, J. (2018). Genomics of experimental adaptation of Staphylococcus aureus to a natural combination of insect antimicrobial

peptides. *Sci. Rep.* 8. doi:10.1038/s41598-018-33593-7.

32. Mendel, R. R., and Leimkühler, S. (2015). The biosynthesis of the molybdenum cofactors. *J. Biol. Inorg. Chem.* 20, 337–347. doi:10.1007/s00775-014-1173-y.

33. Mergaert, P., Kikuchi, Y., Shigenobu, S., and Nowack, E. C. M. (2017). Metabolic Integration of Bacterial Endosymbionts through Antimicrobial Peptides. *Trends Microbiol.* 25, 703–712. doi:10.1016/j.tim.2017.04.007.

34. Mi, H., Muruganujan, A., Ebert, D., Huang, X., and Thomas, P. D. (2019). PANTHER version 14: more genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools. *Nucleic Acids Res.* 47, D419–D426. doi:10.1093/nar/gky1038.

35. Miyafusa, T., Caaveiro, J. M. M., Tanaka, Y., and Tsumoto, K. (2013). Dynamic elements govern the catalytic activity of CapE, a capsular polysaccharide-synthesizing enzyme from *Staphylococcus aureus*. *FEBS Lett.* 587, 3824–3830. doi:10.1016/j.febslet.2013.10.009.

36. Mookherjee, N., Anderson, M. A., Haagsman, H. P., and Davidson, D. J. (2020). Antimicrobial host defence peptides: functions and clinical potential. *Nat. Rev. Drug Discov.* doi:10.1038/s41573-019-0058-8.

37. Ng, V., Kuehne, S. A., and Chan, W. C. (2018). Rational Design and Synthesis of Modified Teixobactin Analogues: In Vitro Antibacterial Activity against Staphylococcus aureus, Propionibacterium acnes and Pseudomonas aeruginosa. *Chem. - A Eur. J.* 24, 9136–9147. doi:10.1002/chem.201801423.

38. Nicolas, P., and Mor, A. (1995). Peptides as weapons against microorganisms in the chemical defense system of vertebrates. *Annu. Rev. Microbiol.* 49, 277–304. doi:10.1146/annurev.mi.49.100195.001425.

39. Noguchi, K., Gel, Y. R., Brunner, E., and Konietschke, F. (2012). nparLD : An R Software Package for the Nonparametric Analysis of Longitudinal Data in Factorial Experiments . *J. Stat. Softw.* 50, 1–23. doi:10.18637/jss.v050.i12.

40. Oku, Y., Kurokawa, K., Ichihashi, N., and Sekimizu, K. (2004). Characterization of the Staphylococcus aureus mprF gene, involved in lysinylation of phosphatidylglycerol. *Microbiology* 150, 45–51. doi:10.1099/mic.0.26706-0.

41. Omari, K. El, Dhaliwal, B., Lockyer, M., Charles, I., Hawkins, A. R., and Stammers, D. K. (2006). Structural Biology and Crystallization Communications Structure of Staphylococcus aureus guanylate monophosphate kinase. *Struct. Commun. Acta Cryst* 62, 949–953. doi:10.1107/S174430910603613X.

42. Perron, G. G., Zasloff, M., and Bell, G. (2006). Experimental evolution of resistance to an antimicrobial peptide. *Proc. Biol. Sci.* 273, 251–6. doi:10.1098/rspb.2005.3301.

43. Perry, W. J., Spraggins, J. M., Sheldon, J. R., Grunenwald, C. M., Heinrichs, D. E., Cassat, J. E., et al. (2019). Staphylococcus aureus exhibits heterogeneous siderophore production within the vertebrate host. *Proc. Natl. Acad. Sci. U. S. A.* 116, 21980–21982. doi:10.1073/pnas.1913991116.

44. Potempa, J., Dubin, A., Korzus, G., and Travis, J. (1988). Degradation of elastin by a cysteine proteinase from Staphylococcus aureus. *J. Biol. Chem.* 263, 2664–2667.

45. Pundir, S., Martin, M. J., and O'Donovan, C. (2016). UniProt Tools. *Curr. Protoc. Bioinforma.* 53, 1.29.1-1.29.15. doi:10.1002/0471250953.bi0129s53.

46. R Core Team (2017). *R: A language and environment for statistical computing.* Vienna, Austria: R Foundation for Statistical Computing.

47. Rappsilber, J., Mann, M., and Ishihama, Y. (2007). Protocol for micro-purification, enrichment, pre-fractionation and storage of peptides for proteomics using StageTips. *Nat. Protoc.* 2, 1896–906. doi:10.1038/nprot.2007.261.

48. Rodríguez-Rojas, A., Baeder, D. Y., Johnston, P., Regoes, R. R., and Rolff, J. (2019). Bacteria primed by antimicrobial peptides develop tolerance and persist. *bioRxiv*, 802207. doi:10.1101/802207.

49. Rodríguez-Rojas, A., Makarova, O., Müller, U., and Rolff, J. (2015). Cationic Peptides Facilitate Iron-induced Mutagenesis in Bacteria. 11, e1005546. doi:10.1371/journal.pgen.1005546.

50. Rodríguez-Rojas, A., Makarova, O., and Rolff, J. (2014). Antimicrobials, stress and mutagenesis. *PLoS Pathog.* 10, e1004445. doi:10.1371/journal.ppat.1004445.

51. Rodríguez-Rojas, A., Moreno-Morales, J., Mason, A. J., and Rolff, J. (2018). Cationic antimicrobial peptides do not change recombination frequency in *Escherichia coli*. *Biol. Lett.* 14, 20180006. doi:10.1098/rsbl.2018.0006.

52. ROGERS, H. J., and PERKINS, H. R. (1960). 5-Fluorouracil and mucopeptide biosynthesis by Staphylococcus aureus. *Biochem. J.* 77, 448–459. doi:10.1042/bj0770448.

53. Scholefield, G., and Murray, H. (2013). YabA and DnaD inhibit helix assembly of the DNA replication initiation protein DnaA. *Mol. Microbiol.* 90, 147–159. doi:10.1111/mmi.12353.

54. Singh, V., Brecik, M., Mukherjee, R., Evans, J. C., Svetlíková, Z., Blaško, J., et al. (2015). The complex mechanism of antimycobacterial action of 5-fluorouracil. *Chem. Biol.* 22, 63–75. doi:10.1016/j.chembiol.2014.11.006.

55. Sitthisak, S., Knutsson, L., Webb, J. W., and Jayaswal, R. K. (2007). Molecular characterization of the copper transport system in Staphylococcus aureus. *Microbiology* 153, 4274–4283. doi:10.1099/mic.0.2007/009860-0.

56. Spohn, R., Daruka, L., Lázár, V., Martins, A., Vidovics, F., Grézal, G., et al. (2019). Integrated evolutionary analysis reveals antimicrobial peptides with limited resistance. *Nat. Commun.* 10. doi:10.1038/s41467-019-12364-6.

57. Stapleton, M. R., Horsburgh, M. J., Hayhurst, E. J., Wright, L., Jonsson, I. M., Tarkowski, A., et al. (2007). Characterization of IsaA and SceD, two putative lytic transglycosylases of Staphylococcus aureus. *J. Bacteriol.* 189, 7316–7325. doi:10.1128/JB.00734-07.

58. Staubitz, P., Neumann, H., Schneider, T., Wiedemann, I., and Peschel, A. (2004). MprF-mediated biosynthesis of lysylphosphatidylglycerol, an important determinant in staphylococcal defensin resistance. *FEMS Microbiol. Lett.* 231, 67–71. doi:10.1016/S0378-1097(03)00921-2.

59. Stickgold, R. A., and Neuhaus, F. C. (1967). On the initial stage in peptidoglycan synthesis. Effect of 5-fluorouracil substitution on phospho-N-acetylmuramyl-pentapeptide translocase (uridine 5′-phosphate). *J. Biol. Chem.* 242, 1331–7.

60. Szklarczyk, D., Franceschini, A., Wyder, S., Forslund, K., Heller, D., Huerta-Cepas, J., et al. (2015). STRING v10: protein–protein interaction networks, integrated over the tree of life. *Nucleic Acids Res.* 43, D447–D452. doi:10.1093/nar/gku1003.

61. Tallarida, R. J. (2006). An overview of drug combination analysis with isobolograms. *J. Pharmacol. Exp. Ther.* 319, 1–7. doi:10.1124/jpet.106.104117.

62. Thomson, J. M., and Lamont, I. L. (2019). Nucleoside analogues as antibacterial agents. *Front. Microbiol.* 10, 952. doi:10.3389/fmicb.2019.00952.

63. Tyanova, S., Temu, T., and Cox, J. (2016a). The MaxQuant computational platform for mass spectrometry-based shotgun proteomics. *Nat. Protoc.* 11, 2301–2319. doi:10.1038/nprot.2016.136.

64. Tyanova, S., Temu, T., Sinitcyn, P., Carlson, A., Hein, M. Y., Geiger, T., et al. (2016b). The Perseus computational platform for comprehensive analysis of (prote)omics data. *Nat. Methods* 13, 731–40. doi:10.1038/nmeth.3901.

65. Tyers, M., and Wright, G. D. (2019). Drug combinations: a strategy to extend the life of antibiotics in the 21st century. *Nat. Rev. Microbiol.* 17, 141–155. doi:10.1038/s41579-018-0141-x.

66. Vitko, N. P., Grosser, M. R., Khatri, D., Lance, T. R., and Richardson, A. R. (2016). Expanded glucose import capability affords Staphylococcus aureus optimized glycolytic flux during infection. *MBio* 7. doi:10.1128/mBio.00296-16.

67. Wang, Z., and Wang, G. (2004). APD: the Antimicrobial peptide database. *Nucleic Acids Res* 32, D590–D592.

68. Weidenmaier, C., Kristian, S., and Peschel, A. (2005). Bacterial Resistance to Antimicrobial Host Defenses - An Emerging Target for Novel Antiinfective Strategies? *Curr. Drug Targets* 4, 643–649. doi:10.2174/1389450033490731.

69. Westerhoff, H. V., Juretic, D., Hendler, R. W., and Zasloff, M. (1989). Magainins and the disruption of membrane-linked free-energy transduction. *Proc. Natl. Acad. Sci. U. S. A.* 86, 6597–6601. doi:10.1073/pnas.86.17.6597.

70. Wiegand, I., Hilpert, K., and Hancock, R. E. W. (2008). Agar and broth dilution methods to determine the minimal inhibitory concentration (MIC) of antimicrobial substances. *Nat. Protoc.* 3, 163–75. doi:10.1038/nprot.2007.521.

71. Yan, H., and Hancock, R. E. W. (2001). Synergistic interactions between mammalian antimicrobial defense peptides. *Antimicrob. Agents Chemother.* 45, 1558–1560. doi:10.1128/AAC.45.5.1558-1560.2001.

72. Yu, G., Baeder, D. Y., Regoes, R. R., and Rolff, J. (2016). Combination Effects of Antimicrobial Peptides. *Antimicrob. Agents Chemother.* 60, 1717–1724. doi:10.1128/AAC.02434-15.

73. Yu, G., Baeder, D. Y., Regoes, R. R., and Rolff, J. (2018). Predicting drug resistance evolution: Insights from antimicrobial peptides and antibiotics. *Proc. R. Soc. B Biol. Sci.* 285. doi:10.1098/rspb.2017.2687.

74. Zasloff, M. (2002). Antimicrobial peptides of multicellular organisms. *Nature* 415, 389–95. doi:10.1038/415389a.

Figure S1. Volcano plots (q-values versus Labal-free quantification diference versus control of LC-MS) showing pexiganan response at different fractions of the minimal inhibitory concentrations. Black dots represent non-differentially expressed proteins while green dots show the upregulated portions and red ones represent the down-regulated fraction.

Figure S2. Heatmap of relative protein expression based on label-free quantification by liquid chromatography/mass spectrometry (LC-MS). Only the 50 most statistically significant down-regulated proteins are shown, taking as a reference the ones from 1xMIC pexiganan concentration (0.125, 0.25, 0.5 and 1 fractions of the minimal inhibitory concentration). Intensity ranges of the log2 fold-changes are given from highest intensity (green) to lowest (red).

Figure S3. Network analysis of pexiganan stress (1xMIC) on essential genes interactome of *S. aureus* SH1000. Pale green nodes indicate upregulated proteins while pale red ones represent down-regulated ones. Grey nodes correspond with genes with a high degree of connectivity with this essential network but they were not labelled. Note that the higher proportion of downregulated genes among essential proteome while majority of unregulated proteins are ribosomal components and they aggregate due to physical interaction. The interaction among nodes shows the proteome-wide impact of pexiganan stress at an inhibitory concentration.

6-Azauracil

Gemcitabine
hydrochloride

5-Fluorouracil

6-Thioguanine

Figure S4. Chemical structure of nucleoside antimetabolites used in this work. Images were obtained from Wikipedia.

Figure S5. Isobologram showing the additive and synergistic activity of pexiganan and different nucleotide antimetabolite combination against *S. aureus* SH1000.

Table S1. Minimum inhibitory concentration (MIC) data for *S. aureus* SH1000 to pexiganan and antimetabolite analogues used in this study.

| Drug | MIC (µg/ml) |
|---|---|
| pexiganan | 8 |
| 6-azauracil | 4 |
| gemcitabine | 0.5 |
| 5-fluorouracil | 2 |
| 6-thioguanine | 8 |

Table S2. Output table (excel file) of the proteomic experiment reporting *S. aureus* SH1000 response to pexiganan addition (0.125, 0.25, 0.5 and 1x MIC). Bacteria were sampled after 30 minutes of the treatment. Each treatment group consisted of six independent replicates and bacteria before treatment (T0) were used as control. Statistical analysis used student t-test and false discovery rate for correction of the p-values (data analysis using Maxquant (Tyanova et al., 2016a) and Perseus software (Tyanova et al., 2016b) for label-free quantification of proteins with LC-MS).

Table S3. Fractional inhibitory concentration (FIC) data for *S. aureus* SH1000 to Pexiganan and antimetabolite analogues used in this study.

| Drug combination | FIC index | Interaction type |
|---|---|---|
| pexiganan+gemcitabine | 1 | Aynergistic |
| pexiganan+6-azauracil | 2 | Additive |
| pexiganan+5-fluorouracil | 1.5 | Additive |
| pexiganan+6-thiogunaine | 2 | Additive |