1 **Assessing the impact of the threatened crucian carp (*Carassius***

2 ***carassius*) on pond invertebrate diversity - a comparison of**

3 **conventional and molecular tools**

5 **Lynsey R. Harper[1,2*], Lori Lawson Handley[1], Carl D. Sayer[3], Daniel S. Read[4],**

6 **Marco Benucci[1], Rosetta C. Blackman[1,5,6], Matthew J. Hill[7], and Bernd Hänfling[1]**

7
8
[1] Department of Biological and Marine Sciences, University of Hull, Hull, HU6 7RX, UK

[2] Illinois Natural History Survey, Prairie Research Institute, University of Illinois at Urbana-Champaign, Champaign,

IL, USA

[3] Pond Restoration Research Group, Environmental Change Research Centre, Department of Geography, University

College London, London, WC1E 6BT, UK

[4] Centre for Ecology & Hydrology (CEH), Benson Lane, Crowmarsh Gifford, Wallingford, Oxfordshire, OX10 8BB, UK

[5] Department of Aquatic Ecology, Eawag: Swiss Federal Institute of Aquatic Science and Technology, Dübendorf,

Switzerland

[6] Department of Evolutionary Biology and Environmental Studies, University of Zurich, Winterthurerstrasse 190,

8057 Zurich, Switzerland.

[7] School of Applied Sciences, University of Huddersfield, Queensgate, Huddersfield, HD1 3DH, UK

20
21
22 ***Corresponding author:**

23 Email: lynsey.harper2@gmail.com

25 **Running title:** impact assessment using molecular tools

26
27

Abstract

Fishes stocked for recreation and angling can damage freshwater habitats and negatively impact

biodiversity. The pond-associated crucian carp (*Carassius carassius*) is rare across its native

European range and has been stocked for conservation management in England despite plausible

non-native status. However, its impacts on English pond biota have not been broadly assessed.

Freshwater invertebrates comprise a large proportion of aquatic biodiversity, encompassing many

rare and endemic species, but small size and high abundance complicates their assessment.

Practitioners have typically employed sweep-netting and kick-net sampling with microscopy, but

specimen size/quality and experience can bias identification. DNA and eDNA metabarcoding now

offer alternative means of invertebrate assessment. We compared invertebrate diversity in ponds

(*N* = 18) with and without crucian carp using sweep-netting and microscopy, DNA metabarcoding,

and eDNA metabarcoding. Five 2 L water samples and 4 minute sweep-net samples were collected

at each pond. Inventories produced by morphotaxonomic identification of netted samples, DNA

metabarcoding of bulk tissue samples, and eDNA metabarcoding of water samples were compared.

Alpha diversity was greatest with DNA or eDNA metabarcoding, depending on whether methods

were considered as standard or with potential biases removed. DNA metabarcoding reflected

morphotaxonomic identification, whereas eDNA metabarcoding produced markedly different

communities. Therefore, these complementary tools should be combined for comprehensive

freshwater invertebrate assessment. Crucian carp presence minimally reduced alpha diversity in

ponds, but positively influenced beta diversity through taxon turnover (i.e. ponds with crucian carp

contained different invertebrate taxa to fishless ponds). Concerning pond invertebrates, the crucian

carp can be considered a naturalised, non-impactful species in England, supporting continued

conservation management in this region. Our results show that molecular tools can enhance

freshwater invertebrate assessment and facilitate development of more accurate and ecologically

effective pond management strategies.

Key words: community DNA, environmental DNA (eDNA), invertebrates, metabarcoding,

morphotaxonomic identification, monitoring

1. Introduction

Freshwater ecosystems comprise <1% of the Earth's surface, but represent major biodiversity hotspots and provide vital ecosystem services (Dudgeon et al., 2006). Ponds in particular provide critical habitat for biodiversity in a fragmented landscape (Céréghino, Biggs, Oertli, & Declerck, 2008), supporting many rare, unique or specialist species not found in other waterbodies (Biggs, von Fumetti, & Kelly-Quinn, 2016; Wood, Greenwood, & Agnew, 2003). These highly diverse and species-rich ecosystems contribute considerably to regional-scale diversity due to high habitat heterogeneity and associated species turnover (Davies et al., 2008; Williams et al., 2003). Aquatic invertebrates are a crucial and abundant component of this diversity, and occupy the vast range of microhabitats made available in ponds by their broad-ranging physicochemical properties and habitat complexity (Davies et al., 2008; Williams et al., 2003; Wood et al., 2003).

In Europe, lakes and ponds are commonly stocked with fish for angling and recreation, despite often having negative effects on invertebrates and amphibians (Gledhill, James, & Davies, 2008; Wood, Greenwood, Barker, & Gunn, 2001). Fish can alter community structure (Schilling, Loftin, & Huryn, 2009a, 2009b; Wood et al., 2001), reduce diversity (Lemmens et al., 2013; Wood et al., 2001), and reduce abundance and biomass (Marklund, Sandsten, Hansson, & Blindow, 2002; Schilling et al., 2009a) of invertebrates. These effects may manifest through direct predation by fish, altered water quality and loss of macrophyte diversity via foraging activity of fish, or management practices associated with angling activity (Lemmens et al., 2013; Maceda-Veiga, López, & Green, 2017; Schilling et al., 2009a; Wood et al., 2001). However, the impact of fish stocking can be negligible or even beneficial to invertebrate diversity, particularly at regional-scale, provided that fish species are carefully selected and managed (Gee, Smith, Lee, & Griffiths, 1997; Hassall, Hollinshead, & Hull, 2011; Lemmens et al., 2013; Stefanoudis et al., 2017).

The crucian carp (*Carassius carassius*) is one of few fishes strongly associated with small ponds in Europe, but has suffered heavy declines and local extinctions in the last century (Copp & Sayer, 2010; Sayer et al., 2011) due to habitat loss, species displacement by the invasive

goldfish (*Carassius auratus*) and gibel carp (*Carassius gibelio*) (Copp, Warrington, & Wesley,

2008; Sayer et al., 2011), and genetic introgression through hybridisation with goldfish and

common carp (*Cyprinus carpio*) (Hänfling, Bolton, Harley, & Carvalho, 2005). In 2010, the

crucian carp was designated as a Biodiversity Action Plan (BAP) species in the county of Norfolk,

eastern England (Copp & Sayer, 2010). A key objective of this BAP is to increase the number of

viable crucian carp populations across Norfolk through pond restoration and species

reintroduction. Many Norfolk ponds have since been stocked with crucian carp to realise this

objective (Sayer et al., *in review*), but continued stocking is controversial in light of genetic

evidence that indicates the crucian carp is not native to the UK (Jeffries et al., 2017).

Nonetheless, there is support for UK conservation efforts to continue to protect the genetic

integrity of the crucian carp at the European level, and provide a natural stronghold for the

species (Harper et al., 2019; Jeffries et al., 2017; Stefanoudis et al., 2017) in the face of

persistent declines throughout its native range of Northwest and Central Europe (Copp et al.,

2008; Sayer et al., 2011).

The impact of stocking crucian carp on lentic biodiversity has been largely ignored, and

little is known about interactions between this benthic fish and other pond species. Existing

research suggests crucian carp are characteristic of ponds rich in invertebrates with extensive

macrophyte cover (Copp et al., 2008; Sayer et al., 2011), and play an important ecological role

(along with other pond-associated fishes) by increasing landscape-scale diversity across pond

networks (Stefanoudis et al., 2017). Yet to our knowledge, only one study has assessed

biodiversity (specifically macrophytes, zooplankton, and water beetles) in ponds with and

without pond-associated fish in the UK (Stefanoudis et al., 2017), and no studies have

specifically focused on crucian carp. Consequently, there is a need to survey and compare

fishless ponds to ponds stocked with crucian carp to assess the impact of this species on

invertebrate diversity more broadly. However, obtaining species resolution data for

invertebrates is complicated by the level of taxonomic expertise needed for accurate

morphotaxonomic identification as well as the cost and time required to survey pond networks

(Briers & Biggs, 2003; Haase, Pauls, Schindehütte, & Sundermann, 2010; Hill et al., 2018).

Metabarcoding offers a rapid, high-resolution, cost-effective approach to biodiversity

assessment, where entire communities can be identified using High-Throughput Sequencing (HTS) in conjunction with community DNA from bulk tissue samples (DNA metabarcoding), or environmental DNA (eDNA) from environmental samples (eDNA metabarcoding), such as soil or water (Deiner et al., 2017; Taberlet, Coissac, Pompanon, Brochmann, & Willerslev, 2012). DNA metabarcoding of aquatic invertebrate samples has proven relatively successful, with applications in biomonitoring (Andújar et al., 2017; Elbrecht, Vamos, Meissner, Aroviita, & Leese, 2017a; Emilson et al., 2017; Lobo, Shokralla, Costa, Hajibabaei, & Costa, 2017). Use of eDNA metabarcoding for invertebrate assessment in freshwater rivers (Blackman et al., 2017; Carew, Kellar, Pettigrove, & Hoffmann, 2018; Deiner, Fronhofer, Mächler, Walser, & Altermatt, 2016; Klymus, Marshall, & Stepien, 2017), streams (Macher et al., 2018), and lakes (Klymus et al., 2017) is also gaining traction, but there are currently no published studies that have used metabarcoding for pond invertebrates.

We assessed invertebrate diversity in UK ponds with and without crucian carp using metabarcoding in conjunction with morphotaxonomic identification. The entire taxon inventories generated by each monitoring tool (standard methods) were compared to evaluate which tool provides the most holistic assessment of invertebrate diversity. We also compared the taxon inventories produced by each method when potential biases were removed, i.e. taxa with no reference sequences and microinvertebrates that would not be captured by a 1 mm mesh net (unbiased methods). The effect of crucian carp presence on invertebrate diversity was then determined using the taxon inventories from the standard or unbiased methods individually and combined. We hypothesised that DNA metabarcoding and morphotaxonomic identification would produce congruent invertebrate communities, whereas eDNA metabarcoding would reveal taxa not identified by these tools. DNA and eDNA metabarcoding were expected to enable species resolution data for problematic taxa that cannot be morphotaxonomically identified to species-level using standard keys. We hypothesised that alpha diversity would be lower in ponds with crucian carp, but beta diversity would be enhanced due to heterogeneity induced by crucian carp presence in the pond network. This pattern was expected regardless of the monitoring tool used. We provide recommendations for the application of molecular tools to freshwater invertebrate assessment as well as impact

assessment and conservation management of the non-native crucian carp alongside pond

biodiversity.

2. Methods

2.1 Study sites

We surveyed nine ponds containing crucian carp across Norfolk (*n* = 6) and East Yorkshire (*n* =

3), and nine fishless ponds in Norfolk (Fig. S1). The fish status of ponds was determined from

prior fyke-net surveys (Sayer et al., 2011; Sayer et al., *in review*). Roach (*Rutilus rutilus*) and

rudd (*Scardinius erythrophthalmus*) were the only other fishes that occurred in three of the

Norfolk ponds, but at lower density. All study ponds were selected to be similar in terms of

morphometry and habitat structure, being <1 ha in area, <5 m in depth, macrophyte-

dominated, with a largely open-canopy and thus minimal shading (≤20%) of the water surface.

Ponds were mainly surrounded by arable fields, excluding one pond located in woodland.

Samples for morphotaxonomic identification and metabarcoding were collected in autumn

2016, a time when invertebrate diversity often peaks (Hill, Sayer, & Wood, 2016). Data on

physical (area, depth, estimated percentages of perimeter with emergent vegetation, emergent

macrophyte cover, submerged macrophyte cover, and shading) and chemical (conductivity

measured with a HACH HQ30d meter) properties of ponds were collected between May and

August from 2010 to 2017.

2.2 Sweep-netting and morphotaxonomic identification

Sweep-netting was performed in accordance with the UK National Pond Survey methodology

(Biggs, Fox, & Nicolet, 1998), using a 1 mm mesh long-handled net (0.3 m square bag), to

generate a conventional taxonomic inventory of lentic macroinvertebrates. Sampling time at

each pond comprised a 3 min sweep-net sample and 1 min hand search (see Supporting

Information). Samples were deposited in a 1.2 L sterile Whirl-Pak® stand-up bag (Cole-Palmer,

London) and stored at -20 $^{\circ}$C until laboratory processing and sorting. Samples were thawed and

passed through sieves of 8 mm, 2 mm, and 0.25 mm in succession to remove large items of

vegetation and detrital matter. For each sample, invertebrate specimens were sorted into three

body size categories on laminated millimetre graph paper: small (<2.5 x 5 mm), medium (2.5 x 5

mm up to 5 x 10 mm), and large (>10 mm and up to 10 x 20 mm) (Elbrecht, Peinert, & Leese,

2017b). Where possible, size-sorted specimens were identified under a light microscope to

species-level using Freshwater Biological Association publications (Bass, 1998; Brooks & Cham,

2014; Edington & Hildrew, 1995; Elliott, 2009; Elliott & Humpesch, 2010; Elliott & Dobson,

2015; Friday, 1988; Gledhill, Sutcliffe, & Williams, 1993; Macan, 1960; Savage, 1989; Wallace,

Wallace, & Philipson, 1990). Terrestrial taxa and empty cases/shells were discarded. The

laminated paper was sterilised with 50% v/v chlorine-based commercial bleach solution (Elliot

Hygiene Ltd, UK) and 80% v/v ethanol solution between measuring specimens from different

ponds to minimise cross-contamination risk. Specimens were preserved in sterile 15 mL falcon

tubes (SARSTED, Germany) containing 100% ethanol according to size category and pond

sampled, and stored at -20 $^{\circ}$C until DNA extraction.

2.3 DNA samples

We followed the DNA extraction procedure from Blackman *et al.* (2017). Size categories from

each pond were dried overnight on FisherBrand cellulose filter paper (Fisher Scientific UK Ltd,

UK) in sterile glass funnels and conical flasks to remove excess ethanol. Size categories were

lysed (3 × 30 sec) using a TissueLyser II (Qiagen®, Germany) with DigiSol (50 mM Tris, 20 M

EDTA, 120 mM NaCl and 1% SDS). The adapter sets held 1.5 g of dried tissue and corresponding

volume of DigiSol. If the dry tissue weight of any size category exceeded 1.5 g, we processed

the size category in batches until all tissue had been lysed. The lysates from all batches were

pooled to recreate size categories. The size categories were incubated overnight at 55 °C with

SDS and Proteinase K (Bioline®, UK), following which 200 μL of lysate from each size category

was used for extraction with the DNeasy Blood & Tissue Kit (Qiagen®, Germany) according to

the manufacturer's protocol. Consequently, 16 bulk tissue samples were each represented by

three DNA extracts and two bulk tissue samples each represented by two DNA extracts that

were stored at -20 °C until metabarcoding.

2.4 eDNA samples

We repurposed eDNA samples from Norfolk ponds that were collected by Harper et al. (2019)

to validate a quantitative PCR (qPCR) assay for crucian carp. Additional samples were taken

from East Yorkshire ponds for the present study. Details of water sample collection, filtration,

and DNA extraction are provided in the Supporting Information. Briefly, five 2 L surface water

samples were collected at equidistant points around the pond perimeter where access

permitted. For each pond, a full process blank (1 L molecular grade water) was taken into the

field, transported, filtered, and extracted alongside samples to identify contamination. Filters

were stored at -20 °C until DNA extraction with the PowerWater® DNA Isolation Kit (MO BIO

Laboratories, CA, USA) following the manufacturer's protocol. eDNA extracts were stored at -20

°C until metabarcoding.

2.5 Metabarcoding workflow

Our metabarcoding workflow is fully described in the Supporting Information. In short, a

comprehensive list of UK invertebrate species living in or associated with freshwater habitats

was used to create custom, curated reference databases for UK aquatic invertebrates,

excluding Diptera (Appendix 1). Species without database representation are listed in Table S1

(see also Figs. S2, S3). Published metazoan primers mICOIintF (Leray et al., 2013) and

jgHCO2198 (Geller, Meyer, Parker, & Hawk, 2013), which amplify a 313 bp fragment of the

mitochondrial cytochrome c oxidase subunit I (*COI*) gene, were validated *in silico* using ecoPCR

software (Ficetola et al., 2010) against the reference databases. Parameters set allowed a 250-

350 bp fragment and maximum of three mismatches between each primer and each reference

sequence (Appendix 2). Primers were validated *in vitro* against 38 invertebrate species,

representing 38 families and 10 major groups, and compared to published macroinvertebrate

metabarcoding primers (Elbrecht & Leese, 2017; Vamos, Elbrecht, & Leese, 2017) (Figs. S4, S5,

S6).

PCR conditions were optimised (Figs. S7, S8) before two independent libraries were

constructed for DNA and eDNA metabarcoding using a two-step PCR protocol (Appendix 1).

During the first PCR (performed in triplicate), the target region was amplified using

oligonucleotides comprising the locus primers, sequencing primers, and pre-adapters. DNA

from the exotic, terrestrial two-spotted assassin bug (*Platymeris biguttatus*) was used for PCR

positive controls (tissue DNA *n* = 9, eDNA *n* = 11) as this species is not found in the UK, whilst

sterile molecular grade water (Fisher Scientific UK Ltd, UK) substituted template DNA for PCR

negative controls (tissue DNA *n* = 9, eDNA *n* = 11). PCR products were individually purified using

a magnetic bead clean-up (VWR International Ltd, UK), following a double size selection

protocol from Bronner *et al*. (2009). The second PCR bound Multiplex Identification (MID) tags

to the purified products. PCR products were pooled according to PCR run and the pooled PCR

product purified using a magnetic bead clean-up, following a double size selection protocol

from Bronner *et al*. (2009). Each purified PCR product was quantified on a Qubit™ 3.0

fluorometer using a Qubit™ dsDNA HS Assay Kit (Invitrogen, UK) and normalised according to

concentration and sample number to produce a pooled volume of 20 µL.

The pooled libraries were quantified by qPCR using the NEBNext® Library Quant Kit for

Illumina® (New England Biolabs® Inc., MA, USA). An Agilent 2200 TapeStation and High

Sensitivity D1000 ScreenTape (Agilent Technologies, CA, USA) were used to verify removal of

secondary products from libraries by bead purification and that a fragment of the expected size

(531 bp) remained. A total of 52 bulk tissue subsamples sequenced in triplicate (*n* = 156), 12

extraction blanks, and 18 PCR controls were included alongside samples from other projects (*N*

= 188) in the DNA library. A total of 90 eDNA samples, 18 full process blanks, and 22 PCR

controls were included alongside samples from other projects (*N* = 140) in the eDNA library.

The DNA and eDNA libraries were sequenced on an Illumina® MiSeq using 2 x 300 bp V3

chemistry with 10% and 20% PhiX Sequencing Control respectively (Illumina, Inc, CA, USA).

Raw sequences were converted to taxonomic assignments using metaBEAT v0.97.11, a

custom pipeline for reproducible analysis of metabarcoding data (https://github.com/HullUni-
bioinformatics/metaBEAT). After quality filtering, trimming, merging, chimera detection, and
clustering, non-redundant query sequences were compared against our reference databases
using BLAST (Zhang, Schwartz, Wagner, & Miller, 2000). Putative taxonomic identity was
assigned using a lowest common ancestor (LCA) approach based on the top 10% BLAST matches
for any query matching with at least 90% identity to a reference sequence across more than
80% of its length. Unassigned sequences were subjected to a separate BLAST against the
complete NCBI nucleotide (nt) database at 90% identity to determine the source via LCA as
described above. Bioinformatic settings (Appendix 1) were chosen based on comprehensive
exploration of the parameter space and comparison of metaBEAT taxonomic assignments to
morphotaxonomic inventories. Reference databases in GenBank format and the bioinformatic
analysis have been deposited in a dedicated GitHub repository, which has been permanently
archived for reproducibility (https://doi.org/10.5281/zenodo.3733127).

2.6 Data analysis

All analyses were performed in R v.3.4.3 (R Core Team, 2017), with data and R scripts deposited
in the GitHub repository. Manipulation of the DNA and eDNA metabarcoding datasets for
analysis is described in the Supporting Information. For each method, two presence-absence
datasets (species-level and family-level) were generated using the *decostand* function in vegan
v2.4-6 (Oksanen et al., 2018) for downstream analyses. This is because potential amplification
bias during PCR can prevent reliable abundance or biomass estimation from sequence reads
produced by DNA or eDNA metabarcoding (Elbrecht et al., 2017b). The entire species and
family inventories produced by each method (hereafter standard methods) were compared.
We also compared inventories from each method after removing potential biases: taxa without
reference sequences were removed from the morphotaxonomic dataset, and
microinvertebrates (i.e. taxa that would escape a 1 mm mesh net) were removed from the
metabarcoding datasets (hereafter unbiased methods). The effect of crucian carp on
invertebrate diversity was then assessed at species and family-level according to standard

methods (main text) and unbiased methods (Supporting Information) individually and

combined. Finally, the impact of crucian carp alongside environmental variables on invertebrate

diversity was investigated using the combined standard method (main text) and unbiased

method (Supporting Information) datasets.

We define alpha diversity as taxon richness within individual ponds, and beta diversity

as the difference between communities present at each pond whilst accounting for taxon

identity (Baselga & Orme, 2012). For each standard or unbiased dataset, the following analyses

were performed. Alpha diversity (response variable) was obtained using the *specnumber*

function in vegan v2.4-6 and modelled against sampling method (explanatory variable), then

crucian carp presence-absence (explanatory variable). Using the combined method datasets,

alpha diversity of the major invertebrate groups, i.e. orders, classes, or phyla (Dobson, Pawley,

Fletcher, & Powell, 2012), in each pond (response variable), was modelled against crucian carp

presence-absence (explanatory variable). All alpha diversity comparisons were made using

negative binomial Generalized Linear Models (GLMs), and Pairwise Tukey's HSD tests were used

to assess significance of differences. We used betapart v1.5.0 (Baselga & Orme, 2012) to

estimate total beta diversity, partitioned by nestedness-resultant and turnover, across sampling

methods and ponds with the *beta.multi* function. These three beta diversity components were

estimated for inventories produced by each sampling method, and ponds with or without

crucian carp using the *beta.pair* function (pairwise beta diversity scores). For each beta diversity

component, we compared community heterogeneity in each group (sampling method or

crucian carp presence-absence) by calculating homogeneity of multivariate dispersions

(MVDISP) using the *betadisper* function from vegan v2.4-6. Variation in MVDISP was statistically

tested using an analysis of variance (ANOVA), and pairwise Tukey's HSD tests used to determine

if there were significant differences between the groups. The effect of sampling method and

crucian carp on each beta diversity component was visualised using Non-metric

Multidimensional Scaling (NMDS) ordination plots with the *metaMDS* function, and tested

statistically using permutational multivariate analysis of variance (PERMANOVA) with the

adonis function in vegan v2.4-6. Pre-defined cut-off values were used for effect size, where

PERMANOVA results were interpreted as moderate and strong effects if $R^2 > 0.09$ and $R^2 > 0.25$

respectively. These values are broadly equivalent to correlation coefficients of r = 0.3 and 0.5

which represent moderate and strong effects accordingly (Macher et al., 2018; Nakagawa &

Cuthill, 2007). Jaccard dissimilarity was used as a measure of beta diversity for all analyses.

We tested whether the species and family-level invertebrate communities produced by

the combined standard or unbiased methods were influenced by pond properties in

conjunction with crucian carp presence-absence. Redundancy Analysis (RDA) was selected for

constrained ordination as it analyses variation in biological communities in response to

explanatory variables (Legendre & Legendre, 2012). Principal Coordinate Analysis (PCoA) was

performed using the *pcoa* function in ape v5.0 (Paradis & Schliep, 2018) on the turnover,

nestedness-resultant, and total beta diversity matrices. The Lingoes correction was employed

to account for negative eigenvalues (Legendre, 2014). The resultant PCoA eigenvectors

(principle coordinates) for each distance matrix were used as the response variable in variance

partitioning. Our variables were grouped as biotic (crucian carp presence-absence) or abiotic

(pond conductivity, area, depth, percentages of perimeter with emergent vegetation, emergent

macrophyte cover, submerged macrophyte cover, and shading). Abiotic variables were log10

transformed to eliminate their physical units (Legendre & Birks, 2012). Significant abiotic

variables influencing each component of beta diversity were identified using the *ordiR2step*

function in vegan v2.4-6 to perform separate RDA analyses under a forward selection

procedure. Where applicable, the relative contributions of the biotic and abiotic variables on

turnover, nestedness-resultant, and total beta diversity for the species and family-level

invertebrate communities were assessed by variance partitioning (Borcard, Legendre, &

Drapeau, 1992) using the *varpart* function from vegan v2.4-6. For each beta diversity

component, RDA was performed using biotic and significant abiotic variables, and variance

partitioning used to divide the total percentage of variation explained into unique and shared

contributions for biotic and abiotic predictor groups. The *anova* function in vegan v2.4-6 was

used to examine the statistical significance of the full model and the unique contributions of

each predictor group. We report the adjusted R^2-fractions as they are widely recommended

and unbiased (Peres-Neto, Legendre, Dray, & Borcard, 2006).

3. Results

3.1 Comparison of methods for freshwater invertebrate assessment

Summaries of taxon detection by standard and unbiased methods are reported in the Supporting Information (Appendix 2; Tables S2, S3, S4, S5; Figs. S9, S10). Briefly, 1,404 specimens belonging to 91 species and 2,281 specimens belonging 38 families were identified by microscopy. With DNA metabarcoding, 2,454,206 and 2,907,021 sequence reads were assigned to 140 species and 56 families. With eDNA metabarcoding, 389,766 and 831,073 sequence reads were assigned to 160 species and 92 families respectively. Only 17 species (Figs. 1ai, bi) and 21 families (Figs. 1aii, bii) were shared by all three methods (standard or unbiased). eDNA metabarcoding detected the most unique species and families, whereas DNA metabarcoding and morphotaxonomic identification were more comparable (Fig. 1). The number of taxa that morphotaxonomic identification shared with DNA and eDNA metabarcoding was unchanged using standard or unbiased methods. Morphotaxonomic identification shared one family with eDNA metabarcoding, but 36 species and 13 families with DNA metabarcoding (Fig. 1). Morphotaxonomic identification detected 19 species without reference sequences, whereas DNA and eDNA metabarcoding detected 4 and 57 microinvertebrate species respectively (Fig. 1a). The metabarcoding approaches either shared 49 species and 18 families (standard methods, Fig. 1a) or 29 species and 7 families (unbiased methods, Fig. 1b).

Standard sampling method influenced alpha diversity of ponds at species (GLM χ^2_2 = 36.999, P < 0.001) and family-level (GLM χ^2_2 = 54.800, P < 0.001). Significant differences between the alpha diversity means for morphotaxonomic identification and DNA metabarcoding (species-level -0.452 ± 0.131, Z = -3.446, P = 0.002; family-level -0.295 ± 0.109, Z = -2.719, P = 0.018), morphotaxonomic identification and eDNA metabarcoding (species-level -0.779 ± 0.128, Z = -6.069, P < 0.001; family-level -0.729 ± 0.102, Z = -7.122, P < 0.001), and DNA and eDNA metabarcoding (species-level 0.327 ± 0.123, Z = 2.671, P = 0.021; family-level 0.434 ±

0.096, Z = 4.519, P < 0.001) were observed. Conversely, unbiased sampling method influenced

alpha diversity of ponds at species-level (GLM χ^2_2 = 20.767, P < 0.001), but not family-level

(GLM χ^2_2 = 3.832, P = 0.147). Significant differences between the species-level alpha diversity

means of morphotaxonomic identification and DNA metabarcoding (species-level -0.443 ±

0.126, Z = -3.506, P = 0.001; family-level -0.134 ± 0.106, Z = -1.267, P = 0.414),

morphotaxonomic identification and eDNA metabarcoding (species-level 0.086 ± 0.134, Z =

0.639, P = 0.798; family-level 0.072 ± 0.110, Z = 0.650, P = 0.792), and DNA and eDNA

metabarcoding (species-level -0.529 ± 0.128, Z = -4.132, P < 0.001; family-level -0.206 ± 0.108, Z

= -1.915, P = 0.134) were observed. With standard methods, species and family-level alpha

diversity was highest using eDNA metabarcoding, followed by DNA metabarcoding then

morphotaxonomic identification (Figs. 2ai, aii). With unbiased methods, DNA metabarcoding

detected more species than morphotaxonomic identification and eDNA metabarcoding, but

alpha diversity was comparable at family-level (Figs. 2bi, bii). Regarding beta diversity, MVDISP

only significantly differed between unbiased methods for family-level turnover and total beta

diversity (Table S6). Both standard and unbiased sampling method had moderate and strong

positive effects on turnover and total beta diversity at species (Figs. 3ai, aiii, ci, ciii) and family-

level (Figs. 3aii, aiv, cii, civ) respectively, but not nestedness-resultant (Table S7).

3.2 Impact of crucian carp stocking on pond invertebrates

Here, we report the impact of crucian carp stocking assessed using standard methods as these

provide the most comprehensive evaluation of invertebrate diversity. Crucian carp impact

assessment using unbiased methods is reported in the Supporting Information (Appendix 2;

Tables S9, S10, S11, S12; Figs. S13, S14, S15, S16). Independently and combined, standard

methods showed alpha diversity of invertebrates to be marginally reduced in ponds containing

crucian carp at species (Figs. 4ai-iv) and family-level (Figs. 4bi-iv), but not significantly so (P >

0.05, Table 1). Detailed examination of alpha diversity within the major invertebrate groups

identified by all three methods combined revealed that Coleoptera and Mollusca diversity was

significantly reduced in ponds with crucian carp at species-level (GLM χ^2_{24} = 30.029, P = 0.184;

Coleoptera -0.517 ± 0.255, Z = -2.025, P = 0.043; Mollusca -0.815 ± 0.268, Z = -3.046, P = 0.002),

but not family-level (GLM χ^2_{24} = 1.503, P = 0.220; Coleoptera -0.511 ± -0.298, Z = -1.713, P =

0.087; Mollusca -0.442 ± -0.302, Z = -1.462, P = 0.144). However, differences in alpha diversity

between ponds with or without crucian carp were not significant for other invertebrate groups

at either taxonomic rank (Fig. S11). Alpha diversity relationships held true for the unbiased

methods (Table S9; Fig. S13, S14).

Total beta diversity of ponds was consistently high at species and family-level for

independent and combined methods (standard or unbiased). Variation in invertebrate

community composition was driven by turnover rather than nestedness-resultant (Tables 2,

S10). MVDISP differed between ponds for species-level nestedness-resultant with

morphotaxonomic identification, species-level total beta diversity with eDNA metabarcoding,

and species and family-level total beta diversity with all methods combined. Bar nestedness-

resultant with morphotaxonomic identification, fishless ponds had significantly lower

dispersion than ponds with crucian carp (Table S8). MVDISP patterns were not consistent across

standard and unbiased methods (Table S11). At species-level, morphotaxonomic identification,

eDNA metabarcoding, and all methods combined revealed a weak or moderate positive

influence of crucian carp presence on turnover (Table 3; Figs. 5ai, iii, iv) and total beta diversity

(Table 3; Figs. 5ci, iii, iv) between ponds, but not nestedness-resultant (Table 3; Figs. 5bi, iii, iv).

At family-level, only eDNA metabarcoding revealed a weak or moderate positive influence of

crucian carp presence on turnover (Table 3; Fig. S12aiii) and total beta diversity (Table 3; Fig.

S12ciii) between ponds, but not nestedness-resultant (Table 3; Fig. S12biii). Crucian carp

presence did not influence any beta diversity component with DNA metabarcoding (Table 3;

Figs. 5aii, bii, cii). Less insights on beta diversity were obtained using unbiased methods (Table

S12; Figs. S15, S16) than standard methods.

Additional analyses undertaken on the combined method dataset supported an effect of

crucian carp presence and excluded the influence of abiotic variables on invertebrate diversity.

At species-level, only pond area was identified as a significant abiotic variable for turnover and

total beta diversity by forward selection. Variance partitioning analysis was undertaken for

turnover and total beta diversity using crucian carp presence-absence and pond area. Using the

adjusted R^2 values, biotic and abiotic variables explained 6.30% and 4.33% of the total variation

in species turnover and total beta diversity respectively (Fig. 6). Crucian carp presence-absence

significantly contributed to turnover (Fig. 6a; adjusted R^2 = 5.15%, F_1 = 1.947, P = 0.001) and

total beta diversity (Fig. 6b; adjusted R^2 = 3.79%, F_1 = 1.679, P = 0.001), whereas pond area

explained less variance and did not influence turnover (Fig. 6a; adjusted R^2 = 2.57%, F_1 = 1.197,

P = 0.170) or total beta diversity (Fig. 6b; adjusted R^2 = 1.72%, F_1 = 1.090, P = 0.276). RDA of

nestedness-resultant without abiotic data indicated no influence of crucian carp presence-

absence (F_1 = 0.364, P = 0.844). At family-level, forward selection did not identify any significant

abiotic variables for turnover, nestedness-resultant, or total beta diversity. Therefore, variance

partitioning was not undertaken for any component of beta diversity. RDA of each beta

diversity component minus abiotic data revealed crucian carp presence-absence influenced

total beta diversity (F_1 = 1.430, P = 0.047), but not turnover (F_1 = 1.410, P = 0.111) or

nestedness-resultant (F_1 = 1.793, P = 0.178). Variance partitioning results using the unbiased

methods are presented in Appendix 2.

4. Discussion

4.1 Comparison of methods for freshwater invertebrate assessment

DNA (unbiased methods) or eDNA (standard methods) metabarcoding generated the highest

species and family-level alpha diversity. Differences in alpha diversity were more pronounced

between standard methods, but unique species were detected nonetheless. Morphotaxonomic

identification and DNA metabarcoding produced more similar communities, and performed

best for Coleoptera, Hemiptera, Hirudinea, Megaloptera, and Odonata. eDNA metabarcoding

generated a markedly different community, and detected aquatic taxa overlooked in pond net

samples (Arachnida, Crustacea, Lepidoptera, Psocoptera, Trichoptera) alongside singleton

terrestrial taxa (Hymenoptera, Thysanoptera). Both metabarcoding approaches performed well

for Annelida, Collembola, Diptera, and Ephemeroptera, particularly at species-level.

Critically, we used a 1 mm mesh net which is designed to sample macroinvertebrates,

whereas a 500 µm mesh net would also sample the microinvertebrates detected by

metabarcoding. Microinvertebrates may inflate metabarcoding diversity estimates against

morphotaxonomic identification, but may also prevent PCR amplification of

macroinvertebrates. To mitigate bias, we performed all analyses excluding microinvertebrates

not captured by netting and excluding morphotaxonomically identified taxa without reference

sequences that metabarcoding would not recover. Alpha diversity results from the standard

and unbiased method comparisons were variable, but beta diversity results were more

consistent. Both method comparisons echo other studies where metabarcoding captured

greater diversity than morphotaxonomic approaches and resolved problematic groups (e.g.

Diptera) for species-level morphotaxonomic identification (Andújar et al., 2017; Carew et al.,

2018; Clarke, Beard, Swadling, & Deagle, 2017; Elbrecht et al., 2017a; Emilson et al., 2017;

Klymus et al., 2017; Lobo et al., 2017). Standard and unbiased methods broadly revealed the

same ecological relationships, although standard methods may yield more insights. All three

methods will be most powerful when used in combination for ecological investigation.

Each method has inherent biases that influence performance for freshwater

invertebrate assessment. The UK National Pond Survey methodology recommends that pond

net samples are placed into buckets/zip-lock bags, followed by sorting and identification in the

laboratory as soon as possible. Samples are typically preserved in ethanol or refrigerated to

prolong processing time (Biggs et al., 1998), but this standard methodology does not lend itself

to molecular applications. After collection, samples were transported on ice then frozen at -20

°C to prevent predation within samples, minimise organismal decay and subsequent DNA

degradation, and allow samples to be processed as required in the laboratory. Using this

strategy, small or inconspicuous dead specimens would not have been recovered during sorting

(Biggs et al., 1998) and therefore excluded from morphotaxonomic and DNA metabarcoding

inventories. These losses are surplus to the 29% of specimens already overlooked during sorting

due to smaller body size (Haase et al., 2010). Sorting may also damage or completely destroy

some recovered specimens, influencing morphotaxonomic identification (Lobo et al., 2017;

Zizka, Leese, Peinert, & Geiger, 2018).

Human error is another source of discrepancy between morphotaxonomic identification

and metabarcoding. Some taxa were identified to genus or family-level and omitted from our

species-level inventories, whereas other taxa may have been falsely identified and inventoried

(Carew, Pettigrove, Metzeling, & Hoffmann, 2013; Elbrecht et al., 2017a; Haase et al., 2010).

Losses incurred by the preservation and sorting treatments for morphotaxonomic identification

cannot be mitigated, and human error can only be reduced with taxonomic expertise.

Conversely, taxon recovery by DNA metabarcoding can be improved with protocols that

preserve and extract DNA from intact, unsorted samples (Elbrecht et al., 2017a). For example,

ethanol used to preserve specimens (Zizka et al., 2018), or replacement of ethanol with DNA

extraction buffer followed by incubation then specimen removal (Carew, Coleman, &

Hoffmann, 2018), offer alternative starting materials to bulk tissue for DNA metabarcoding.

However, these alternative DNA sources tend to produce false negatives for schlerotised

groups, such as Coleoptera and Trichoptera (Carew et al., 2018; Zizka et al., 2018).

Despite successful application of DNA metabarcoding to bulk tissue samples for

biomonitoring (Andújar et al., 2017; Carew et al., 2018; Elbrecht et al., 2017a; Emilson et al.,

2017; Macher et al., 2018), recurring issues have been encountered. Arguably, the most

pressing issue is size bias, where DNA from large and/or high biomass taxa can outcompete

DNA of smaller and/or low biomass taxa during PCR amplification and sequencing (Elbrecht et

al., 2017b). Although we used the size-sorting approach of Elbrecht *et al*. (2017b) and

sequenced body size categories independently with data pooling downstream, DNA

metabarcoding failed to detect some taxa that have reference sequences and were reliably

identified by microscopy, including several Coleopterans (*Agabus sturmii*, *Hydroporus*

*erythrocephalu*s, *Rhantus frontalis*, *Haliplus confinis*, *Haliplus ruficollis*), a small Mollusc

(*Gyraulus crista*), a medium Hirudinean (*Erpobdella lineata*), and two large Anisopterans

(*Aeshna mixta*, *Anax imperator*). Non-delivery of desired results is problematic as size-sorting is

time-consuming, labour-intensive, potentially increases cross-contamination risk between

samples (Elbrecht et al., 2017b), and still requires samples to be sorted from vegetation and

substrate (Elbrecht et al., 2017a). Size-sorting may therefore be a drain on resources and time

allocated to DNA metabarcoding projects, but other sources of false negatives must be

excluded.

In contrast, eDNA metabarcoding preferentially amplifies DNA from small, floating
organisms (Deiner et al., 2016; Macher et al., 2018), and these organisms are often retained on
filter membranes used for eDNA capture (*pers. obs.*). Consequently, their DNA is abundant in
samples and overwhelms DNA from other taxa during sequencing. Pre-filtering with or only
using large pore size filter membranes may reduce this bias (Macher et al., 2018). eDNA
metabarcoding may also detect different taxa due to species variability in eDNA production and
shedding rates. The species infrequently detected by eDNA metabarcoding were those that
possess thicker exoskeletons composed of chitin and occasionally calcium carbonate, e.g.
Coleoptera, Hemiptera. Exoskeletons may restrict the release of DNA into the water column
(Tréguier et al., 2014) as opposed to organisms that are filter-feeders or produce slime (e.g.
Crustacea, Mollusca), ectoparasites (i.e. Acari) that feed on blood or skin of other species, or
use external versus internal fertilisation (Wacker et al., 2019). Furthermore, species may differ
in their habitat preferences and utilisation, potentially resulting in highly localised distributions
(Klymus et al., 2017). Therefore, more samples or greater volumes may be required to improve
detection of pond biota (Harper et al., 2019).

Marker choice and primer design can substantially influence amplification success and
taxonomic assignment. Although the *COI* region offers species resolution, has extensive
database representation, and is used as standard in DNA barcoding (Elbrecht, Hebert, &
Steinke, 2018), it lacks conserved primer-binding sites as a protein-coding gene (Clarke et al.,
2017). This introduces high risk of primer mismatch and subsequent bias during metabarcoding
(Clarke et al., 2017; Elbrecht et al., 2018), where preferential DNA amplification for some taxa
may prevent recovery of all taxa present (Elbrecht et al., 2016; Lobo et al., 2017). Consequently,
most metabarcoding primers designed to target the *COI* region in metazoans (Elbrecht & Leese,
2017; Leray et al., 2013; Meusnier et al., 2008; Zeale, Butlin, Barker, Lees, & Jones, 2011) are
degenerate to allow primers to bind at highly variable sites. However, highly degenerate
primers may bind non-target regions (Elbrecht et al., 2018) and experience biased amplification
toward non-metazoans, e.g. bacteria, algae, fungi (Brandon-Mong et al., 2015; Macher et al.,
2018). This bias can occur even when primers are designed for macroinvertebrates (Elbrecht &

Leese, 2017a; Elbrecht et al., 2017b), and is more pronounced in sequences obtained from

eDNA samples (Macher et al., 2018). Invertebrate metabarcoding may benefit from the use of

different (e.g. 16S ribosomal RNA) or multiple markers (Deiner et al., 2016; Elbrecht et al.,

2016). Unfortunately, alternative markers often provide less taxonomic resolution due to lack

of reference database representation (Clarke et al., 2017; Elbrecht et al., 2016). Use of multiple

markers will also increase PCR and sequencing costs, preventing metabarcoding uptake by

freshwater monitoring schemes, but these are expected to subside (Elbrecht et al., 2017a).

We obtained an excessive number (~60%) of unassigned reads from eDNA

metabarcoding, despite using a relaxed BLAST identity (90%) against custom and public

reference databases. This suggests that sequences were of poor quality, could not be assigned

to reference sequences, or lacked reference database representation (Macher et al., 2018).

Elbrecht *et al*. (2018) demonstrated that degenerate primers like mlCOIintF can experience

primer slippage and produce variable length sequences when they bind to low diversity DNA

regions. This differs across species with implications for bioinformatic processing and

taxonomic assignment. Public reference database records may be few, mislabelled, or have

limited geographic coverage (Curry, Gibson, Shokralla, Hajibabaei, & Baird, 2018; Elbrecht et al.,

2017a; Klymus et al., 2017). We morphotaxonomically identified 19 species (Hemiptera,

Mollusca, Odonata) without reference sequences whose DNA may have been sequenced, but

would not have been taxonomically assigned to species-level. Future studies must develop

more specific primers that have binding and flanking regions of high nucleotide diversity

(Elbrecht et al., 2018) and/or target particular invertebrate orders or families (Klymus et al.,

2017) as well as procure reference sequences for different markers (Curry et al., 2018; Elbrecht

et al., 2016). These are essential steps to improve the reliability and accuracy of molecular

monitoring in freshwater ecosystems.

4.2 Impact of crucian carp stocking on pond invertebrates

Crucian carp had a negligible influence on alpha diversity and a positive influence on beta

diversity of pond invertebrates. Alpha diversity in ponds with crucian carp was marginally

reduced compared to alpha diversity in fishless ponds, but this difference was not significant
across methods used at either taxonomic rank. Within the major invertebrate groups identified
by all methods combined, species-level alpha diversity of Coleoptera and Mollusca was
significantly reduced in ponds containing crucian carp as opposed to fishless ponds. This may be
because Coleopterans are large, active swimmers that are visible and easy to hunt, and
Molluscs are slow-moving prey. Across ponds, total beta diversity of invertebrate communities
was driven by turnover (taxon replacement) rather than nestedness-resultant (taxon subsets)
(Baselga & Orme, 2012). Our results revealed crucian carp to positively influence turnover and
total beta diversity between ponds. Therefore, taxa in fishless ponds were replaced by different
taxa in ponds with crucian carp, resulting in dissimilar community composition. Removal of top
predators and dominant species, such as Coleopterans, by crucian carp may allow other prey
species to colonise but further research is needed.

Our results both echo and contradict Stefanoudis *et al*. (2017), where the presence of
fish (including crucian carp) in ponds altered macrophyte and cladoceran community
composition, but not water beetle composition. Hassall *et al*. (2011) also found that fish
presence in ponds had a positive effect on species richness of most invertebrate taxa, but a
negative effect on Coleoptera. Gee *et al*. (1997) observed no influence of fish stocking on
macrophyte and macroinvertebrate species richness, although Odonata richness was lower and
Trichoptera richness higher in stocked ponds. Conversely, other research found that
managed/stocked ponds, some of which contained crucian carp, had lower invertebrate
diversity than unmanaged sites, which were characterised by Trichoptera, Coleoptera and
Zygoptera larvae (Wood et al., 2001). Similarly, large, active and free-swimming taxa
(Notonectidae, Corixidae, Gyrinidae, Dytiscidae, Aeshnidae, Libellulidae and Chaoboridae) were
strongly associated with fish absence as well as more diverse and abundant in fishless lakes
(Bendell & McNicol, 1995; Schilling et al., 2009a, 2009b).

Critically, only two of the aforementioned studies accounted for the identity of fish
species assemblages present (Stefanoudis et al., 2017; Wood et al., 2001) and just one was
based on fish surveys (Stefanoudis et al., 2017), whereas other studies only considered the
presence of a particular species (Schilling et al., 2009a) or fish presence-absence generally

(Bendell & McNicol, 1995; Gee et al., 1997; Hassall et al., 2011; Schilling et al., 2009b). The

contrasting results produced by these studies and our own indicate that the impact of fish on

pond biodiversity will likely depend on the species present, population density, and

management strategy. Wetland fishes vary in dietary preference and consume different

proportions of invertebrate taxa, thus different fishes will suppress numbers of and confer

benefits to different invertebrate taxa (Batzer, Pusateri, & Vetter, 2000). Invasive fish species

may be more detrimental than non-invasive species, for example, the mosquitofish (*Gambusia*

affinis) reduced zooplankton abundance and macroinvertebrate density by 90% and 50%

respectively after introduction in a wetland ecosystem experiment (Preston et al., 2017).

Additionally, duration of stocking was found to substantially reduce invertebrate species

richness and abundance (Schilling et al., 2009a). Therefore, local- and regional-scale diversity

may benefit most from pond mosaics that maintain fish-free and stocked (at low-moderate

densities) ponds to prevent increases in fish biomass and predation pressure over time

(Lemmens et al., 2013).

The effects of environmental variables on alpha and beta diversity of pond invertebrates

must also be considered (Hassall et al., 2011; Hill, Heino, Thornhill, Ryves, & Wood, 2017). Our

study ponds were selected to have similar physical and chemical properties, which may explain

the minimal contribution of abiotic factors to variance in invertebrate community structure.

Although pond area was retained by model selection, it did not significantly influence

community structure and explained less variance than crucian carp presence-absence. Likewise,

other studies have shown a weak or no effect of pond area on invertebrate species richness and

community composition (Gee et al., 1997; Gledhill et al., 2008; Oertli et al., 2002; Wood et al.,

2001). Critically, our environmental data were collected over a 7-year period, whereas

contemporary data may have better explained variance in community structure. We omitted

variables that experience high temporal variation from our analyses, but these may be linked to

the Coleoptera and Mollusca alpha diversity reductions and different invertebrate communities

observed in crucian carp ponds, such as water temperature, pH, nutrient concentration, and

surface dissolved oxygen (Bendell & McNicol, 1995; Hassall et al., 2011; Menetrey, Sager, Oertli,

& Lachavanne, 2005). Investigations examining contemporary physicochemical variables

alongside crucian carp presence-absence will help disentangle the effects of stocking and

habitat. The degree of connectivity between ponds may also substantially influence

invertebrate community structure, both independently and in combination with crucian carp

presence (Hill et al., 2017).

Crucian carp might be assumed to have negative impacts on pond biodiversity by having

"carp" in its name. Common carp foraging activities are known to reduce invertebrate density

and macrophyte cover, with knock-on effects for waterfowl species richness and abundance

(Haas et al., 2007; Maceda-Veiga et al., 2017). However, all of our crucian carp ponds were

dominated by floating and/or submerged macrophyte beds as in Stefanoudis *et al*. (2017), and

we observed no evidence for negative impacts on vegetation. Our study demonstrates that

crucian carp can be an important, positive driver of invertebrate community heterogeneity and

subsequently beta diversity in ponds. Our findings indicate that crucian carp presence enhances

invertebrate diversity across pond networks, and that current conservation management of

stocked populations in England is appropriate. English waters are absent of gibel carp and

subsequently hybridisation with this species that is threatening the crucian carp across Europe,

thus England is an important epicentre for crucian carp conservation (Jeffries et al., 2017; Sayer

et al., *in review*). Crucian carp may contribute to taxon turnover and subsequent community

heterogeneity in ponds by providing colonisation opportunities for smaller invertebrates

through predator (e.g. Coleopterans) removal. We conclude that management strategies to

support high landscape-scale pond biodiversity should consider a patchwork of fish-containing

and fishless ponds, with the crucian carp likely to be an important species in this respect.

However, the impact of crucian carp on pond biodiversity, especially amphibians, must be

studied more broadly with respect to crucian carp population density and the aforementioned

environmental variables to determine whether stocking is invariably beneficial.

Acknowledgements

This work was funded by the University of Hull. We thank the many landowners who allowed

access to ponds. We are grateful to the Norfolk Crucian Carp Project for providing data

associated with Norfolk ponds. We thank Michael Lee and Matt Smith (Environment Agency)

for selecting East Yorkshire ponds and providing associated data, and Jane Colley (Askham

Bryan College) for providing invertebrate samples from these ponds. We are grateful to

Jianlong Li and Cristina Di Muri (University of Hull) for assistance with water sampling and

filtration as well as advice on library preparation and sequencing.

Data accessibility

Raw sequence reads have been archived on the NCBI Sequence Read Archive (Study:

SRP163672; BioProject: PRJNA494857; BioSamples: SAMN10181701 - SAMN10182084 [bulk

tissue DNA] and SAMN10187732 - SAMN10188115 [eDNA]; SRA accessions: SRR7969394 -

SRR796977 [bulk tissue DNA] and SRR7985814 - SRR7986197 [eDNA]). Jupyter notebooks, R

scripts and corresponding data are deposited in a dedicated GitHub repository

(https://github.com/HullUni-

bioinformatics/Harper_et_al_2020_crucian_carp_impact_invertebrates_conventional_molecul

ar_tools) which has been permanently archived (https://doi.org/10.5281/zenodo.3733127).

Author contributions

L.R.H, B.H, and L.L.H conceived and designed the study. C.D.S selected the study ponds in

Norfolk and provided associated environmental data. L.R.H, M.B, and R.C.B built the custom

reference sequence database for invertebrates. L.R.H and R.C.B sampled the Norfolk ponds for

invertebrates. L.R.H and M.B collected and filtered water samples from all study ponds. D.S.R

assisted with water filtration of eDNA samples and provided advice on primers, PCR protocols,

and sequencing. L.R.H. performed laboratory work and analysed the data. L.R.H. wrote the

manuscript, which all authors revised.

References

Andújar, C., Arribas, P., Gray, C., Bruce, C., Woodward, G., Yu, D. W., & Vogler, A. P. (2017). Metabarcoding of freshwater invertebrates to detect the effects of a pesticide spill. *Molecular Ecology*, *27*, 146–166. https://doi.org/10.1111/mec.14410

Baselga, A., & Orme, C. D. L. (2012). betapart: an R package for the study of beta diversity : Betapart package. *Methods in Ecology and Evolution*, *3*, 808–812. https://doi.org/10.1111/j.2041-210X.2012.00224.x

Bass, J. (1998). *Last-instar larvae and pupae of the Simuliidae of Britain and Ireland: A key with brief ecological notes* (Vol. 18). Ambleside, Freshwater Biological Association.

Batzer, D. P., Pusateri, C. R., & Vetter, R. (2000). IMPACTS OF FISH PREDATION ON MARSH INVERTEBRATES: DIRECT AND INDIRECT EFFECTS. *Wetlands*, *20*, 307–312. https://doi.org/10.1672/0277-5212(2000)020[0307:IOFPOM]2.0.CO;2

Bendell, B. E., & McNicol, D. K. (1995). Lake acidity, fish predation, and the distribution and abundance of some littoral insects. *Hydrobiologia*, *302*, 133–145. https://doi.org/10.1007/BF00027038

Biggs, J., Fox, G., & Nicolet, P. (1998). *A guide to the methods of the National Pond Survey*. Oxford, Pond Action.

Biggs, J., von Fumetti, S., & Kelly-Quinn, M. (2016). The importance of small waterbodies for biodiversity and ecosystem services: implications for policy makers. *Hydrobiologia*, *793*, 3–39. https://doi.org/10.1007/s10750-016-3007-0

Blackman, R. C., Constable, D., Hahn, C., Sheard, A. M., Durkota, J., Hänfling, B., & Lawson Handley, L. (2017). Detection of a new non-native freshwater species by DNA metabarcoding of environmental samples--first record of *Gammarus fossarum* in the UK. *Aquatic Invasions*, *12*, 177–189. https://doi.org/10.3391/ai.2017.12.2.06

Borcard, D., Legendre, P., & Drapeau, P. (1992). Partialling out the Spatial Component of Ecological Variation. *Ecology*, *73*, 1045–1055. https://doi.org/10.2307/1940179

Brandon-Mong, G.-J., Gan, H.-M., Sing, K.-W., Lee, P.-S., Lim, P.-E., & Wilson, J.-J. (2015). DNA metabarcoding of insects and allies: an evaluation of primers and pipelines. *Bulletin of Entomological Research*, *105*, 717–727. https://doi.org/10.1017/S0007485315000681

Briers, R. A., & Biggs, J. (2003). Indicator taxa for the conservation of pond invertebrate diversity. *Aquatic Conservation: Marine and Freshwater Ecosystems*, *13*, 323–330. https://doi.org/10.1002/aqc.576

Bronner, I. F., Quail, M. A., Turner, D. J., & Swerdlow, H. (2009). Improved Protocols for Illumina Sequencing. *Current Protocols in Human Genetics*, *18*, 18.2.1–18.2.42. https://doi.org/10.1002/0471142905.hg1802s79

Brooks, S., & Cham, S. (2014). *Field Guide to the Dragonflies & Damselflies of Great Britain and Ireland*. London, Bloomsbury Publishing Plc.

Carew, M. E., Coleman, R. A., & Hoffmann, A. A. (2018). Can non-destructive DNA extraction of bulk invertebrate samples be used for metabarcoding? *PeerJ*, *6*, e4980. https://doi.org/10.7717/peerj.4980

Carew, M. E., Kellar, C. R., Pettigrove, V. J., & Hoffmann, A. A. (2018). Can high-throughput
sequencing detect macroinvertebrate diversity for routine monitoring of an urban river?
Ecological Indicators, *85*, 440–450. https://doi.org/10.1016/j.ecolind.2017.11.002
Carew, M. E., Pettigrove, V. J., Metzeling, L., & Hoffmann, A. A. (2013). Environmental
monitoring using next generation sequencing: rapid identification of macroinvertebrate
bioindicator species. *Frontiers in Zoology*, *10*, 45. https://doi.org/10.1186/1742-9994-10-45
Céréghino, R., Biggs, J., Oertli, B., & Declerck, S. (2008). The ecology of European ponds:
defining the characteristics of a neglected freshwater habitat. *Hydrobiologia*, *597*, 1–6.
https://doi.org/10.1007/s10750-007-9225-8
Clarke, L. J., Beard, J. M., Swadling, K. M., & Deagle, B. E. (2017). Effect of marker choice and
thermal cycling protocol on zooplankton DNA metabarcoding studies. *Ecology and Evolution*,
7, 873–883. https://doi.org/10.1002/ece3.2667
Copp, G. H., & Sayer, C. D. (2010). *Norfolk Biodiversity Action Plan--Local Species Action Plan for*
Crucian Carp (Carassius carassius). Lowestoft, Fisheries & Aquaculture Science.
Copp, G. H., Warrington, S., & Wesley, K. J. (2008). Management of an ornamental pond as a
conservation site for a threatened native fish species, crucian carp Carassius carassius.
Hydrobiologia, *597*, 149–155. https://doi.org/10.1007/978-90-481-9088-1_13
Curry, C. J., Gibson, J. F., Shokralla, S., Hajibabaei, M., & Baird, D. J. (2018). Identifying North
American freshwater invertebrates using DNA barcodes: are existing COI sequence libraries
fit for purpose? *Freshwater Science* , *37*, 178–189. https://doi.org/10.1086/696613
Davies, B., Biggs, J., Williams, P., Whitfield, M., Nicolet, P., Sear, D., Bray, S., & Maund, S. (2008).
Comparative biodiversity of aquatic habitats in the European agricultural landscape.
Agriculture, Ecosystems & Environment, *125*, 1–8.
https://doi.org/10.1016/j.agee.2007.10.006
Deiner, K., Bik, H. M., Mächler, E., Seymour, M., Lacoursière-Roussel, A., Altermatt, F., Creer, S.,
Bista, I., Lodge, D. M., de Vere, N., Pfrender, M. E., & Bernatchez, L. (2017). Environmental
DNA metabarcoding: Transforming how we survey animal and plant communities. *Molecular*
Ecology, *26*, 5872–5895. https://doi.org/10.1111/mec.14350
Deiner, K., Fronhofer, E. A., Mächler, E., Walser, J.-C., & Altermatt, F. (2016). Environmental
DNA reveals that rivers are conveyer belts of biodiversity information. *Nature*
Communications, *7*, 12544. https://doi.org/10.1038/ncomms12544
Dobson, M., Pawley, S., Fletcher, M., & Powell, A. (2012). *Guide to Freshwater Invertebrates*.
Ambleside, Freshwater Biological Association.
Dudgeon, D., Arthington, A. H., Gessner, M. O., Kawabata, Z. -I., Knowler, D. J., Lévêque, C.,
Naiman, R. J., Prieur-Richard, A. -H., Soto, D., Stiassny, M. L. J., & Sullivan, C. A. (2006).
Freshwater biodiversity: importance, threats, status and conservation challenges. *Biological*
Reviews of the Cambridge Philosophical Society, *81*, 163–182.
https://doi.org/10.1017/S1464793105006950
Edington, J. M., & Hildrew, A. G. (1995). *A revised key to the caseless caddis larvae of the British*
Isles with notes on their ecology. Ambleside, Freshwater Biological Association.
Elbrecht, V., Hebert, P. D. N., & Steinke, D. (2018). Slippage of degenerate primers can cause
variation in amplicon length. *Scientific Reports*, *8*, 10999. https://doi.org/10.1038/s41598-
018-29364-z

Elbrecht, V., & Leese, F. (2017). Validation and Development of COI Metabarcoding Primers for
Freshwater Macroinvertebrate Bioassessment. *Frontiers in Environmental Science*, *5*.
https://doi.org/10.3389/fenvs.2017.00011
Elbrecht, V., Peinert, B., & Leese, F. (2017). Sorting things out: Assessing effects of unequal
specimen biomass on DNA metabarcoding. *Ecology and Evolution*, *7*, 6918–6926.
https://doi.org/10.1002/ece3.3192
Elbrecht, V., Taberlet, P., Dejean, T., Valentini, A., Usseglio-Polatera, P., Beisel, J. -N., Coissac, E.,
Boyer, F., & Leese, F. (2016). Testing the potential of a ribosomal 16S marker for DNA
metabarcoding of insects. *PeerJ*, 4, e1966. https://doi.org/10.7717/peerj.1966
Elbrecht, V., Vamos, E.E., Meissner, K., Aroviita, J. & Leese, F. (2017). Assessing strengths and
weaknesses of DNA metabarcoding-based macroinvertebrate identification for routine
stream monitoring. *Methods in Ecology and Evolution*, *8*, 1265–1275.
https://doi.org/10.1111/2041-210X.12789
Elliott, J. M. (2009). *Freshwater Megaloptera and Neuroptera of Britain and Ireland: keys to*
adults and larvae, and a review of their ecology. Ambleside, Freshwater Biological
Association.
Elliott, J. M., & Dobson, M. (2015). *Freshwater leeches of Britain and Ireland: keys to the*
Hirudinea and a review of their ecology. Ambleside, Freshwater Biological Association.
Elliott, J. M., & Humpesch, U. H. (2010). *Mayfly larvae (Ephemeroptera) of Britain and Ireland:*
keys and a review of their ecology. Ambleside, Freshwater Biological Association.
Emilson, C.E., Thompson, D.G., Venier, L.A., Porter, T.M., Swystun, T., Chartrand, D., Capell, S.,
& Hajibabaei, M. (2017). DNA metabarcoding and morphological macroinvertebrate metrics
reveal the same changes in boreal watersheds across an environmental gradient. *Scientific*
Reports, *7*, 12777. https://doi.org/10.1038/s41598-017-13157-x
Ficetola, G.F., Coissac, E., Zundel, S., Riaz, T., Shehzad, W., Bessière, J., Taberlet, P., &
Pompanon, F. (2010). An *In silico* approach for the evaluation of DNA barcodes. *BMC*
Genomics, *11*, 434. https://doi.org/10.1186/1471-2164-11-434
Friday, L. E. (1988). A key to the adults of British water beetles. *Field Studies*, *7*, 1–151.
Gee, J. H. R., Smith, B. D., Lee, K. M., & Griffiths, S. W. (1997). The ecological basis of freshwater
pond management for biodiversity. *Aquatic Conservation: Marine and Freshwater*
Ecosystems, *7*, 91–104. https://doi.org/10.1002/(SICI)1099-0755(199706)7:2<91::AID-
AQC221>3.0.CO;2-O
Geller, J., Meyer, C., Parker, M., & Hawk, H. (2013). Redesign of PCR primers for mitochondrial
cytochrome c oxidase subunit I for marine invertebrates and application in all-taxa biotic
surveys. *Molecular Ecology Resources*, *13*, 851–861. https://doi.org/10.1111/1755-
0998.12138
Gledhill, D. G., James, P., & Davies, D. H. (2008). Pond density as a determinant of aquatic
species richness in an urban landscape. *Landscape Ecology*, *23*, 1219–1230.
https://doi.org/10.1007/s10980-008-9292-x
Gledhill, T., Sutcliffe, D. W., & Williams, W. D. (1993). *British freshwater Crustacea*
Malacostraca: a key with ecological notes. Ambleside, Freshwater Biological Association.

Haase, P., Pauls, S. U., Schindehütte, K., & Sundermann, A. (2010). First audit of
macroinvertebrate samples from an EU Water Framework Directive monitoring program:
human error greatly lowers precision of assessment results. *Journal of the North American*
Benthological Society, *29*, 1279–1291. https://doi.org/10.1899/09-183.1

Haas, K., Köhler, U., Diehl, S., Köhler, P., Dietrich, S., Holler, S., Jaensch, A., Niedermaier, M., &
Vilsmeier, J. (2007). Influence of fish on habitat choice of water birds: a whole system
experiment. *Ecology*, *88*, 2915–2925. https://doi.org/10.1890/06-1981.1

Hänfling, B., Bolton, P., Harley, M., & Carvalho, G. R. (2005). A molecular approach to detect
hybridisation between crucian carp (*Carassius carassius*) and non-indigenous carp species
(*Carassius* spp. and *Cyprinus carpio*). *Freshwater Biology*, *50*, 403–417.
https://doi.org/10.1111/j.1365-2427.2004.01330.x

Harper, L. R., Griffiths, N. P., Lawson Handley, L., Sayer, C. D., Read, D. S., Harper, K. J.,
Blackman, R. C., Li, J., & Hänfling, B. (2019). Development and application of environmental
DNA surveillance for the threatened crucian carp (*Carassius carassius*). *Freshwater Biology*,
64, 93–107. https://doi.org/10.1111/fwb.13197

Hassall, C., Hollinshead, J., & Hull, A. (2011). Environmental correlates of plant and invertebrate
species richness in ponds. *Biodiversity and Conservation*, *20*, 3189–3222.
https://doi.org/10.1007/s10531-011-0142-9

Hill, M. J., Hassall, C., Oertli, B., Fahrig, L., Robson, B. J., Biggs, J., Samways, M. J., Usio, N.,
Takamura, N., Krishnaswamy, J., & Wood, P. J. (2018). New policy directions for global pond
conservation. *Conservation Letters*, *142*, e12447. https://doi.org/10.1111/conl.12447

Hill, M. J., Heino, J., Thornhill, I., Ryves, D. B., & Wood, P. J. (2017). Effects of dispersal mode on
the environmental and spatial correlates of nestedness and species turnover in pond
communities. *Oikos*, *126*, 1575–1585. https://doi.org/10.1111/oik.04266

Hill, M. J., Sayer, C. D., & Wood, P. J. (2016). When is the best time to sample aquatic
macroinvertebrates in ponds for biodiversity assessment? *Environmental Monitoring and*
Assessment, *188*, 194. https://doi.org/10.1007/s10661-016-5178-6

Jeffries, D. L., Copp, G. H., Maes, G. E., Lawson Handley, L., Sayer, C. D., & Hänfling, B. (2017).
Genetic evidence challenges the native status of a threatened freshwater fish (*Carassius*
carassius) in England. *Ecology and Evolution*, *7*, 2871–2882.
https://doi.org/10.1002/ece3.2831

Klymus, K. E., Marshall, N. T., & Stepien, C. A. (2017). Environmental DNA (eDNA)
metabarcoding assays to detect invasive invertebrate species in the Great Lakes. *PLoS ONE*,
12, e0177643. https://doi.org/10.1371/journal.pone.0177643

Legendre, P. (2014). Interpreting the replacement and richness difference components of beta
diversity. *Global Ecology and Biogeography: A Journal of Macroecology*, *23*, 1324–
1334. https://doi.org/10.1111/geb.12207

Legendre P., & Birks H. J. B. (2012). From Classical to Canonical Ordination. In: Birks H.,
Lotter A., Juggins S., Smol J. (eds) Tracking Environmental Change Using Lake Sediments.
Developments in Paleoenvironmental Research, vol 5. Dordrecht, Springer.
https://doi.org/10.1007/978-94-007-2745-8_8

Legendre, P., & Legendre, L. F. J. (2012). *Numerical Ecology*. The Netherlands, Elsevier Science.

Lemmens, P., Mergeay, J., De Bie, T., Van Wichelen, J., De Meester, L., & Declerck, S. A. J.
(2013). How to maximally support local and regional biodiversity in applied conservation?
Insights from pond management. *PLoS ONE*, *8*, e72538.
https://doi.org/10.1371/journal.pone.0072538
Leray, M., Yang, J. Y., Meyer, C. P., Mills, S. C., Agudelo, N., Ranwez, V., Boehm, J. T., & Machida,
R. J. (2013). A new versatile primer set targeting a short fragment of the mitochondrial COI
region for metabarcoding metazoan diversity: application for characterizing coral reef fish
gut contents. *Frontiers in Zoology*, *10*, 34. https://doi.org/10.1186/1742-9994-10-34
Lobo, J., Shokralla, S., Costa, M. H., Hajibabaei, M., & Costa, F. O. (2017). DNA metabarcoding
for high-throughput monitoring of estuarine macrobenthic communities. *Scientific Reports*,
7, 15618. https://doi.org/10.1038/s41598-017-15823-6
Macan, T. T. (1960). *A key to the British fresh-and brackish-water gastropods: with notes on
their ecology*. Ambleside, Freshwater Biological Association.
Maceda-Veiga, A., López, R., & Green, A. J. (2017). Dramatic impact of alien carp *Cyprinus carpio*
on globally threatened diving ducks and other waterbirds in Mediterranean shallow lakes.
Biological Conservation, *212*, 74–85. https://doi.org/10.1016/j.biocon.2017.06.002
Macher, J. -N., Vivancos, A., Piggott, J. J., Centeno, F. C., Matthaei, C. D., & Leese, F. (2018).
Comparison of environmental DNA and bulk-sample metabarcoding using highly degenerate
COI primers. *Molecular Ecology Resources*, *18*, 1456–1468. https://doi.org/10.1111/1755-
0998.12940
Marklund, O., Sandsten, H., Hansson, L. -A., & Blindow, I. (2002). Effects of waterfowl and fish
on submerged vegetation and macroinvertebrates. *Freshwater Biology*, *47*, 2049–2059.
https://doi.org/10.1046/j.1365-2427.2002.00949.x
Menetrey, N., Sager, L., Oertli, B., & Lachavanne, J. -B. (2005). Looking for metrics to assess the
trophic state of ponds. Macroinvertebrates and amphibians. *Aquatic Conservation: Marine
and Freshwater Ecosystems*, *15*, 653–664. https://doi.org/10.1002/aqc.746
Meusnier, I., Singer, G. A. C., Landry, J. -F., Hickey, D. A., Hebert, P. D. N., & Hajibabaei, M.
(2008). A universal DNA mini-barcode for biodiversity analysis. *BMC Genomics*, *9*, 214.
https://doi.org/10.1186/1471-2164-9-214
Nakagawa, S., & Cuthill, I. C. (2007). Effect size, confidence interval and statistical significance: a
practical guide for biologists. *Biological Reviews of the Cambridge Philosophical Society*, *82*,
591–605. https://doi.org/10.1111/j.1469-185X.2007.00027.x
Oertli, B., Joye, D. A., Castella, E., Juge, R., Cambin, D., & Lachavanne, J.-B. (2002). Does size
matter? The relationship between pond area and biodiversity. *Biological Conservation*, *104*,
59–70. https://doi.org/10.1016/S0006-3207(01)00154-9
Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R.,
O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., & Wagner, H. (2011)
vegan: Community Ecology Package. R package version 2.5-4. https://CRAN.R-
project.org/package=vegan
Paradis, E., & Schliep, K. (2018). ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics*, *35*, 526–528.
https://doi.org/10.1093/bioinformatics/bty633

Peres-Neto, P. R., Legendre, P., Dray, S., & Borcard, D. (2006). Variation partitioning of species
data matrices: estimation and comparison of fractions. *Ecology*, *87*, 2614–2625.
https://doi.org/10.1890/0012-9658(2006)87[2614:VPOSDM]2.0.CO;2
Preston, D. L., Hedman, H. D., Esfahani, E. R., Pena, E. M., Boland, C. E., Lunde, K. B., & Johnson,
P. T. J. (2017). Responses of a wetland ecosystem to the controlled introduction of invasive
fish. *Freshwater Biology*, *62*, 767–778. https://doi.org/10.1111/fwb.12900
R Core Team (2017). R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. https://www.R-project.org/
Savage, A. A. (1989). *Adults of the British aquatic Hemiptera Heteroptera: a key with ecological
notes*. Ambleside, Freshwater Biological Association.
Sayer, C. D., Copp, G. H., Emson, D., Godard, M. J., Zięba, G., & Wesley, K. J. (2011). Towards the
conservation of crucian carp *Carassius carassius*: understanding the extent and causes of
decline within part of its native English range. *Journal of Fish Biology*, *79*, 1608–
1624. https://doi.org/10.1111/j.1095-8649.2011.03059.x
Schilling, E. G., Loftin, C. S., & Huryn, A. D. (2009a). Effects of introduced fish on
macroinvertebrate communities in historically fishless headwater and kettle lakes. *Biological
Conservation*, *142*, 3030–3038. https://doi.org/10.1016/j.biocon.2009.08.003
Schilling, E. G., Loftin, C. S., & Huryn, A. D. (2009b). Macroinvertebrates as indicators of fish
absence in naturally fishless lakes. *Freshwater Biology*, *54*, 181–202.
https://doi.org/10.1111/j.1365-2427.2008.02096
Stefanoudis, P. V., Sayer, C. D., Greaves, H. M., Davidson, T. A., Robson, H., Almeida, D., &
Smith, E. (2017). Consequences of fish for cladoceran, water beetle and macrophyte
communities in a farmland pond landscape: implications for conservation. *Fundamental and
Applied Limnology*, *190*, 141–156. https://doi.org/10.1127/fal/2017/1004
Taberlet, P., Coissac, E., Pompanon, F., Brochmann, C., & Willerslev, E. (2012). Towards next-
generation biodiversity assessment using DNA metabarcoding. *Molecular Ecology*, *21*, 2045–
2050. https://doi.org/10.1111/j.1365-294X.2012.05470.x
Tréguier, A., Paillisson, J.-M., Dejean, T., Valentini, A., Schlaepfer, M. A., & Roussel, J.-M. (2014).
Environmental DNA surveillance for invertebrate species: advantages and technical
limitations to detect invasive crayfish *Procambarus clarkii* in freshwater ponds. *Journal of
Applied Ecology*, *51*, 871–879. https://doi.org/10.1111/1365-2664.12262
Vamos, E.E., Elbrecht, V., & Leese, F. (2017). Short COI markers for freshwater
macroinvertebrate metabarcoding. *Metabarcoding and Metagenomics*, *1*, e14625.
https://doi.org/10.3897/mbmg.1.14625
Wacker, S., Fossøy, F., Larsen, B. M., Brandsegg, H., Sivertsgård, R., & Karlsson, S. (2019).
Downstream transport and seasonal variation in freshwater pearl mussel (*Margaritifera
margaritifera*) eDNA concentration. *Environmental DNA*, *1*, 64–73.
https://doi.org/10.1002/edn3.10
Wallace, I. D., Wallace, B., & Philipson, G. N. (1990). *A key to the case-bearing caddis larvae of
Britain and Ireland*. Ambleside, Freshwater Biological Association.
Williams, P., Whitfield, M., Biggs, J., Bray, S., Fox, G., Nicolet, P., & Sear, D. (2003). Comparative
biodiversity of rivers, streams, ditches and ponds in an agricultural landscape in Southern
England. *Biological Conservation*, *115*, 329–341. https://doi.org/10.1016/S0006-
3207(03)00153-8

Wood, P. J., Greenwood, M. T., & Agnew, M. D. (2003). Pond biodiversity and habitat loss in the
UK. *Area*, *35*, 206–216. https://doi.org/10.1111/1475-4762.00249
Wood, P. J., Greenwood, M. T., Barker, S. A., & Gunn, J. (2001). The effects of amenity
management for angling on the conservation value of aquatic invertebrate communities in
old industrial ponds. *Biological Conservation*, *102*, 17–29. https://doi.org/10.1016/S0006-
3207(01)00087-8
Zeale, M. R. K., Butlin, R. K., Barker, G. L. A., Lees, D. C., & Jones, G. (2011). Taxon-specific PCR
for DNA barcoding arthropod prey in bat faeces. *Molecular Ecology Resources*, *11*, 236–244.
https://doi.org/10.1111/j.1755-0998.2010.02920.x
Zhang, Z., Schwartz, S., Wagner, L., & Miller, W. (2000). A greedy algorithm for aligning DNA
sequences. *Journal of Computational Biology*, *7*, 203–214.
https://doi.org/10.1089/10665270050081478
Zizka, V. M. A., Leese, F., Peinert, B., & Geiger, M. F. (2018). DNA metabarcoding from sample
fixative as a quick and voucher preserving biodiversity assessment method. *Genome*, *62*,
122–136. https://doi.org/10.1139/gen-2018-0048
966

967 **Table 1.** Summary of analyses (GLM) statistically comparing alpha diversity at species-level and
968 family-level between ponds with and without crucian carp using independent and combined
969 standard methods.

		Species-level						Family-level				
		GLM			LRT			GLM			LRT	
	df	Estimate ± SE	Z	P	χ^2	P	df	Estimate ± SE	Z	P	χ^2	P
Sweep-netting and microscopy	1	-0.139 ± 0.186	-0.746	0.456	0.556	0.456	1	-0.098 ± 0.158	-0.620	0.535	0.385	0.535
DNA metabarcoding	1	0.047 ± 0.141	0.332	0.740	0.110	0.740	1	-0.015 ± 0.121	-0.121	0.904	0.015	0.904
eDNA metabarcoding	1	-0.045 ± 0.198	-0.227	0.820	0.052	0.820	1	0.147 ± 0.141	-1.043	0.297	1.089	0.297
Combined methods	1	-0.049 ± 0.121	-0.409	0.683	0.167	0.683	1	-0.117 ± 0.096	-1.225	0.220	1.503	0.220

971
972

973 **Table 2.** Relative contribution of taxon turnover and nestedness-resultant to total beta diversity
974 (Jaccard dissimilarity) using independent and combined standard methods. A value of 1
975 corresponds to all sites containing different species.
976

	Species-level			Family-level		
	Turnover	Nestedness-resultant	Total beta diversity	Turnover	Nestedness-resultant	Total beta diversity
Sweep-netting and microscopy	0.935 (98.01%)	0.019 (1.99%)	0.954 (100%)	0.867 (94.75%)	0.048 (5.25%)	0.915 (100%)
DNA metabarcoding	0.939 (98.63%)	0.013 (1.37%)	0.952 (100%)	0.885 (96.83%)	0.029 (3.17%)	0.914 (100%)
eDNA metabarcoding	0.917 (97.24%)	0.026 (2.76%)	0.943 (100%)	0.870 (95.29%)	0.043 (4.71%)	0.913 (100%)
Combined methods	0.927 (98.41%)	0.015 (1.59%)	0.942 (100%)	0.870 (96.35%)	0.033 (3.65%)	0.903 (100%)

978 **Table 3.** Summary of analyses (PERMANOVA) statistically examining variation in community
979 composition between ponds with and without crucian carp at species-level and family-level
980 using independent and combined standard methods.
981

| | | Community similarity (PERMANOVA) | | | | | | |
| | Species-level | | | | Family-level | | | |
	df	F	R^2	P	df	F	R^2	P
Sweep-netting and microscopy								
Turnover	1	1.673	0.095	**0.022**	1	1.198	0.070	0.371
Nestedness-resultant	1	-3.454	-0.275	0.954	1	-0.103	-0.007	0.649
Total beta diversity	1	1.369	0.079	**0.029**	1	1.136	0.066	0.311
DNA metabarcoding								
Turnover	1	1.271	0.074	0.154	1	1.668	0.094	0.106
Nestedness-resultant	1	-1.082	-0.073	0.748	1	-0.113	-0.007	0.731
Total beta diversity	1	1.121	0.066	0.224	1	1.259	0.073	0.213
eDNA metabarcoding								
Turnover	1	2.484	0.134	**0.002**	1	1.850	0.104	**0.033**
Nestedness-resultant	1	-2.136	-0.154	0.963	1	-0.015	-0.001	0.687
Total beta diversity	1	1.841	0.103	**0.003**	1	1.521	0.087	**0.032**
Combined methods								
Turnover	1	1.924	0.107	**0.001**	1	1.494	0.085	0.106
Nestedness-resultant	1	-1.449	-0.099	0.954	1	2.621	0.141	0.162
Total beta diversity	1	1.669	0.095	**0.003**	1	1.430	0.082	0.051

983
984
985 **Figure 1.** Venn diagram which summarises taxon detection by standard **(a)** and unbiased **(b)**
986 methods of invertebrate assessment. The number of invertebrate species **(i)** and families **(ii)**
987 detected across the 18 study ponds by sweep-netting and microscopy (green circle), DNA
988 metabarcoding (purple circle), and eDNA metabarcoding (orange circle) is displayed. Overlap in
989 species or family detections between methods is shown within circle intersections.
990

Figure 2. Mean alpha diversity of invertebrates in ponds across Norfolk and East Yorkshire. Alpha diversity is displayed according to standard **(a)** and unbiased **(b)** methods of invertebrate assessment. Species **(i)** and family **(ii)** richness generated by netting and microscopy (green points), DNA metabarcoding (purple points), and eDNA metabarcoding (orange points) is shown. Boxes show 25th, 50th, and 75th percentiles, and whiskers show 5th and 95th percentiles.

1000
1001

Figure 3. Non-metric Multidimensional Scaling (NMDS) plots of invertebrate communities (Jaccard dissimilarity) produced by sweep-netting and microscopy (green points/ellipse), DNA metabarcoding (purple points/ellipse), and eDNA metabarcoding (orange points/ellipse) for the 18 study ponds. The turnover **(a)** and nestedness-resultant **(b)** partitions of total beta diversity **(c)** are shown at species-level and family-level for standard **(i, ii)** and unbiased **(iii, iv)** methods.

Figure 4. Mean alpha diversity of invertebrates in ponds with crucian carp (blue points) and without fish (grey points) across Norfolk and East Yorkshire. Alpha diversity at species-level **(a)** and family-level **(b)** is shown according to standard methods of invertebrate assessment: sweep-netting and microscopy **(i)**, DNA metabarcoding **(ii)**, eDNA metabarcoding **(iii)**, and all methods combined **(iv)**. Boxes show 25th, 50th, and 75th percentiles, and whiskers show 5th and 95th percentiles.

(a) Turnover **(b) Nestedness** **(c) Total β Diversity**

Crucian carp — Absent — Present

Figure 5. Non-metric Multidimensional Scaling (NMDS) plots of species-level invertebrate communities (Jaccard dissimilarity) from ponds with crucian carp (blue points/ellipse) and without fish (grey points/ellipse) across Norfolk and East Yorkshire. The turnover **(a)** and nestedness-resultant **(b)** partitions of total beta diversity **(c)** are shown according to standard methods of invertebrate assessment: netting and microscopy **(i)**, DNA metabarcoding **(ii)**, eDNA metabarcoding **(iii)**, and all methods combined **(iv)**.

Figure 6. The relative contribution of biotic (crucian carp presence-absence) and abiotic variables (pond area) to species-level turnover **(a)** and total beta diversity **(b)** when the combined invertebrate data from all three standard sampling methods were considered. Values within circles and circle intersections represent the adjusted R^2 values. Significant variables are indicated by asterisks (* = $P < 0.05$).