1 **Assessing the impact of the threatened crucian carp (*Carassius***

2 ***carassius*) on pond invertebrate diversity - a comparison of**

3 **conventional and molecular tools**

5 **Lynsey R. Harper[1,2,3]*, Lori Lawson Handley[1], Carl D. Sayer[4], Daniel S. Read[5],**

6 **Marco Benucci[1], Rosetta C. Blackman[1,6,7], Matthew J. Hill[8], and Bernd Hänfling[1]**

7
8

[1] Department of Biological and Marine Sciences, University of Hull, Hull, HU6 7RX, UK

[2] Illinois Natural History Survey, Prairie Research Institute, University of Illinois at Urbana-Champaign, Champaign,

IL, USA

[3] School of Biological and Environmental Sciences, Liverpool John Moores University, L3 3AF, UK

[4] Pond Restoration Research Group, Environmental Change Research Centre, Department of Geography, University

College London, London, WC1E 6BT, UK

[5] Centre for Ecology & Hydrology (CEH), Benson Lane, Crowmarsh Gifford, Wallingford, Oxfordshire, OX10 8BB, UK

[6] Department of Aquatic Ecology, Eawag: Swiss Federal Institute of Aquatic Science and Technology, Dübendorf,

Switzerland

[7] Department of Evolutionary Biology and Environmental Studies, University of Zurich, Winterthurerstrasse 190,

8057 Zurich, Switzerland.

[8] School of Applied Sciences, University of Huddersfield, Queensgate, Huddersfield, HD1 3DH, UK

21
22

23 ***Corresponding author:**

24 Email: lynsey.harper2@gmail.com

26 **Running title:** Impact assessment using molecular tools

27
28

Abstract

Fishes stocked for recreation and angling can damage freshwater habitats and negatively impact biodiversity. The pond-associated crucian carp (*Carassius carassius*) is rare across Europe and stocked for conservation management in England, but impacts on pond biota are understudied. Freshwater invertebrates contribute substantially to aquatic biodiversity, encompassing many rare and endemic species, but their small size and high abundance complicates their assessment. Practitioners have employed sweep-netting and kick-sampling with microscopy (morphotaxonomy), but specimen size/quality and experience can bias identification. DNA and eDNA metabarcoding offer alternative means of invertebrate assessment. We compared invertebrate diversity in ponds (*N* = 18) with and without crucian carp using morphotaxonomic identification, DNA metabarcoding, and eDNA metabarcoding. Five 2-L water samples and 3-minute sweep-net samples were collected at each pond. Inventories produced by morphotaxonomic identification of netted samples, DNA metabarcoding of bulk tissue samples, and eDNA metabarcoding of water samples were compared. Alpha diversity was greatest with DNA or eDNA metabarcoding, depending on whether standard or unbiased methods were considered. DNA metabarcoding reflected morphotaxonomic identification, whereas eDNA metabarcoding produced markedly different communities. These complementary tools should be combined for comprehensive invertebrate assessment. Crucian carp presence minimally reduced alpha diversity in ponds, but positively influenced beta diversity through taxon turnover (i.e. ponds with crucian carp contained different invertebrates to fishless ponds). Crucian carp presence contributes to landscape-scale invertebrate diversity, supporting continued conservation management in England. Our results show that molecular tools can enhance freshwater invertebrate assessment and facilitate development of more accurate and ecologically effective pond management strategies.

Key words: community DNA, environmental DNA (eDNA), invertebrates, metabarcoding, morphotaxonomic identification, monitoring

1. Introduction

Freshwater ecosystems comprise <1% of the Earth's surface, but represent major biodiversity hotspots and provide vital ecosystem services (Dudgeon et al., 2006). Ponds (see definition in Biggs, von Fumetti, & Kelly-Quinn, 2016) in particular provide critical habitat for biodiversity in a fragmented landscape (Céréghino, Biggs, Oertli, & Declerck, 2008), supporting many rare and specialist species not found in other waterbodies (Biggs et al., 2016; Wood, Greenwood, & Agnew, 2003). These highly diverse and species-rich ecosystems contribute considerably to regional-scale diversity due to high habitat heterogeneity and associated species turnover (Davies et al., 2008; Williams et al., 2003). Aquatic invertebrates are a crucial and abundant component of this diversity, and occupy the vast range of microhabitats made available in ponds by their broad-ranging physicochemical properties and habitat complexity (Davies et al., 2008; Williams et al., 2003; Wood et al., 2003).

In Europe, lakes and ponds are commonly stocked with fish for angling and recreation purposes, despite often having negative effects on invertebrates and amphibians (Gledhill, James, & Davies, 2008; Wood, Greenwood, Barker, & Gunn, 2001). Fish can alter community structure (Schilling, Loftin, & Huryn, 2009a, 2009b; Wood et al., 2001), reduce diversity (Lemmens et al., 2013; Wood et al., 2001), and reduce the abundance and biomass (Marklund, Sandsten, Hansson, & Blindow, 2002; Schilling et al., 2009a) of invertebrates. These effects may manifest through direct predation by fish, altered water quality and loss of macrophyte diversity and cover via foraging activity of fish, or management practices (e.g. commercial farming, artificial feeding, pond depth and bank alteration, removal of aquatic vegetation and sediment) associated with angling activity (Lemmens et al., 2013; Maceda-Veiga, López, & Green, 2017; Schilling et al., 2009a; Wood et al., 2001). However, the impact of fish stocking can be negligible or even beneficial to invertebrate diversity, particularly at a regional-scale, provided that fish species are carefully selected and managed (Gee, Smith, Lee, & Griffiths, 1997; Hassall, Hollinshead, & Hull, 2011; Lemmens et al., 2013; Stefanoudis et al., 2017).

The crucian carp (*Carassius carassius*) is one of few fish species strongly associated with small ponds in Europe, but has suffered heavy declines and local extinctions in the last century

(Copp & Sayer, 2010; Sayer et al., 2011) due to habitat loss, species displacement by the

invasive goldfish (*Carassius auratus*) and gibel carp (*Carassius gibelio*) (Copp, Warrington, &

Wesley, 2008; Sayer et al., 2011), and genetic introgression through hybridisation with goldfish,

gibel carp, and common carp (*Cyprinus carpio*) (Hänfling, Bolton, Harley, & Carvalho, 2005). In

2010, the crucian carp was designated as a Biodiversity Action Plan (BAP) species in the county

of Norfolk in eastern England (Copp & Sayer, 2010). A key objective of this BAP is to increase

the number of viable crucian carp populations across Norfolk through pond restoration and

species reintroduction. Many Norfolk ponds have since been restored and stocked with crucian

carp to realise this objective (Sayer et al., 2020), but maintaining populations and continued

stocking requires justification in light of genetic evidence that indicates the crucian carp is not

native to the UK (Jeffries et al., 2017). Nonetheless, there is support for UK conservation efforts

to continue to protect the genetic integrity of the crucian carp at the European level, and

provide a natural stronghold for the species (Harper et al., 2019; Jeffries et al., 2017;

Stefanoudis et al., 2017) in the face of persistent declines throughout its native range of

Northwest and Central Europe (Copp et al., 2008; Sayer et al., 2011; Sayer et al., 2020).

The impact of stocking crucian carp on lentic biodiversity has been largely ignored, and

little is known about its interactions with other pond species. Existing research suggests crucian

carp are characteristic of ponds rich in invertebrates with extensive macrophyte cover (Copp et

al., 2008; Sayer et al., 2011), and play an important ecological role (along with other pond-

associated fishes) by increasing landscape-scale diversity across pond networks (Stefanoudis et

al., 2017). Yet to our knowledge, only one study has assessed biodiversity (specifically

macrophytes, zooplankton, and water beetles) in ponds with and without pond-associated

fishes in the UK (Stefanoudis et al., 2017), and no studies have specifically focused on crucian

carp. Consequently, there is a need to survey and compare fishless ponds to ponds containing

crucian carp to assess the impact of maintaining existing crucian carp populations and

conservation-based stocking of this species on invertebrate diversity more broadly.

Assessments of invertebrate diversity in pond networks have relied upon

morphotaxonomic identification using sweep-netting together with microscopy, but this

approach requires extensive resources and taxonomic expertise for accurate species-level

identification (Briers & Biggs, 2003; Haase, Pauls, Schindehütte, & Sundermann, 2010; Hill et al.,

2018). Metabarcoding offers a rapid, high-resolution, cost-effective approach to biodiversity

assessment, where entire communities can be identified using High-Throughput Sequencing

(HTS) in conjunction with community DNA from bulk tissue samples (DNA metabarcoding), or

environmental DNA (eDNA) from environmental samples (eDNA metabarcoding), such as soil or

water (Deiner et al., 2017; Taberlet, Coissac, Pompanon, Brochmann, & Willerslev, 2012). DNA

metabarcoding of aquatic invertebrate samples has proven relatively successful, with

applications in biomonitoring (Andújar et al., 2017; Elbrecht, Vamos, Meissner, Aroviita, &

Leese, 2017a; Emilson et al., 2017; Lobo, Shokralla, Costa, Hajibabaei, & Costa, 2017). Use of

eDNA metabarcoding for invertebrate assessment in freshwater rivers (Blackman et al., 2017;

Carew, Kellar, Pettigrove, & Hoffmann, 2018; Deiner, Fronhofer, Mächler, Walser, & Altermatt,

2016; Klymus, Marshall, & Stepien, 2017; Leese et al., 2020), streams (Macher et al., 2018), and

lakes (Klymus et al., 2017) is also gaining traction, but there are currently few published studies

that have used metabarcoding for small lake or pond invertebrates (Beentjes, Speksnijder,

Schilthuizen, Hoogeveen, & van der Hoorn, 2019).

In this study, we assessed invertebrate diversity in UK ponds with and without crucian

carp using metabarcoding in conjunction with morphotaxonomic identification. The entire

taxon inventories generated by each monitoring tool (standard methods) were compared to

evaluate which tool provided the most holistic assessment of invertebrate diversity. We also

compared the taxon inventories produced by each method when potential biases were

removed, i.e. taxa with no reference sequences and meiofauna that would not be captured by a

1 mm mesh net (unbiased methods). The effect of crucian carp presence on invertebrate

diversity was then determined using the taxon inventories from the standard or unbiased

methods individually and combined. We hypothesised that: 1) DNA metabarcoding and

morphotaxonomic identification would produce congruent invertebrate communities (e.g.

Carew et al., 2018; Emilson et al., 2017; Nichols et al. 2020), whereas eDNA metabarcoding

would reveal taxa not identified by these tools (e.g. Macher et al., 2018; Mächler et al., 2019);

2) DNA and eDNA metabarcoding would enable species resolution data for problematic taxa

that cannot be morphotaxonomically identified to species-level using standard keys (e.g.

Blackman et al., 2017; Elbrecht et al., 2017a; Lobo et al., 2017; Nichols et al., 2020); and 3)

alpha diversity would be lower in ponds with crucian carp due to direct predation or altered

habitat quality (e.g. Lemmens et al., 2013; Maceda-Veiga, López, & Green, 2017; Schilling et al.,

2009a; Wood et al., 2001), but beta diversity would be enhanced due to community

heterogeneity induced by crucian carp presence in the pond network (e.g. Stefanoudis et al.,

2017). The third pattern was expected regardless of the monitoring tool used. We provide

recommendations for the application of molecular tools to freshwater invertebrate assessment

as well as impact assessment and conservation management of the non-native crucian carp

alongside pond biodiversity.

2. Methods

2.1 Study sites

We surveyed nine ponds containing crucian carp across Norfolk (n = 6) and East Yorkshire (n =

3), and nine fishless ponds in Norfolk (Fig. S1). Crucian carp were either present in ponds as

well-established populations, or individuals that were restocked between 2010 and 2017 as

part of the species' BAP (Copp & Sayer, 2010) because ponds contained populations between

1970-1980 according to local anglers (Sayer et al., 2011; Sayer et al., 2020). The fish status of

ponds was confirmed by prior fyke-net surveys (Sayer et al., 2011; Sayer et al., 2020). For ponds

containing crucian carp, the last fyke-net survey undertaken was 2016 (n = 7), 2015 (n = 1), and

2013 (n = 1). Roach (*Rutilus rutilus*) and rudd (*Scardinius erythrophthalmus*) were the only other

fishes that occurred in three and one of the Norfolk ponds respectively, but at much lower

density (see Table S1). All study ponds were selected to be similar in terms of morphometry and

habitat structure, being <1 ha in area, <5 m in depth, dominated by submerged and floating-

leaved macrophytes, with a largely open-canopy and thus minimal shading (≤20%) of the water

surface. Ponds were mainly surrounded by arable fields, excluding one pond located in

woodland. As invertebrate diversity often peaks in autumn (Hill, Sayer, & Wood, 2016) but large

and more easily identifiable odonate larvae are typically present in summer prior to emergence

(Brooks & Cham, 2014), invertebrate specimens were collected for morphotaxonomic

identification and DNA metabarcoding alongside water samples for eDNA metabarcoding in late

August 2016. Data on physical (area, depth, estimated percentages of perimeter with emergent

vegetation, emergent macrophyte cover, submerged macrophyte cover, and shading) and

chemical (conductivity measured with a HACH HQ30d meter) properties of the ponds were

collected between May and August from 2010 to 2017.

2.2 Sweep-netting and morphotaxonomic identification

Sweep-netting was performed in accordance with the UK National Pond Survey methodology

(Biggs, Fox, & Nicolet, 1998), using a 1 mm mesh long-handled net (0.3 m square bag), to

generate a conventional taxonomic inventory of lentic macroinvertebrates. Sampling time at

each pond comprised a 3 min sweep-net sample and 1 min hand search (see Supporting

Information). Samples were deposited in a 1.2 L sterile Whirl-Pak® stand-up bag (Cole-Palmer,

London) and stored at -20 °C until laboratory processing and sorting. Samples were thawed and

passed through sieves of 8 mm, 2 mm, and 0.25 mm in succession to remove large items of

vegetation and detrital matter. For each sample, invertebrate specimens were sorted into three

body size categories on laminated millimetre graph paper: small (<2.5 x 5 mm), medium (2.5 x 5

mm up to 5 x 10 mm), and large (>10 mm and up to 10 x 20 mm) (Elbrecht, Peinert, & Leese,

2017b). Where possible, size-sorted specimens were identified under a light microscope to

species-level using Freshwater Biological Association publications (Bass, 1998; Brooks & Cham,

2014; Edington & Hildrew, 1995; Elliott, 2009; Elliott & Humpesch, 2010; Elliott & Dobson,

2015; Friday, 1988; Gledhill, Sutcliffe, & Williams, 1993; Macan, 1960; Savage, 1989; Wallace,

Wallace, & Philipson, 1990). Specimens that could not be identified to species-level were

identified to genus or family-level, except Hydrachnidea, Tabanini, Tanytarsini, Oligochaeata,

and Collembola which were recorded as such. Terrestrial taxa and empty cases/shells were

discarded. The laminated paper was sterilised with 50% v/v chlorine-based commercial bleach

solution (Elliot Hygiene Ltd, UK) and 80% v/v ethanol solution between measuring specimens

from different ponds to minimise cross-contamination risk. Specimens were preserved in sterile

15 mL falcon tubes (SARSTED, Germany) containing 100% ethanol according to size category

and pond sampled, and stored at -20 °C until DNA extraction.

2.3 DNA samples

We followed the DNA extraction procedure from Blackman *et al.* (2017). Size categories from

each pond were dried overnight on FisherBrand cellulose filter paper (Fisher Scientific UK Ltd,

UK) in sterile glass funnels and conical flasks to remove excess ethanol. Size categories were

lysed (3 × 30 sec) using a TissueLyser II (Qiagen®, Germany) with DigiSol (50 mM Tris, 20 M

EDTA, 120 mM NaCl and 1% SDS). The adapter sets held 1.5 g of dried tissue and corresponding

volume of DigiSol. If the dry tissue weight of any size category exceeded 1.5 g, we processed

the size category in batches until all tissue had been lysed. The lysates from all batches were

pooled to recreate size categories. The size categories were incubated overnight at 55 °C with

SDS and Proteinase K (Bioline®, UK), following which 200 µL of lysate from each size category

was used for extraction with the DNeasy Blood & Tissue Kit (Qiagen®, Germany) according to

the manufacturer's protocol. Specimens collected from two ponds were categorised as medium

or large only, whereas specimens collected from other study ponds were categorised as small,

medium, and large (Table S2). Consequently, 16 bulk tissue samples were each represented by

three DNA extracts and two bulk tissue samples each represented by two DNA extracts that

were stored at -20 °C until metabarcoding.

2.4 eDNA samples

We repurposed eDNA samples from Norfolk ponds that were collected by Harper et al. (2019)

to validate a quantitative PCR (qPCR) assay for crucian carp. Additional samples were taken

from East Yorkshire ponds for the present study. Details of water sample collection, filtration,

and DNA extraction are provided in the Supporting Information. Briefly, five 2 L surface water

samples were collected at equidistant points around the pond perimeter where access

permitted. For each pond, a full process blank (1 L molecular grade water) was taken into the

field, transported, filtered, and extracted alongside samples to identify contamination. Filters

were stored at -20 °C until DNA extraction with the PowerWater® DNA Isolation Kit (MO BIO

Laboratories, CA, USA) following the manufacturer's protocol. eDNA extracts were stored at -20

°C until metabarcoding.

2.5 Metabarcoding workflow

Our metabarcoding workflow is fully described in the Supporting Information. In short, a

comprehensive list of UK invertebrate species living in or associated with freshwater habitats

(expected fauna) was used to create custom, curated reference databases for UK aquatic

invertebrates, excluding Diptera (Appendix 1). Public GenBank records for Diptera were missing

record features (e.g. 'gene' or 'CDS') and/or names were not in the format required for custom

reference database construction using the selected bioinformatic tools. Assignments to Diptera

were therefore made using the entire NCBI nucleotide database (see below). For the remaining

taxonomic groups, 1483 species were represented in our custom reference database (78.97% of

the expected fauna), although representation varied strongly across groups (Fig. S2). Species

without database representation are listed in Table S3 (see also Figs. S2, S3). Published

metazoan primers mICOIintF (Leray et al., 2013) and jgHCO2198 (Geller, Meyer, Parker, &

Hawk, 2013), which amplify a 313 bp fragment of the mitochondrial cytochrome c oxidase

subunit I (*COI*) gene, were validated *in silico* using ecoPCR software (Ficetola et al., 2010)

against the reference databases. Parameters set allowed a 250-350 bp fragment and maximum

of three mismatches between each primer and each reference sequence (Appendix 2). Primers

were validated *in vitro* against 38 invertebrate species, representing 38 families and 10 major

groups, and compared to published macroinvertebrate metabarcoding primers (Elbrecht &

Leese, 2017; Vamos, Elbrecht, & Leese, 2017) (Figs. S4, S5, S6).

PCR conditions were optimised (Figs. S7, S8) before two independent libraries were

constructed for DNA and eDNA metabarcoding using a two-step PCR protocol (Appendix 1).

During the first PCR (performed in triplicate), the target region was amplified using

oligonucleotides comprising the locus primers, sequencing primers, and pre-adapters. DNA

from the exotic, terrestrial two-spotted assassin bug (*Platymeris biguttatus*) was used for PCR

positive controls (tissue DNA *n* = 9, eDNA *n* = 11) as this species is not found in the UK, whilst

sterile molecular grade water (Fisher Scientific UK Ltd, UK) substituted template DNA for PCR

negative controls (tissue DNA *n* = 9, eDNA *n* = 11). PCR products were individually purified using

a magnetic bead clean-up (VWR International Ltd, UK), following a double size selection

protocol from Bronner et al. (2009). The second PCR bound Multiplex Identification (MID) tags

to the purified products. PCR products were pooled according to PCR run and the pooled PCR

product purified using a magnetic bead clean-up, following a double size selection protocol

from Bronner et al. (2009). Each pooled purified PCR product was quantified on a Qubit™ 3.0

fluorometer using a Qubit™ dsDNA HS Assay Kit (Invitrogen, UK) and normalised according to

concentration and sample number to produce a pooled volume of 20 µL.

The pooled libraries were quantified by qPCR using the NEBNext® Library Quant Kit for

Illumina® (New England Biolabs® Inc., MA, USA). An Agilent 2200 TapeStation and High

Sensitivity D1000 ScreenTape (Agilent Technologies, CA, USA) were used to verify removal of

secondary products from libraries by bead purification and that a fragment of the expected size

(531 bp) remained. A total of 52 bulk tissue subsamples sequenced in triplicate (*n* = 156), 12

extraction blanks, and 18 PCR controls were included alongside samples from other projects (*N*

= 188) in the DNA library. A total of 90 eDNA samples, 18 full process blanks, and 22 PCR

controls were included alongside samples from other projects (*N* = 140) in the eDNA library.

The DNA and eDNA libraries were sequenced on an Illumina® MiSeq using 2 x 300 bp V3

chemistry with 10% and 20% PhiX Sequencing Control respectively (Illumina, Inc, CA, USA).

Raw sequences were converted to taxonomic assignments using metaBEAT v0.97.11, a

custom pipeline for reproducible analysis of metabarcoding data (https://github.com/HullUni-

bioinformatics/metaBEAT). After quality filtering, trimming, merging, chimera detection, and

clustering, non-redundant query sequences were compared against our reference databases

using BLAST (Zhang, Schwartz, Wagner, & Miller, 2000). Putative taxonomic identity was

assigned using a lowest common ancestor (LCA) approach based on the top 10% BLAST matches

for any query matching with at least 90% identity to a reference sequence across more than

80% of its length. Unassigned sequences were subjected to a separate BLAST against the

complete NCBI nucleotide (nt) database at 90% identity to determine the source via LCA as

described above. Bioinformatic settings (Appendix 1) were chosen based on comprehensive

exploration of the parameter space and comparison of metaBEAT taxonomic assignments to

morphotaxonomic inventories. Reference databases in GenBank or fasta format and the

bioinformatic analysis have been deposited in a dedicated GitHub repository, which has been

permanently archived for reproducibility (https://doi.org/10.5281/zenodo.3993125).

2.6 Data analysis

All analyses were performed in R v.3.6.3 (R Core Team, 2017), with data and R scripts deposited

in the GitHub repository. Manipulation of the DNA and eDNA metabarcoding datasets for

analysis is described in the Supporting Information. For each method, two presence-absence

datasets (species-level and family-level) were generated using the *decostand* function in vegan

v2.5-6 (Oksanen et al., 2018) for downstream analyses. This is because potential amplification

bias during PCR can prevent reliable abundance or biomass estimation from sequence reads

produced by DNA or eDNA metabarcoding (Elbrecht et al., 2017b). The entire species and

family inventories produced by each method (hereafter standard methods) were compared.

We also compared inventories from each method after correcting for known biases: taxa

without reference sequences (i.e. those which cannot be detected with metabarcoding) were

removed from the morphotaxonomic dataset, and meiofauna (i.e. taxa that would escape a 1

mm mesh net) were removed from the metabarcoding datasets (hereafter unbiased methods).

The effect of crucian carp on invertebrate diversity was then assessed at species and family-

level according to standard methods (main text) and unbiased methods (Supporting

Information) individually and combined. Finally, the impact of crucian carp alongside

environmental variables on invertebrate diversity was investigated using the combined

standard method (main text) and unbiased method (Supporting Information) datasets. For

species-level analyses, only species records were used, whereas family-level analyses included

species, genus and family records grouped at the taxonomic rank of family.

We define alpha diversity as taxon richness (species or families) within individual ponds,

and beta diversity as the difference between communities present at each pond whilst

accounting for taxon identity (Baselga & Orme, 2012). Jaccard dissimilarity was used as a

measure of beta diversity for all analyses. For each standard or unbiased dataset, the following

analyses were performed. Alpha diversity (response variable) was obtained using the

specnumber function in vegan v2.5-6 and modelled against sampling method (explanatory

variable), then modelled against crucian carp presence-absence (explanatory variable). Using

the combined method datasets, alpha diversity of the major invertebrate groups, i.e. orders,

classes, or phyla (Dobson, Pawley, Fletcher, & Powell, 2012), in each pond (response variable),

was modelled against crucian carp presence-absence (explanatory variable). All alpha diversity

comparisons were made using negative binomial Generalized Linear Models (GLMs), and

Pairwise Tukey's HSD tests were used to assess significance of differences. We used betapart

v1.5.1 (Baselga & Orme, 2012) to estimate total beta diversity, partitioned into nestedness-

resultant and turnover components, across sampling methods and ponds using the *beta.multi*

function. These three beta diversity components were estimated for inventories produced by

each sampling method, and ponds with or without crucian carp using the *beta.pair* function

(pairwise beta diversity scores). For each beta diversity component, we compared community

heterogeneity in each group (sampling method or crucian carp presence-absence) by

calculating homogeneity of multivariate dispersions (MVDISP) using the *betadisper* function

from vegan v2.5-6. Variation in MVDISP was statistically tested using an analysis of variance

(ANOVA), and pairwise Tukey's HSD tests used to determine if there were significant

differences between the groups. The effect of sampling method and crucian carp on each beta

diversity component was visualised using Non-metric Multidimensional Scaling (NMDS)

ordination plots with the *metaMDS* function, and tested statistically using permutational

multivariate analysis of variance (PERMANOVA) with the *adonis* function in vegan v2.5-6. Pre-

defined cut-off values were used for effect size, where PERMANOVA results were interpreted as

moderate and strong effects if $R^2 > 0.09$ and $R^2 > 0.25$ respectively. These values are broadly

equivalent to correlation coefficients of $r = 0.3$ and 0.5 which represent moderate and strong

effects accordingly (Macher et al., 2018; Nakagawa & Cuthill, 2007).

We tested whether the species and family-level invertebrate communities produced by

the combined standard or unbiased methods were influenced by pond properties in

conjunction with crucian carp presence-absence. Redundancy Analysis (RDA) was selected for

constrained ordination as it analyses variation in biological communities in response to

explanatory variables (Legendre & Legendre, 2012). Principal Coordinate Analysis (PCoA) was

performed using the *pcoa* function in ape v5.3 (Paradis & Schliep, 2018) on the turnover,

nestedness-resultant, and total beta diversity matrices. The Lingoes correction was employed

to account for negative eigenvalues (Legendre, 2014). The resultant PCoA eigenvectors

(principle coordinates) for each distance matrix were used as the response variable in variance

partitioning. Our variables were grouped as biotic (crucian carp presence-absence) or abiotic

(pond conductivity, area, depth, percentages of perimeter with emergent vegetation, emergent

macrophyte cover, submerged macrophyte cover, and shading). Abiotic variables were log10

transformed to eliminate their physical units (Legendre & Birks, 2012). Significant abiotic

variables influencing each component of beta diversity were identified using the *ordiR2step*

function in vegan v2.5-6 to perform separate RDA analyses under a forward selection

procedure. Where applicable, the relative contributions of the biotic and abiotic variables on

turnover, nestedness-resultant, and total beta diversity for the species and family-level

invertebrate communities were assessed by variance partitioning (Borcard, Legendre, &

Drapeau, 1992) using the *varpart* function in vegan v2.5-6. For each beta diversity component,

RDA was performed using biotic and significant abiotic variables, and variance partitioning used

to divide the total percentage of variation explained into unique and shared contributions for

biotic and abiotic predictor groups. The *anova* function in vegan v2.5-6 was used to examine

the statistical significance of the full model and the unique contributions of each predictor

group. We report the adjusted R^2-fractions as they are widely recommended and unbiased

(Peres-Neto, Legendre, Dray, & Borcard, 2006).

3. Results

3.1 Comparison of methods for freshwater invertebrate assessment

Summaries of taxon detection by standard and unbiased methods are reported in the Supporting Information (Appendix 2; Tables S4, S5, S6, S7; Figs. S9, S10). Briefly, using microscopy, we identified 2,281 specimens belonging to 38 families, of which 1,404 specimens belonged to 91 species. With DNA metabarcoding, 2,906,869 sequence reads were assigned to 55 families, of which 2,448,078 sequence reads were assigned to 131 species. With eDNA metabarcoding, 813,376 sequence reads were assigned to 90 families, of which 346,163 sequence reads were assigned to 145 species. Only 19 species (Figs. 1ai, bi) and 22 families (Figs. 1aii, bii) were shared by all three methods (standard or unbiased). Standard methods detected more distinct taxa than unbiased methods, especially morphotaxonomic identification and eDNA metabarcoding. eDNA metabarcoding detected the most unique species and families, whereas DNA metabarcoding and morphotaxonomic identification were more comparable (Fig. 1). The number of taxa that morphotaxonomic identification shared with DNA and eDNA metabarcoding was unchanged using standard or unbiased methods. Detection of one family was exclusive to morphotaxonomic identification and eDNA metabarcoding, whereas detection of 36 species and 13 families was exclusive to morphotaxonomic identification and DNA metabarcoding (Fig. 1). Morphotaxonomic identification detected 19 species without reference sequences, whereas DNA and eDNA metabarcoding detected 4 and 41 meiofaunal species respectively (Fig. 1a). The metabarcoding approaches either shared 44 species and 16 families (standard methods, Fig. 1a) or 33 species and 8 families (unbiased methods, Fig. 1b).

Standard sampling method influenced alpha diversity of ponds at species (GLM χ^2_2 = 26.058, P < 0.001) and family-level (GLM χ^2_2 = 48.238, P < 0.001). Significant differences (P < 0.05) between the alpha diversity means for morphotaxonomic identification and DNA metabarcoding, morphotaxonomic identification and eDNA metabarcoding, and DNA and eDNA metabarcoding (family-level only) were observed at species and family-level (Table 1).

Conversely, unbiased sampling method influenced alpha diversity of ponds at species-level
(GLM χ^2_2 = 14.614, P < 0.001), but not family-level (GLM χ^2_2 = 5.214, P = 0.074). Significant
differences between the species-level alpha diversity means of morphotaxonomic identification
and DNA metabarcoding, and DNA and eDNA metabarcoding were observed (Table 1). With
standard methods, species and family-level alpha diversity was highest using eDNA
metabarcoding, followed by DNA metabarcoding then morphotaxonomic identification (Figs.
2ai, aii). With unbiased methods, DNA metabarcoding detected more species than
morphotaxonomic identification and eDNA metabarcoding, but alpha diversity was comparable
at family-level (Figs. 2bi, bii). For beta diversity, MVDISP did not significantly differ between
standard or unbiased methods for turnover, nestedness-resultant or total beta diversity at
either taxonomic rank (Table S8). Both standard and unbiased sampling method had moderate
and strong positive effects on turnover and total beta diversity at species (Figs. 3ai, aiii, ci, ciii)
and family-level (Figs. 3aii, aiv, cii, civ) respectively, but not nestedness-resultant (Table S9).

3.2 Impact of crucian carp stocking on pond invertebrates

Here, we report the impact of crucian carp stocking assessed using standard methods as these
provide the most comprehensive evaluation of invertebrate diversity. Crucian carp impact
assessment using unbiased methods is reported in the Supporting Information (Appendix 2;
Tables S10, S11, S12, S13; Figs. S11, S12, S13, S14). Independently and combined, standard
methods showed alpha diversity of invertebrates to be marginally lower in ponds containing
crucian carp at species (Figs. 4ai-iv) and family-level (Figs. 4bi-iv), but this difference was not
statistically significant (P > 0.05, Table 2). Detailed examination of alpha diversity within the
major invertebrate groups identified by all three methods combined revealed that Coleoptera
and Mollusca diversity was reduced (Coleoptera borderline significant and Mollusca highly
significant) in ponds with crucian carp at species-level (GLM χ^2_{21} = 28.033, P = 0.139; Coleoptera
-0.481 ± 0.254, Z = -1.897, P = 0.058; Mollusca -0.925 ± 0.278, Z = -3.324, P < 0.001), but not
family-level (GLM χ^2_{22} = 18.366, P = 0.684; Coleoptera -0.544 ± -0.296, Z = -1.834, P = 0.067;
Mollusca -0.442 ± -0.302, Z = -1.462, P = 0.144). However, differences in alpha diversity

between ponds with or without crucian carp were not significant for other invertebrate groups

at either taxonomic rank (Fig. S15). Alpha diversity relationships held true for the unbiased

methods (Table S10; Fig. S11, S12).

Total beta diversity of ponds was consistently high at species and family-level for

independent and combined methods (standard or unbiased). Variation in invertebrate

community composition was driven by turnover rather than nestedness-resultant (Tables 3,

S11). MVDISP differed between ponds for species-level total beta diversity with eDNA

metabarcoding, and species and family-level total beta diversity with all methods combined.

Fishless ponds had significantly lower dispersion than ponds with crucian carp (Table S14).

MVDISP patterns were not consistent across standard and unbiased methods (Table S12). At

species-level, morphotaxonomic identification, eDNA metabarcoding, and all methods

combined revealed a weak or moderate positive influence of crucian carp presence on turnover

(Table 4; Figs. 5ai, iii, iv) and total beta diversity (Table 4; Figs. 5ci, iii, iv) between ponds, but

not nestedness-resultant (Table 4; Figs. 5bi, iii, iv). At family-level, eDNA metabarcoding and all

methods combined revealed a weak positive influence of crucian carp presence on total beta

diversity (Table 4; Fig. S16ciii, iv) between ponds, but not turnover (Table 4; Fig. S16aiii, iv) or

nestedness-resultant (Table 4; Fig. S16biii, iv). Crucian carp presence did not influence any beta

diversity component produced by DNA metabarcoding (Table 4; Figs. 5aii, bii, cii). Similar beta

diversity patterns were obtained using unbiased eDNA metabarcoding and combined methods

(Table S13; Figs. S13, S14).

Additional analyses undertaken on the combined method datasets supported an effect

of crucian carp presence and excluded the influence of abiotic variables on invertebrate

diversity. At species-level, only pond area was identified by forward selection as a significant

abiotic variable for turnover and total beta diversity. Variance partitioning analysis was

undertaken for turnover and total beta diversity using crucian carp presence-absence and pond

area. Using the adjusted R^2 values, biotic and abiotic variables explained 7.04% and 4.75% of

the total variation in species turnover and total beta diversity respectively (Fig. 6). Crucian carp

presence-absence significantly contributed to turnover (Fig. 6a; adjusted R^2 = 5.55%, F_1 = 2.031,

P = 0.001) and total beta diversity (Fig. 6b; adjusted R^2 = 4.05%, F_1 = 1.730, P = 0.001), whereas

pond area explained less variance and did not influence turnover (Fig. 6a; adjusted R^2 = 2.94%,

F_1 = 0.340, P = 0.852) or total beta diversity (Fig. 6b; adjusted R^2 = 1.93%, F_1 = 1.118, P = 0.218).

RDA of nestedness-resultant without abiotic data indicated no influence of crucian carp

presence-absence (F_1 = 0.340, P = 0.850). At family-level, forward selection did not identify any

significant abiotic variables for turnover, nestedness-resultant, or total beta diversity.

Therefore, variance partitioning was not undertaken for any component of beta diversity. RDA

of each beta diversity component minus abiotic data showed that crucian carp presence-

absence influenced total beta diversity (F_1 = 1.474, P = 0.034), but not turnover (F_1 = 1.399, P =

0.106) or nestedness-resultant (F_1 = 1.970, P = 0.141). Variance partitioning results from the

unbiased methods are presented in Appendix 2.

4. Discussion

4.1 Comparison of methods for freshwater invertebrate assessment

DNA (unbiased methods) or eDNA (standard methods) metabarcoding generated the highest

species and family-level alpha diversity. Differences in alpha diversity were more pronounced

between standard methods, but unique species were detected with unbiased methods

nonetheless. Morphotaxonomic identification and DNA metabarcoding produced more similar

communities, and performed best for Coleoptera, Hemiptera, Hirudinea, Megaloptera, and

Odonata. eDNA metabarcoding generated a markedly different community, and detected

aquatic taxa overlooked in pond net samples (Crustacea, Lepidoptera [i.e. Crambidae species

with aquatic larvae such as *Cataclysta lemnata*], Trichoptera) alongside singleton terrestrial

taxa (Arachnida, Hymenoptera, Psocoptera, Thysanoptera). Both metabarcoding approaches

performed well for Annelida, Collembola, Diptera and Ephemeroptera, particularly at species-

level.

Critically, we used a 1 mm mesh net which is designed to sample macroinvertebrates,

whereas a 500 μm mesh net would also sample the meiofauna detected by metabarcoding.

Meiofauna may inflate metabarcoding diversity estimates compared to morphotaxonomic

identification, but may also prevent PCR amplification of macroinvertebrates (Beentjes et al.,

2019; Deiner et al., 2016; Macher et al., 2018). To mitigate bias, we performed all analyses

excluding meiofauna not captured by netting and excluding morphotaxonomically identified

taxa without reference sequences that metabarcoding would not recover. Alpha diversity

results from the standard and unbiased method comparisons were variable, but beta diversity

results were more consistent. Both method comparisons echo other studies where

metabarcoding captured greater diversity than morphotaxonomic approaches and resolved

problematic groups (e.g. Diptera) for species-level morphotaxonomic identification (Andújar et

al., 2017; Carew et al., 2018; Clarke, Beard, Swadling, & Deagle, 2017; Elbrecht et al., 2017a;

Emilson et al., 2017; Klymus et al., 2017; Lobo et al., 2017). Standard and unbiased methods

broadly revealed the same ecological relationships, although standard methods may yield more

insights. Regardless, all three methods will be most powerful when used in combination for

ecological investigation.

Each method has inherent biases that influence performance for freshwater

invertebrate assessment. The UK National Pond Survey methodology recommends that pond

net samples are placed into buckets/zip-lock bags, followed by sorting and identification in the

laboratory as soon as possible. Samples are typically preserved in ethanol or refrigerated to

prolong processing time (Biggs et al., 1998), but this standard methodology does not lend itself

to molecular applications. After collection, samples were transported on ice then frozen at -20

°C to prevent predation within samples, minimise organismal decay and subsequent DNA

degradation, and allow samples to be processed as required in the laboratory. Using this

strategy, small or inconspicuous dead specimens would not have been recovered during sorting

(Biggs et al., 1998) and therefore excluded from morphotaxonomic and DNA metabarcoding

inventories. These losses are surplus to the estimated 29% of specimens already overlooked

during sorting due to smaller body size (Haase et al., 2010). Sorting may also damage or

completely destroy some recovered specimens, influencing morphotaxonomic identification

(Lobo et al., 2017; Zizka, Leese, Peinert, & Geiger, 2018).

Human error is another source of discrepancy between morphotaxonomic identification
and metabarcoding (Carew, Pettigrove, Metzeling, & Hoffmann, 2013; Elbrecht et al., 2017a;
Haase et al., 2010). Some taxa were identified to genus or family-level and omitted from our
species-level inventories, whereas other taxa may have been falsely identified and inventoried
despite caution exercised to only assign taxa to species-level where there was high confidence
in identification. Losses incurred by the preservation and sorting treatments for
morphotaxonomic identification cannot be mitigated, and human error can only be reduced
with taxonomic expertise. Conversely, taxon recovery by DNA metabarcoding can be improved
with protocols that preserve and extract DNA from intact, unsorted samples (Elbrecht et al.,
2017a; Nichols et al., 2020). For example, ethanol used to preserve specimens (Zizka et al.,
2018), or replacement of ethanol with DNA extraction buffer followed by incubation then
specimen removal (Carew, Coleman, & Hoffmann, 2018), offer alternative starting materials to
bulk tissue for DNA metabarcoding. However, these alternative DNA sources tend to produce
false negatives for schlerotised groups, such as Coleoptera, Trichoptera, and Mollusca (Carew et
al., 2018; Zizka et al., 2018). More recently, Nichols et al. (2020) demonstrated that unsorted
samples, including the sample matrix, recovered the same taxa as sorted samples, thus sorting
may be unnecessary.

Despite successful application of DNA metabarcoding to bulk tissue samples for
biomonitoring (Andújar et al., 2017; Carew et al., 2018; Elbrecht et al., 2017a; Emilson et al.,
2017; Macher et al., 2018), recurring issues have been encountered. Arguably, the most
pressing issue is size bias, where DNA from large and/or high biomass taxa can outcompete
DNA of smaller and/or low biomass taxa during PCR amplification and sequencing (Elbrecht et
al., 2017b). Although we used the size-sorting approach of Elbrecht *et al*. (2017b) and
sequenced body size categories independently with data pooling downstream, DNA
metabarcoding failed to detect several taxa which have reference sequences and were reliably
identified by microscopy, including multiple coleopterans (*Agabus sturmii*, *Hydroporus*
*erythrocephalu*s, *Rhantus frontalis*, *Haliplus confinis*, *Haliplus ruficollis*), a small mollusc
(*Gyraulus crista*), a medium-sized hirudinean (*Erpobdella lineata*), and two large anisopterans
(*Aeshna mixta*, *Anax imperator*). Primer bias may be partly responsible for these nondetections

(see below), but non-delivery of desired results by size-sorting is problematic as it is a time-
consuming, labour-intensive process that potentially increases cross-contamination risk
between samples (Elbrecht et al., 2017b), and still requires samples to be sorted from
vegetation and substrate (Elbrecht et al., 2017a). Size-sorting may therefore be a drain on
resources and time allocated to DNA metabarcoding projects, but other sources of false
negatives must be excluded.

In contrast, eDNA metabarcoding preferentially amplifies DNA from small, floating
organisms (Beentjes et al., 2019; Deiner et al., 2016; Leese et al., 2020; Macher et al., 2018),
and these organisms are often retained on filter membranes used for eDNA capture (*pers.*
obs.). Consequently, their DNA is abundant in samples and overwhelms DNA from other taxa
during sequencing. Pre-filtering with or only using large pore size filter membranes may reduce
this bias (Macher et al., 2018). eDNA metabarcoding may also detect different taxa due to
species variability in eDNA production and shedding rates. The species infrequently detected by
eDNA metabarcoding were those that possess thicker exoskeletons composed of chitin and
occasionally calcium carbonate, e.g. Coleoptera, Hemiptera. Exoskeletons may restrict the
release of DNA into the water column (Tréguier et al., 2014) as opposed to organisms that are
filter-feeders or produce slime (e.g. Mollusca), ectoparasites (i.e. Acari) that feed on blood or
skin of other species, or use external versus internal fertilisation (Mächler et al., 2019; Wacker
et al., 2019). Furthermore, species may differ in their habitat preferences and utilisation,
potentially resulting in highly localised distributions (Klymus et al., 2017; Mächler et al., 2019).
Therefore, more samples or greater volumes may be required to improve detection of pond
biota (Harper et al., 2019).

Marker choice and primer design can substantially influence amplification success and
taxonomic assignment. Although the *COI* region offers species resolution, has extensive
database representation, and is used as standard in DNA barcoding (Elbrecht, Hebert, &
Steinke, 2018), it lacks conserved primer-binding sites as a protein-coding gene (Clarke et al.,
2017; Leese et al., 2020). This introduces high risk of primer mismatch and subsequent bias
during metabarcoding (Clarke et al., 2017; Elbrecht et al., 2016; Elbrecht et al., 2018; Lobo et
al., 2017). Most *COI* metabarcoding primers (Elbrecht & Leese, 2017; Leray et al., 2013;

Meusnier et al., 2008; Zeale, Butlin, Barker, Lees, & Jones, 2011) are degenerate to enable

binding, but degenerate primers may bind non-target regions (Elbrecht et al., 2018) and amplify

non-metazoans, e.g. bacteria, algae, fungi (Brandon-Mong et al., 2015; Leese et al., 2020;

Macher et al., 2018). Degenerate primers can experience primer slippage and produce variable

length sequences across species, which has implications for bioinformatic processing and

taxonomic assignment (Elbrecht et al., 2018). Using different (e.g. 16S ribosomal RNA) or

multiple markers may counter these issues, but different markers often lack reference database

representation, hindering taxonomic resolution, and use of multiple markers will increase PCR

and sequencing costs (Clarke et al., 2017; Deiner et al., 2016; Elbrecht et al., 2016; Elbrecht et

al., 2017a).

We obtained an excessive number (~60%) of unassigned reads from eDNA

metabarcoding, despite using a relaxed BLAST identity (90%) against custom and public

reference databases. This suggests that sequences were of poor quality, could not be assigned

to reference sequences, or lacked reference database representation (Macher et al., 2018).

Public reference database records may be few, mislabelled, or have limited geographic

coverage (Curry, Gibson, Shokralla, Hajibabaei, & Baird, 2018; Elbrecht et al., 2017a; Klymus et

al., 2017; Weigand et al., 2019). We morphotaxonomically identified 19 species (Hemiptera,

Mollusca, Odonata) without reference sequences whose DNA may have been sequenced, but

would not have been taxonomically assigned to species-level. Future studies must develop

more specific primers that have binding and flanking regions of high nucleotide diversity

(Elbrecht et al., 2018; Leese et al., 2020) and/or target particular invertebrate orders or families

(Klymus et al., 2017) as well as procure reference sequences for different markers (Curry et al.,

2018; Elbrecht et al., 2016; Weigand et al., 2019). These are essential steps to improve the

reliability and accuracy of molecular monitoring in freshwater ecosystems.

4.2 Impact of crucian carp stocking on pond invertebrates

Our findings indicate that the crucian carp can be an important driver of invertebrate

community heterogeneity in ponds. Crucian carp had a negligible influence on alpha diversity

and a positive influence on beta diversity of pond invertebrates. Alpha diversity in ponds with

crucian carp was marginally reduced compared to alpha diversity in fishless ponds, but this

difference was not significant across methods used at either taxonomic rank. Within the major

invertebrate groups identified by all methods combined, species-level alpha diversity of

Coleoptera and Mollusca was reduced in ponds containing crucian carp as opposed to fishless

ponds. This may be because coleopterans are large, active swimmers that are visible and easy

to hunt, and because molluscs are slow-moving prey. Across ponds, total beta diversity of

invertebrate communities was driven by turnover (taxon replacement) rather than nestedness-

resultant (taxon subsets) (Baselga & Orme, 2012). Our results revealed crucian carp to

positively influence turnover and total beta diversity between ponds. Therefore, taxa in fishless

ponds were replaced by different taxa in ponds with crucian carp, resulting in dissimilar

community composition. Removal of top predators and dominant species, such as

coleopterans, by crucian carp may allow other prey species to colonise but further research is

needed to test this hypothesis.

Our results both echo and contradict Stefanoudis et al. (2017), where the presence of

fish (including crucian carp) in ponds altered macrophyte and cladoceran community

composition, but not water beetle composition. Hassall et al. (2011) also found that fish

presence in ponds had a positive effect on species richness of most invertebrate taxa, but a

negative effect on Coleoptera. Gee et al. (1997) observed no influence of fish stocking on

macrophyte and macroinvertebrate species richness, although Odonata richness was lower and

Trichoptera richness higher in stocked ponds. Conversely, other research found that

managed/stocked ponds, some of which contained crucian carp, had lower invertebrate

diversity than unmanaged sites (i.e. no human intervention to slow succession), which were

characterised by Trichoptera, Coleoptera and Zygoptera larvae (Wood et al., 2001). Similarly,

large, active and free-swimming taxa (Notonectidae, Corixidae, Gyrinidae, Dytiscidae,

Aeshnidae, Libellulidae and Chaoboridae) were strongly associated with fish absence as well as

more diverse and abundant in fishless lakes (Bendell & McNicol, 1995; Schilling et al., 2009a,

2009b).

Critically, only two of the aforementioned studies accounted for the identity of fish

species assemblages present (Stefanoudis et al., 2017; Wood et al., 2001) and just one was

based on standardised fish surveys (Stefanoudis et al., 2017), whereas other studies only

considered the presence of a particular species (Schilling et al., 2009a) or fish presence-absence

generally (Bendell & McNicol, 1995; Gee et al., 1997; Hassall et al., 2011; Schilling et al., 2009b).

The contrasting results produced by these studies and our own indicate that the impact of fish

on pond biodiversity will likely depend on the species present, population density, and

management strategy. Wetland fishes vary in dietary preference and consume different

proportions of invertebrate taxa, thus different fishes will suppress numbers of and confer

benefits to different invertebrate taxa (Batzer, Pusateri, & Vetter, 2000). Invasive fish species

may be more detrimental than non-invasive species, for example, the mosquitofish (*Gambusia*

affinis) reduced zooplankton abundance and macroinvertebrate density by 90% and 50%

respectively after introduction in a wetland ecosystem experiment (Preston et al., 2017).

Additionally, invertebrate species richness and abundance was found to decrease as duration of

stocking increased (Schilling et al., 2009a). Therefore, local- and regional-scale diversity may

benefit most from pond mosaics that maintain fish-free and fish-containing (at low-moderate

densities) ponds to prevent increases in fish biomass and predation pressure over time

(Lemmens et al., 2013).

The effects of environmental variables on alpha and beta diversity of pond invertebrates

must also be considered (Hassall et al., 2011; Hill, Heino, Thornhill, Ryves, & Wood, 2017). Our

study ponds were selected to have similar physical and chemical properties, which may explain

the minimal contribution of abiotic factors to variance in invertebrate community structure.

Although pond area was retained by model selection, it did not significantly influence

community structure and explained less variance than crucian carp presence-absence. Likewise,

other studies have shown a weak or no effect of pond area on invertebrate species richness and

community composition (Gee et al., 1997; Gledhill et al., 2008; Oertli et al., 2002; Wood et al.,

2001). Critically, our environmental data were collected over a 7-year period, whereas

contemporary data may have better explained variance in community structure. We omitted

variables that experience high temporal variation from our analyses, but these may be linked to

the Coleoptera and Mollusca alpha diversity reductions and different invertebrate communities

observed in crucian carp ponds, such as water temperature, pH, nutrient concentration, and

surface dissolved oxygen (Bendell & McNicol, 1995; Hassall et al., 2011; Menetrey, Sager, Oertli,

& Lachavanne, 2005). Investigations examining contemporary physicochemical variables

alongside crucian carp presence-absence will help disentangle the effects of stocking and

habitat. The degree of connectivity between ponds may also substantially influence

invertebrate community structure, both independently and in combination with crucian carp

presence (Hill et al., 2017).

The crucian carp might be assumed to have negative impacts on pond biodiversity by

having "carp" in its name and likely being non-native (Jeffries et al., 2017). Foraging activities of

introduced common carp are known to reduce invertebrate density and macrophyte cover,

with knock-on effects for amphibian and waterfowl species richness and abundance (Chan,

2011; Haas et al., 2007; Maceda-Veiga et al., 2017). However, all crucian carp ponds studied

here were dominated by floating and/or submerged macrophyte beds as in Stefanoudis et al.

(2017), and we observed no evidence for negative impacts on vegetation. Additionally, no

impact of crucian carp on amphibian presence, oviposition, larval behaviour, or recruitment

success has been found (Chan, 2011; Harper et al., 2019; Jarvis, 2012). Despite its probable non-

native status, cumulative evidence suggests that the crucian carp does not have invasive

potential. Instead, the crucian carp is naturalised and faces the same threats as native UK pond

biodiversity, such as the great crested newt (*Triturus cristatus*), including terrestrialisation,

dessication and infilling of ponds as well as predation by other fishes, e.g. pike (*Esox lucius*) and

three-spined stickleback (*Gasterosteus aculeatus*) (Sayer et al., 2020). Additionally, the crucian

carp is threatened by hybridisation with goldfish, common carp, and gibel carp across Europe

(Hänfling et al., 2005). Gibel carp are though to be absent from UK waters and given the severe

threat that this species poses to crucian carp persistence in ponds, England can be regarded as

an important centre for crucian carp conservation (Jeffries et al., 2017; Sayer et al., 2020).

Moreover, pond restoration as part of crucian carp conservation efforts in Norfolk has

substantially benefitted macrophyte, invertebrate, amphibian, and farmland bird diversity

(Sayer et al., 2013; Lewis-Phillips et al., 2019).

Our study demonstrates that crucian carp presence enhances invertebrate diversity

across pond networks, and that current conservation management of established and stocked
populations in England is appropriate. Crucian carp may contribute to taxon turnover and
subsequent community heterogeneity in ponds by providing colonisation opportunities for
smaller invertebrates through predator (e.g. coleopterans) removal. We conclude that
management strategies to support high landscape-scale pond biodiversity should consider a
patchwork of fish-containing and fishless ponds, although stocking should only occur in
artificially created ponds or in ponds where fish were previously known to occur (Sayer et al.,
2020). The crucian carp is likely to be an important species in this respect, but more research is
required to verify that crucian carp presence increases community heterogeneity without
contributing to the decline of endangered invertebrates, albeit this is unlikely. Furthermore, the
impact of crucian carp on all pond biodiversity, especially amphibians, must be studied more
broadly with respect to crucian carp population density, seasonality, and the aforementioned
environmental variables to determine whether maintaining populations and stocking is
invariably beneficial.

Acknowledgements

This work was funded by the University of Hull. Additionally, D.S.R was supported by the
Natural Environment Research Council award number NE/R016429/1 as part of the UK-SCAPE
programme delivering National Capability. We thank the many landowners who allowed access
to ponds. We are grateful to the Norfolk Crucian Project for providing data associated with
Norfolk ponds. We thank Michael Lee and Matt Smith (Environment Agency) for selecting East
Yorkshire ponds and providing associated data, and Jane Colley (Askham Bryan College) for
providing invertebrate samples from these ponds. We are grateful to Jianlong Li and Cristina Di
Muri (University of Hull) for assistance with water sampling and filtration as well as advice on
library preparation and sequencing.

Data accessibility

Raw sequence reads have been archived on the NCBI Sequence Read Archive (Study: SRP163672; BioProject: PRJNA494857; BioSamples: SAMN10181701 - SAMN10182084 [bulk tissue DNA] and SAMN10187732 - SAMN10188115 [eDNA]; SRA accessions: SRR7969394 - SRR796977 [bulk tissue DNA] and SRR7985814 - SRR7986197 [eDNA]). Jupyter notebooks, R scripts and corresponding data are deposited in a dedicated GitHub repository (https://github.com/HullUni-bioinformatics/Harper_et_al_2020_crucian_carp_impact_invertebrates_conventional_molecular_tools) which has been permanently archived (https://doi.org/10.5281/zenodo.3993125).

Author contributions

L.R.H, B.H, and L.L.H conceived and designed the study. C.D.S selected the study ponds in Norfolk and provided associated environmental data. L.R.H, M.B, and R.C.B built the custom reference sequence database for invertebrates. L.R.H and R.C.B sampled the Norfolk ponds for invertebrates. L.R.H and M.B collected and filtered water samples from all study ponds. D.S.R assisted with water filtration of eDNA samples and provided advice on primers, PCR protocols, and sequencing. L.R.H performed laboratory work, and analysed the data with assistance from M.J.H. L.R.H wrote the manuscript, which all authors revised.

References

Andújar, C., Arribas, P., Gray, C., Bruce, C., Woodward, G., Yu, D. W., & Vogler, A. P. (2017). Metabarcoding of freshwater invertebrates to detect the effects of a pesticide spill. *Molecular Ecology*, *27*, 146–166. https://doi.org/10.1111/mec.14410

Baselga, A., & Orme, C. D. L. (2012). betapart: an R package for the study of beta diversity : Betapart package. *Methods in Ecology and Evolution*, *3*, 808–812. https://doi.org/10.1111/j.2041-210X.2012.00224.x

Bass, J. (1998). *Last-instar larvae and pupae of the Simuliidae of Britain and Ireland: A key with brief ecological notes* (Vol. 18). Ambleside, Freshwater Biological Association.

Batzer, D. P., Pusateri, C. R., & Vetter, R. (2000). IMPACTS OF FISH PREDATION ON MARSH INVERTEBRATES: DIRECT AND INDIRECT EFFECTS. *Wetlands*, *20*, 307–312. https://doi.org/10.1672/0277-5212(2000)020[0307:IOFPOM]2.0.CO;2

Beentjes, K. K., Speksnijder, A. G. C., Schilthuizen, M., Hoogeveen, M., & van der Hoorn, B.B. (2019) The effects of spatial and temporal replicate sampling on eDNA metabarcoding. *PeerJ*, *7*, e7335. https://doi.org/10.7717/peerj.7335 Bendell, B. E., & McNicol, D. K. (1995). Lake acidity, fish predation, and the distribution and abundance of some littoral insects. *Hydrobiologia*, *302*, 133–145. https://doi.org/10.1007/BF00027038

Biggs, J., Fox, G., & Nicolet, P. (1998). *A guide to the methods of the National Pond Survey*. Oxford, Pond Action.

Biggs, J., von Fumetti, S., & Kelly-Quinn, M. (2016). The importance of small waterbodies for biodiversity and ecosystem services: implications for policy makers. *Hydrobiologia*, *793*, 3–39. https://doi.org/10.1007/s10750-016-3007-0

Blackman, R. C., Constable, D., Hahn, C., Sheard, A. M., Durkota, J., Hänfling, B., & Lawson Handley, L. (2017). Detection of a new non-native freshwater species by DNA metabarcoding of environmental samples--first record of *Gammarus fossarum* in the UK. *Aquatic Invasions*, *12*, 177–189. https://doi.org/10.3391/ai.2017.12.2.06

Borcard, D., Legendre, P., & Drapeau, P. (1992). Partialling out the Spatial Component of Ecological Variation. *Ecology*, *73*, 1045–1055. https://doi.org/10.2307/1940179

Brandon-Mong, G.-J., Gan, H.-M., Sing, K.-W., Lee, P.-S., Lim, P.-E., & Wilson, J.-J. (2015). DNA metabarcoding of insects and allies: an evaluation of primers and pipelines. *Bulletin of Entomological Research*, *105*, 717–727. https://doi.org/10.1017/S0007485315000681

Briers, R. A., & Biggs, J. (2003). Indicator taxa for the conservation of pond invertebrate diversity. *Aquatic Conservation: Marine and Freshwater Ecosystems*, *13*, 323–330. https://doi.org/10.1002/aqc.576

Bronner, I. F., Quail, M. A., Turner, D. J., & Swerdlow, H. (2009). Improved Protocols for Illumina Sequencing. *Current Protocols in Human Genetics*, *18*, 18.2.1–18.2.42. https://doi.org/10.1002/0471142905.hg1802s79

Brooks, S., & Cham, S. (2014). *Field Guide to the Dragonflies & Damselflies of Great Britain and Ireland*. London, Bloomsbury Publishing Plc.

Carew, M. E., Coleman, R. A., & Hoffmann, A. A. (2018). Can non-destructive DNA extraction of bulk invertebrate samples be used for metabarcoding? *PeerJ*, *6*, e4980. https://doi.org/10.7717/peerj.4980

Carew, M. E., Kellar, C. R., Pettigrove, V. J., & Hoffmann, A. A. (2018). Can high-throughput sequencing detect macroinvertebrate diversity for routine monitoring of an urban river? *Ecological Indicators*, *85*, 440–450. https://doi.org/10.1016/j.ecolind.2017.11.002

Carew, M. E., Pettigrove, V. J., Metzeling, L., & Hoffmann, A. A. (2013). Environmental monitoring using next generation sequencing: rapid identification of macroinvertebrate bioindicator species. *Frontiers in Zoology*, *10*, 45. https://doi.org/10.1186/1742-9994-10-45

Céréghino, R., Biggs, J., Oertli, B., & Declerck, S. (2008). The ecology of European ponds: defining the characteristics of a neglected freshwater habitat. *Hydrobiologia*, *597*, 1–6. https://doi.org/10.1007/s10750-007-9225-8

Chan, K. (2011) *Can Great Crested Newts (Triturus Cristatus) Coexist with Fish?* Masters Thesis, University College London (UCL).

Clarke, L. J., Beard, J. M., Swadling, K. M., & Deagle, B. E. (2017). Effect of marker choice and thermal cycling protocol on zooplankton DNA metabarcoding studies. *Ecology and Evolution*, *7*, 873–883. https://doi.org/10.1002/ece3.2667

Copp, G. H., & Sayer, C. D. (2010). *Norfolk Biodiversity Action Plan--Local Species Action Plan for Crucian Carp (Carassius carassius)*. Lowestoft, Fisheries & Aquaculture Science.

Copp, G. H., Warrington, S., & Wesley, K. J. (2008). Management of an ornamental pond as a conservation site for a threatened native fish species, crucian carp Carassius carassius. *Hydrobiologia*, *597*, 149–155. https://doi.org/10.1007/978-90-481-9088-1_13

Curry, C. J., Gibson, J. F., Shokralla, S., Hajibabaei, M., & Baird, D. J. (2018). Identifying North American freshwater invertebrates using DNA barcodes: are existing COI sequence libraries fit for purpose? *Freshwater Science* , *37*, 178–189. https://doi.org/10.1086/696613

Davies, B., Biggs, J., Williams, P., Whitfield, M., Nicolet, P., Sear, D., Bray, S., & Maund, S. (2008). Comparative biodiversity of aquatic habitats in the European agricultural landscape. *Agriculture, Ecosystems & Environment*, *125*, 1–8. https://doi.org/10.1016/j.agee.2007.10.006

Deiner, K., Bik, H. M., Mächler, E., Seymour, M., Lacoursière-Roussel, A., Altermatt, F., Creer, S., Bista, I., Lodge, D. M., de Vere, N., Pfrender, M. E., & Bernatchez, L. (2017). Environmental DNA metabarcoding: Transforming how we survey animal and plant communities. *Molecular Ecology*, *26*, 5872–5895. https://doi.org/10.1111/mec.14350

Deiner, K., Fronhofer, E. A., Mächler, E., Walser, J.-C., & Altermatt, F. (2016). Environmental DNA reveals that rivers are conveyer belts of biodiversity information. *Nature Communications*, *7*, 12544. https://doi.org/10.1038/ncomms12544

Dobson, M., Pawley, S., Fletcher, M., & Powell, A. (2012). *Guide to Freshwater Invertebrates*. Ambleside, Freshwater Biological Association.

Dudgeon, D., Arthington, A. H., Gessner, M. O., Kawabata, Z. -I., Knowler, D. J., Lévêque, C., Naiman, R. J., Prieur-Richard, A. -H., Soto, D., Stiassny, M. L. J., & Sullivan, C. A. (2006). Freshwater biodiversity: importance, threats, status and conservation challenges. *Biological Reviews of the Cambridge Philosophical Society*, *81*, 163–182. https://doi.org/10.1017/S1464793105006950

Edington, J. M., & Hildrew, A. G. (1995). *A revised key to the caseless caddis larvae of the British Isles with notes on their ecology*. Ambleside, Freshwater Biological Association.

Elbrecht, V., Hebert, P. D. N., & Steinke, D. (2018). Slippage of degenerate primers can cause variation in amplicon length. *Scientific Reports*, *8*, 10999. https://doi.org/10.1038/s41598-018-29364-z

Elbrecht, V., & Leese, F. (2017). Validation and Development of COI Metabarcoding Primers for Freshwater Macroinvertebrate Bioassessment. *Frontiers in Environmental Science*, *5*. https://doi.org/10.3389/fenvs.2017.00011

Elbrecht, V., Peinert, B., & Leese, F. (2017). Sorting things out: Assessing effects of unequal specimen biomass on DNA metabarcoding. *Ecology and Evolution*, *7*, 6918–6926. https://doi.org/10.1002/ece3.3192

Elbrecht, V., Taberlet, P., Dejean, T., Valentini, A., Usseglio-Polatera, P., Beisel, J. -N., Coissac, E.,
Boyer, F., & Leese, F. (2016). Testing the potential of a ribosomal 16S marker for DNA
metabarcoding of insects. *PeerJ*, 4, e1966. https://doi.org/10.7717/peerj.1966
Elbrecht, V., Vamos, E.E., Meissner, K., Aroviita, J. & Leese, F. (2017). Assessing strengths and
weaknesses of DNA metabarcoding-based macroinvertebrate identification for routine
stream monitoring. *Methods in Ecology and Evolution*, *8*, 1265–1275.
https://doi.org/10.1111/2041-210X.12789
Elliott, J. M. (2009). *Freshwater Megaloptera and Neuroptera of Britain and Ireland: keys to*
adults and larvae, and a review of their ecology. Ambleside, Freshwater Biological
Association.
Elliott, J. M., & Dobson, M. (2015). *Freshwater leeches of Britain and Ireland: keys to the*
Hirudinea and a review of their ecology. Ambleside, Freshwater Biological Association.
Elliott, J. M., & Humpesch, U. H. (2010). *Mayfly larvae (Ephemeroptera) of Britain and Ireland:*
keys and a review of their ecology. Ambleside, Freshwater Biological Association.
Emilson, C.E., Thompson, D.G., Venier, L.A., Porter, T.M., Swystun, T., Chartrand, D., Capell, S.,
& Hajibabaei, M. (2017). DNA metabarcoding and morphological macroinvertebrate metrics
reveal the same changes in boreal watersheds across an environmental gradient. *Scientific*
Reports, *7*, 12777. https://doi.org/10.1038/s41598-017-13157-x
Ficetola, G.F., Coissac, E., Zundel, S., Riaz, T., Shehzad, W., Bessière, J., Taberlet, P., &
Pompanon, F. (2010). An *In silico* approach for the evaluation of DNA barcodes. *BMC*
Genomics, *11*, 434. https://doi.org/10.1186/1471-2164-11-434
Friday, L. E. (1988). A key to the adults of British water beetles. *Field Studies*, *7*, 1–151.
Gee, J. H. R., Smith, B. D., Lee, K. M., & Griffiths, S. W. (1997). The ecological basis of freshwater
pond management for biodiversity. *Aquatic Conservation: Marine and Freshwater*
Ecosystems, *7*, 91–104. https://doi.org/10.1002/(SICI)1099-0755(199706)7:2<91::AID-
AQC221>3.0.CO;2-O
Geller, J., Meyer, C., Parker, M., & Hawk, H. (2013). Redesign of PCR primers for mitochondrial
cytochrome c oxidase subunit I for marine invertebrates and application in all-taxa biotic
surveys. *Molecular Ecology Resources*, *13*, 851–861. https://doi.org/10.1111/1755-
0998.12138
Gledhill, D. G., James, P., & Davies, D. H. (2008). Pond density as a determinant of aquatic
species richness in an urban landscape. *Landscape Ecology*, *23*, 1219–1230.
https://doi.org/10.1007/s10980-008-9292-x
Gledhill, T., Sutcliffe, D. W., & Williams, W. D. (1993). *British freshwater Crustacea*
Malacostraca: a key with ecological notes. Ambleside, Freshwater Biological Association.
Haase, P., Pauls, S. U., Schindehütte, K., & Sundermann, A. (2010). First audit of
macroinvertebrate samples from an EU Water Framework Directive monitoring program:
human error greatly lowers precision of assessment results. *Journal of the North American*
Benthological Society, *29*, 1279–1291. https://doi.org/10.1899/09-183.1
Haas, K., Köhler, U., Diehl, S., Köhler, P., Dietrich, S., Holler, S., Jaensch, A., Niedermaier, M., &
Vilsmeier, J. (2007). Influence of fish on habitat choice of water birds: a whole system
experiment. *Ecology*, *88*, 2915–2925. https://doi.org/10.1890/06-1981.1

Hänfling, B., Bolton, P., Harley, M., & Carvalho, G. R. (2005). A molecular approach to detect
hybridisation between crucian carp (*Carassius carassius*) and non-indigenous carp species
(*Carassius* spp. and *Cyprinus carpio*). *Freshwater Biology*, *50*, 403–417.
https://doi.org/10.1111/j.1365-2427.2004.01330.x
Harper, L. R., Griffiths, N. P., Lawson Handley, L., Sayer, C. D., Read, D. S., Harper, K. J.,
Blackman, R. C., Li, J., & Hänfling, B. (2019). Development and application of environmental
DNA surveillance for the threatened crucian carp (*Carassius carassius*). *Freshwater Biology*,
64, 93–107. https://doi.org/10.1111/fwb.13197
Harper, L.R., Lawson Handley, L., Hahn, C., Boonham, N., Rees, H.C., Lewis, E., Adams, I.P.,
Brotherton, P., Phillips, S. & Hänfling, B. (2019) Generating and testing ecological hypotheses
at the pondscape with environmental DNA metabarcoding: A case study on a threatened
amphibian. *Environmental DNA*, *2*, 184–199. https://doi.org/10.1002/edn3.57
Hassall, C., Hollinshead, J., & Hull, A. (2011). Environmental correlates of plant and invertebrate
species richness in ponds. *Biodiversity and Conservation*, *20*, 3189–3222.
https://doi.org/10.1007/s10531-011-0142-9
Hill, M. J., Hassall, C., Oertli, B., Fahrig, L., Robson, B. J., Biggs, J., Samways, M. J., Usio, N.,
Takamura, N., Krishnaswamy, J., & Wood, P. J. (2018). New policy directions for global pond
conservation. *Conservation Letters*, *142*, e12447. https://doi.org/10.1111/conl.12447
Hill, M. J., Heino, J., Thornhill, I., Ryves, D. B., & Wood, P. J. (2017). Effects of dispersal mode on
the environmental and spatial correlates of nestedness and species turnover in pond
communities. *Oikos*, *126*, 1575–1585. https://doi.org/10.1111/oik.04266
Hill, M. J., Sayer, C. D., & Wood, P. J. (2016). When is the best time to sample aquatic
macroinvertebrates in ponds for biodiversity assessment? *Environmental Monitoring and
Assessment*, *188*, 194. https://doi.org/10.1007/s10661-016-5178-6
Jarvis, L. E. (2010). Non-consumptive effects of predatory three-spined sticklebacks
(*Gasterosteus aculeatus*) on great crested newt (*Triturus cristatus*) embryos. *The
Herpetological Journal*, *20*, 271–275.
Jeffries, D. L., Copp, G. H., Maes, G. E., Lawson Handley, L., Sayer, C. D., & Hänfling, B. (2017).
Genetic evidence challenges the native status of a threatened freshwater fish (*Carassius
carassius*) in England. *Ecology and Evolution*, *7*, 2871–2882.
https://doi.org/10.1002/ece3.2831
Klymus, K. E., Marshall, N. T., & Stepien, C. A. (2017). Environmental DNA (eDNA)
metabarcoding assays to detect invasive invertebrate species in the Great Lakes. *PLoS ONE*,
12, e0177643. https://doi.org/10.1371/journal.pone.0177643
Leese, F., Sander, M., Buchner, D., Elbrecht, V., Haase, P., & Zizka, V. M. A. (2020) Improved
freshwater macroinvertebrate detection from eDNA through minimized non-target
amplification. *bioRxiv*, https://doi.org/10.1101/2020.04.27.063545
Legendre, P. (2014). Interpreting the replacement and richness difference components of beta
diversity. *Global Ecology and Biogeography: A Journal of Macroecology*, *23*, 1324–
1334. https://doi.org/10.1111/geb.12207

Legendre P., & Birks H. J. B. (2012). From Classical to Canonical Ordination. In: Birks H.,
Lotter A., Juggins S., Smol J. (eds) Tracking Environmental Change Using Lake Sediments.
Developments in Paleoenvironmental Research, vol 5. Dordrecht, Springer.
https://doi.org/10.1007/978-94-007-2745-8_8

Legendre, P., & Legendre, L. F. J. (2012). *Numerical Ecology*. The Netherlands, Elsevier Science.

Lemmens, P., Mergeay, J., De Bie, T., Van Wichelen, J., De Meester, L., & Declerck, S. A. J.
(2013). How to maximally support local and regional biodiversity in applied conservation?
Insights from pond management. *PLoS ONE*, *8*, e72538.
https://doi.org/10.1371/journal.pone.0072538

Leray, M., Yang, J. Y., Meyer, C. P., Mills, S. C., Agudelo, N., Ranwez, V., Boehm, J. T., & Machida,
R. J. (2013). A new versatile primer set targeting a short fragment of the mitochondrial COI
region for metabarcoding metazoan diversity: application for characterizing coral reef fish
gut contents. *Frontiers in Zoology*, *10*, 34. https://doi.org/10.1186/1742-9994-10-34

Lewis-Phillips, J., Brooks, S., Sayer, C. D., McCrea, R., Siriwardena, G., & Axmacher, J. C.
(2019). Pond management enhances the local abundance and species richness of farmland
bird communities. *Agriculture, Ecosystems and Environment*, 273, 130-140.
http://doi.org/10.1016/j.agee.2018.12.015

Lobo, J., Shokralla, S., Costa, M. H., Hajibabaei, M., & Costa, F. O. (2017). DNA metabarcoding
for high-throughput monitoring of estuarine macrobenthic communities. *Scientific Reports*,
7, 15618. https://doi.org/10.1038/s41598-017-15823-6

Macan, T. T. (1960). *A key to the British fresh-and brackish-water gastropods: with notes on
their ecology*. Ambleside, Freshwater Biological Association.

Maceda-Veiga, A., López, R., & Green, A. J. (2017). Dramatic impact of alien carp *Cyprinus carpio*
on globally threatened diving ducks and other waterbirds in Mediterranean shallow lakes.
Biological Conservation, *212*, 74–85. https://doi.org/10.1016/j.biocon.2017.06.002

Macher, J. -N., Vivancos, A., Piggott, J. J., Centeno, F. C., Matthaei, C. D., & Leese, F. (2018).
Comparison of environmental DNA and bulk-sample metabarcoding using highly degenerate
COI primers. *Molecular Ecology Resources*, *18*, 1456–1468. https://doi.org/10.1111/1755-
0998.12940

Mächler, E., Little, C. J., Wüthrich, R., Alther, R., Fronhofer, E. A., Gounand, I., Harvey, E.,
Hürlemann, S., Walser, J., & Altermatt, F. (2019) Assessing different components of diversity
across a river network using eDNA. *Environmental DNA*, 1, 290–301.
https://doi.org/10.1002/edn3.33

Marklund, O., Sandsten, H., Hansson, L. -A., & Blindow, I. (2002). Effects of waterfowl and fish
on submerged vegetation and macroinvertebrates. *Freshwater Biology*, *47*, 2049–2059.
https://doi.org/10.1046/j.1365-2427.2002.00949.x

Menetrey, N., Sager, L., Oertli, B., & Lachavanne, J. -B. (2005). Looking for metrics to assess the
trophic state of ponds. Macroinvertebrates and amphibians. *Aquatic Conservation: Marine
and Freshwater Ecosystems*, *15*, 653–664. https://doi.org/10.1002/aqc.746

Meusnier, I., Singer, G. A. C., Landry, J. -F., Hickey, D. A., Hebert, P. D. N., & Hajibabaei, M.
(2008). A universal DNA mini-barcode for biodiversity analysis. *BMC Genomics*, *9*, 214.
https://doi.org/10.1186/1471-2164-9-214

Nakagawa, S., & Cuthill, I. C. (2007). Effect size, confidence interval and statistical significance: a
practical guide for biologists. *Biological Reviews of the Cambridge Philosophical Society*, *82*,
591–605. https://doi.org/10.1111/j.1469-185X.2007.00027.x

Nichols, S. J., Kefford, B. J., Campbell, C. D., Bylemans, J., Chandler, E., Bray, J.P., Shackleton, M.,
Robinson, K. L., Carew, M. E., & Furlan, E. M. (2020) Towards routine DNA metabarcoding of
macroinvertebrates using bulk samples for freshwater bioassessment: Effects of debris and
storage conditions on the recovery of target taxa. *Freshwater Biology*, *65*, 607–620.
https://doi.org/10.1111/fwb.13443

Oertli, B., Joye, D. A., Castella, E., Juge, R., Cambin, D., & Lachavanne, J.-B. (2002). Does size
matter? The relationship between pond area and biodiversity. *Biological Conservation*, *104*,
59–70. https://doi.org/10.1016/S0006-3207(01)00154-9

Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R.,
O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., & Wagner, H. (2011)
vegan: Community Ecology Package. R package version 2.5-4. https://CRAN.R-
project.org/package=vegan

Paradis, E., & Schliep, K. (2018). ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics*, *35*, 526–528.
https://doi.org/10.1093/bioinformatics/bty633

Peres-Neto, P. R., Legendre, P., Dray, S., & Borcard, D. (2006). Variation partitioning of species
data matrices: estimation and comparison of fractions. *Ecology*, *87*, 2614–2625.
https://doi.org/10.1890/0012-9658(2006)87[2614:VPOSDM]2.0.CO;2

Preston, D. L., Hedman, H. D., Esfahani, E. R., Pena, E. M., Boland, C. E., Lunde, K. B., & Johnson,
P. T. J. (2017). Responses of a wetland ecosystem to the controlled introduction of invasive
fish. *Freshwater Biology*, *62*, 767–778. https://doi.org/10.1111/fwb.12900

R Core Team (2017). R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. https://www.R-project.org/

Savage, A. A. (1989). *Adults of the British aquatic Hemiptera Heteroptera: a key with ecological*
notes. Ambleside, Freshwater Biological Association.

Sayer, C. D., Copp, G. H., Emson, D., Godard, M. J., Zięba, G., & Wesley, K. J. (2011). Towards the
conservation of crucian carp *Carassius carassius*: understanding the extent and causes of
decline within part of its native English range. *Journal of Fish Biology*, *79*, 1608–
1624. https://doi.org/10.1111/j.1095-8649.2011.03059.x

Sayer, C., Shilland, E., Greaves, H., Dawson, B., Patmore, I., Emson, D., Alderton, E., Robinson,
P., Andrews, K., Axmacher, J., & Wiik, E. (2013) Managing Britain's ponds - conservation
lessons from a Norfolk farm. *British Wildlife*, *25*, 21–28.

Sayer, C.D., Emson, D., Patmore, I.R., Greaves, H.M., West, W.P., Payne, J., Davies, G.D., Tarkan,
A.S., Wiseman, G., Cooper, B., Grapes, T., Cooper, G. & Copp, G.H. (2020) Recovery of the
crucian carp *Carassius carassius* (L.): Approach and early results of an English conservation
project. *Aquatic Conservation: Marine and Freshwater Ecosystems*.

Schilling, E. G., Loftin, C. S., & Huryn, A. D. (2009a). Effects of introduced fish on
macroinvertebrate communities in historically fishless headwater and kettle lakes. *Biological*
Conservation, *142*, 3030–3038. https://doi.org/10.1016/j.biocon.2009.08.003

Schilling, E. G., Loftin, C. S., & Huryn, A. D. (2009b). Macroinvertebrates as indicators of fish absence in naturally fishless lakes. *Freshwater Biology*, *54*, 181–202. https://doi.org/10.1111/j.1365-2427.2008.02096

Stefanoudis, P. V., Sayer, C. D., Greaves, H. M., Davidson, T. A., Robson, H., Almeida, D., & Smith, E. (2017). Consequences of fish for cladoceran, water beetle and macrophyte communities in a farmland pond landscape: implications for conservation. *Fundamental and Applied Limnology*, *190*, 141–156. https://doi.org/10.1127/fal/2017/1004

Taberlet, P., Coissac, E., Pompanon, F., Brochmann, C., & Willerslev, E. (2012). Towards next-generation biodiversity assessment using DNA metabarcoding. *Molecular Ecology*, *21*, 2045–2050. https://doi.org/10.1111/j.1365-294X.2012.05470.x

Tréguier, A., Paillisson, J.-M., Dejean, T., Valentini, A., Schlaepfer, M. A., & Roussel, J.-M. (2014). Environmental DNA surveillance for invertebrate species: advantages and technical limitations to detect invasive crayfish *Procambarus clarkii* in freshwater ponds. *Journal of Applied Ecology*, *51*, 871–879. https://doi.org/10.1111/1365-2664.12262

Vamos, E.E., Elbrecht, V., & Leese, F. (2017). Short COI markers for freshwater macroinvertebrate metabarcoding. *Metabarcoding and Metagenomics*, *1*, e14625. https://doi.org/10.3897/mbmg.1.14625

Wacker, S., Fossøy, F., Larsen, B. M., Brandsegg, H., Sivertsgård, R., & Karlsson, S. (2019). Downstream transport and seasonal variation in freshwater pearl mussel (*Margaritifera margaritifera*) eDNA concentration. *Environmental DNA*, *1*, 64–73. https://doi.org/10.1002/edn3.10

Wallace, I. D., Wallace, B., & Philipson, G. N. (1990). *A key to the case-bearing caddis larvae of Britain and Ireland*. Ambleside, Freshwater Biological Association.

Weigand, H., Beermann, A. J., Čiampor, F., Costa, F. O., Csabai, Z., Duarte, S., Geiger, M. F., Grabowski, M., Rimet, F., Rulik, B., Strand, M., Szucsich, N., Weigand, A. M., Willassen, E., Wyler, S. A., Bouchez, A., Borja, A., Čiamporová-Zaťovičová, Z., Ferreira, S., Dijkstra, K. -D. B., Eisendle, U., Freyhof, J., Gadawski, P., Graf, W., Haegerbaeumer, A., van der Hoorn, B. B., Japoshvili, B., Keresztes, L., Keskin, E., Leese, F., Macher, J. N., Mamos, T., Paz, G., Pešić, V., Pfannkuchen, D. M., Pfannkuchen, M. A., Price, B. W., Rinkevich, B., Teixeira, M. A. L., Várbíró, G. & Ekrem, T. (2019) DNA barcode reference libraries for the monitoring of aquatic biota in Europe: Gap-analysis and recommendations for future work. *The Science of the Total Environment*, *678*, 499–524. https://doi.org/10.1016/j.scitotenv.2019.04.247

Williams, P., Whitfield, M., Biggs, J., Bray, S., Fox, G., Nicolet, P., & Sear, D. (2003). Comparative biodiversity of rivers, streams, ditches and ponds in an agricultural landscape in Southern England. *Biological Conservation*, *115*, 329–341. https://doi.org/10.1016/S0006-3207(03)00153-8

Wood, P. J., Greenwood, M. T., & Agnew, M. D. (2003). Pond biodiversity and habitat loss in the UK. *Area*, *35*, 206–216. https://doi.org/10.1111/1475-4762.00249

Wood, P. J., Greenwood, M. T., Barker, S. A., & Gunn, J. (2001). The effects of amenity management for angling on the conservation value of aquatic invertebrate communities in old industrial ponds. *Biological Conservation*, *102*, 17–29. https://doi.org/10.1016/S0006-3207(01)00087-8

1047 Zeale, M. R. K., Butlin, R. K., Barker, G. L. A., Lees, D. C., & Jones, G. (2011). Taxon-specific PCR
1048  for DNA barcoding arthropod prey in bat faeces. *Molecular Ecology Resources*, *11*, 236–244.
1049  https://doi.org/10.1111/j.1755-0998.2010.02920.x
1050 Zhang, Z., Schwartz, S., Wagner, L., & Miller, W. (2000). A greedy algorithm for aligning DNA
1051  sequences. *Journal of Computational Biology*, *7*, 203–214.
1052  https://doi.org/10.1089/10665270050081478
1053 Zizka, V. M. A., Leese, F., Peinert, B., & Geiger, M. F. (2018). DNA metabarcoding from sample
1054  fixative as a quick and voucher preserving biodiversity assessment method. *Genome*, *62*,
1055  122–136. https://doi.org/10.1139/gen-2018-0048
1056

1057 **Table 1.** Summary of analyses (Pairwise Tukey's HSD Test) statistically comparing alpha diversity
1058 at species-level and family-level between standard and unbiased methods.
1059

	Species-level			Family-level		
	Pairwise Tukey's HSD Test			Pairwise Tukey's HSD Test		
	Estimate ± SE	*Z*	*P*	Estimate ± SE	*Z*	*P*
Standard methods						
Sweep-netting and microscopy :DNA metabarcoding	-0.399 ± 0.129	-3.087	**0.006**	-0.284 ± 0.108	-2.630	**0.023**
Sweep-netting and microscopy : eDNA metabarcoding	-0.645 ± 0.127	-5.087	**<0.001**	-0.686 ± 0.102	-6.712	**<0.001**
DNA metabarcoding : eDNA metabarcoding	0.246 ± 0.121	2.030	0.105	0.402 ± 0.096	4.183	**<0.001**
Unbiased methods						
Sweep-netting and microscopy : DNA metabarcoding	-0.463 ± 0.128	-3.632	**<0.001**	-0.134 ± 0.106	-1.267	0.414
Sweep-netting and microscopy : eDNA metabarcoding	0.128 ± 0.132	-0.973	0.594	0.072 ± 0.110	0.650	0.792
DNA metabarcoding : eDNA metabarcoding	-0.335 ± 0.125	-2.671	**0.021**	-0.206 ± 0.108	-1.915	0.134

1060

1061 **Table 2.** Summary of analyses (GLM) statistically comparing alpha diversity at species-level and
1062 family-level between ponds with and without crucian carp using independent and combined
1063 standard methods.
1064

		Species-level						Family-level				
		GLM			LRT			GLM			LRT	
	df	Estimate ± SE	Z	P	χ²	P	df	Estimate ± SE	Z	P	χ²	P
Sweep-netting and microscopy	1	-0.139 ± 0.186	-0.746	0.456	0.556	0.456	1	-0.098 ± 0.158	-0.620	0.535	0.385	0.535
DNA metabarcoding	1	0.060 ± 0.141	0.429	0.668	0.184	0.668	1	-0.007 ± 0.122	-0.061	0.952	0.004	0.952
eDNA metabarcoding	1	-0.072 ± 0.194	-0.375	0.708	0.141	0.708	1	-0.173 ± 0.140	-1.234	0.217	1.524	0.217
Combined methods	1	-0.062 ± 0.122	-0.506	0.613	0.256	0.613	1	-0.128 ± 0.095	-1.349	0.177	1.822	0.177

1065
1066

1067 **Table 3.** Relative contribution of taxon turnover and nestedness-resultant to total beta diversity
1068 (Jaccard dissimilarity) using independent and combined standard methods. A value of 1
1069 corresponds to all sites containing different species.
1070

	Species-level			Family-level		
	Turnover	Nestedness-resultant	Total beta diversity	Turnover	Nestedness-resultant	Total beta diversity
Sweep-netting and microscopy	0.935 (98.01%)	0.019 (1.99%)	0.954 (100%)	0.867 (94.75%)	0.048 (5.25%)	0.915 (100%)
DNA metabarcoding	0.938 (98.53%)	0.014 (1.47%)	0.952 (100%)	0.885 (96.83%)	0.029 (3.17%)	0.914 (100%)
eDNA metabarcoding	0.917 (97.24%)	0.026 (2.76%)	0.943 (100%)	0.871 (95.29%)	0.043 (4.71%)	0.914 (100%)
Combined methods	0.927 (98.41%)	0.015 (1.59%)	0.942 (100%)	0.871 (96.46%)	0.032 (3.54%)	0.903 (100%)

1071

1072 **Table 4.** Summary of analyses (PERMANOVA) statistically examining variation in community
1073 composition between ponds with and without crucian carp at species-level and family-level
1074 using independent and combined standard methods.
1075

	Community similarity (PERMANOVA)							
	Species-level				Family-level			
	df	*F*	*R²*	*P*	df	*F*	*R²*	*P*
Sweep-netting and microscopy								
Turnover	1	1.673	0.095	**0.018**	1	1.198	0.070	0.358
Nestedness-resultant	1	-3.454	-0.275	0.949	1	-0.103	-0.007	0.620
Total beta diversity	1	1.369	0.079	**0.033**	1	1.136	0.066	0.306
DNA metabarcoding								
Turnover	1	1.305	0.075	0.137	1	1.668	0.094	0.098
Nestedness-resultant	1	-0.578	-0.038	0.664	1	-0.116	-0.007	0.717
Total beta diversity	1	1.163	0.068	0.176	1	1.277	0.074	0.197
eDNA metabarcoding								
Turnover	1	2.374	0.129	**0.001**	1	1.551	0.088	0.097
Nestedness-resultant	1	-2.515	-0.187	0.967	1	0.586	0.035	0.560
Total beta diversity	1	1.790	0.101	**0.004**	1	1.502	0.086	**0.041**
Combined methods								
Turnover	1	1.999	0.111	**0.001**	1	1.452	0.083	0.136
Nestedness-resultant	1	-1.663	-0.116	0.966	1	3.103	0.162	0.128
Total beta diversity	1	1.717	0.097	**0.001**	1	1.474	0.084	**0.027**

1076

Figure 1. Venn diagrams summarising taxon detection by standard **(a)** and unbiased **(b)** methods of invertebrate assessment. The number of invertebrate species **(i)** and families **(ii)** detected across the 18 study ponds by sweep-netting and microscopy (green circle), DNA metabarcoding (purple circle), and eDNA metabarcoding (orange circle) is displayed. Overlap in species or family detections between methods is shown within circle intersections.

Figure 2. Mean alpha diversity of invertebrates in ponds across Norfolk and East Yorkshire. Alpha diversity is displayed according to standard **(a)** and unbiased **(b)** methods of invertebrate assessment. Species **(i)** and family **(ii)** richness generated by sweep-netting and microscopy (green points), DNA metabarcoding (purple points), and eDNA metabarcoding (orange points) is shown. Boxes show 25th, 50th, and 75th percentiles, and whiskers show 5th and 95th percentiles.

Figure 3. Non-metric Multidimensional Scaling (NMDS) plots of invertebrate communities (Jaccard dissimilarity) produced by sweep-netting and microscopy (green points/ellipse), DNA metabarcoding (purple points/ellipse), and eDNA metabarcoding (orange points/ellipse) for the 18 study ponds. The turnover **(a)** and nestedness-resultant **(b)** partitions of total beta diversity **(c)** are shown at species-level and family-level for standard **(i, ii)** and unbiased **(iii, iv)** methods.

Figure 4. Mean alpha diversity of invertebrates in ponds with crucian carp (blue points) and without fish (grey points) across Norfolk and East Yorkshire. Alpha diversity at species-level **(a)** and family-level **(b)** is shown according to standard methods of invertebrate assessment: sweep-netting and microscopy **(i)**, DNA metabarcoding **(ii)**, eDNA metabarcoding **(iii)**, and all methods combined **(iv)**. Boxes show 25th, 50th, and 75th percentiles, and whiskers show 5th and 95th percentiles.

1111
1112
Figure 5. Non-metric Multidimensional Scaling (NMDS) plots of species-level invertebrate
communities (Jaccard dissimilarity) from ponds with crucian carp (blue points/ellipse) and
without fish (grey points/ellipse) across Norfolk and East Yorkshire. The turnover **(a)** and
nestedness-resultant **(b)** partitions of total beta diversity **(c)** are shown according to standard
methods of invertebrate assessment: netting and microscopy **(i)**, DNA metabarcoding **(ii)**, eDNA
metabarcoding **(iii)**, and all methods combined **(iv)**.
1119

Figure 6. The relative contribution of biotic (crucian carp presence-absence) and abiotic variables (pond area) to species-level turnover **(a)** and total beta diversity **(b)** when the combined invertebrate data from all three standard sampling methods were considered. Values within circles and circle intersections represent the adjusted R^2 values. Significant variables are indicated by asterisks (* = $P < 0.05$).