Article

Title: Recombination and purifying selection preserves covariant movements of mosaic

SARS-CoV-2 protein S

Authors: Massimiliano S. Tagliamonte[1,2#], Nabil Abid[3,4#], David A. Ostrov[2], Giovanni

Chillemi[5,6], Sergei L. Kosakovsky Pond[7], Marco Salemi[1,2*], Carla Mavian[1,2*]

Affiliations: 1. Emerging Pathogen Institute, University of Florida, Gainesville, FL, USA;

2. Department of Pathology, immunology and Laboratory Medicine, University of Florida,

Gainesville, FL, USA; 3. Laboratory of Transmissible Diseases and Biological Active

Substances LR99ES27, Faculty of Pharmacy, University of Monastir, Rue Ibn Sina,

5000, Monastir, Tunisia

4. High Institute of Biotechnology of Sidi Thabet, Department of Biotechnology,

University of Manouba, BP-66, 2020, Ariana-Tunis, Tunisia; 5. Department for

Innovation in Biological, Agro-food and Forest systems, DIBAF, University of Tuscia, via

S. Camillo de Lellis s.n.c., 01100 Viterbo, Italy; 6. Institute of Biomembranes,

Bioenergetics and Molecular Biotechnologies, IBIOM, CNR, Via Giovanni Amendola,

122/O, 70126, Bari, Italy; 7 Department of Biology, Temple University, Philadelphia, PA

19122,USA

#= authors contributed equally to this work

*= corresponding author, cmavian@ufl.edu

*= corresponding author, salemi@pathology.ufl.edu

Keywords: SARS-CoV-2, COVID-19, recombination, negative selection, pre-fusion,

pangolin, bat, furin-like cleavage site, hACE2, receptor binding domain

Abstract

In depth evolutionary and structural analyses of severe acute respiratory syndrome
coronavirus 2 (SARS-CoV-2) isolated from bats, pangolins, and humans are necessary
to assess the role of natural selection and recombination in the emergence of the current
pandemic strain. The SARS-CoV-2 S glycoprotein unique features have been
associated with efficient viral spread in the human population. Phylogeny-based and
genetic algorithm methods clearly show that recombination events between viral
progenitors infecting animal hosts led to a mosaic structure in the S gene. We identified
recombination coldspots in the S glycoprotein and strong purifying selection. Moreover,
although there is little evidence of diversifying positive selection during host-switching,
structural analysis suggests that some of the residues emerged along the ancestral
lineage of current pandemic strains may contribute to enhanced ability to infect human
cells. Interestingly, recombination did not affect the long-range covariant movements of
SARS-CoV-2 S glycoprotein monomer in pre-fusion conformation but, on the contrary,
could contribute to the observed overall viral efficiency. Our dynamic simulations
revealed that the movements between the host cell receptor binding domain (RBD) and
the novel furin-like cleavage site are correlated. We identified threonine 333 (under
purifying selection), at the beginning of the RBD, as the hinge of the opening/closing
mechanism of the SARS-CoV-2 S glycoprotein monomer functional to hACE2 binding.
Our findings support a scenario where ancestral recombination and fixation of amino
acid residues in the RBD of the S glycoprotein generated a virus with unique features,
capable of extremely efficient infection of the human host.

Introduction

Coronaviruses (CoVs) are single strand, positive sense single-stranded RNA (+ssRNA)
viruses, with diverse tropism, able to infect respiratory, enteric, and hepatic tissues of
several species (Fehr and Perlman 2015). Within the CoVs family, Beta CoVs have
repeatedly proven the ability to shift from their natural reservoir and adapt to human
hosts (Su, et al. 2016). Several CoVs, such as HCoV-229E, HCoV-OC43, HCoV-HKU1,
and HcoV-NL63, are mostly associated with mild symptoms (Bucknall, et al. 1972; Woo,
et al. 2005). Few CoVs caused limited outbreaks of more severe illness, such as the
severe acute respiratory syndrome (SARS) and Middle East respiratory syndrome
(MERS) (Leung, et al. 2004; Zaki, et al. 2012), and the recently emerged SARS-CoV-2
(Wu, et al. 2020)). First reported in China in December 2019, CoV disease 2019

(COVID-19) quickly turned into a pandemic, with nearly 7,000,000 cases and hundreds of thousands of deaths reported worldwide as of June 2020 (Kamel Boulos and Geraghty 2020). An early study hypothesized SARS-CoV-2 emergence from snakes (Ji, et al. 2020), but further phylogenetic analyses traced its origins to CoVs circulating in bat (Zhou, Chen, et al. 2020; Zhou, Yang, et al. 2020) and pangolin (Xiao, et al. 2020). Contacts between different animal species prior to human exposure might facilitate host shift, possibly through adaptation to intermediate hosts, such as camels in the case of MERS-CoV (Dudas and Rambaut 2016; Su, et al. 2016) or civets or pigs in the case of SARS-CoV-1 (Wang and Eaton 2007).

While most RNA viruses use low-fidelity RNA polymerases, CoV genomes encode a proofreading and mismatch correction pathway, resulting in less error prone replication and translating into lower mutation rates (Ferron, et al. 2018; Li, Wang, et al. 2020). This feature might ultimately limit the mutagenic variability of the virus, and recombination might prove a prominent driving factor in the evolution and expansion of species range and cellular tropism. Recombination can take place whenever a single host is infected with multiple, genetically different, strains, and is an important factor driving viral diversity (Simon-Loriere and Holmes 2011; Mavian, et al. 2017). Recombination has been reported as a possible mechanism favoring cross-species transmission of CoVs and increasing their adaptability to new host (Graham and Baric 2010; Su, et al. 2016). There is copious evidence of such events having occurred in the ancestral history of both MERS-CoV and SARS-CoV-1 (Lau, et al. 2015; Dudas and Rambaut 2016; Anthony, et al. 2017) and CoVs in general (Lai, et al. 1985; Decaro, et al. 2009; Terada, et al. 2014; Tian, et al. 2014), but evidence regarding the recombinant origin of SARS-CoV-2 is not conclusive (Ji, et al. 2020; Liu, et al. 2020; Paraskevis, et al. 2020; Zhou, Chen, et al. 2020). Paraskevis *et al.*, who made the case for a non-recombinant origin of the virus, used a limited dataset, which did not include pangolin CoVs (Paraskevis, et al. 2020). Based on genetic similarity plots, Ji *et al.* and Liu *et al.* hypothesized a possible recombination event with an undetermined viral isolate in the ancestry of SARS-CoV-2 (Ji, et al. 2020; Liu, et al. 2020). A very recent publication by Li *et al.* (Li, Giorgi, Marichannegowda, et al. 2020) also hypothesized a recombination event between bat and pangolin viruses as progenitor of the human CoV-2. While recent studies have used simple recombination detection algorithms, leading to various results that may depend on the assumptions of the specific algorithm and the data set of choice (Boni, et al. 2020; Li, Giorgi, Marichann, et al. 2020; Wong, et al. 2020), we performed an in-depth

analysis of SARS-CoV-2 and closely related SARS-CoV-1 and SARS-like CoVs lineages

using seven different methods, including phylogeny-based and genetic algorithms (see

Methods section) (Kosakovsky Pond, et al. 2006; Martin, et al. 2015), with the purpose

of achieving higher sensitivity (Jia, et al. 2018). Our analysis in fact revealed that the

recombination pattern of the S glycoprotein is far more complex than previously thought.

Because of the lack of sensitive recombination methods (Li, Giorgi, Marichannegowda,

et al. 2020) and the comparison with only close relative of SARS-CoV-2, i.e. bat-SARS-

CoV-2 RaTG13 and RmYN02 (Zhou, Chen, et al. 2020), it is possible that the likely

mosaic nature of the SARS-CoV-2 genome was masked.

The SARS-CoV-2 S glycoprotein trimer is responsible for the crucial step of fusion of the

viral membrane with the host cell membrane (Gallagher and Buchmeier 2001; Simmons,

et al. 2013). Complex structural rearrangements follow the interaction of RBD (that must

be in an exposed state referred as "up" conformation) with the peptidase domain (PD) of

hACE2 (Wrapp, et al. 2020; Yan, et al. 2020). Conversely, the "down" conformation

corresponds to the receptor-inaccessible state. The RBD up/down rotation is made

possible by a hinge region at the N-term of RBD (Wrapp, Wang et al. 2020) and our

Molecular Dynamics (MD) simulations identified threonine 333 as the hinge residue. A

two-step sequential protease cleavage model has been proposed for activation of CoV S

proteins, as shown for SARS-CoV and MERS-CoV (Belouzard, et al. 2009; Millet and

Whittaker 2014), priming cleavage at the S_1/S_2 domain junction (Li 2016; Gui, et al.

2017), allowing the projection of the fusion peptide toward the target cell membrane

(Bosch, et al. 2008) and activating cleavage on S2' site (Ou, et al. 2020).

An important peculiarity of SARS-CoV-2 as compared to other CoVs is the presence of a

furin-like cleavage sequence (Madu, et al. 2009; Lai, et al. 2017). Exposure of this

cleavage site to the soluble furin protease is an important step in the viral infection

(Wrapp, et al. 2020), and explain, at least in part, the enhanced infectivity and

pathogenicity of the virus (Bosch, et al. 2003).

Results

On the origin of SARS-CoV-2 lineage: recombination in bat and pangolin

Since the human CoV isolates are ~96% identical to bat-SARS-CoV-2 (Andersen, et al.

2020), but ACE2

binding residues F486, Q493, S494 and N501 (Figure 1c) are shared with pangolin

SARS-CoV-2 lineage b isolates, we investigated the evolutionary relationships among

available full genome sequences (bat and pangolin-related SARS-CoV-2 isolates, bat-

SL-CoVs, SARS-CoV-1, and SARS-like isolates) using a neighbor network (NNet) based

algorithm (Bryant and Moulton 2004). Unlike standard phylogenetic inference methods,

network approaches can naturally represent the presence of conflicting phylogenetic

signal in the data (which is a hallmark of recombination), as we found in the analysis of

the coronavirus genomes (Figure 1a). Conflicting phylogenetic signal in the data,

however, is not sufficient to infer recombination (Huson and Bryant 2006). To test

whether the NNet topology is due to recombination we used the pairwise homoplasy

index (PHI) test (Bruen, et al. 2006), and the Benjamini-Hochberg correction for multiple

testing (Benjamini and Hochberg 1995). Recombination signal ($q \ll 10^{-6}$) was detected

among all included isolates (Figure 1a). SARS-CoV-1 and human SARS-CoV-2

monophyletic clades do not display net-like topology, and recombination signal of a sub-

dataset including only these lineages was not significant ($q = 0.14$). SARS-like and bat-

SL-CoVs, on the other hand, show a high degree of conflicting phylogenetic signals

reflected by intricate networks (Figure 1a). SARS-like isolates are ancestors of SARS-

CoV-1 isolates, bat-SL-CoVs stand as intermediate link between SARS-CoV-1 and

SARS-CoV-2 isolates. Bat-SARS-CoV-2 RaTG13 (NCBI accession no. MN996532;

GISAID id EPI_ISL_402131, isolated from bat in 2013 in China) and pangolin-SARS-

CoV-2 isolates (EPI_ISL_410538-43 isolates denoted with "a" ; and EPI_ISL_412860

and EPI_ISL_410721 isolates indicated as "b", were isolated in 2017 and 2019 in China

respectively, Figure 1a) are ancestors of SARS-CoV-2 circulating in humans. When

scanning sub-sets of the alignment including only SARS-CoV-2 and its ancestors,

recombination signal was highly significant ($q < 1 \times 10^{-6}$). A signal for recombination was

also detected for SARS-CoV-2 with pangolin-SARS-CoV-2 ancestral isolates ($q = 0.03$),

and when testing SARS-CoV-2 bat (EPI_ISL_402131) and pangolin isolates ($q = 0.002$).

We next scanned the SARS-CoV-2 genomes for putative recombination breakpoints

using RDP, GENECOV, MAXCHI, CHIMAERA, SISCAN, and 3SEQ, implemented in

RDP4 (Martin, et al. 2015). This pipeline allows for the identification of recombinants as

well as potential their parental isolates - if present in the sampled sequences. Sixty-eight

potential recombination events distributed along the genome were found in bat-SL-

CoVs, SARS-CoVs, and SARS-like CoVs isolates (Table S1). We found that human

SARS-CoV-2 isolates were potentially either parental or recombinant sequences in

recombination events with pangolin-SARS-CoV-2 isolated in 2019 and bat-SARS-CoV-2

genomes. Uncertainty in the estimation (the parental may be the recombinant

sequences, or vice versa) suggests missing links in SARS-CoV-2 lineages, which is

likely, given the sparse sampling of the wildlife coronavirus diversity, or low sensitivity of

the recombination algorithms. Because the recombination event between CoVs of

pangolin "b" and human origin involved the region of the S glycoprotein that binds to the

cell receptor (Figure 1b-c and Table S1), exactly matching the ACE2 binding region

where residues are shared between pangolin and human (Liu, et al. 2020), we conclude

that the SARS-CoV-2 human lineage is the result of recombination between a strain

belonging to the pangolin "b" lineage, potential minor parental, with a strain belonging to

the bat lineage close to bat-SARS-CoV-2 RaTG13.

We corroborated the potential recombination event by phylogenetic inference based on

the recombinant and non-recombinant genome fragments from human, bat, and

pangolin CoV-2 sequences, after assessing for the presence of phylogenetic signal

(Table S2 and S3). Trees inferred using the recombinant region (part of the S

glycoprotein) supported pangolin and CoV-2 ancestral relationship, while the segment

derived by the major parent kept the CoV-2 clade clustering with the bat sequence

(Figure S1).

We investigated further recombination patterns with a very sensitive genetic algorithm,

GARD, based on a likelihood model selection procedure that searches multiple

sequence alignments for evidence of recombination breakpoints (Kosakovsky Pond, et

al. 2006). Extensive simulations have shown that such an in-depth analysis can detect

recombination events with higher power and accuracy than tools based on phylogeny

discordance among large genomic segments. Indeed, GARD detected a more complex

mosaic-like structure of the S gene (Figure 1d), comprised of at least four distinct non-

recombinant segments, in addition to a segment (S gene nucleotide position 1502-2262)

with statistically significant evidence of further intra-segment recombination (Table S4).

The mosaic is the result of multiple independent events involving different ancestral

lineages during the evolutionary process that eventually gave rise to the current

pandemic strain. The first two non-recombinant segments of the S gene (nt 1-906 and nt

907-1316) show a human/bat ancestor (that subsequently splits in Bat-SL-CoVs and

SARS-CoV-2) sharing a common ancestor with Pangolin A and Pangolin B CoVs,

respectively. The third segment (nt 1317-1501), which overlaps with the one identified by

RDP4 analyses involving the RBD domain (Table S4), confirms its SARS-CoV-

2/pangolin B CoVs common ancestry. The fourth segment (nt 1502-2262) appears to be

itself a complex mosaic, although exact location of each putative breakpoint could not be

calculated given the complexity of all possible patterns. The fifth segment (nt 1502-2262)

shows, again, the human/bat ancestor sharing a common ancestor with Pangolin B

CoVs. Overall, we hypothesize that the evolutionary past of the currently circulating

pandemic strains included multiple recombination events between an ancestor of human

SARS-CoV-2, or a closely related to bat-SARS-CoV-2, and an ancestor of pangolin-

SARS-CoV-2 lineage b (but not necessarily in a pangolin host). Given that some regions

in the SARS-CoV-2 genome show similarity to these clades (Figure S2), it is likely that

one or more missing links among these strains have yet to be sampled in the animal

reservoir.

Prediction of recombination hot- and cold- spots in SARS-CoV-2

Recombination hotspots analysis was performed through a permutation test in RDP4,

which randomly distributes the inferred breakpoints across the genome on the variable

nucleotide positions between the three sequences involved in the event. From the

permutations, density plots are drawn, and the distribution of observed breakpoints is

compared to the permutations results. Permuting the detected breakpoints relative to

variable positions takes into account the fact that breakpoints might be easier/more

difficult to detect based on the local nucleotide diversity. Segments that had more

observed breakpoints than the local maximum number of breakpoints of 99% of 1,000

permutations were considered hotspots (p=0.01); in the same fashion, segments that

had fewer breakpoints than the local minimum of 99% of permutations were considered

coldspots (Heath, et al. 2006). The algorithm uses a sliding window of a set size, which

is moved across the genome one nucleotide at the time. The size of the sliding window

may affect the sensitivity of the test, since smaller window sizes might detect tightly

clustered hotspots, but might miss dispersed hotspots or coldspots. Thus, we used the

windows size range 100-200 as suggested by the program authors. We plotted the

locations of inferred recombination breakpoints and analyzed their distribution through

permutations to predict hot- and coldspots of recombination across the coronavirus

genomes (Figure 2).

We identified nine potential recombination hotspots based on an analysis with sliding

windows of size 100 bp. Seven of them were located in the polyprotein segment (Figure

2), which spans three quarters of the viral RNA; the remaining two involved protein E,

orf8, and protein N. It has been reported that the *orf8* gene plays a role in regulating the

viral replication in the host cell (Muth, et al. 2018), and was probably acquired by the

whole lineage of SARS-CoV-1 from the SARS-related CoV from Greater Horseshoe bats

(SARSr-Rf-BatCoVs) following a recombination event with a Chinese horseshoe bat

coronavirus (SARSr-Rs-BatCoV) (Lau, et al. 2015). Three recombination coldspot

regions were detected using a sliding window of size 200 (Figure 2). Despite the S

glycoprotein being recombinant in nature, the N-terminal region holds two coldspots

away from the recombination breakpoints: the first is located between residues R34 to

D53, while the second between residues F201 to L229, (p ≤0.01). Another coldspot was

found in the N-terminal region of Pp1ab. Coldspots might be possibly related to

maintaining specific segments of the protein, or to the function of the viral transcription

regulatory sequences (Kim, et al. 2020).

SARS-CoV-2 spike (S) glycoprotein acquired unique features driven by

recombination and purifying selection

We provide a full description of all residues unique to the S glycoprotein of SARS-CoV-2

isolates circulating among humans by comparison with SARS-CoV-2 from bat and two

pangolin (lineage "a" isolates isolated from pangolin in 2017 in China; and "b" isolates

isolated from pangolin in 2019 in China) (Table S5). The overall structure of the SARS-

CoV-2 has significant structural homology to its SARS-CoV-1 counterpart (Wang, et al.

2020). In comparison with the latter available structure (Gui, et al. 2017), and a recently

published study (Lan, et al. 2020), the SARS-CoV-2 S glycoprotein is composed of the

S_1 subunit, that binds to hACE2 (Li 2016), and the S_2 subunit which plays a role in

membrane fusion (Bosch, et al. 2003). In particular, S_1 is composed of an N-terminal

domain (NTD, residues 14-294 in SARS-CoV-1 and 14-303 in SARS-CoV-2) and three

C-terminal domains (CTD1, CTD2, and CTD3) (Figure 3). The CTD1 (residues 320-516

for SARS-CoV-1 and residue 333-529 for SARS-CoV-2, see Methods) functions as the

RBD, which allows SARS-CoV-2 to directly bind to the PD of the hACE2 (Li, et al. 2005).

For simplicity, we show canonical hACE2 binding residues F486, Q493, S494 and N501,

as reported by Andersen *et al* (Andersen, et al. 2020). The S_2 subunit is located

downstream of S_1 after the S_1/S_2 cleavage site (residue 667 for SARS-CoV-1 and

residue 684 for SARS-CoV-2). The majority of the residues that differentiate human

SARS-CoV-2 S glycoprotein from bat or pangolin ones were found within the S_1 subunit

(Figure 3a, Table S6). While the S glycoproteins of MERS-CoV, HcoV-OC43, and HKU1

display the canonical (R/K)-(2X)n-(R/K)* (or RXXR*SA) motif for the S1/S2 cleavage (Li

2016; Gui, et al. 2017), the S glycoprotein of SARS-CoV-2 presents a solvent exposed

furin-like cleavage sequence (PRRARS*V) in positions 681-684 (Andersen, et al. 2020),

similar to the pattern found in a recently collected bat isolate (Zhou, Chen, et al. 2020)

(Figure 3). The cleavage site within the S1/S2 junction of the S glycoprotein is the target

for furin protease and essential for SARS-CoV-2 infection of humans (Hoffmann, et al.

2020). The high activity of the human furin, compared to bat, may enhance the kinetic of

the infection, as the cleavage of S1 and S2 is a crucial step for virus entry into cells.

For selection analyses, we retained only unique sequences (removing all other identical

sequences) resulting in an alignment of 119 sequences. Because of the negative impact

of recombination events on selection analyses (Schierup and Hein 2000; Posada and

Crandall 2002; Shriner, et al. 2003), we used the additional partitioning identified by

GARD for the selection analyses. On lineages that separate host clades (Figure 4), the

S glycoprotein is under strong purifying selection (mean dN/dS = 0.0165), and 582

variable residues out of 838 found to be evolving with dN/dS < 1 (p-value ≤ 0.1)

(Kosakovsky Pond and Frost 2005), including 106 residues out of 197 located in the

RBD (Table S7). The recombination coldspots hosted a proportion of sites under

purifying selection similar to the rest of the alignment. There was no evidence of

episodic diversifying selection, either at the level of the entire alignment (dN/dS = 0.0

with weight 86.71%, dN/dS = 0.09 with weight 13.28%, p = 0.5) (Murrell, et al. 2015), or

at individual sites (p < 0.1) (Murrell, et al. 2012), and no evidence was found of lineage

specific episodic adaptation (p < 0.05). Focusing on internal branches for selection

analyses removes the biasing effects of unresolved intra-host evolution or sequencing

errors (Smith, et al. 2015). To identify sites which may be evolving adaptively in the

SARS-CoV-2 clade, we tested for episodic diversifying selection only for internal

branches in the human clade (Murrell, et al. 2012) and identified twenty-one such sites,

eight in S_1 (p < 0.1) (Figure 4, Table S6). Next, we investigated whether residues were

evolving under the same selective pressure (Poon, et al. 2007). A total of five pairs of

codons were found to coevolve together at posterior probability ≥ 0.9 (Figure S3); some

of these residues are involved in epistatic interactions between residues in the RBD and

external residues.

Polybasic furin-like cleavage and prefusion conformation correlate with "up"

conformation

The RBD domain of SARS-CoV-2 S glycoprotein for the pre-fusion conformation rotates

outward from center of the triangular head to an "up" position, while this conformation for

SARS-CoV is "down"(Gui, et al. 2017; Wrapp, et al. 2020) (Figure 3d). The recognition

step between SARS-CoV-2 RBD and hACE2 is a dynamic process in which the protein

surface of both partners adapts one to the other, in line with cryo-EM evidences that

shows how the receptor binding can open up RBD (Song, et al. 2018). In general, the

RBD of each S monomer, prior to CoV infection, is buried in the inactive "down"

conformation (prefusion state) and cannot bind to hACE2 due to steric clash (Wrapp, et

al. 2020; Yan, et al. 2020). In the course of infection, one RBD monomer turned "up" to

expose enough space to hACE2 (postfusion state), inducing further conformational open

and loose for proteolysis (Gui, et al. 2017; Song, et al. 2018). Homology modeling of

SARS-CoV-2 S glycoprotein showed a "up" conformation of one SARS-CoV-2 S

monomer during the prefusion state, which was not showed for SARS-CoV-1.

We performed dynamic simulations (MD) to understand whether the genotype changes

that we identified earlier played an important role in shaping the structure of the S

glycoprotein of SARS-CoV-2, fundamental for viral binding to human cells. We carried

out essential dynamics analysis on the MD trajectories, in order to separate the large

collective protein movements connected to functional properties from the small and

uninteresting motions (Amadei 1993). This analysis is based on the diagonalization of

the covariance matrix built from the atomic fluctuations after the removal of the

translational and rotational movement. It is usually applied on the c-alpha atoms only,

since they describe the motion of the protein main chain. Interestingly, projection of the

MD molecular trajectory along the first eigenvector (i.e. the direction that contains the

majority of the protein total motion as shown by eigenvector values) is dominated by the

up/down (opening/closing) motion of the RBD (see Figure 5a-b and Suppl. Movies 1-2).

We identified threonine 333 (corresponding to T333 and T331 in bat and pangolin,

respectively) as the hinge of this movement. Noticeably, we verified that this motion is

conserved within the SARS-CoV-2 lineage, and therefore is observed in human, bat, and

pangolin S glycoproteins (Figure 5f-g and h-i, respectively). It is also worth noting that

residues 325-344 (mostly under negative/purifying selection) are nearly completely

conserved between SARS-CoV-2, bat and pangolin CoVs (Figure 3a), in line with the

key role of this protein region in guiding the up/down RBD movement.

The conformational flexibility of the S protein during the sampled 100 ns trajectory was

assessed by means of root mean square fluctuations (RMSF), calculated after

superimposing each individual structure of a trajectory onto the initial structure by means

of least-squares fitting, to remove rotational and translational motions. Projection of the

MD molecular trajectory along the first three eigenvectors (that cumulatively contain 90%

of the total protein motion) allowed to highlight co-variant movements (simultaneously

changing) among long distance S glycoprotein regions, functional to the protein

rearrangement during the pre-fusion step (Figure 5c-e). In detail, projection along

eigenvector 1 (Figure 5c) shows a correlated movement between RBD and the region

828-852 (highlighted by cyan lines in Figure 5c), encompassing the C-terminal portion of

the fusion peptide, defined by residues 816-833. The up conformation of RBD, in

particular, corresponds to a more solvent exposed conformation of the fusion peptide,

that is functional to the fusion of the viral and the host cell membranes (Figure S4 and

Suppl. Movie 3). Projection along eigenvector 2 (Figure 5d) shows a correlated

movement between a more restricted portion of RBD (residues 470-494, highlighted by

red lines in Figure 5d), and again the previously defined region 828-852. A third peak in

RMSF along this eigenvector is observed in the region 620-640, in CTD3. It is significant

that the indicated portion of RBD includes three out of the four hACE2 binding residues,

i.e. F486, Q493 and S494, demonstrating a direct structural communication between the

ACE2 binding region and the fusion peptide (see Figure S5 and Suppl. Movie 4).

Projection along eigenvector 3 (Figure 5e), finally, showed a correlated movement

between residues 244-265 in the NTD region (blue lines in Figure 5e), residues 460-494

in CTD1 (red lines in Figure 5e), and residues 673-692 (orange lines in Figure 5e), which

encompass the furin-like cleavage site typical of SARS-CoV-2. Also in this case, the up

conformation corresponds to a more solvent exposed surface of the furin-like cleavage

site. A peak of fluctuation for the furin-like cleavage site is observed also along

eigenvector 2 (Figure 5d), even though of minor amplitude as compared to the

highlighted three co-variant regions (i.e. hACE2 binding residues, fusion peptide and res

620-640 in CTD3) along eigenvector1. The long-range effects between the hACE2

binding residues and the furin-like cleavage site (see Figure S6 and Suppl Movie 5),

might therefore be functional to the novel acquired capabilities of this virus. This motion,

paired with the open/up conformation of SARS-CoV-2 glycoprotein that exposes the

whole interaction interface of RBD and the fusion peptide to hACE2 and host cell

membrane, might explain why SARS-CoV-2 has been shown greater infectivity and

spread than previous CoVs.

Discussion

Recombination is a hallmark of CoV evolution (Graham and Baric 2010). Although our

results clearly point to several recombination events occurred between ancestral

pangolin- and bat- SARS-CoV-2 lineages, the limited sampling of these animal hosts

makes difficult to draw definitive conclusions about its origin. According to our results,

one of these events led to the current configuration of SARS-CoV-2 ACE2 binding

residues, including recently described residues forming hydrogen bonds with the host
cell receptor (Yan, et al. 2020).
Current algorithms cannot fully distinguish recombinant sequences from the parental
ones, at least for some recombination events. Yet, the lack of positive diversifying
selection on the branch, in the S glycoprotein phylogeny, that leads to the recent human
CoV-2 lineage, and the strong purifying negative selection exerted on the majority of the
S glycoprotein codons, support our hypothesis that recombination constitutes the main
drive to escape a local fitness maximum (Weinreich and Chao 2005), successfully
expanding host tropism. It is likely that the ancestral lineages that actually recombined is
part of a yet unsampled population. A more extensive sampling of wildlife CoVs would
help in clarifying the events that led to the jump to humans. Our analyses were based on
sequences obtained from strains circulating relatively early in the epidemic; future work
including later strains' genomes might discover marks of further adaptation to the new
host, as new mutations emerge and get fixed in the human population.
Despite the frequency of recombination events involving the S glycoprotein, the N-
terminal region appears to host two recombination coldspots. While
mutation/recombination events in the glycoprotein RBD could result in expanded viral
tropism, recombination coldspots and negative/diversifying selective pressure limit
variability, thus maintaining essential structure and function. It must be noted that the
high diversity between CoVs in the glycoprotein region, as compared to the rest of the
alignment, might have an impact on this analysis, thus *in vitro* studies might be needed
to verify these findings. Yet, while RBD is definitively a good target for therapeutic
development, regions in coldspots or those undergoing negative/diversifying selective
pressure may be helpful for successful vaccine design.
SARS-CoV-2 S glycoprotein gained specific features that have facilitated the successful
spread of the virus as compared to its predecessors.
SARS-CoV-2 S glycoprotein presents two notable changes: a favorable furin-like
cleavage site (Andersen, et al. 2020), and RBD up in pre-fusion mode (Wrapp, et al.
2020). Furin is ubiquitously expressed at different levels in all tissues (Schalken, et al.
1987; Hatsuzawa, et al. 1990), but kinetics of furin-like enzyme activity are different
among species – e.g., human lung cells display quicker kinetic than bat (Pteropus
alecto) (El Najjar, et al. 2015) – and may not fully explain SARS-CoV-2 boosted
infectivity. Thus, the assumption that the insertion of a furin-cleavage site allowed a bat
CoV to gain the ability to infect humans facilitating the host specie jump once the

conditions were favorable cannot be excluded. On the contrary S glycoprotein of

previously circulating SARS-CoV isolates do not have a favorable furin-like cleavage site

(Andersen, et al. 2020) and remains un-cleaved after biosynthesis (Belouzard, et al.

2009). Furin has the potential to cleave specifically viral envelope glycoproteins, to

enhance viral fusion with host cell membranes (Izaguirre 2019), thereby any

variation/mutational changes in amino acid composition at cleavage sites may impact

tissue and cell tropism, host range, and pathogenesis (Klenk and Garten 1994), as

reported for other respiratory viruses (Klenk and Garten 1994; Peng, et al. 2017;

Izaguirre 2019). MD simulations correlated more solvent-exposed surface in furin-like

cleavage site and fusion peptide the S protein up conformation, identifying a conserved

region that acts as a hinge in the opening/closing mechanism of all the SARS-CoV-2

lineage. Development of inhibitors of such motion may reduce substantially pre-fusion

conformation and therefore the infectivity of SARS-CoV-2. Finally, the presence of

coevolving sites might derive from host tropism; recent *in silico* and *in vitro* studies have

already found evidence of epistatic effects in coronavirus genes (Donaldson, et al. 2007;

Stobart, et al. 2012; Forni, et al. 2016). These connections might be linked to host

adaptation (Bhatt, et al. 2013; Longdon, et al. 2014; Bedhomme, et al. 2015). Further *in*

vitro research is needed to verify these findings and investigate the driving biological

mechanisms behind it.

In summary, our study provided novel molecular insights on SARS-CoV-2 glycoprotein

features and placed SARS-CoV-2 lineage in a new context within the evolutionary

trajectory that has led to the current pandemic. Extensive recombination of isolates

circulating in bat, pangolin and/or potentially other unsampled hosts such as civets or

pigs, has been a crucial factor facilitating the emergence of a highly infectious virus,

capable both to elude the immune system (Yuan, et al. 2020) and infect target cells

more efficiently (Andersen, et al. 2020). The landscape of CoVs diversity still needs to

be exhaustively explored, and future emergence of novel recombinant CoVs, and other

viruses as well, should be an urgent concern. Strengthening surveillance of viral

populations and improving our understanding of their dynamics will help the prevention

of new zoonotic outbreaks and spread of emerging pathogens.

Materials and Methods

Genetic data

Publicly available genomic CoVs sequences were obtained from GenBank and GISAID

(Table S5 and S7). To gather closely related CoVs BLASTn was performed using the

ViPR database and keeping the first 100 hits. Genome sequences were aligned using

MAFFT (Katoh and Standley 2016) and refined manually. Alignments for the

glycoprotein were extracted from the genome alignment.

Recombination breakpoint and hotspot analyses

The presence of conflicting phylogenetic signals (i.e. distinct tree topologies compatible

with the same set of aligned sequences) in the full genome data set was investigated,

first, by inferring networks using neighbor-net (Bryant and Moulton 2004) implemented in

SplitsTree4 (Huson and Bryant 2006). Significant presence of recombination signal was

then assessed with the pairwise homoplasy index (Phi) test (Bruen, et al. 2006).

Identification of putative recombinant events and associated breakpoints was performed

using the RDP(Martin and Rybicki 2000), GENECOV (Padidam, et al. 1999), MAXCHI

(Smith 1992), CHIMAERA (Posada and Crandall 2001), SISCAN (Gibbs, et al. 2000)

and 3SEQ (Boni, et al. 2007) algorithms implemented in RDP4 (Martin, et al. 2015)

software (http://web.cbio.uct.ac.za/~darren/rdp.html), v.4.97. Statistical evidence of

recombination was indicated by p-values < 0.05, after Bonferroni correction for multiple

comparisons. A putative recombination events were considered as such if supported by

at least five of the six algorithms used. Full table with recombination results id given in

Table S1. BOOTSCAN (Salminen, et al. 1995) plots were generated using RDP4.

Phylogenetic signal for the subset alignment including bat, human and pangolin CoV-2

sequences was l was confirmed with TREE-PUZZLE(Schmidt, et al. 2002), and

weighted SH and AU tests (Shimodaira 2002) implemented in PAUP* v.4.04a build 167

(Hancock 2014). Neighbor joining, maximum likelihood and Bayesian trees were

calculated respectively with Q-Tree v.2.0 (Nguyen, et al. 2015) and MrBayes (Ronquist,

et al. 2012); best models for the alignments were chosen using Model Finder

(Kalyaanamoorthy, et al. 2017).

Recombination hotspots analysis was also performed using RDP4 through a

permutation test, which randomly distributes the observed breakpoints across the

genome on the variable nucleotide positions between the three sequences involved in

the event. From the permutations, density plots are drawn, and the distribution of

observed breakpoints is compared to the permutations results. Two iterations were

performed, with sliding windows of size 100 and 200, moved one nucleotide at the time,

with different sensitivity to hotspots and coldspots. Segments that had more observed

breakpoints than the local maximum number of breakpoints of 99% of 1,000

permutations were considered hotspots (p=0.01); in the same fashion, segments that

had less breakpoints than the local minimum of 99% of permutations were considered

coldspots (Heath, et al. 2006).

To identify the recombinant structure used to correct selection analyses, we used a

genetic algorithm, GARD (Kosakovsky Pond, et al. 2006), on the subset of sequences

including all bat and pangolin isolates, as well as five randomly chosen SARS-CoV-2

sequences; the reason for this subsampling is that for a fixed number of sites, GARD

loses power when there are many nearly related sequences.

Signature pattern and selection analyses

To identify new features of CoV-2, we used a higher number of isolates permit to explore

a larger evolutionary landscape. VESPA algorithm (Korber and Myers 1992) was used

with a dataset comprising the 728 genomes of SARS-CoV-2, and SARS-CoV-2 isolates

from the closely related bat (EPI_ISL_402131, 2013, Accession MN996532.1) and

pangolin lineage "a" (EPI_ISL_410538-43 isolates isolated in 2017 in China) and lineage

"b" (EPI_ISL_412860 and EPI_ISL_410721 isolates isolated in 2019 in China) (Table

S8). Positions are numbered with SARS-CoV-2 isolate Wuhan-Hu-1 (Accession

MN908947.3) as reference (Table S9).

For selection analyses, we used GARD results to partition the 119 unique sequences

subset. We inferred ML trees separately for each segment using *raxml-ng* and the

GTR+G+I model with 5 random parsimony starting trees. In these trees, we further

identified three branches that included host-switching events and the branch separating

2017 and 2019 pangolin isolates (see Figure 4).

We used BUSTED (Murrell, et al. 2015) to assess the presence of selection on the gene

S partitions. We ran FEL (Kosakovsky Pond and Frost 2005) and MEME (Murrell, et al.

2012) methods on this partitioned alignment to look for pervasive (FEL) or episodic

(MEME) diversifying positive selection affecting the four inter-clade branches, and, in a

separate analysis, affecting only SARS-CoV-2 clade. To look for lineage specific

selection on inter-clade branches we used the aBSREL (Smith, et al. 2015) method

separately on each partition (since it cannot be applied to multi-partition data). Finally, to

look for co-evolution between sites, we applied BGM (Poon, et al. 2007) to each partition

separately, focusing only on internal branches and sites that accumulated at least two

substitutions on internal branches. All selection analyses were carried out in HyPhy
v2.5.14 (Pond, et al. 2005).
505

Structure homology modeling
As a first step, the amino acid sequences of SARS-CoV-2 Wuhan-Hu-1 (Accession
QHD43416) and EPI_ISL_402131 (bat, 2013, Accession MN996532.1) were submitted
to Swiss-model (Waterhouse, et al. 2018) . The best models showing high values of
Global Model Quality Estimation (GMQE) were selected and used as templates. Amino
acid alignments of targets and templates were performed using EMBOSS Needle
(Madeira, et al. 2019). The 3D homology homo-trimer models were created with the
Modeller v9. 23 (Eswar, et al. 2006; Webb and Sali 2014) automodel class using the
alignment as a guide. Then Discreet Optimised Protein Energy Score (DOPE) based
model selection (Shen and Sali 2006) and refinements were conducted using Modeller
v9.23 scripts. The final structure model validation was conducted with ProSA (Sippl
1993; Wiederstein and Sippl 2007) and QMEAN servers (Benkert, et al. 2009; Benkert,
et al. 2011). Visualization of the atomic models, including figures and movies, is made
with Chimera v1.12 (Pettersen, et al. 2004) and VMD v1.9.2 (Humphrey, et al. 1996).
520

Molecular Dynamics simulation
The structure of SARS-CoV-2, bat and pangolin, obtained for homology modeling, were
used as starting structures for Molecular Dynamics (MD) simulations. In detail, topology
file was generated with the pdb2gmx Gromacs tool (Páll 2015), using the amber99sb
forcefield (Lindahl, et al. 2017)..The proteins were embedded in a triclinic box, extending
up to 15 Å from the solute, and immersed in TIP3P water molecules (Jorgensen 1983).
Counter ions were added to neutralize the overall charge with the genion gromacs tool.
After energy minimizations, the systems were slowly relaxed for 5 ns by applying
positional restraints of 1000 kJ mol^{-1} nm^{-2} to the protein atoms. Then unrestrained MD
simulations were carried out for a length of 100 ns for each system, with a time step of 2
fs, using GROMACS 2018.3 simulation package (Páll 2015) (supercomputer Galileo,
CINECA, Bologna, Italy). V-rescale temperature coupling was employed to keep the
temperature constant at 300 K (Bussi 2007). The Particle-Mesh Ewald method was used
for the treatment of the long-range electrostatic interactions (Darden 1993). The first 5 ns
portion of each trajectory was excluded from the analysis.

We carried out the Essential Dynamics analysis to identify the main 3N directions along

which the majority of the protein motion is defined (Amadei 1993). This analysis is based

on the diagonalization of the covariance matrix built from the atomic fluctuations after the

removal of the translational and rotational movement and we applied it on the c-alpha

atoms of each monomer (residue 27-1146 in our model of SARS-CoV-2), after

concatenating their trajectories (255 ns total sampling).

Figures and Tables

Figure 1. Recombination analysis across CoV genomes. (a) NNet graphs obtained

using split-decomposition and based on the full genome sequence to detect

recombination. P-value of PHI test of recombination is reported. Pangolin a taxa:

EPI_ISL_410538-43 isolates isolated in 2017 in China; pangolin b taxa:

EPI_ISL_412860 and EPI_ISL_410721 isolates isolated in 2019 in China. (b)

Recombinant event involving human, bat and pangolin SARS-CoV-2 glycoprotein S. (c)

Detail of the RBD recombinant fragment amino acid alignment. Dots represent identity

with he human CoV-2. (d) Mosaic structure of SARS-CoV-2 glycoprotein S. The 4th

partition (1501-2262), as identified by GARD analysis, still presents some recombination

signal (see table S4 for details).

Figure 2. Recombination hot- and cold- spots prediction across CoV genomes.

Hotspot (upper graph, sliding window size = 100) or coldspot (lower graph, sliding

window size = 200) analysis. Detected recombination breakpoints are marked on the

top of the panel. P-values ≤ 0.01 above the 0 mark indicate presence of hotspots (upper

graph, "peaks"); in the bottom panel, p-values ≤ 0.01 below the 0 mark ("valleys")

indicate the presence of coldspots.

Figure 3. Signature residues in the S glycoprotein of SARS-CoV-2 and ribbon

representations of the S protomer. (a) Signature residues that distinguish human

SARS-COV-2 from closely related bat and pangolin (lineage 'a' isolates isolated in 2017

in China; 'b' isolates isolated in 2019 in China) isolates; residues are colored according

their charge: hydrophilic (RKDENQ) residues in blue, neutral (SGHTAP) in green, and

hydrophobic (YVMCLFIW) in black. Numbers correspond to the residue position in

Wuhan-Hu-1 MN908947.3 isolate. In the box below we show a schematic representation

of the spike glycoprotein is given: S_1 and the S_2 subunits of the S glycoprotein are

indicated in grey and fuchsia, while CTD1 is shown green, CTD2 in red, CTD3 in blue.

Blue boxes show coldspots. Fuchsia box around the residues between 333-516 are

indicating the residues involved into rotation of RBD, red dotted box indicates residues

340-511 that SARS-CoV-2 obtained from the recombination event with pangolin lineage

"b". Residues unique to human SARS-CoV-2 are shaded in cyan, residues that form the

hACE receptor binding are shaded in orange, and the acquired polybasic residues

(PRRA) in the S_1/S_2 cleavage site in yellow. (b) The monomer of SARS-CoV-2 S

glycoprotein and its three domains of the S_1 subunit (CTD1, CTD2, and CTD3) and S_2

subunit (pink) are shown in different colors: CTD1 in green, CTD2 in red, CTD3 in blue.

The N-terminal domain was shown is colored tan. The residues unique to human SARS-

CoV-2, hACE2 biding site residues, and S_1/S_2 cleavage sites were shown with cyan,

orange and yellow spheres, respectively. (c) Homo-trimer structure of the modeled

SARS-CoV-2. Residues, S_1 (NTD, CTD1, CTD2, and CTD3) and the S_2 subunits are

colored corresponding to panel b. (d) The superposition of the modeled S protomer of

SARS-CoV-2 with the S protomer of SARS-CoV (PDB 5WRG). The missing NTD

domain for SARS-CoV was not shown whereas the NTD domain of SARS-CoV-2 is

shown by tan color. Residues and domains are colored corresponding to panel b.

Figure 4. Selection analyses of Spike glycoprotein. Selection analyses were applied

to a partitioned alignment of 119 CoV sequences, with partitions indicated by alternating

white and grey rectangles. A maximum likelihood tree (rooted on the bat isolate for

clarity) for each partition is shown, and the four inter-clade branches representing

potential zoonotic events depicted in dark red. The branch ancestral to SARS-CoV-2

isolates has an arrow pointing to it. Branches longer that 0.25 subs/site (under the MG94

codon model in aBSREL analyses) are censored at 0.25 subs/site and shown in dashed

lines. The impact of selective forces at individual sites is shown in two vertical bars at the

top, where MEME p-values are shown either for the SARS-CoV-2 clade (top bar) or the

inter-clade branches (bottom bar) as trail plots. Tick-marks between the bars correspond

to the location of sites that were inferred to be subject to diversifying positive selection in

the SARS-CoV-2 clade (there were none in the inter-clade branches), and those subject

to pervasive purifying selection along the inter-clade branches.

Figure 5. Essential Dynamics analysis of SARS-CoV-2 in comparison with CoV from bat and pangolin. (a-b) Projection of the SARS-CoV-2 S glycoprotein MD trajectory along Essential Dynamics (ED) eigenvector 1 (see also Supplementary movies 1-2). The two extreme conformations are shown in (a) and (b). Domain colors as in panel Figure 3b. The opening/closing mechanism of the SARS-CoV-2 S glycoprotein monomer by means of the rotation of RBD is evident. (c) Per-residue Root Mean Square Fluctuations (RMSF) in nm of the SARS-CoV-2 S glycoprotein filtered trajectory along ED eigenvectors 1. The region with higher fluctuations is the CTD1, highlighted by red lines, demonstrating that its rotation in the opening/closing mechanism is the most correlated motion in the protein. The peak in fluctuations involving residues 828-852, highlighted by cyan lines, demonstrates that when the RBD is in the up conformation, the fusion peptide visits a more solvent exposed conformation (see Figure S4 and Supplementary movie 3); (d) as in (c) but along eigenvector 2. The peak of fluctuations highlighted by red, purple and cyan lines are indicative of a direct structural communication between the hACE2 binding region, residue 620-640 in CTD3 and the fusion peptide (see Figure S5 and Supplementary movie 4); (e) as in (c) but along

630 eigenvector 3. The peak of fluctuations highlighted by blue, red and orange lines are

631 indicative of long range communications between residue 244-265 in NTD, the hACE2

632 binding residues and the furin-like cleavage site (see Figure S6 and Suppl Movie 5).

634 **Acknowledgements**
635
MS is supported in part by the Stephany W. Holloway University Chair in AIDS Research. This
work was supported by the Tunisian Ministry of Higher Education and Scientific Research, the
'Departments of Excellence-2018' Program (Dipartimenti di Eccellenza) of the Italian Ministry of
Education, University and Research, DIBAF-Department of University of Tuscia, Project
'Landscape 4.0–food, wellbeing and environment'. We thank all those who have contributed
SARS-CoV-2 genome sequences to the GISAID database (https://www.gisaid.org/). We
acknowledge Cineca and ELIXIR-IIB for computing resources We would like to thank Dr. Darren
Martin for his advice on the recombination hotspot analysis.
644

References

Amadei A, Linssen,A.B. and Berendsen,H.J. . 1993. Essential dynamics of proteins. .
Proteins, 17, :412–425.
Andersen KG, Rambaut A, Lipkin WI, Holmes EC, Garry RF. 2020. The proximal
origin of SARS-CoV-2. Nature Medicine.
Anthony SJ, Gilardi K, Menachery VD, Goldstein T, Ssebide B, Mbabazi R, Navarrete-
Macias I, Liang E, Wells H, Hicks A, et al. 2017. Further Evidence for Bats as the
Evolutionary Source of Middle East Respiratory Syndrome Coronavirus. mBio
8:e00373-00317.
Bedhomme S, Hillung J, Elena SF. 2015. Emerging viruses: why they are not jacks of
all trades? Curr Opin Virol 10:1-6.
Belouzard S, Chu VC, Whittaker GR. 2009. Activation of the SARS coronavirus spike
protein via sequential proteolytic cleavage at two distinct sites. Proc Natl Acad Sci U
S A 106:5871-5876.
Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate - a Practical
and Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society
Series B-Statistical Methodology 57:289-300.
Benkert P, Biasini M, Schwede T. 2011. Toward the estimation of the absolute
quality of individual protein structure models. Bioinformatics 27:343-350.
Benkert P, Kunzli M, Schwede T. 2009. QMEAN server for protein model quality
estimation. Nucleic Acids Res 37:W510-514.
Bhatt S, Lam TT, Lycett SJ, Leigh Brown AJ, Bowden TA, Holmes EC, Guan Y, Wood JL,
Brown IH, Kellam P, et al. 2013. The evolutionary dynamics of influenza A virus
adaptation to mammalian hosts. Philos Trans R Soc Lond B Biol Sci 368:20120382.
Boni MF, Lemey P, Jiang X, Lam TT-Y, Perry B, Castoe T, Rambaut A, Robertson DL.
2020. Evolutionary origins of the SARS-CoV-2 sarbecovirus lineage responsible for
the COVID-19 pandemic. bioRxiv:2020.2003.2030.015008.
Boni MF, Posada D, Feldman MW. 2007. An exact nonparametric method for
inferring mosaic structure in sequence triplets. Genetics 176:1035-1047.

Bosch BJ, Bartelink W, Rottier PJ. 2008. Cathepsin L functionally cleaves the severe
acute respiratory syndrome coronavirus class I fusion protein upstream of rather
than adjacent to the fusion peptide. J Virol 82:8887-8890.
Bosch BJ, van der Zee R, de Haan CA, Rottier PJ. 2003. The coronavirus spike protein
is a class I virus fusion protein: structural and functional characterization of the
fusion core complex. J Virol 77:8801-8811.
Bruen TC, Philippe H, Bryant D. 2006. A simple and robust statistical test for
detecting the presence of recombination. Genetics 172:2665-2681.
Bryant D, Moulton V. 2004. Neighbor-net: an agglomerative method for the
construction of phylogenetic networks. Mol Biol Evol 21:255-265.
Bucknall RA, King LM, Kapikian AZ, Chanock RM. 1972. Studies with human
coronaviruses. II. Some properties of strains 229E and OC43. Proc Soc Exp Biol Med
139:722-727.
Bussi GD, D.; Parrinello, M. 2007. Canonical sampling through velocity rescaling. . J
Chem Phys 126, 014101, doi:10.1063/1.2408420.
Darden TY, D.; Pedersen, L. . 1993. Particle mesh Ewald: An N log(N) method for
Ewald sums in large systems. J Chem Phys 98, 10089, doi: 10.1063/1.464397.
Decaro N, Mari V, Campolo M, Lorusso A, Camero M, Elia G, Martella V, Cordioli P,
Enjuanes L, Buonavoglia C. 2009. Recombinant canine coronaviruses related to
transmissible gastroenteritis virus of Swine are circulating in dogs. J Virol 83:1532-
1537.
Donaldson EF, Graham RL, Sims AC, Denison MR, Baric RS. 2007. Analysis of murine
hepatitis virus strain A59 temperature-sensitive mutant TS-LA6 suggests that nsp10
plays a critical role in polyprotein processing. J Virol 81:7086-7098.
Dudas G, Rambaut A. 2016. MERS-CoV recombination: implications about the
reservoir and potential for adaptation. Virus Evol 2:vev023.
El Najjar F, Lampe L, Baker ML, Wang LF, Dutch RE. 2015. Analysis of cathepsin and
furin proteolytic enzymes involved in viral fusion protein activation in cells of the
bat reservoir host. PLoS One 10:e0115736.
Eswar N, Webb B, Marti-Renom MA, Madhusudhan MS, Eramian D, Shen MY, Pieper
U, Sali A. 2006. Comparative protein structure modeling using Modeller. Curr Protoc
Bioinformatics Chapter 5:Unit-5 6.
Fehr AR, Perlman S. 2015. Coronaviruses: an overview of their replication and
pathogenesis. Methods Mol Biol 1282:1-23.
Ferron F, Subissi L, Silveira De Morais AT, Le NTT, Sevajol M, Gluais L, Decroly E,
Vonrhein C, Bricogne G, Canard B, et al. 2018. Structural and molecular basis of
mismatch correction and ribavirin excision from coronavirus RNA. Proc Natl Acad
Sci U S A 115:E162-E171.
Forni D, Cagliani R, Mozzi A, Pozzoli U, Al-Daghri N, Clerici M, Sironi M. 2016.
Extensive Positive Selection Drives the Evolution of Nonstructural Proteins in
Lineage C Betacoronaviruses. J Virol 90:3627-3639.
Gallagher TM, Buchmeier MJ. 2001. Coronavirus spike proteins in viral entry and
pathogenesis. Virology 279:371-374.
Gibbs MJ, Armstrong JS, Gibbs AJ. 2000. Sister-scanning: a Monte Carlo procedure
for assessing signals in recombinant sequences. Bioinformatics 16:573-582.

Graham RL, Baric RS. 2010. Recombination, reservoirs, and the modular spike:
mechanisms of coronavirus cross-species transmission. J Virol 84:3134-3146.
Gui M, Song W, Zhou H, Xu J, Chen S, Xiang Y, Wang X. 2017. Cryo-electron
microscopy structures of the SARS-CoV spike glycoprotein reveal a prerequisite
conformational state for receptor binding. Cell Res 27:119-129.
Hancock JM, Zvelebil, M.J. and Cummings, M.P. . 2014. PAUP* (Phylogenetic Analysis
Using Parsimony (and Other Methods)). In. Dictionary of Bioinformatics and
Computational Biology: In Dictionary of Bioinformatics and Computational Biology
(eds J.M. Hancock and M.J. Zvelebil).
Hatsuzawa K, Hosaka M, Nakagawa T, Nagase M, Shoda A, Murakami K, Nakayama K.
1990. Structure and expression of mouse furin, a yeast Kex2-related protease. Lack
of processing of coexpressed prorenin in GH4C1 cells. J Biol Chem 265:22075-
22078.
Heath L, van der Walt E, Varsani A, Martin DP. 2006. Recombination patterns in
aphthoviruses mirror those found in other picornaviruses. J Virol 80:11827-11832.
Hoffmann M, Kleine-Weber H, Pohlmann S. 2020. A Multibasic Cleavage Site in the
Spike Protein of SARS-CoV-2 Is Essential for Infection of Human Lung Cells. Mol Cell
78:779-784 e775.
Humphrey W, Dalke A, Schulten K. 1996. VMD: visual molecular dynamics. J Mol
Graph 14:33-38, 27-38.
Huson DH, Bryant D. 2006. Application of phylogenetic networks in evolutionary
studies. Mol Biol Evol 23:254-267.
Izaguirre G. 2019. The Proteolytic Regulation of Virus Cell Entry by Furin and Other
Proprotein Convertases. Viruses 11.
Ji W, Wang W, Zhao X, Zai J, Li X. 2020. Homologous recombination within the spike
glycoprotein of the newly identified coronavirus may boost cross-species
transmission from snake to human. J Med Virol.
Jia L, Hu F, Li H, Li L, Tang X, Liu Y, Deng H, Han J, Li J, Cai W. 2018. Characterization
of small genomic regions of the hepatitis B virus should be performed with more
caution. Virol J 15:188.
Jorgensen WLC, J.; Madura,J.D.; Impey,R.W.; Klein,M.L. . 1983. Comparison of simple
potential functions for simulating liquid water. J. Chem. Phys. 79:79, 926–935.
Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017.
ModelFinder: fast model selection for accurate phylogenetic estimates. Nat Methods
14:587-589.
Kamel Boulos MN, Geraghty EM. 2020. Geographical tracking and mapping of
coronavirus disease COVID-19/severe acute respiratory syndrome coronavirus 2
(SARS-CoV-2) epidemic and associated events around the world: how 21st century
GIS technologies are supporting the global fight against outbreaks and epidemics.
Int J Health Geogr 19:8.
Katoh K, Standley DM. 2016. A simple method to control over-alignment in the
MAFFT multiple sequence alignment program. Bioinformatics 32:1933-1942.
Kim D, Lee JY, Yang JS, Kim JW, Kim VN, Chang H. 2020. The Architecture of SARS-
CoV-2 Transcriptome. Cell 181:914-921 e910.
Klenk HD, Garten W. 1994. Host cell proteases controlling virus pathogenicity.
Trends Microbiol 2:39-43.

Korber B, Myers G. 1992. Signature pattern analysis: a method for assessing viral
sequence relatedness. AIDS Res Hum Retroviruses 8:1549-1560.
Kosakovsky Pond SL, Frost SD. 2005. Not so different after all: a comparison of
methods for detecting amino acid sites under selection. Mol Biol Evol 22:1208-1222.
Kosakovsky Pond SL, Posada D, Gravenor MB, Woelk CH, Frost SD. 2006. GARD: a
genetic algorithm for recombination detection. Bioinformatics 22:3096-3098.
Lai AL, Millet JK, Daniel S, Freed JH, Whittaker GR. 2017. The SARS-CoV Fusion
Peptide Forms an Extended Bipartite Fusion Platform that Perturbs Membrane
Order in a Calcium-Dependent Manner. J Mol Biol 429:3875-3892.
Lai MM, Baric RS, Makino S, Keck JG, Egbert J, Leibowitz JL, Stohlman SA. 1985.
Recombination between nonsegmented RNA genomes of murine coronaviruses. J
Virol 56:449-456.
Lan J, Ge J, Yu J, Shan S, Zhou H, Fan S, Zhang Q, Shi X, Wang Q, Zhang L, et al. 2020.
Structure of the SARS-CoV-2 spike receptor-binding domain bound to the ACE2
receptor. Nature 581:215-220.
Lau SK, Feng Y, Chen H, Luk HK, Yang WH, Li KS, Zhang YZ, Huang Y, Song ZZ, Chow
WN, et al. 2015. Severe Acute Respiratory Syndrome (SARS) Coronavirus ORF8
Protein Is Acquired from SARS-Related Coronavirus from Greater Horseshoe Bats
through Recombination. J Virol 89:10532-10547.
Leung GM, Hedley AJ, Ho LM, Chau P, Wong IO, Thach TQ, Ghani AC, Donnelly CA,
Fraser C, Riley S, et al. 2004. The epidemiology of severe acute respiratory
syndrome in the 2003 Hong Kong epidemic: an analysis of all 1755 patients. Ann
Intern Med 141:662-673.
Li F. 2016. Structure, Function, and Evolution of Coronavirus Spike Proteins. Annu
Rev Virol 3:237-261.
Li F, Li W, Farzan M, Harrison SC. 2005. Structure of SARS coronavirus spike
receptor-binding domain complexed with receptor. Science 309:1864-1868.
Li X, Giorgi EE, Marichann MH, Foley B, Xiao C, Kong X-p, Chen Y, Korber B, Gao F.
2020. Emergence of SARS-CoV-2 through Recombination and Strong Purifying
Selection. bioRxiv:2020.2003.2020.000885.
Li X, Giorgi EE, Marichannegowda MH, Foley B, Xiao C, Kong X-P, Chen Y,
Gnanakaran S, Korber B, Gao F. 2020. Emergence of SARS-CoV-2 through
recombination and strong purifying selection. Science Advances.
Li X, Wang W, Zhao X, Zai J, Zhao Q, Li Y, Chaillon A. 2020. Transmission dynamics
and evolutionary history of 2019-nCoV. J Med Virol 92:501-511.
Lindahl V, Villa A, Hess B. 2017. Sequence dependency of canonical base pair
opening in the DNA double helix. PLoS Comput Biol 13:e1005463.
Liu P, Jiang JZ, Wan XF, Hua Y, Li L, Zhou J, Wang X, Hou F, Chen J, Zou J, et al. 2020.
Are pangolins the intermediate host of the 2019 novel coronavirus (SARS-CoV-2)?
PLoS Pathog 16:e1008421.
Longdon B, Brockhurst MA, Russell CA, Welch JJ, Jiggins FM. 2014. The evolution and
genetics of virus host shifts. PLoS Pathog 10:e1004395.
Madeira F, Park YM, Lee J, Buso N, Gur T, Madhusoodanan N, Basutkar P, Tivey ARN,
Potter SC, Finn RD, et al. 2019. The EMBL-EBI search and sequence analysis tools
APIs in 2019. Nucleic Acids Res 47:W636-W641.

Madu IG, Roth SL, Belouzard S, Whittaker GR. 2009. Characterization of a highly
conserved domain within the severe acute respiratory syndrome coronavirus spike
protein S2 domain with characteristics of a viral fusion peptide. J Virol 83:7411-
7421.
Martin D, Rybicki E. 2000. RDP: detection of recombination amongst aligned
sequences. Bioinformatics 16:562-563.
Martin DP, Murrell B, Golden M, Khoosal A, Muhire B. 2015. RDP4: Detection and
analysis of recombination patterns in virus genomes. Virus Evol 1:vev003.
Mavian C, Rife BD, Dollar JJ, Cella E, Ciccozzi M, Prosperi MCF, Lednicky J, Morris JG,
Capua I, Salemi M. 2017. Emergence of recombinant Mayaro virus strains from the
Amazon basin. Sci Rep 7:8718.
Millet JK, Whittaker GR. 2014. Host cell entry of Middle East respiratory syndrome
coronavirus after two-step, furin-mediated activation of the spike protein. Proc Natl
Acad Sci U S A 111:15214-15219.
Murrell B, Weaver S, Smith MD, Wertheim JO, Murrell S, Aylward A, Eren K, Pollner
T, Martin DP, Smith DM, et al. 2015. Gene-wide identification of episodic selection.
Mol Biol Evol 32:1365-1371.
Murrell B, Wertheim JO, Moola S, Weighill T, Scheffler K, Kosakovsky Pond SL. 2012.
Detecting individual sites subject to episodic diversifying selection. PLoS Genet
8:e1002764.
Muth D, Corman VM, Roth H, Binger T, Dijkman R, Gottula LT, Gloza-Rausch F,
Balboni A, Battilani M, Rihtaric D, et al. 2018. Attenuation of replication by a 29
nucleotide deletion in SARS-coronavirus acquired during the early stages of human-
to-human transmission. Sci Rep 8:15177.
Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and
effective stochastic algorithm for estimating maximum-likelihood phylogenies. Mol
Biol Evol 32:268-274.
Ou X, Liu Y, Lei X, Li P, Mi D, Ren L, Guo L, Guo R, Chen T, Hu J, et al. 2020.
Characterization of spike glycoprotein of SARS-CoV-2 on virus entry and its immune
cross-reactivity with SARS-CoV. Nat Commun 11:1620.
Padidam M, Sawyer S, Fauquet CM. 1999. Possible emergence of new geminiviruses
by frequent recombination. Virology 265:218-225.
Páll SA, M.J.; Kutzner, C.; Hess, B.; Lindahl, E. . 2015. Tackling exascale software
challenges in molecular dynamics simulati ons with GROMACS. In Solving Software
Challenges for Exascale (eds. Markadis, S. & Laure, E.). Springer International
Publishing: Switzerland, London Volume 8759, :pp. 3–27.
Paraskevis D, Kostaki EG, Magiorkinis G, Panayiotakopoulos G, Sourvinos G,
Tsiodras S. 2020. Full-genome evolutionary analysis of the novel corona virus
(2019-nCoV) rejects the hypothesis of emergence as a result of a recent
recombination event. Infect Genet Evol 79:104212.
Peng G, Yang Y, Pasquarella JR, Xu L, Qian Z, Holmes KV, Li F. 2017. Structural and
Molecular Evidence Suggesting Coronavirus-driven Evolution of Mouse Receptor. J
Biol Chem 292:2174-2181.
Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE.
2004. UCSF Chimera--a visualization system for exploratory research and analysis. J
Comput Chem 25:1605-1612.

Pond SL, Frost SD, Muse SV. 2005. HyPhy: hypothesis testing using phylogenies.
Bioinformatics 21:676-679.
Poon AF, Lewis FI, Pond SL, Frost SD. 2007. An evolutionary-network model reveals
stratified interactions in the V3 loop of the HIV-1 envelope. PLoS Comput Biol
3:e231.
Posada D, Crandall KA. 2002. The effect of recombination on the accuracy of
phylogeny estimation. J Mol Evol 54:396-402.
Posada D, Crandall KA. 2001. Evaluation of methods for detecting recombination
from DNA sequences: computer simulations. Proc Natl Acad Sci U S A 98:13757-
13762.
Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Hohna S, Larget B, Liu
L, Suchard MA, Huelsenbeck JP. 2012. MrBayes 3.2: efficient Bayesian phylogenetic
inference and model choice across a large model space. Syst Biol 61:539-542.
Salminen MO, Carr JK, Burke DS, McCutchan FE. 1995. Identification of breakpoints
in intergenotypic recombinants of HIV type 1 by bootscanning. AIDS Res Hum
Retroviruses 11:1423-1425.
Schalken JA, Roebroek AJ, Oomen PP, Wagenaar SS, Debruyne FM, Bloemers HP, Van
de Ven WJ. 1987. fur gene expression as a discriminating marker for small cell and
nonsmall cell lung carcinomas. J Clin Invest 80:1545-1549.
Schierup MH, Hein J. 2000. Consequences of recombination on traditional
phylogenetic analysis. Genetics 156:879-891.
Schmidt HA, Strimmer K, Vingron M, von Haeseler A. 2002. TREE-PUZZLE:
maximum likelihood phylogenetic analysis using quartets and parallel computing.
Bioinformatics 18:502-504.
Shen MY, Sali A. 2006. Statistical potential for assessment and prediction of protein
structures. Protein Sci 15:2507-2524.
Shimodaira H. 2002. An approximately unbiased test of phylogenetic tree selection.
Syst Biol 51:492-508.
Shriner D, Nickle DC, Jensen MA, Mullins JI. 2003. Potential impact of recombination
on sitewise approaches for detecting positive natural selection. Genet Res 81:115-
121.
Simmons G, Zmora P, Gierer S, Heurich A, Pohlmann S. 2013. Proteolytic activation
of the SARS-coronavirus spike protein: cutting enzymes at the cutting edge of
antiviral research. Antiviral Res 100:605-614.
Simon-Loriere E, Holmes EC. 2011. Why do RNA viruses recombine? Nature
Reviews Microbiology 9:617-626.
Sippl MJ. 1993. Recognition of errors in three-dimensional structures of proteins.
Proteins 17:355-362.
Smith JM. 1992. Analyzing the mosaic structure of genes. J Mol Evol 34:126-129.
Smith MD, Wertheim JO, Weaver S, Murrell B, Scheffler K, Kosakovsky Pond SL.
2015. Less is more: an adaptive branch-site random effects model for efficient
detection of episodic diversifying selection. Mol Biol Evol 32:1342-1353.
Song W, Gui M, Wang X, Xiang Y. 2018. Cryo-EM structure of the SARS coronavirus
spike glycoprotein in complex with its host cell receptor ACE2. PLoS Pathog
14:e1007236.

Stobart CC, Lee AS, Lu X, Denison MR. 2012. Temperature-sensitive mutants and
revertants in the coronavirus nonstructural protein 5 protease (3CLpro) define
residues involved in long-distance communication and regulation of protease
activity. J Virol 86:4801-4810.
Su S, Wong G, Shi W, Liu J, Lai ACK, Zhou J, Liu W, Bi Y, Gao GF. 2016. Epidemiology,
Genetic Recombination, and Pathogenesis of Coronaviruses. Trends Microbiol
24:490-502.
Terada Y, Matsui N, Noguchi K, Kuwata R, Shimoda H, Soma T, Mochizuki M, Maeda
K. 2014. Emergence of pathogenic coronaviruses in cats by homologous
recombination between feline and canine coronaviruses. PLoS One 9:e106534.
Tian PF, Jin YL, Xing G, Qv LL, Huang YW, Zhou JY. 2014. Evidence of recombinant
strains of porcine epidemic diarrhea virus, United States, 2013. Emerg Infect Dis
20:1735-1738.
Wang LF, Eaton BT. 2007. Bats, civets and the emergence of SARS. Curr Top
Microbiol Immunol 315:325-344.
Wang Q, Zhang Y, Wu L, Niu S, Song C, Zhang Z, Lu G, Qiao C, Hu Y, Yuen KY, et al.
2020. Structural and Functional Basis of SARS-CoV-2 Entry by Using Human ACE2.
Cell 181:894-904 e899.
Waterhouse A, Bertoni M, Bienert S, Studer G, Tauriello G, Gumienny R, Heer FT, de
Beer TAP, Rempfer C, Bordoli L, et al. 2018. SWISS-MODEL: homology modelling of
protein structures and complexes. Nucleic Acids Res 46:W296-W303.
Webb B, Sali A. 2014. Protein structure modeling with MODELLER. Methods Mol
Biol 1137:1-15.
Weinreich DM, Chao L. 2005. Rapid evolutionary escape by large populations from
local fitness peaks is likely in nature. Evolution 59:1175-1182.
Wiederstein M, Sippl MJ. 2007. ProSA-web: interactive web service for the
recognition of errors in three-dimensional structures of proteins. Nucleic Acids Res
35:W407-410.
Wong MC, Javornik Cregeen SJ, Ajami NJ, Petrosino JF. 2020. Evidence of
recombination in coronaviruses implicating pangolin origins of nCoV-2019.
bioRxiv:2020.2002.2007.939207.
Woo PC, Lau SK, Tsoi HW, Huang Y, Poon RW, Chu CM, Lee RA, Luk WK, Wong GK,
Wong BH, et al. 2005. Clinical and molecular epidemiological features of coronavirus
HKU1-associated community-acquired pneumonia. J Infect Dis 192:1898-1907.
Wrapp D, Wang N, Corbett KS, Goldsmith JA, Hsieh CL, Abiona O, Graham BS,
McLellan JS. 2020. Cryo-EM structure of the 2019-nCoV spike in the prefusion
conformation. Science 367:1260-1263.
Wu F, Zhao S, Yu B, Chen YM, Wang W, Song ZG, Hu Y, Tao ZW, Tian JH, Pei YY, et al.
2020. A new coronavirus associated with human respiratory disease in China.
Nature 579:265-269.
Xiao K, Zhai J, Feng Y, Zhou N, Zhang X, Zou JJ, Li N, Guo Y, Li X, Shen X, et al. 2020.
Isolation of SARS-CoV-2-related coronavirus from Malayan pangolins. Nature.
Yan R, Zhang Y, Li Y, Xia L, Guo Y, Zhou Q. 2020. Structural basis for the recognition
of SARS-CoV-2 by full-length human ACE2. Science 367:1444-1448.

Yuan M, Wu NC, Zhu X, Lee CD, So RTY, Lv H, Mok CKP, Wilson IA. 2020. A highly
conserved cryptic epitope in the receptor-binding domains of SARS-CoV-2 and
SARS-CoV. Science.
Zaki AM, van Boheemen S, Bestebroer TM, Osterhaus AD, Fouchier RA. 2012.
Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. N Engl
J Med 367:1814-1820.
Zhou H, Chen X, Hu T, Li J, Song H, Liu Y, Wang P, Liu D, Yang J, Holmes EC, et al.
2020. A novel bat coronavirus reveals natural insertions at the S1/S2 cleavage site
of the Spike protein and a possible recombinant origin of HCoV-19.
bioRxiv:2020.2003.2002.974139.
Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, Si HR, Zhu Y, Li B, Huang CL, et
al. 2020. A pneumonia outbreak associated with a new coronavirus of probable bat
origin. Nature 579:270-273.
958

Codon in the Spike glycoprotein alignment

SARS-CoV-2 clade
Inter-clade branches
MEME p-value

Positively selected in SARS-CoV-2
Negatively selected in SARS-CoV-2

SARS-COV-2
Yunnan Bat 13
Pangolin 2017
Pangolin 2019

a

b

human

c eigenvector 1

d eigenvector 2

e eigenvector 3

RSMF (nm)

Residue number

f

g

bat

h

i

pangolin