Mapping the coupled human and natural disturbance regimes of Europe's forests

Cornelius Senf[1,2] * and Rupert Seidl[1,2,3]

[1] Institute for Silviculture, University of Natural Resources and Life Sciences (BOKU), Vienna, Austria

[2] Ecosystem dynamics and forest management group, Technical University of Munich, Hans-Carl-von-Carlowitz-Platz 2, 85354 Freising, Germany

[3] Berchtesgaden National Park, Doktorberg 6, 83471, Berchtesgaden, Germany

*Corresponding author: Cornelius.senf@tum.de

Abstract: Forest disturbance shape ecosystem composition and structure, and changes in

forest disturbances can have strong consequences for carbon storage and biodiversity. Yet we

currently lack consistent quantitative data on Europe's forest disturbance regimes and their

changes over time. Here we used satellite data to map three decades (1986-2016) of forest

disturbances across continental Europe, covering 35 countries and a forest area of 210 Mill.

ha at a spatial grain of 30 m, and analyzed the patterns and trends in disturbance size,

frequency and severity. Between 1986 and 2016, 17% of Europe's forest area was disturbed

by anthropogenic or natural causes, totaling to 25 Mill. individual disturbance patches with a

mean patch size 1.09 ha (range between 1^{st} and 99^{th} percentile 0.18 – 10.10 ha). On average

0.52 (0.02 – 3.01) disturbances occurred per km² every year, removing on average 77% (22 –

100%) of the canopy. While spatial patterns of disturbance were highly variable, disturbance

frequency consistently increased, and disturbance severity decreased since 1986. Both social

and ecological factors are needed to explain the observed patterns and trends in forest

disturbance. We thus conclude that in order to understand and manage the changes in

Europe's forest disturbance regimes a coupled human and natural systems perspective is

needed.

Keywords: Coupled Human and Environmental System; Disturbance regime; Remote

sensing; Forest management; Resilience

Forests cover 33 % of Europe's total land area and provide important services to society, ranging from carbon sequestration to the filtration of water, protection of soil from erosion, and human infrastructure from natural hazards [1]. Europe's forests have expanded in recent decades [2] and accumulated substantial amounts of biomass due to intensive post-war reforestation programs, changes in management systems, and timber harvesting rates that remained below increment [3]. This success story of 20th century forestry in Europe, however, also has side effects, as the resultant changes in forest structure have – in combination with climate change – led to an episode of increasing forest disturbances in recent decades [4–7]. Increasing forest disturbances have the potential to erode Europe's carbon storage potential [8,9] and also impact other important services provided by Europe's forests [10,11]. Given a predicted increase in the demand for wood [1] and an expected future intensification of forest dieback under climate change [12], it is fundamental to increase the resilience of Europe's forests to changing disturbances [13–15].

Developing resilient management strategies requires a robust quantitative understanding of forest ecosystem dynamics [16]. In particular, it is essential to understand the disturbance regimes of Europe's forests [17]. Disturbance regimes characterize the cumulative effects of all disturbance events occurring in a given area and time period, and understanding them is fundamental to understanding the current state and future trajectories of forest ecosystems [18]. In Europe, however, forests have been utilized by humans for centuries, transforming species composition and structure [19–21], and consequently also the natural disturbance regimes of forests. In addition to this indirect effect, human land-use is directly disturbing forest canopies through timber harvesting, altering the rate and spatial patterns of forest disturbances compared to natural systems [22]. Human land-use also interacts with natural disturbances, e.g. by salvaging disturbed timber [23] and shortening early seral stages through planting [24]. More broadly, forest management alters biological legacies and landscape structure [23,25], with feedbacks on subsequent disturbances. Due to the intricate linkages

between natural and human processes driving forest disturbances in Europe, analyzing them

in the context of coupled human and natural systems [26,27] is a promising approach. Given that

relevant drivers of future changes in forest disturbance regimes are social-ecological (e.g.,

anthropogenic climate change, novel disturbances caused by the introduction of non-native

disturbance agents, land-use change) applying such a perspective could considerably increase

the inferential potential on current and future changes of forest dynamics cross Europe's

forests.

68   Despite the widespread and long-lasting impacts of changing forest disturbances on

forest ecosystems [10] little quantitatively information on disturbance regimes and their changes

through time exist for Europe. We, for instance, do not know how disturbance size, frequency

and severity (i.e., the main descriptors of disturbance regimes; [28]) are distributed throughout

Europe. Furthermore, while recent studies indicate an increase in disturbance rates across

Europe's natural and managed forests [4,6], it remains unknown whether this change is mainly

driven by changes in disturbance frequency (i.e., more disturbance events) or disturbance size

(i.e., larger individual disturbance patches). Likewise, our quantitative knowledge on changes

in disturbance severity is scant, and it remains unclear whether disturbances in Europe have

become more severe in recent decades (e.g., through increased burn severities; [29]) or whether

recent changes in forest policy (e.g., the adoption of "close-to-nature" silviculture; [30]) have

reduced disturbance severity [4].

80   Here, our aim was to map the patterns and trends of recent (1986-2016) forest

disturbance regimes in Europe. Our specific research questions were: (I) What is the size,

frequency and severity of forest disturbances across Europe's forests? And (II) How did size,

frequency and severity of forest disturbances change over the past three decades? We address

these first two questions by mapping forest disturbance occurrence and severity continuously

for continental Europe (35 countries covering 210 Mill. ha of forest) at a spatial grain of

30 m. We subsequently analyze the disturbance regimes of Europe's forests in a coupled

human and natural systems framework, asking (III) how strongly patterns and trends in

disturbance size, frequency and severity are explained by variation in the natural template vs.

variation in forest policy. We address this third question by comparing the patterns and trends

in disturbance regime indicators among ecoregions (i.e., an aggregate proxy of the spatial

differences in the drivers of natural forest development) and countries (as the spatial entities

encapsulating the variation in forest policies in Europe [31]), hypothesizing that under a coupled

human and natural system perspective both social and ecological factors are needed to explain

patterns and trends in disturbance regimes across Europe's forests.

Results

We identified a total of 25 Mill. individual disturbance patches occurring across Europe

between 1986-2016, equaling a disturbed forest area of 4 Mill. ha or 17 % of Europe's forests

(Figure 1). The overall accuracy of our map was 92.5 %, with a disturbance commission error

of 14.6 % and a disturbance omission error of 32.8 %, but we refer the reader to

Supplementary Note 1 for further information on map accuracies. The average patch size of

forest disturbances in Europe was 1.09 ha, but the disturbance size distribution was highly

left-skewed (Figure 2 B). The median disturbance size was only 0.45 ha, with 78 % of the

disturbances being smaller than 1 ha and 99 % of the disturbances being smaller than 10 ha

(Supplement Table 2). The largest disturbance patch mapped across Europe was a 16,617 ha

large forest fire occurring in 2012 in southern Spain. The average disturbance frequency was

0.52 patches per km^2 per year (median of 0.37 patches per km^2), with highest frequencies

(highest 1 %) ranging from 3 to 31 patches per km^2 (Supplement Table 2). Disturbance

severity, that is a measure between 0 and 100 indicating the loss of canopy during disturbance

(see Figure 1C), ranged from 23 % to 100 %, with an average of 77 % of canopy loss within a

disturbed patch (median of 83 %; Supplement Table 2).

Fig. 1: Forest disturbances in Europe 1986-2016. Disturbance maps were derived from manually interpreting more than 30,000 Landsat images systematically distributed across Europe. Panel A shows the occurrence of disturbances across Europe. Panels B show the

disturbance year and panels C show the disturbance severity – a measure of canopy cover loss

– for three selected areas: (1) Bark beetle outbreak of varying severity in and around Harz

National Park (Germany); (2) wind disturbance in an intensively managed plantation forest in

the Landes of Gascony (France) with very high disturbance severities; and (3) fire

disturbances on the Peloponnese peninsula (Greece), with variable burn-severities.

Spatial variability in the size, frequency and severity of forest disturbances is high

across Europe (Figure 2 A). Disturbance patches are generally larger in Northern and

Southern Europe compared to Central Europe. Also, Eastern Europe has larger disturbance

patches compared to Western Europe (Figure 2 A). Above-average disturbance frequencies

were found in Central Europe, the hemi-boreal zone, parts of France and the Iberian Peninsula

(Figure 2 A). The highest disturbance frequencies (i.e., above 3 patches per km^2) occurred

almost exclusively in Portugal. Disturbance severity was more evenly distributed than the

other two disturbance regime indicators (Figure 2 A), with a tendency towards higher

severities in the Atlantic forests of Ireland and the United Kingdom, the Iberian Peninsula, the

Po-Valley, and the Pannonian Basin. In contrast, low disturbance severities were recorded for

South-Eastern Europe along the Dinaric mountain range, as well as in the Apennine

mountains of Italy.

Fig. 2: (A) Maps of average disturbance size, frequency and severity calculated for hexagons on a 50 km grid across continental Europe. (B) Distribution of disturbance size, frequency and severity across Europe.

Disturbance regimes changed profoundly between 1986 and 2016, with 65 % of Europe's forests experiencing an increase in disturbance size, and 74 % an increase in disturbance frequency (Figure 3 B). Hot spots of increasing disturbance size were in the Baltic states, the United Kingdom, Ireland, and Italy (Figure 3 A), whereas trends were largely negative in Eastern Germany, western Poland and southeastern Europe (Figure 3 A). Hot spots of increasing disturbance frequency were located in Central and Eastern Europe (Figure 3), whereas negative trends in disturbance frequency were recorded for Belarus, western Europe, and northern Scandinavia (Figure 3 A). Disturbance severity decreased in

85 % of the European forest area (Figure 3), with particularly strong trends in Central and

Southeastern Europe.

While the mean disturbance size generally increased across Europe (Figure 3) the

median disturbance size was more stable (no change in median disturbance size in 81 % of

European forests; Supplementary Table 3). Hence, changes in disturbance size were driven by

a widening of the disturbance size distribution, with approximately 50 % of Europe's forests

showing an increase in large disturbance patches (i.e., in the 75% quantile and maximum of

the disturbance patch size distribution; Supplement Table 3). Overall, changes in disturbance

frequency explained 71 % of the variability in changing disturbance rates (i.e., the trend in the

annual percent of forest area disturbed), whereas changes in disturbance size only accounted

for 24 % (see Supplementary Figure 8). Thus, changes in disturbance rates are primarily

driven by changes in disturbance frequency, and not disturbance size, in Europe's forests.

Fig. 3: (A) Maps of trends in disturbance size, frequency and severity calculated at a 50 km

hexagon grid across continental Europe. (B) Distribution of forest area among trend classes.

Spatial variability in all three disturbance regime indicators varied significantly with

ecoregion (i.e., a coarse filter proxy of spatial differences in the drivers of natural forest

development), and ecoregions alone explained 15, 28 and 39 % of the continental-scale

spatial variation in disturbance size, frequency and severity, respectively (Table 1). Including

within-ecoregion variability by country (i.e., a coarse filter proxy for differences in forest

policy and management) further increased the variance explained by 16, 24 and 16 percentage

points, respectively (Table 1).

Table 1: Variance explained by ecoregions and countries nested within ecoregions derived

from linear mixed effect models testing for differences in averages and trends in disturbance

size, frequency and severity measured across 3,240 50 km^2 hexagons. Also shown is the

amount of variance explained by random variation among years (for averages), as well as the

residual variance (i.e., variance not explained by the two coarse filter variables used here).

Indicator	Variance explained			Residual variance
	Ecoregions	Ecoregions/countries	Year	
Average				
Mean size (log)	0.15	0.16	0.01	0.67
Frequency (log)	0.28	0.24	0.03	0.45
Severity	0.39	0.16	0.07	0.37
Trend				
Mean size	0.12	0.11	-	0.77
Frequency	0.09	0.24	-	0.67
Severity	0.19	0.14	-	0.68

Trends in size and severity were also determined by a combination of ecoregions and

policy differences within ecoregions, with the two factors explaining roughly equal parts of

the spatial variation in disturbances trends (Table 1). For changes in frequency ecoregion

alone was a poor predictor, whereas country explained 24 % of the variance in frequency

trends (Table 2). Hence, while the ecological template determines the general spatial pattern

of the disturbance regimes in Europe, human activity modulates this pattern considerably (for

estimates see Supplementary Figure S9, S10), underlining the coupled human and natural

nature of forest disturbances in Europe.

Discussion

We here provide the first quantitative characterization of Europe's forest disturbance regimes,

highlighting the wide variability in disturbance sizes, frequencies and severities prevailing

across the European continent. We show that this variability is determined by the combined

effects of natural processes (i.e., variation in climate and soil, resulting in different forest

development trajectories) and human activity (i.e., variation in forest policy, resulting in

different management regimes). Differences in forest policy can be large even among

neighboring countries in Europe, especially among countries of variable socio-political

histories (e.g., the iron curtain dividing the continent into two geopolitical spheres). These

differences in forest policy result in different management intensities and silvicultural

practices [31] and variable land use legacies [21]. For example, spatial patterns of timber

extraction can vary widely between countries, resulting in contrasting disturbance sizes and

frequencies within ecoregions (Supplementary Figure S11). Another example for the impact

of forest policy on disturbance regimes is the varying share of plantation forests with non-

native tree species [1], such as *Eucalyptus* sp. in Portugal or Black locust (*Robinia*

pseudoacacia) in Hungary (Supplementary Figure S12). Human activity thus has a profound

impact on the forest disturbance regimes of Europe, on the one hand altering disturbance

regimes directly via timber harvesting, and on the other hand indirectly modifying disturbance

processes through changing species composition and forest structure. Our results hence

support the hypothesis that the European forest disturbance regime is a coupled human and

natural system, driven by the complex interplay between social and ecological forces.

The disturbance regimes of Europe's forests are changing profoundly. We here show

that the previously reported increase in disturbance activity [4,6,7] is primarily an effect of

increasing disturbance frequency, while disturbance patch size distributions are becoming

more variable and disturbance severities are decreasing. Variation in forest policy was a more

important predictor of changing disturbance frequencies than the variation in the natural

template, suggesting human resource use as a major driver of change [1,4,7,32]. In contrast, the

widening of the patch size distribution likely results from the combined effects of

management changes (towards smaller intervention sizes) and increased natural disturbance

activity (resulting in large areas of canopy removal). The same developments also result in

decreasing disturbance severity at the continental scale, with management systems being

optimized to reduce impact [33] and natural disturbance events being frequently characterized

by mixed severities [34]. We conclude that recent trends in Europe's forest disturbance regimes

are strongly driven by the interaction of social and ecological forces, with human resource use

feedbacking to the natural system and *vice versa* (Figure 4).

Fig. 4: Europe's disturbance regimes as coupled human and natural system.

Our results have important implications for the future of Europe's forests. First, by

characterizing the recent disturbance regimes of Europe we provide a baseline to assess future

changes. This is important as disturbances are expected to change considerably in the coming

decades [22,35], yet a sound quantitative baseline for quantifying these changes has been missing

to date. Second, we highlight that both socio-ecological shocks (such as reforestation waves

after the Second World War [3] or the collapse of the Soviet Union [36]) and changes in the

biophysical environment (such as climate change) drive changes in forest disturbance regimes

(Figure 4). This has important consequences for managing the resultant negative

consequences of increasing disturbances on ecosystem service provisioning [10]. In particular,

our analyses suggest that efforts focusing on both the mitigation of climate change and the

compensation of increasing canopy openings in forest policy and management are needed to

bend the curve and stabilize forest canopy turnover at sustainable rates [37]. A further focus

should be on steering the intricate social-ecological interactions driving disturbance regimes

towards dampening feedbacks (Figure 4). For instance, management should focus on creating

land use legacies that reduce the propensity of natural disturbances in the future [28,38], which

will reduce the need for management techniques that cause harm to important ecosystem

services and biodiversity (i.e., salvage logging) [25,39]. Overall, we suggest that acknowledging

complex social-ecological interactions is elementary for managing Europe's forests and key

to creating resilient systems that will sustain important functions and services under an

uncertain future.

Materials and Methods

Reference data

Materials and Methods

Reference data

Acquiring consistent reference data across large areas – such as continental Europe – is

challenging and we therefore make use of manual interpretation of satellite data, serving as

valuable alternative to field-based data [40]. Manual interpretation of satellite data for

calibrating and validating Landsat-based forest change maps is a well-established approach

and has been used in numerous studies previously [41–44]. In essence, an interpreter inspects the

temporal profile of a Landsat pixel and, with the help of Landsat image chips and very high-

resolution imagery available in Google Earth, makes a well-informed call whether the

trajectory represents a stable forest or a forest experiencing a mortality event [45]. We here used

a previously established set of 19,996 interpreted Landsat pixels [4,7]. The initial sample was

drawn at random within forests of Europe, with samples stratified by country. Yet, as

interpreters might declare a plot as no-forest during interpretation (caused by errors in the

automatically generated forest mask), the realized sample size slightly varied (Table S1).

Please note that samples for six Central European countries stem from another study using the

same image interpretation approach but have higher sampling densities. The response design

followed well-documented protocols developed and published previously [4]. Manual

interpretation was done by a total of nine interpreters using established software tools [45], and

the data is freely accessible under following repository:

https://doi.org/10.5281/zenodo.3561925

The reference sample set only consisted of forest pixels and there was thus need for

substituting the sample with non-forest reference pixels. We therefore drew a country-

stratified sample of non-forest pixels using a Landsat-based land cover map from [46]. Each

countries sample size was chosen to match the forest proportion of the respective country

(based on data from the FAOSTATS database), that is the total sample of each country

equaled a random sample across its terrestrial forested and non-forested land surface (see

Table S1). In total we drew 46,461 non-forest reference pixel that, paired with the 19,996

forest reference pixels manually interpreted, totaled to 66,457 reference pixels used for

calibration and validation. From the full reference sample, we randomly drew a sub-sample of

5,000 pixels for map validation, and the remaining 61,457 pixels were used for model

calibration. The validation sub-sample was drawn proportionally to the size of each country to

ensure a consistent and unbiased estimation of map accuracies for the final European map

product.

Mapping disturbances

At the core of our mapping workflow we rely on an established time-series segmentation

approach called LandTrendr [47], implemented in the high-performance cloud-computing

environment Google Earth Engine [48]. In essence, LandTrendr segments annual Landsat pixel

time series into linear features from which a set of metrics can be extracted. We here do not

provide details on the underlying LandTrendr routines but focus on the salient details of our

mapping workflow (please see Figure S2 for a graphical outline of our mapping workflow).

The workflow was based on code published in Kennedy et al. [48].

In a first step we screened all available Tier 1 Landsat 4, 5, 7 and 8 images in the

United States Geological Survey archive. Tier 1 images are delivered as ready-to-use surface

reflectance images including a cloud mask, yet we used coefficients from Roy et al. [49] to

spectrally align the varying sensor types used onboard Landsat 4/5 (Thematic Mapper),

Landsat 7 (Enhanced Thematic Mapper Plus), and Landsat 8 (Operational Land Imager).

After spectral alignment we filtered all available images for summer-season acquisition dates

(1^{st} of June to 30^{th} September) and built annual medoid composites following Flood [50].

Second, we ran LandTrendr for two spectral bands (shortwave infrared I and II) and

two spectral indices commonly used for forest disturbance and mortality mapping [43,44,51–53]:

The Tasseled Cap wetness (TCW) and the Normalized Burn Ration (NBR). We used a

standard parameter set for LandTrendr with no filtering or thresholding and thus allowing for

maximum sensitivity in detecting changes (i.e., allowing for a high commission error).

Third, we extracted the greatest change segment from each pixel's LandTrendr fit to

both spectral bands and both spectral indices. From the greatest change segment we derived a

set of three metrics describing the magnitude, duration and rate of change [51]; as well as a

measure of the signal-to-noise ratio described in Cohen et al. [52]. We further derived the

spectral band/index value prior to, and the rate of change following the greatest change

segment. Similar metrics as the ones used here have been applied also in many previous

studies mapping forest cover changes [42,44,53].

Fourth, we used the set of metrics derived from the greatest change segment for the

two spectral bands and the two spectral indices, the calibration data outlined in the previous

section, and random forest classification [54] to classify each pixel into either no-forest,

undisturbed forest or disturbed forest (i.e., at least one disturbance event during the study

period). This last step filtered out commission errors by LandTrendr and thus greatly

improves mapping accuracy compared to purely automatic algorithms [55]. Yet, we experienced

difficulties in correctly separating forest and no-forest areas solely based on LandTrendr

outputs. This was due to high spectral changes in agricultural areas, which were identified as

disturbances by LandTrendr. To tackle this problem, we added a three-year Tasseled Cap

Brightness, Greenness and Wetness median composite centered on 1985 and 2018,

respectively, to the classification stack. The additional six bands delivered more detailed

spectral information on stable forest and no-forest pixels. Finally, we applied the trained

random forest model to the full classification stack (i.e., LandTrendr metrics from the two

spectral bands and two spectral indices plus the Tasseled Cap composite from 1985 and 2018)

to consistently map the categories no forest, undisturbed forest and forest disturbances across

continental Europe. We validated the final map using the validation sub-sample described in

the previous section. We derived a confusion matrix and report overall accuracy, errors of

commission and errors of omission following best-practice recommendations given in [40].

Fifth, while the thus derived map indicates that a mortality event has happened, they

do not indicate when the mortality event happened. We therefore calculated the disturbance

onset year (i.e., the year of the greatest spectral change) from all spectral bands and spectral

indices using a majority vote. If there was a tie (e.g., all four bands/indices indicated a

different year), we reverted to the median value. To validate this processing step, we

compared the year assigned from LandTrendr to the manually interpreted year of disturbance

for the 19,996 forest reference plots.

Spatial filtering

The last step in creating disturbance maps for continental Europe was to apply a set of spatial

filters for smoothing the resulting disturbance maps and enhancing spatial pattern analysis.

We first set a minimum mapping unit of two 30 × 30 m pixels (i.e., 0.18 ha) and removed all

patches smaller than the minimum mapping unit. In a second filtering step, we identified all

annual patches smaller than the minimum mapping unit and assigned them to the year of the

surrounding pixels, thus smoothing the definition of patches (see Figure S3). In a final

filtering step, we removed holes within mortality patches smaller than the minimum mapping

unit by filling them with the year of the surrounding pixels. While the spatial filtering was

done to improve the spatial analysis described in the following section, we note that the

filtering was applied after the accuracy assessment. The accuracy assessment thus reports the

raw classification performance without additional filtering.

Characterizing disturbance regimes and their changes

From the annual forest disturbance maps we calculated three disturbance regime indicators

based on Turner [17] and Johnstone et al. [28]: disturbance size, frequency and severity.

Disturbance size and severity were first calculated at the patch level and then aggregated to

the landscape level, while disturbance frequency was calculated at the landscape level

directly. Patch size is the number of disturbed pixels for each individual patch (patches were

defined annually using a rook-contiguity) multiplied by pixel size (0.09 ha). For calculating

disturbance frequency, we sub-divided the total study area into a 50 * 50 km hexagon grid

(here representing the landscape scale), totaling to 3,240 hexagons across Europe's land area.

We chose hexagons over squares, as hexagons minimize the spatial difference to the more

complex border of countries and ecoregions used in later analysis. For each hexagon, we then

counted the number of individual patches per year and divided this number by the total forest

area within the hexagon, yielding a measure of the number of patches per km^2 forest area per

year as in indicator of disturbance frequency.

356   For quantifying disturbance severity, we made use of the spectral change magnitude

provided by the LandTrendr analysis. The spectral change magnitude during disturbance is

well correlated with changes in vegetation cover [51,56–59]. Consequently, we here use it as

measure of canopy cover change within a disturbed patch. To combine the spectral change

magnitude from all four spectral bands/indices into one index of canopy cover change we

used logistic regression to predict the occurrence of stand-replacing disturbances from the

four spectral change magnitudes. Data on stand replacing disturbances was generated from

the reference sample by analyzing the manually interpreted land cover after a disturbance

segment. If the land cover switched to non-treed following a disturbance segment (e.g., after

clear-cut harvest or high intensity fire), the disturbance is assigned as stand-replacing. If the

land cover remains treed following a disturbance segment (e.g., following a thinning

operation or a low intensity windthrow), the disturbance is classified as non-stand-replacing.

The method is based on Senf et al. [4] who showed that visual interpretation of post-disturbance

land cover is an accurate measure for separating stand-replacing from non-stand-replacing

disturbances. By predicting the occurrence of stand-replacing disturbances (i.e., complete

removal of the canopy), we scale the spectral change magnitudes to a value between 0 and 1

(or 0 % and 100%), where 1 (or 100 %) indicates complete canopy loss. We validated this

measure of disturbance severity by separating stand-replacing and non-stand-replacing

disturbances solely based on disturbance severity, expecting a high discriminatory power in

separating the different disturbance types.

For spatially visualizing disturbance size, frequency and severity, as well as for

calculating and visualizing trends, we finally aggregated the patch-based metrics (i.e.,

disturbance size and severity) to the landscape level (i.e., the hexagon) by calculating the

arithmetic mean. We report the mean over the median as it is sensitive to changes in both the

central tendency and the spread of the distribution, but we also report other statistics in the

Supplement. Trends in disturbance size, frequency and severity were quantified using a non-

parametric Theil–Sen estimator, which is a non-parametric measure of monotonic trends in

time series insensitive to outliers [60].

Statistical analysis of differences between ecoregions and countries

We used linear mixed effect models (LMM) to test for differences in averages and trends in

disturbance size, frequency and severity among ecoregions and countries nested within

ecoregions. Both ecoregions and countries nested within ecoregions were modelled as random

effects, assuming the individual averages and trends emerging from a common underlying

distribution [61]. For the average modes we also considered years as random effect. From the

fitted LMMs we calculated the variance partition coefficient, that is the amount of variance

explained by ecoregions, countries within ecoregions, and the residual variance within

countries nested within ecoregions (i.e., variance not explained).

References

1. Forest Europe. *State of Europe's forests 2015*. (2015).

2. Fuchs, R., Herold, M., Verburg, P. H., Clevers, J. G. P. W. & Eberle, J. Gross changes in reconstructions of historic land cover/use for Europe between 1900 and 2010. *Global Change Biology* **21**, 299–313 (2015).

3. Ciais, P. *et al.* Carbon accumulation in European forests. *Nature Geoscience* **1**, 425–429 (2008).

4. Senf, C. *et al.* Canopy mortality has doubled across Europe's temperate forests in the last three decades. *Nature Communications* **9**, 4978 (2018).

5. Seidl, R., Schelhaas, M.-J. & Lexer, M. J. Unraveling the drivers of intensifying forest disturbance regimes in Europe. *Global Change Biology* **17**, 2842–2852 (2011).

6. Senf, C. & Seidl, R. Natural disturbances are spatially diverse but temporally synchronized across temperate forest landscapes in Europe. *Global Change Biology* **24**, 1201–1211 (2018).

7. Senf, C., Sebald, J. & Seidl, R. Increasing canopy mortality challenges the future of Europe's forests. (In revision).

8. Nabuurs, G.-J. *et al.* First signs of carbon sink saturation in European forest biomass. *Nature Climate Change* **3**, 792–796 (2013).

9. Seidl, R., Schelhaas, M. J., Rammer, W. & Verkerk, P. J. Increasing forest disturbances in Europe and their impact on carbon storage. *Nature Climate Change* **4**, 806–810 (2014).

10. Thom, D. & Seidl, R. Natural disturbance impacts on ecosystem services and biodiversity in temperate and boreal forests. *Biological Reviews* **91**, 760–81 (2016).

11. Lindner, M. *et al.* Climate change impacts, adaptive capacity, and vulnerability of European forest ecosystems. *Forest Ecology and Management* **259**, 698–709 (2010).

12. Allen, C. D. *et al.* A global overview of drought and heat-induced tree mortality

reveals emerging climate change risks for forests. *Forest Ecology and Management* **259**, 660–

684 (2010).

13.   Trumbore, S., Brando, P. & Hartmann, H. Forest health and global change. *Science*

349, 814–8 (2015).

14.   Millar, C. I., Stephenson, N. L. & Stephens, S. L. CLIMATE CHANGE AND

FORESTS OF THE FUTURE: MANAGING IN THE FACE OF UNCERTAINTY.

Ecological Applications **17**, 2145–2151 (2007).

15.   Seidl, R. The Shape of Ecosystem Management to Come: Anticipating Risks and

Fostering Resilience. *BioScience* **64**, 1159–1169 (2014).

16.   Bengtsson, J., Nilsson, S. G., Franc, A. & Menozzi, P. Biodiversity, disturbances,

ecosystem function and management of European forests. *Forest Ecology and Management*

132, 39–50 (2000).

17.   Turner, M. G. Disturbance and landscape dynamics in a changing world. *Ecology* **91**,

2833–49 (2010).

18.   Asner, G. P. Geography of forest disturbance. *Proceedings of the National Academy of*

Sciences of the United States of America **110**, 3711–2 (2013).

19.   Bebi, P. *et al.* Changes of forest cover and disturbance regimes in the mountain forests

of the Alps. *Forest Ecology and Management* **388**, 43–56 (2017).

20.   Kulakowski, D., Bebi, P. & Rixen, C. The interacting effects of land use change,

climate change and suppression of natural disturbances on landscape forest structure in the

Swiss Alps. *Oikos* **120**, 216–225 (2011).

21.   Munteanu, C. *et al.* Legacies of 19th century land use shape contemporary forest

cover. *Global Environmental Change-Human and Policy Dimensions* **34**, 83–94 (2015).

22.   Sommerfeld, A. *et al.* Patterns and drivers of recent disturbances across the temeprate

forest biome. *Nature Communications* **9**, (2018).

23.   Lindenmayer, D. B. *et al.* Salvage harvesting policies after natural disturbance.

Science **303**, 1303 (2004).

24.   Senf, C., Müller & Seidl, R. Post-disturbance recovery of forest cover and tree height

differ with management in Central Europe. *Landscape Ecology* (2019).

25.   Thorn, S. *et al.* Impacts of salvage logging on biodiversity: A meta-analysis. *Journal*

of Applied Ecology **55**, 279–289 (2018).

26.   Jianguo Liu *et al.* Coupled Human and Natural Systems. *AMBIO: A Journal of the*

Human Environment **36**, 639–649 (2007).

27.   Liu, J. *et al.* Complexity of Coupled Human and Natural Systems. *Science* **317**, 1513–

1516 (2007).

28.   Johnstone, J. F. *et al.* Changing disturbance regimes, ecological memory, and forest

resilience. *Frontiers in Ecology and the Environment* **14**, 369–378 (2016).

29.   Stephens, S. L. *et al.* Temperate and boreal forest mega-fires: characteristics and

challenges. *Frontiers in Ecology and the Environment* **12**, 115–122 (2014).

30.   Brang, P. *et al.* Suitability of close-to-nature silviculture for adapting temperate

European forests to climate change. *Forestry* **87**, 492–503 (2014).

31.   Levers, C. *et al.* Drivers of forest harvesting intensity patterns in Europe. *Forest*

Ecology and Management **315**, 160–172 (2014).

32.   Curtis, P. G., Slay, C. M., Harris, N. L., Tyukavina, A. & Hansen, M. C. Classifying

drivers of global forest loss. *Science* **361**, 1108–1111 (2018).

33.   Mori, A. S. & Kitagawa, R. Retention forestry as a major paradigm for safeguarding

forest biodiversity in productive landscapes: A global meta-analysis. *Biological Conservation*

175, 65–73 (2014).

34.   Meigs, G. W. *et al.* More ways than one: Mixed-severity disturbance regimes foster

structural complexity via multiple developmental pathways. *Forest Ecology and Management*

406, 410–426 (2017).

35.   Dale, V. H. *et al.* Climate Change and Forest Disturbances. *BioScience* **51**, 723–734

(2001).

36.   Kuemmerle, T. *et al.* Post-Soviet farmland abandonment, forest recovery, and carbon

sequestration in western Ukraine. *Global Change Biology* **17**, 1335–1349 (2011).

37.   Lindenmayer, D., Messier, C. & Sato, C. Avoiding ecosystem collapse in managed

forest ecosystems. *Frontiers in Ecology and the Environment* **14**, 561–568 (2016).

38.   Seidl, R., Donato, D. C., Raffa, K. F. & Turner, M. G. Spatial variability in tree

regeneration after wildfire delays and dampens future bark beetle outbreaks. *Proc Natl Acad*

Sci U S A **113**, 13075–13080 (2016).

39.   Dobor, L. *et al.* Is salvage logging effectively dampening bark beetle outbreaks and

preserving forest carbon stocks? *Journal of Applied Ecology* **n/a**,.

40.   Olofsson, P. *et al.* Good practices for estimating area and assessing accuracy of land

change. *Remote Sensing of Environment* **148**, 42–57 (2014).

41.   Potapov, P. V. *et al.* Eastern Europe's forest cover dynamics from 1985 to 2012

quantified from the full Landsat archive. *Remote Sensing of Environment* **159**, 28–43 (2015).

42.   Senf, C., Pflugmacher, D., Hostert, P. & Seidl, R. Using Landsat time series for

characterizing forest disturbance dynamics in the coupled human and natural systems of

Central Europe. *ISPRS J Photogramm Remote Sens* **130**, 453–463 (2017).

43.   Kennedy, R. E. *et al.* Spatial and temporal patterns of forest disturbance and regrowth

within the area of the Northwest Forest Plan. *Remote Sensing of Environment* **122**, 117–133

(2012).

44.   Hermosilla, T., Wulder, M. A., White, J. C., Coops, N. C. & Hobart, G. W. An

integrated Landsat time series protocol for change detection and generation of annual gap-free

surface reflectance composites. *Remote Sensing of Environment* **158**, 220–234 (2015).

45.   Cohen, W. B., Yang, Z. & Kennedy, R. Detecting trends in forest disturbance and

recovery using yearly Landsat time series: 2. TimeSync — Tools for calibration and

validation. *Remote Sensing of Environment* **114**, 2911–2924 (2010).

46.   Pflugmacher, D., Rabe, A., Peters, M. & Hostert, P. Mapping pan-European land

cover using Landsat spectral-temporal metrics and the European LUCAS survey. *Remote*

Sensing of Environment (2018).

47.   Kennedy, R. E., Yang, Z. & Cohen, W. B. Detecting trends in forest disturbance and

recovery using yearly Landsat time series: 1. LandTrendr — Temporal segmentation

algorithms. *Remote Sensing of Environment* **114**, 2897–2910 (2010).

48.   Kennedy, R. *et al.* Implementation of the LandTrendr Algorithm on Google Earth

Engine. *Remote Sensing* **10**, (2018).

49.   Roy, D. P. *et al.* Characterization of Landsat-7 to Landsat-8 reflective wavelength and

normalized difference vegetation index continuity. *Remote Sensing of Environment* **185**, 57–

70 (2016).

50.   Flood, N. Seasonal Composite Landsat TM/ETM+ Images Using the Medoid (a Multi-

Dimensional Median). *Remote Sensing* **5**, 6481–6500 (2013).

51.   Pflugmacher, D., Cohen, W. B. & E. Kennedy, R. Using Landsat-derived disturbance

history (1972–2010) to predict current forest structure. *Remote Sensing of Environment* **122**,

146–165 (2012).

52.   Cohen, W. B., Yang, Z., Healey, S. P., Kennedy, R. E. & Gorelick, N. A LandTrendr

multispectral ensemble for forest disturbance detection. *Remote Sensing of Environment* **205**,

131–140 (2018).

53.   Senf, C., Pflugmacher, D., Wulder, M. A. & Hostert, P. Characterizing spectral–

temporal patterns of defoliator and bark beetle disturbances using Landsat time series. *Remote*

Sensing of Environment **170**, 166–177 (2015).

54.   Breiman, L. Random Forests. *Machine Learning* **45**, 5–32 (2001).

55.   Cohen, W. *et al.* How Similar Are Forest Disturbance Maps Derived from Different

Landsat Time Series Algorithms? *Forests* **8**, 98 (2017).

56.   Bright, B. C., Hudak, A. T., Kennedy, R. E. & Meddens, A. J. H. Landsat Time Series

and Lidar as Predictors of Live and Dead Basal Area Across Five Bark Beetle-Affected

Forests. *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*

7, 3440–3452 (2014).

57.   Zald, H. S. J. *et al.* Integrating Landsat pixel composites and change metrics with lidar

plots to predictively map forest structure and aboveground biomass in Saskatchewan, Canada.

Remote Sensing of Environment **176**, 188–201 (2016).

58.   Meigs, G. W., Kennedy, R. E., Gray, A. N. & Gregory, M. J. Spatiotemporal

dynamics of recent mountain pine beetle and western spruce budworm outbreaks across the

Pacific Northwest Region, USA. *Forest Ecology and Management* **339**, 71–86 (2015).

59.   Meddens, A. J. H. & Hicke, J. A. Spatial and temporal patterns of Landsat-based

detection of tree mortality caused by a mountain pine beetle outbreak in Colorado, USA.

Forest Ecology and Management **322**, 78–88 (2014).

60.   Wilcox, R. R. *Fundamentals of modern statistical methods: Substantially improving*

power and accuracy. (Springer, 2010).

61.   Gelman, A. Multilevel (Hierarchical) Modeling: What It Can and Cannot Do.

Technometrics **48**, 432–435 (2006).

Acknowledgements

C. Senf acknowledges funding from the Austrian Science Fund (FWF) Lise-Meitner Program

(Nr. M2652). R. Seidl acknowledges funding from FWF START grant Y895-B25. We thank

Justin Brasten (Oregon State University) for making the code of LandTrendr open source,

which greatly helped in implementing this research.

Author contribution

CS and RS designed the research; CS performed all computations and analyses; CS wrote the

manuscript with input from RS.

Data availability

All maps presented in this research paper will be made publicly available after peer-review.

Code availability

The code used for processing the Landsat data is available at https://github.com/eMapR/LT-

GEE. The research code used for creating the maps will be made available after peer-review.