The adaptive response to long term nitrogen starvation in *Escherichia coli* requires the

breakdown of allantoin

Amy Switzer, Lynn Burchell, Josh McQuail and Sivaramesh Wigneshweraraj[1]*

[1]MRC Centre for Molecular Bacteriology and Infection, Imperial College London, London,

SW7 2AZ, UK

*Corresponding author: E-mail: s.r.wig@imperial.ac.uk; Tel.: +44 207 594 1867.

ABSTRACT

Bacteria initially respond to conditions that attenuate their growth by eliciting large-scale transcriptional changes. The accompanying changes in gene expression and metabolism allow the bacterial cells to effectively adapt to the growth attenuated state. How the transcriptome subsequently changes as growth attenuation ensues is not well understood. We used nitrogen (N) starvation as a model nutrient starvation condition to study the transcriptome of growth attenuated *Escherichia coli*. The results reveal that the transcriptome of nitrogen starvation-induced growth attenuated *E. coli* remains dynamic and perturbations to it compromise the viability of growth attenuated bacteria and their ability to effectively recover growth when N starvation conditions become alleviated. We further reveal that, over time, N starvation-induced growth attenuated bacteria rely on the degradation of allantoin for optimal growth recovery when N becomes replenished. This study provides insights into the temporally coordinated adaptive responses that occur in *E. coli* experiencing sustained N starvation.

IMPORTANCE

Bacteria in their natural environments seldom encounter conditions that support continuous growth. Hence, many bacteria spend the majority of their time in states of little or no growth due to starvation of essential nutrients. To cope with prolonged periods of nutrient starvation, bacteria have evolved several strategies, primarily manifesting themselves through changes in how the information in their genes is accessed. How these coping strategies change over time under nutrient starvation is not well understood and this knowledge is not only important to broaden our understanding of bacterial cell function, but also to potentially find ways to manage harmful bacteria. This study provides insights into how nitrogen starved *Escherichia coli* bacteria rely on different genes during long term nitrogen starvation.

KEYWORDS

Escherichia coli, nitrogen regulation, bacterial stress response, transcriptome, allantoin

INTRODUCTION

The process of RNA synthesis, transcription, is often the first decision-making and regulatory point in the flow of genetic information and is catalysed by the multisubunit enzyme, RNA polymerase (RNAP). The transcriptional responses that establish and maintain specific cellular states are dynamic and this plasticity is often controlled by many transcription regulatory factors that communicate with and influence the activity of the RNAP. Conditions that sustain constant bacterial growth are seldom found in nature. Whether in a host or the environment, bacteria frequently encounter nutrient deprivation, which can attenuate bacterial growth. Hence, the growth attenuated state is now widely acknowledged as an important physiological state in bacterial cell function. However, although the transcriptional basis of *bacterial growth* is widely studied, our understanding of the transcriptome of growth attenuated bacteria remains in its infancy. In fact, several recent studies on different bacteria have revealed that growth attenuated bacteria are transcriptionally active. For example: Gefen et al showed that a batch culture of *E. coli* cells retain transcriptional activity in the stationary phase of growth over a period of nearly 2 days (1); Stapels et al showed that non-growing intracellular *Salmonella* in host macrophages are transcriptionally active and this activity is important to shape the host environment to benefit bacterial survival (2); similarly, non-replicating *Mycobacterium tuberculosis* cells retained transcriptional activity in the lungs of mice (3); and more recently, Gray et al reported that *Bacillus subtilis* cells under oligotrophic conditions are transcriptionally active and replicate at an extremely slow rate (4). Therefore, understanding the transcriptional landscape in growth attenuated bacteria is important as this knowledge could inform and inspire new antibacterial strategies, such as disrupting genetic networks and

products that are activated during growth attenuation, which subsequently become essential for

survival and/or optimal growth recovery.

In this study, we used nitrogen (N) starvation as a model condition to understand how

the transcriptional landscape in nutrient starvation-induced growth attenuated *E. coli* affects

bacterial survival and ability to recover growth when nutrients (i.e. N) become replenished.

Nitrogen (N) is an integral nutrient to bacterial cell function since N is an essential element of

most macromolecules in the bacterial cell (proteins, metabolites, nucleic acids and cell wall

components). Therefore, an exponentially growing batch culture of *E. coli* rapidly becomes

growth attenuated upon N run-out. The initial adaptive response to N run-out, the N regulation

(Ntr) stress response, results in the large scale and multilevel reprogramming of the

transcriptome, involving ~40% of all *E. coli* genes, to allow *E. coli* to adjust to N starvation.

Since *E. coli* perceive N run-out as a decrease in the internal concentration of glutamine, the

genes expressed as part of the Ntr response are predominantly involved in the transport and

assimilation of nitrogenous compounds into glutamine or glutamate, yielding these

catabolically or sparing the requirement for them in cellular processes (5, 6). However,

although our understanding of the initial adaptive transcriptional response to N starvation is

well established, how the transcriptome changes as N starvation-induced growth attenuation

(NSIGA) ensues is far from clear.

RESULTS AND DISCUSSION

Perturbation of the transcriptional response during NSIGA compromises survival

and growth recovery of *E. coli*. To setup an experimental system to study how the

transcriptome changes during NSIGA, we grew a batch culture of *E. coli* in a defined minimal

growth medium in the presence of a limiting amount (3 mM) of ammonium chloride as the sole

N source (5). Under these conditions, when ammonium chloride runs out, bacteria enter a

complete state of N starvation, and subsequent growth attenuation (5). From the onset of N

starvation (N-), the viability of the population of growth attenuated *E. coli*, as measured by the

number of colony forming units (c.f.u.) in the population, increases at a slow rate over the

initial 24 h under N starvation (doubling time ~39 h, which is ~80-fold less than during

exponential growth when N is abundant; N+) (Fig. 1A, dark orange area). However, as N

starvation ensues beyond 24 h (N-24), the population ceases growth before the number of

viable cells gradually begins to decline after 48 h under N starvation (Fig. 1A, light orange

area). Thus, it seems that 24 h following N run-out, which we refer to as the N starved adaptive

phase (NAP), represents an important period in which metabolic processes, including

transcription, are temporally adjusted in order to balance maintenance of growth (albeit at a

very slow rate) and maximising chances of survival and growth recovery when N starvation

stress is alleviated.

To evaluate the importance of the transcriptional changes that occur during the NAP,

we determined how perturbation of the transcriptome during the NAP affects cell survival and

growth recovery when N starvation stress is alleviated. To do this, we treated the bacteria with

the bacteriostatic antibiotic rifamycin (Rif) at ~1 h following N run-out (i.e. N-1) for ~23 h.

Rif was used at 150 μg/ml as, under our experimental conditions, this concentration of Rif

prevents transcription elongation by the RNAP by trapping it at promoters (5). We initially

compared the proportion of viable cells in the Rif treated and untreated population of bacteria

at N-24 (i.e. ~23 h following Rif treatment at N-1). As shown in Fig. 1B, at N-24, only ~50%

cells in the Rif treated culture were viable compared to the Rif untreated culture. In control

experiments, the addition of the same amount of Rif (150 μg/ml) to bacteria at N+ (i.e. during

exponential growth when N was available) did not affect cell viability. This suggests that ~150

μg/ml of Rif is not bactericidal under our experimental conditions and that the reduction in

viability of the Rif treated population during the NAP in Fig. 1B is due to Rif-induced

perturbations to the transcriptome during the NAP. In other words, Rif-induced perturbation of the transcriptional changes during NAP has a detrimental effect on cell viability. Rif treatment also had a profound effect on the ability of bacteria from N-24 to resume growth compared to untreated bacteria: as shown in Fig. 1C and 1D, the lag time to growth recovery of the Rif treated bacteria (following c.f.u. correction to ensure an equal number of viable treated and untreated bacteria were used; Fig. 1B) from N-24 was delayed by ~4.8 h compared to untreated bacteria. To discount the possibility that the difference in lag time between Rif treated and untreated bacteria was due to the presence of a subpopulation of bacteria that resumed growth heterogeneously, we measured the lag time distribution of appearance of individual colonies during growth recovery using ScanLag (7). As shown in Fig. 1E, no heterogeneity in the populations of Rif treated and untreated bacteria from N-24 was detected. Further, the ScanLag data also substantiates the data in Fig. 1C and 1D and demonstrates that Rif treated cells have an increased lag time to growth recovery compared to untreated cells. Although, once growth resumed, the doubling time of Rif treated and untreated bacteria did not markedly differ (Fig. 1D). Overall, we conclude that perturbation of the transcriptome during the NAP has a detrimental effect on the ability of *E. coli* experiencing NSIGA to maximise chances of viability and optimal growth recovery when N becomes available.

The transcriptome *E. coli* in the NAP is dynamic. To study the transcriptional changes that occur during the NAP, we compared the transcriptomes of *E. coli* during exponential growth (N+), following initial N run-out (N-), and at N-24. We defined differentially expressed genes as those with expression levels changed ≥ 2-fold with a false discovery rate adjusted P-value < 0.05. As previously reported by us and others, the initial transcriptional response to N starvation involves differential expression of 1787 genes primarily required for scavenging and catabolism of alternative N sources (6, 8, 9) and thus,

we did not investigate these genes further (Phase A in Fig. 2A). However, during the NAP,

upon transition from N- to N-24 (Phase B in Fig. 2A), 451 genes (~10% of *E. coli* genes) are

differentially expressed, of which 105 genes (~2% of *E. coli* genes) become differentially

expressed in Phase B, that were not differentially expressed in Phase A (Fig. 2B). Functional

categorisation of the genes uniquely differentially expressed in Phase B, hereafter called NAP

genes, revealed that they are involved in diverse processes including transcription, amino acid

transport, metabolism and several genes of unknown function (Fig. 2C). Eleven (*abgR*, *bfd*,

ileY, *leuO*, *ompN*, *sibE*, *tauA*, *ycaD*, *yedL*, *yhdV*, and *yihO*) of the upregulated and nine (*evgA*,

glnA, *gltI*, *hemH*, *ivy*, *pfkB*, *ybjX*, *yieP*, and *yjeJ*) of the downregulated NAP genes were known

to be regulated by either NtrC (*glnA* and *gltI*) or Nac (all other genes listed above) (9, 10) - the

master transcription regulators that activate the initial adaptive response to N starvation.

However, the absence of either *glnG* (which encodes for NtrC) or *nac* did not detectably affect

viability of bacteria under NSIGA or the ability of NSIGA bacteria to recover growth when

inoculated into fresh N replete minimal growth media (Fig. 2D and 2E). Overall, we conclude

that the transcriptome of *E. coli* during the NAP is dynamic, and this is consistent with the

finding above that Rif treatment during the NAP has a detrimental effect on the viability of the

E. coli population under NSIGA and its ability to recover growth when N becomes replenished.

However, it seems that the adaptive transcriptional changes that occur during the NAP happen

subsequently to the initial adaptive changes to NISGA and hence do not depend on NtrC (and

Nac), which coordinates the initial adaptive transcriptional response to NSIGA.

Four upregulated NAP genes are important for optimal growth recovery from

NSIGA. We next investigated how the upregulated NAP genes affected viability under NSIGA

and growth recovery from NSIGA. Of the 51 51 upregulated NAP genes 44 are available in

the Keio single gene knockout collection (11). Thus, we determined how the absence of any of

the upregulated NAP genes affected growth recovery from NSIGA. To do this, we grew the

library of upregulated NAP gene mutant bacteria in minimal growth media containing excess

(10 mM) ammonium chloride as the sole N source and measured cell density after 16 h of

growth. The growth of the library of upregulated NAP gene mutant strains was done in a plate

reader incubator which allowed continuous monitoring of growth of each mutant strain

simultaneously to allow better side-by-side comparison. The results shown in Fig. 3A indicate

that absence of any of the NAP genes did not markedly affect growth under N replete

conditions. However, we cannot exclude the possibility that the relative dynamics of growth of

the individual mutant strains could have differed from that of the wild-type (WT) strain over

the 16 h period of time. We note that bacteria in which the tryptophan biosynthesis genes are

deleted, lack γ subunit of formate dehydrogenase-N (*fdnI*) or ferric rhodotorulic acid outer

membrane transporter (*fhuE*), are unable to grow in our minimal growth media, most likely

due to auxotrophy (12). Next, we reinoculated bacteria that grew under N replete conditions

into fresh minimal growth media containing limiting, i.e. 3 mM, ammonium chloride and

determined how the absence of the upregulated NAP genes affected viability at N-24, i.e.

following 24 h under NSIGA. The results in Fig. 3B show that the viabilities of the upregulated

NAP gene mutants were similar to that of the WT strain. However, when the 24 h N starved

mutant bacteria were inoculated into fresh N replete minimal growth media, the dynamics of

growth recovery for the majority of the mutant strains were similar to that of the WT strain

(Fig. 3C). However, Δ*allB*, Δ*leuO*, Δ*ynfP*, and Δ*ybcJ* mutant strains displayed noticeably

compromised growth recovery properties compared to that of the WT strain (Fig. 3C). These

genes encode an allantoinase (*allB*), a DNA-binding transcription dual regulator (*leuO*), a

small, possibly fragmented, protein of unknown function (*ynfP*), and a putative RNA-binding

protein (*ybcJ*) (12). As shown in Fig. 3D, although the doubling time of Δ*allB*, Δ*leuO*, Δ*ynfP*,

and Δ*ybcJ* mutant strains were similar to that of the WT strain, their lag time to growth recovery

was increased by ~1.5-4 h compared to that of the WT strain. Thus, it seems that the

transcription of *allB*, *leuO*, *ynfP*, and *ybcJ* during the NAP are required for optimal growth

recovery from NSIGA. As the Δ*allB* mutant strain displayed the most pronounced

compromised ability to recover growth (Fig. 3C) when compared to the WT strain, we decided

to focus on the role of *allB* during the NAP and recovery from NSIGA.

The degradation of allantoin during the NAP is required for optimal growth

recovery from NSIGA. To discount the possibility that the difference in lag time during

growth recovery between the WT and Δ*allB* bacteria was not due to the presence of a

subpopulation of slow growing Δ*allB* bacteria, we measured the lag time distribution of

appearance of individual colonies during growth recovery using ScanLag (7) as described in

Fig. 1E. As shown in Fig. 4A, we observed the expected lag time in growth recovery for the

Δ*allB* bacteria, but no evidence of a slow growing subpopulation was seen. Further, we did

not detect a lag in growth recovery of WT and Δ*allB* bacteria sampled from N- when

reinoculated into N replete fresh growth medium (Fig. 4B). This result underscores the view

that *allB* has a specific role in *E. coli* during the NAP. To establish that the increased lag time

in growth recovery of Δ*allB* bacteria was directly due to the absence of *allB*, we complemented

the Δ*allB* bacteria with plasmid-borne *allB*. As shown in Fig. 4C and 4D, the lag time in growth

recovery of Δ*allB* bacteria reverted to that of WT bacteria when *allB* was supplied via a

plasmid. The product of *allB* is allantoinase, a tetrameric enzyme that catalyses the hydrolytic

conversion of (*S*)-(+)-allantoin to allantoate. A previous study (13) reported that substitutions

at invariant amino acid positions N94 and S317 (in both cases to D and V), which are located

proximal to the active centre, rendered allantoinase inactive (Fig. 4E). As shown in Fig. 4C

and 4D, in the context of the complementation experiments, plasmid-borne *allB*-N94V and

allB-S317V failed to fully decrease the lag time in growth recovery of Δ*allB* bacteria as seen

with the WT version of *allB*. In contrast, plasmid-borne *allB*-N94D and to a larger extent *allB*-

S317D reverted the lag time in growth recovery of Δ*allB* bacteria as seen with the WT version

of *allB*. Although we are unable to explain why specifically substitution by D but not V at N94

or S317 renders allantoinase catalytically inactive under our experimental conditions, it seems

that the degradation of allantoin occurs at some point in *E. coli* experiencing NSIGA. This is

conceivable because, as shown in Fig. 4F, the conversion of allantoate (the product of

allantoinase) to S-ureidoglycine by allantoate amidohydrolase (AllC) and the subsequent

conversion of (*S*)-ureidoglycine to (S)-ureidoglycolate by (*S*)-ureidoglycine aminohydrolase

(AllE) produces ammonium, which may be needed for maintenance of cellular functions

required for optimal growth recovery from NSIGA. Consistent with this view, we observed

that 24 h N starved Δ*allC* and Δ*allE* bacteria displayed a similarly increased lag time to growth

recovery as the Δ*allB* bacteria (Fig. 4G and 4H) when grown in fresh N replete growth media.

As allantoin is a product of the purine degradation pathway in N starved *E. coli* (14), we

considered whether the requirement for *allB* is a consequence of purine degradation in *E. coli*

during the NAP. In other words, we considered whether purines are degraded to replenish

diminishing N sources during the NAP to sustain cellular functions via the allantoin

degradation pathway (as in Fig. 4F). To indirectly determine whether purines are degraded in

E. coli during the NAP, we measured the integrity of total RNA as a surrogate for purine

degradation in *E. coli* as a function of time in the NAP by chip-based capillary electrophoresis

(see Materials and Methods). As shown in Fig. 4J, strikingly, we observed a gradual

degradation of 16S and 23S ribosomal RNA as the NAP ensued. This observation further

underscores the view that RNA landscape and turnover during the NAP is plastic and consistent

with the view that the transcriptome of *E. coli* under NSIGA is dynamic. Collectively, the

results are support the view that *allB* contributes to optimal growth recovery from NSIGA by

the degradation of allantoin during the NAP in *E. coli*, which most probably accumulates as a

result of purine degradation.

CONCLUSIONS

N is central to all metabolic process in the bacterial cell. As such, bacteria can utilize a wide

range of N containing compounds as sole sources of cellular N. Upon N run-out, *E. coli* cells

activate the transcription of genes associated with N assimilatory metabolic pathways

necessary for utilization of N from the extracellular environment and catabolic biosynthetic

pathways for intracellular assimilation of N. We have presented evidence that the growth a

population of N starved *E. coli* does not completely cease upon N run-out and that *E. coli* are

transcriptionally active for a period of up to 24 h – the NAP - following entry into NSIGA. It

seems that NAP represents an important adaptive phase in *E. coli* experiencing NSIGA to

maximise chances of viability and optimise growth recovery should N become replenished.

We have also presented evidence that the adaptive transcriptional response to NSIGA in *E. coli*

during the NAP occurs in at least two distinct temporally coordinated phases: The NtrC

dependent initial response to NSIGA is followed by – to a large extent – a NtrC independent

transcriptional response. We have uncovered that an aspect of the adaptive transcriptional

response that occurs in the NAP subsequent to the initial adaptive transcription response in *E.*

coli under NSIGA involves the degradation of allantoin. As ammonium is a by-product of

degradation of allantoin, we propose that allantoin degradation allows the assimilation of

ammonium into glutamine and glutamate to maintain cellular processes during the NAP to

maximise chances of viability and optimise growth recovery when N becomes replenished.

Previous studies have shown that allantoin can *only* be used as the sole N source by *E. coli*

grown under anaerobic conditions (15, 16), and consistent with these results, we were unable

to grow ΔallB and WT bacteria under our conditions when ammonium chloride (the sole N

source used in our experiments) was replaced by allantoin - in both, the initial growth and

recovery growth experiments (data not shown). Therefore, as the end-product of allantoin

degradation under aerobic conditions is 3-phosphoglycerate, in addition to the by-product

ammonium, we propose that the breakdown of allantoin, which likely occurs as a consequence

of purine catabolism during the NAP in *E. coli* under NSIGA, serves to maintain as some sort

of 'metabolic threshold' to allow optimal growth recovery of *E. coli* from NSIGA under

aerobic conditions when N becomes replenished. Although future work will focus on

deciphering the metabolic landscape to unravel the metabolic adaptive responses that define

the NAP in *E. coli* under NSIGA, this work has underscored the importance of studying the

transcriptional changes that occur in growth attenuated bacteria to further our understanding of

bacterial stress physiology.

MATERIALS AND METHODS

Bacterial strains and plasmids, growth conditions and viability measurements.

WT and mutant *E. coli* BW25113 strains were used in this study (11). Bacteria were grown as

described in Figueira et al (17) in 10 mM NH_4Cl (for overnight cultures) or 3 mM NH_4Cl (for

inducing NSIGA) Gutnick minimal medium, consisting of 33.8 mM KH_2PO_4 , 77.5 mM

K_2HPO_4 , 5.74 mM K_2SO_4 , 0.41 mM $MgSO_4$, supplemented with Ho-LE trace elements and

0.4 % (w/v) glucose, using NH_4Cl as the sole N source (9). Where relevant, rifampicin was

added at a final concentration of 150 μg/ml at N-1 or N+ as stated, and L-arabinose was added

to complementation strains containing the pBAD18 plasmid at 0.2% (w/v) for induction of

gene expression in growth cultures. Ampicillin was added at a final concentration of 100 μg/ml

to all cultures containing derivatives of the pBAD18 plasmid. Measurement of growth was

taken by measuring the optical density (OD_{600nm}) of bacterial cultures over time to generate

growth curves. To determine population viability, the number of viable cells was calculated by

conducting serial dilutions of cultures at designated time points as shown in relevant figures,

grown overnight on Luria-Bertani (LB) agar plates, and measuring colony forming units (c.f.u.)

per ml. The pBAD18-*allB* plasmid was made by traditional restriction enzyme cloning to

incorporate the ribosomal binding site of pET33b(+) with the *allB* gene to allow arabinose-

inducible expression of the AllB protein. The N94D, N94V, S317D and S317V mutations in

allB were done by standard site-directed mutagenesis using pBAD18-*allB* as the template.

ScanLag. Bacteria we grown in 3 mM Gutnick minimal medium to N-24, washed twice

in 1 ml PBS, and diluted in PBS. One hundred microlitres of diluted culture was spread on an

LB agar plate and incubated at 33 °C for 48 h with pictures of plates taken every 20 minutes.

Analysis of colony growth rate and appearance time was modified from Levin-Reisman et al

(18) by Dr Miles Priestman and is available at

https://github.com/mountainpenguin/NQBMatlab.

RNA-sequencing (RNA-seq). The RNA-seq data set was previously generated and

analysed by Switzer et al (8). The analysis of differential gene expression during NAP as well

as the statistical analyses were also conducted in the same way as described previously. The

sequencing data can be accessed under accession number E-MTAB-6491 in the ArrayExpress

database at EMBL-EBI (www.ebi.ack.uk/arrayexpress).

Total RNA analysis. Cultures were grown as above and sampled at the following time

points: at the onset of N starvation (N-) and 3 h (N-3), 6 h (N-6), 9 h (N-9), 12 h (N-12), 18 h

(N-18), and 24 h (N-24) under N starvation. At least three biological replicates were taken at

each time point. RNA was acquired and stabilized at specified time points using Qiagen RNA

Protect reagent (Qiagen, 76526). RNA extraction was carried out using the PureLink RNA

mini kit (Invitrogen, 12183025) and DNase Set (Invitrogen, 12185010) as per manufacturer's

protocol. Electropherograms of total RNA integrity were generated using an Agilent

Bioanalyser 2100 instrument (Agilent, G2939BA) and RNA 6000 Nano kit (Agilent, 5067-

1511) as per the manufacturer's protocol.

Statistical analysis and growth calculations. Unless otherwise stated in the text, all

growth curves and c.f.u. data show the mean average of three independent experiments and

variation calculated by the standard error of the mean (SEM). The student's t-test or one-way

ANOVA was used to determine statistical significance if the probability (P) value was < 0.05

(* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$). Lag and doubling times were calculated by first

generating a straight-line graph of the exponential growth section of each growth curve. Lag

time was defined as the intersection of the straight line graph with the x axis, while the doubling

time was determined during exponential growth (19).

FIGURE LEGENDS

Fig 1. Perturbation of the transcriptional response during NSIGA compromises survival

and growth recovery of *E. coli*. (A) Graph showing viability, measured by counting colony

forming units (c.f.u.) of *E. coli* over 5 days under NSIGA. Inset table shows doubling time

between different growth periods. (B) Graph showing proportion of viable cells (measured by

c.f.u.) in the population of *E. coli* under NSIGA with (red) and without (blue) Rif treatment at

N-1; the dashed red line indicates addition of Rif to N+ cells (see text for details). (C) Graph

showing recovery growth in fresh N replete medium as measured by OD_{600nm} readings. The

time points and conditions from which bacteria were sampled for recovery growth experiments

are indicated and discussed in the text. (D) Graph showing lag time and doubling time during

growth recovery experiments from (C). (E) ScanLag analysis of colony growth rate and

appearance time after plating of 24 h N starved bacteria on LB agar after Rif treatment (red)

and no Rif treatment (blue) at N-1. The various shades of the same colour represent distinct

biological replicates, where n=3. In (A)-(D), error bars represent SEM, where n=3 and the

schematics describe the different experiments for ease of understanding.

Fig 2. The transcriptome of NSIGA *E. coli* is dynamic. (A) Graphs showing the $\log_2$ fold

change in gene expression at N- compared to N+ (Phase A), and N-24 compared to N- (Phase

B). Genes in blue (Phase A) or red (Phase B = NAP) represent significant differential

expression above a $\log_2$ fold change of 1 and a false discovery rate adjusted *P* value < 0.05.

Genes in black (Phase B) represent the NAP genes, defined as having significant differential

gene expression unique to Phase B (NAP), but not Phase A. (B) Venn diagrams showing genes

significantly upregulated (top) and downregulated (bottom) in Phase A compared to Phase B.

(C) Graph showing Clusters of Orthologous Groups (COG) categorisation of NAP genes, and

the number of genes within each category whose expression is up- or downregulated from N-.

(D) Graph showing viability as measured by c.f.u. counting of viable cells in the population of

WT (blue), Δ*glnG* (red) and Δ*nac* (green) bacteria over 48 h of N starvation (*left*). Graph

showing growth recovery of N-24 WT (blue), Δ*glnG* (red) and Δ*nac* (green) bacteria in N

replete fresh media as measured by OD_{600nm} readings. In both graphs, error bars represent SEM,

where n=3.

Fig 3. Four upregulated NAP genes are important for optimal growth recovery from

NSIGA. (A) Graph showing overnight growth of NAP gene mutant bacteria as measured by

OD_{600nm} compared to WT (blue) bacteria in minimal growth media containing excess (10 mM)

N (ammonium chloride). (B) Graph showing viability as measured by c.f.u. counts of wild-

type (blue) and NAP gene mutant bacteria (black) bacteria at N-24. (C) Graph showing

recovery growth of 24 h N starved NAP gene mutant bacteria in N replete fresh media as

measured by OD_{600nm} readings. The recovery growth curves of WT (blue), Δ*allB* (red), Δ*leuO*

(dark green), Δ*ynfP* (purple), and Δ*ybcJ* (orange) are highlighted. (D) Graph showing lag time

(blue) and doubling time (red) of WT, Δ*allB*, Δ*leuO*, Δ*ynfP*, and Δ*ybcJ* bacteria extracted from

the growth curves in (C). In (C) and (D), the error bars represent SEM, where *n*=2. Statistical

analyses were performed using the Student's *t*-test (*P < 0.05, **P < 0.01).

Fig. 4. The degradation of allantoin during the NAP is required for optimal growth

recovery from NSIGA. (A) ScanLag analysis of colony growth rate and appearance time after

plating of 24 h N starved WT (blue) and Δ*allB* (red) bacteria on LB agar. Various shades of

the same colour represent distinct biological replicates, where n=3. (B) Graph showing the

recovery growth curves of WT (blue) and Δ*allB* (red) bacteria from the onset of N starvation

(N-) in fresh N replete media as measured by OD_{600nm} readings of the cultures. (C) Graph

showing the recovery growth curves of 24 h N starved WT + pBAD18 (blue), Δ*allB* + pBAD18

(red), and Δ*allB* + pBAD18-*allB* (green) bacteria in fresh N replete media as measured by

OD_{600nm} readings of the cultures. (D) Graph showing lag time to growth recovery of WT +

pBAD18 (blue), Δ*allB* + pBAD18 (red), Δ*allB* + pBAD18-*allB* (bright green), Δ*allB* +

pBAD18-*allB* N94D, Δ*allB* + pBAD18-*allB* N94V, Δ*allB* + pBAD18-*allB* S317D and Δ*allB*

+ pBAD18-*allB* S317V in fresh N replete media as extracted by OD_{600nm} readings of the

cultures. (E) Cartoon representation of the structure of a monomer of AllB in which the

positions of the N94 and S317 are indicated in orange relative to the two Fe moieties at the

active site of the enzyme (indicated in red). (F) A schematic showing the allantoin degradation

pathway indicating the steps where ammonia is produced. (G) As in (C), but the experiment

was conducted with Δ*allC* and Δ*allE* bacteria. (H) Graph showing lag time (blue) and doubling

time (red) of WT, Δ*allB*, Δ*allC* and Δ*allE* bacteria extracted from the growth curves in (G). (I)

Electropherogram of total RNA from bacterial cultures taken at indicated timepoints. Graph

shows representative data from at least three biological replicates. 16S and 23S rRNA peaks

are indicated. In (B), (C), (D), (G) and (H) error bars represent SEM, where n=3.

ACKNOWLEDGMENTS

A Wellcome Trust Investigator award WT100958MA and a Leverhulme Trust project grant

(RPG-2017-431) funded this work. J. M. is an MRC funded PhD student.

REFERENCES

1.   Gefen O, Fridman O, Ronin I, Balaban NQ. 2014. Direct observation of single
stationary-phase bacteria reveals a surprisingly long period of constant protein
production activity. Proc Natl Acad Sci U S A 111:556-61.
2.   Stapels DAC, Hill PWS, Westermann AJ, Fisher RA, Thurston TL, Saliba AE,
Blommestein I, Vogel J, Helaine S. 2018. Salmonella persisters undermine host
immune defenses during antibiotic treatment. Science 362:1156-1160.
3.   Manina G, Dhar N, McKinney JD. 2015. Stress and host immunity amplify
Mycobacterium tuberculosis phenotypic heterogeneity and induce nongrowing
metabolically active forms. Cell Host Microbe 17:32-46.
4.   Gray DA, Dugar G, Gamba P, Strahl H, Jonker MJ, Hamoen LW. 2019. Extreme
slow growth as alternative strategy to survive deep starvation in bacteria. Nat
Commun 10:890.
5.   Brown DR, Barton G, Pan Z, Buck M, Wigneshwararaj S. 2014. Nitrogen stress
response and stringent response are coupled in Escherichia coli. Nat Commun 5:4115.
6.   Zimmer DP, Soupene E, Lee HL, Wendisch VF, Khodursky AB, Peter BJ, Bender
RA, Kustu S. 2000. Nitrogen regulatory protein C-controlled genes of Escherichia
coli: scavenging as a defense against nitrogen limitation. Proc Natl Acad Sci U S A
97:14674-9.
7.   Levin-Reisman I, Fridman O, Balaban NQ. 2014. ScanLag: high-throughput
quantification of colony growth and lag time. J Vis Exp doi:10.3791/51456.
8.   Switzer A, Evangelopoulos D, Figueira R, de Carvalho LPS, Brown DR,
Wigneshwararaj S. 2018. A novel regulatory factor affecting the transcription of
methionine biosynthesis genes in Escherichia coli experiencing sustained nitrogen
starvation. Microbiology 164:1457-1470.
9.   Brown DR, Barton G, Pan Z, Buck M, Wigneshwararaj S. 2014. Combinatorial stress
responses: direct coupling of two major stress responses in Escherichia coli. Microb
Cell 1:315-317.
10. Aquino P, Honda B, Jaini S, Lyubetskaya A, Hosur K, Chiu JG, Ekladious I, Hu D,
Jin L, Sayeg MK, Stettner AI, Wang J, Wong BG, Wong WS, Alexander SL, Ba C,
Bensussen SI, Bernstein DB, Braff D, Cha S, Cheng DI, Cho JH, Chou K, Chuang J,
Gastler DE, Grasso DJ, Greifenberger JS, Guo C, Hawes AK, Israni DV, Jain SR,
Kim J, Lei J, Li H, Li D, Li Q, Mancuso CP, Mao N, Masud SF, Meisel CL, Mi J,
Nykyforchyn CS, Park M, Peterson HM, Ramirez AK, Reynolds DS, Rim NG, Saffie

JC, Su H, Su WR, et al. 2017. Coordinated regulation of acid resistance in Escherichia
coli. BMC Syst Biol 11:1.
11.   Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita
M, Wanner BL, Mori H. 2006. Construction of Escherichia coli K-12 in-frame,
single-gene knockout mutants: the Keio collection. Mol Syst Biol 2:2006 0008.
12.   Keseler IM, Mackie A, Santos-Zavaleta A, Billington R, Bonavides-Martinez C,
Caspi R, Fulcher C, Gama-Castro S, Kothari A, Krummenacker M, Latendresse M,
Muniz-Rascado L, Ong Q, Paley S, Peralta-Gil M, Subhraveti P, Velazquez-Ramirez
DA, Weaver D, Collado-Vides J, Paulsen I, Karp PD. 2017. The EcoCyc database:
reflecting new knowledge about Escherichia coli K-12. Nucleic Acids Res 45:D543-
D550.
13.   Kim K, Kim MI, Chung J, Ahn JH, Rhee S. 2009. Crystal structure of metal-
dependent allantoinase from Escherichia coli. J Mol Biol 387:1067-74.
14.   Xi H, Schneider BL, Reitzer L. 2000. Purine catabolism in Escherichia coli and
function of xanthine dehydrogenase in purine salvage. J Bacteriol 182:5332-41.
15.   Rintoul MR, Cusa E, Baldoma L, Badia J, Reitzer L, Aguilar J. 2002. Regulation of
the Escherichia coli allantoin regulon: coordinated function of the repressor AllR and
the activator AllS. J Mol Biol 324:599-610.
16.   Cusa E, Obradors N, Baldoma L, Badia J, Aguilar J. 1999. Genetic analysis of a
chromosomal region containing genes required for assimilation of allantoin nitrogen
and linked glyoxylate metabolism in Escherichia coli. J Bacteriol 181:7479-84.
17.   Figueira R, Brown DR, Ferreira D, Eldridge MJ, Burchell L, Pan Z, Helaine S,
Wigneshweraraj S. 2015. Adaptation to sustained nitrogen starvation by Escherichia
coli requires the eukaryote-like serine/threonine kinase YeaG. Sci Rep 5:17524.
18.   Levin-Reisman I, Gefen O, Fridman O, Ronin I, Shwa D, Sheftel H, Balaban NQ.
2010. Automated imaging with ScanLag reveals previously undetectable bacterial
growth phenotypes. Nat Methods 7:737-9.
19.   Powell EO. 1956. Growth rate and generation time of bacteria, with special reference
to continuous culture. J Gen Microbiol 15:492-511.

Figure 1

Figure 2

Figure 3

518
519
520

Figure 4

521
522

523
524
525
526 **Figure 4 – cont.**