**The adaptive response to long-term nitrogen starvation in *Escherichia coli* requires**

**the breakdown of allantoin**

Amy Switzer, Lynn Burchell, Josh McQuail and Sivaramesh Wigneshweraraj[1]*

[1]MRC Centre for Molecular Bacteriology and Infection, Imperial College London,

London, SW7 2AZ, UK

*Corresponding author: E-mail: s.r.wig@imperial.ac.uk; Tel.: +44 207 594 1867.

**ABSTRACT**

Bacteria initially respond to nutrient starvation by eliciting large-scale transcriptional

changes. The accompanying changes in gene expression and metabolism allow the bacterial

cells to effectively adapt to the nutrient starved state. How the transcriptome subsequently

changes as nutrient starvation ensues is not well understood. We used nitrogen (N) starvation

as a model nutrient starvation condition to study the transcriptional changes in *Escherichia*

*coli* experiencing long-term N starvation. The results reveal that the transcriptome of N

starved *E. coli* undergoes changes that are required to maximise chances of viability and to

effectively recover growth when N starvation conditions become alleviated. We further

reveal that, over time, N starved *E. coli* cells rely on the degradation of allantoin for optimal

growth recovery when N becomes replenished. This study provides insights into the

temporally coordinated adaptive responses that occur in *E. coli* experiencing sustained N

starvation.

**IMPORTANCE**

Bacteria in their natural environments seldom encounter conditions that support continuous

growth. Hence, many bacteria spend the majority of their time in states of little or no growth

due to starvation of essential nutrients. To cope with prolonged periods of nutrient starvation,

bacteria have evolved several strategies, primarily manifesting themselves through changes in

how the information in their genes is accessed. How these coping strategies change over time

under nutrient starvation is not well understood and this knowledge is not only important to

broaden our understanding of bacterial cell function, but also to potentially find ways to

manage harmful bacteria. This study provides insights into how nitrogen starved *Escherichia*

*coli* bacteria rely on different genes during long term nitrogen starvation.

**KEYWORDS**

*Escherichia coli*, nitrogen regulation, bacterial stress response, transcriptome, allantoin

**INTRODUCTION**

The process of RNA synthesis, transcription, is often the first decision-making and regulatory

point in the flow of genetic information and is catalysed by the multisubunit enzyme, RNA

polymerase (RNAP) (1). The transcriptional responses that establish and maintain specific

cellular states are dynamic and are often controlled by many transcription regulatory factors

that influence the activity of the RNAP (1). Conditions that sustain constant bacterial growth

are seldom found in nature. Whether in a host or the environment, bacteria frequently

encounter nutrient deprivation, which can attenuate bacterial growth (2-4). Hence, the growth

attenuated state is now widely acknowledged as an important physiological state in bacterial

cell function. However, although the transcriptional basis of bacterial growth is widely

studied, our understanding of the transcriptome of growth attenuated bacteria remains in its

infancy. In fact, several recent studies on different species of bacteria have revealed that

growth attenuated bacteria are transcriptionally active. For example: Gefen et al showed that

a batch culture of *E. coli* cells retain transcriptional activity in the stationary phase of growth

over a period of nearly 2 days (5); Stapels et al showed that non-growing intracellular

*Salmonella* in host macrophages are transcriptionally active and this activity is important to

shape the host environment to benefit bacterial survival (6); similarly, non-replicating

*Mycobacterium tuberculosis* cells retained transcriptional activity in the lungs of mice (7);

and more recently, Gray et al reported that *Bacillus subtilis* cells under oligotrophic

conditions are transcriptionally active and replicate at an extremely slow rate (8). Therefore,

understanding the transcriptional landscape in growth attenuated bacteria is important as this

knowledge could inform and inspire new antibacterial strategies, such as disrupting genetic

networks and products that are activated during growth attenuation, which subsequently

become essential for survival and/or optimal growth recovery.

In this study, we used nitrogen (N) starvation as a model condition to understand how

the transcriptional landscape in nutrient starvation-induced growth attenuated *E. coli* affects

bacterial survival and ability to recover growth when nutrients (i.e. N) become replenished.

N is an integral nutrient to bacterial cell function since N is an essential element of most

macromolecules in the bacterial cell (proteins, metabolites, nucleic acids and cell wall

components). Therefore, an exponentially growing batch culture of *E. coli* rapidly attenuates

growth upon N run-out (9-11). The initial adaptive response to N run-out, the N regulation

(Ntr) stress response, results in the large scale and multilevel reprogramming of the

transcriptome, involving ~40% of all *E. coli* genes, to allow *E. coli* to adjust to N starvation

(12-14). Since *E. coli* perceive N run-out as a decrease in the internal concentration of

glutamine, the genes expressed as part of the Ntr response are predominantly involved in the

transport and assimilation of nitrogenous compounds into glutamine or glutamate, yielding

these catabolically or sparing the requirement for them in cellular processes (12, 13).

Although our understanding of the initial adaptive transcriptional response to N starvation is

well established, how the transcriptome changes during long-term N starvation is far from

clear.

**RESULTS AND DISCUSSION**

**Inhibition of transcription in N starved *E. coli* compromises survival and growth**

**recovery.** To setup an experimental system to study how the transcriptome changes during

long-term N starvation, we grew a batch culture of *E. coli* in a defined minimal growth

medium in the presence of a limiting amount (3 mM) of ammonium chloride as the sole N

source and excess (22 mM) glucose as the carbon (C) source. As shown in Fig. 1A (left  and

middle panels), under these conditions, when ammonium chloride becomes depleted (but

sufficient glucose is still available to support growth), bacteria attenuate growth due to N

starvation (12). From the onset of N starvation (referred to as N-), the number of cells in the

population of growth attenuated *E. coli*, as measured by the number of colony forming units

(c.f.u.) in the population, increases at a slow rate over the initial 24 h under N starvation

(doubling time ~39 h, which is ~80-fold less than during exponential growth when N is

abundant; referred to as N+) (Fig. 1A, right panel, orange area). This growth seems to occur

despite the absence of any detectable ammonium chloride in the growth medium (Fig. 1A,

left and middle panels). However, as N starvation ensues beyond 24 h (referred to as N-24),

the population ceases growth before the number of viable cells gradually begins to decline

after 48 h under N starvation (Fig. 1A, red area). Therefore, the first 24 h period following N-

, which we refer to as the N starved adaptive phase (NSAP), seems to be an important

adaptive period during long-term N starvation. It is likely that transcriptional changes occur

during NSAP and we sought to determine how the inhibition of transcription during NSAP

affects cell survival and growth recovery when N starvation stress is alleviated. To do this,

we treated the bacteria with 150 µg/ml of the bacteriostatic antibiotic rifamycin (Rif) at ~1 h

following N- for ~23 h. As previously shown by us, under these conditions, 150 µg/ml Rif

prevents transcription elongation by the RNAP by trapping it at promoters (12). We initially

compared the proportion of viable cells in the Rif treated and untreated population of bacteria

at N-24 (i.e. ~23 h following Rif treatment at N-1). As shown in Fig. 1B, at N-24, only ~50%

cells in the Rif treated culture were viable compared to the Rif untreated culture. Rif

treatment also had a profound effect on the ability of bacteria from N-24 to resume growth

compared to untreated bacteria. As shown in Fig. 1C and 1D, the lag time to growth recovery

of the Rif treated bacteria (following c.f.u. correction to ensure an equal number of viable

treated and untreated bacteria were used; based on data in Fig. 1B) from N-24 was delayed by

~4.8 h compared to untreated bacteria. We considered whether this difference in lag time

between Rif treated and untreated bacteria was due to the presence of a slower growing

subpopulation(s) of bacteria in the Rif treated population that resumed growth slowly. To

investigate this, we used a method called ScanLag (15), which allows measurement of

appearance time of individual colonies on N replete solid growth media as a function of time

following inoculation. We expected that if the Rif treated population consisted of one or more

subpopulations compared to the untreated population, then the appearance of individual

colonies of bacteria treated with Rif would be more heterogenous than that of untreated

bacteria. As shown in Fig. 1E, no heterogeneity in growth in the populations of Rif treated

and untreated bacteria from N-24 was detected. However, the ScanLag data substantiated the

data in Fig. 1C and 1D and demonstrated that Rif treated cells have an increased lag time to

growth recovery compared to untreated cells. Once growth resumed, the doubling time of Rif

treated, and untreated bacteria did not markedly differ (Fig. 1D). Overall, we conclude that

inhibition of transcription at the onset of NSAP has a detrimental effect on the ability of *E.*

*coli* to adapt to long-term N starvation in terms of maximising chances of viability and

optimal growth recovery when N becomes available.

**The transcriptome *E. coli* undergoes changes during the NSAP.** To study the

transcriptional changes that occur during the NSAP, we compared, as indicated in Fig. 2A,

the transcriptomes of *E. coli* during exponential growth (N+), following initial N run-out (N-

), and at N-24. We defined differentially expressed genes as those with expression levels

changed ≥ 2-fold with a false discovery rate adjusted *P*-value < 0.05. As previously reported

by us and others, the initial transcriptional response to N starvation involves differential

expression of 1787 genes primarily required for scavenging and catabolism of alternative N

sources (13, 14, 16) and thus, we did not investigate these genes further (Phase A in Fig. 2A).

However, during the NSAP, upon transition from N- to N-24 (Phase B in Fig. 2A), 451 genes

(~10% of *E. coli* genes) are differentially expressed, of which 105 genes (~2% of *E. coli*

genes) become differentially expressed in Phase B, that were not differentially expressed in

Phase A (Fig. 2B). Functional categorisation of the genes uniquely differentially expressed in

Phase B, hereafter called NSAP genes, revealed that they are involved in diverse processes

including transcription, amino acid transport, metabolism and several genes of unknown

function (Fig. 2C). Eleven (*abgR*, *bfd*, *ileY*, *leuO*, *ompN*, *sibE*, *tauA*, *ycaD*, *yedL*, *yhdV*, and

*yihO*) of the upregulated and nine (*evgA*, *glnA*, *gltI*, *hemH*, *ivy*, *pfkB*, *ybjX*, *yieP*, and *yjeJ*) of

the downregulated NSAP genes were known to be regulated by either NtrC (*glnA* and *gltI*) or

Nac (all other genes listed above) (16, 17) - the master transcription regulators that activate

the initial adaptive response to N starvation. However, the absence of either *glnG* (which

encodes for NtrC) or *nac* did not detectably affect viability of bacteria following 24 h under

N starvation or the ability of 24 h N starved bacteria to recover growth when inoculated into

fresh N replete minimal growth media (Fig. 2D). Overall, we conclude that the transcriptome

of *E. coli* undergoes changes during the NSAP, and this is consistent with the finding above

that 150 μg/ml of Rif treatment during the NSAP has a detrimental effect on the viability of

the *E. coli* population following 24 h under N starvation and its ability to recover growth

when N becomes replenished (Fig. 1B-D) . It therefore seems that the adaptive transcriptional

changes that occur during the NSAP happen subsequently to the initial adaptive changes to N

starvation and do not depend on NtrC (and Nac), which coordinates the initial adaptive

transcriptional response to N run-out.

**Four upregulated NSAP genes are important for optimal growth recovery from**

**long-term N starvation.** We next investigated how the upregulated NSAP genes affected

viability and growth recovery of long-term N starved *E. coli*. Of the 51 upregulated NSAP

genes, 44 are available in the Keio single gene knockout collection (18). Thus, we determined

how the absence of any of the upregulated NSAP genes affected growth recovery following

24 under N starvation. To do this, we grew the library of upregulated NSAP gene mutant

bacteria in minimal growth media containing excess (10 mM) ammonium chloride as the sole

N source and measured cell density after 16 h of growth. The growth of the library of

upregulated NSAP gene mutant strains was done in a plate reader incubator which allowed

continuous monitoring of growth of each mutant strain simultaneously to allow better side-

by-side comparison. The results shown in Fig. 3A indicate that absence of any of the NSAP

genes did not markedly affect growth under N replete conditions. However, we cannot

exclude the possibility that the relative dynamics of growth of the individual mutant strains

could have differed from that of the wild-type (WT) strain over the 16 h period of time. We

note that bacteria in which the tryptophan biosynthesis genes are deleted, lack the γ subunit

of formate dehydrogenase-N (*fdnI*), or ferric rhodotorulic acid outer membrane transporter

(*fhuE*), are unable to grow in our minimal growth media, most likely due to auxotrophy (19).

Next, we reinoculated bacteria that grew under N replete conditions into fresh minimal

growth media containing limiting, i.e. 3 mM, ammonium chloride and determined how the

absence of the upregulated NSAP genes affected viability at N-24, i.e. following 24 h under

N starvation. The results in Fig. 3B show that the viabilities of the upregulated NSAP gene

mutants were similar to that of the WT strain. When the 24 h N starved mutant bacteria were

inoculated into fresh N replete minimal growth media, the dynamics of growth recovery for

the majority of the mutant strains were similar to that of the WT strain (Fig. 3C). However,

Δ*allB*, Δ*leuO*, Δ*ynfP*, and Δ*ybcJ* mutant strains displayed noticeably compromised growth

recovery properties compared to that of the WT strain (Fig. 3C). These genes encode an

allantoinase (*allB*), a DNA-binding transcription dual regulator (*leuO*), a small, possibly

fragmented, protein of unknown function (*ynfP*), and a putative RNA-binding protein (*ybcJ*)

(19). As shown in Fig. 3D, although the doubling time of Δ*allB*, Δ*leuO*, Δ*ynfP*, and Δ*ybcJ*

mutant strains were similar to that of the WT strain, their lag time to growth recovery was

increased by ~1.5-4 h compared to that of the WT strain. Thus, it seems that the transcription

of *allB*, *leuO*, *ynfP*, and *ybcJ* during the NSAP are required for optimal growth recovery from

long-term N starvation. As the Δ*allB* mutant strain displayed the most pronounced

compromised ability to recover growth when compared to the WT strain (Fig. 3C), we

decided to focus on the role of *allB* during the NSAP and recovery from long-term N

starvation.

**Degradation of allantoin during the NSAP is required for optimal growth**

**recovery from long-term N starvation.** To discount the possibility that the difference in lag

time during growth recovery between the WT and Δ*allB* bacteria was not due to the presence

of a subpopulation of slow growing Δ*allB* bacteria, we measured the lag time distribution of

appearance of individual colonies during growth recovery using ScanLag (15) as described in

Fig. 1E. As shown in Fig. 4A, we observed the expected lag time in growth recovery for the

Δ*allB* bacteria, but no evidence of a slow growing subpopulation was seen. Further, we did

not detect a lag in growth recovery of WT and Δ*allB* bacteria sampled from N- when

reinoculated into N replete fresh growth medium (Fig. 4B). This result underscores the view

that *allB* has a specific role in *E. coli* during the NSAP. To establish that the increased lag

time in growth recovery of Δ*allB* bacteria was directly due to the absence of *allB*, we

complemented the Δ*allB* bacteria with plasmid-borne *allB*. As shown in Fig. 4C and 4D, the

lag time in growth recovery of Δ*allB* bacteria reverted to that of WT bacteria when *allB* was

supplied via a plasmid. The product of *allB* is allantoinase: a tetrameric enzyme that catalyses

the hydrolytic conversion of (*S*)-(+)-allantoin to allantoate. A previous study (20) reported

that substitutions at invariant amino acid positions N94 and S317 (in both cases to D and V),

which are located proximal to the active centre, rendered allantoinase inactive (Fig. 4E). As

shown in Fig. 4C and 4D, in the context of the complementation experiments, plasmid-borne

*allB*-N94V and *allB*-S317V failed to fully decrease the lag time in growth recovery of Δ*allB*

bacteria as seen with the WT version of *allB*.  In contrast, plasmid-borne *allB*-N94D and to a

larger extent *allB*-S317D reverted the lag time in growth recovery of Δ*allB* bacteria as seen

with the WT version of *allB*. Although we are unable to explain why specifically substitution

by D but not V at N94 or S317 renders allantoinase catalytically inactive under our

experimental conditions, it seems that the degradation of allantoin occurs at some point

during the NSAP in *E. coli*. This is conceivable because, as shown in Fig. 4F, the conversion

of allantoate (the product of allantoinase) to (*S*)-ureidoglycine by allantoate amidohydrolase

(AllC) and the subsequent conversion of (*S*)-ureidoglycine to (*S*)-ureidoglycolate by (*S*)-

ureidoglycine aminohydrolase (AllE) produces ammonium, which may be needed for

maintenance of cellular functions required for optimal growth recovery from long-term N

starvation. Intriguingly, compared to *allB*, transcripts corresponding to *allC* and *allE* were

only moderately upregulated (~1.2-1.5 fold) in phases A and B in Fig. 2A and thus did not

fall into the category of NSAP genes. However, we observed that 24 h N starved Δ*allC* and

Δ*allE* bacteria displayed a similarly increased lag time to growth recovery as the Δ*allB*

bacteria (Fig. 4G and 4H) when grown in fresh N replete growth media. This suggests that

the activity of the genes of the allantoin degradation pathway have a role in the adaptive

response to long-term N starvation in *E. coli*. As allantoin is a product of the purine

degradation pathway in N starved *E. coli* (21), we considered whether the requirement for

*allB* is a consequence of purine degradation in *E. coli* during the NSAP to sustain cellular

functions via the allantoin degradation pathway (as in Fig. 4F). To indirectly determine

whether purines are degraded in *E. coli* during the NSAP, we measured the integrity of total

RNA as a surrogate for purine degradation in *E. coli* as a function of time in the NSAP by

chip-based capillary electrophoresis (see Materials and Methods). As shown in Fig. 4I,

strikingly, we observed a gradual degradation of 16S and 23S ribosomal RNA as the NSAP

ensued. This observation further underscores the view that the RNA landscape and turnover

during the NSAP is plastic and consistent with the view that the transcriptome of *E. coli*

undergoes changes during long-term N starvation. Collectively, the results support the view

that *allB* contributes to optimal growth recovery from long-term N starvation by the

degradation of allantoin during the NSAP in *E. coli*, which likely accumulates as a result of

purine degradation.

**CONCLUSIONS**

N is central to all metabolic process in the bacterial cell. As such, bacteria can utilize a wide

range of N containing compounds as sole sources of cellular N. Upon N run-out, *E. coli* cells

activate the transcription of genes associated with N assimilatory metabolic pathways

necessary for utilization of N from the extracellular environment and catabolic biosynthetic

pathways for intracellular assimilation of N. We have presented evidence that the growth of a

population of N starved *E. coli* does not completely cease upon N run-out and that *E. coli*

cells continue to divide over a period of up to 24 h – the NSAP - following entry into N

starvation. It seems that the NSAP represents an important adaptive phase in *E. coli*

experiencing long-term N starvation to maximise chances of viability and optimise growth

recovery should N become replenished. We have also presented evidence that the adaptive

transcriptional response to N starvation in *E. coli* occurs in at least two distinct temporally

coordinated phases, where the NtrC dependent initial response to N starvation is followed by

– to a large extent – a NtrC independent transcriptional response. We have uncovered that an

aspect of the adaptive transcriptional response, which occurs in the NSAP subsequent to the

initial adaptive transcriptional response in *E. coli* under long-term N starvation, involves the

degradation of allantoin. As ammonium is a by-product of degradation of allantoin, we

propose that allantoin degradation allows the assimilation of ammonium into glutamine and

glutamate to maintain cellular processes during the NSAP to maximise chances of viability

and optimise growth recovery when N becomes replenished. Previous studies have shown

that allantoin can *only* be used as the sole N source by *E. coli* grown under anaerobic

conditions (22, 23), and consistent with these results, we were unable to grow Δ*allB* and WT

bacteria under our conditions when ammonium chloride (the sole N source used in our

experiments) was replaced by allantoin - in both, the initial growth and recovery growth

experiments (data not shown). Therefore, as the end-product of allantoin degradation under

aerobic conditions is 3-phosphoglycerate, in addition to the by-product ammonium, we

propose that the breakdown of allantoin, which likely occurs as a consequence of purine

catabolism during the NSAP in *E. coli* under long-term N starvation, serves to maintain some

sort of 'metabolic threshold' to allow optimal growth recovery of *E. coli* from long-term N

starvation under aerobic conditions when N becomes replenished. In support of this view, the

current study and previous work by us (12, 14) and others (13) has also shown that genes

associated with the degradation of pyrimidine (e.g. the *rut* operon) become upregulated at the

onset of N starvation and remain upregulated throughout the NSAP (14) Thus, collectively, it

seems that the degradation of nucleotides is an essential component of the adaptive response

to N starvation in *E. coli*. Although future work will focus on deciphering the metabolic and

nucleotide landscape to unravel in depth the adaptive responses that define the NSAP in *E.*

*coli*, this work has underscored the importance of studying the transcriptional changes that

occur in long-term nutrient starved bacteria to further our understanding of bacterial stress

physiology.

**MATERIALS AND METHODS**

**Bacterial strains and plasmids, growth conditions and viability measurements.**
WT and mutant *E. coli* BW25113 strains were used in this study (18). Bacteria were grown
as described in Figueira et al (24) in 10 mM $NH_4Cl$ (for overnight cultures) or 3 mM $NH_4Cl$
(for N starvation experiments) Gutnick minimal medium, consisting of 33.8 mM $KH_2PO_4$ ,
77.5 mM $K_2HPO_4$ , 5.74 mM $K_2SO_4$ , 0.41 mM $MgSO_4$ , supplemented with Ho-LE trace
elements and 0.4 % (w/v) glucose, using $NH_4Cl$ as the sole N source (16). Where relevant,
rifampicin was added at a final concentration of 150 μg/ml at N-1 or N+ as stated, and L-
arabinose was added to complementation strains containing the pBAD18 plasmid at 0.2%
(w/v) for induction of gene expression in growth cultures. Ampicillin was added at a final
concentration of 100 μg/ml to all cultures containing derivatives of the pBAD18 plasmid.
Measurement of growth was taken by measuring the optical density ($OD_{600nm}$) of bacterial
cultures over time to generate growth curves. Ammonium and glucose concentrations in the
media were determined using the Aquaquant ammonium quantification kit and glucose assay
kit, respectively, (both by Merck Millipore, UK) according to the manufacturer's instructions.
To determine population viability, the number of viable cells was calculated by conducting
serial dilutions of cultures at designated time points as shown in relevant figures, grown
overnight on Luria-Bertani (LB) agar plates, and measuring colony forming units (c.f.u.) per
ml. The pBAD18-*allB* plasmid was made by traditional restriction enzyme cloning to
incorporate the ribosomal binding site of pET33b(+) with the *allB* gene to allow arabinose-
inducible expression of the AllB protein. The N94D, N94V, S317D and S317V mutations in
*allB* were done by standard site-directed mutagenesis using pBAD18-*allB* as the template.

**ScanLag**. Bacteria were grown in 3 mM Gutnick minimal medium to N-24, washed
twice in 1 ml PBS, and diluted in PBS. One hundred microlitres of diluted culture was spread
on an LB agar plate and incubated at 33 °C for 48 h with pictures of plates taken every 20

minutes. Analysis of colony growth rate and appearance time was modified from Levin-
Reisman et al (25) by Dr Miles Priestman and is available at
https://github.com/mountainpenguin/NQBMatlab.

**RNA-sequencing (RNA-seq).** The RNA-seq data set was previously generated and
analysed by Switzer et al (14). The analysis of differential gene expression during NSAP as
well as the statistical analyses were also conducted in the same way as described previously.
The sequencing data can be accessed under accession number E-MTAB-6491 in the
ArrayExpress database at EMBL-EBI (www.ebi.ack.uk/arrayexpress).

**Total RNA analysis.** Cultures were grown as above and sampled at the following
time points: at the onset of N starvation (N-) and 3 h (N-3), 6 h (N-6), 9 h (N-9), 12 h (N-12),
18 h (N-18), and 24 h (N-24) under N starvation. At least three biological replicates were
taken at each time point. RNA was acquired and stabilized at specified time points using
Qiagen RNA Protect reagent (Qiagen, 76526). RNA extraction was carried out using the
PureLink RNA mini kit (Invitrogen, 12183025) and DNase Set (Invitrogen, 12185010) as per
manufacturer's protocol. Electropherograms of total RNA integrity were generated using an
Agilent Bioanalyser 2100 instrument (Agilent, G2939BA) and RNA 6000 Nano kit (Agilent,
5067-1511) as per the manufacturer's protocol.

**Statistical analysis and growth calculations.** Unless otherwise stated in the text, all
growth curves and c.f.u. data show the mean average of three independent experiments and
variation calculated by the standard error of the mean (SEM). The student's t-test or one-way
ANOVA was used to determine statistical significance if the probability ($P$) value was $< 0.05$
($* P < 0.05$, $** P < 0.01$, $*** P < 0.001$). Lag and doubling times were calculated by first
generating a straight-line graph of the exponential growth section of each growth curve. Lag
time was defined as the intersection of the straight line graph with the x axis, while the
doubling time was determined during exponential growth (26).

**FIGURE LEGENDS**

**Fig 1. Inhibition of transcription in N starved *E. coli* compromises survival and growth**

**recovery.** (A) Graph showing a representative growth curve (measured by $OD_{600nm}$ readings)

of *E. coli* and concentration (mM) of ammonium chloride in the growth medium as a function

of growth (left panel). The time points referred to as N+, N- and N-24 in the text are

indicated on the growth curve. The bar graph shows the concentration (mM) of ammonium

chloride and glucose at N+, N- and N-24 (middle panel). Graph showing viability, measured

by counting colony forming units (c.f.u.) of *E. coli* over 5 days under N starvation (right

panel). Inset table shows doubling time between different growth periods and the NSAP is

indicated.  (B) Bar  graph showing the proportion of viable cells (measured by c.f.u.) in the

population of *E. coli* at N-1 (black), following 24 h under N starvation with (red) and without

(blue) Rif treatment at N-1. (C) Graph showing recovery growth in fresh N replete medium

as measured by $OD_{600nm}$ readings. The time points and conditions from which bacteria were

sampled for recovery growth experiments are indicated and discussed in the text.  (D) Graph

showing lag time and doubling time during growth recovery experiments from (C). (E)

ScanLag analysis of colony growth rate and appearance time after plating of 24 h N starved

bacteria on LB agar after Rif treatment (red) and no Rif treatment (blue) at N-1. The various

shades of the same colour represent distinct biological replicates, where n=3. In (A)-(D),

error bars represent SEM, where n=3 and the schematics describe the different experiments

for ease of understanding.

**Fig 2. The transcriptome *E. coli* undergoes changes during the NSAP.** (A) Graph

showing a representative growth curve (measured by $OD_{600nm}$ readings) of *E. coli* and

concentration (mM) of ammonium chloride in the growth medium as a function of growth

(left panel). The points at which samples were taken for RNA-seq analysis (N+, N- and N-24)

and the phases referred to as A and B in the text are indicated. Graphs showing the $\log_2$ fold

change in gene expression at N- compared to N+ (Phase A), and N-24 compared to N- (Phase

B). Genes in blue (Phase A) or red (Phase B = NSAP) represent significant differential

expression above a $\log_2$ fold change of 1 and a false discovery rate adjusted $P$ value < 0.05

(right panel). Genes in black (Phase B) represent the NSAP genes, defined as having

significant differential gene expression unique to Phase B (NSAP), but not Phase A. (B)

Venn diagrams showing genes significantly upregulated (top) and downregulated (bottom) in

Phase A compared to Phase B. (C) Graph showing Clusters of Orthologous Groups (COG)

categorisation of NSAP genes, and the number of genes within each category whose

expression is up- or downregulated from N-. (D) Graph showing viability as measured by

c.f.u. counting of viable cells in the population of WT (blue), Δ*glnG* (red) and Δ*nac* (green)

bacteria over 48 h of N starvation (left panel).  Graph showing growth recovery of N-24 WT

(blue), Δ*glnG* (red) and Δ*nac* (green) bacteria in N replete fresh media as measured by

$OD_{600nm}$ readings (right panel). In both graphs, error bars represent SEM, where n=3.

**Fig 3. Four upregulated NSAP genes are important for optimal growth recovery from**

**long-term N starvation.** (A) Graph showing overnight growth of NSAP gene mutant

bacteria as measured by $OD_{600nm}$ compared to WT (blue) bacteria in minimal growth media

containing excess (10 mM) N (ammonium chloride). (B) Graph showing viability as

measured by c.f.u. counts of wild-type (blue) and NSAP gene mutant bacteria (black)

bacteria at N-24. (C) Graph showing recovery growth of 24 h N starved NSAP gene mutant

bacteria in N replete fresh media as measured by $OD_{600nm}$ readings. The recovery growth

curves of WT (blue), Δ*allB* (red), Δ*leuO* (dark green), Δ*ynfP* (purple), and Δ*ybcJ* (orange)

are highlighted. (D) Graph showing lag time (blue) and doubling time (red) of WT, $\Delta allB$,

$\Delta leuO$, $\Delta ynfP$, and $\Delta ybcJ$ bacteria extracted from the growth curves in (C). In (C) and (D),

the error bars represent SEM, where $n=2$. Statistical analyses were performed using the

Student's $t$-test ($*P < 0.05$, $**P < 0.01$).

**Fig. 4. Degradation of allantoin during the NSAP is required for optimal growth**

**recovery from long-term N starvation.** (A) ScanLag analysis of colony growth rate and

appearance time after plating of 24 h N starved WT (blue) and $\Delta allB$ (red) bacteria on LB

agar. Various shades of the same colour represent distinct biological replicates, where n=3.

(B) Graph showing the recovery growth curves of WT (blue) and $\Delta allB$ (red) bacteria from

the onset of N starvation (N-) in fresh N replete media as measured by $OD_{600nm}$ readings of

the cultures. (C) Graph showing the recovery growth curves of 24 h N starved WT +

pBAD18 (blue), $\Delta allB$ + pBAD18 (red), and $\Delta allB$ + pBAD18-$allB$ (green) bacteria in fresh

N replete media as measured by $OD_{600nm}$ readings of the cultures. (D) Graph showing lag

time to growth recovery of WT + pBAD18 (blue), $\Delta allB$ + pBAD18 (red), $\Delta allB$ + pBAD18-

$allB$ (bright green), $\Delta allB$ + pBAD18-$allB$ N94D, $\Delta allB$ + pBAD18-$allB$ N94V,  $\Delta allB$ +

pBAD18-$allB$ S317D and $\Delta allB$ + pBAD18-$allB$ S317V in fresh N replete media as extracted

by $OD_{600nm}$ readings of the cultures. (E) Cartoon representation of the structure of a monomer

of AllB in which the positions of the N94 and S317 are indicated in orange relative to the two

Fe moieties at the active site of the enzyme (indicated in red). (F) A schematic showing the

allantoin degradation pathway indicating the steps where ammonia is produced. (G) As in

(C), but the experiment was conducted with $\Delta allC$ and $\Delta allE$ bacteria. (H) Graph showing lag

time (blue) and doubling time (red) of WT, $\Delta allB$, $\Delta allC$ and $\Delta allE$ bacteria extracted from

the growth curves in (G). (I) Electropherogram of total RNA from bacterial cultures taken at

indicated timepoints. Graph shows representative data from at least three biological

replicates. 16S and 23S rRNA peaks are indicated. In (B), (C), (D), (G) and (H) error bars

represent SEM, where n=3.

**ACKNOWLEDGMENTS**

A Wellcome Trust Investigator award WT100958MA and a Leverhulme Trust project grant

(RPG-2017-431) funded this work. J. M. is an MRC funded PhD student.

**REFERENCES**

1.   Browning DF, Butala M, Busby SJW. 2019. Bacterial Transcription Factors:
Regulation by Pick "N" Mix. J Mol Biol 431:4067-4077.
2.   Peterson CN, Mandel MJ, Silhavy TJ. 2005. Escherichia coli starvation diets:
essential nutrients weigh in distinctly. J Bacteriol 187:7549-53.
3.   Zhang YJ, Rubin EJ. 2013. Feast or famine: the host-pathogen battle over amino
acids. Cell Microbiol 15:1079-87.
4.   Navarro Llorens JM, Tormo A, Martinez-Garcia E. 2010. Stationary phase in gram-
negative bacteria. FEMS Microbiol Rev 34:476-95.
5.   Gefen O, Fridman O, Ronin I, Balaban NQ. 2014. Direct observation of single
stationary-phase bacteria reveals a surprisingly long period of constant protein
production activity. Proc Natl Acad Sci U S A 111:556-61.
6.   Stapels DAC, Hill PWS, Westermann AJ, Fisher RA, Thurston TL, Saliba AE,
Blommestein I, Vogel J, Helaine S. 2018. Salmonella persisters undermine host
immune defenses during antibiotic treatment. Science 362:1156-1160.
7.   Manina G, Dhar N, McKinney JD. 2015. Stress and host immunity amplify
Mycobacterium tuberculosis phenotypic heterogeneity and induce nongrowing
metabolically active forms. Cell Host Microbe 17:32-46.
8.   Gray DA, Dugar G, Gamba P, Strahl H, Jonker MJ, Hamoen LW. 2019. Extreme
slow growth as alternative strategy to survive deep starvation in bacteria. Nat
Commun 10:890.
9.   Schumacher J, Behrends V, Pan Z, Brown DR, Heydenreich F, Lewis MR, Bennett
MH, Razzaghi B, Komorowski M, Barahona M, Stumpf MP, Wigneshweraraj S,
Bundy JG, Buck M. 2013. Nitrogen and carbon status are integrated at the
transcriptional level by the nitrogen regulator NtrC in vivo. mBio 4:e00881-13.
10.  Brauer MJ, Yuan J, Bennett BD, Lu W, Kimball E, Botstein D, Rabinowitz JD. 2006.
Conservation of the metabolomic response to starvation across two divergent
microbes. Proc Natl Acad Sci U S A 103:19302-7.
11.  Reitzer L. 2003. Nitrogen assimilation and global regulation in Escherichia coli. Annu
Rev Microbiol 57:155-76.
12.  Brown DR, Barton G, Pan Z, Buck M, Wigneshweraraj S. 2014. Nitrogen stress
response and stringent response are coupled in Escherichia coli. Nat Commun 5:4115.
13.  Zimmer DP, Soupene E, Lee HL, Wendisch VF, Khodursky AB, Peter BJ, Bender
RA, Kustu S. 2000. Nitrogen regulatory protein C-controlled genes of Escherichia

| 467 | | coli: scavenging as a defense against nitrogen limitation. Proc Natl Acad Sci U S A |
| 468 | | 97:14674-9. |
| 469 | 14. | Switzer A, Evangelopoulos D, Figueira R, de Carvalho LPS, Brown DR, |
| 470 | | Wigneshweraraj S. 2018. A novel regulatory factor affecting the transcription of |
| 471 | | methionine biosynthesis genes in Escherichia coli experiencing sustained nitrogen |
| 472 | | starvation. Microbiology 164:1457-1470. |
| 473 | 15. | Levin-Reisman I, Fridman O, Balaban NQ. 2014. ScanLag: high-throughput |
| 474 | | quantification of colony growth and lag time. J Vis Exp doi:10.3791/51456. |
| 475 | 16. | Brown DR, Barton G, Pan Z, Buck M, Wigneshweraraj S. 2014. Combinatorial stress |
| 476 | | responses: direct coupling of two major stress responses in Escherichia coli. Microb |
| 477 | | Cell 1:315-317. |
| 478 | 17. | Aquino P, Honda B, Jaini S, Lyubetskaya A, Hosur K, Chiu JG, Ekladious I, Hu D, |
| 479 | | Jin L, Sayeg MK, Stettner AI, Wang J, Wong BG, Wong WS, Alexander SL, Ba C, |
| 480 | | Bensussen SI, Bernstein DB, Braff D, Cha S, Cheng DI, Cho JH, Chou K, Chuang J, |
| 481 | | Gastler DE, Grasso DJ, Greifenberger JS, Guo C, Hawes AK, Israni DV, Jain SR, |
| 482 | | Kim J, Lei J, Li H, Li D, Li Q, Mancuso CP, Mao N, Masud SF, Meisel CL, Mi J, |
| 483 | | Nykyforchyn CS, Park M, Peterson HM, Ramirez AK, Reynolds DS, Rim NG, Saffie |
| 484 | | JC, Su H, Su WR, et al. 2017. Coordinated regulation of acid resistance in Escherichia |
| 485 | | coli. BMC Syst Biol 11:1. |
| 486 | 18. | Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita |
| 487 | | M, Wanner BL, Mori H. 2006. Construction of Escherichia coli K-12 in-frame, |
| 488 | | single-gene knockout mutants: the Keio collection. Mol Syst Biol 2:2006 0008. |
| 489 | 19. | Keseler IM, Mackie A, Santos-Zavaleta A, Billington R, Bonavides-Martinez C, |
| 490 | | Caspi R, Fulcher C, Gama-Castro S, Kothari A, Krummenacker M, Latendresse M, |
| 491 | | Muniz-Rascado L, Ong Q, Paley S, Peralta-Gil M, Subhraveti P, Velazquez-Ramirez |
| 492 | | DA, Weaver D, Collado-Vides J, Paulsen I, Karp PD. 2017. The EcoCyc database: |
| 493 | | reflecting new knowledge about Escherichia coli K-12. Nucleic Acids Res 45:D543- |
| 494 | | D550. |
| 495 | 20. | Kim K, Kim MI, Chung J, Ahn JH, Rhee S. 2009. Crystal structure of metal- |
| 496 | | dependent allantoinase from Escherichia coli. J Mol Biol 387:1067-74. |
| 497 | 21. | Xi H, Schneider BL, Reitzer L. 2000. Purine catabolism in Escherichia coli and |
| 498 | | function of xanthine dehydrogenase in purine salvage. J Bacteriol 182:5332-41. |
| 499 | 22. | Rintoul MR, Cusa E, Baldoma L, Badia J, Reitzer L, Aguilar J. 2002. Regulation of |
| 500 | | the Escherichia coli allantoin regulon: coordinated function of the repressor AllR and |
| 501 | | the activator AllS. J Mol Biol 324:599-610. |
| 502 | 23. | Cusa E, Obradors N, Baldoma L, Badia J, Aguilar J. 1999. Genetic analysis of a |
| 503 | | chromosomal region containing genes required for assimilation of allantoin nitrogen |
| 504 | | and linked glyoxylate metabolism in Escherichia coli. J Bacteriol 181:7479-84. |
| 505 | 24. | Figueira R, Brown DR, Ferreira D, Eldridge MJ, Burchell L, Pan Z, Helaine S, |
| 506 | | Wigneshweraraj S. 2015. Adaptation to sustained nitrogen starvation by Escherichia |
| 507 | | coli requires the eukaryote-like serine/threonine kinase YeaG. Sci Rep 5:17524. |
| 508 | 25. | Levin-Reisman I, Gefen O, Fridman O, Ronin I, Shwa D, Sheftel H, Balaban NQ. |
| 509 | | 2010. Automated imaging with ScanLag reveals previously undetectable bacterial |
| 510 | | growth phenotypes. Nat Methods 7:737-9. |
| 511 | 26. | Powell EO. 1956. Growth rate and generation time of bacteria, with special reference |
| 512 | | to continuous culture. J Gen Microbiol 15:492-511. |
| 513 | | |

**Figure 1**

**Figure 2**

**Figure 3**

**A**

OD$_{600nm}$

- wild-type
- Δ*allB*

Appearance Time (h)

**B**

0.6

OD$_{600nm}$

— WT
— Δ*allB*

Time (h)

**C**

0.6

OD$_{600nm}$

Time (h)

— wild-type + pBAD18
— Δ*allB* + pBAD18
— Δ*allB*+ pBAD18-*allB*

**D**

Lag time (h)

wild-type +pBAD18
Δ*allB*+ pBAD18
Δ*allB*+ pBAD18-*allB*
Δ*allB*+ pBAD18-*allB-N94D*
Δ*allB*+ pBAD18-*allB-N94V*
Δ*allB*+ pBAD18-*allB-S317D*
Δ*allB*+ pBAD18-*allB-S317V*

**E**

N

C

Ser317

Asn94

**F**

(*S*)-(+)-allantoin

$H_2O$
$H^+$    allantoinase: AllB

allantoate

$H_2O + 2H^+$
$CO_2 + NH_4^+$    allantoate amidohydrolase: AllC

(*S*)-ureidoglycine

$H_2O$
$NH_4^+$    (*S*)-ureidoglycine aminohydrolase: AllE

(*S*)-ureidoglycolate

**G**

0.8

OD$_{600nm}$

— Wild-type
— Δ*allB*
— Δ*allC*
— Δ*allE*

Time (h)

**H**

Time (h)

■ Lag time
■ Doubling time

Wild-type   Δ*allB*   Δ*allC*   Δ*allE*

**I**

Flourescence units

16S
23S

N-
N-3
N-6
N-9
N-12
N-18
N-24

Fragment size (nucleotides)

**Figure 4**