1 **S100A8/A9 modulates inflammatory collateral tissue damage during**

2 **intraperitoneal origin systemic candidiasis**

<u>Running title:</u>

Calprotectin modulates collateral tissue damage

<u>Authors:</u>

Madhu Shankar[*1,4], Nathalie Uwamahoro[*1], Sandra Holmberg[2], Maria Joanna Niemiec[1,5],

Johannes Roth[3], Thomas Vogl[3], Constantin F. Urban[1#]

[*] Equally contributing first authors

[#] Corresponding author

<u>Affiliations of authors:</u>

[1]Department of Clinical Microbiology, Umeå Centre for Microbial Research (UCMR) &

Molecular Infection Medicine Sweden (MIMS), Umeå University, Umeå 90185, Sweden.

[2]Department of Medical Chemistry and Biophysics, Umeå University, Umeå 90187, Sweden.

[3]Institute of Immunology, Universitätsklinikum Münster & University of Münster, 48149

Münster, Germany

<u>Current addresses:</u>

[4] Department of Biochemistry & Molecular Biology, Monash Biomedicine Discovery Institute,

Melbourne, Australia

[5] Leibniz Institute for Natural Product Research and Infection Biology, Hans Knöll Institute

(HKI) & Center for Sepsis Control and Care, Jena, Germany

Full postal addresses of corresponding author:

Dr. Constantin F. Urban, Universitetssjukhuset by 6M, Department of Clinical Microbiology,

Immunology, Plan 4, Umeå universitet, 901 85 Umeå, Sweden, Phone: +46 90 785 27 02, Fax:

+46 90 12 99 05, Email: constantin.urban@umu.se

Keywords:

Sepsis, host-pathogen interactions, peritonitis, *Candida albicans*, inflammation

Abstract

Peritonitis is a leading cause of severe sepsis in surgical intensive care units, as over 70% of

patients diagnosed with peritonitis develop septic shock. A critical role of the immune system

is to return to homeostasis after combating infection. S100A8/A9 (calprotectin) is an

antimicrobial, pro-inflammatory protein complex often used as a biomarker for diagnosis of

disease activities in many inflammatory disorders. Here we describe the role of S100A8/A9 on

inflammatory collateral tissue damage (ICTD).

We performed an *in vivo Candida albicans* disseminated peritonitis mouse model using WT

and S100A9-deficient mice and stimulated primary macrophages with recombinant S100A8/A9

in the presence or absence of the compound paquinimod, a specific inhibitor of S100A9. In

addition, the effects on ICTD and fungal clearance were investigated. S100A9-deficient mice

developed less ICTD than wildtype mice. Restoration of S100A8/A9 in S100A9 knockout mice

resulted in increased ICTD and fungal clearance comparable to wildtype levels. Treatment with

paquinimod abolished ICTD.

The data indicated that S100A8/A9 controls ICTD levels and host antimicrobial modulation at

a systemic level during intra-abdominal candidiasis (IAC).

Introduction

Peritonitis frequently results in severe sepsis in surgical patients, especially in the intensive care unit (ICU) (1), as more than 70% of patients often succumb to death within 72h (2). Hence, treatment options are required that extend life expectancy to allow proper treatment (3). Peritonitis is characterized by inflammation of the membrane in the abdominal cavity, the peritoneum. Peritonitis often occurs upon disruption of physical barriers or can become spontaneous in severe organ failure, causing alterations of the physiologic flora residing in the gastrointestinal tract (1,4). These alterations prompt an inflammatory response that targets the removal of contaminants from the peritoneal cavity into circulation. There, pathogens induce activation of host immune responses and the release of pro-inflammatory mediators including Interleukin-6 (IL6), macrophage inflammatory protein -1α (MIP-1α) and tumor necrosis factor −α (TNFα) to recruit phagocytes to the peritoneum (5). The attempt to restrict infection promotes abscess development through production of fibrinous exudate. Failure to confine peritonitis may lead to organ failure, coma and death (6).

The inflammatory response of the innate immune system can either have intentional (pathogen clearance) or collateral outcomes (pathological side effect, called inflammatory collateral tissue damage -ICTD) (7). The focus of most studies on detection and elimination of pathogens has neglected host tolerance to disease through the resolution of collateral outcomes of inflammation in an attempt to establish homeostasis (8).

Acute or systemic inflammatory pathological states are often associated with intra-abdominal surgery (9). Surgical intervention disrupting the natural barriers of the gastrointestinal tract leads to deep-seated microbial infection by gut colonizing organisms and frequently *Candida albicans* infection. As a result, the most common non-mucosal fungal diseases among hospitalized patients are *Candida* peritonitis, also referred to as intra-abdominal candidiasis

(IAC). IAC is often challenging to diagnose and hence results in high mortality rates ranging

from 25% - 60% (9).

The resolution of infection is an active process involving reprogramming of cells and

modulation of mediators (10) such as the S100A8/A9 heterodimeric complex (calprotectin) also

known as alarmin or DAMP (11–13). Here, we used an experimental model for IAC to study

the contribution of S100A8/A9 to ICTD.

Participating as a signaling molecule that binds to Toll-like receptor-4 (TLR4) to induce pro-

inflammatory chemokines and cytokines, the heterodimer is the physiologically relevant and

active form that is secreted by activated, stressed or necrotic cells. The prolonged presence of

S100A8/A9, however, leads to a tolerated state of the immune system as a counter mechanism

or stress tolerance (12). Under high calcium conditions, such as present in the extracellular

milieu and culture medium, S100A8/A9 heterodimers inactivate itself by tetramer formation,

to restrict their activity locally and to avoid overwhelming immune reactions (13). Thus,

S100A8 homodimers, which cannot tetramerise, are widely accepted experimental stimuli to

mimic heterodimer activity.

Furthermore, S100A8/A9 binds micronutrients including zinc, manganese, and calcium, and is

often deployed by leukocytes as a mechanism to either deprive microbes of the nutrients or

poison the microbes in high quantities (14). Neutrophils release the heterodimer during the

formation of neutrophil extracellular traps to bind or capture *C. albicans* (15). In addition to the

antimicrobial benefit, S100A8/A9 is used in diagnostics to monitor neutrophil elevation in

various inflammatory diseases (16). The q-compound paquinimod, an immunomodulatory

compound that prevents the binding of S100A9 to TLR4, was designed to target chronic

inflammatory S100A8/A9 dependent diseases (17–22). Using a murine peritonitis model with

C. albicans, we show here that the inflammatory response failed to contain the pathogen in the

peritoneum but instead led to detrimental ICTD dependent on the presence of S100A8/A9.

Treatment of mice with paquinimod abrogated S100A8/A9-induced ICTD suggesting

paquinimod as promising adjunct therapy option during severe IAC.

Materials and methods

Ethical statement

98   Animal experiments and isolation of cells were carried out following the recommendations in

99   the Guide for the Care and Use of Laboratory Animals, conformed to Swedish animal protection

100   laws and applicable guidelines (djurskyddslagen 1988:534; djurskyddsförordningen 1988:539;

101   djurskyddsmyndigheten DFS 2004:4) in a protocol approved by the local Ethical Committee

102   (Umeå djurförsöksetiska nämnd, Permit number A12-13, A80-14 and A79-14.

Statistical analyses

Statistical analysis was conducted using Graphpad Prism 6 software and P values less than 0.5

were considered significant. All two-group comparisons in *Candida* CFU data, inflammatory

score, ALT data, and ELISA data were conducted using the unpaired, two-tailed student's t-

test. Comparisons of WT and *S100A9$^{-/-}$* mice data were performed using One-way ANOVA

multiple comparisons analysis as specified in figure legends. In all comparisons, the sample

size is specified in figure legends, and a $P<0.05$ was considered significant. *p<0.05; **p<0.01;

***p<0.001; ****p<0.0001.

Yeast strains and growth conditions

112   *C. albicans* clinical isolate strain - SC5314 was cultured overnight in YPD (1% yeast extract,

113   2% bacto-peptone and 2% glucose) at 30°C. The *Candida* cells were washed three times in PBS

114   prior use in all assays. Cell numbers were calculated using Vi-CELL Cell Viability Analyzer

115   (Beckman Coulter AB).

Animal infections and isolation of bone marrow-derived macrophages, and tissue analyses

All mice were maintained according to a previous report (23) at Umeå Centre for Comparative Biology (UCCB), Umeå University, Umeå, Sweden. If not otherwise stated mice were infected intraperitoneally with 3×10^6 *C. albicans* cells per g mouse from an overnight culture in YPD. For intravenous infection (only Fig. S4) mice were challenged intravenously with 2.5×10^3 *C. albicans* cells per g mouse.

Primary macrophages (BMDMs) cells and differentiated as described in a previous report (24). BMDMs flow cell cytometry (FACs) was conducted using BD LSR II flow cytometer (BD Biosciences, San Jose, CA), using propidium iodide (24), on paquinimod treated cells at indicated concentrations.

For tissue damage and fungal load, analyses were conducted according to a previous report (25), with 3×10^6 *C. albicans* cells intraperitoneal injection per g of mouse after 24 hours of infection (26). Blood ALT levels were measured using vetscan VS2^{TM} (SCIL animal care company) as previously reported (27–29).

Histological preparations and inflammatory score analyses were conducted as in previous reports (15,30). For the inflammatory score, Whole sections were analyzed for inflammation and scored under the supervision of a specialized animal pathologist. The sections from each animal were scored as zero if they had no inflammatory cells present in the tissue, one for a few inflammatory cells (1–20 cells), two for moderate cell infiltration (21–40 cells), three for a large number of inflammatory cells (41–60 cells), and four if inflammation was spread all over in the tissue (61 cells) (30).

Cytokine and chemokine quantification

BMDMs were seeded at 1×10^5 cells per well in 96-well microplates and then infected with *C. albicans* at an MOI of 1 for 24 hours. Cell-free culture supernatants were harvested after *C.*

albicans infection. Supernatants were analyzed for indicated cytokines of chemokines by

ELISA (Biolegend-ELISA MAX™ 9727 Pacific Heights Blvd, San Diego, CA, USA), or Pro-

Mouse cytokine BioPlex® 200 multiplex (Bio-Rad Laboratories) according to the

manufacturer's instructions.

Generation of recombinant S100A8

The S100A8 and S100A9 gene encoding for monomers of the mouse dimer S100A8/A9

(UniProt P27005, P31725) was synthesized, codon harmonized and purchased from DNA2.0.

The gene were GST tagged and cloned into *E. coli* BL21 strain. The cloned cells were auto-

induced overnight. The cells were pelleted and suspended in 10 mg/ml of 1×PBS supplemented

with DNase and protease inhibitors. The cells were lysed on ice by sonication (Branson Digital

sonifier; 10 mm horn, 50% power) for 6 min with alternating 10 s pulses and pauses. The lysate

was clarified by centrifugation at 23,000g for 20 min at 4˚C. The clarified lysate was filtered

through a 0.45 µm syringe filter and batch bound to 2.0 ml GST sepharose (~25mg protein/ml)

for 2h at 4˚C gravity flow over column and washed with 30 column volumes of PBS. Samples

were eluted in 4-6 ml fractions with 50 mM Tris pH 8 and 10 mM glutathione. For each fraction,

the resin was incubated with elution buffer for 10 min prior to collecting the flow through. The

fractions were analyzed at A280. The pooled fractions of protein were cleaved (1:100) using

protease 3c (PreScission®protease) to remove the GST tag and the fractions of GST tag free

protein were pooled. The protein was then separated by size-exclusion chromatography (SEC)

using a Superdex 75 16/600 column (GE Healthcare Life Sciences, UK) equilibrated with PBS

pH 7, at a flow rate of 0.5 ml/min. The fractions containing purified S100A8 (rA8) and S100A9

(rA9) were concentrated using an Amicon Ultra-4 centrifugal filter device with a 3 kDa

molecular-weight cutoff (Millipore). The primary sequence, the intact mass and the presence

of product were confirmed by mass spectrometry using an ABI 4800 MALDI tandem time-of-

flight mass spectrometer. Recombinant proteins were screened for endotoxin contamination and

levels were below 0.6 pg per µg protein, as previously recommended (31).

Results

A disseminated fungal peritonitis model to determine roles of S100A8/A9

During IAC, fungal cells dissemination reach the liver and other organs via lymphatics or

bloodstream (6). Liver tissue damage and leukocytosis are hallmarks of deep-seated and

systemic *C. albicans* infection (32). To establish a fungal IAC model and determine the role of

S100A8/A9 in systemic inflammation, we used *S100A9*-deficient mice (*S100A9$^{-/-}$*) that also

lack the S100A8 protein on the protein level despite normal S100A8 RNA level and thus

actually represent a functional S100A8/A9 double knockout mouse strain (33). Normally

expressed S100A8 protein in S100A9$^{-/-}$ mice is rapidly degraded in the proteasome, avoiding

systemic overwhelming immune responses. However, under chronic TNF conditions, this

degradation process is insufficient leading to severe phenotypes in artificial mouse strains (TTP^{-}

$^{/-}$/S100A9$^{-/-}$ or itghTNF/S100A9$^{-/-}$) indicating that heterodimer activity must be regulated

tightly to restrict operating range (13).

Using wildtype (WT) and *S100A9$^{-/-}$* mice, *C. albicans* intraperitoneal injections were

administered, liver tissues collected after 24 hours of infection and indicators of sepsis were

measured. The infection was determined to be systemic using phenotypic analysis of mice and

histopathological analysis from hematoxylin-eosin stained (H & E stain) liver sections after 24

hours. The development of eye exudates was an indication of fungal dissemination from the

peritoneal cavity to other loci in the mouse (Fig. 1A). WT liver sections indicated higher

numbers of leukocyte infiltration zones compared to *S100A9$^{-/-}$* sections (Fig. 1A, top panel).

Conversely, there was visual evidence of more fungal cells in the *S100A9$^{-/-}$* livers compared to

WT (Fig. 1A, bottom panel). The inflammatory score (Fig. 1B) showed a large number of

inflammatory cell infiltrates (level 3) in WT *C. albicans* infected samples compared to lack of

inflammation (level 1) observed in *S100A9^-/-* mice.

Blood alanine aminotransferase (ALT) levels in the blood are an indicator of liver damage

(1,29). *C. albicans* infected WT mice (101.6U/L) showed elevated levels compared to *S100A9^-*

^/- mice (37.2U/L) which showed similar ALT levels to uninfected mice, suggesting lack of

systemic tissue damage in the S100A8/A9-deficient animals (Fig. 2A). The ability of the host

to clear the infection or organ microbial load is related to the colony-forming units (CFUs) (34).

S100A9^-/- mice had a significantly higher fungal load (2.5 fold of average levels) compared to

WT mice (Fig. 2B).

Macrophages are sentinels for immune signaling that leads to leukocytosis but may cause

problems when uncontrolled (35). The pro-inflammatory cytokine response of WT and *S100A9^-*

^/- was determined using bone marrow-derived macrophages (BMDMs) to monitor the ability of

primary macrophages to induce TNFα (Fig. 2C). Higher levels of TNFα (3.1 fold of average

levels measured) upon *C. albicans* infection were induced by WT BMDMs (570 pg/ml)

compared to *S100A9^-/-* BMDMs (185 pg/ml), suggesting a defect in TNFα induction in cells

lacking S100A8/A9.

S100A8/A9 enhances pro-inflammatory cytokine release of macrophages

The implications of the use of S100A8/A9 in recombinant protein therapy is unknown. To

determine whether the S100A8 activity would aid the *S100A9^-/-* mutant in eliciting an

appropriate cytokine response *in vitro*, BMDMs derived from S100A9-/- mice were infected

with *C. albicans* and treated with rS100A8. Murine *S100A8* was expressed in *E. coli,* purified

(Fig. S1) and verified by mass spectrometry to obtain functionally active homodimers as a

substitute for the heterodimer as previously reported (36). We included analyses for cytokines

typically released during early inflammatory responses (Fig. 3A-F).

Considerable increase of TNFα levels was obtained in $S100A9^{-/-}$ BMDMs treated with

rS100A8 (Fig. 3A) comparable to WT levels upon *C. albicans* infection (Fig. 2C), and IL-6

was also strongly enhanced (Fig. 3B). In addition, $S100A9^{-/-}$ BMDMs infected with *C. albicans*

and treated with rS100A8 released higher levels of chemokines *MIP1α* (*CCL3*) and *MIP1β*

(*CCL4*), crucial for the recruitment of various leukocyte subpopulations (Fig. 3C and 3D) as

compared to infected and untreated control macrophages. Also monitored was the induction of

the anti-inflammatory cytokine IL10, which showed low but significant levels induced when

mice were treated with rS100A8, while CXCl-1 levels declined (Fig. 3E and 3F). Induction of

pro-inflammatory cytokines and modulation of tissue damage by rS100A8 suggest that

S100A8/A9 heterodimers are main contributors of the acute pro-inflammatory response during

C. albicans infection.

Paquinimod reduces cytokine release of C. albicans-infected macrophages

To potentially reduce ICTD during IAC pharmacological intervention blocking the pro-

inflammatory activity of S100A8/A9 could be used. Hence. the activity of paquinimod, a novel

anti-inflammatory compound initially developed against Systemic Lupus Erythematosus (SLE)

that targets S100A9, one subunit of the S100A8/A9 complex, in liver, lung, heart and skin (37),

was tested *in vitro*. The percentage of dead cells (propidium iodide positive cells) using flow

cytometry analysis (FACS) was assessed on treatment of WT BMDMs with various drug

concentrations of paquinimod. Among the concentrations used, no significant toxic effects were

observed (Fig. S2). However, there was significant TNFα reduction at 300 µg/ml and 930 µg/ml

paquinimod concentrations compared to untreated *C. albicans* infected BMDMs (Fig. 4A).

CCL-3 and IL-10 were also reduced upon addition of 930 µg/ml paquinimod confirming the

activity for other cytokines and chemokines. Thus, paquinimod is likely able to block

S100A8/A9-mediated cytokine release during *C. albicans* infection.

S100A8/A9 is responsible for ICTD during disseminated peritonitis

The active role of S100A8/A9 as alarmin during IAC is currently unknown. To test the usefulness of recombinant protein therapy, we determined the effects of recombinant S100A8 protein (rS100A8) on disseminated *C. albicans* infection. Purified rS100A8 protein (100μl of a 100 μg/ml solution) was injected into *C. albicans* infected WT and *S100A9*$^{-/-}$ mice. Treatment of *S100A9*$^{-/-}$ mice with rS100A8 showed a higher level of ALT (5.9 fold of average levels measured) compared to untreated with levels not significantly different to WT suggesting that constitutively active S100A8 homodimers mimic heterodimers to cause WT levels of liver damage (Fig. 5A) and that the S100A8/A9-mediated effect on ICTD is direct rather than indirect. S100A8/A9 is an antimicrobial protein against *C. albicans* (15). In this context, the fungal clearance defect observed in *C. albicans* infected *S100A9*$^{-/-}$ mice was remedied, reducing fungal load after treatment with rS100A8 (Fig. 5B).

Paquinimod therapy reduces ICTD induced by disseminated peritonitis

Targeted deletion of *S100A9* improves survival in mouse models of bacterial-induced sepsis (11,38). Higher lethality of a disseminated infection is often due to the inability to contain inflammation-induced organ and tissue damage by the host (1). There are no specific immunotherapies against disseminated infections, such as sepsis, and often management focuses on containing the infection through source control and antibiotics or antifungals plus organ function support (39). We hypothesized that using an anti-inflammatory drug in WT mice may mimic the beneficial aspect (lack of tissue damage) observed in the *S100A9*$^{-/-}$ mutant might extend host survival from disseminated IAC. Treatment of intraperitoneally infected mice with paquinimod led to complete elimination of ICTD as indicated by reduced ALT plasma levels in infected mice to levels of uninfected mice (Fig. 5C). As expected, paquinimod treatment did not have an effect on ALT levels in infected *S100A9*$^{-/-}$ mice (Fig. 5C). In addition, paquinimod treatment affected fungal burden in the liver of infected WT and *S100A9*$^{-/-}$ mice only moderately

(Fig. 5D), probably due to reduced S100A8/A9-dependent activation of fungi-eliminating

immune cells. Of note, the CFU count in livers of *S100A9^{-/-}* mice was significantly higher than

in livers of WT mice, suggesting that ICTD after 24 h of infection is not strictly correlated to

fungal burden, but depends in this setting to a significant proportion on S100A8/A9 activity.

Paquinimod and other quinoline-3-carboxamides have been described as specific binders of

S100A9 (40). As rS100A9 does not form functional homodimers *in vitro* and hence could not

lead to a reduction of ICTD in experimental *C. albicans* peritonitis (Fig. S3), we used rS100A8.

Interestingly, the effect of rS100A8 injection on ALT plasma levels in infected *S100A9^{-/-}* mice

could be reverted by treatment with paquinimod (Fig. 5E), but expectedly had no effect on

fungal burden in the liver (Fig. 5F). It is possible that paquinimod binds recombinant mouse

S100A8 with somewhat higher affinity than recombinant human S100A8 which possibly could

explain the described effect.

Paquinimod has a moderate effect on survival in disseminated peritonitis

To determine whether paquinimod could be used to alleviate sepsis derived from fungal

peritonitis, infected wildtype mice were treated with paquinimod every 24 hours with

paquinimod (Fig. 6A). The treatment resulted in a moderately increased survival rate after five

days post-infection with a statistically significant effect using log-rank test (Fig. 6B). Although

the compound did not prevent all mice from succumbing to systemic infection, between 24 and

48 hours treated mice survived longer compared to untreated mice (Fig. 6B). Despite the similar

weight loss in infected mice observed between treated and untreated groups (Fig. 6C), treated

mice were generally more active during *C. albicans* infection. Hence, our data suggest that anti-

S100A8/A9 therapy could be a useful strategy to increase the therapeutic window for antifungal

treatment during IAC, but probably could not serve as a standalone therapeutic approach.

Discussion

This work characterizes the role of immune-modulating S100A8/A9 on host resolution of inflammation from a peritoneal-derived disseminated *C. albicans* infection. The gastrointestinal commensal nature of *C. albicans* requires that mucosal damage and neutropenia are achieved for *C. albicans* dissemination (41). Most experimental models studying systemic fungal diseases use intravenous (IV) injection of fungal cells which bypass mucosal host defenses and establish an infection predominantly in the kidney and in the brain (42). This study utilized an IAC model to induce systemic inflammation mimicking a severe clinical concern of postoperative *Candida* peritonitis (43). Similar to IV, IP-induced infection allows rapid blood dissemination of pathogens with exposure to an active population of phagocytes, complement cascade and the potential for abscess formation in the peritoneal cavity (43). The peritonitis infection model presented here aimed to breach the immune barriers in the peritoneal cavity, and this was phenotypically clear in the dissemination of *C. albicans* to the liver and to the eyes of infected animals (Fig. 1 and 2). The IP injection route allows pathogen exposure to active phagocyte populations in the peritoneum and the potential for the host to contain the infection through the formation of abscesses (6). In intravenous infection, in contrast to the IP injection route, ALT plasma level did not increase above uninfected control within the first 48 h of infection (Fig. S4) and hence the peritonitis infection model is suitable to describe acute and severe onset of disseminated infection leading to ICTD.

WT liver fungal clearance was ~3 times more efficient when compared to the S100A8/A9-deficient mice (Fig. 2B). The various leucocyte infiltration zones observed in WT liver tissues compared to *S100A9*$^{-/-}$ mice (Fig. 1A) indicated that the recruitment of higher levels of leukocytes coincides with fungal clearance in WT mice. These findings suggested that the presence of S100A8/A9 is required for fungal clearance during disseminated infection, and supports our previous findings that implicated antimicrobial properties of S100A8/A9 during

candidiasis (15). Notably, presence of S100A8/A9 in WT mice resulted to significantly more

liver damage compared to *S100A9*[-/-] mice as indicated by ALT plasma levels despite higher

fungal load in the liver (Fig. 2A and B), implicating that the induction of S100A8/A9

inflammatory responses modulates processes which contribute to tissue damage. These findings

support the association of the S100 family of proteins with inflammatory disorders (13), and

our data indicate that S100A8/A9 activity depends on modulation of both antimicrobial and

inflammatory activity. These insights would especially be critical in the case of

immunocompromised patients.

The IP treatment with rS100A8 of *C. albicans*-infected *S100A9*[-/-] mice resulted in fungal

clearance similar to WT mice. Since rS100A8 showed significant *in vivo* activity, we used

S100A9[-/-] bone marrow-derived macrophages in a *C. albicans ex vivo* infection assay, and

showed that rS100A8 mediates macrophage activation as indicated by the higher levels of the

chemokines CCL-3 and CCL-4, pro-inflammatory cytokines IL-6 and TNFα, as well as the

anti-inflammatory cytokine IL10 (Fig. 3). The induction of IL10 by rS100A8 is consistent with

homeostatic management of systemic inflammation (44). Balancing of pro- and anti-

inflammatory cytokines is a common theme of host factors with local and system-wide effects

as exemplified by interferon signaling pathways during viral infections (45). Albeit

administration of rS100A8 led to enhanced fungal clearance in the liver of infected S100A8/A9-

deficient mice, the activity of S100A8 has the potential to induce dire effects on host-mediated

tissue damage which probably do not warrant recombinant therapy with rS100A8 in

immunocompromised individuals.

Our data demonstrate that ICTD, which was alleviated in S100A8/A9-deficient mice, could be

fully reintroduced by injection of recombinant protein (rS100A8). Thus, the reported effect on

ICTD in an experimental model of IAC is, at least to a large proportion, dependent on

S100A8/A9 and pharmacological targeting of the protein complex could be beneficial in

patients with IAC. To demonstrate this, we used paquinimod; a compound applied for treatment

of chronic inflammatory diseases (17,20). As paquinimod is a specific binder for S100A9 (40),

it blocks the pro-inflammatory activity of the S100A8/A9 complex *in vivo*. Surprisingly, we

found that paquinimod also reverted ICTD in infected S100A9-/- mice previously treated with

rS100A8 (Fig. 5E). Bjork *et al*., showed strong binding of paquinimod to human and mouse

S100A9 and negligible binding to human S100A8 *in vitro*; however, binding to mouse S100A8

was not investigated (40). Notably, it may also be possible that paquinimod exerts other

immune-modulatory effects that give rise to the reduction seen in Fig. 5E. No toxicity was

observed against bone marrow-derived macrophages (BMDMs) at the concentration used in

animal infection experiments and paquinimod activity affected the inflammatory modulators as

observed in the reduction of CCL-3 and IL-10. No adverse effects on fungal load in livers of

infected animals were seen either (Fig. 5D). Hence, paquinimod represents a potential adjuvant

therapeutic option that dampens the host responses to avoid tissue damage and increases the

therapeutic window (Fig. 6B) for targeting the pathogen with antifungal drugs, in particular,

since the compound had no adverse effect on fungal clearance.

This study showed the dilemma of the protective and harmful roles of S100A8/A9 in *Candida*

albicans-induced fungal peritonitis (Fig. 6). Fungal peritonitis that is not contained by host

defenses leads to pathogen infiltration of host organs (Fig. 7.1). Fungal-host interaction induces

the systemic release of S100A8/A9 (15) for fungal clearance and ALTs into the systemic

circulation (Fig. 7.2-7.3). Local innate immune cells (represented in this study as primary

macrophages respond with a cytokine 'storm' of pro-inflammatory cytokines (Fig. 7.4-7.5), that

leads to increased leukocyte infiltrates systemically (Fig. 7.6), and promote increased tissue

damage in the effort of fungal clearance (Fig. 7.7). Hence, our study presents both the potential

for a systemically functional protein, evidence for potential antimicrobial recombinant protein

therapy, and adjuvant treatment as paquinimod-mediated inhibition of S100A8/A9 showed

reduction of ICTD (Fig. 7). Considering the high mortality (>50%) due to do fungal sepsis (1),

future adjuvant therapies similar to paquinimod could be the key against peritonitis and other

severe inflammatory malignancies.

Acknowledgments

The authors thank José Pedro Lopes and Sujan Yellagunda for support with animal experiments. We acknowledge essential advice and the kind gift of paquinimod from Active Biotech. We acknowledge support for heterologous protein expression of the Protein Expertise Platform (PEP) at Umeå University. Furthermore, we are grateful for funding provided to CFU by the Swedish research council VR-M 2014-02281 and VR-M 2017-01681, the Kempe Foundation SMK-1453, the Åke Wiberg Foundation M14-0076 and M15-0108, and the Medical Faculty of Umeå University 316-886-10. NU and MS held a postdoctoral fellowship received in competition from UCMR. The funders had no rule in study design nor analysis and interpretation of the results.

Author Contribution

CFU provided funding and together with MS and NU designed the study, MS and NU collected and analyzed the data. TV and JR contributed to the conceptualization of the study. Manuscript drafting by NU, MS and CFU. SH contributed preparation of histological sections. MJN provided data from intravenous infection. All authors contributed to data analysis, drafting and critically revising the paper, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

Conflict of interest

The authors declare no conflict of interest.

References

1. Fiore M, Maraolo AE, Leone S, Gentile I, Cuomo A, Schiavone V, Bimonte S, Pace MC, Cascella M. Spontaneous peritonitis in critically ill cirrhotic patients: a diagnostic algorithm for clinicians and future perspectives. *Ther Clin Risk Manag* (2017) **13**:1409–1414. doi:10.2147/TCRM.S144262

2. Paoli C, Reynolds M, Sinha M, Gitlin M. 1486: Current burden, Outcomes, and Costs of Sepsis Management in United States hospitals. *Crit Care Med* (2018) **46**:

3. Denning DW, Perlin DS, Muldoon EG, Colombo AL, Chakrabarti A, Richardson MD, Sorrell TC. Delivering on Antimicrobial Resistance Agenda Not Possible without Improving Fungal Diagnostic Capabilities. *Emerg Infect Dis J* (2017) **23**:177. doi:10.3201/eid2302.152042

4. de Jong PR, González-Navajas JM, Jansen NJG. The digestive tract as the origin of systemic inflammation. *Crit Care* (2016) **20**:279. doi:10.1186/s13054-016-1458-3

5. Yao V, Platell C, Hall JC. Role of peritoneal mesothelial cells in peritonitis. *BJS (British J Surgery)* (2003) **90**:1187–1194. doi:10.1002/bjs.4373

6. Skipworth RJE, Fearon KCH. Acute abdomen: peritonitis. *Surg* (2008) **26**:98–101. doi:https://doi.org/10.1016/j.mpsur.2008.01.004

7. Wang A, Huen SC, Luan HH, Yu S, Zhang C, Gallezot J-D, Booth CJ, Medzhitov R. Opposing Effects of Fasting Metabolism on Tissue Tolerance in Bacterial and Viral Inflammation. *Cell* (2016) **166**:1512-1525.e12. doi:10.1016/j.cell.2016.07.026

8. Hamidzadeh K, Christensen SM, Dalby E, Chandrasekaran P, Mosser DM. Macrophages and the Recovery from Acute and Chronic Inflammation. *Annu Rev Physiol* (2017) **79**:567–592. doi:10.1146/annurev-physiol-022516-034348

9. Lichtenstern C, Schmidt J, Knaebel HP, Martin E, Büchler MW, Weigand MA. Postoperative Bacterial/Fungal Infections: A Challenging Problem in Critically Ill Patients after Abdominal Surgery. *Dig Surg* (2007) **24**:1–11. doi:10.1159/000099009

10. Netea MG, Balkwill F, Chonchol M, Cominelli F, Donath MY, Giamarellos-Bourboulis EJ, Golenbock D, Gresnigt MS, Heneka MT, Hoffman HM, et al. A guiding map for inflammation. *Nat Immunol* (2017) **18**:826–831. doi:10.1038/ni.3790

11. Vogl T, Tenbrock K, Ludwig S, Leukert N, Ehrhardt C, van Zoelen MAD, Nacken W, Foell D, van der Poll T, Sorg C, et al. Mrp8 and Mrp14 are endogenous activators of Toll-like receptor 4, promoting lethal, endotoxin-induced shock. *Nat Med* (2007) **13**:1042–1049. doi:10.1038/nm1638

12. Austermann J, Friesenhagen J, Fassl SK, Ortkras T, Burgmann J, Barczyk-Kahlert K, Faist E, Zedler S, Pirr S, Rohde C, et al. Alarmins MRP8 and MRP14 Induce Stress Tolerance in Phagocytes under Sterile Inflammatory Conditions. *Cell Rep* (2014) **9**:2112–2123. doi:https://doi.org/10.1016/j.celrep.2014.11.020

13. Vogl T, Stratis A, Wixler V, Völler T, Thurainayagam S, Jorch SK, Zenker S, Dreiling A, Chakraborty D, Fröhling M, et al. Autoinhibitory regulation of S100A8/S100A9 alarmin activity locally restricts sterile inflammation. *J Clin Invest* (2018) **128**:1852–1866. doi:10.1172/JCI89867

14. Zackular JP, Chazin WJ, Skaar EP. Nutritional Immunity: S100 Proteins at the Host-Pathogen Interface. *J Biol Chem* (2015) **290**:18991–18998. doi:10.1074/jbc.R115.645085

15. Urban CF, Ermert D, Schmid M, Abu-Abed U, Goosmann C, Nacken W, Brinkmann V, Jungblut PR, Zychlinsky A. Neutrophil extracellular traps contain calprotectin, a cytosolic protein complex involved in host defense against Candida albicans. *PLoS Pathog* (2009)

5:e1000639–e1000639. doi:10.1371/journal.ppat.1000639

16.   Chan JK, Roth J, Oppenheim JJ, Tracey KJ, Vogl T, Feldmann M, Horwood N, Nanchahal J.
Alarmins: awaiting a clinical response. *J Clin Invest* (2012) **122**:2711–2719.
doi:10.1172/JCI62423

17.   Fransén Pettersson N, Deronic A, Nilsson J, Hannibal TD, Hansen L, Schmidt-Christensen A,
Ivars F, Holmberg D. The immunomodulatory quinoline-3-carboxamide paquinimod reverses
established fibrosis in a novel mouse model for liver fibrosis. *PLoS One* (2018) **13**:e0203228–
e0203228. doi:10.1371/journal.pone.0203228

18.   Kraakman MJ, Lee MK, Al-Sharea A, Dragoljevic D, Barrett TJ, Montenont E, Basu D,
Heywood S, Kammoun HL, Flynn M, et al. Neutrophil-derived S100 calcium-binding proteins
A8/A9 promote reticulated thrombocytosis and atherogenesis in diabetes. *J Clin Invest* (2017)
127:2133–2147. doi:10.1172/JCI92450

19.   Masouris I, Klein M, Dyckhoff S, Angele B, Pfister HW, Koedel U. Inhibition of DAMP
signaling as an effective adjunctive treatment strategy in pneumococcal meningitis. *J
Neuroinflammation* (2017) **14**:214. doi:10.1186/s12974-017-0989-0

20.   Schelbergen RF, Geven EJ, van den Bosch MHJ, Eriksson H, Leanderson T, Vogl T, Roth J,
van de Loo FAJ, Koenders MI, van der Kraan PM, et al. Prophylactic treatment with S100A9
inhibitor paquinimod reduces pathology in experimental collagenase-induced osteoarthritis.
Ann Rheum Dis (2015) **74**:2254 LP – 2258. doi:10.1136/annrheumdis-2014-206517

21.   van den Bosch MH, Blom AB, Schelbergen RF, Koenders MI, van de Loo FA, van den Berg
WB, Vogl T, Roth J, van der Kraan PM, van Lent PL. Alarmin S100A9 Induces
Proinflammatory and Catabolic Effects Predominantly in the M1 Macrophages of Human
Osteoarthritic Synovium. *J Rheumatol* (2016) **43**:1874 LP – 1884. doi:10.3899/jrheum.160270

22.   Yun J, Xiao T, Zhou L, Beuerman RW, Li J, Zhao Y, Hadayer A, Zhang X, Sun D, Kaplan HJ,
et al. Local S100A8 Levels Correlate With Recurrence of Experimental Autoimmune Uveitis
and Promote Pathogenic T Cell Activity. *Invest Ophthalmol Vis Sci* (2018) **59**:1332–1342.
doi:10.1167/iovs.17-23127

23.   Hobbs JAR, May R, Tanousis K, McNeill E, Mathies M, Gebhardt C, Henderson R, Robinson
MJ, Hogg N. Myeloid cell function in MRP-14 (S100A9) null mice. *Mol Cell Biol* (2003)
23:2564–2576. doi:10.1128/mcb.23.7.2564-2576.2003

24.   Uwamahoro N, Verma-Gaur J, Shen H-H, Qu Y, Lewis R, Lu J, Bambery K, Masters SL,
Vince JE, Naderer T, et al. The Pathogen Candida albicans Hijacks Pyroptosis for Escape from
Macrophages. *MBio* (2014) **5**:e00003-14. doi:10.1128/mBio.00003-14

25.   Lopes JP, Stylianou M, Backman E, Holmberg S, Jass J, Claesson R, Urban CF. Evasion of
Immune Surveillance in Low Oxygen Environments Enhances Candida albicans Virulence.
MBio (2018) **9**:e02120-18. doi:10.1128/mBio.02120-18

26.   Vethanayagam RR, Almyroudis NG, Grimm MJ, Lewandowski DC, Pham CTN, Blackwell
TS, Petraitiene R, Petraitis V, Walsh TJ, Urban CF, et al. Role of NADPH Oxidase versus
Neutrophil Proteases in Antimicrobial Host Defense. *PLoS One* (2011) **6**:e28149. Available at:
https://doi.org/10.1371/journal.pone.0028149

27.   Parasuraman S, Raveendran R, Kesavan R. Blood sample collection in small laboratory
animals. *J Pharmacol Pharmacother* (2010) **1**:87–93. doi:10.4103/0976-500X.72350

28.   Röhm M, Grimm MJ, D'Auria AC, Almyroudis NG, Segal BH, Urban CF. NADPH oxidase
promotes neutrophil extracellular trap formation in pulmonary aspergillosis. *Infect Immun*
(2014) **82**:1766–1777. doi:10.1128/IAI.00096-14

29.   Vatsalya V, Avila D, Frimodig JC, Barve SS, McClain CJ, Gobejishvili L. Liver Injury

| | 475 | | Assessment by Vetscan VS2 Analyzer and Most Frequently Used ALT/GTP Reagent. |
| 476 | | | *Gastroenterol Hepatol (Bartlesville, Okla)* (2016) **4**:107. doi:10.15406/ghoa.2016.04.00107 |

30. Bieghs V, Vlassaks E, Custers A, Van Gorp PJ, Gijbels MJ, Bast A, Bekers O, Zimmermann LJI, Lütjohann D, Voncken J-W, et al. Chorioamnionitis Induced Hepatic Inflammation and Disturbed Lipid Metabolism in Fetal Sheep. *Pediatr Res* (2010) **68**:466–472. doi:10.1203/PDR.0b013e3181f70eeb

31. Schwarz H, Schmittner M, Duschl A, Horejs-Hoeck J. Residual endotoxin contaminations in recombinant proteins are sufficient to activate human CD1c+ dendritic cells. *PLoS One* (2014) **9**:e113840–e113840. doi:10.1371/journal.pone.0113840

32. Fiore M, Cascella M, Bimonte S, Maraolo AE, Gentile I, Schiavone V, Pace MC. Liver fungal infections: an overview of the etiology and epidemiology in patients affected or not affected by oncohematologic malignancies. *Infect Drug Resist* (2018) **11**:177–186. doi:10.2147/IDR.S152473

33. Manitz M-P, Horst B, Seeliger S, Strey A, Skryabin B V, Gunzer M, Frings W, Schönlau F, Roth J, Sorg C, et al. Loss of S100A9 (MRP14) results in reduced interleukin-8-induced CD11b surface expression, a polarized microfilament system, and diminished responsiveness to chemoattractants in vitro. *Mol Cell Biol* (2003) **23**:1034–1043. doi:10.1128/mcb.23.3.1034-1043.2003

34. Bayer AS, Edwards JE, Guze LB. Experimental intraabdominal candidiasis in rabbits: therapy with low-total-dose intravenous amphotericin B. *Antimicrob Agents Chemother* (1981) **19**:179 LP – 184. doi:10.1128/AAC.19.1.179

35. Grainger JR, Konkel JE, Zangerle-Murray T, Shaw TN. Macrophages in gastrointestinal homeostasis and inflammation. *Pflugers Arch* (2017) **469**:527–539. doi:10.1007/s00424-017-1958-2

36. Futami J, Atago Y, Azuma A, Putranto EW, Kinoshita R, Murata H, Sakaguchi M. An efficient method for the preparation of preferentially heterodimerized recombinant S100A8/A9 coexpressed in Escherichia coli. *Biochem Biophys reports* (2016) **6**:94–100. doi:10.1016/j.bbrep.2016.03.009

37. Li X, Zhu L, Wang B, Yuan M, Zhu R. Drugs and Targets in Fibrosis. *Front Pharmacol* (2017) **8**:855. doi:10.3389/fphar.2017.00855

38. Kingsley MK, Bhat BV, Badhe BA, Dhas BB, Parija SC. Narciclasine improves outcome in sepsis among neonatal rats via inhibition of calprotectin and alleviating inflammatory responses. *Sci Rep* (2020) **10**:2947. doi:10.1038/s41598-020-59716-7

39. Rello J, Valenzuela-Sánchez F, Ruiz-Rodriguez M, Moyano S. Sepsis: A Review of Advances in Management. *Adv Ther* (2017) **34**:2393–2411. doi:10.1007/s12325-017-0622-8

40. Björk P, Björk A, Vogl T, Stenström M, Liberg D, Olsson A, Roth J, Ivars F, Leanderson T. Identification of human S100A9 as a novel target for treatment of autoimmune disease via binding to quinoline-3-carboxamides. *PLoS Biol* (2009) **7**:e97–e97. doi:10.1371/journal.pbio.1000097

41. Koh AY, Köhler JR, Coggshall KT, Van Rooijen N, Pier GB. Mucosal Damage and Neutropenia Are Required for Candida albicans Dissemination. *PLOS Pathog* (2008) **4**:e35. Available at: https://doi.org/10.1371/journal.ppat.0040035

42. Hohl TM. Overview of vertebrate animal models of fungal infection. *J Immunol Methods* (2014) **410**:100–112. doi:10.1016/j.jim.2014.03.022

43. Hall JC, Heel KA, Papadimitriou JM, Platell C. The pathobiology of peritonitis. *Gastroenterology* (1998) **114**:185–196. doi:10.1016/S0016-5085(98)70646-8

521   44.   Shouval DS, Ouahed J, Biswas A, Goettel JA, Horwitz BH, Klein C, Muise AM, Snapper SB.
522   Interleukin 10 receptor signaling: master regulator of intestinal mucosal homeostasis in mice
523   and humans. *Adv Immunol* (2014) **122**:177–210. doi:10.1016/B978-0-12-800267-4.00005-5

524   45.   Majer O, Bourgeois C, Zwolanek F, Lassnig C, Kerjaschki D, Mack M, Müller M, Kuchler K.
525   Type I interferons promote fatal immunopathology by regulating inflammatory monocytes and
526   neutrophils during Candida infections. *PLoS Pathog* (2012) **8**:e1002811–e1002811.
527   doi:10.1371/journal.ppat.1002811

Figure Legends

Figure 1. Disseminated *C. albicans* peritonitis mouse model depicting the necessity of

S100A8/A9 in inflammation. Intraperitoneal (IP) *C. albicans* infection (3 x 10^6 cells per g

mouse) in WT or S100A8/A9-deficient mice after 24 hours. **A)** Fungal eye exudate as evidence

for peritoneal cavity failure to contain C. albicans infection after 24. Shown is the clear eye of

uninfected (PBS) compared to the white pus in infected wild type (WT) mouse. **B)**

Representative photomicrographs of hematoxylin-eosin (H&E) stained liver sections of

uninfected and infected wildtype and s100A9 mutant used in quantifications in **C)**. Zoomed

images show one of many observed increased inflammatory cell infiltrates in WT infected cells

(top panel), and arrows indicate yeast or hyphal cells (bottom panel) at 20x magnification. CV:

central vein, PV: portal vein. **C)** An inflammatory score of H&E-stained liver sections of

uninfected and infected WT and S100A9-/- mutant mice. A score of 2 = moderate cell

infiltration, >3 = large number of infiltrates, 4 = full tissue inflammatory infiltration. Data from

10x microscopy image fields of 2 sections from mice n=2 (WT and *S100A9 $^{-/-}$*) and n=6 (WT

+ Ca and *S100A9 $^{-/-}$* + Ca). Error bars indicate standard deviation and ****p< 0.0001

Figure 2. S100A8/A9 is required for inflammatory induced collateral tissue damage

(CTD) and fungal clearance in the liver. **A-B)** WT and S100A9-/- mice were infected with 3

x 10^6 cells per g of mice. A) Alanine transferase levels (ALTs) used as an indicator of liver

tissue damage as a result of *C. albicans* infection. Plasma ALT levels were analyzed from 100µl

of plasma in a vetscan rotor 24 h post-infection. **B)** Organ levels of colony-forming units

(CFUs) as a measure of host fungal burden. *C. albicans* CFUs in homogenized infected livers

of indicated mice were quantified. **A-B)** Box- and whisker plots show the smallest observation,

lower quartile, mean, upper quartile and largest observation. n = 7 (WT and S100A9-/-), and

n= 10 (WT and S100A9-/- + Ca) total mice. **** p-value =<0.0001. **C)** *C. albicans in vitro*

infection of immune cells induces the production of less pro-inflammatory cytokines. WT and

S100A9-/- bone marrow-derived macrophages (BMDMs) were infected with *C. albicans* (MOI

1). TNFα levels were measured from supernatants at 24-hour post-infection. ** p-value =<0.01,

n = 3.

Figure 3. Recombinant S100A8 protein restores pro-inflammatory cytokine release of

S100A9$^{-/-}$ bone marrow-derived macrophages upon *in vitro C. albicans* infection. Cytokine

levels **A)** TNFα, **B)** LI-6, **C)** CCL3, **D)** CCL-4, **E)** IL-10, and **F)** CXCL-1 in supernatants of

bone marrow-derived macrophages (BMDMs) infected with *C. albicans* (MOI 1) were

measured 24-hour post-infection. A Pro-Mouse cytokine BioPlex® 200 multiplex array (Bio-

Rad, Hercules, CA) was used to detect and quantify *S100A9$^{-/-}$* mouse cytokines infected with

C. albicans with and without additional treatment with rS100A8 (10 µg/ml). **A-F)** n=3, *p-

value =<0.05 ****p-value =<0.0001.

Figure 4. Quinoline-3-carboxamide paquinimod (Paq) reduces pro-inflammatory

cytokine release of WT bone marrow-derived macrophages upon *in vitro C. albicans*

infection. A) WT BMDM were infected and treated with different Paq concentrations

(930µg/ml – 12.5 µg/ml). TNFα was measured from supernatants 24 h post-infection. **B-C)** WT

BMDM were infected and treated with 930 µg/ml or 0 µg/ml Paq and release of B) CCL-3 and

C) IL-10 was measured from supernatants after 24 h. **A-C)** n=3. *p-value =<0.05, ****p-value

=<0.0001.

Figure 5. Recombinant S100A8 and paquinimod modify collateral tissue damage in

experimental disseminated *C. albicans* peritonitis. A). Effect of recombinant soluble

rS100A8 on *C. albicans* induced collateral tissue damage. Protein was intraperitoneally

administered (approximately 10 µg per mouse) to WT and *S100A9$^{-/-}$* mice intraperitoneally

infected with *C. albicans* (3 x 10^6 cells per g mouse) and shown are plasma ALT levels. **B)**

Effect of rS100A8 on host fungal clearance. Shown are CFUs from homogenized infected livers

of indicated mice **A-B)** Data are presented as a box- and whisker plots showing the smallest

observation, lower quartile, mean, upper quartile and largest observation, statistical significance

was analyzed by one-way ANOVA n = 5 mice per group in 3 separate experiments, **** *p*-

value =<0.0001. **C-D)** Effects of paquinimod (Paq) on *C. albicans*-induced collateral tissue

damage and fungal clearance in intraperitoneally infected mice. Paq was administered

intraperitoneally (30mg/kg). Shown are plasma ALT levels **C)** and CFUs from homogenized

infected livers **D)** from *C. albicans* infected WT and *S100A9*$^{-/-}$ mice. E-F) Effect of Paq on

recombinant S100A8/A9 induced collateral tissue damage during *C. albicans* infection. Note

that **E)** contains *S100A9*$^{-/-}$ controls initially presented in **A)** as experiments were conducted

together. Shown are plasma ALT levels **E)** and CFUs from homogenized infected livers **F)**

from *C. albicans* infected *S100A9*$^{-/-}$ mice treated with rS100A8 (10µg per mouse) with our

without Paq treatment. Data are presented as a box- and whisker plots showing the smallest

observation, lower quartile, mean, upper quartile and largest observation, statistical significance

was analyzed by student t-test n = 5 mice per group, **p*-value =<0.05, **** *p*-value =<0.0001.

Figure 6. Inhibition of S100A8/A9 by paquinimod moderately increases host survival

from disseminated *C. albicans* peritonitis A-D) WT mice infected with *C. albicans* were

treated with Paq (30mg/kg) at 24-hour intervals up to 5 days, and mice weight and survival

were monitored. **A)** Paq mice treatment strategy. **B** Survival B) and sequential body weight C)

of WT mice infected and treated at the indicated timeline. Survival is moderately but

significantly increased. n = 8 mice per group. For survival log-rank test *p*-value = 0.0169, for

mouse weight grams lost over time is shown. **p-value <0.01. Shown are mean and SD.

Figure 7. Study summary. Using a systemic fungal peritonitis mouse model (**1**), findings

implicate S100A8/A9 as a systemically essential protein (**2-6, i-vi)**) for the control of cytokine

and chemokine modulation (**4-5**), organ leukocyte recruitment (**6**), host tissue damage (**7**) and

fungal clearance (**vi**). In systemic intra-abdominal originating candidiasis, the cost of fungal

clearance (antimicrobial/active S100A8/A9 signaling) a feedback-loop that increases tissue

damage, while the cost of decreased tissue damage (anti-inflammatory/ inactive S100A8/A9

signaling) is a higher fungal burden (Fig. 1-5). Both are lethal if not controlled (Fig. 5B). Future

studies should focus on fine-tuning recombinant therapy in cases where active S100A8/A9

signaling is inadequate, and anti-inflammatory treatments like paquinimod, in conjunction with

antifungal treatments to increase host survival time to aid treatment.

Figure 1 (Shankar et al. 2020)

Figure 2 (Shankar et al. 2020)

Figure 4 (Shankar et al. 2020)

Figure 5 (Shankar et al. 2020)

Figure 7 (Shankar et al. 2020)