A way to break bones? The weight of intuitiveness

D. Vettese[1,2] (Corresponding author), T. Stavrova[1], A. Borel[1,3], J. Marin[1], M.-H. Moncel[1], M. Arzarello[2], C. Daujeard[1].

[1]Histoire Naturelle de l'Homme Préhistorique (HNHP, UMR 7194), Sorbonne Universités, Muséum national d'Histoire naturelle (MNHN), Homme et Environnement, CNRS, Institut de Paléontologie Humaine, Paris, France
[2]Università degli Studi di Ferrara, Dipartimento degli Studi Umanistici, Sezione di Scienze Preistoriche e Antropologiche, Corso Ercole I d'Este 32, 44121 Ferrara, Italy
[3]Institute of Archaeological Sciences, Eötvös Loránd University, Múzeum krt. 4/b, 1088 Budapest, Hungary.

Delphine Vettese (corresponding author): delphine.vettese@mnhn.fr, https://orcid.org/0000-0001-6441-6054

Trajanka Stavrova: stavrovatrajanka@gmail.com

Antony Borel: antony.borel@mnhn.fr

Juan Marin: juan.marin.hernando@gmail.com, http://orcid.org/0000-0002-5698-602X

Marie-Hélène Moncel: marie-helene.moncel@mnhn.fr

Marta Arzarello: marta.arzarello@unife.it

Camille Daujeard: camille.daujeard@mnhn.fr, https://orcid.org/0000-0001-7489-8691

Abstract:

The essential relationship to fat in the Middle Paleolithic, and especially to the yellow marrow, explains the importance of addressing this issue of butchery cultural practices through the study of bone fracturing gestures and techniques. In view of the quasi-systematization of bone marrow extraction in many anthropized archaeological levels, this butchery activity had to be recurrent, standardized and counter-intuitive. Thus, the highlighting of butcher traditions made possible by the analysis of the distribution of percussion marks within fossil assemblages, in particular by opposition at patterns resulting from an intuitive practice. With this in mind, we carried out an experiment that focus at the intuitive way of fracturing bones to extract marrow, involving volunteers with no previous experience in this butchery activity. The objective of this experiment was to highlight the presence or absence of a distribution pattern for percussion marks in an intuitive context by comparing several long bones and individuals.

Thus, we wanted to evaluate the influence of the morphological specificity of the element and the specific characteristics of volunteers on the distribution of percussion marks during marrow extraction. Indeed, a previous study was able to show the possible existence of intuitive patterns of distribution of these traces according to the elements (Stavrova et al. 2019). In addition, the comparison of the different behaviours of volunteers during bone fracturing with the production of remains and marks on bone surfaces highlighted the variables that most influence the creation of an intuitive model. We selected twelve from a larger experiment that resulted in the fracturing of more than 360 long bones. Each of the experimenters broke a series of ten long bones, always the same element. Subsequently, we compared the data collected during the experiment with the data from the laboratory study of the remains. Then, we applied an innovative GIS (Geographic Information System) method to analyze the distribution of percussion marks to highlight recurrent patterns.

One of the most significant results of our article shows the existence of significant concentrations of percussion marks, regardless of the volunteers' behaviour during bone fracturing. The predominance of two factors explains the distribution patterns that emerge from our analysis: for humerus, radio ulnas and tibias, the morphology of each element seems to constrain the location of percussion marks, while for femurs, individual choices have more weight in this distribution. In addition, we have observed that at different levels of our analyses, the bone response to fracturing may be totally opposite (quantity of bone marrow, marrow quality, number of blows, difficulty felt, number of fragments produced, type of marks registered), particularly regarding the results for radio-ulnas and tibias. Subsequently, it would be most interesting to compare the intuitive models that we were able to highlight through spatial analysis

54 with the distributions of percussion marks registered in fossil assemblages. It would thus be possible to

55 propose new hypotheses on butchering practices based on the results presented in this work.

56 **Keywords**: GIS; bone breakage; experimental archaeology; intuitiveness; percussion marks;

57 spatial analyses

 Introduction

Lipids are essential nutrients for the human organism. During the Palaeolithic, fat was even

more important, due to its role in the gluconeogenesis metabolism, synthesizing glucose from non-

carbohydrate precursors (Costamagno and Rigaud, 2014; Holt et al., 2018; Speth, 2010). At that time,

various flora and fauna provided lipid resources but their accessibility was largely influenced by

environmental contexts. Lipids were found in oilseeds, such as hazelnuts. However, cold and dry periods

were characterized by a low vegetal biomass during which animal resources represented the most

important food resources (Benazzi et al., 2015; Binford, 1981; O'Connell et al., 1992; Speth, 2010). The

red marrow present in bone epiphyses and yellow marrow from the diaphysis were both important

resources of fat from animal carcasses. Marrow was even the most widely available fat resource during

the winter (Abe, 2005; Binford, 1978; Brugal and Defleur, 1989; Costamagno and David, 2009; Hardy

et al., 2006; Kuntz et al., 2016; Speth and Spielman, 1983). Indeed, the recovery of yellow marrow was

quasi-systematic among Palaeolithic hunter-gatherers, and particularly for Neanderthals (Noe-Nygaard,

1977; Patou-Mathis, 1985; Speth, 1989; Speth et al., 2012). Prehistorians noted widespread evidence of

marrow extraction through long bone breakage at Middle Palaeolithic sites (e.g.,: Blasco and Fernández

Peris, 2012; Daujeard et al., 2017, 2012; Joana et al., 2014; Marín et al., 2017; Romandini et al., 2014;

Valensi et al., 2013).

The traces left on bones by marrow recovery are mainly percussion marks. For several decades,

percussion marks and long bone breakage methods have been extensively studied (Boulestin, 1999;

Capaldo and Blumenschine, 1994; Fisher, 1995; Gifford-Gonzalez, 2018; Patou-Mathis, 1985; White,

1992. Since the beginning of the twentieth century and up until now, archaeological experiments on

long bone breakage have been carried out to characterize percussion marks and their location (e.g.,:

(Blasco et al., 2014; Blumenschine and Selvaggio, 1988; Brugal and Defleur, 1989; Galán et al., 2009;

Martin, 1910; Moclán and Domínguez-Rodrigo, 2018; Noe-Nygaard, 1977; Patou-Mathis, 1985;

Pickering and Egeland, 2006). Fracture patterns have also been extensively studied on different states

of bone preservation: fresh, dry, weathered, fossilized but also heated, frozen, boiled (Alhaique, 1997;

Karr and Outram, 2012; Lacroix, 2011; Outram, 2002; Villa and Mahieu, 1991). The authors based their

experimental protocols on ethnographic works studying current populations of hunter-gatherers (e.g.,:

Abe, 2005; Binford, 1981; Costamagno and David, 2009; Enloe, 1993; Oliver, 1993; ; Stiner et al., 2011;

Yravedra et al., 2017). More recently, the use of new methodological approaches (e.g., geometric

morphometrics, 3D modelling or GIS analyses of distribution along the diaphysis…) presents new

challenges for the experimental study of marrow recovery.

Some recent studies have focused on the distribution of percussion marks on long bones to

approach the subsistence behaviours of past hominins (Blasco et al., 2013; Masset et al., 2016; Moclán

and Domínguez-Rodrigo, 2018; Vettese et al., 2017). One of these studies highlighted for the first time

a pattern of non-random bone breakage and interpreted it as a product of butchery traditions among

Neanderthals (Blasco et al., 2013). Through comparisons with an experiment performed by non-trained

volunteers, the authors established that this systematic pattern differed from intuitive patterns. In

addition, Moclan and colleagues (2018) proved that the morphology of the skeletal element from which

marrow is extracted could influence the distribution of percussion marks.

We conducted a large-scale experiment to test how non-trained individuals recover marrow.

Indeed, in order to identify butchery traditions in Palaeolithic sites, it is necessary to differentiate know-

how from intuitiveness. The definition of a butchery tradition is a systematic and counterintuitive pattern

shared by a same group. The immediate apprehension of the non-trained butcher to break a bone is

influenced by numerous variables, including anatomical constraints (Blasco et al., 2013; Moclán and

Domínguez-Rodrigo, 2018). Individuals who regularly break long bones acquire an empirical approach

and develop specific skills that enhance efficiency (Pickering and Egeland, 2006). Hence, their skills

include habits and preferences gained by experience and/or group traditions. This know-how cannot be

assessed without differentiating between physical bone features and socio-cultural practices.

Thus, our aim is to experimentally test whether the inter-individual specificity of long bone

morphology may have some influence on the distribution of percussion marks. Based on (GIS) spatial

analysis of percussion marks on the bone surface, we also test the existence of a preferential pattern

regarding the intuitive breakage of long bones. Moreover, we intend to verify whether non-experienced

volunteers develop their own method to improve efficiency, such as an auto-learning process. Finally,

it should be possible to grasp the influence of each bone structure by comparing the elements between

them and by assessing the performance of individuals on the same types of bones. This allows for the

comparison of behaviours and the influence of behaviour on the production of bone remains and the

marks recorded.

Material and Methods

Material

The studied sample includes 120 limb bones (humeri, radio-ulnas, femurs and tibias) from adult

cows at least 24 months old. A slaughterhouse supplied this experimental series, with 30 specimens of

each element, both left and right (Table 1). Professional butchers defleshed the carcasses. During the

process, they cut the metapodials with cutting pliers. Thus, these bones are absent from the present

experiment. After the reception of the bones, they were stored for less than a week in a fridge at 4 °C.

In addition, the elements broken by individuals 11 and 12 were frozen for 40 days and thawed for three

days in the same fridge before the experiment (temperature: 4°C).

Table 1: Data about the bone element broken by each experimenter (number of ID, element used, side). * These bones were
frozen.

Individual number	Element	Right	Left
1	Humerus	3	7
2	Radio-ulna	6	4
3	Femur	6	4
4	Tibia	2	8
5	Femur	4	6
6	Tibia	4	6
7	Humerus	5	5
8	Femur	6	4
9	Radio-ulna	6	4
10	Tibia	3	7
11	Humerus*	4	6
12	Radio-ulna*	3	7

The experimental series involved 12 volunteers (Table 2) without any experience in bone
breakage: eight men (mean age =34 years old; SD = 11.1 years) and four women (mean age = 31 years
old; SD = 7.1 years). Five individuals had theoretical knowledge of long bone anatomy and one
(individual number 1) was used to flaking lithic tools. Each experimenter broke a series of 10 long bones
of the same element. The number of tests is defined by the order in which each bone is broken one after
the other in a series of 10. To avoid selection biases, the bones were stacked in a disordered pile when
they were presented to the experimenters. Experimenters were isolated from each other so they could
not observe how the others broke the bones. No demonstration was performed. Before the experiment,
they only received one instruction: break the bone to extract the highest quantity and quality of yellow
marrow suitable for consumption. The breakage activity lasted for two to three hours depending on the
bone element and the individual. Experimenters had at their disposal a non-retouched quartzite
hammerstone weighing about 2 kg, a limestone anvil, a plate to deposit the marrow and a wooden stick.
The periosteum was not removed before breakage. Experimenters stopped once they had extracted as
much bone marrow as they could, and then collected it in the plate and weighed it (Vettese 2020).

153 *Table 2: Data about the volunteers for each series (W: Women; M: Men;,Stand. Dev.: Standard Deviation).*

Individual number	Age	Size (cm)	Weight (kg)	Sex	Used hand	Sport Practice	Bone knowledge	Broken bone element
1	29	167	53	W	Right	Yes	Yes	Humerus
2	40	174	82	M	Right	No	Yes	Radio-ulna
3	25	185	78	M	Right	Yes	Yes	Femur
4	30	186	80	M	Right	Yes	No	Tibia
5	51	160	59	W	Left	No	Yes	Femur
6	22	190	95	M	Left	Yes	No	Tibia
7	39	169	63	M	Right	Yes	No	Humerus
8	23	182	95	M	Right	No	Yes	Femur
9	23	167	58	W	Right	Yes	No	Radio-ulna
10	46	169	74	M	Right	Yes	No	Tibia
11	25	158	73	W	Right	Yes	No	Humerus
12	29	178	80	M	Right	Yes	No	Radio-ulna
Average	31.8	173.8	74.2					
Stand. dev.	9.8	10.4	13.7					
Maximum	51	190	95					
Minimum	22	158	53					

Our experiment involved volunteers with no empirical knowledge of recovering marrow using
long bone breakage. Some were archaeologists or palaeontologists, the rest of the experimenters worked
in other fields with no relation to skeletal anatomy or bone tissue properties. The first category of
experimenters were familiar with bone morphology and structure as a study object only. Thus, the
approach to the long bones during marrow recovery was as intuitive as possible, independently of the
experimenters' age or sex. Besides, the volunteers broke a series of ten bones one after another. This
process allowed novice experimenters to self-assess and eventually learn from their mistakes.

After breakage, all the fragmented remains were grouped together in a bag with an identification
code. The bones were boiled during 2 or 3 hours, to remove the grease and to soften the flesh, and then
placed in an oven at 40°C in a solution of water and Papain (papaya enzyme) for 48 hours. Then, they
were soaked in a solution of water and sodium perborate for 24 hours, before being air-dried. The
material is currently kept in the Institut de Paléontologie Humaine (Paris).

Data acquisition

During the experiments, an observer recorded the following variables: the series number,
element laterality, position of the individual, the way the hammerstone was grasped, the position of the
bone, the use of the anvil and the number and location of blows. For these latter, we defined an area as
a long bone portion associated with a side (Figure 1). In order to evaluate the progression of volunteers
during the experiment, from the first try to the end of the operation, we recorded task difficulty

evaluation and the number of blows. We asked the experimenters to auto-evaluate the difficulty encountered during marrow recovery on a scale from 1 to 5, (1 = very easy; 5= very hard), considering mainly the opening of the medullary cavity and marrow extraction. This auto-evaluation resulted in an empirical comparison between long bones. The aim of this auto-evaluation was to assess whether volunteers progressed and developed an efficient method to extract marrow or if tiredness along with the long period of effort affected their capabilities. For quality control purposes, the experimenter and observer evaluated results on a scale from 1 to 5, 1 referring to contaminated marrow containing abundant splinters and 5 being very clean marrow.

Figure 1: Bone areas by portions and sides. Portion 1 (p1): proximal articular end; Portion 2 (p2): proximal diaphysis; Portion 3 (p3): medial diaphysis; Portion 4 (p4): distal diaphysis; Portion 5 (p5): distal articular end and Anterior (a), Posterior (p), Medial (m) and Lateral (l) side (after the drawing by Castel (2010), reference A. Grunwald - © 2016 https://allisongrunwald.wordpress.com/).

The number of identified specimens (NISP), the number of undetermined specimens (NUSP) and the total number of specimens (NSP) were recorded. As the exact location of the percussion marks is necessary, only identified fragments for each bone element were included. Besides, throughout the treatment and the study of each bone, all the remains of one element were kept together, but the NUSP were splinters that we could not refit. Remains that could not be refitted were not considered in the following analysis. The width and length of fragments were measured using a digital caliper. The location of the percussion marks on the bone fragments was analysed according to the GIS protocol developed by Stavrova et al. (2019). The outline of the cortical surface of each identified fragment was

digitized using georeferenced photographic images representing the four-sided visualization of each
bone as a base.

Bone surface modifications were identified and recorded with the naked eye and 15-20x lens.
The centre of each percussion mark was recorded in ArcGIS with a point symbol, based on the
anatomical landmarks of each bone (e.g.,: foramen, crest) (Stavrova et al. 2019). In accordance with the
terminology of Vettese et al. (submitted), five types of percussion marks were recorded: adhering flakes,
crushing marks, flakes, percussion notches, ovoid or triangular percussion pits and grooves. The number
of percussion marks (NPM) and the marks related to percussion (i.e., striations, microstriations and
pseudo-notches) were also noted. These last marks were not taken into account in the total number of
percussion marks. In addition, we excluded the flake because if we had taken into account the flake and
the notch, which were respectively positive and negative for the same percussion mark, we would have
counted the percussion point twice. In this analysis, percussion marks were divided into two groups: 1)
crushing marks, notches and adhering flakes (called "CNA") and 2) pits and grooves (called "Pits").
The "CNA" grouped the percussion marks related to cracking and fractures whereas "Pits" were only
produced by incipient percussion or the rebound effect.

Distances between percussion marks were calculated in ArcGIS and the "Optimized Hot Spot
analysis" tool was used to evaluate and highlight zones with high concentrations of percussion marks.
This tool identifies statistically significant spatial clusters of high concentrations (hot spots) and low
concentration of percussion marks (cold spots). The number of percussion marks recorded for each
series complies with the minimum of 30 features required to conduct the analysis (Stavrova et al. 2019).

The data collected during the experiment were compared to the data analysed in the laboratory
in order to understand which variables of the experimenters' behaviour could influence fragmentation
and the production of percussion marks. Spearman's rho was used to test the correlation between the
data collected during the experiment (i.e., number of blows and marrow weight) and the data recorded
on the bones after treatment (i.e., number of remains and number of percussion marks). Moreover, in
order to evaluate auto-learning and individual improvement, we tested the Spearman correlation of the
number of attempts according to the number of blows for each individual. We presented ρ of Spearman.
To highlight the evolution of the number of blows between the beginning and end of the experiment for
each of the volunteers, we compared, on the one hand, the two averages of the first three and the last
three trials and, on the other hand, the first five and the last five trials using a non-parametric Mann-
Whitney U test. Indeed, the t-test hypothesis was never respected for our data (i.e., the normality and
equivalence of the variance of data). We propose an Efficiency Index calculated by dividing the mass
of marrow extracted from each bone by the number of blows. This Efficiency Index represents the
relationship between a form of expended energy and a form of recovered energy. The higher the
Efficiency Index, the more marrow was extracted with the least number of blows. To test a possible

evolution of the Efficiency Index between the three first and the last three tries and also between the

five first and the last five attempts, we compared the two averages using the Mann-Whitney U Test,

when assumptions for the t-test were not respected. We used the chi-squared test for the scale of auto-

assessment. As Cochran's rule was not respected, we used the Fisher exact test to check the independence

between the scale of marrow quality of the different elements. The correlation coefficients and the tests

were computed using R software (R Core Team, 2019). We use an alpha of 0.05 for all the tests.

Results

Data recorded during the experiment

Bone and individual positions: different ways to break a bone

The position of the bone elements and the position of the volunteers themselves varied during

the breaking process (Table 3). Five different positions chosen by the experimenters were observed. The

majority chose the squatting position (45%), followed by the kneeling position (43%). The standing

position was rarely selected. Only one individual broke bones in a seated cross-legged position

(individual n°2). A minority of the experimenters adjusted their positions during breakage (five

individuals: n°2, 4, 8, 9 and 11; 42%). Individual n°7 remained in the squatting position during the entire

experiment, but nonetheless tried the standing position once at the eighth try, without repeating it.

Volunteer n°10 was the only one who did not adopt and maintain a position. By contrast, four

experimenters developed some kind of habit after a certain number of attempts. Individual n°3, after

three tries, chose the seated position and individual n°6 selected a kneeling position after two tries.

Volunteer n°5 developed a routine from the sixth attempt onwards. For each attempt, she started in a

standing position, then kneeled down, and she finished in a squatting position. Individual n°1 squatted

for all the attempts and half the time, she stood to hit the bone on the anvil. Only one volunteer tested

almost all the positions during the experiment (individual n°12). Nonetheless, after two tries, he

switched back to a squatting or kneeling position.

Regarding the choice of hand to grasp the hammerstone, only one individual used the left

hand, although two volunteers were left-handed and one was ambidextrous. Therefore, the use of both

hands did not seem to be dependent on laterality. All the volunteers breaking the radio-ulna used both

hands at least once to grasp the pebble, which was never the case for the volunteers who broke the tibias.

The individuals who used both hands were both men and women.

We distinguished five ways to position the long bone in order to open its medullar (Table 3).

Ten of the volunteers used the anvil at least once. Two experimenters (individuals n°8 and n°11) did not

use the anvil at all, but positioned the bones directly on the ground. One individual (individual n°7)

selected the batting technique. He hit the bone directly on the anvil with his hands grasping each

epiphysis in order to separate portion 1 from the diaphysis. Then, he laid the largest fragment on the anvil and pursued breakage using the pebble to recover as much marrow as possible. He began using this mixed technique after two tries and repeated it for the next attempts until the end of the series. He judged this mixed technique more efficient than only the hammerstone on anvil technique. One-third (33%) of the volunteers always positioned their bones in the same way during the experiment. Therefore, the majority of the experimenters varied bone position during each try. We noticed that volunteers experienced some difficulties in stabilizing the bone in order to hit it, especially for the radio-ulnas. Sometimes the bone slipped and the blow was less efficient. Some experimenters used an extra stone, their knees or the anvil to block the bone and keep it from slipping.

Table 3: Experimenter and bone position for each bone element, percentages were performed by 30 attempts by elements, and by the 120 attempts for the total.

	Squatting position	Standing position	Seated position	Two kneels postition	One kneel position	Right grasp hand	Left grasp hand	Both grasp hand	On the anvil	One side on the anvil	Batting	On anvil and ground	On the ground
Femur	7 23.3%	4 13.3%	10 33.3%	13 43.3%	2 6.7%	20 66.7%	0 0%	12 40%	18 60%	0 0%	0 0%	1 3.3%	16 53.3%
3			10	3	2	10		2	10				
5	6	4						10	8			1	6
8	1			10		10							10
Humérus	20 66.7%	1 3.3%	3 10%	8 26.7%	0 0%	30 100%	0 0%	9 30%	20 66.7%	3 10%	9 30%	0 0%	10 33.3%
1	9	1		2		10			10				
7	10					10		8	10	3	9		
11	1		1	8		10		1					10
Radio-Ulna	16 53.3%	0 0%	11 36.7%	5 16.7%	3 10%	23 76.7%	0 0%	13 43.3%	16 53.3%	0 0%	0 0%	14 46.7%	10 33.3%
2			10			10		2	10				
9	10					3		10	1			5	10
12	6		1	5	3	10		1	5			9	
Tibia	20 66.7%	0 0%	0 0%	10 33.3%	6 20%	20 66.7%	10 33.3%	0 0%	26 86.7%	2 6.7%	0 0%	11 36.7%	0 0%
4	10					10			10	2			
6				10			10		10			1	
10	10				6	10			6			10	
Total général	63 52.5%	5 4.2%	24 20%	36 30%	11 9.2%	93 77.5%	10 8.3%	34 28.3%	80 66.7%	5 4.2%	9 7.5%	26 21.7%	36 30.0%

The marrow quantity and quality

In order to extract marrow after breakage, most of the volunteers used the wooden stick we provided, but some preferred selecting their own. Only one individual (n°7) used the bone splinters created during breakage. Some also used their fingers. The radio-ulna was the long bone that yielded the smallest quantity of marrow; it represents 13% of the total collected marrow (Table 4). The quantity of marrow extracted from the femur is twice as high (26%). The volunteers who broke the humeri and the tibias extracted a slightly higher quantity of marrow (around 30%). The total quantity of yellow marrow recovered represented 18 kg. We did not notice high variation among volunteers, except for individual n°5 who recovered a significantly smaller quantity of marrow from the femurs than the other experimenters for the femur series (Supplementary Information 1, 2, 3 and 4).

Element	N°indiv	MW (g)		NSP		NISP		NUSP		NB		NPM		Npit		NCNA	
Femur	3	1861	10.20%	259	9.50%	130	9.30%	129	9.80%	641	10.30%	263	5.30%	190	4.80%	73	7.40%
Femur	5	1030	5.60%	202	7.40%	137	9.80%	65	4.90%	496	7.90%	326	6.50%	282	7.10%	44	4.40%
Femur	8	1780	9.80%	222	8.20%	113	8.10%	109	8.20%	179	2.90%	221	4.40%	165	4.10%	56	5.70%
Femur	**Total**	**4671**	**25.60%**	**683**	**25.10%**	**380**	**27.10%**	**303**	**22.90%**	**1316**	**21.10%**	**810**	**16.20%**	**637**	**15.90%**	**173**	**17.50%**
Humerus	1	1710	9.40%	290	10.70%	109	7.80%	181	13.70%	573	9.20%	194	3.90%	72	1.80%	122	12.30%
Humerus	7	1737	9.50%	131	4.80%	77	5.50%	54	4.10%	244	3.90%	304	6.10%	217	5.40%	87	8.80%
Humerus	11	1941	10.60%	195	7.20%	84	6.00%	111	8.40%	377	6.00%	367	7.40%	301	7.50%	66	6.70%
Humerus	**Total**	**5388**	**29.50%**	**616**	**22.60%**	**270**	**19.30%**	**346**	**26.20%**	**1194**	**19.10%**	**865**	**17.30%**	**590**	**14.80%**	**275**	**27.80%**
Radio-ulna	2	799	4.40%	263	9.70%	119	8.50%	144	10.90%	808	12.90%	180	3.60%	49	1.20%	131	13.20%
Radio-ulna	9	775	4.20%	113	4.20%	80	5.70%	33	2.50%	873	14.00%	503	10.10%	428	10.70%	75	7.60%
Radio-ulna	12	811	4.40%	189	6.90%	73	5.20%	116	8.80%	822	13.20%	942	18.90%	894	22.40%	48	4.80%
Radio-ulna	**Total**	**2385**	**13.10%**	**565**	**20.80%**	**272**	**19.40%**	**293**	**22.20%**	**2503**	**40.10%**	**1625**	**32.60%**	**1371**	**34.30%**	**254**	**25.70%**
Tibia	4	1755	9.60%	439	16.10%	192	13.70%	247	18.70%	285	4.60%	280	5.60%	138	3.50%	144	14.50%
Tibia	6	1759	9.60%	175	6.40%	120	8.60%	55	4.20%	753	12.10%	738	14.80%	673	16.80%	65	6.60%
Tibia	10	2290	12.50%	244	9.00%	166	11.90%	78	5.90%	189	3.00%	669	13.40%	590	14.80%	79	8.00%
Tibia	**Total**	**5804**	**31.80%**	**858**	**31.50%**	**478**	**34.10%**	**380**	**28.70%**	**1227**	**19.70%**	**1687**	**33.80%**	**1401**	**35.00%**	**288**	**29.10%**
	Total	**18248**		**2722**		**1400**		**1322**		**6240**		**4987**		**3999**		**990**	

The evaluation of the quality of collected marrow shows a low proportion of bad or very bad assessments (10 %) (Table 5). In addition, most of those who made this judgment broke radio-ulnas. Most (2/3) of the recovered marrow was evaluated as good or very good to consume. None of the volunteers who broke the humerus series evaluated the marrow as very consumable. However, more than a half of the volunteers estimated the tibia marrow to be very good. The Fisher exact tests show that the scales of the tested elements were independent (df = 12, $p < 0.05$).

We noted a significant negative correlation between the quantity of marrow and the number of blows. Indeed, the radio-ulna was the element that received the most blows during the experiment and where a smaller quantity of marrow was recovered. Concerning the radio-ulna, we observed a significant negative correlation between the NISP and the quantity of marrow collected. Besides, we noticed a significant positive correlation between the quantity of recovered marrow and both the NUSP and the number of CNA.

Felt scale	All		Femur		Humerus		Radio-ulna		Tibia	
Very easy	14	11.7%	4	13.3%	2	6.7%	1	3.3%	7	23.3%
Easy	53	44.2%	11	36.7%	21	70%	5	16.7%	16	53.3%
Moderate	28	23.3%	9	30%	4	13.3%	11	36.7%	4	13.3%
Hard	16	13.3%	4	13.3%	1	3.3%	8	26.7%	3	10%
Very hard	9	7.5%	2	6.7%	2	6.7%	5	16.7%	0	0%
Marrow quality scale	All		Femur		Humerus		Radio-ulna		Tibia	
Very bad	3	2.5%	0	0%	3	10%	0	0%	0	0.0%
Bad	9	7.5%	1	3.3%	0	0%	8	26.7%	0	0.0%
Neutral	29	24.2%	7	23%	15	50%	6	20%	1	3.3%
Good	55	45.8%	17	56.7%	12	40%	14	46.7%	12	40%
Very good	24	20%	5	16.7%	0	0%	2	6.7%	17	57%

The level of difficulty experienced during the experiment and the number of blows

In summary, the experimenters found the activity very easy 14 times, easy 53 times, moderate
28 times, difficult 16 times and very difficult only 9 times. Volunteers who broke the humeri did not
encounter any difficulty in most cases (77% easy or very easy). The radio-ulna seems to be more difficult
to break (43% hard or very hard and 37% moderate). None of the volunteers who broke a tibia estimated
the task to be very difficult. They did not find the task any easier or harder as they progressed. In other
words, they did not find the task harder at the beginning because they did not know how to go about it
and then easier because they found an efficient technique. Likewise, exhaustion from the activity did
not influence whether volunteers experienced more difficulty at the end of the breakage series (Table
5). The difficulty encountered according to the bone element varied significantly (Chi-squared test: df
=12, X^2=73.00362, $p<0.01$).

The results of Spearman's correlation analysis between the number of attempts according to the
number of blows for each individual completed these observations. For five individuals (individuals n°1
and n°7 (humerus), individual n°3 (femur) and (individuals n°6 and n°9 (tibia)), the correlation analysis
shows a significant negative relationship (Spearman's rank correlation coefficient results respectively ρ
= -0.81; -0.77; -0.73; -0.93; -0.69). Regarding the other individual series, the results were not significant
(Spearman's rank correlation coefficient results respectively ρ: n°2 = -0.29; n°4 = 0.54; n°5 = -0.2; n°8
= 0.51; n°10 = 0.41; n°11 = -0.44; n°12 = 0.32).

The results of Spearman's correlation analysis between the number of attempts according to the
number of blows for each individual completed these observations. For five individuals (individuals n°1
and n°7 (humerus), individual n°3 (femur) and (individuals n°6 and n°9 (tibia)), the correlation analysis
shows a significant negative relationship (Spearman's rank correlation coefficient results respectively ρ
= -0.81; -0.77; -0.73; -0.93; -0.69). Regarding the other individual series, the results were not significant
(Spearman's rank correlation coefficient results respectively ρ: n°2 = -0.29; n°4 = 0.54; n°5 = -0.2; n°8
= 0.51; n°10 = 0.41; n°11 = -0.44; n°12 = 0.32).

The proportion of the number of blows varied over the last attempts depending on individuals.
Experimenters n° 1, 3, 7, 9 reduced their number of hits by half, on average, n°6 by six between the
three first attempts and the last three (Supplementary Information 1). Volunteer n°12 increased the
average number of blows after the three first tries. However, if we consider the same individual breaking
the same element during the whole experiment, we did not observe a significant difference between the
five first and the last five attempts. We found similar results between the three first and the last three,
except for the humerus (p-value = 0.0254) (Supplementary Information 2).

The bi-plots showing the number of blows and marrow weight did not display any linearity in
the data regarding each element (Supplementary Information 3). When we examine bone element type,
the volunteers who broke the radio-ulnas reveal reduced Efficiency Index compared to the other
elements (Figure 2; EFI all). On the other hand, the experimenters who broke the tibias had a high
average Efficient Index (>5). There was marked variation between the volunteers who broke the humeri
and the femurs, especially for the first three attempts (Figure 2; EFI 3). Between the first three and the
last seven tries, the EFI of the volunteers who broke the tibias varied, like individual n°8. At the
individual level, two-thirds of the volunteers increased their EFI between the first five and last five tries
(n°3, 5, 1, 7, 11, 9, 6, 10). Similarly, more than half the individuals augmented their EFI between the
first and last three (n°3, 1, 7, 11, 2, 9, 6) (Supplementary Information 5). However, regarding all the
experimenters, the Wicoxon signed rank test did not show a significant difference between the first five
and the last five tries or between the first three and the last three tries (p<0.05) (Supplementary
Information 6). We observed similar results regarding each element (p<0.05).

Figure 2: Box plots of the Efficiency Index by Individual during the experiment regarding the third first tries, the last seventh tries and all tries.

Data recorded on the faunal assemblage obtained

Bone fragmentation

The experiments yielded 2,722 bone remains, among which 1,400 were refitted (Table 4). The NSP of the tibias (32%) is higher than for the other elements, which are almost identical. Likewise, the tibia NISP is the highest (34%) and the NISP of the humerus and radio-ulna are the lowest (19%). Nonetheless, we noticed that the average NUSP is around 25% for all the elements. We did not observe marked variability between individuals regarding the NISP, NUSP and the NSP (Supplementary Information 4). However, volunteer n°4 produced more numerous NUSP, which also affected the NSP. According to our application of the Spearman's rank correlation test, the NSP was positively correlated with both the NUSP and the NISP for all the elements (all elements: $\rho = 0.44$ and $\rho = 0.76$; humerus: ρ

= 0.57 and ρ = 0.79; radio-ulna: ρ = 0.39 and ρ = 0.68; femur: ρ = 0.51 and ρ = 0.8; tibia: ρ = 0.51 and

ρ = 0.8). The NUSP and the NISP were also positively correlated between them (all elements: ρ = 0.91;

humerus: ρ = 0.93; radio-ulna: ρ = 0.91; femur: ρ = 0.9; tibia: ρ = 0.9) (Supplementary Information 7).

Number of percussion marks

For one-third of the series, we recorded more percussion marks than the number of blows given

by the experimenters (Table 4). This anomaly was observed for one individual for each bone element.

Three of the individuals who produced a high number of percussion marks used an anvil, but this was

not the case for volunteer n°8. The tibia and radio-ulna series presented numerous percussion marks

(around 33%), corresponding to almost twice as many percussion marks as the femurs and humeri. We

recorded the most abundant pits and grooves on the tibia and the radio-ulna series. The other percussion

marks represented around 27% for all elements, with slightly less on the femurs (18%). Furthermore,

these marks display a lower standard deviation (SD = 33) than the pits (SD = 263) or all the percussion

marks combined (SD = 245). We observed an inter-individual difference, in particular for the first series

of the humeri, radio-ulnas and tibias. A different observer recorded these series.

According to our application of the Spearman's rank correlation test, we noted a significant

positive correlation between percussion marks with the number of pits and grooves on one hand, and

the number of crushing marks, adhering flakes and notches ("CNA"), on the other, for all the long bones

except the tibias ($\rho_{NPM/Npit}$: humerus = 0.95; radio-ulna = 0.96; femur = 0.95; tibia = 0.93; all elements =

0.93; $\rho_{NPM/NCAN}$: humerus = 0.35; radio-ulna = 0.7; femur = 0.35; tibia = -0.07; all elements =0.48). The

number of pits and grooves in the tibia series depended on the number of percussion marks, whereas the

other marks were independent. The number of percussion marks causing a fracture ("CNA") was not

significantly correlated with the other percussion marks (pits) ($\rho_{NCAN/Npit}$: humerus = 0.16; radio-ulna =

0.59; femur = 0.29; tibia = -0.32; all elements = 0.24). However, we observed a positive correlation for

the radio-ulna series and a negative one for the tibia series. The number of percussion marks producing

fractures ("CNA") was positively correlated with the NSP for the humerus series (ρ = 0.34). We

observed a significant positive correlation between the "CNA" percussion marks and the weight of

marrow and the NUSP for femurs and tibias (respectively, femurs: ρ = 0.5; ρ = 0.38 and tibias: ρ = 0.62;

ρ = 0.92). In addition, the number of pits and grooves on the tibias was correlated positively with the

NISP and the NSP (respectively: ρ = 0.5; ρ = 0.38). Finally, the radio-ulnas showed a positive correlation

between percussion marks causing breakage ("CNA") and the number of blows (ρ = 0.39)

(Supplementary data Figure 5).

The graph for the PCA analyses showed more differences between the tibia series on one hand

and the radio-ulna and humerus series, on the other, in particular on the second dimension (Figure 3).

These differences are mainly due to the NSP and the Efficiency Index. On the first dimension, radio-

ulnas and humeri are opposed by the number of blows and by the number of tries.

Figure 3: PCA regarding the number of remains (NSP), the number of percussion marks (NPM), the Efficiency Index and the number of attempts (Series Number) with ellipses at 0.95.

Percussion mark types and spatial distribution

We observed around 5,000 percussion marks on the identified fragments (Table 6). Pits were
the most numerous (78%) and grooves were the least frequent marks (2.6%). Both triangular and ovoid
pits were present, but the former were less numerous. We recorded more pits on the tibias and radio-
ulnas than on the other bone types. Adhering flakes and notches represented 15% of all the percussion
marks. Crushing marks were one of the least numerous traces (4.9 %), and were often located close to
the articular portions. We recorded fewer crushing marks and more grooves on tibias compared to the
other long bones. Conversely, we documented the most numerous crushing marks and adhering flakes
on the humeri. We noted high variation in the number of pits and adhering flakes depending on the
observers who documented the percussion marks. This difference in recording did not influence the
number of other marks to the same degree. We also observed inter-individual variability, in particular
for the radio-ulna series.

Table 6: Type of percussion marks by experimenter and by element on NISP (Total PM: Percussion marks total).

Element/ N°indiv	Percussion notch		Adhering flake		Triangular pit		Ovoid pit		Grooves		Crushing marks		Total PM
Femur	**85**	**10.5%**	**26**	**3.2%**	**41**	**5.1%**	**574**	**70.9%**	**22**	**2.7%**	**62**	**7.7%**	**810**
3	45	17.1%	10	3.8%	9	3.4%	173	65.8%	8	3.0%	18	6.8%	263
5	15	4.6%	5	1.5%	14	4.3%	257	78.8%	11	3.4%	24	7.4%	326
8	25	11.3%	11	5.0%	18	8.1%	144	65.2%	3	1.4%	20	9.0%	221
Humerus	**80**	**9.2%**	**93**	**10.8%**	**95**	**11.0%**	**462**	**53.4%**	**33**	**3.8%**	**102**	**11.8%**	**865**
1	39	20.1%	66	34.0%	8	4.1%	47	24.2%	17	8.8%	17	8.8%	194
7	21	6.9%	18	5.9%	44	14.5%	166	54.6%	7	2.3%	48	15.8%	304
11	20	5.4%	9	2.5%	43	11.7%	249	67.8%	9	2.5%	37	10.1%	367
Radio-ulna	**120**	**7.4%**	**77**	**4.7%**	**625**	**38.5%**	**717**	**44.1%**	**29**	**1.8%**	**57**	**3.5%**	**1625**
2	66	36.7%	39	21.7%	9	5.0%	33	18.3%	7	3.9%	26	14.4%	180
9	39	7.8%	17	3.4%	141	28.0%	275	54.7%	12	2.4%	19	3.8%	503
12	15	1.6%	21	2.2%	475	50.4%	409	43.4%	10	1.1%	12	1.3%	942
Tibia	**174**	**10.3%**	**86**	**5.1%**	**411**	**24.4%**	**943**	**55.9%**	**47**	**2.8%**	**24**	**1.4%**	**1687**
4	93	33.2%	42	15.0%	26	9.3%	93	33.2%	19	6.8%	5	1.8%	280
6	23	3.1%	31	4.2%	224	30.4%	435	58.9%	14	1.9%	11	1.5%	738
10	58	8.7%	13	1.9%	161	24.1%	415	62.0%	14	2.1%	8	1.2%	669
Total	**459**	**9.2%**	**282**	**5.7%**	**1172**	**23.5%**	**2696**	**54.1%**	**131**	**2.6%**	**245**	**4.9%**	**4987**

The crushing marks were often situated close to the articular portions (Figure 4 and 5). We
documented some crushing marks on the medial diaphysis of the humeri and the radio-ulnas and the
tibias of volunteer n°4. Furthermore, we recorded most of the percussion marks on the diaphysis; only
the pits and grooves were located on the articular portions. It was difficult to observe a concentrated
area of percussion marks on the template on which we drew the marks. We merged all the percussion
marks derived from the 10 bone elements into a single series and performed an optimized hot spot
analysis to evaluate high concentration zones of percussion marks (Figure 6, 8, 10, 12). For each long
bone series, we tested all the percussion marks together, apart from percussion marks causing a fracture
("CNA") and the others (pits and grooves) (Figure 7, 9, 11, 13).

Figure 4 : Distribution of percussion marks along the humerus and the radio-ulnas divided by type of percussion mark (red
star: crushing marks, white triangle: pits and grooves and grey circle: notches and adhering flakes); A- Individual n°1; B-
Individual n°7; C- Individual n°11; D- Merged humerus series; F- Individual n°2; G- Individual n°9; H- Individual n°12; I-
Merged radio-ulnas series.

Figure 5: Distribution of percussion marks along the femurs divided by type of percussion mark (red star: crushing marks, white triangle: pits and grooves and grey circle: notches and adhering flakes); A- Individual n°3; B- Individual n°5; C- Individual n°7; D-: Merged femurs series; G- Individual n°4; H- Individual n°6; I- Individual n°10; J- Merged tibias series.

The analyses taking into consideration all the percussion marks generally highlighted cold spots on the articular portions, for all the studied long bones. Conversely, hot spots were generally highlighted on shaft portions. In particular, most of the time the ulna was a cold spot for all the series of radio-ulnas. Likewise, for almost all the series, the tibia crest was a cold spot. The majority of the cold spots were observed on proximal and distal articular portions with the exception of the proximal portion of the radius in the "CNA" analyses) and on the tibia crest and ulnas (Figure 4 and 5).

The spatial analyses show that the majority of the high confidence hot spot zones for the humerus series were on the medial and lateral sides (Figure 6). Experimenter n°11 was an exception with hot spot areas on portion 4 of the posterior and anterior sides. The merged humerus series showed four high confidence hot spot zones on the medial and lateral sides. The hot spot zone on the medial side of portion 2 is the largest, and the one on the lateral side of portion 2 is the smallest. The last ones are located on portion 4 (Figure 6). When we consider the two types of percussion marks, we observe some differences in the location of the high confidence hot spot zone. The "CNA" were on the lateral and medial sides of portion 2 , the pits and grooves were also on the medial side of portion 2, but also on all sides of portion 4 (Figure 7).

Figure 6: Optimized Hot spot analysis of combined percussion marks from the humerus in each series: A- Individual n°1; B- Individual n°7; C- Individual n°11; D- Merged humerus series

467
Figure 7: Optimized Hot spot analysis of combined percussion marks from the humerus in each series: A-E- Individual n°1;
B-F- Individual n°5; C-G- Individual n°8; D-H- Merged humerus series; A, B, C and D: "CNA" Percussion marks and E, F,
G and H: pits and grooves

For the radio-ulna series, we observed hot spot areas on the anterior and posterior sides for all
the individuals (Figure 8). Individual n°9 was the only one who frequently hit portion 3-4 on the medial
side. The analysis of the combined radius-ulna series highlighted a high confidence hot spot zone on the
posterior side of diaphysis portions 2 and 3. This result reflected the general tendencies observed for all
volunteers. The analysis defined a reduced area, portion 2 on the anterior side, as a hot spot zone, which
was influenced by individual n°9. The results of the analysis showed that the high confidence hot spot
zone for the merged radio-ulna was on portion 3 of the posterior side, and on the anterior side portion
3-4 for "CNA" marks only (Figure 9).

Figure 8: Optimized Hot spot analysis of combined percussion marks from the radio-ulnas in each series: A- Individual n°2;
B- Individual n°9; C- Individual n°12; D- Merged radio-ulnas series

483
484 *Figure 9: Optimized Hot spot analysis of combined percussion marks from the radio-ulnas in each series: A-E- Individual n°2;*
485 *B-F- Individual n°9; C-G- Individual n°11; D-H- Merged radio-ulnas series; A, B, C and D: "CNA" Percussion marks and E,*
486 *F, G and H: pits and grooves*

The spatial analyses of the femur series show diversified distribution of the hot spot zones along the diaphysis, depending on the experimenter (Figure 10). For the three series, all sides were a hot spot area at least once, reflecting high inter-individual variability. However, the spatial analysis of the merged femur series shows three confident hot spot zones, quite similar to individual n°8, on portions 2 and 4 on the anterior side, and on portions 3 and 4 on the medial side. High confidence hot spot zones for both percussion mark types are more numerous and dispersed on the merged femur than on the other bones (Figure 11).

Figure 10 Optimized Hot spot analysis of combined percussion marks from the femurs in each series: A- Individual n°3; B-Individual n°5; C- Individual n°7; D- Merged femurs series

Figure 11: Optimized Hot spot analysis of combined percussion marks from the femurs in each series: A-E- Individual n°3;
B-F- Individual n°5; C-G- Individual n°7; D-H- Merged femurs series; A, B, C and D: "CNA" Percussion marks and E, F, G
and H: pits and grooves.

For the tibia series, we observed high confidence hot spot zones on anterior and posterior sides
for each volunteer, in particular on portions 3 and 4 (Figure 12). The lateral side highlighted hot spot
areas for individuals n°6 and n°10, especially on portion 2. The merged tibia series reflects the dispersion
of the high confidence hot spot zones observed for each series. The spatial analyses showed six hot spot
zones, two of which very limited in size. However, the "CNA" area of high confidence hot spot was
only on the posterior side of portions 3 and 4. Zones with pits and grooves were numerous, distributed
on the diaphysis of the lateral, anterior and posterior sides (Figure 13).

Figure 12: Optimized Hot spot analysis of combined percussion marks from the tibias in each series: A- Individual n°4; B-
individual n°6; C- Individual n°10; D- Merged tibias series

Figure 13: Optimized Hot spot analysis of combined percussion marks from the tibias in each series: A-E- Individual n°4; B-
F- Individual n°6; C-G- Individual n°10; D-H- Merged tibias series; A, B, C and D: "CNA" Percussion marks and E, F, G
and H: pits and grooves.

 Discussion

The most significant information highlighted by our experiments is related to the variability

observed during the breakage process and that observed among the bone elements produced during

experimentation. Indeed, some individual behavioural tendencies emerge during the breakage process,

as well as patterns in the production and distribution of percussion marks. Our results also brought to

light the influence of long bone morphology in an intuitiveness context.

How do individual gesture alternatives and bone elements influence breakage methods?

Some volunteers applied reduced force when they hit the bones, especially during the first

attempts. Initially, experimenters tested the resistance of the bones and some were afraid of hurting

themselves. In these cases, they hit the bones with less force. Likewise, at the end of the experiment,

tiredness resulted in a reduction of the force applied. After the activity, most of the experimenters had a

shaking hand, and all the volunteers had aches and pains the following days. This shows that marrow

extraction requires a good physical condition and/or practice and/or good technical skill. During the

activity, some volunteers progressed or developed habits or a routine. However, they did not seem to

become more efficient at optimally breaking bone to extract the maximum quantity of marrow. In

addition, during the experiment, the volunteers mentioned inter-element differences, whereby some

bones were easier to break than others, irrespective of the experience acquired or the degree of

exhaustion. Moreover, it is important to note that sometimes the hammerstone slipped on the cortical

surface or slipped out of the experimenter's hand. This happened especially when the hammerstone and

hands were covered in grease after several tries. When an individual broke several bones, he/she had to

clean the tools or his/her hands to pursue marrow extraction.

One of the most significant results of our experiment was the variability of individual behaviour

during marrow recovery. One volunteer adopted an unexpected posture: a cross-legged seated position.

However, the observed variability only comprised five positions. In addition to these postural variations,

we documented some non-linear behaviour during the experiment. Our experiment showed that non-

trained experimenters could change their position at different moments during both the marrow

extraction from one bone and throughout the ten bone series. However, the majority of the experimenters

kept the same position or developed some kind of habit or routine in their practice. This routine

concerned the individual and bone position and the way of grasping the bone and percussor. The auto-

learning developed by experimenters to be as efficient as possible involved their position and their way

of grasping the bone and percussor. The position influences the amplitude of the gestures, the force

involved in breakage and the location of the blows. The standing position was only selected by women,

and not by all of them. Sex did not influence the choice of posture or position of the bone. No link was

found between the gender of the experimenter and how consistent they were in their posture choice or

in positioning the bone throughout the experiment. Likewise, sex did not seem to influence the way the

percussor was grasped or the use of an anvil. These results could provide some evidence regarding the

organization of butchery task distribution during the Palaeolithic. We noted that the posture adopted by

individuals did not vary in relation to the skeletal element. However, the type of long bones seemed to

affect practices and gestures, in particular for tibias and radio-ulnas. This observation could be related

to the high number of blows applied to these two bone types by novice experimenters.

Although experimenters proceeded differently depending on the elements, we noticed some

divergences in our results, in particular between radio-ulnas and tibias. The experimenters who broke

the radio-ulna series had a very low yield in relation to the number of blows. On the whole, the opposite

is true for the tibia series, for which individuals had a high yield. Thus, volunteers who broke radio-

ulnas found the activity quite difficult and considered the marrow to be of relatively poor quality. On

the other hand, the volunteers who broke tibias found the activity quite easy and the extracted marrow

was considered to be of relatively good quality. Furthermore, experimenters used two hands at least

once for breaking radio-ulnas, but never for tibias. The use of both hands to grasp the pebble was an

expression of the difficulty felt by the individuals breaking radio-ulnas. Regarding the NSP produced

for both these elements, we recorded the highest number of blows and the lowest NSP for the radio-

ulnas and the exact opposite for the tibias of our sample. This was why we observed a high number of

pits and the lowest number of "CNA" percussion marks on radio-ulnas. Contrary to the tibia, the

breakage of a radio-ulna of an animal such as a cow requires some mastery. Numerous studies have

highlighted differences between the volumes of the medullar cavity of these two long bones. Tibias and

femurs comprise large medullary cavities. On the other hand, medullar cavities are smaller for radio-

ulnas and humeri and thus contain a lesser quantity of yellow marrow (Binford, 1981, 1978; Jones and

Metcalfe, 1988). Therefore, the EFI (Efficiency Index) highlighted differences between the energy spent

by counting the blows and the quantity of marrow extracted, in part because of the difference in volume

between the radio-ulna and tibia series. However, medullary cavity volume alone cannot account for all

the differences observed between bones. In terms of density, tibias and radiuses show the highest

diaphysis density of all the long bones used in our experiment (Lam et al., 1999; Lyman, 1994).

However, they seem to react differently to blows. This difference was highlighted by the PCA, showing

an opposition between the radio-ulna and the tibia series regarding the NSP, the Efficiency Index and

the number of percussion marks.

What correlation can be made between blows and percussion marks?

Various researchers have stressed that the aim of mastering percussion breakage was to reduce

the number of blows required to remove all the marrow from the medullary cavity (Capaldo and

Blumenschine, 1994; Galán et al., 2009; Moclán and Domínguez-Rodrigo, 2018). For this reason, it was

expected that novice experimenters would average more blows than experts. Indeed, we noted a high

number of blows, ranging from 6-279, with an average of 52, for the entire bone experiment. If we

compare our experiment with previous works, we observe a huge difference between the average
number and range of blows (Table 7). The average number of blows by volunteers in our experiment
was higher than the maximum in other studies. In particular, for their first tries, two individuals hit the
bones more than 200 times. Besides, none of the experimenters stopped hitting the bones after the first
fracture. The presence of the periosteum on the bones was not sufficient to explain the very high number
of recorded blows. Indeed, during some experiments quoted in Table 7, the periosteum was not removed
(Martin, 1910; Pickering and Egeland, 2006; Rovira Formento, 2010). Novice volunteers tried to break
the element further in order to recover the maximum amount of marrow. The weighing of the marrow
after each attempt motivated some of them. At the individual scale, we observed some increase or
decrease in the number of blows at the beginning and the end of the experiments, but overall, we did not
highlight significant progress for all the experimenters or a particular element. The only exception was
the humerus, for which the number of blows decreased for all the volunteers. The number of blows and
the experimenters' decision to pursue breakage after the first fracture can increase the NSP. Intensive
breakage can reflect a high number of blows in order to access the yellow marrow. In addition, we
observed the quasi-systematic conservation of the epiphyses in their entirety. Indeed, it is not necessary
to break the articular portions for the extraction of yellow marrow only.

Table 7: Listing of different experiment of bone breakage for marrow extraction and comparing data.

References	Species	Element	Range of number of blows	Mean of number of blows	NSP/NME=AverageNSP
(Rovira Formento, 2010)	Horse	Femurs	1-27	8.8	-
(Galán et al., 2009)	Cow	Humerus Femurs Radio-ulna	- - -	- - -	145/22=6.5
(Pickering and Egeland, 2006)	White deer	Radius Humerus	2-15 1-4	6 2	811/38=21.3 472/36=13.1
(Thiébaut et al., 2007)	Cow	Femurs	1-8	3.5	189/13=14.53
(De la Torre et al., 2013)	Cow	Limb bones	-	20	-
(Martin, 1910)	Horse	Femurs Tibia	1-3 3	2 3	-

Interestingly, the number of blows is not correlated with the number of percussion marks.
Comparing the two values, we noted that for two-thirds of the individual bone elements, the number of
blows is higher than the number of percussion marks and more surprisingly, for one-third of the bone
elements, the number of percussion marks is higher than the number of blows. For one experimenter per

bone element (humerus: n°7, radio-ulna: n°12, femur: n°8 and tibia: n°10), the number of blows was

inferior to the number of percussion marks.

In most cases (more than 95%), the number of pits and grooves or "CNA" marks was different

to the number of hits. In an archaeological context, it can thus be impossible to infer the number of

blows from the number of percussion marks. The use of an anvil or the hammerstone could explain the

differences recorded between the number of marks and blows. In their experiment, (Pickering and

Egeland, 2006) also observed a majority of pits among the recorded percussion marks. They concluded

that most of these pits were produced by the counterblow of the anvil. In our experiment, pits were

generally more numerous than the other marks (67 %). Thus, pits and groves result from the rebound

effect caused by the hammerstone or by the anvil due to the counterblow. In this last case, the

counterblow is not a mark; but rather the result of the rebound effect of the bone on the anvil. The marks

due to counterblows include pits, microstriations, outer conchoidal scars or notches (e.g.,: Boulestin,

1999; Capaldo and Blumenschine, 1994; Gifford-Gonzalez, 2018; Johnson, 1985; Lyman, 1994;

Outram, 2002), located on the opposite side to the blow. Due to the multiplicity of traces and the

overlapping of the traces produced by the hammerstone or when the long bone is batted against a hard

surface, it is difficult to discriminate blow marks from counterblow marks. In addition, we recorded

some triangular pits and grooves even though experimenters only used non-retouched pebbles. These

marks, characterized by straight and sharp edges, could be the product of the rebound effect from the

anvil. Therefore, the use of an anvil could create supplementary marks during the blow. Two volunteers

(n°8 and 11) did not use the anvil at all, but simply laid the bones on the ground. Nonetheless, some

triangular pits and grooves were identified (around 10 %). However, we sometimes recorded more marks

than blows. Thus, the counterblow is not the sole explanation for the difference between the number of

blows and the number of percussion marks. It is possible that the shape of the hammerstone, the impact

with the ground or the collision of bones between them during the activity created several pits. More

than one-quarter of the recorded pits were very small and superficial, with a diameter of 10 mm or less

(28%). In an archaeological context, the different taphonomical processes, such as trampling, for

example (Domínguez-Rodrigo et al., 2010), can erase these pits or produce others. For the remaining

two-thirds of the broken elements, we recorded a lower number of percussion marks than blows. In our

protocol, the periosteum was never removed. The presence of the periosteum, which absorbed some of

the force of the blows, could partly explain this difference.

Percussion marks distribution

The areas where the least percussion marks were observed, apart from the articular portions,

were the anterior crest of the tibia and the diaphysis of the ulna. The distribution of percussion marks

showed that novice experimenters avoided shape constraints, such as bone protuberances. They seemed

to prefer relatively flat areas. Thus, the spatial analyses for tibias and radio-ulnas highlighted significant

hot spot zones for the merged series, where bone shape was the flattest. Furthermore, for the "CNA"

marks, a single hot spot area was most impacted on the tibias. This area was located on the posterior side of the distal diaphysis. It was one of the flattest and smoothest zones of the tibias with no muscular lines. Moreover, the distal diaphysis is the least dense portion of the tibia (index density = 44) (Lyman, 1994). Regarding the "CNA" marks, two specific high confidence hot spot zones were identified on the radio-ulnas. They were located on relatively flat and easily accessible areas to blows. The first one is situated on the anterior side of the medial shaft, which is relatively flat considering the globally very arched shape of the radio-ulna. The radio-ulna section in this area is flattened compared with portions 3 and 4, which present a semi-lunar crescent shape. When the bone was placed on a small anvil, one of the best ways to stabilize the radio-ulna was to place the medial shaft on the posterior side. Thus, the anterior side was exposed and very accessible. The other hot spot area was on the exact opposite side because, on the ground, it was easier to place the bone on the anterior side to avoid interference from the ulna. In addition, when the radio-ulna was positioned on the anterior side, this area was located at the inflexion point of the bone curve and was the most accessible to hit. Furthermore, these areas were located on portions 2 and 3, which are the densest part of the radius (marrow index = 0.56 & 0.62).

Unlike zeugopods, the diaphyseal section of humeri and femurs is rather cylindrical. Their diaphyses are relatively straight in comparison to the radio-ulna. The articular portion shows important differences in terms of stability. The humerus is more stable on its lateral and medial sides whereas the femur is relatively stable on all sides. Indeed, the very rounded shape of the humerus head causes instability when placed on its posterior side. This is why the distribution of percussion marks, highlighted by the hot spot analyses, is concentrated on the lateral and medial sides. Pit and groove marks are generally distributed on the distal part of the bone. This part of the humerus diaphysis is denser than the others (density index = 0.48). Conversely, the hot spot area of "CNA" marks is only located on the proximal shaft portion, in the least dense part of the bone (index density = 0.25). These "CNA" marks represent one of the most the numerous hot spot zones. In addition, these hot spot areas were similar for the three merged spatial analyses of the femurs (all percussion marks, "CNA" marks and pits and groove marks). All the diaphysis portions were hot spot zones at least once, independently of density variability.

Thus, percussion mark location is influenced by the morphological constraints of bones, in particular for the radio-ulna, the tibia and the humerus, in an intuitive context. Volunteers were faced with bone marrow extraction for the first time and were highly influenced by shape constraints. These constraints were so strong that volunteers continued hitting these zones after many attempts, as bone stability and the accessibility of these areas facilitated breakage and therefore marrow recovery. Nonetheless, the absence of morphological constraints seems to highlight novice choices. Therefore, based on the results of our analyses, in an intuitive context, femurs seem to be the bones for which individual variability was highest.

Archaeological, experimental and ethnographical data comparisons: some convergences

Binford (1981) described the practices and gestures of the Nunamiut in order to infer past hunter-gatherer behaviour from archaeological assemblages. His observations describe systematic breakage, hitting the same bone portions each time. Moreover, in the same way as experiments, ethnographical studies highlight wide-ranging variables, which are hardly accessible in archaeological contexts.

We highlighted the existence of a preferential pattern for the intuitive breakage of long bones, using the spatial analysis of the location of percussion marks on radio-ulnas, tibias and humeri. Archaeological and ethnographical studies focusing on the distribution of percussion marks along long bone diaphyses have revealed the existence of preferentially impacted areas (Binford, 1981; Blasco et al., 2013; Masset et al., 2016; Moclán and Domínguez-Rodrigo, 2018; Vettese et al., 2017). The comparison between these results and ours underlines the influence of bone shape on some systematic practices of extant and past hunter-gatherers (Figure 14). For radio-ulnas, archaeologists record percussion marks on the anterior side, in particular in the faunal assemblages of level XVIIc of Bolomor Cave (Blasco et al., 2013) and at the Abri du Maras layer 4.1 (Vettese et al., 2017), where some tendencies were observed (Middle Palaeolithic), Combe Saunière and Cuzoul de Vers (Castel, 1999) and the "upper unit" of Saint-Germain-la-Rivière (Costamagno, 1999; Masset et al., 2016) and the Rond du Barry (Upper Palaeolithic), and also among the extant Nunamiut (Binford, 1981). The areas highlighted by hot spot analyses were similar to the areas of percussion mark concentrations in these comparative examples. These faunal assemblages represent practically all the Palaeolithic sites for which percussion mark distribution has been studied. The aforementioned faunal assemblages are dominated by different species: reindeer, saiga antelope, ibex, red deer or larger-sized ungulates, like horse or bison, and in our experiment, cow. This comparison suggests that the high shape constraints of the radio-ulnas strongly influenced the distribution of percussion marks on whole long bone remains in an archaeological level independently of ungulate size. Furthermore, for ungulates, the fact that the radius is merged with the ulna constrains the morphological shape of the bones. However, level IV of Bolomor Cave and the "lower unit" of Saint-Germain-la-Rivière showed different patterns on the lateral side for the former site and both lateral and medial sides for the latter (Figure 14).

For the humerus, the distribution of percussion marks was quite similar to our observations. Almost all the sites presented in the work of Masset et al. (2016) reveal a pattern on the medial and/or lateral sides, with the sole exception of Bolomor Cave level IV. Compared to the other sites, we noted variability in the exact area impacted by percussion marks, but we observed a preference for portion 4 in the majority of the assemblages. A minority of the studied tibia series showed a systematic pattern or strong tendencies. The posterior sides displayed the most areas of percussion mark concentration.

Finally, most of the femur series from the archaeological sites studied do not seem to show systematic distribution, with the exception of level VI of Bolomor, which indicates two preferentially impacted zones, and the Abri du Maras, where one area was more impacted than the other. We observe similarities with the Nunamiut for the anterior side.

Figure 14 : Location of percussion marks (grey areas) for the humerus, radio-ulna, femur, and tibia for sites from the Middle Paleolithic: Bolomor cave (ungulates: Blasco et al., 2013b.) and Abri du Maras (middle sized-ungulates: Vettese et al. 2017); the Upper Paleolithic: Combe Saunière (reindeer, Solutrean: Castel, 1999), Cuzoul de Vers (reindeer, Badegoulian: Castel, 1999), Saint-Germain-la-Rivière, Lower Unit (saiga antelope, Lower Magdalenian: Costamagno, 1999) and Upper Unit (saiga antelope: Masset et al. 2016), Rond du Barry (ibex, Upper Magdalenian: Costamagno, 1999), for these series the data regarding the femurs were missing and most of the tibias did not present higher tendencies; Current: Nunamiut people of Alaska (reindeer: Binford, 1981 and the results of our experiment regarding the CNA).

Conclusion

For the purposes of testing intuitiveness in bone breakage, we experimentally tested the influence of individual behaviour and long bone morphology on percussion mark location. We observed a high variability of gestures during the breakage process. Most of the volunteers developed their own routine. This routine included several positions and different techniques, involving variability in the applied force and different bone element responses. Nevertheless, our results highlight similarities in percussion mark distribution despite the specificity of each experimenter, for most of the long bones studied here. Indeed, one of the most important results is the identification of specific hot spot areas, regardless of the variability of experimenters' behaviour during the breakage process. These similarities can be considered as intuitive patterns.

The location of percussion marks is influenced by numerous variables depending on: the experimenters' behaviour and skills, the techniques employed, the location of the blows, the morphology of the skeletal element and bone position during breakage. Nevertheless, our results highlighted the

predominance of two main factors in an intuitive context: long bone morphology for the humerus, radio-
ulna and tibia and experimenters' behaviour for the femur.

From a different perspective, we have shown that bone response can be very different in an
intuitive context, notably for the radio-ulna and the tibia. These differences can be observed at almost
all the different levels of the analysed data: volunteer behaviour or perception, marrow quality and
quantity, number of blows and remains produced and type of percussion marks recorded. These results
highlight the diametrically opposed differences between tibias and radio-ulnas in an intuitive context.

Our results shed new light on the importance of gestures in the breakage process. It could prove
interesting to model these results and evaluate their amplitude and the force involved. Numerous studies
focus on the force required to break long bones but generally concentrate on pressure methods.
Consequently, data pertaining to the percussion process are lacking. In addition to measuring the force
applied by the blow, it is essential to evaluate the force of the counter-blow caused by different materials,
such as stone, bone or wood.

Percussion marks are very instructive in an archaeological context to access past subsistence
behaviour. Our results show that percussion traces on fossil bones are only the tip of the iceberg. They
only reveal a small window into past bone marrow extraction practices. This experiment highlighted the
variability of volunteers' behaviour and the different bone responses during fracturing in order to
propose new hypotheses on past butchering practices based on the comparison of the different data
analysed in this work.

To conclude, the influence of bone morphology on the distribution of percussion marks in the
archaeological record should be taken into account before inferring cultural traditions. It is essential to
compare the intuitive patterns highlighted by spatial analysis with the patterns identified in
archaeological levels. It could also prove interesting in archaeological contexts to carefully consider
percussion mark distribution on the femur as this could be the most instructive bone element, in terms
of marrow recovery methods and group specificity.

Acknowledgements

We express our gratitude to CHARAL S.A.S. who kindly provided all the bones used in the preparation

of this experiment. We are grateful for the funding of the Fondation Nestlé France. Many thanks to Eric

Pellé and Zoé Thalaud of the Service de Préparation Ostéologique et Taxidermique (SPOT) from the

Muséum national d'Histoire naturelle in Paris, for the osteological preparation of all the long bones. We

also wish to thank Laetitia Demay for her precious advice during proofreading and her valuable help

during the experiment. Laurent Crépin and colleagues provided helping logistical advice and assistance

during the experiment. We are deeply grateful to the volunteers for their selfless efforts in helping with

our experiment. We also thank Lou Albessard, Jeremy Duveau and Guilhem Mauran for their precious

advice. L. Byrne, an official translator and native English speaker, edited the English manuscript.

Financial Disclosure

This project was supported by the Fondation Nestlé France (SJ 671-16) (https://fondation.nestle.fr/); the

Centre d'Information des Viandes – Viande, sciences et société (SJ 334-17); the Muséum national

d'Histoire naturelle.

Disclosure statement

No potential conflict of interest was reported by the authors.

References

Abe, Y., 2005. Hunting and Butchery Patterns of the Evenki in Northern Transbaikalia, Russia. Stony
brook University.

Alhaique, F., 1997. DO PATTERNS OF BONE BREAKAGE DIFFER BETWEEN COOKED AND UNCOOKED
BONES ? AN EXPERIMENTAL APPROACH. Anthropozoologica 25–26, 49–56.

Benazzi, S., Blasco, R., Fiorenza, L., Benazzi, S., Henry, A.G., Salazar-garc, D.C., 2015. To Meat or Not to
Meat ? New Perspectives on Neanderthal Ecology. Yearb. Phys. Anthropol. 156, 43–71.
https://doi.org/10.1002/ajpa.22659

Binford, L.R., 1981. Bones: Ancient Men and Modern Myths., Academic P. ed. New-York.

Binford, L.R., 1978. Nunamiut: Ethnoarchaeology, Academic P. ed. New-York, Chicago, London.

Blasco, R., Domínguez-Rodrigo, M., Arilla, M., Camarós, E., Rosell, J., 2014. Breaking Bones to Obtain
Marrow: A Comparative Study between Percussion by Batting Bone on an Anvil and Hammerstone
Percussion. Archaeometry 56, 1085–1104. https://doi.org/10.1111/arcm.12084

Blasco, R., Fernández Peris, J., 2012. Small and large game: Human use of diverse faunal resources at
Level IV of Bolomor Cave (Valencia, Spain). Comptes Rendus - Palevol 11, 265–282.
https://doi.org/10.1016/j.crpv.2012.01.003

Blasco, R., Rosell, J., Domínguez-Rodrigo, M., Lozano, S., Pastó, I., Riba, D., Vaquero, M., Peris, J.F.,
Arsuaga, J.L., de Castro, J.M.B., Carbonell, E., 2013. Learning by Heart: Cultural Patterns in the Faunal
Processing Sequence during the Middle Pleistocene. PLoS One 8, e55863.
https://doi.org/10.1371/journal.pone.0055863

Blumenschine, R.J., Selvaggio, M.M., 1988. Percussion marks on bone surfaces as a new diagnostic of
hominid behaviour. Nature 333, 763–765.

Brugal, J.P., Defleur, A., 1989. Approche expérimentale de la fracturation des os des membres de
grands mammifère. Artefacts 7, 14–20.

Capaldo, S.D., Blumenschine, R.J., 1994. A Quantitative Diagnosis of Notches Made by Hammerstone
Percussion and Carnivore Gnawing on Bovid Long Bones. Am. Antiq. 59, 724–748.

Castel, J.C., 1999. Castel, J. C. (1999). Comportements de subsistance au Solutréen et au Badegoulien
d'après les faunes de Combe Saunière (Dordogne) et du Cuzoul de Vers (Lot). Université Bordeaux I.

Costamagno, S., 1999. Stratégies de Chasse et Fonction des Sites au Magdalénien dans le Sud de la
France. Unniversité Bordeaux I.

Costamagno, S., David, F., 2009. Comparaison des pratiques bouchères et culinaires de différents
groupes sibériens vivant de la renniculture. Archaeofauna 18, 9–25.

Costamagno, S., Rigaud, J.-P., 2014. L'exploitation de la graisse au Paléolithique. Hist. l'alimentation
Hum. entre choix contraintes 134–152.

Daujeard, C., Fernandes, P., Guadelli, J.L., Moncel, M.H., Santagata, C., Raynal, J.P., 2012. Neanderthal
subsistence strategies in Southeastern France between the plains of the Rhone Valley and the mid-
mountains of the Massif Central (MIS 7 to MIS 3). Quat. Int. 252, 32–47.
https://doi.org/10.1016/j.quaint.2011.01.047

Daujeard, C., Vettese, D., Britton, K., Béarez, P., Boulbes, N., Crégut-Bonnoure, E., Desclaux, E., Lateur,
N., Pike-Tay, A., Rivals, F., Allué, E., Chacón, M.G., Puaud, S., Richard, M., Courty, M.-A., Gallotti, R.,
Hardy, B., Bahain, J.J., Falguères, C., Pons-Branchu, E., Valladas, H., Moncel, M.-H., 2017. Neanderthal
selective hunting of reindeer? The case study of Abri du Maras (south-eastern France). Archaeol.

Anthropol. Sci. 1–27. https://doi.org/10.1007/s12520-017-0580-8

De la Torre, I., Benito-calvo, A., Arroyo, A., Zupancich, A., Proffitt, T., 2013. Experimental protocols for the study of battered stone anvils from Olduvai Gorge. J. Archaeol. Sci. 40, 313–332. https://doi.org/10.1016/j.jas.2012.08.007

Domínguez-Rodrigo, M., Pickering, T.R., Bunn, H.T., 2010. Configurational approach to identifying the earliest hominin butchers. Proc. Natl. Acad. Sci. 1–6. https://doi.org/10.1073/pnas.1013711107

Enloe, J.G., 1993. Ethnoarchaeology of marrow cracking: implications for the recognition of prehistoric subsistence organization., in: Hudson, J. (Ed.), From Bones to Behavior: Ethnoarchaeological and Experimental Contributions to the Interpretation of Faunal Remains. Southern Illinois University, Carbondale, pp. 82–100.

Galán, B., Rodríguez, M., de Juana, S., Domínguez-Rodrigo, M., 2009. A new experimental study on percussion marks and notches and their bearing on the interpretation of hammerstone-broken faunal assemblages. J. Archaeol. Sci. 36, 776–784. https://doi.org/10.1016/j.jas.2008.11.003

Hardy, K., Brand-miller, J., Brown, K.D., Thomas, M.G., Copeland, L., 2006. The Importance of Dietary Carbohydrate in Human Evolution. Quaterly Rev. Biol. 90 (3), 251–268.

Holt, S.H., Eaton, S.B., Miller, J.B., Cordain, L., Speth, J.D., Mann, N., 2018. Plant-animal subsistence ratios and macronutrient energy estimations in worldwide hunter-gatherer diets. Am. J. Clin. Nutr. 71, 682–692. https://doi.org/10.1093/ajcn/71.3.682

Joana, M., Cáceres, I., Rosell, J., Saladié, P., Vallverdú, J., 2014. From small bone fragments to Neanderthal activity areas : The case of Level O of the Abric Romaní (Capellades , Barcelona , Spain). Quat. Int. 330, 36–51. https://doi.org/10.1016/j.quaint.2013.12.015

Jones, K.T., Metcalfe, D., 1988. Bare bones archaeology: Bone marrow indices and efficiency. J. Archaeol. Sci. 15, 415–423. https://doi.org/10.1016/0305-4403(88)90039-8

Karr, L.P., Outram, A.K., 2012. Tracking changes in bone fracture morphology over time : environment , taphonomy , and the archaeological record. J. Archaeol. Sci. 39, 555–559. https://doi.org/10.1016/j.jas.2011.10.016

Kuntz, D., Costamagno, S., Feyfant, L., Martin, F., 2016. The exploitation of ungulates in the Magdalenian in the Entre-Deux-Mers (Gironde, France). Quat. Int. 414, 135–158. https://doi.org/10.1016/j.quaint.2015.12.079

Lacroix, M.N., 2011. A Study of the Impact of Weathering Upon the Minimal Force Required To Fracture Bone. Boston.

Lam, Y.M., Chen, X., Pearson, O.M., 1999. Intertaxonomic Variability in Patterns of Bone Density and the Differential Representation of Bovid , Cervid , and Equid Elements in the Archaeological Record. Soc. Am. Archaeol. 64, 343–362. https://doi.org/10.2307/2694204

Lyman, L.R., 1994. Vertebrate Taphonomy. Cambridge University Press, Cambridge.

Marín, J., Saladié, P., Rodríguez-Hidalgo, A., Carbonell, E., 2017. Neanderthal hunting strategies inferred from mortality profiles within the Abric Romaní sequence, Plos One. https://doi.org/10.1371/journal.pone.0186970

Martin, H., 1910. La Percussion osseuse et les esquilles qui en dérivent . Expérimentation. Bull. la Soc. Prehist. Fr. 5, 299–304.

Masset, C., Costamagno, S., Cochard, D., Laroulandie, V., 2016. La fracturation osseuse : du fait technique à l'essai d'interprétation sociétale L'exemple de l'antilope saïga du gisement magadelenien

de Saint-Germain -la-Rivière (Gironde). Bull. la Société préhistorique française 113, 691–712.

Moclán, A., Domínguez-Rodrigo, M., 2018. An experimental study of the patterned nature of
anthropogenic bone breakage and its impact on bone surface modification frequencies. J. Archaeol.
Sci. 96, 1–13. https://doi.org/10.1016/j.jas.2018.05.007

Noe-Nygaard, N., 1977. Butchering and Marrow Fracturing as a Taphonomic Factor in Archaeological.
Paleontol. Soc. 3, 218–237.

O'Connell, J.F., Hawkes, K., Blurton-Jones, N.G., 1992. Patterns in the distribution, site structure and
assemblage composition of Hadza kill-butchering sites. J. Archaeol. Sci. 19, 319–345.
https://doi.org/10.1016/0305-4403(92)90020-4

Oliver, J.S., 1993. Carcass processing by the Hadza: bone breakage from butchery to consumption, in:
Hudson, J. (Ed.), From Bones to Behavior: Ethnoarchaeological and Experimental Contributions to the
Interpretation of Faunal Remains. Occasional Paper No. Center for Archaeological Investigations,
Southern Illinois University, Carbondale, pp. 200–227.

Outram, A.K., 2002. Bone Fracture and Within-bone Nutrients : an Experimentally Based Method for
Investigating Levels of Marrow Extraction, in: Miracle, P., Milner, N. (Eds.), Consuming Passions and
Patterns of Consumption. Cambridge.

Patou-Mathis, M., 2000. La chasse chez les! Kung: San du Nord-Ouest du Kalahari, Botswana.
Anthropol. préhistoire 111, 344–354.

Patou-Mathis, M., 1985. La fracturation des os longs de mammifères: élaboration d'un lexique et d'une
fiche type. Artefacts 1, 11–22.

Pickering, T.R., Egeland, C.P., 2006. Experimental patterns of hammerstone percussion damage on
bones: Implications for inferences of carcass processing by humans. J. Archaeol. Sci. 33, 459–469.
https://doi.org/10.1016/j.jas.2005.09.001

R Core Team, 2019. R: A language and environment for statistical computing. [WWW Document].
Found. Stat. Comput. Vienna, Austria.

Romandini, M., Nannini, N., Tagliacozzo, A., Peresani, M., 2014. The ungulate assemblage from layer
A9 at Grotta di Fumane, Italy: A zooarchaeological contribution to the reconstruction of Neanderthal
ecology. Quat. Int. 337, 11–27. https://doi.org/10.1016/j.quaint.2014.03.027

Rovira Formento, M., 2010. Aproximación experimental a la explotación de huesos largos de grandes
animales para la recuperación de la médula ósea y su aplicación arqueológica al registro faunístico del
Nivel 3 colluvio de Isernia La Pineta (Molise , Italia). Arqueologia del Quaternari i Evolució Humana.

Speth, J.D., 2010. The Paleoanthropology and Archaeology of Big-Game Hunting Protein, Fat, or
Politics?, Springer. ed.

Speth, J.D., 1989. Early hominid hunting and scavenging: the role of meat as an energy source. J. Hum.
Evol. 18, 329–343. https://doi.org/10.1016/0047-2484(89)90035-3

Speth, J.D., Meignen, L., Bar-Yosef, O., Goldberg, P., 2012. Spatial organization of Middle Paleolithic
occupation X in Kebara Cave (Israel): Concentrations of animal bones. Quat. Int. 247, 85–102.
https://doi.org/10.1016/j.quaint.2011.03.001

Speth, J.D., Spielman, K.A., 1983. Energy source, protein metabolism, and hunter-gatherer subsistence
strategies. J. Anthropol. Archaeol. 2 (1), 1–31.

Stiner, M.C., Gopher, A., Barkai, R., 2011. Hearth-side socioeconomics, hunting and paleoecology
during the late Lower Paleolithic at Qesem Cave, Israel. J. Hum. Evol. 60, 213–233.

https://doi.org/10.1016/j.jhevol.2010.10.006

Thiébaut, C., Claud, É., Coudenneau, A., Coumont, M.-P., Asselin, G., Beauval, C., Chacón, G., Sandrine,
C., Daulny, L., Gerbe, M., Mallye, J.-B., Maury, S., Mourre, V., Plisson, H., Provenzano, N., Streit, L.,
2007. Des traces et des Hommes.

Valensi, P., Michel, V., El Guenouni, K., Liouville, M., 2013. New data on human behavior from a 160 ,
000 year old Acheulean occupation level at Lazaret cave , south-east France : An archaeozoological
approach. Quat. Int. 316, 123–139. https://doi.org/10.1016/j.quaint.2013.10.034

Vettese, D., 2020. An archaological experiment focused on the intuitive way of long bones breakage
to extract marrow. **protocols.io**
dx.doi.org/10.17504/protocols.io.bb6airae

Vettese, D., Daujeard, C., Blasco, R., Borel, A., Caceres, I., Moncel, M.H., 2017. Neandertal long bone
breakage process : Standardized or random patterns ? The example of Abri du Maras (Southeastern
France , MIS 3). J. Archaeol. Sci. Reports 13, 151–163. https://doi.org/10.1016/j.jasrep.2017.03.029

Villa, P., Mahieu, E., 1991. Breakage pattern of human long bones. J. Hum. Evol. 21, 27–48.

Yravedra, J., Maté-González, M.Á., Palomeque-González, J.F., Aramendi, J., Estaca-Gómez, V., San Juan
Blazquez, M., García Vargas, E., Organista, E., González-Aguilera, D., Arriaza, M.C., Cobo-Sánchez, L.,
Gidna, A., Uribelarrea Del Val, D., Baquedano, E., Mabulla, A., Domínguez-Rodrigo, M., 2017. A new
approach to raw material use in the exploitation of animal carcasses at BK (Upper Bed II, Olduvai Gorge,
Tanzania): a micro-photogrammetric and geometric morphometric analysis of fossil cut marks. Boreas.
https://doi.org/10.1111/bor.12224